

 Open access • Journal Article • DOI:10.1038/SJ.BDJ.2018.444

The treatment need and associated cost of erosive tooth wear rehabilitation - a service evaluation within an NHS dental hospital. — [Source link](#)

Saoirse O'Toole, Mark Pennington, S Varma, David Bartlett

Institutions: King's College London, Guy's Hospital

Published on: 08 Jun 2018 - British Dental Journal (Nature Publishing Group)

Topics: Tooth wear

Related papers:

- [Basic Erosive Wear Examination \(BEWE\): a new scoring system for scientific and clinical needs](#)
- [Prevalence of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults](#)
- [An index for measuring the wear of teeth](#)
- [Evaluation of the basic erosive wear examination \(BEWE\) for use in general dental practice.](#)
- [The signs and symptoms of tooth wear in a referred group of patients.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/the-treatment-need-and-associated-cost-of-erosive-tooth-wear-1am09bgulj>



King's Research Portal

DOI:

[10.1038/sj.bdj.2018.444](https://doi.org/10.1038/sj.bdj.2018.444)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

O'Toole, S., Pennington, M., Varma, S., & Bartlett, D. W. (2018). The treatment need and associated cost of erosive tooth wear rehabilitation –: - a service evaluation within an NHS dental hospital. *British Dental Journal*, 224(12), 957–961. <https://doi.org/10.1038/sj.bdj.2018.444>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The treatment need and associated cost of erosive tooth wear
rehabilitation – a service evaluation within an NHS dental hospital.

Abstract

Aim: To establish the average treatment need and cost of prosthodontic rehabilitation of severe erosive tooth wear within an NHS hospital setting.

Methods: The clinical notes of patients referred and accepted for treatment to King's College London Dental Institute specialist restorative clinics by their GDP between 01/01/2014 and 01/01/2016 for severe erosive tooth wear were audited. The first 30 patients with completed treatment plans were audited and the following was recorded: age, gender, BEWE score upon presentation, the location and aetiology of the wear, presenting complaint, number of clinical sessions, number of treatment planning sessions, the treatment provided, and materials used. The cost of care including materials and staff costs were estimated. Cost of treatment within a private setting was estimated based on private fees from three practices in the London area.

Results: The average treatment time for patients was 20.8 months (SD 9.6, Range 8-44 months). The average number of clinical visits during this time was 24.3 (SD 12.7; 8-48). The mean total cost per completed treatment plan was £2,371 (SD £1,290: £675-£4,807 which included mean staff costs of £1,333 (SD £697: £439-£2,637) and £1,039 (SD £668: £199-£2,500) for materials and laboratory work. The estimated cost of similar treatment provided by a specialist in private clinic was £13,353 (SD £6,905; £4,737-£31,224) per patient. The only predictor of costs was the presence of wear on both anterior and posterior surfaces.

Conclusion: The prosthodontic rehabilitation of erosive tooth wear is complex, interdisciplinary and costly to the NHS.

Introduction

The treatment of erosive tooth wear is not straightforward. It involves multiple, often interdisciplinary planning visits, prolonged care and usually changing the occlusal vertical dimension (OVD).

Within the UK, patients with severe tooth wear are either referred to secondary specialist care, specialist teaching hospitals or attend a prosthodontist privately, or rely on their general dental practitioner to provide the best care they can. The decision to treat erosive tooth wear is made on relatively arbitrary criteria. A survey of 124 secondary care referrals to a Scottish dental hospital reported that the reasons for patients requesting tooth wear rehabilitation were aesthetics (54%), followed by pain/sensitivity (25%) and then functional problems (12%)¹. Of these referrals, only 8% were accepted for secondary care treatment. On the other hand, severe erosive tooth wear has been shown to impact on quality of life^{2,3} with one study reporting that it was equivalent to that of being edentulous². The treating practitioner and patient are faced with difficult decisions juggling the cost, the complexity and commitment to treatment planning. There is also the need to justify possible further removal of tooth structure for placement of long lasting restorations to achieve patient expectations.

The UK is not alone in this difficult decision-making process. Danish colleagues have devised a “Tooth Wear Evaluation System” which is taught to their dental students to aid in the diagnosis and management planning of the worn dentition⁴. This comprehensive evaluation considers primary factors such as the amount of tooth wear, the surfaces and the number of teeth affected, in addition to secondary factors such as the speed of progression, the age of the patient and the aetiology. However, the authors repeatedly recognise that no clear cut off criteria can be defined.

Monitoring tooth wear is an appropriate care pathway⁴⁻⁶, but there is no objective, quantitative method of measuring erosive tooth wear progression in general dental practice. Often direct composite restorations are seen as non-invasive methods of improving aesthetics and restoring the OVD. However, recent reports have shown that the complication rates can be high as composites are prone to fracture/discolouration; consequently, maintenance levels are high⁶⁻⁸.

If a patient opts to undergo a full mouth rehabilitation, they should be informed about the treatment time and commitment involved. Treatment required is likely to be extensive and hence costly. However, there is little data, aside from anecdotal information, to allow the patient to make a fully informed treatment decision. Both the patient and the practitioner should be aware of these costs prior to commencing an extensive treatment plan.

The aim of this retrospective service evaluation was to establish the average treatment need and cost of prosthodontic rehabilitation of severe erosive tooth wear. We aimed to provide detailed estimates of costs in a primary and tertiary care setting, in addition to estimates of cost had such care been delivered in a private setting.

Methods

This service evaluation was registered with Guy's and St Thomas's Hospital Trust audit bank, project number 7494. The study population were those who had been referred to King's College London Dental Institute specialist restorative clinics by their GDP between 01/01/2014 and 01/01/2016 for severe erosive tooth wear. The inclusion criteria were: minimum of 6 teeth in each jaw, referred by their regular general dental practitioner, acceptance for treatment by postgraduate prosthodontic trainees and treatment plans were completed by the 31/08/17. The exclusion criteria were provision of implants in the treatment planning and underlying dental conditions which would influence severity of wear or long-

term prognosis of restorations eg amelogenesis imperfecta, hypoplastic enamel, hypodontia. The clinical notes of the first 30 patients identified were audited and the following was recorded: the age, gender and BEWE score upon presentation, the location and aetiology of the wear, the presenting complaint, number of clinical sessions, how many of these were treatment planning sessions, the treatment provided and materials used.

Costs were estimated from an NHS perspective for the care delivered in a tertiary care setting. Costs included all consumables and the total cost of NHS staff providing dental care (including overheads, training and administrative support). We did not include any costs borne by the patient and they do not make co-payments towards their care. We also did not include the costs of care provided by dental students in the base case analysis, although we did include costs of supervision of students, and an element of overheads associated with the provision of a teaching clinic. Costs of consumables and equipment were obtained from stock managers. Patient contact time with NHS employees providing dental care was estimated and combined with appropriate unit costs per hour taken from the Unit Costs of Health and Social Care, 2016⁹. This authoritative source provides unit costs per hour which include elements for on-costs, capital overheads, training, and where appropriate, administrative support.

The number of consultations and treatments sessions attended by the patients were recorded from our online appointments system and verified in the clinical notes. The clinics are staffed by a consultant supervising six chairs occupied by up to 7 postgraduate students during each three-hour clinical session, where an average of two patients per session are treated. We assumed that 14 individual treatment sessions for patients were delivered in each clinic.

Clinics were staffed with one NHS consultant dentist at a cost of £135 per hour, and three band 4 nurses, each costing £28 per working hour. An additional university-funded consultant doctor attends the clinic but is excluded in this analysis as their costs are covered by student fees. The resulting staff costs were £46.93 per patient treated. In addition, instrument

sterilisation was approximated at £4 per patient and overheads such as lighting/cleaning also was approximated at £4 per patient. The total amounted to £143.43 per patient per clinical visit.

Additional costs for direct restorations and root canal treatments were estimated by summing the individual cost per use of materials needed to complete the restoration (material capsules, bonding agents, polishing discs, disposable endodontic files etc.) and were provided by the stock managers. The estimates are reported in Table 1.

We also estimated the likely treatment cost to the patient if patients had received the same package of care at a private dental practice. Private fees were estimated using the consultation costs and costs for lab-work as is the typical model for private dental treatment. Consultation and lab-work costs were estimated by averaging practice fees from three mixed NHS/Private clinics based within the Greater London area and two central London private practices. These are provided as a range in Table 2.

We explored the extent of correlation between NHS and private cost estimates and patient characteristics. The characteristics we considered were age; BEWE score; patient concerns regarding progression, aesthetics and tooth shortening; the presence of tooth sensitivity; treatment of one or both anterior and posterior teeth; and diagnosis of more than one cause of tooth wear (from abrasion, attrition and erosion). Spearman's rank correlation was used to test for a correlation between two non-normally distributed continuous variables. The Wilcoxon-Mann-Whitney test was used to test for a correlation between a non-normally distributed continuous variable and a dichotomous variable.

We undertook sensitivity analysis on the estimates of cost in an NHS setting in which we assumed a cost of £59 per hour for dental registrars providing care in treatment sessions

rather than zero (base case). We based this sensitivity analysis on the cost per working hour for a registrar in the UK NHS ⁹.

Results

The clinical notes of 21 males and 9 females were evaluated. The average age at first examination was 45.5 years (SD 12.7, Range 24-76) and the average BEWE score was 14.2 (SD 3.2, Range 5-18). The aetiological factors were primarily erosion (n=13, 43.3%), erosion/attrition (n=12, 40%), attrition (n=3, 10%) and erosion/attrition/abrasion (n=2, 6.7%). The primary source of erosion was diet (n=13, 43.3%), intrinsic sources (n=5, 16.7%) or both (n=8, 26.7%). The most common presenting complaint from the patient was fear of progression or further shortening of teeth (n=23, 76.7%), aesthetics (n=19, 63.3%), dentine hypersensitivity (n=8, 26.7%) and reduced function (n=4, 13.3%).

The average treatment time for patients was 20.8 months (SD 9.6, Range 8-44 months). The average number of clinical visits during this time was 24.3 (SD 12.7; 8-48). Of these an average of 3.8 visits were treatment planning visits, often involving multiple disciplines.

Nine patients received direct composite restorations with no indirect restorations or prosthesis required. For these nine patients, the mean cost of providing direct anterior restorations only was £834 (SD £177; Range £675- £1,247). The mean costs of staff time (excluding postgraduate registrar time) and environment was £580 (SD £165; £439-£989) and materials and laboratory fees were £254 (SD £44; £199-£333). The remaining 21 patients required more extensive treatment. A full list of the most commonly prescribed treatment is presented in Table 1.

In addition, two patients had direct fibre core post restorations placed on 3 anterior teeth. Eight acrylic partial dentures and seven cobalt chrome partial dentures were provided to ten patients. Eight patients received conventional bridges with a total of 15 pontics. Two patients

received gold alloy palatal veneers (total of 11 teeth). All ceramic restorations were placed in two patients, one of whom received 12 ceramic crowns placed and one of whom received a ceramic veneer. One patient had a cast posterior post core made. Finally, 17 participants had either a Michigan (n=14) or a soft acrylic splint (n=3) provided.

The average cost per completed treatment plan for all patients in the base case was £2,371 (SD £1,290: £675-£4,807) for each of the 30 patients. This represented a mean of £1,333 (SD £697: £439-£2,637) for staff costs and £1,039 (SD £668: £199-£2,500) for materials and laboratory work. Sensitivity analysis in which dental registrars were assumed to cost £59 per hour generated mean staff costs of £3,481 (SD £1,820: £1,147-£6,885) and mean total costs of £4,519 (SD £2,382: £1,383-£8,254)

The estimated cost of similar treatment provided by a specialist in private clinic was £13,353 (SD £6,905; £4,737-£31,224) per patient and was 4.1 (SD 1.3) times higher than treatment provided within a hospital setting.

Costs estimated in both NHS and private settings did not follow a Normal distribution.

Likewise, patient age and BEWE were non-Normally distributed. Spearman's rank correlations between age and costs (either NHS or private setting) and between BEWE score and costs (NHS or private) were weak ($\rho < 0.3$ and $p > 0.10$ in all cases). Wilcoxon-Mann-Whitney tests of the correlation between costs and patient's reporting of concerns around progression; concerns around aesthetics; concerns around shortening of teeth; and sensitivity were also weak. Both NHS and private costs were correlated with the provision of treatment to both anterior and posterior teeth as opposed to treatment of anterior or posterior teeth only ($p = 0.006$ in both cases). There was a modest correlation between costs estimated in a private setting and multiple aetiological factors for tooth wear ($p = 0.07$).

Discussion

The breadth of treatments illustrated by this service evaluation demonstrate that the time commitment, cost and scope for erosive tooth wear restorative care is variable. There is no one prescriptive treatment, and planning is driven by patient needs, aiming to maintain functioning for as long as possible. The average patient spent 21 months in treatment requiring a visit at least once a month. Each treatment session was typically 1.5 hours in duration. This required significant patient compliance that was not without cost to the patient and includes taking time off work in addition to travel to and from appointments. Analysis of this was outside the scope of this paper but is worth taking into consideration when planning treatment. Treatment was invasive, often multi-disciplinary and requiring appointments with different practitioners.

The mean cost of a tooth wear rehabilitation case was £2,371 (SD £1290) in the hospital setting, with materials costing a mean of £1149 (SD £698). These financial estimates are probably conservative. The use of postgraduate training students in a tertiary setting enables provision of care at moderate cost to the NHS. Expanding provision from teaching hospitals would almost certainly entail higher costs. In our sensitivity analysis costs rose by an average of 1.9 times when the postgraduate registrars time was accounted for. There are also significant efficiencies of scale possible within a hospital environment which include discounts for materials. However, care may be prolonged in a teaching hospital because of the need for training. Furthermore, we could only estimate cost for treatment written in the clinical notes. Costs of some additional procedures (e.g. special trays) or multiple instruments and missed appointments were not captured. Lastly, it is a pre-requisite for patients who are referred for specialist prosthodontic rehabilitation to have good oral hygiene and stable periodontal health. This may not be reflective of those presenting with severe erosive tooth

wear to the general dental practitioner and stabilisation of periodontal health may further exacerbate treatment time and costs.

The treatment provided for each patient could range between a direct composite build-up of worn teeth to a full mouth rehabilitation involving multiple anterior and posterior crowns.

The interdisciplinary nature of the treatment provided, frequent need for endodontic treatment and extensive treatment planning necessary prior to direct restorative interventions makes high quality treatment of erosive tooth wear difficult under the current NHS reimbursement scheme.

Nine patients received treatment which might be classified as Band 2 NHS treatment using the current guidelines. A direct cost comparison to general dental practice using the English banding structure was too complex to calculate. However, the complexity of care, provision of multiple crowns in the same patient, endodontics and milled crowns around metal based partial dentures makes the financial commitment for general dental services very challenging. Conversely, directly placed plastic restorations remain a treatment option for some but are liable to repeated fractures and maintenance costs are high. In theory, these are possible to deliver in general practice. However, a patient may prefer crowns which become more challenging to provide.

The cost of erosive tooth wear rehabilitation in private dental practice is high and reflective of the extensive treatments required. Multifactorial tooth wear resulting from several aetiological factors were correlated with increased costs in NHS and private practice. This is a novel, but not unexpected finding, due to the complexity of treatment. Although the treatment plans will often be similar, the number of visits required in private practice may be reduced compared to a hospital setting as there is no training requirement. It is also recognised that private costs used and listed within this paper may not be indicative of costs

for every private practitioner in the UK or overseas. However, the aim of this paper was to provide an estimation and comparison, despite these limitations.

The cost of private treatment is likely to be unaffordable for many patients and is likely to contribute to a gap in service provision for those who do not have access to secondary/tertiary care. It is worth remembering that erosive tooth wear does not always require restorative intervention as erosive tooth wear is not always active and may reflect historical damage. If a convenient measure of progression of tooth wear were possible in general practice, patients may be more likely to accept non-invasive management options. The development of practice based quantitative monitoring tools is urgently required to objectively inform if restorative intervention is indicated. It is interesting to note that progression or further shortening of teeth was the main concern for 73% of the patients in this study. Other symptoms can be managed through non-invasive means using fluoride or similar interventions¹⁰. There is a lack of evidence on whether extensive prosthodontic treatment improves quality of life for those with severe erosive tooth wear. If the reason for the reduced quality of life is aesthetics and fear of progression, this may or may not be improved with advanced restorative work. Future work could consider investigating if quality of life is improved with prosthodontic rehabilitation, and the extent to which such improvements justify any additional treatment costs.

Conclusion

The prosthodontic rehabilitation of erosive tooth wear is complex, interdisciplinary and costly to the NHS. Patients should be fully informed and involved in the decision to treat and treatment planning stages. For extensive tooth wear cases, the cost to the NHS general dental practitioner makes offering high quality rehabilitation challenging. The option of private care is unaffordable for many. Costs are measured differently within a hospital setting, although

access is a limiting factor. Further work should investigate whether prosthodontic rehabilitation improves quality of life for those with erosive tooth wear.

Acknowledgments

The authors would like to thank Dr. Pranay Patel for his assistance in the assessment of private dental treatment fees.

References

1. Ahmed KE, Murray CA, Whitters CJ. A prospective survey of secondary care tooth wear referrals: demographics, reasons for concern and referral outcomes. *Br Dent J* 2014; 216: E9.
2. Papagianni CE, van der Meulen MJ, Naeije M, et al. Oral health-related quality of life in patients with tooth wear. *J Oral Rehabil* 2013; 40: 185–190.
3. Li MHM, Bernabé E. Tooth wear and quality of life among adults in the United Kingdom. *J Dent*; In press. Epub ahead of print 2016. DOI: 10.1016/j.jdent.2016.09.013.
4. Wetselaar P, Lobbezoo F. The tooth wear evaluation system: a modular clinical guideline for the diagnosis and management planning of worn dentitions. *J Oral Rehabil* 2016; 43: 69–80.
5. Bartlett DW, Ganss C, Lussi A. Basic Erosive Wear Examination (BEWE): a new scoring system for scientific and clinical needs. *Clin Oral Investig* 2008; 12: S65–S68.
6. Muts E-J, van Pelt H, Edelhoff D, et al. Tooth wear: A systematic review of treatment options. *J Prosthet Dent* 2014; 112: 752–759.
7. Bartlett D, Varma S. A retrospective audit of the outcome of composites used to

restore worn teeth. *Bdj* 2017; 223: 33–36.

8. Gulamali AB, Hemmings KW, Tredwin CJ, et al. Survival analysis of composite Dahl restorations provided to manage localised anterior tooth wear (ten year follow-up). *Br Dent J* 2011; 211: 1–8.
9. Curtis L, Burns A. Unit Costs of Health and Social Care 2016 | PSSRU <http://www.pssru.ac.uk/project-pages/unit-costs/unit-costs-2016/> (accessed 9 October 2017).
10. Lussi A, Carvalho T. The Future of Fluorides and Other Protective Agents in Erosion Prevention. *Caries Res* 2015; 49: 18–29.

Table 1. Lists of most common treatments provided

Treatments provided				
	Number of patients requiring treatment	Average number of units (SD)	Min	Max
Study models	30	3.4 (2.2)	2	12
Diagnostic wax ups	30	10.2 (5.0)	4	20
Crown lengthening surgery	12	2.8 (3.9)	1	11
Endodontics	14	2.4 (1.7)	1	7
Direct anterior restorations	25	6.3 (2.5)	2	12
Direct posterior restorations	16	3.4 (1.8)	1	6
Anterior Metal Ceramic Crowns	14	5.6 (2.5)	1	12
Posterior Gold Crowns	13	4.0 (3.0)	1	12
Anterior Metal Ceramic Crowns	4	4.0 (2.0)	1	8

Table 2: Cost of hospital materials and overheads

	Hospital Costs (£)
Cost per clinical session	54.93
Study models (n)	15
Teeth waxed up (n)	10
Crown lengthening surgery per tooth	40
Surgical Stent	50
Single root endo	70
Multi root endo	110
Acrylic denture	75.69
Cobalt Chrome denture	164
Bridge Abutments	60
Bridge Wings	20
Bridge Pontic	60
Anterior Glass Ionomer	9.20
Anterior Composite	18.20
Posterior Glass Ionomer	9.20
Posterior Composite	21.30
Posterior Amalgam	8.20
Anterior Post Core	31
Anterior Metal Ceramic Crown	61.50
Anterior Ceramic Crown	90
Gold alloy palatal veneer	51.25
Posterior Post Cores	30.75
Posterior Metal Ceramic Crown	61.5
Posterior Gold Crown	51.25
Michigan Splint	85
Soft Splint	20.25

Table 3: Estimated cost of treatment in private practice*

	Private Costs (£)	Range (£)
Treatment planning sessions	133.33	100-150
Study models (n)	76.67	30-100
Teeth waxed up (n)	40.00	20-50
Crown lengthening surgery per tooth	316.67	250-450
Surgical Stent	250.00	50-450
Single root endo	650.00	550-1000
Multi root endo	798.33	600-1000
Acrylic denture	816.67	550-1200
Cobalt Chrome denture	1883.33	1350-2500
Bridge Abutments	733.33	600-800
Bridge Wings	666.67	400-800
Bridge Pontic	733.33	600-800
Anterior Glass Ionomer	73.33	50-85
Anterior Composite	343.33	130-600
Posterior Glass Ionomer	80.00	70-85
Posterior Composite	353.33	160-600
Posterior Amalgam	101.67	70-150
Anterior Post Core	450.00	250-800
Anterior Metal Ceramic Crown	733.33	600-800
Anterior Ceramic Crown	783.33	750-800
Gold alloy palatal veneer	500.00	500
Posterior Post Cores	300.00	300
Posterior Metal Ceramic Crown	733.33	600-800
Posterior Gold Crown	816.67	800-850
Michigan Splint	533.33	500-550
Soft Splint	73.33	60-100

*Based upon the mean costs of 3 private practices in the Greater London Area. May not be indicative of costs in other practices/locations