



The twenty-first century Colorado River hot drought and implications for the future

Item Type	Article
Authors	Udall, Bradley; Overpeck, Jonathan T.
Citation	The twenty-first century Colorado River hot drought and implications for the future 2017, 53 (3):2404 Water Resources Research
DOI	10.1002/2016WR019638
Publisher	AMER GEOPHYSICAL UNION
Journal	Water Resources Research
Rights	© 2017. American Geophysical Union. All Rights Reserved.
Download date	26/08/2022 11:40:49
Item License	http://rightsstatements.org/vocab/InC/1.0/
Version	Final published version
Link to Item	http://hdl.handle.net/10150/624032



RESEARCH ARTICLE

10.1002/2016WR019638

The twenty-first century Colorado River hot drought and implications for the future

Bradley Udall^{1,2}  and Jonathan Overpeck^{2,3} 

¹Colorado Water Institute, Colorado State University, Fort Collins, Colorado, USA, ²Colorado River Research Group, Boulder, Colorado, USA, ³Department of Geosciences and Department of Hydrology and Atmospheric Sciences, Institute for the Environment, University of Arizona, Tucson, Arizona, USA

Key Points:

- Record Colorado River flow reductions averaged 19.3% per year during 2000–2014. One-third or more of the decline was likely due to warming
- Unabated greenhouse gas emissions will lead to continued substantial warming, translating to twenty-first century flow reductions of 35% or more
- More precipitation can reduce the flow loss, but lack of increase to date and large megadrought threat, reinforce risk of large flow loss

Supporting Information:

- Supporting Information S1

Correspondence to:

B. Udall,
Bradley.udall@Colostate.edu

Citation:

Udall, B. and J. Overpeck (2017), The twenty-first century Colorado River hot drought and implications for the future, *Water Resour. Res.*, 53, 2404–2418, doi:10.1002/2016WR019638.

Received 14 AUG 2016

Accepted 14 FEB 2017

Accepted article online 17 FEB 2017

Published online 24 MAR 2017

Abstract Between 2000 and 2014, annual Colorado River flows averaged 19% below the 1906–1999 average, the worst 15-year drought on record. At least one-sixth to one-half (average at one-third) of this loss is due to unprecedented temperatures (0.9°C above the 1906–1999 average), confirming model-based analysis that continued warming will likely further reduce flows. Whereas it is virtually certain that warming will continue with additional emissions of greenhouse gases to the atmosphere, there has been no observed trend toward greater precipitation in the Colorado Basin, nor are climate models in agreement that there should be a trend. Moreover, there is a significant risk of decadal and multidecadal drought in the coming century, indicating that any increase in mean precipitation will likely be offset during periods of prolonged drought. Recently published estimates of Colorado River flow sensitivity to temperature combined with a large number of recent climate model-based temperature projections indicate that continued business-as-usual warming will drive temperature-induced declines in river flow, conservatively –20% by midcentury and –35% by end-century, with support for losses exceeding –30% at midcentury and –55% at end-century. Precipitation increases may moderate these declines somewhat, but to date no such increases are evident and there is no model agreement on future precipitation changes. These results, combined with the increasing likelihood of prolonged drought in the river basin, suggest that future climate change impacts on the Colorado River flows will be much more serious than currently assumed, especially if substantial reductions in greenhouse gas emissions do not occur.

Plain Language Summary Between 2000 and 2014, annual Colorado River flows averaged 19% below the 1906–1999 average, the worst 15-year drought on record. Approximately one-third of the flow loss is due to high temperatures now common in the basin, a result of human caused climate change. Previous comparable droughts were caused by a lack of precipitation, not high temperatures. As temperatures increase in the 21st century due to continued human emissions of greenhouse gasses, additional temperature-induced flow losses will occur. These losses may exceed 20% at mid-century and 35% at end-century. Additional precipitation may reduce these temperature-induced losses somewhat, but to date no precipitation increases have been noted and climate models do not agree that such increases will occur. These results suggest that future climate change impacts on the Colorado River will be greater than currently assumed. Reductions in greenhouse gas emissions will lead to lower future temperatures and hence less flow loss.

1. Introduction

A large number of studies over the last 25 years have considered the future runoff of the Colorado River (Figure 1) under climate change. Nearly all of these studies have cautioned that future warming will deplete the flow of the river, but the results have varied from minor to major [Nash and Gleick, 1991; Christensen et al., 2004; Milly et al., 2005; Brekke et al., 2007; Christensen and Lettenmaier, 2007; National Research Council, 2007; Seager et al., 2007; Barnett and Pierce, 2008; Ray et al., 2008; Barnett and Pierce, 2009; Rajagopalan et al., 2009; Cayan et al., 2010; Reclamation, 2013; Harding et al., 2012; Seager et al., 2012; Vano et al., 2012; Ficklin et al., 2013; Vano et al., 2014; Ayers et al., 2016; Milly and Dunne, 2016]. In contrast, the latest U.S. Government assessment implies little or no change is likely because precipitation increases will be sufficient to maintain temperature-depleted flows [Reclamation, 2016]. Fifteen years into the twenty-first century, the emerging reality is that climate change is already depleting




Figure 1. Map of the Colorado River Basin. Lower and Upper Basins, major U.S. cities receiving Colorado River water, major tributaries, and Lakes Mead and Powell are shown. The Central Arizona Project canal in red.

Colorado River water supplies at the upper end of the range suggested by previously published projections. Record setting temperatures are an important and underappreciated component of the flow reductions now being observed.

Between the start of the drought in 2000 and the end of 2014, our analysis period, annual flow reductions averaged 19.3% below the 1906–1999 normal period, and Lakes Mead and Powell, the nation’s two largest reservoirs, ended the period at approximately 40% of maximum volume despite starting the period nearly full [Wines, 2014; *Colorado River Basin Stakeholders*, 2015] (Figure 2a). This drought has continued into 2015 and 2016 with higher, but still below normal, flows estimated at 94% in 2015 and 94% in 2016 with unusual late season May and June precipitation in both years that raised runoff by nearly 20% [Alcorn, 2015, 2016]. Despite these smaller recent reductions, Lake Mead continues to decline and in May 2016 it hit a level not seen since its initial filling in the 1930s [James, 2016]. The overall Colorado River reservoir system stores 4 times the annual flow of the river, one of the largest ratios in the world. This storage provides a large drought buffer when full. However, when the reservoirs are low, shortage risk can be high for years because high demands, now equal to twentieth century average flow, make it difficult to refill system storage [Reclamation, 2012]. While the multiyear California drought has been garnering more national attention, the more slowly unfolding Colorado River drought is every bit as serious and also has national and international ramifications [Wines, 2014].

The Colorado River Basin encompasses seven states and northern Mexico and is home to 22 federally recognized tribes. The river provides municipal and industrial water for 40 m people distributed across every major Southwestern city both within and without the basin, including Los Angeles, San Diego, Las Vegas, Phoenix, Tucson, Salt Lake City, Denver and the entire Front Range of Colorado, Albuquerque, and Santa Fe [Reclamation, 2012].

Continued low flows would result in additional declines at Lake Mead, eventually requiring Lower Basin (Arizona, California, Nevada) water delivery shortages with mandatory cutbacks imposed primarily on Arizona, but also Nevada and Mexico [Verburg, 2011]. At the same time, Upper Basin (Colorado, New Mexico, Utah, Wyoming) water users would continue to endure physical shortages from a lack of water. These initial Lower Basin Lake Mead delivery shortages and Upper Basin physical shortages are manageable to a point; however, under current operating rules with continued low flows during the next 6 to 8 years Lake Mead would drop to elevation 305 m (1000 feet) above sea level, resulting in a number of serious and unprecedented problems [Collum and McCann, 2014].

In the Lower Basin, Arizona could theoretically lose its water allocation for the entire Central Arizona Project canal, a critical \$4.4B, 530 km cross-state 2 bcm/yr water source for 4.7 m people, multiple sovereign Indian




Figure 2. (a) Lakes Mead and Powell combined monthly contents. Upper Basin annual Colorado River (b) runoff at Lees Ferry from 1906 to 2014, (c) precipitation and (d) temperatures from 1896 to 2014. Mead first filled in 1935, Powell in 1963 (supporting information Text S1). Two 15-year drought periods, 1953–1967 and 2000–2014, are highlighted and discussed in main text.

nations, and over 120,000 irrigated hectares [Glennon, 1995; Colorado River Basin Stakeholders, 2015]. This canal currently relies on occasional but uncertain “equalization” releases from Lake Powell that only occur with irregular and rare large Powell inflows. The extra water is delivered when Lake Powell reaches levels substantially higher than Lake Mead, a use allowed under the 1922 Colorado River Compact section III (e) and formalized most recently under rules established in a 2007 Record of Decision for coordinated operations of Lakes Powell and Mead and for shortage sharing in the Lower Basin [Department of Interior, 2007].

Under normal operating rules, without these extra inflows, Lake Mead has excess outflows of 1.5 bcm per year, the so-called Lower Basin “structural deficit” [Collum and McCann, 2014]. The structural deficit was created in 1968 when Congress authorized the Central Arizona Project (CAP). In order to obtain the support of

the large California Congressional delegation, Arizona agreed to rely on this unused, but in the long run unreliable water, because there was not enough remaining unallocated Lower Basin water. The CAP had long been a desire of Arizona and the state was willing to make this bargain despite its flaws [Johnson, 1977]. This same water is first available for use by the Upper Basin under the Colorado River Compact, but heretofore has not been developed for Upper Basin use. A plan to augment the Colorado River with flows from outside the basin, discussed during the hearings on the legislation, but not included in the final package due to opposition from potential source areas, was never revisited by Congress. Reclamation in 2011 said that such augmentation was now unlikely.

The structural deficit only became a problem when the CAP was fully completed in the mid-1990s combined with the drought that began in 2000. Upper Basin demand growth has also played a small role, although Upper Basin demands are still much less than forecast in 1968 for the year 2000 [Tipton and Kalmbach, Inc., 1965; Johnson, 1977]. The recent Lake Mead declines are strongly influenced by this imbalance, and solutions to this deficit have been a recent focus of the Basin states and federal government [Central Arizona Project, 2016; Davis, 2016].

The Upper Basin also has serious issues, one of which ripples into the Lower Basin. When the surface of Lake Mead declines to an elevation 305 m (1000 feet) above sea level, Lake Powell will also be below its minimum power pool 75% of the time [Collum and McCann, 2014]. This occurs in part because low Mead levels make “equalization” releases from Powell more likely thus driving Powell lower. Hydropower losses at Lake Powell could result in substantial rate increases for irrigators who rely on the reservoirs for long term lower cost power contracts, and would also dry up funding for basin-wide programs necessary for water delivery environmental compliance [Adler, 2007; Collum and McCann, 2014]. Under such low reservoir conditions, there is also a high likelihood that the Upper Basin states would have to curtail existing water deliveries to cities such as Denver, Colorado Springs, Albuquerque and Salt Lake City in order to make required deliveries to Lake Mead. Heretofore, largely because of the structure of the Colorado River Compact, the Upper Basin and Lower Basin have been managed separately. With permanent flow declines of approximately 20%, however, the required deliveries to Lake Mead would become a hardship on the Upper Basin, as well as create Lower Basin delivery shortages [Reclamation, 2007; Barnett and Pierce, 2009; Rajagopalan et al., 2009]. The original compact, signed during one of the wettest periods in the last 450 years [Woodhouse et al., 2006], did not envision how large scale flow declines would be managed between the basins, and such declines could cause an allocation crisis between the Upper and Lower Basins [Adler, 2008].

Understanding the cause of, and reacting properly to, the ongoing drought is critical to the future of the Southwest. Herein we investigate the role of precipitation versus temperatures as causes of the current drought, provide temperature-based and precipitation-based twenty-first century flow projections and provide policy implications of these findings. Our approach separates the impacts of high-confidence temperature projections from those associated with the much lower-confidence projections of future precipitation using a simple but powerful sensitivity technique. Moreover, we make a novel—and important—case that there is a high likelihood that the impacts of continued atmospheric warming will overwhelm any future increases in precipitation because prolonged dry periods lasting multiple decades are likely to negate the beneficial impacts of additional precipitation during other times.

2. Causes of the 2000–2014 Drought

The 2000–2014 drought is defined by the lowest average annual flows for any 15-year period in the historical record. To analyze this drought, gridded 4×4 km temperature and precipitation data from 1896–2014 for the area above Lees Ferry were obtained from the Precipitation-Elevation Regression on Independent Slopes (PRISM) model [Daly et al., 1994; Guentchev et al., 2010; Oyster et al., 2015a, 2015b; Rangwala et al., 2015]. In addition, we obtained reservoir contents and natural flows at Lees Ferry from the U.S. Bureau of Reclamation (Reclamation) (Text S1). Lees Ferry is situated just below Lake Powell and is the Compact dividing line between the Upper and Lower Basins. Approximately 85% of the flow originates above Lees Ferry [Christensen and Lettenmaier, 2007].

Historically, Upper Colorado River Basin precipitation has been the main Colorado River runoff driver such that high flow years (1920s, 1980s) were associated with high precipitation and low flow years (1930s, 1950s) with low precipitation (Figures 2b and 2c). The current drought (our study period is 2000–2014, but

Table 1. Winter/Summer/Annual Upper Basin Mean Water Year Precipitation

	1953–1967			2000–2014			1896–2014	
	mm			mm			mm	
	Total	Anomaly	Anomaly % of Mean (%)	Total	Anomaly	Anomaly % of Mean (%)	Mm	% Avg
Winter (Oct to Mar)	176	–16	–8.6	187	–5	–2.7	192	100
Summer (Apr to Sep)	184	–7	–3.6	179	–12	–6.4	191	100
Total	359	–23	–6.1	365	–17	–4.6	383	100

the drought is still on-going), with its modest –4.6% precipitation decline and –19.3% flow decline, stands in stark contrast to the second-lowest 15-year flow period (1953–1967), a precipitation-driven drought with averaged precipitation reductions of –6.1% per year and flow reductions of –18.1% per year (Figures 2b and 2c and Table 1). Compared to the 1950s drought, the 2000s feature much more (near normal) winter precipitation (–8.6% 1950s decline versus –2.7% 2000s) and significantly less summer precipitation (–3.6% 1950s decline versus –6.4% 2000s). The 2000s precipitation decline is only 75% of the decline in the 1950s, thus begging the question of why the recent drought was more serious. What has changed is that temperatures in the runoff producing Upper Basin are now 0.9°C above the 1896–1999 average and are the highest in the gaged record; whereas temperatures during the 1953–1967 drought were much cooler and only slightly above the 1896–1999 average (Figure 2d and Table 2). This makes the current drought unprecedented in the gaged record.

In contrast to the more precipitation-driven current California drought [Differbaugh et al., 2015; Williams et al., 2015], lack of precipitation is only partially to blame for the Colorado River runoff declines during the last 15 years. Instead, approximately a third, or more, of the recent Colorado River flow reduction is most likely a result of record-setting warmth. Since 1988 an increase in the frequency of warm years has been strongly associated with lower flows than expected [Woodhouse et al., 2016], suggesting an important role for temperature in flow losses. Such temperature-driven droughts have been termed “global-change type droughts” and “hot drought,” with higher temperatures turning what would have been modest droughts into severe ones, and also increasing the odds of drought in any given year or period of years [Breshears et al., 2005; Overpeck, 2013]. Higher temperatures increase atmospheric moisture demand, evaporation from water bodies and soil, sublimation from snow, evapotranspiration (ET) from plants, and also increase the length of the growing season during which ET occurs [Pitman, 2003; Weiss et al., 2009; Seneviratne et al., 2010; Seager et al., 2015a]. Warm season (April to September) warming has been identified by models as especially important in reducing Colorado River flows because of the increases in ET from longer growing seasons [Das et al., 2011]. Increases in measured vapor pressure deficits in the Southwest caused by warming and a decrease in water vapor provide strong support for higher ET during the recent drought [Seager et al., 2015b]. As increasing temperatures drive further drying, additional positive feedbacks are possible in the form of lower humidity and less evaporative cooling, decreased cloudiness and increased incident radiation, as well as decreased snow cover and more radiative heating [Betts et al., 1996; Brubaker and Entekhabi, 1996; Pitman, 2003; Seneviratne et al., 2010]. In the twentieth century, droughts were associated almost exclusively with a lack of precipitation. In this century, however, high temperatures alone can lead to anomalously dry conditions.

Table 2. Upper Basin Water Year Flows and Temperatures

Period	Average Annual Flow		Average Annual Temperature	
	bcm	% 1906–1999	°C	°C Anomaly to 1896–1999
1953–1967	15.38	81.9	7.0	0.2
2000–2014	15.15	80.7	7.7	0.9
1906–1999	18.77	100.0	6.8	0.0
1906–2014	18.27	97.3	6.9	0.1

3. Estimates of 2000–2014 Temperature-Induced Flow Loss

Over the last several years several studies specific to the Colorado River Basin have investigated the specific relationships among temperatures, precipitation and flow in the basin using the concepts of temperature

sensitivity and precipitation elasticity [McCabe and Wolock, 2007; Nowak et al., 2012; Vano et al., 2012, 2014; Vano and Lettenmaier, 2014]. Temperature sensitivity is defined as the percent change in annual flow per degree rise in annual temperature. Precipitation elasticity is defined as the fractional change in annual flow divided by the fractional change in annual precipitation [Vano et al., 2012]. Note that elasticity has been studied for both increases and decreases in precipitation, whereas sensitivity is typically investigated only for temperature increases. These numbers can be determined empirically and through model studies.

Previous studies on temperature sensitivity and precipitation elasticity show that future impacts to streamflow from increases in temperatures and changes in precipitation can be considered separately using sensitivity and elasticity, and then added together to produce flow estimates [Vano et al., 2014; Vano and Lettenmaier, 2014]. Considering these effects separately and additively is a powerful conceptual tool for investigating climate change impacts because of the ease in measuring the two variables for current impacts and the wide availability of temperature and precipitation projections from global climate models for assessing future impacts. In addition, the large differences in certainty associated with future changes in the two variables (temperature will surely increase, whereas precipitation may increase or decrease—see below) helps to set apart the risk of future changes in flow associated with each variable.

Vano et al. [2012, 2014], McCabe and Wolock [2007], and Nowak et al. [2012] provide multiple estimates of the flow sensitivity of the Colorado River flow to temperature using three different methods. Vano et al. [2012, 2014] utilized six high-resolution, commonly used hydrology models and two different temperature adjustment methods to obtain Lees Ferry temperature sensitivities. They report an average sensitivity of $-6.5\%/^{\circ}\text{C}$ warming with a one standard deviation range from -3.0% to $-10.0\%/^{\circ}\text{C}$ for the Upper Basin. Approximately 50% models show increasing sensitivity and 50% decreasing sensitivity as temperatures warm so we elect to use a constant sensitivity over all future temperatures. McCabe and Wolock [2007] constructed a simple water balance model that infers an average temperature sensitivity of $-8.9\%/^{\circ}\text{C}$ and Nowak et al. [2012] found an empirical temperature sensitivity of $-13.8\%/^{\circ}\text{C}$.

We use the complete one standard deviation range ($-3\%/^{\circ}\text{C}$ to $-10\%/^{\circ}\text{C}$) of the Vano et al. [2012, 2014] temperature sensitivity estimates as they were the most conservative and rigorous of the three studies we investigated. Using this range, we found that recent warming of 0.9°C has likely already reduced river flows from -2.7% to -9% from the mean 1906–1999 flow. This represents approximately one-sixth to one-half (average of one-third) of the total flow loss during the 2000–2014 drought.

The higher temperature sensitivities of the two other studies suggest the actual Colorado River temperature sensitivities are near the upper end and possibly exceed the Vano et al. [2012, 2014] estimates. These higher sensitivities imply much greater temperature-induced losses during the current drought (-7.9% to -12.3% versus -2.7% to -9%). Empirical results from the 2000 to 2014 drought also point to mid to high temperature sensitivities. Vano et al. [2012] report precipitation elasticities ranging from 2 to 3 at Lees Ferry. Thus, using a midrange precipitation elasticity of 2.5, the 2000–2014 annual -4.6% precipitation decline implies runoff reductions of -11.4% , leaving the remaining -7.9% decline to be explained by other causes. If temperature were the sole cause of this remaining decline, the inferred temperature sensitivity is $-8.8\%/^{\circ}\text{C}$. Using a precipitation elasticity of 3.0 implies a temperature sensitivity of $-6.2\%/^{\circ}\text{C}$, very close to the midrange Vano et al., sensitivity. These temperature sensitivities imply large losses as temperatures rise, the subject of the next section.

4. Twenty-First Century Flow Response to Changing Temperatures and Precipitation

For the analysis on how future temperatures and precipitation would affect runoff, and for investigating how well current linked climate-hydrology models can reproduce the current drought, we used Reclamation's climate projection data sets [Brekke et al., 2013, 2014]. These data sets use Coupled Model Intercomparison Project 3 and 5 (CMIP3, CMIP5 after the class of climate models used) climate model projection data linked to the Variable Infiltration Capacity hydrology model to produce flows from 1950 to 2099 (supporting information Text S2, Figures S2, and S3) [Liang et al., 1996; Meehl et al., 2007; Moss et al., 2010; Taylor et al., 2012].

The same temperature sensitivity and precipitation elasticity numbers discussed above can be used to estimate future flow reductions using climate model outputs under high (business-as-usual, SRES A2 and

RCP8.5) and moderate (somewhat reduced by mitigation, SRES A1B and RCP4.5) greenhouse gas emissions to the atmosphere. By 2050, moderate and high emissions are projected to yield Upper Basin *mean* warming of 2.6–2.8°C (Figure 3), three times recent warming, and by 2100, warming of 3.6°C under moderate emissions and 5.4°C under high emissions. This warming implies total multimodel mean temperature-induced flow losses at midrange sensitivity of $-6.5\%/^{\circ}\text{C}$ of about -17% by midcentury and -25% to -35% at end-century (Figures 4 and 5). The multimodel mean complete flow loss *range* over both periods and both emissions is approximately -8% to -55% using the lower and upper temperature sensitivities (Figures 4 and 5). As discussed above, there is little empirical evidence that the true temperature sensitivity of flow to temperature increase is near the low sensitivity.

Temperature-induced losses may be somewhat buffered by projected additional precipitation that can increase runoff by 2–3% for every 1% change in precipitation [Vano *et al.*, 2012]. At midcentury precipitation increases of $+4\text{--}+11\%$ given a midrange elasticity of 2.5 would balance the range of temperature-induced flow losses at a midrange $-6.5\%/^{\circ}\text{C}$ sensitivity (Figure 5, right y axis). At end-century, with the same sensitivity and elasticity, additional precipitation increases of $+4\text{--}+20\%$ would balance the range of possible temperature-driven losses. At a higher $-10\%/^{\circ}\text{C}$ sensitivity, the balancing precipitation would need to be as great as $+15\%$ or more at midcentury and $+22\%$ or more at end-century. While these may seem like relatively small increases in precipitation, and thus possible, they would represent a major and unprecedented change in precipitation regime compared to the observed historical variation in precipitation (Figure 2c). During the twentieth century, for example, the wettest 10-year period (1983–1997) had only a $+8\%$ precipitation increase. This unusual period was marked by major floods downstream of Lakes Powell and Mead due to uncontrolled reservoir spilling and the near catastrophic loss of the spillways at Glen Canyon Dam [Udall, 1983].

Vano and Lettenmaier [2014] argue that the sensitivity-based approach used in our projections provides similar estimates of future streamflow to those generated with more computationally intensive coupled-model methods, except for some (i.e., 10%) overstatement of flow reductions at the highest levels of possible warming by 2100 (e.g., the business-as-usual SRES A2 scenario used in the CMIP3 projections and the RCP8.5 in the CMIP5 projections). This would reduce the end of century high emissions mean flow reductions shown in Figure 5 to a still very significant -45% by 2100.

Recent studies have suggested that CO₂ fertilization may increase plant water efficiency thus reducing future evapotranspiration which could serve to mitigate our projected losses [Milly and Dunne, 2016; Swann *et al.*, 2016]. Both studies call into question results that show large portions of the globe drying in the twenty-first century [e.g., Dai, 2012; Cook *et al.*, 2014]. However, Milly and Dunne [2016] and Swann *et al.* [2016] show that, despite this increase in plant water use efficiency, the Southwestern US will still dry, a finding that is consistent with multiple global assessments showing substantial drying risk to midlatitude areas such as the Colorado River Basin. Moreover, a recent Australian study found that higher




Figure 3. Probability density functions of Upper Colorado River Basin temperature projections for midcentury and end-century under moderate (SRES A1B and RCP4.5) and high (SRES A2 and RCP8.5) emissions.




Figure 4. Probability density functions of Upper Colorado River Basin temperature-induced flow reductions for midcentury and end-century with the three temperature sensitivities (−3%, −6.5%, −10%) and the two levels of emissions (Moderate: SRES A1B and RCP4.5 and High: SRES A2 and RCP8.5).

evapotranspiration associated with the increased plant growth stimulated by higher CO₂ outweighed any CO₂-related water-use efficiency effect, and served to reduce streamflows in semiarid regions [Ukkola *et al.*, 2015], a trend that must be exacerbated by the temperature-induced lengthening of the growing season. These results suggest that plant physiological responses are likely consistent with our results, and in any case, do not invalidate them.

5. Megadrought Risks to Flows

Megadroughts lasting decades in the Colorado River Basin have occurred in the past, with resulting substantial flow reductions [Meko *et al.*, 2007]. Multiple papers now suggest there is high twenty-first century risk for megadrought in the American Southwest and that the risk will increase as temperatures rise [Ault *et al.*, 2014; Cook *et al.*, 2015; Ault *et al.*, 2016]. In addition, current GCMs underrepresent the frequency of megadrought [Ault *et al.*, 2012, 2013]. These findings provide additional support for large flow reductions during at least multidecadal drought periods and suggest that current twenty-first century flow projections underrepresent this risk.

Significant Colorado River flow losses occurred during previous multidecadal megadroughts. During the twelfth century, flow reductions of approximately −16% occurred during one 25-year period [Meko *et al.*,




Figure 5. Temperature-induced flow losses by model run (one per dot) with temperature increases shown on horizontal axis. For each period (midcentury, end-century) and emissions type (moderate, high), flow losses for each model run are shown with the 3 (low = $-3\%/^{\circ}\text{C}$, medium = $-6.5\%/^{\circ}\text{C}$, high = $-10\%/^{\circ}\text{C}$) temperature sensitivities. Black dots/circles are averages/medians for each sensitivity. Precipitation increases needed to counteract flow losses at right are based on 2.5 precipitation elasticity. Range for the temperature-induced losses during 2000–2014 drought are shown in shaded brown at the top (supporting information Text S5).

2007]. Evidence indicates that hemispheric and Southwest temperature anomalies were significantly smaller during past megadroughts than the rapid on-going current warming that could easily exceed 4–5°C by the end of century under business-as-usual emissions [Salzer and Kipfmüller, 2005; Mann et al., 2009; Salzer et al., 2014] (Figure 5). Using the additivity concepts discussed above, additional warming of 1°C, 2°C, or 3°C beyond the historic twelfth century megadrought temperatures would have reduced the -16% flow declines by an additional -6.5% , -13% , or -19.5% at medium temperature sensitivity. These additional reductions would have thus turned a -16% flow decline into declines of -21.5% , -28% , or -34.5% , losses near the middle of our projections.

There is recent strong evidence that continued warming over the next 80 years could increase the risk of multidecadal drought [Ault et al., 2014, 2016; Cook et al., 2015]. Independent of the added drought risk due

to continued warming, the risk of a 35-year precipitation-deficit drought later in this century exceeds 15% within a 50-year period [Ault *et al.*, 2014]. In contrast, with continued anthropogenic warming, the risk of multidecadal megadrought in the Southwest increases to over 90% over this century if there is no increase in mean precipitation; even if modest precipitation increases do occur, the risk will still exceed 70% [Ault *et al.*, 2014, 2016]. At medium warming (4°C), 20–30% precipitation increases will be needed to reduce megadrought risk below 50% and at high amounts of warming (>6°C), it will take a ~40% increase in precipitation to reduce megadrought risk below 50% [Ault *et al.*, 2016]. These changes in precipitation are huge and unlikely, and they would still only reduce megadrought risk to below 50%.

Both the CMIP3 and CMIP5 Global Climate Models may not adequately reproduce the frequency of occurrence of known past decadal and multidecadal precipitation droughts [Ault *et al.*, 2012, 2013]. In the Colorado River Basin empirical evidence of this problem can be found in the linked GCM-hydrology model results from Reclamation's projections for the basin [Brekke *et al.*, 2014]. Approximately half of the CMIP5 models and one-quarter of the CMIP3 models cannot simulate the 2000–2014 drought at any point in the twenty-first century (supporting information Text S3 and Tables S1–S4). This wet bias significantly affects the mean flows of drought-capable and nondrought capable models. At the end of the twenty-first century, the models unable to simulate the current drought are much wetter (109% of twentieth century average Lees Ferry runoff for CMIP3, 113% for CMIP5) than the models that are able to simulate the current drought (85% of average runoff for CMIP3, 91% CMIP5) (supporting information Tables S1–S4). These flow differences are greater than 20%, and represent the difference between serious management challenges and significant oversupply.

6. Risk-Based Framing of Future Runoff Projections

At present, some outputs from global climate models are ready to support reliable risk-based policy while others are not as ready. A key novel aspect of our research is to provide more insight into where confidence is warranted, and where it is not, with respect to projections of future climate and flow change in the Colorado River Basin. In the case of the Basin, every single moderate and high emissions model simulation agrees that temperatures will continue to rise significantly with continued emissions of greenhouse gases to the atmosphere—this result is robust, highly certain and well-suited for informing policy choices. The fact that observations also show substantial warming only strengthens this assertion.

On the other hand, simulated future precipitation change in the Basin is clouded with much greater uncertainty due to substantial disagreement among models and a highly uncertain ability to simulate realistic change in key phenomena such as storm-track position or decadal and longer-scale drought. Whereas climate models are in general agreement that cool season (warm season much less certain) precipitation declines are likely in the Lower Colorado River Basin, these same models disagree when it comes to the sign and amount of precipitation change that is likely in the Upper Basin. This is because precipitation change in the Upper Basin will depend heavily on the exact changes in the position of cool season jet stream and storm-tracks, two aspects of climate change that are not simulated with confidence by global climate models [Collins *et al.*, 2013].

Moreover, there is strong evidence that the mean positions of both the jet stream and storm-tracks are likely to push poleward, expanding the area of aridity in the Colorado River Basin, but the amount of this expansion is poorly constrained [Collins *et al.*, 2013]. Multiple studies, including some focused on the American Southwest, suggest that the proximate cause of this drying, Hadley Cell expansion, is already well underway and will continue [Seager *et al.*, 2007; Scheff and Frierson, 2012; Feng and Fu, 2013; Norris *et al.*, 2016; Prein *et al.*, 2016].

Our results regarding future changes in Colorado River flows agree with many previous studies in suggesting climate change translates to flow reductions, although our work is generally not directly comparable because we separate out high confidence temperature-related impacts from the possible effects of much less certain and highly variable precipitation projections. However, our work, as well as this larger body of literature, appears to be at odds with the recent Reclamation projections for the Colorado River Basin, which are widely cited and used. Reclamation's projections use a global climate model output that is downscaled to drive a hydrology model. It is worth understanding why our results emphasize substantially greater risks along with apparently greater flow losses.

The 2011 CMIP3 climate change flow projections by Reclamation indicate a modest multimodel median flow decline of -9% by 2060 for the river, but with a wide range of outcomes from flow increases to flow decreases [Reclamation, 2012] (supporting information Table S1). Reclamation's most recent CMIP5 projections show no change in mean and median basin-wide flow by 2070s [Reclamation, 2016], but also embody a wide range of results. Compared to CMIP3, the CMIP5 results show increased precipitation, especially in the northern parts of the basin including Northeast Utah, Northwest Colorado's Yampa River and the Green River in Wyoming [Brekke *et al.*, 2014; Ayers *et al.*, 2016] (supporting information Tables S1 and S3). The increased precipitation in the CMIP5 model runs compared to CMIP3 can be attributed to more southerly storm tracks in CMIP5 that occur in late spring [Brekke *et al.*, 2014].

Another issue arises in both the CMIP3 and CMIP5 data sets when GCM precipitation is adjusted by the downscaling techniques necessary for off-line hydrology models. The first step in Reclamation's downscaling is a bias correction step. This step can add approximately 5% more precipitation to the raw GCM precipitation, and this increase appears to not have a physical basis [Reclamation, 2013; Brekke *et al.*, 2013]. The final downscaling step, spatial downscaling, also increases GCM precipitation, although there is at least a plausible physical explanation for some of the increase: higher elevations in the Rockies receive large amounts of precipitation, but these elevations are not properly modeled by the GCMs. In one study of the CMIP5 data set after downscaling, dry and average models show precipitation increases of approximately $+5\%$ from the raw GCM output, but the wettest models show $+10\%$ increases, doubling future precipitation increases from $+10\%$ to $+20\%$ [Lukas *et al.*, 2014]. This extra precipitation is manifested in a number of hydrology model runs that project huge and implausible flow increases in some years that are 150% of the highest known flows in the twentieth century (supporting information Text S4, Figures S2, and S3). The downscaling wetness problem has been identified, but has not been resolved [Lukas *et al.*, 2014]. Reclamation acknowledges that the newer CMIP5 projections have not been determined to be better or more reliable [Brekke *et al.*, 2014]. It is noteworthy that internally consistent GCM-only Southwest runoff projections almost uniformly produce significant declines in both CMIP3 and CMIP5 runs [Milly *et al.*, 2005; Seager *et al.*, 2007, 2012; Koirala *et al.*, 2014; Milly and Dunne, 2016].

Our results are generally comparable to Reclamation's most recent results when considering the full range of our analysis when both precipitation and temperatures are included. However, our focus and emphasis is on the large near-certain temperature-induced flow declines with a separate analysis of precipitation. Reclamation, by contrast, has focused on climate multimodel-ensemble median declines, including medians calculated across emission scenarios [Reclamation, 2013, 2012]. Decision makers often treat these median outcomes as a proxy for risk despite the fact that the median obscures the wide range of results and lumps wet and dry, warm and hot, large and small emission increases and, most critically, near certain temperature increases and very uncertain precipitation changes.

We assert that the large precipitation increases necessary to offset substantial temperature-induced flow decreases appear unlikely to occur for a number of reasons. These reasons include the potential for storm tracks to go north of the basin due to Hadley Cell expansion, the high potential for megadrought to increase evaporation while reducing precipitation and runoff for extended periods, the large size of the needed precipitation increases, especially when compared to decadal historical increases, the consistent identification by global assessments of the Southwest as an area likely to dry, and finally the lack of any trend over the last century or last 16 years (Figure 2c). Hence, we choose to focus on highly likely temperature-induced declines with separate analysis of the precipitation needed to offset these declines.

7. Policy Implications and Solutions

The climate science take-home messages for Colorado River managers are thus: (1) there is little doubt (i.e., high confidence) that temperatures will continue to increase as long as the emissions of greenhouse gases to the atmosphere continue; (2) there is also high confidence that continued temperature increases will cause river flows to decline, ranging from -11% to as much as -55% by end of century under moderate to high emissions (Figures 4 and 5); (3) there is only low confidence associated with the possibility of storms and precipitation in the Upper Basin increasing enough to even partially offset the temperature-driven declines in river flows; (4) the risk of multidecadal megadrought in the Basin is significant even in the absence of continued anthropogenic climate change, and this risk rises substantially with continued global

warming; (5) the likelihood of drought and megadrought means that there will likely be decades-long periods with anomalously low runoff even if there is an increase in precipitation relative to the historical mean during some other periods due to anthropogenic climate change.

Temperature-driven threats to the flows of the Colorado are thus large and real. The only way to curb substantial risk of long term mean declines in Colorado River flow is thus to work toward aggressive reductions in the emissions of greenhouse gases into the atmosphere. Our work shows that modest (e.g., RCP4.5) reductions in greenhouse gas emissions, while having better outcomes than the business-as-usual future (e.g., RCP8.5), still imply large Colorado River flow losses.

The record warm nature of the on-going Colorado River drought indicates that this drought is not just a natural drought, and our work demonstrates that flows are unlikely to return to the twentieth century averages if we only wait. Unusually wet periods like the 1920s and 1990s will still continue to occur, but they will co-occur with higher temperatures that will increase water demand from plants, soil, snow, and humans.

Climate models and theory suggest that flow reductions would be more severe in the Southern portions of the Upper Colorado Basin affecting tributaries such as the San Juan, Dolores, and Gunnison more severely, with smaller impacts to more northerly tributaries such as the Yampa and Green [Ayers *et al.*, 2016]. Such spatial distribution would provide additional water management challenges in that the more southerly basins have in general more people, infrastructure, and uses. Such a distribution would create new localized water supply shortages in addition to the overall basin-wide issues.

Other known threats to streamflows include the potential large scale loss of conifers [Breshears *et al.*, 2005; Adams *et al.*, 2009; Allen *et al.*, 2010, 2015], and the impacts of dust on snow [Painter *et al.*, 2010; Deems *et al.*, 2013]. These factors along with the observed and projected temperature-induced Colorado River flow declines, the inability of many linked climate-hydrology models to simulate persistent droughts, and the increasing likelihood of hot drought and megadrought, all imply that future Colorado River water supply risk is high. It is imperative that decision-makers begin to consider seriously the policy implications of potential large-scale future flow declines. Stable twentieth century Colorado River flow regimes may not reoccur for many centuries—the time scale of climate system readjustment to the complete cessation of greenhouse gas emissions [Solomon *et al.*, 2009; Collins *et al.*, 2013].

The Colorado River declines do not stand alone as the only warming-related threat to Southwestern water supplies. The Rio Grande also has a grim prognosis [Reclamation, 2013; Elias *et al.*, 2015]. The drought in California has garnered national attention, and multiple studies have strongly implicated increasing temperatures as a contributor to these woes [Griffin and Anchukaitis, 2014; Belmecheri *et al.*, 2016; Diffenbaugh *et al.*, 2015; Mann and Gleick, 2015; Seager *et al.*, 2015a]. Southern California is particularly at risk, with a critical economy and a very large population, all coupled with a large reliance on both climate-threatened in-state, as well as Colorado River, water.

Adjusting to the new reality of rapid climate change will not be an easy or fast task; water management and water policy change slowly. The Colorado River is managed by a complex set of agreements, interstate compacts approved by Congress, international agreements, legislation, and court decrees set in place over the last 100 years [Verburg, 2011]. Most agreements were derived from twentieth century state-based negotiations with win/lose policy prescriptions that minimized basin-wide considerations of economic prosperity and potential harm [Adler, 2008]. None expressly includes climate change risk management, nor the provision for flow reductions that will be relentless on decadal timescales. New agreements often take years to put in place [Department of Interior, 2007]. The recently proposed structural deficit solution [Central Arizona Project, 2016], while important and laudable for the short term, will not solve the problem of large scale flow losses. With reduced water supplies, much will have to change in these agreements to address equity, economics, and social concerns on regional, state, basin-wide, and even national levels. Climate change threats to western water supplies are very real, and should prompt great concern and urgency among both water managers and the citizens of the Southwest.

References

- Adams, H. D., M. Guardiola-Claramonte, G. A. Barron-Gafford, J. C. Villegas, D. D. Breshears, C. B. Zou, P. A. Troch, and T. E. Huxman (2009), Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global-change-type drought, *Proc. Natl. Acad. Sci. U. S. A.*, 106(17), 7063–7066.

Acknowledgments

We thank the U.S. Bureau of Reclamation for data assistance and comments, and M. Kelly, D. Kenney, E. Kuhn, D. Lettenmaier, J. Robison, P. Troch, and C. Woodhouse for comments and discussion. A portion of this research was supported by the Colorado Water Institute, National Science Foundation, the NOAA Climate Assessment for the Southwest, and the U.S. Geological Survey Southwest Climate Science Center. We acknowledge the Downscaled CMIP3 and CMIP5 Climate and Hydrology Projections archive at: http://gdo-dcp.ucllnl.org/downscaled_cmip_projections as well as the World Climate Research Programme's Working Group on Coupled Modeling and the U.S. Department of Energy, Office of Science. The views expressed in this article are those solely of the authors. All data used in the analysis and conclusions in this paper are available from links in supporting information.

- Adler, R. W. (2007), *Restoring Colorado River Ecosystems: A Troubled Sense of Immensity*, Island Press, Wash.
- Adler, R. W. (2008), Revisiting the Colorado River Compact: Time for a change, *J. Land Resour. Environ. Law*, **28**, 19.
- Alcorn, B. (2015), *Colorado Basin River Forecast Center Final Reservoir Forecasts*, NOAA, Salt Lake City, Utah.
- Alcorn, B. (2016), *Colorado Basin River Forecast Center Final Reservoir Forecasts*, NOAA, Salt Lake City, Utah.
- Allen, C. D., et al. (2010), A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *For. Ecol. Manage.*, **259**(4), 660–684, doi:10.1016/j.foreco.2009.09.001.
- Allen, C. D., D. D. Breshears, and N. G. McDowell (2015), On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene, *Ecosphere*, **6**(8), 129.
- Ault, T. R., J. E. Cole, and S. St. George (2012), The amplitude of decadal to multidecadal variability in precipitation simulated by state-of-the-art climate models, *Geophys. Res. Lett.*, **39**, L21705, doi:10.1029/2012GL053424.
- Ault, T. R., J. E. Cole, J. T. Overpeck, G. T. Pederson, S. St. George, B. Otto-Bliessner, C. A. Woodhouse, and C. Deser (2013), The continuum of hydroclimate variability in Western North America during the Last Millennium, *J. Clim.*, **26**(16), 5863–5878, doi:10.1175/JCLI-D-11-00732.1.
- Ault, T. R., J. E. Cole, J. T. Overpeck, G. T. Pederson, and D. M. Meko (2014), Assessing the risk of persistent drought using climate model simulations and paleoclimate data, *J. Clim.*, **27**(20), 7529–7549, doi:10.1175/JCLI-D-12-00282.1.
- Ault, T. R., J. S. Mankin, B. I. Cook, and J. E. Smerdon (2016), Relative impacts of mitigation, temperature, and precipitation on 21st-century megadrought risk in the American Southwest, *Sci. Adv.*, **2**(10), e1600873, doi:10.1126/sciadv.1600873.
- Ayers, J., D. L. Ficklin, I. T. Stewart, and M. Strunk (2016), Comparison of CMIP3 and CMIP5 projected hydrologic conditions over the Upper Colorado River Basin, *Int. J. Climatol.*, **36**(11), 3807–3818, doi:10.1002/joc.4594.
- Barnett, T. P., and D. W. Pierce (2008), When will Lake Mead go dry?, *Water Resour. Res.*, **44**, W03201, doi:10.1029/2007WR006704.
- Barnett, T. P., and D. W. Pierce (2009), Sustainable water deliveries from the Colorado River in a changing climate, *Proc. Natl. Acad. Sci. U. S. A.*, **106**(18), 7334–7338.
- Belmecheri, S., F. Babst, E. R. Wahl, D. W. Stahle, and V. Trouet (2016), Multi-century evaluation of Sierra Nevada snowpack, *Nat. Clim. Change*, **2–3**.
- Betts, A. K., J. H. Ball, A. C. M. Beljaars, M. J. Miller, and P. A. Viterbo (1996), The land surface-atmosphere interaction: A review based on observational and global modeling perspectives, *J. Geophys. Res.*, **101**(D3), 7209–7225.
- Brekke, L., B. Harding, T. Piechota, B. Udall, C. Woodhouse, and D. Yates (2007), *Review of Science and Methods for Incorporating Climate Change Information into Reclamation's Colorado River Basin Planning Studies*, Reclamation, Washington, D. C.
- Brekke, L., B. Thrasher, Maurer, Edwin P., and T. Pruitt (2013), *Downscaled CMIP3 and CMIP5 Climate Projections*, Reclamation, Washington, D.C.
- Brekke, L., A. Wood, and T. Pruitt (2014), *Downscaled CMIP3 and CMIP5 Hydrology Projections*, Reclamation, Washington, D. C.
- Breshears, D. D., et al. (2005), Regional vegetation die-off in response to global-change-type drought, *Proc. Natl. Acad. Sci. U. S. A.*, **102**(42), 15,144–15,148.
- Brubaker, K. L., and D. Entekhabi (1996), Analysis of feedback mechanisms in land-atmosphere interaction, *Water Resour. Res.*, **32**(5), 1343–1357.
- Cayan, D. R., T. Das, D. W. Pierce, T. P. Barnett, M. Tyree, and A. Gershunov (2010), Future dryness in the southwest US and the hydrology of the early 21st century drought, *Proc. Natl. Acad. Sci. U. S. A.*, **107**(50), 21,271–21,276, doi:10.1073/pnas.0912391107.
- Central Arizona Project (2016), *Lower Basin Drought Contingency Plan*, Central Arizona Project, Phoenix, AZ.
- Christensen, N. S., and D. P. Lettenmaier (2007), A multimodel ensemble approach to assessment of climate change impacts on the hydrology and water resources of the Colorado River Basin, *Hydrol. Earth Syst. Sci. Discuss.*, **11**(4), 1417–1434.
- Christensen, N. S., A. W. Wood, N. Voisin, D. P. Lettenmaier, and R. N. Palmer (2004), The effects of climate change on the hydrology and water resources of the Colorado River basin, *Clim. Change*, **62**(1–3), 337–363.
- Collins, M., et al. (2013), Long-term climate change: Projections, commitments and irreversibility, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker et al., Cambridge Univ. Press, Cambridge, U. K.
- Collum, C., and T. McCann (2014), *Central Arizona Project Board Report Agenda Number 9*, Central Arizona Project, Phoenix, AZ.
- Colorado River Basin Stakeholders (2015), *Colorado River Basin Stakeholders Moving Forward to Address Challenges Identified in the Colorado River Basin Water Supply and Demand Study, Phase 1 Report*, Bureau of Reclamation, Washington, D. C.
- Cook, B. I., J. E. Smerdon, R. Seager, and S. Coats (2014), Global warming and 21st century drying, *Clim. Dyn.*, **43**(9–10), 2607–2627.
- Cook, B. I., T. R. Ault, and J. E. Smerdon (2015), Unprecedented 21st century drought risk in the American Southwest and Central Plains, *Sci. Adv.*, **1**(1), e1400082, doi:10.1126/sciadv.1400082.
- Dai, A. (2012), Increasing drought under global warming in observations and models, *Nat. Clim. Change*, **3**(1), 52–58, doi:10.1038/nclimate1633.
- Daly, C., R. P. Neilson, and D. L. Phillips (1994), A statistical-topographic model for mapping climatological precipitation over mountainous terrain, *J. Appl. Meteorol.*, **33**(2), 140–158.
- Das, T., D. W. Pierce, D. R. Cayan, J. A. Vano, and D. P. Lettenmaier (2011), The importance of warm season warming to western U.S. streamflow changes, *Geophys. Res. Lett.*, **38**, L23403, doi:10.1029/2011GL049660.
- Davis, T. (2016), Big CAP cuts coming as 3-state water agreement nears, *Ariz. Dly. Star*. [Available at http://tucson.com/news/local/big-cap-cuts-coming-as-state-water-agreement-nears/article_876e3aa6-6cf0-53ec-bd0c-95be8c6468ae.html.]
- Deems, J. S., T. H. Painter, J. J. Barsugli, J. Belnap, and B. Udall (2013), Combined impacts of current and future dust deposition and regional warming on Colorado River Basin snow dynamics and hydrology, *Hydrol. Earth Syst. Sci.*, **17**(11), 4401–4413, doi:10.5194/hess-17-4401-2013.
- Department of Interior (2007), *Record of Decision Colorado River Interim Guidelines for Lower Basin Shortages and the Coordinated Operations for Lake Powell and Lake Mead*, Dep. of Inter., Washington, D. C.
- Diffenbaugh, N. S., D. L. Swain, and D. Touma (2015), Anthropogenic warming has increased drought risk in California, *Proc. Natl. Acad. Sci. U. S. A.*, **112**(13), 3931–3936, doi:10.1073/pnas.1422385112.
- Elias, E. H., A. Rango, C. M. Steele, J. F. Mejia, and R. Smith (2015), Assessing climate change impacts on water availability of snowmelt-dominated basins of the Upper Rio Grande basin, *J. Hydrol. Reg. Stud.*, **3**, 525–546, doi:10.1016/j.ejrh.2015.04.004.
- Feng, S., and Q. Fu (2013), Expansion of global drylands under a warming climate, *Atmos. Chem. Phys.*, **13**(19), 10,081–10,094, doi:10.5194/acp-13-10081-2013.
- Ficklin, D. L., I. T. Stewart, and E. P. Maurer (2013), Climate change impacts on streamflow and subbasin-scale hydrology in the Upper Colorado River Basin, edited by V. Shah, *PLoS One*, **8**(8), e71297, doi:10.1371/journal.pone.0071297.
- Glennon, R. J. (1995), Coattails of the past: Using and financing the Central Arizona Project, *Ariz. State Law J.*, **27**, 677–756.
- Griffin, D., and K. J. Anchukaitis (2014), How unusual is the 2012–2014 California drought?, *Geophys. Res. Lett.*, **41**, 9017–9023, doi:10.1002/2014GL062433.

- Guentchev, G., J. J. Barsugli, and J. Eischeid (2010), Homogeneity of gridded precipitation datasets for the Colorado River Basin, *J. Appl. Meteorol. Climatol.*, *49*(12), 2404–2415, doi:10.1175/2010JAMC2484.1.
- Harding, B. L., A. W. Wood, and J. R. Prairie (2012), The implications of climate change scenario selection for future streamflow projection in the Upper Colorado River Basin, *Hydrol. Earth Syst. Sci.*, *16*(11), 3989–4007, doi:10.5194/hess-16-3989-2012.
- James, I. (2016), Lake Mead declines to lowest level in history, *Desert Sun*. [Available at <http://www.desertsun.com/story/news/environment/2016/05/19/lake-mead-declines-new-record-low/84597120/>]
- Johnson, R. (1977), *The Central Arizona Project, 1918–1968*, Univ. of Arizona Press, Tucson, Ariz.
- Koirala, S., Y. Hirabayashi, R. Mahendran, and S. Kanae (2014), Global assessment of agreement among streamflow projections using CMIP5 model outputs, *Environ. Res. Lett.*, *9*(6), 64,017, doi:10.1088/1748-9326/9/6/064017.
- Liang, X., D. P. Lettenmaier, and E. F. Wood (1996), One-dimensional statistical dynamic representation of subgrid spatial variability of precipitation in the two-layer variable infiltration capacity model, *J. Geophys. Res.*, *101*(D16), 21,403–21,422.
- Lukas, J., J. Barsugli, N. Doesken, I. Rangwala, and K. Wolter (2014), *Climate Change in Colorado A Synthesis to Support Water Resources Management and Adaptation*, Western Water Assessment, Boulder, CO.
- Mann, M. E., and P. H. Gleick (2015), Climate change and California drought in the 21st century, *Proc. Natl. Acad. Sci. U. S. A.*, *112*(13), 3858–3859, doi:10.1073/pnas.1503667112.
- Mann, M. E., Z. Zhang, S. Rutherford, R. S. Bradley, M. K. Hughes, D. Shindell, C. Ammann, G. Faluvegi, and F. Ni (2009), Global signatures and dynamical origins of the little ice age and medieval climate anomaly, *Science*, *326*(5957), 1256–1260, doi:10.1126/science.1177303.
- McCabe, G. J., and D. M. Wolock (2007), Warming may create substantial water supply shortages in the Colorado River basin, *Geophys. Res. Lett.*, *34*, L22708, doi:10.1029/2007GL031764.
- Meehl, G. A., C. Covey, K. E. Taylor, T. Delworth, R. J. Stouffer, M. Latif, B. McAvaney, and J. F. B. Mitchell (2007), THE WCRP CMIP3 multimodel dataset: A new era in climate change research, *Bull. Am. Meteorol. Soc.*, *88*(9), 1383–1394, doi:10.1175/BAMS-88-9-1383.
- Meko, D. M., C. A. Woodhouse, C. A. Baisan, T. Knight, J. J. Lukas, M. K. Hughes, and M. W. Salzer (2007), Medieval drought in the upper Colorado River Basin, *Geophys. Res. Lett.*, *34*, L10705, doi:10.1029/2007GL029988.
- Milly, P. C. D., and K. A. Dunne (2016), Potential evapotranspiration and continental drying, *Nat. Clim. Change*, *6*(10), 946–949, doi:10.1038/nclimate3046.
- Milly, P. C. D., K. A. Dunne, and A. V. Vecchia (2005), Global pattern of trends in streamflow and water availability in a changing climate, *Nature*, *438*(7066), 347–350, doi:10.1038/nature04312.
- Moss, R. H., et al. (2010), The next generation of scenarios for climate change research and assessment, *Nature*, *463*(7282), 747–756.
- Nash, L. L., and P. H. Gleick (1991), Sensitivity of streamflow in the Colorado basin to climatic changes, *J. Hydrol.*, *125*(3), 221–241.
- National Research Council (2007), *Colorado River Basin Water Management: Evaluating and Adjusting to Hydroclimatic Variability*, The Natl. Acad. Press, Washington, D. C.
- Norris, J. R., R. J. Allen, A. T. Evan, M. D. Zelinka, C. W. O'Dell, and S. A. Klein (2016), Evidence for climate change in the satellite cloud record, *Nature*, *536*(7614), 72–75, doi:10.1038/nature18273.
- Nowak, K., M. Hoerling, B. Rajagopalan, and E. Zagona (2012), Colorado River Basin hydroclimatic variability, *J. Clim.*, *25*(12), 4389–4403, doi:10.1175/JCLI-D-11-00406.1.
- Overpeck, J. T. (2013), Climate science: The challenge of hot drought, *Nature*, *503*(7476), 350–351.
- Oyler, J. W., S. Z. Dobrowski, A. P. Ballantyne, A. E. Klene, and S. W. Running (2015a), Artificial amplification of warming trends across the mountains of the western United States, *Geophys. Res. Lett.*, *42*, 153–161, doi:10.1002/2014GL02803.
- Oyler, J. W., A. Ballantyne, K. Jencso, M. Sweet, and S. W. Running (2015b), Creating a topoclimatic daily air temperature dataset for the conterminous United States using homogenized station data and remotely sensed land skin temperature, *Int. J. Climatol.*, *35*(9), 2258–2279, doi:10.1002/joc.4127.
- Painter, T. H., J. S. Deems, J. Belnap, A. F. Hamlet, C. C. Landry, and B. Udall (2010), Response of Colorado River runoff to dust radiative forcing in snow, *Proc. Natl. Acad. Sci. U. S. A.*, *107*(40), 17,125–17,130.
- Pitman, A. J. (2003), The evolution of, and revolution in, land surface schemes designed for climate models, *Int. J. Climatol.*, *23*(5), 479–510, doi:10.1002/joc.893.
- Prein, A. F., G. J. Holland, R. M. Rasmussen, M. P. Clark, and M. R. Tye (2016), Running dry: The U.S. Southwest's drift into a drier climate state, *Geophys. Res. Lett.*, *43*, 1272–1279, doi:10.1002/2015GL066727.
- Rajagopalan, B., K. Nowak, J. Prairie, M. Hoerling, B. Harding, J. Barsugli, A. Ray, and B. Udall (2009), Water supply risk on the Colorado River: Can management mitigate? *Water Resour. Res.*, *45*, W08201, doi:10.1029/2008WR007652.
- Rangwala, I., T. Bardsley, M. Pescinski, and J. Miller (2015), *SNOTEL Sensor Upgrade has Caused Temperature Record Inhomogeneities for the Intermountain West: Implications for Climate Change Impact Assessments*, West. Water Assess., Boulder, Colo.
- Ray, A. J., J. J. Barsugli, K. B. Averyt, K. Wolter, M. Hoerling, N. Doesken, B. Udall, and R. S. Webb (2008), *Climate change in Colorado: a synthesis to support water resources management and adaptation*, Western Water Assessment, Boulder, Colo.
- Reclamation (2007), *Final Environmental Impact Statement Colorado River Interim Guidelines for Lower Basin Shortages and Coordinated Operations for Lake Powell and Lake Mead*, Bureau of Reclamation, Boulder City, NV.
- Reclamation (2012), *Colorado River Basin Water Supply and Demand Study*, U.S. Dep. of the Inter., Bur. of Reclam., Tech. Serv. Cent., Denver, Colo.
- Reclamation (2013), *West-Wide Climate Risk Assessment: Upper Rio Grande Impact Assessment*, U.S. Department of the Interior, Bureau of Reclamation, Albuquerque, NM.
- Reclamation (2016), *SECURE Water Act Section 9503(c) – Reclamation Climate Change and Water*, U.S. Department of the Interior, Bureau of Reclamation, Denver, Colo.
- Salzer, M. W., and K. F. Kipfmüller (2005), Reconstructed temperature and precipitation on a millennial timescale from tree-rings in the southern Colorado Plateau, USA, *Clim. Change*, *70*(3), 465–487.
- Salzer, M. W., A. G. Bunn, N. E. Graham, and M. K. Hughes (2014), Five millennia of paleotemperature from tree-rings in the Great Basin, USA, *Clim. Dyn.*, *42*(5–6), 1517–1526.
- Scheff, J., and D. M. Frierson (2012), Robust future precipitation declines in CMIP5 largely reflect the poleward expansion of model subtropical dry zones, *Geophys. Res. Lett.*, *39*, L18704, doi:10.1029/2012GL052910.
- Seager, R., et al. (2007), Model projections of an imminent transition to a more arid climate in Southwestern North America, *Science*, *316*(5828), 1181–1184, doi:10.1126/science.1139601.
- Seager, R., M. Ting, C. Li, N. Naik, B. Cook, J. Nakamura, and H. Liu (2012), Projections of declining surface-water availability for the southwestern United States, *Nat. Clim. Change*, *3*(5), 482–486, doi:10.1038/nclimate1787.
- Seager, R., M. Hoerling, S. Schubert, H. Wang, B. Lyon, A. Kumar, J. Nakamura, and N. Henderson (2015a), Causes of the 2011–14 California Drought*, *J. Clim.*, *28*(18), 6997–7024.

- Seager, R., A. Hooks, A. P. Williams, B. Cook, J. Nakamura, and N. Henderson (2015b), Climatology, variability, and trends in the US vapor pressure deficit, an important fire-related meteorological quantity*, *J. Appl. Meteorol. Climatol.*, *54*(6), 1121–1141.
- Seneviratne, S. I., T. Corti, E. L. Davin, M. Hirschi, E. B. Jaeger, I. Lehner, B. Orlowsky, and A. J. Teuling (2010), Investigating soil moisture–climate interactions in a changing climate: A review, *Earth Sci. Rev.*, *99*(3), 125–161.
- Solomon, S., G.-K. Plattner, R. Knutti, and P. Friedlingstein (2009), Irreversible climate change due to carbon dioxide emissions, *Proc. Natl. Acad. Sci. U. S. A.*, *106*(6), 1704–1709, doi:10.1073/pnas.0812721106.
- Swann, A. L. S., F. M. Hoffman, C. D. Koven, and J. T. Randerson (2016), Plant responses to increasing CO₂ reduce estimates of climate impacts on drought severity, *Proc. Natl. Acad. Sci. U. S. A.*, *113*, 10,019–10,024, doi:10.1073/pnas.1604581113.
- Taylor, K. E., R. J. Stouffer, and G. A. Meehl (2012), An overview of Cmp5 and the experiment design, *Bull. Am. Meteorol. Soc.*, *93*(4), 485–498.
- Tipton and Kalmbach, Inc. (1965), *Water Supplies of the Colorado River*, Upper Colo. River Comm., Denver, Colo.
- Udall, J. (1983), Floods reveal water policy chaos, *High Ctry. News*, *15*(18), 1–10.
- Ukkola, A. M., I. C. Prentice, T. F. Keenan, A. I. J. M. van Dijk, N. R. Viney, R. B. Myneni, and J. Bi (2015), Reduced streamflow in water-stressed climates consistent with CO₂ effects on vegetation, *Nat. Clim. Change*, *6*(1), 75–78, doi:10.1038/nclimate2831.
- Vano, J. A., and D. P. Lettenmaier (2014), A sensitivity-based approach to evaluating future changes in Colorado River discharge, *Clim. Change*, *122*(4), 621–634, doi:10.1007/s10584-013-1023-x.
- Vano, J. A., T. Das, and D. P. Lettenmaier (2012), Hydrologic sensitivities of Colorado River runoff to changes in precipitation and temperature*, *J. Hydrometeorol.*, *13*(3), 932–949, doi:10.1175/JHM-D-11-069.1.
- Vano, J. A., et al. (2014), Understanding uncertainties in future Colorado River streamflow, *Bull. Am. Meteorol. Soc.*, *95*(1), 59–78.
- Verburg, K. O. (2011), *Colorado River Documents 2008*, Reclamation, Denver, Colo.
- Weiss, J. L., C. L. Castro, and J. T. Overpeck (2009), Distinguishing pronounced droughts in the Southwestern United States: Seasonality and effects of warmer temperatures, *J. Clim.*, *22*(22), 5918–5932, doi:10.1175/2009JCLI2905.1.
- Williams, A. P., R. Seager, J. T. Abatzoglou, B. I. Cook, J. E. Smerdon, and E. R. Cook (2015), Contribution of anthropogenic warming to California drought during 2012–2014, *Geophys. Res. Lett.*, *42*, 6819–6828, doi:10.1002/2015GL064924.
- Wines, M. (2014), Colorado River drought forces a painful reckoning for states, *New York Times*, 6th January.
- Woodhouse, C. A., S. T. Gray, and D. M. Meko (2006), Updated streamflow reconstructions for the Upper Colorado River basin, *Water Resour. Res.*, *42*, W05415, doi:10.1029/2005WR004455.
- Woodhouse, C. A., G. T. Pederson, K. Morino, S. A. McFee, and G. J. McCabe (2016), Increasing influence of air temperature on upper Colorado River streamflow, *Geophys. Res. Lett.*, *43*, 2174–2181, doi:10.1002/2015GL067613.