

R E V I E W



The type I interferon response during viral infections: a “SWOT” analysis

Giel R. Gaajetaan, Cathrien A. Bruggeman and Frank R. Stassen*

Department of Medical Microbiology, Maastricht University Medical Center, The Netherlands

SUMMARY

The type I interferon (IFN) response is a strong and crucial moderator for the control of viral infections. The **strength** of this system is illustrated by the fact that, despite some temporary discomfort like a common cold or diarrhea, most viral infections will not cause major harm to the healthy immunocompetent host. To achieve this, the immune system is equipped with a wide array of pattern recognition receptors and the subsequent coordinated type I IFN response orchestrated by plasmacytoid dendritic cells (pDCs) and conventional dendritic cells (cDCs). The production of type I IFN subtypes by dendritic cells (DCs), but also other cells is crucial for the execution of many antiviral processes. Despite this coordinated response, morbidity and mortality are still common in viral disease due to the ability of viruses to exploit the **weaknesses** of the immune system. Viruses successfully evade immunity and infection can result in aberrant immune responses. However, these weaknesses also open **opportunities** for improvement via clinical interventions as can be seen in current vaccination and antiviral treatment programs. The application of IFNs, Toll-like receptor ligands, DCs, and antiviral proteins is now being investigated to further limit viral infections. Unfortunately, a common **threat** during stimulation of immunity is the possible initiation or aggravation of autoimmunity. Also the translation from animal models to the human situation remains difficult. With a **Strengths–Weaknesses–Opportunities–Threats** (“SWOT”) analysis, we discuss the interaction between host and virus as well as (future) therapeutic options, related to the type I IFN system. Copyright © 2011 John Wiley & Sons, Ltd.

Received: 15 April 2011; Revised: 26 August 2011; Accepted: 31 August 2011

INTRODUCTION

For centuries, infectious diseases have been the most common cause of morbidity and mortality

worldwide. Due to achievements like vaccination and antimicrobial drugs, many infectious diseases can now be prevented or controlled. Most striking in this respect is the development of a vaccine against smallpox, a lethal virus that globally claimed millions of lives. Although the vaccination procedure was already developed in the 18th century, it lasted until the end of the 20th century before the world was declared smallpox-free. Based on this success, there was great confidence that viral infections could be conquered definitely, either by vaccination or by antiviral drug treatment. Inspired by these successes, the US Surgeon General William Stewart stated in 1967 that “The time has come to close the book on infectious diseases”. Unfortunately, the future has shown otherwise.

In 1983, the HIV was discovered as the AIDS causing agent. Despite massive efforts, HIV is still a major problem worldwide [1,2]. In addition, the rise of new (variants of) viruses like influenza A strains [3,4] and severe acute respiratory syndrome corona virus (SARS CoV) and their potential pandemic threat is a general and realistic concern [5]. Furthermore,

*Correspondence author: Dr. F. R. Stassen, Department of Medical Microbiology, Maastricht University Medical Center, P.Debyelaan 25, 6202 AZ, Maastricht, The Netherlands. E-mail: f.stassen@maastrichtuniversity.nl

Abbreviations

APOBEC3G, apolipoprotein B-mRNA-editing enzyme-catalytic polypeptide-like 3 G; BST-2, bone marrow stromal antigen-2; cDC, conventional dendritic cell; CLR, C-type lectin receptor; COPD, chronic obstructive pulmonary disease; CpG ODN, CpG oligodeoxynucleotides; DAI, DNA-dependent activator of IFN-regulatory factors; DC, dendritic cell; dsRNA, double stranded RNA; HPV, human papilloma virus; IFNAR, interferon α/β receptor; IRF, interferon response factor; ISG, interferon-stimulated genes; ISG15, IFN-stimulated protein of 15 kDa; LPS, lipopolysaccharide; MDA5, melanoma differentiation-associated gene 5; miRNA, microRNA; Mx1, myxovirus resistance 1; NF- κ B, nuclear factor κ B; NK, natural killer; NS1, non-structural 1; OAS1, oligoadenylate synthetase 1; PAMP, pathogen-associated molecular patterns; pDC, plasmacytoid dendritic cell; PKR, protein kinase R; Poly(I:C), polyinosinic:polycytidylic acid; PRR, pattern recognition receptor; RIG-I, retinoic acid inducible gene 1; RNaseL, ribonuclease L; SARS CoV, severe acute respiratory syndrome corona virus; SLE, systemic lupus erythematosus; ssRNA, single stranded RNA; SWOT, strengths-Weaknesses-Opportunities-Threats; TLR, Toll-like receptor; TRIM5 α , tripartite motif 5 α .

seasonal respiratory viral infections and various other viruses can cause major inconvenience in healthy people and can be life-threatening in the immunocompromised [5,6]. Thus, despite vaccines and antiviral drugs, viral disease is still common and requires development of additional therapeutics.

In this review, we apply a Strengths–Weaknesses–Opportunities–Threats (SWOT) analysis to discuss virus-immune interactions and speculate on (im) possibilities how to use these interactions in view of new treatment options.

STRENGTHS

Once the virus has been able to cross first barriers like the skin or mucosa, the **strength** of the host's natural defense system will determine the outcome of the infection. In the succeeding text, we will briefly discuss some of the initial key steps involved in the antiviral response (see also Figure 1).

Recognition: pattern recognition receptors

Before an appropriate immune response can be generated, the virus needs to be recognized. For this, immune cells are equipped with different groups of receptors, which are able to sense microbial intruders including viruses. These pattern recognition receptors (PRRs) recognize pathogen-associated molecular patterns (PAMPs), which are fundamentally different from host structures. One of the first discovered and best characterized PRRs are the Toll-like receptors (TLRs) [7–10], which are mostly present on antigen-presenting cells like macrophages and dendritic cells (DCs) [8,9], but also on non-immune cells like fibroblasts and epithelial cells [10]. These transmembrane receptors are located on the cell surface or at the endosome [7,9–11]. The cell surface-located TLRs recognize mainly lipids and proteins from bacteria and yeasts [10]. Viruses, on the other hand, are intracellular parasites, which




Figure 1. Schematic overview of different signal transduction pathways that are activated in plasmacytoid dendritic cells (pDCs) and conventional dendritic cell (cDCs) following viral encounters. In general, pDCs endocytose the virus and subsequently Toll-like receptor 7 (TLR7) and/or TLR9 is stimulated. Interferon response factor 7 (IRF7) is activated and induces transcription of IFN α/β . Besides execution of many antiviral functions, autocrine signaling via the interferon α/β receptor (IFNAR) also induces more type I IFN production. In contrast, infection of or endocytosis by cDCs results in activation of the cytoplasmic pattern recognition receptors, TLR3, and TLR8. Accordingly, IRF3 and nuclear factor- κ B (NF- κ B) facilitate transcription of IFN β and proinflammatory cytokines. Via IFNAR, IRF7 is activated and induces production of type IFN α/β . Red indicates major routes, dotted arrows indicate minor routes

may explain the endosomal localization of the viral nucleic acid-recognizing TLR3, TLR7, TLR8, and TLR9 (Figure 1) [11–19]. Also, this endosomal location of the TLRs probably serves to ensure tolerance for “self” molecules and to promote ligand accessibility [10,14]. Interestingly, in addition to the well-known lipopolysaccharides from Gram-negative bacteria, cell surface TLRs have also been associated with viral recognition. TLR4 has been shown to recognize the fusion protein of RSV [10,20]. Likewise, next to the recognition of Gram-positive bacteria, TLR2 is involved in detection of various DNA viruses like HSV1 and 2, measles virus, vaccinia virus, and CMV [21–23]. Interestingly, this TLR2-dependent detection seems to be regulated especially by monocytes [21,22,24].

In addition to the well-described TLRs, other PRRs also play an important role in viral recognition. The cytoplasmic PRRs, such as retinoic acid-inducible gene I (RIG-I), melanoma differentiation-associated gene 5 (MDA5), and DNA-dependent activator of IFN-regulatory factors (DAI), recognize viral nucleic acids [25,26] and are, in contrast to TLRs, expressed in all cells. RNA viruses are differentially recognized by RIG-I and MDA5, but activate similar pathways (Figure 1) [26–28]. Although RIG-I can respond to both positive and negative strand RNA viruses, MDA5 senses mainly picornaviruses like rhinovirus and poliovirus [29,30]. Earlier data suggested that MDA5 preferentially binds long dsRNA (picornaviruses), whereas shorter fragments of dsRNA and other specific nucleotide sequences are sensed by RIG-I [30,31]. However, some viruses can be detected by both receptors [29]. Also, the recently discovered receptor DAI is important for intracellular detection of viral DNA [32,33].

C-type lectin receptors (CLRs) and NOD-like receptors (NLRs) also belong to the large family of PRRs. CLRs are present on DCs and recognize carbohydrate structures present on pathogens [34,35] and are especially important for induction of antigen presentation to T cells, but also in modulating TLR responses [36]. NLRs, a group of cytoplasmic proteins formerly thought to detect only bacterial PAMPs, also sense RNA [37–39] and DNA viruses [33,40,41]. This induces the production of the proinflammatory cytokines IL-1 β and IL-18 via the inflammasome, a complex composed of NLRs, and leads to the recruitment of immune cells to the site of infection [42,43].

Taken together, the innate immune system is equipped with a large variety of PRRs and this extended array is essential to sense the various microbial components and to prevent or limit viral spread as much as possible [7,44,45].

Implementation of antiviral immunity: conventional and plasmacytoid dendritic cells

After recognition of a virus, a cell-dependent signaling cascade will be initiated. Infection of non-immune cells usually results in detection of viral DNA/RNA or their intermediates by the cytoplasmic PRRs and the production of IFN β , which is required to limit the infection. This antiviral cytokine also primes cells to produce other type I IFNs, which comprise all IFN α subtypes, IFN β , and various other IFN types, essential to initiate production of antiviral proteins [46].

Dendritic cells are better equipped than non-immune cells for the initiation of an antiviral response. Conventional dendritic cells (cDCs) recognize viral invaders with both extracellular (TLR 4 and CLRs) and intracellular PRR (TLR3, 8, RIG-I, MDA5), which are highly expressed on cDCs (Figure 1) [11,12,47,48]. As in infection of non-immune cells, viral nucleic acids need to be detected before IFN β and other type I IFNs can be produced.

For the successful eradication or control of the virus, the intervention of plasmacytoid dendritic cells (pDCs) is indispensable. The pDC is one of the few cells that express both TLR7 and TLR9 (Figure 1), allowing detection of an extended repertoire of viruses. To initiate the antiviral response, viruses or virus-infected cells are first internalized by endocytosis or phagocytosis, respectively, and subsequently recruited to the endolysosomes of the pDC [49]. The acidic environment disassembles the virus, and viral nucleic acids are subsequently recognized by TLR7 or TLR9 [50,51]. Ultimately, massive amounts of type I IFN are produced. In contrast to cDCs and non-immune cells, in pDCs the TLRs contribute significantly more to viral recognition than the cytoplasmic PRRs RIG-I and MDA5 [26,52,53]. Consequently, pDCs are less dependent on steps in the viral life cycle for recognition, which significantly accelerates the response to an infection in these DCs.

The difference in response time between pDC and cDC is also because of marked differences in

intracellular signaling cascades that are activated following PRR stimulation. In cDCs, viral components stimulate the TLRs (apart from the cytoplasmic PRRs) resulting in phosphorylation of interferon-regulatory factor 3 (IRF3). IRF3 is essential for the production of proinflammatory cytokines and IFN β (the first wave IFN) and is constitutively expressed, not only in cDCs but in most cell types [10,54]. Next, because of autocrine or paracrine signaling through the interferon- α/β receptor (IFNAR), IRF7 is activated, leading to the production of all type I IFNs including the various IFN α subtypes (the second wave IFN) [55,56]. Alternatively, in pDCs IRF7 is constitutively expressed and activated immediately after stimulation of TLR7 or TLR9, and thus no prior phosphorylation of IRF3 or autocrine/paracrine signaling is required (Figure 1) [48,52,57–60]. Accordingly, a robust antiviral response is initiated that, in contrast to the response seen in cDCs, is rapid and characterized by the production of high amounts of type I IFNs [61,62].

Consequently, the pDC is clearly the major antiviral cell type due to its rapid and abundant IFN α production. Yet, the cDC is indispensable for clearance of a viral infection. This can be illustrated by the function of TLR8 expressed by cDCs. This receptor is similar to TLR7 in pDCs and also recognizes viral ssRNA. Interestingly, stimulation of TLR8 on cDCs and TLR7 on pDCs results in entirely different responses [63]. Although the pDC produces mainly IFN α , the cDC induces a pro-inflammatory profile in which nuclear factor- κ B (NF- κ B) is activated for the production of TNF- α and IL-6 [64]. More importantly, IL-12 is produced (Figure 1). This cytokine augments the cytolytic activity of natural killer (NK) cells and also induces the production of the immunoregulatory cytokine IFN γ by T and NK cells [65]. Thus, although both DC subsets use different antiviral pathways, they are certainly not mutually exclusive in their response to viral infection. Because of their different cytokine patterns, pDCs and cDCs respond collaboratively to viral infection and connect innate and adaptive immunity [66]. Communication and cooperation between these two DC subsets are vital to induce appropriate immune responses towards invading pathogens.

Effector: Type I interferon

The type I IFNs are key effector molecules of the innate immune system and are essential for the

antiviral response towards a plethora of viruses. In humans, the type I IFN family comprises 13 IFN α subtypes, IFN β , IFN κ , IFN ϵ , IFN ω , IFN τ , and IFN δ , and all these molecules engage the ubiquitously expressed IFNAR. Binding to IFNAR then stimulates more than 300 interferon-stimulated genes (ISGs) [67,68], which subsequently induce an antiviral state. The antiviral state is a collective term for limitation of viral replication, viral resistance of neighboring cells, and apoptosis of virally infected cells.

Although IFNAR signaling induces the transcription of more than 300 ISGs, surprisingly, few of these genes encode proteins with direct antiviral effects [69]. Those proteins target viruses in many different ways (Figure 1). For example, the protein ISG15 (IFN-stimulated protein of Mr 15 000) has been reported to prevent virus-mediated degradation of IRF3 [70], to enhance NF- κ B signaling [71], and to modulate the immune response [72]. Myxovirus resistance 1 (Mx1) proteins target viral nucleocapsid-like structures [73] and mediate vesicle trafficking in the ER to effectively trap essential viral components and subsequently degrade them [74,75]. The enzyme 2',5'-oligoadenylate synthetase 1 (OAS1) accumulates after signaling through the IFNAR by type I IFN. When exposed to dsRNA, this enzyme gains activity that eventually leads to the activation of ribonuclease L (RNaseL), concomitantly enabling cleavage of cellular and viral RNAs [69,76]. Protein kinase R (PKR) is also initially inactive. Type I IFN induces accumulation of PKR and dsRNA activates PKR to inhibit translation [77]. For a more detailed overview of the ISG function, we would like to refer to the excellent review recently published by Sadler *et al.* [69].

Interferons also induce antiviral proteins termed restriction factors. A good example is the bone marrow stromal antigen-2 (BST-2) protein, which restricts the release of fully formed progeny virions from infected cells. This tetherin protein showed activity against various viruses, including HIV [78–80]. Another restriction factor is apolipoprotein B mRNA-editing enzyme-catalytic polypeptide-like 3G (APOBEC3G), which leads to degradation of HIV DNA [81,82]. The restriction factor tripartite motif (TRIM) 5 α seems to counteract capsid formation by HIV (reviewed by Sastri *et al.*) [83].

In addition, many proteins stimulated by type I IFN are involved in IFN signaling (IRF7, RIG-I, MDA5, TLRs), thereby amplifying the IFN response (positive feedback). IFNs also induce or modulate

adaptive immune responses by upregulating MHC class I and II, to facilitate T and B cell stimulation [84,85]. Finally, IFNs promote leukocyte accumulation at sites of infection by promoting vascular adhesion molecule expression and induction of chemokines, which are essential in leukocyte recruitment [86].

Recently, a new type I IFN-dependent antiviral pathway has been suggested. Pedersen *et al.* demonstrated that IFN β rapidly induced the expression of several microRNAs (miRNAs) both in a hepatocarcinoma cell line (Huh cells) and primary hepatocytes [87]. These small non-coding RNA molecules are post-transcriptional regulators that inhibit gene expression by translational repression, mRNA cleavage and deadenylation [87,88]. Intriguingly, eight of these IFN β -induced miRNAs showed sequence-predicted targets within the HCV genomic RNA. Moreover, application of synthetic miRNA-mimics resulted in antiviral effects similar to those induced by IFN β , whereas anti-miRNA markedly reduced the IFN β -mediated antiviral effect [87]. In addition, it has recently been shown that hepatic miRNA expression might be a useful tool for predicting the therapeutic outcome of a pegylated IFN/ribavirin combination therapy, further emphasizing the potential role of miRNAs in IFN-mediated antiviral effects [89].

In conclusion, the presence of a wide variety of PRRs enables the detection of multiple viral ligands present during infection. Activation of the PRR-DC-type I IFN axis (and especially the TLR7/9-pDC-IFN α axis) induces a rapid response to the virus. The many ISGs and the diversity of the type I IFNs that can be stimulated or produced, respectively, enables a coordinated response to the various viral infections, leading to control or elimination of the viral intruder.

WEAKNESSES

In the previous section, we described how well-equipped the immune system is to protect the host against viral infections. Nevertheless, viruses can evade or influence the immune response by targeting certain weaknesses of the immune system resulting in (severe) disease.

Modulation of the type I interferon response by viruses

Because of the strong antiviral and immunoregulatory role of type I IFN, viruses developed a large variety of anti-type I IFN mechanisms. Consequently, nearly all steps of the type I IFN pathway can be blocked or manipulated by different viruses for their own benefit (Table 1) [90,91]. For example, PRR signaling can be suppressed by inhibition of downstream signaling or by sequestration of typical viral nucleic acids like dsRNA [90]. In this way, viral recognition is inhibited. Alternatively, viruses interfere with the production of type I IFN by targeting the transcription factors IRF3 and IRF7. The proteins involved in IRF activation are inactivated or IRF mimics are synthesized, which compete with the host IRFs [90,92,93]. Also, binding of IFN to IFNAR can be prevented by a virally-encoded type I IFN receptor, as observed during vaccinia virus infection [94,95]. Finally, the antiviral or immuno-regulatory effects of type I IFN are inhibited by targeting various ISGs and thereby facilitating viral replication and preventing immune recognition [96–99].

Alternatively, virus-related morbidity and mortality are not only due to virus-induced immune evasion, which facilitates extensive viral replication, but may also result from a concomitant, an inappropriate, and an exaggerated response of

Table 1. Viral inhibition of the type I IFN pathway

General target	Specific target	Virus examples	References
PRR signaling	almost all proteins	Ebola, influenza, HCV	[90,91]
Transcription	IRF3, IRF7,	Paramyxoviruses, Rabies	[90,92,93]
Cytokine receptors	IFNAR	Vaccinia	[94,95]
ISGs	ISG15, mx1, OAS1, PKR, for example	SARS, influenza, HCV	[96–99]

PRR, pattern recognition receptor; ISGs, interferon-stimulated genes; IRF, interferon response factor; IFNAR, interferon α/β receptor; SARS, severe acute respiratory syndrome.

the immune system with devastating consequences for the host. A typical example of a combination of efficient inhibition of the type I IFN response together with an exaggerated immune response is provided by the highly pathogenic avian H5N1 influenza strain. The non-structural 1 (NS1) protein of H5N1 is an effective antagonist of the type I IFN pathway [100–102]. This results not only in high viral replication but also in an inflammatory response characterized by high levels of cytokines like TNF α [103]. This “hypercytokinemia” or “cytokine storm” results in excessive infiltration of inflammatory cells into the lungs [103–106]. Also, higher plasma levels of inflammatory mediators were detected in deceased H5N1 patients compared with survivors [107]. The deregulation of type I IFN by H5N1 is also observed in the highly virulent 1918 H1N1 influenza strain and the Ebola and Marburg viruses [108–112], in which both viral and immune pathology result in severe disease [6]. Thus, the increased resistance to the antiviral effects of IFN enhances viral replication and evokes an aberrant proinflammatory response characterized by high levels of cytokines and chemokines, which induces the pulmonary injury observed in H5N1 patients.

Likewise, the devastating effects of an HIV infection may also result from such a combination. HIV infection results in progressive immune deficiency, impaired adaptive responses, low CD4 T cell counts and increases susceptibility to opportunistic infections. One of the earliest findings during the AIDS epidemic was a deficient IFN α production in HIV-infected patients. Next to a lower number of IFN-producing cells, also each cell produced less IFN α in response to HIV [113,114]. The decrease in IFN α can be due to the Vpr protein of HIV, which strongly inhibits type I IFN production by pDCs [115]. In addition, the effects of IFN α are antagonized by the HIV protein Vpu, which induces degradation of the restriction factor BST2 [79,116]. However, during the chronic phase of HIV infection, it is hypothesized that IFN α contributes to the decline of the immune system by inducing apoptosis of CD4 T cells. Because of the non-infectious interaction between the HIV-bound gp120 protein and the CD4 receptor on pDCs, IFN α is produced and this results in killing (possibly by pDCs) of uninfected CD4 T cells [114–117]. Thus, although apoptosis of infected cells is usually a protective mechanism to prevent viral spread [118,119], here, it results in a distinct advantage for the virus due to the decreased immune control by CD4 T cells.

Thus, despite the strength of the type I IFN system, viruses have evolved mechanisms to evade or manipulate the system to guarantee their survival. Among others, this is predominantly accomplished through interfering with PRR signaling, inhibition of IRF3 and IRF7 activation and targeting ISGs.

OPPORTUNITIES

The search for therapies has led to the development of vaccines and antiviral drugs, which resulted in an impressive reduction in virus-related morbidity and mortality. Unfortunately, both vaccination and antiviral drugs are not sufficient to prevent or control all viral infections, which make it imperative to develop novel therapies. As a result, immune-based therapies are currently under development as new treatment methods. This may provide new opportunities for the treatment of acute or chronic viral infections (Figure 2).




Figure 2. Antiviral therapy options. Current therapy involves antiviral drugs, vaccination, and IFN α therapy for treatment of HCV patients. In addition to these therapies, treatment with type I and III IFNs can counteract acute and local infections, TLR ligands have shown to be beneficial in various viral infections, and DC transfer could be attractive where dysfunctional or limited numbers of DCs contribute to the pathogenesis

Interferon therapy revisited

A plausible approach to treat virally infected patients is the administration of type I IFN. Indeed, pegylated IFN α in combination with the antiviral drug ribavirin is commonly used in treating patients with a chronic HCV infection. Although this therapy is effective in nearly 50% of the cases, the administration of pegylated IFN α is associated with severe side effects [120–122]. Normally, during viral infections, type I IFN gives the “sick-signal” that results in fever. Patients treated with type I IFN have to endure these feverish periods for prolonged periods of time. In addition, hematologic and psychological problems have been frequently reported during treatment periods. Also with respect to HIV, positive effects of IFN-treatment have been reported both *in vitro* [123,124], as in clinical trials [125–129]. On the other hand, (excessive) IFN α can contribute to the immunopathogenesis (reviewed by Herbeuval *et al.*) [117]. Thus, it remains controversial whether IFN α is beneficial or detrimental in HIV, because both underproduction and overproduction of IFN α can induce severe effects in the host.

Nonetheless, because of their strong antiviral effects, type I IFNs remain attractive drugs for antiviral therapy. In particular during acute (respiratory) infections, IFNs may be an interesting therapy. This requires no systemic and chronic application of IFNs as observed in HCV patients, which may, therefore, significantly reduce the observed side effects. Local application, for example, by a nasal spray, has been shown to be effective in the prevention of seasonal respiratory infections without causing severe side effects [86,130]. This administration route might be, particularly, attractive for the prevention of virus-induced exacerbations in chronic obstructive pulmonary disease and asthmatic patients in which impaired IFN production may be an important mechanism contributing to virus-induced exacerbations [131,132]. IFN α also showed promising effects in severe acute respiratory syndrome (SARS) [133–135] and can be very important to induce an adequate immune response and possibly suppress excessive inflammatory responses observed in SARS [136–139]. Interestingly, also other members of the IFN family can be used to prevent or treat viral respiratory infections. The recently discovered type III IFNs (or IFN λ 1 and IFN λ 2/3) show strong antiviral effects against respiratory viral infections [131,140–142], especially when given prophylactically [142,143].

Toll-like receptor ligands

Because stimulation of TLRs by antigenic microbial epitopes is sufficient to induce a full-blown immune response, TLRs seem a likely target for antimicrobial therapy. Indeed, synthetic variants of the microbial structures have been shown to induce natural responses without the need for infection, and this quality has been used extensively to improve the efficacy of vaccines. For example, vaccines composed of a mixture of TLR ligand and antigens have been shown to be more effective than antigens alone [144–148]. Moreover, TLR ligands covalently linked to peptides are even superior in their ability to induce specific CD8+ T cells [149].

When a direct antiviral response is required, the use of synthetic TLR3, TLR7, TLR8, or TLR9 ligands can be considered. Both *in vitro* and *in vivo* studies have shown that prophylactic treatment with the dsRNA mimic polyinosinic:polycytidylic acid (poly (I:C)) and CpG oligodeoxynucleotides (CpG ODNs) specific for TLR3 and TLR9, respectively, is protective during viral infection [150–153]. Depending on virus and cell type, different types of CpG ODNs can be applied to initiate an appropriate response [154–156]. Also, TLR7 and TLR8 may be therapeutic targets. For stimulation of these TLRs, imidazoquinolones (e.g. resiquimod and imiquimod) are the best known ligands, and these small molecular weight compounds have indeed been shown to possess antiviral properties [15,16,157–160], although their immunostimulatory and antiviral effect may be limited compared with poly (I:C) and CpG ODNs [161]. Interestingly, the use of imiquimod as a cream to treat human papillomavirus-induced genital warts has already been approved [16,145,162]. TLR ligands can also reduce HCV viremia [163–165] and even HIV could be targeted [166]. Besides stimulation of type I IFN production, TLR ligands also initiate immunoregulatory mechanisms [167]. This is particularly important for the generation of the adaptive immune response and immunological memory. Nonetheless, at this time, few TLR ligands have been approved for clinical application in treating viral disease [13,148].

Dendritic cell transfer

During various viral infections, pDCs (and cDCs) are less functional or are present in lower numbers [168–170]. This is, for example, observed in HCV-infected [171,172] and HIV-infected patients, where the number of pDCs (partially) predicts the clinical

outcome [173–175]. Therefore, adoptive transfer of pDCs (and cDCs) can be used to reach the required level of pDCs and the subsequent initiation of the type I IFN response. Moreover, this will increase the efficacy of TLR ligands as they require their appropriate receptors that are predominantly present on DCs. As shown by Wang *et al.*, adoptive transfer of pDCs was used to successfully activate the antiviral response and limit RSV replication [176]. Thus, the administration of (stimulated) pDCs (in concert with cDCs) to restore DC function and/or numbers can activate the innate immune system to reach the required level of immune activation to control the viral infection, but this is probably dependent on the individual, the type of viral infection (chronic) and the stage of infection.

Other options

As observed in many viral infections, the (concomitant) proinflammatory response can contribute significantly to the disease. Therefore, anti-inflammatory drugs [177] are attractive to suppress symptoms during viral disease. Also, the use of antiviral drugs for specific inhibition of viral replication remains attractive as therapy, especially in combination with other treatments (like IFN α treatment and ribavirin in HCV patients). Furthermore, although TLR ligands and IFNs can induce production of restriction factors, these might also be applied directly to limit viral replication. On the other hand, IFN-inhibitor proteins of viruses can be targeted to restore immune functions [178] and make additional restriction factors or immunotherapy more effective.

Taken together, although viruses are well able to subvert or manipulate the type I IFN response, the IFN system can also be used or stimulated to strengthen the response towards viral infections. IFNs themselves are already used in HCV treatment, and promising effects have been shown in respiratory viral infections. Moreover, the therapeutic use of TLR ligands is currently under intense investigation as they have shown to have great potency to stimulate those immune cells critically involved in the antiviral immune response. This stimulates the production of antiviral proteins or inhibitors of viral evasion proteins, which can also be used independently of TLR stimulation or IFN application. Finally, the transfer of (stimulated) pDCs for gradual production of type I IFN and other cytokines (in combination with cDCs for

induction of adaptive immunity) might be an option to limit symptoms or even control virus replication.

THREATS

In the previous section, we revealed among others the opportunities related to TLR ligands as potential antiviral drugs. Yet, although promising results with TLR ligands have been reported during the last decade, there are also several **threats**.

Autoimmunity

Endosomal TLRs usually only respond to DNA/RNA derived from pathogens while immune responses to host genetic material are prevented in different ways. First, DNA (and RNA) from apoptotic or necrotic host cells is removed by DNases (and RNases, respectively). Second, the nucleic acids from microbes are fundamentally different from host nucleic acids. Viral and bacterial DNA contain unmethylated CpG motifs, whereas in host DNA heavy methylation and fewer CpG motifs are common [179]. Furthermore, the TLRs that bind (microbial) nucleic acids are endosomally located [7,14]. Because of this intracellular localization, self-nucleic acids cannot stimulate these TLRs. Finally, regulatory receptors are present on pDCs, which limit type I IFN responses [180].

Sometimes, however, these barriers are not sufficient, and aberrant immune responses arise ultimately resulting in autoimmune diseases like systemic lupus erythematosus (SLE) [181–183], an autoimmune disorder that especially affects the skin. In SLE, it is assumed that apoptotic or necrotic material containing nucleic acids are phagocytosed by pDCs and cDCs. The pDCs respond with production of type I IFN and other cytokines resulting in activation of the cDCs, which then stimulate autoreactive T and B cells. After differentiation of B cells into plasma cells, autoantibodies are produced and complex with the nucleic acids from necrotic cells. Subsequent binding to the Fc receptor for IgG (Fc γ RIIIa) on pDCs [184] and cDCs results in further type I IFN production and B-cell stimulation [185]. This vicious cycle can be evoked or aggravated by the administration of TLR ligands. The reason why these pDCs respond to the host-derived nucleic acids is still unclear.

Thus, concerns about instigating or enhancing autoimmune diseases are important reason why TLR ligands are not extensively administered in

the clinic. Despite promising results in the last decade with these ligands in antiviral therapy, precautionary measures to prevent induced autoimmune responses are definitely necessary.

Species differences

Much of what we know comes from animal experiments, but translating experimental results from laboratory animals to humans is often problematic. This is also the case with the translation of our knowledge from the immune response of well-studied mouse models to humans. For example, the response to certain viruses can be entirely different in both hosts, due to adaptation of the virus to its host [186]. Moreover, important differences in antiviral mechanisms between mice and humans have been observed.

First, there are differences in the TLR-induced response. Studies indicate that murine pDCs are able to produce IL-12p70 in addition to IFN α post-TLR9 stimulation, whereas human pDCs do not [60,62,63]. Secondly, the location of TLR9 is different in mice than in humans. In humans, TLR9 is exclusively expressed in pDCs and B cells [187] while mice express TLR9 on cDCs, B-cells, macrophages, and monocytes [188]. Thus, a TLR9 ligand can induce entirely different responses in both species. Another major difference is the function of TLR8. TLR8 stimulation induces IL-12 production in humans [189], but this receptor appears to be non-functional in mice, although this is still a matter of debate [190]. Finally, the cytokine flt-3 ligand is used to differentiate murine hematopoietic stem cells into DCs with a relatively high percentage of pDCs [191,192]. This does not reflect the human situation in which most experiments are performed with PBMCs, containing a very low number of pDCs [154–156,193] that are probably at a different stage of maturation.

Hence, as stimulation of the type I IFN response can improve immunity toward viral infection, it can also evoke or aggravate aberrant immune responses (autoimmunity), thereby limiting clinical application of TLR ligands and IFNs. Furthermore, although animal experiments have been extremely helpful in deciphering antiviral responses, these are not an exact representation of the human type I IFN response, further hindering clinical application.

CONCLUSION

In this review, we provided a condensed overview of the molecular pathways involved in the most potent antiviral part of the innate immune system, the type I IFN response. Moreover, we reviewed the cells and receptors that are intimately involved in this type I IFN system. Also, we evaluated the (im)possibilities of new ways to modulate the type I IFN response, for example, by TLR ligands or adoptive DC transfer, as promising future antiviral therapies. Nonetheless, although strong antiviral effects of IFNs, TLR ligands, DCs, and restriction factors have been shown by many studies, the clinical application of these immune-based therapies is unfortunately still limited, which might be related to concern for eventual undesired side effects like autoimmune diseases. Therefore, to be clinically successful, perhaps a more personalized approach is required. The application of these immune-based therapies can then be considered based on the individual, virus, stage of infection, and symptoms, thereby fine-tuning the type I IFN response and preventing side effects as much as possible.

CONFLICT OF INTEREST

The authors have no competing interest.

REFERENCES

1. Barre-Sinoussi F. HIV: a discovery opening the road to novel scientific knowledge and global health improvement. *Virology* 2010; **397**(2): 255–259. S00426-822(09)00531-5 [pii]10.1016/j.virol.2009.08.033
2. Kilmarx PH. Global epidemiology of HIV. *Current Opinion in HIV and AIDS* 2009; **4**(4): 240–246. 10.1097/COH.0b013e32832c06db0122929-200907000-00004 [pii]
3. Smith GJ, Vijaykrishna D, Bahl J, *et al.* Origins and evolutionary genomics of the 2009 swine-origin H1N1 influenza A epidemic. *Nature* 2009; **459**(7250): 1122–1125. nature08182 [pii] 10.1038/nature08182
4. Neumann G, Noda T, Kawaoka Y. Emergence and pandemic potential of swine-origin H1N1 influenza virus. *Nature* 2009; **459**(7249): 931–939. nature08157 [pii] 10.1038/nature08157
5. Osterhaus AD. New respiratory viruses of humans. *The Pediatric Infectious Disease Journal* 2008; **27**(10 Suppl): S71–S74. 10.1097/INF.0b013e3181684d7c 00006454-200810001-00007 [pii]
6. Tregoning JS, Schwarze J. Respiratory viral infections in infants: causes, clinical symptoms, virology, and immunology. *Clinical Microbiology Reviews* 2010; **23**(1): 74–98. 23/1/74 [pii] 10.1128/CMR.00032-09

7. Blasius AL, Beutler B. Intracellular toll-like receptors. *Immunity* 2010; **32**(3): 305–315. S1074-7613(10)00095-6 [pii] 10.1016/j.immuni.2010.03.012
8. Aderem A, Ulevitch RJ. Toll-like receptors in the induction of the innate immune response. *Nature* 2000; **406**(6797): 782–787. 10.1038/35021228
9. Akira S, Takeda K, Kaisho T. Toll-like receptors: critical proteins linking innate and acquired immunity. *Nature Immunology* 2001; **2**(8): 675–680. 10.1038/90609 90609 [pii]
10. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nature Immunology* 2010; **11**(5): 373–384. ni.1863 [pii] 10.1038/ni.1863
11. Ito T, Wang YH, Liu YJ. Plasmacytoid dendritic cell precursors/type I interferon-producing cells sense viral infection by Toll-like receptor (TLR) 7 and TLR9. *Springer Seminars in Immunopathology* 2005; **26**(3): 221–229.
12. Trinchieri G, Sher A. Cooperation of Toll-like receptor signals in innate immune defence. *Nature Reviews Immunology* 2007; **7**(3): 179–190. nri2038 [pii] 10.1038/nri2038
13. Cao W, Liu YJ. Innate immune functions of plasmacytoid dendritic cells. *Current Opinion in Immunology* 2007; **19**(1): 24–30. S0952-7915(06)00236-6 [pii] 10.1016/j.coi.2006.11.004
14. Barton GM, Kagan JC, Medzhitov R. Intracellular localization of Toll-like receptor 9 prevents recognition of self DNA but facilitates access to viral DNA. *Nature Immunology* 2006; **7**(1): 49–56. ni1280 [pii] 10.1038/ni1280
15. Lund JM, Alexopoulou L, Sato A, et al. Recognition of single-stranded RNA viruses by Toll-like receptor 7. *Proceedings of the National Academy of Sciences of the United States of America* 2004; **101**(15): 5598–5603. 10.1073/pnas.0400937101 0400937101 [pii]
16. Diebold SS, Kaisho T, Hemmi H, Akira S, Reis e Sousa C. Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science* 2004; **303**(5663): 1529–1531. 10.1126/science.1093616 1093616 [pii]
17. Krug A, French AR, Barchet W, et al. TLR9-dependent recognition of MCMV by IPC and DC generates coordinated cytokine responses that activate antiviral NK cell function. *Immunity* 2004; **21**(1): 107–119.
18. Lund J, Sato A, Akira S, Medzhitov R, Iwasaki A. Toll-like receptor 9-mediated-recognition of herpes simplex virus-2 by plasmacytoid dendritic cells. *The Journal of Experimental Medicine* 2003; **198**(3): 513–520. 10.1084/jem.20030162 jem.20030162 [pii]
19. Weber F, Wagner V, Rasmussen SB, Hartmann R, Paludan SR. Double-stranded RNA is produced by positive-strand RNA viruses and DNA viruses but not in detectable amounts by negative-strand RNA viruses. *Journal of Virology* 2006; **80**(10): 5059–5064. 80/10/5059 [pii] 10.1128/JVI.80.10.5059-5064.2006
20. Kurt-Jones EA, Popova L, Kwinn L, et al. Pattern recognition receptors TLR4 and CD14 mediate response to respiratory syncytial virus. *Nature Immunology* 2000; **1**(5): 398–401. 10.1038/80833
21. Sato A, Linehan MM, Iwasaki A. Dual recognition of herpes simplex viruses by TLR2 and TLR9 in dendritic cells. *Proceedings of the National Academy of Sciences of the United States of America* 2006; **103**(46): 17343–17348. 0605102103 [pii] 10.1073/pnas.0605102103
22. Barbalat R, Lau L, Locksley RM, Barton GM. Toll-like receptor 2 on inflammatory monocytes induces type I interferon in response to viral but not bacterial ligands. *Nature Immunology* 2009; **10**(11): 1200–1207. ni.1792 [pii] 10.1038/ni.1792
23. Bochud PY, Magaret AS, Koelle DM, Aderem A, Wald A. Polymorphisms in TLR2 are associated with increased viral shedding and lesion rate in patients with genital herpes simplex virus Type 2 infection. *Journal of Infectious Diseases* 2007; **196**(4): 505–509. JID38155 [pii] 10.1086/519693
24. Wang JP, Kurt-Jones EA, Shin OS, Manchak MD, Levin MJ, Finberg RW. Varicella-zoster virus activates inflammatory cytokines in human monocytes and macrophages via Toll-like receptor 2. *Journal of Virology* 2005; **79**(20): 12658–12666. 79/20/12658 [pii] 10.1128/JVI.79.20.12658-12666.2005
25. Diebold SS. Recognition of viral single-stranded RNA by Toll-like receptors. *Advanced Drug Delivery Reviews* 2008; **60**(7): 813–823.
26. Kato H, Takeuchi O, Sato S, et al. Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature* 2006; **441**(7089): 101–105. nature04734 [pii] 10.1038/nature04734
27. Vandevenne P, Sadzot-Delvaux C, Piette J. Innate immune response and viral interference strategies developed by human herpesviruses. *Biochemical Pharmacology* 2010; **80**(12): 1955–1972. S0006-2952(10)00487-9 [pii] 10.1016/j.bcp.2010.07.001
28. Wilkins C, Gale M Jr. Recognition of viruses by cytoplasmic sensors. *Current Opinion in Immunology* 2010; **22**(1): 41–47. S0952-7915(09)00226-X [pii] 10.1016/j.coi.2009.12.003
29. Loo YM, Fornek J, Crochet N, et al. Distinct RIG-I and MDA5 signaling by RNA viruses in innate immunity. *Journal of Virology* 2008; **82**(1): 335–345. JVI.01080-07 [pii] 10.1128/JVI.01080-07
30. Saito T, Gale M Jr. Differential recognition of double-stranded RNA by RIG-I-like receptors in antiviral immunity. *The Journal of Experimental Medicine* 2008; **205**(7): 1523–1527. 10.1084/jem.20081210 jem.20081210 [pii]
31. Kato H, Takeuchi O, Mikamo-Satoh E, et al. Length-dependent recognition of double-stranded ribonucleic acids by retinoic acid-inducible gene-I and melanoma differentiation-associated gene 5. *The Journal of Experimental Medicine* 2008; **205**(7): 1601–1610. 10.1084/jem.20080091 jem.20080091 [pii]
32. Takaoka A, Wang Z, Choi MK, et al. DAI (DLM-1/ZBP1) is a cytosolic DNA sensor and an activator of innate immune response. *Nature* 2007; **448**(7152): 501–505. nature06013 [pii] 10.1038/nature06013
33. Hornung V, Latz E. Intracellular DNA recognition. *Nature Reviews Immunology* 2010; **10**(2): 123–130. nri2690 [pii] 10.1038/nri2690
34. van Vliet SJ, Garcia-Vallejo JJ, van Kooyk Y. Dendritic cells and C-type lectin receptors: coupling innate to adaptive immune responses. *Immunology and Cell Biology* 2008; **86**(7): 580–587. icb200855 [pii] 10.1038/icb.2008.55
35. Diebold SS. Activation of dendritic cells by toll-like receptors and C-type lectins.

- Handbook of Experimental Pharmacology* 2009; **188**: 3–30. 10.1007/978-3-540-71029-5_1
36. Geijtenbeek TB, Gringhuis SI. Signalling through C-type lectin receptors: shaping immune responses. *Nature Reviews Immunology* 2009; **9**(7): 465–479. nri2569 [pii] 10.1038/nri2569
 37. Ichinohe T, Lee HK, Ogura Y, Flavell R, Iwasaki A. Inflammasome recognition of influenza virus is essential for adaptive immune responses. *The Journal of Experimental Medicine* 2009; **206**(1): 79–87. jem.20081667 [pii] 10.1084/jem.20081667
 38. Thomas PG, Dash P, Aldridge JR Jr, *et al.* The intracellular sensor NLRP3 mediates key innate and healing responses to influenza A virus via the regulation of caspase-1. *Immunity* 2009; **30**(4): 566–575. S1074-7613(09)00140-X [pii] 10.1016/j.immuni.2009.02.006
 39. Allen IC, Scull MA, Moore CB, *et al.* The NLRP3 inflammasome mediates in vivo innate immunity to influenza A virus through recognition of viral RNA. *Immunity* 2009; **30**(4): 556–565. S1074-7613(09)00139-3 [pii] 10.1016/j.immuni.2009.02.005
 40. Muruve DA, Petrilli V, Zais AK, *et al.* The inflammasome recognizes cytosolic microbial and host DNA and triggers an innate immune response. *Nature* 2008; **452** (7183):103–107. nature06664 [pii] 10.1038/nature06664
 41. Delaloye J, Delaloye J, Roger T, Steiner-Tardivel QG, *et al.* Innate immune sensing of modified vaccinia virus Ankara (MVA) is mediated by TLR2-TLR6, MDA-5 and the NALP3 inflammasome. *PLoS Pathogens* 2009; **5**(6): e1000480. 10.1371/journal.ppat.1000480
 42. Bryant C, Fitzgerald KA. Molecular mechanisms involved in inflammasome activation. *Trends in Cell Biology* 2009; **19**(9): 455–464. S0962-8924(09)00138-X [pii] 10.1016/j.tcb.2009.06.002
 43. Rathinam VA, Jiang Z, Waggoner SN, *et al.* The AIM2 inflammasome is essential for host defense against cytosolic bacteria and DNA viruses. *Nature Immunology* 2010; **11**(5): 395–402. ni.1864 [pii] 10.1038/ni.1864
 44. Paludan SR, Bowie AG, Horan KA, Fitzgerald KA. Recognition of herpesviruses by the innate immune system. *Nature Reviews Immunology* 2011; **11**(2): 143–154. nri2937 [pii] 10.1038/nri2937
 45. Pichlmair A, Reis e Sousa C. Innate recognition of viruses. *Immunity* 2007; **27**(3): 370–383. S1074-7613(07)00418-9 [pii] 10.1016/j.immuni.2007.08.012
 46. Yeow WS, Au WC, Juang YT, *et al.* Reconstitution of virus-mediated expression of interferon alpha genes in human fibroblast cells by ectopic interferon regulatory factor-7. *Journal of Biological Chemistry* 2000; **275**(9): 6313–6320.
 47. Liu YJ. IPC: professional type 1 interferon-producing cells and plasmacytoid dendritic cell precursors. *Annual Review of Immunology* 2005; **23**: 275–306.
 48. Fitzgerald-Bocarsly P, Feng D. The role of type I interferon production by dendritic cells in host defense. *Biochimie* 2007; **89**(6–7): 843–855.
 49. Lee HK, Lund JM, Ramanathan B, Mizushima N, Iwasaki A. Autophagy-dependent viral recognition by plasmacytoid dendritic cells. *Science* 2007; **315** (5817): 1398–1401. 1136880 [pii] 10.1126/science.1136880
 50. Ewald SE, Lee BL, Lau L, *et al.* The ectodomain of Toll-like receptor 9 is cleaved to generate a functional receptor. *Nature* 2008; **456** (7222): 658–662. nature07405 [pii] 10.1038/nature07405
 51. Park B, Brinkmann MM, Spooner E, Lee CC, Kim YM, Ploegh HL. Proteolytic cleavage in an endolysosomal compartment is required for activation of Toll-like receptor 9. *Nature Immunology* 2008; **9**(12): 1407–1414. ni.1669 [pii] 10.1038/ni.1669
 52. Kawai T, Akira S. Toll-like receptor and RIG-I-like receptor signaling. *Annals of the New York Academy of Sciences* 2008; **1143**: 1–20. NYAS1143020 [pii] 10.1196/annals.1443.020
 53. Kato H, Sato S, Yoneyama M, *et al.* Cell type-specific involvement of RIG-I in antiviral response. *Immunity* 2005; **23**(1): 19–28. S1074-7613(05)00142-1 [pii] 10.1016/j.immuni.2005.04.010
 54. Wang J, Basagoudanavar SH, Wang X, *et al.* NF-kappa B RelA subunit is crucial for early IFN-beta expression and resistance to RNA virus replication. *Journal of Immunology* 2010; **185**(3): 1720–1729. jimmunol.1000114 [pii] 10.4049/jimmunol.1000114
 55. Sato M, Suemori H, Hata N, *et al.* Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN-alpha/beta gene induction. *Immunity* 2000; **13**(4): 539–548. S1074-7613(00)00053-4 [pii]
 56. Taniguchi T, Takaoka A. The interferon-alpha/beta system in antiviral responses: a multimodal machinery of gene regulation by the IRF family of transcription factors. *Current Opinion in Immunology* 2002; **14**(1): 111–116. S0952791501003053 [pii]
 57. Kerkmann M, Rothenfusser S, Hornung V, *et al.* Activation with CpG-A and CpG-B oligonucleotides reveals two distinct regulatory pathways of type I IFN synthesis in human plasmacytoid dendritic cells. *Journal of Immunology* 2003; **170**(9): 4465–4474.
 58. Prakash A, Smith E, Lee CK, Levy DE. Tissue-specific positive feedback requirements for production of type I interferon following virus infection. *Journal of Biological Chemistry* 2005; **280**(19): 18651–18657. M501289200 [pii] 10.1074/jbc.M501289200
 59. Izaguirre A, Barnes BJ, Amrute S, *et al.* Comparative analysis of IRF and IFN-alpha expression in human plasmacytoid and monocyte-derived dendritic cells. *Journal of Leukocyte Biology* 2003; **74**(6): 1125–1138. 10.1189/jlb.0603255 jlb.0603255 [pii]
 60. Ito T, Kanzler H, Duramad O, Cao W, Liu YJ. Specialization, kinetics, and repertoire of type 1 interferon responses by human plasmacytoid predendritic cells. *Blood* 2006; **107**(6): 2423–2431. 2005-07-2709 [pii] 10.1182/blood-2005-07-2709
 61. Siegal FP, Kadowaki N, Shodell M, *et al.* The nature of the principal type 1 interferon-producing cells in human blood. *Science* 1999; **284**(5421): 1835–1837.
 62. Colonna M, Trinchieri G, Liu YJ. Plasmacytoid dendritic cells in immunity. *Nature Immunology* 2004; **5**(12): 1219–1226. ni1141 [pii] 10.1038/ni1141
 63. Ito T, Amakawa R, Kaisho T, *et al.* Interferon-alpha and interleukin-12 are induced differentially by Toll-like receptor 7 ligands in human blood dendritic cell subsets. *The Journal of Experimental Medicine* 2002; **195**(11): 1507–1512.
 64. Gordon KB, Gorski KS, Gibson SJ, *et al.* Synthetic TLR agonists reveal functional differences between human TLR7 and

- TLR8. *Journal of Immunology* 2005; **174**(3): 1259–1268. 174/3/1259 [pii]
65. Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nature Reviews Immunology* 2003; **3**(2): 133–146. 10.1038/nri1001 nri1001 [pii]
66. Kramer M, Schulte BM, Eleveld-Trancikova D, et al. Cross-talk between human dendritic cell subsets influences expression of RNA sensors and inhibits picornavirus infection. *Journal of Innate Immunity* 2010; **2**(4): 360–370. 000300568 [pii] 10.1159/000300568
67. Der SD, Zhou A, Williams BR, Silverman RH. Identification of genes differentially regulated by interferon alpha, beta, or gamma using oligonucleotide arrays. *Proceedings of the National Academy of Sciences of the United States of America* 1998; **95**(26): 15623–15628.
68. O'Garra A, Trinchieri G. Are dendritic cells afraid of commitment? *Nature Immunology* 2004; **5**(12): 1206–1208. ni1204-1206 [pii] 10.1038/ni1204-1206
69. Sadler AJ, Williams BR. Interferon-inducible antiviral effectors. *Nature Reviews Immunology* 2008; **8**(7): 559–568. nri2314 [pii] 10.1038/nri2314
70. Lu G, Reinert JT, Pitha-Rowe I, et al. ISG15 enhances the innate antiviral response by inhibition of IRF-3 degradation. *Cellular and Molecular Biology (Noisy-le-Grand, France)* 2006; **52**(1): 29–41. 29 [pii]
71. Takeuchi T, Kobayashi T, Tamura S, Yokosawa H. Negative regulation of protein phosphatase 2Cbeta by ISG15 conjugation. *FEBS Letters* 2006; **580**(18): 4521–4526. S0014-5793(06)00871-4 [pii] 10.1016/j.febslet.2006.07.032
72. D'Cunha J, Ramanujam S, Wagner RJ, Witt PL, Knight E Jr. Borden, EC. In vitro and in vivo secretion of human ISG15, an IFN-induced immunomodulatory cytokine. *Journal of Immunology* 1996; **157**(9): 4100–4108.
73. Kochs G, Haller O. Interferon-induced human MxA GTPase blocks nuclear import of Thogoto virus nucleocapsids. *Proceedings of the National Academy of Sciences of the United States of America* 1999; **96**(5): 2082–2086.
74. Accola MA, Huang B, Al Masri A, McNiven MA. The antiviral dynamin family member, MxA, tubulates lipids and localizes to the smooth endoplasmic reticulum. *Journal of Biological Chemistry* 2002; **277**(24): 21829–21835. 10.1074/jbc.M201641200 M201641200 [pii]
75. Kochs G, Janzen C, Hohenberg H, Haller O. Antivirally active MxA protein sequesters La Crosse virus nucleocapsid protein into perinuclear complexes. *Proceedings of the National Academy of Sciences of the United States of America* 2002; **99**(5): 3153–3158. 10.1073/pnas.052430399 99/5/3153 [pii]
76. Nakanishi M, Goto Y, Kitade Y. 2-5A induces a conformational change in the ankyrin-repeat domain of RNase L. *Proteins* 2005; **60**(1): 131–138. 10.1002/prot.20474
77. Dey M, Cao C, Dar AC, et al. Mechanistic link between PKR dimerization, autophosphorylation, and eIF2alpha substrate recognition. *Cell* 2005; **122**(6): 901–913. S0092-8674(05)00693-8 [pii] 10.1016/j.cell.2005.06.041
78. Sakuma T, Noda T, Urata S, Kawaoka Y, Yasuda J, et al. Inhibition of Lassa and Marburg virus production by tetherin. *Journal of Virology* 2009; **83**(5): 2382–2385. JVI.01607-08 [pii] 10.1128/JVI.01607-08
79. Tokarev A, Skasko M, Fitzpatrick K, Guatelli J. Antiviral activity of the interferon-induced cellular protein BST-2/tetherin. *AIDS Research and Human Retroviruses* 2009; **25**(12): 1197–1210. 10.1089/aid.2009.0253
80. Jouvenet N, Neil SJ, Zhadina M, et al. Broad-spectrum inhibition of retroviral and filoviral particle release by tetherin. *Journal of Virology* 2009; **83**(4): 1837–1844. JVI.02211-08 [pii] 10.1128/JVI.02211-08
81. Peng G, Lei KJ, Jin W, Greenwell-Wild T, Wahl SM. Induction of APOBEC3 family proteins, a defensive maneuver underlying interferon-induced anti-HIV-1 activity. *The Journal of Experimental Medicine* 2006; **203**(1): 41–46. jem.20051512 [pii] 10.1084/jem.20051512
82. Mangeat B, Turelli P, Caron G, Friedli M, Perrin L, Trono D. Broad antiretroviral defence by human APOBEC3G through lethal editing of nascent reverse transcripts. *Nature* 2003; **424**(6944): 99–103. 10.1038/nature01709 nature01709 [pii]
83. Sastri J, Campbell EM. Recent insights into the mechanism and consequences of TRIM5alpha retroviral restriction. *AIDS Research and Human Retroviruses* 2011; **27**(3): 231–238. 10.1089/AID.2010.0367
84. Adalid-Peralta L, Godot V, Colin C, et al. Stimulation of the primary anti-HIV antibody response by IFN-alpha in patients with acute HIV-1 infection. *Journal of Leukocyte Biology* 2008; **83**(4): 1060–1067. jlb.1007675 [pii] 10.1189/jlb.1007675
85. Marrack P, Kappler J, Mitchell T. Type I interferons keep activated T cells alive. *The Journal of Experimental Medicine* 1999; **189**(3): 521–530.
86. Borden EC, Sen GC, Uze G, et al. Interferons at age 50: past, current and future impact on biomedicine. *Nature Reviews. Drug Discovery* 2007; **6**(12): 975–990. nrd2422 [pii] 10.1038/nrd2422
87. Pedersen IM, Cheng G, Wieland S, et al. Interferon modulation of cellular microRNAs as an antiviral mechanism. *Nature* 2007; **449**(7164): 919–922. nature06205 [pii] 10.1038/nature06205
88. Witwer KW, Sisk JM, Gama L, Clements JE. MicroRNA regulation of IFN-beta protein expression: rapid and sensitive modulation of the innate immune response. *Journal of Immunology* 2010; **184**(5): 2369–2376. jimmunol.0902712 [pii] 10.4049/jimmunol.0902712
89. Murakami Y, Tanaka M, Toyoda H, et al. Hepatic microRNA expression is associated with the response to interferon treatment of chronic hepatitis C. *BMC Medical Genomics* 2010; **3**(48).
90. Bowie AG, Unterholzner L. Viral evasion and subversion of pattern-recognition receptor signalling. *Nature Reviews Immunology* 2008; **8**(12): 911–922.
91. Garcia-Sastre A. Mechanisms of inhibition of the host interferon alpha/beta-mediated antiviral responses by viruses. *Microbes and Infection* 2002; **4**(6): 647–655. S1286457902015836 [pii]
92. Lu LL, Puri M, Horvath CM, Sen GC. Select paramyxoviral V proteins inhibit IRF3 activation by acting as alternative substrates for inhibitor of kappaB kinase epsilon (IKKe)/TBK1. *Journal of Biological Chemistry* 2008; **283**(21): 14269–14276. M710089200 [pii] 10.1074/jbc.M710089200
93. Brzozka K, Finke S, Conzelmann KK. Identification of the rabies virus alpha/beta interferon antagonist: phosphoprotein P interferes with phosphorylation of interferon regulatory factor 3. *Journal of Virology* 2005;

- 79(12): 7673–7681. 79/12/7673 [pii] 10.1128/JVI.79.12.7673-7681.2005
94. Alcamí A, Symons JA, Smith GL. The vaccinia virus soluble alpha/beta interferon (IFN) receptor binds to the cell surface and protects cells from the antiviral effects of IFN. *Journal of Virology* 2000; **74**(23): 11230–11239.
95. Colamonici OR, Domanski P, Sweitzer SM, Larner A, Buller RM. Vaccinia virus B18R gene encodes a type I interferon-binding protein that blocks interferon alpha transmembrane signaling. *Journal of Biological Chemistry* 1995; **270**(27): 15974–15978.
96. Clementz MA, Chen Z, Banach BS, *et al.* Deubiquitinating and interferon antagonism activities of coronavirus papain-like proteases. *Journal of Virology* 2010; **84**(9): 4619–4629. JVI.02406-09 [pii] 10.1128/JVI.02406-09
97. Yuan W, Krug RM. Influenza B virus NS1 protein inhibits conjugation of the interferon (IFN)-induced ubiquitin-like ISG15 protein. *EMBO Journal* 2001; **20**(3): 362–371. 10.1093/emboj/20.3.362
98. Garaigorta U, Chisari FV. Hepatitis C virus blocks interferon effector function by inducing protein kinase R phosphorylation. *Cell Host & Microbe* 2009; **6**(6): 513–522. S1931-3128(09)00381-3 [pii] 10.1016/j.chom.2009.11.004
99. Gale M Jr, Katze MG. Molecular mechanisms of interferon resistance mediated by viral-directed inhibition of PKR, the interferon-induced protein kinase. *Pharmacology & Therapeutics* 1998; **78**(1): 29–46. S0163-7258(97)00165-4 [pii]
100. Schmolke M, Garcia-Sastre A. Evasion of innate and adaptive immune responses by influenza A virus. *Cell Microbiol* 2010; **12**(7): 873–880. CMI1475 [pii] 10.1111/j.1462-5822.2010.01475.x
101. Hale BG, Randall RE, Ortin J, Jackson D. The multifunctional NS1 protein of influenza A viruses. *J Gen Virol* 2008; **89**(Pt 10): 2359–2376. 89/10/2359 [pii] 10.1099/vir.0.2008/004606-0
102. Seo SH, Hoffmann E, Webster RG. Lethal H5N1 influenza viruses escape host antiviral cytokine responses. *Nat Med* 2002; **8**(9): 950–954. 10.1038/nm757 nm757 [pii]
103. Cheung CY, Poon LL, Lau AS, *et al.* Induction of proinflammatory cytokines in human macrophages by influenza A (H5N1) viruses: a mechanism for the unusual severity of human disease? *Lancet* 2002; **360**(9348): 1831–1837. S0140673602117727 [pii]
104. de Jong MD, Simmons CP, Thanh TT, *et al.* Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med* 2006; **12**(10): 1203–1207. nm1477 [pii] 10.1038/nm1477
105. Chan M.C., Cheung CY, Chui WH, *et al.* Proinflammatory cytokine responses induced by influenza A (H5N1) viruses in primary human alveolar and bronchial epithelial cells. *Respir Res* 2005; **6**: 135. 1465-9921-6-135 [pii] 10.1186/1465-9921-6-135
106. Korteweg C, Gu J. Pathology, molecular biology, and pathogenesis of avian influenza A (H5N1) infection in humans. *Am J Pathol* 2008; **172**(5): 1155–1170. S0002-9440(10)61876-7 [pii] 10.2353/ajpath.2008.070791
107. Beigel JH, Farrar J, Han AM, *et al.* Avian influenza A (H5N1) infection in humans. *N Engl J Med* 2005; **353**(13): 1374–1385. 353/13/1374 [pii] 10.1056/NEJMra052211
108. Taubenberger JK. The origin and virulence of the 1918 "Spanish" influenza virus. *Proc Am Philos Soc* 2006; **150**(1): 86–112.
109. Kobasa D, Jones SM, Shinya K, *et al.* Aberrant innate immune response in lethal infection of macaques with the 1918 influenza virus. *Nature* 2007; **445**(7125): 319–323. nature05495 [pii] 10.1038/nature05495
110. Kash JC, Tumpey TM, Proll SC, *et al.* Genomic analysis of increased host immune and cell death responses induced by 1918 influenza virus. *Nature* 2006; **443**(7111): 578–581. nature05181 [pii] 10.1038/nature05181
111. Kash JC, Muhlberger E, Carter V, *et al.* Global suppression of the host antiviral response by Ebola- and Marburgviruses: increased antagonism of the type I interferon response is associated with enhanced virulence. *J Virol* 2006; **80**(6): 3009–3020. 80/6/3009 [pii] 10.1128/JVI.80.6.3009-3020.2006
112. Gupta M, Mahanty S, Ahmed R, Rollin PE. Monocyte-derived human macrophages and peripheral blood mononuclear cells infected with ebola virus secrete MIP-1alpha and TNF-alpha and inhibit poly-IC-induced IFN-alpha in vitro. *Virology* 2001; **284**(1): 20–25. 10.1006/viro.2001.0836 S0042-6822(01)90836-0 [pii]
113. Howell DM, Feldman SB, Kloser P, Fitzgerald-Bocarsly P. Decreased frequency of functional natural interferon-producing cells in peripheral blood of patients with the acquired immune deficiency syndrome. *Clin Immunol Immunopathol* 1994; **71**(2): 223–230. S0090122984710762 [pii]
114. Fitzgerald-Bocarsly P, Jacobs ES. Plasmacytoid dendritic cells in HIV infection: striking a delicate balance. *J Leukoc Biol* 2010; **87**(4): 609–620. jlb.0909635 [pii] 10.1189/jlb.0909635
115. Hong HS, Bhatnagar N, Ballmaier M, *et al.* Exogenous HIV-1 Vpr disrupts IFN-alpha response by plasmacytoid dendritic cells (pDCs) and subsequent pDC/NK interplay. *Immunol Lett* 2009; **125**(2): 100–104. S0165-2478(09)00158-8 [pii] 10.1016/j.imlet.2009.06.008
116. Van Damme N, Goff D, Katsura C, *et al.* The interferon-induced protein BST-2 restricts HIV-1 release and is down-regulated from the cell surface by the viral Vpu protein. *Cell Host Microbe* 2008; **3**(4): 245–252. S1931-3128(08)00086-3 [pii] 10.1016/j.chom.2008.03.001
117. Herbeval JP, Shearer GM. HIV-1 immunopathogenesis: how good interferon turns bad. *Clin Immunol* 2007; **123**(2): 121–128. S1521-6616(06)00905-3 [pii] 10.1016/j.clim.2006.09.016
118. Vilcek J. Boosting p53 with interferon and viruses. *Nat Immunol* 2003; **4**(9): 825–826. 10.1038/ni0903-825 ni0903-825 [pii]
119. Herbeval JP, Nilsson J, Boasso A, *et al.* Differential expression of IFN-alpha and TRAIL/DR5 in lymphoid tissue of progressor versus nonprogressor HIV-1-infected patients. *Proc Natl Acad Sci U S A* 2006; **103**(18): 7000–7005. 0600363103 [pii] 10.1073/pnas.0600363103
120. Fried MW. Side effects of therapy of hepatitis C and their management. *Hepatology* 2002; **36**(5 Suppl 1): S237–S244. S0270913902001945 [pii] 10.1053/jhep.2002.36810

121. Medina J, Garcia-Bueyl L, Moreno-Monteagudo JA, Trapero-Marugan M, Moreno-Otero R. Combined antiviral options for the treatment of chronic hepatitis C. *Antiviral Research* 2003; **60**(2): 135–143.
122. Lee J, Wu CC, Lee KJ, et al. Activation of anti-hepatitis C virus responses via Toll-like receptor 7. *Proceedings of the National Academy of Sciences of the United States of America* 2006; **103**(6): 1828–1833. 0510801103 [pii] 10.1073/pnas.0510801103
123. Baca-Regen L, Heinzinger N, Stevenson M, Gendelman HE. Alpha interferon-induced antiretroviral activities: restriction of viral nucleic acid synthesis and progeny virion production in human immunodeficiency virus type 1-infected monocytes. *Journal of Virology* 1994; **68**(11): 7559–7565.
124. Pitha PM. Multiple effects of interferon on the replication of human immunodeficiency virus type 1. *Antiviral Research* 1994; **24**(2–3): 205–219.
125. Lane HC, Davey V, Kovacs JA, et al. Interferon-alpha in patients with asymptomatic human immunodeficiency virus (HIV) infection. A randomized, placebo-controlled trial. *Annals of Internal Medicine* 1990; **112**(11): 805–811.
126. Hatzakis A, Gargalianos P, Kiosses V, et al. Low-dose IFN-alpha monotherapy in treatment-naive individuals with HIV-1 infection: evidence of potent suppression of viral replication. *Journal of Interferon & Cytokine Research* 2001; **21**(10): 861–869. 10.1089/107999001753238114
127. Rivero J, Fraga M, Cancio I, Cuervo J, Lopez-Saura P. Long-term treatment with recombinant interferon alpha-2b prolongs survival of asymptomatic HIV-infected individuals. *Biotherapy* 1997; **10**(2): 107–113.
128. Frissen PH, de Wolf F, Reiss P, et al. High-dose interferon-alpha2a exerts potent activity against human immunodeficiency virus type 1 not associated with antitumor activity in subjects with Kaposi's sarcoma. *Journal of Infectious Diseases* 1997; **176**(3): 811–814.
129. Skillman DR, Malone JL, Decker CF, et al. Phase I trial of interferon alfa-n3 in early-stage human immunodeficiency virus type 1 disease: evidence for drug safety, tolerance, and antiviral activity. *Journal of Infectious Diseases* 1996; **173**(5): 1107–1114.
130. Gao L, Yu S, Chen Q, et al. A randomized controlled trial of low-dose recombinant human interferons alpha-2b nasal spray to prevent acute viral respiratory infections in military recruits. *Vaccine* 2010; **28**(28): 4445–4451. S0264-410X(10)00462-7 [pii] 10.1016/j.vaccine.2010.03.062
131. Contoli M, Message SD, Laza-Stanca V, et al. Role of deficient type III interferon-lambda production in asthma exacerbations. *Nature Medicine* 2006; **12**(9): 1023–1026. nm1462 [pii] 10.1038/nm1462
132. Mallia P, Message SD, Gielen V, et al. Experimental rhinovirus infection as a human model of chronic obstructive pulmonary disease exacerbation. *American Journal of Respiratory and Critical Care Medicine* 2010. 201006-0833OC [pii] 10.1164/rccm.201006-0833OC
133. Stockman LJ, Bellamy R, Garner P. SARS: systematic review of treatment effects. *PLoS Medicine* 2006; **3**(9): e343. 06-PLME-RA-0015R2 [pii] 10.1371/journal.pmed.0030343
134. Zorzitto J, Galligan CL, Ueng JJ, Fish EN, et al. Characterization of the antiviral effects of interferon-alpha against a SARS-like coronavirus infection in vitro. *Cell Research* 2006; **16**(2): 220–229. 7310030 [pii] 10.1038/sj.cr.7310030
135. Loutfy MR, Blatt LM, Siminovitch KA, et al. Interferon alfacon-1 plus corticosteroids in severe acute respiratory syndrome: a preliminary study. *JAMA* 2003; **290**(24): 3222–3228. 10.1001/jama.290.24.3222 290/24/3222 [pii]
136. Thiel V, Weber F. Interferon and cytokine responses to SARS-coronavirus infection. *Cytokine & Growth Factor Reviews* 2008; **19**(2): 121–132. S1359-6101(08)00002-6 [pii] 10.1016/j.cytogfr.2008.01.001
137. Cameron MJ, Bermejo-Martin JF, Danesh A, Muller MP, Kelvin DJ, et al. Human immunopathogenesis of severe acute respiratory syndrome (SARS). *Virus Research* 2008; **133**(1): 13–19. S0168-1702(07)00054-8 [pii] 10.1016/j.virusres. 2007.02.014
138. Lau YL, Peiris JS. Pathogenesis of severe acute respiratory syndrome. *Current Opinion in Immunology* 2005; **17**(4): 404–410. S0952-7915(05)00075-0 [pii] 10.1016/j.coi.2005.05.009
139. Wong CK, Lam CW, Wu AK, et al. Plasma inflammatory cytokines and chemokines in severe acute respiratory syndrome. *Clinical and Experimental Immunology* 2004; **136**(1): 95–103. 10.1111/j.1365-2249.2004.02415.x CEI2415 [pii]
140. Jewell NA, Cline T, Mertz SE, et al. Lambda interferon is the predominant interferon induced by influenza A virus infection in vivo. *Journal of Virology* 2010; **84**(21): 11515–11522. JVI.01703-09 [pii] 10.1128/JVI.01703-09
141. Khaitov MR, Laza-Stanca V, Edwards MR, et al. Respiratory virus induction of alpha-beta- and lambda-interferons in bronchial epithelial cells and peripheral blood mononuclear cells. *Allergy* 2009; **64**(3): 375–386. ALL1826 [pii] 10.1111/j.1398-9995.2008.01826.x
142. Mordstein M, Neugebauer E, Ditt V, et al. Lambda interferon renders epithelial cells of the respiratory and gastrointestinal tracts resistant to viral infections. *Journal of Virology* 2010; **84**(11): 5670–5677. JVI.00272-10 [pii] 10.1128/JVI.00272-10
143. Svetlikova D, Kabat P, Ohradnova A, Pastorek J, Betakova T, et al. Influenza A virus replication is inhibited in IFN-lambda2 and IFN-lambda3 transfected or stimulated cells. *Antiviral Research* 2010; **88**(3): 329–333. S0166-3542(10)00766-7 [pii] 10.1016/j.antiviral.2010.10.005
144. Krieg AM. Therapeutic potential of Toll-like receptor 9 activation. *Nature Reviews. Drug Discovery* 2006; **5**(6): 471–484. nrd2059 [pii] 10.1038/nrd2059
145. Meyer T, Stockfleth E. Clinical investigations of Toll-like receptor agonists. *Expert Opinion on Investigational Drugs* 2008; **17**(7): 1051–1065. 10.1517/13543784.17.7.1051
146. Zhu Q, Egelston C, Gagnon S, et al. Using 3 TLR ligands as a combination adjuvant induces qualitative changes in T cell responses needed for antiviral protection in mice. *The Journal of Clinical Investigation* 2010; **120**(2): 607–616. 39293 [pii] 10.1172/JCI39293
147. Pulendran B, Ahmed R. Translating innate immunity into immunological memory: implications for vaccine development. *Cell* 2006; **124**(4): 849–863. S0092-8674(06)00194-2 [pii] 10.1016/j.cell.2006.02.019
148. Kanzler H, Barrat FJ, Hessel EM, Coffman RL. Therapeutic targeting of innate immunity with Toll-like receptor agonists and

- antagonists. *Nature Medicine* 2007; **13**(5): 552–559.
149. Khan S, Weterings JJ, Britten CM, *et al.* Chirality of TLR-2 ligand Pam3CysSK4 in fully synthetic peptide conjugates critically influences the induction of specific CD8+ T-cells. *Molecular Immunology* 2009; **46**(6): 1084–1091. S0161-5890(08)00705-0 [pii]10.1016/j.molimm.2008.10.006
150. Svensson A, Bellner L, Magnusson M, Eriksson K. Role of IFN- α / β signaling in the prevention of genital herpes virus type 2 infection. *Journal of Reproductive Immunology* 2007; **74**(1–2): 114–123.
151. Wong JP, Nagata LP, Christopher ME, Salazar AM, Dale RM. Prophylaxis of acute respiratory virus infections using nucleic acid-based drugs. *Vaccine* 2005; **23** (17–18): 2266–2268. S0264-410X(05)00040-X [pii]10.1016/j.vaccine.2005.01.037
152. Boivin N, Sergerie Y, Rivest S, Boivin G. Effect of pretreatment with toll-like receptor agonists in a mouse model of herpes simplex virus type 1 encephalitis. *Journal of Infectious Diseases* 2008; **198**(5): 664–672. 10.1086/590671
153. Gill N, Deacon PM, Lichty B, Mossman KL, Ashkar AA. Induction of innate immunity against herpes simplex virus type 2 infection via local delivery of Toll-like receptor ligands correlates with beta interferon production. *Journal of Virology* 2006; **80**(20): 9943–9950. 80/20/9943 [pii]10.1128/JVI.01036-06
154. Hartmann G, Battiany J, Poech H, *et al.* Rational design of new CpG oligonucleotides that combine B cell activation with high IFN- α induction in plasmacytoid dendritic cells. *European Journal of Immunology* 2003; **33**(6): 1633–1641. 10.1002/eji.200323813
155. Marshall JD, Fearon K, Abbate C, *et al.* Identification of a novel CpG DNA class and motif that optimally stimulate B cell and plasmacytoid dendritic cell functions. *Journal of Leukocyte Biology* 2003; **73**(6): 781–792.
156. Vollmer J, Weeratna R, Payette P, *et al.* Characterization of three CpG oligodeoxynucleotide classes with distinct immunostimulatory activities. *European Journal of Immunology* 2004; **34**(1): 251–262. 10.1002/eji.200324032
157. Hammerbeck DM, Bursleson GR, Schuller CJ, *et al.* Administration of a dual toll-like receptor 7 and toll-like receptor 8 agonist protects against influenza in rats. *Antiviral Research* 2007; **73**(1): 1–11.
158. Gibson SJ, Lindh JM, Riter TR, *et al.* Plasmacytoid dendritic cells produce cytokines and mature in response to the TLR7 agonists, imiquimod and resiquimod. *Cellular Immunology* 2002; **218**(1–2): 74–86.
159. Hemmi H, Kaisho T, Takeuchi O, *et al.* Small anti-viral compounds activate immune cells via the TLR7 MyD88-dependent signaling pathway. *Nature Immunology* 2002; **3**(2): 196–200.
160. Heil F, Hemmi H, Hochrein H, *et al.* Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* 2004; **303**(5663): 1526–1529. 10.1126/science.1093620 1093620 [pii]
161. McCluskie MJ, Cartier JL, Patrick AJ, *et al.* Treatment of intravaginal HSV-2 infection in mice: a comparison of CpG oligodeoxynucleotides and resiquimod (R-848). *Antiviral Research* 2006; **69**(2): 77–85. S0166-3542(05)00237-8 [pii]10.1016/j.antiviral.2005.10.007
162. Chang YC, Madkan V, Cook-Norris R, Sra K, Tyring S. Current and potential uses of imiquimod. *Southern Medical Journal* 2005; **98**(9): 914–920.
163. Horsmans Y, Berg T, Desager JP, *et al.* Isatoribine, an agonist of TLR7, reduces plasma virus concentration in chronic hepatitis C infection. *Hepatology* 2005; **42**(3): 724–731.
164. Pockros PJ, Guyader D, Patton H, *et al.* Oral resiquimod in chronic HCV infection: safety and efficacy in 2 placebo-controlled, double-blind phase IIa studies. *Journal of Hepatology* 2007; **47**(2): 174–182. S0168-8278(07)00256-5 [pii]10.1016/j.jhep.2007.02.025
165. McHutchison JG, Bacon BR, Gordon SC, *et al.* Phase 1B, randomized, double-blind, dose-escalation trial of CPG 10101 in patients with chronic hepatitis C virus. *Hepatology* 2007; **46**(5): 1341–1349. 10.1002/hep.21773
166. Gonzalez VD, Landay AL, Sandberg JK. Innate immunity and chronic immune activation in HCV/HIV-1 co-infection. *Clinical Immunology* 2010; **135**(1): 12–25. S1521-6616(09)00913-9 [pii]10.1016/j.clim.2009.12.005
167. Lore K, Betts MR, Brenchley JM, *et al.* Toll-like receptor ligands modulate dendritic cells to augment cytomegalovirus- and HIV-1-specific T cell responses. *Journal of Immunology* 2003; **171**(8): 4320–4328.
168. Soumelis V, Scott I, Gheyas F, *et al.* Depletion of circulating natural type 1 interferon-producing cells in HIV-infected AIDS patients. *Blood* 2001; **98**(4): 906–912.
169. Duan XZ, Wang M, Li HW, Zhuang H, Xu D, Wang FS. Decreased frequency and function of circulating plasmacytoid dendritic cells (pDC) in hepatitis B virus infected humans. *Journal of Clinical Immunology* 2004; **24**(6): 637–646. 10.1007/s10875-004-6249-y
170. Kanto T, Inoue M, Miyatake H, *et al.* Reduced numbers and impaired ability of myeloid and plasmacytoid dendritic cells to polarize T helper cells in chronic hepatitis C virus infection. *Journal of Infectious Diseases* 2004; **190**(11): 1919–1926. JID32860 [pii]10.1086/425425
171. Salem ML, El-Demellawy M, El-Azm AR. The potential use of Toll-like receptor agonists to restore the dysfunctional immunity induced by hepatitis C virus. *Cellular Immunology* 2010; **262**(2): 96–104. S0008-8749(10)00064-X [pii]10.1016/j.cellimm.2010.03.002
172. Averill L, Lee WM, Karandikar NJ. Differential dysfunction in dendritic cell subsets during chronic HCV infection. *Clinical Immunology* 2007; **123**(1): 40–49. S1521-6616(06)00966-1 [pii]10.1016/j.clim.2006.12.001
173. Sabado RL, O'Brien M, Subedi A, *et al.* Evidence of dysregulation of dendritic cells in primary HIV infection. *Blood* 2010; **116**(19): 3839–3852. blood-2010-03-273763 [pii]10.1182/blood-2010-03-273763
174. Donaghy H, Pozniak A, Gazzard B, *et al.* Loss of blood CD11c(+) myeloid and CD11c(–) plasmacytoid dendritic cells in patients with HIV-1 infection correlates with HIV-1 RNA virus load. *Blood* 2001; **98**(8): 2574–2576.
175. Finke JS, Shodell M, Shah K, Siegal FP, Steinman RM. Dendritic cell numbers in the blood of HIV-1 infected patients before and after changes in antiretroviral therapy. *Journal of Clinical Immunology* 2004; **24**(6): 647–652. 10.1007/s10875-004-6250-5
176. Wang H, Peters N, Schwarze J. Plasmacytoid dendritic cells limit viral replication, pulmonary inflammation, and airway hyperresponsiveness in respiratory syncytial virus infection. *Journal of Immunology* 2006; **177**(9): 6263–6270.

177. O'Neill LA. Targeting signal transduction as a strategy to treat inflammatory diseases. *Nature Reviews. Drug Discovery* 2006; **5**(7): 549–563. nrd2070 [pii]10.1038/nrd2070
178. Johnson CL, Owen DM, Gale M Jr. Functional and therapeutic analysis of hepatitis C virus NS3.4A protease control of antiviral immune defense. *Journal of Biological Chemistry* 2007; **282**(14): 10792–10803. M610361200 [pii]10.1074/jbc.M610361200
179. Krieg AM. CpG motifs in bacterial DNA and their immune effects. *Annual Review of Immunology* 2002; **20**: 709–760. 10.1146/annurev.immunol.20.100301.064842 100301.064842 [pii]
180. Gilliet M, Cao W, Liu YJ. Plasmacytoid dendritic cells: sensing nucleic acids in viral infection and autoimmune diseases. *Nature Reviews Immunology* 2008; **8**(8): 594–606.
181. Banchereau J, Pascual V. Type I interferon in systemic lupus erythematosus and other autoimmune diseases. *Immunity* 2006; **25**(3): 383–392. S1074-7613(06)00395-5 [pii]10.1016/j.immuni.2006.08.010
182. Ronnblom LE, Alm GV, Oberg KE. Possible induction of systemic lupus erythematosus by interferon-alpha treatment in a patient with a malignant carcinoid tumour. *Journal of Internal Medicine* 1990; **227**(3): 207–210.
183. Baccala R, Hoebe K, Kono DH, Beutler B, Theofilopoulos AN, et al. TLR-dependent and TLR-independent pathways of type I interferon induction in systemic autoimmunity. *Nature Medicine* 2007; **13**(5): 543–551.
184. Bave U, Magnusson M, Eloranta ML, Perers A, Alm GV, Ronnblom L. Fc gamma RIIa is expressed on natural IFN-alpha-producing cells (plasmacytoid dendritic cells) and is required for the IFN-alpha production induced by apoptotic cells combined with lupus IgG. *Journal of Immunology* 2003; **171**(6): 3296–3302.
185. Ronnblom L, Eloranta ML, Alm GV. Role of natural interferon-alpha producing cells (plasmacytoid dendritic cells) in autoimmunity. *Autoimmunity* 2003; **36**(8): 463–472.
186. Fornek JL, Korth MJ, Katze MG. Use of functional genomics to understand influenza-host interactions. *Advances in Virus Research* 2007; **70**: 81–100. S0065-3527(07)70003-9 [pii] 10.1016/S0065-3527(07)70003-9
187. Weeratna RD, Makinen SR, McCluskie MJ, Davis HL. TLR agonists as vaccine adjuvants: comparison of CpG ODN and resiquimod (R-848). *Vaccine* 2005; **23**(45): 5263–5270. S0264-410X(05)00635-3 [pii] 10.1016/j.vaccine.2005.06.024
188. Hemmi H, Takeuchi O, Kawai T, et al. A Toll-like receptor recognizes bacterial DNA. *Nature* 2000; **408**(6813): 740–745. 10.1038/35047123
189. Martinson JA, Roman-Gonzalez A, Tenorio AR, et al. Dendritic cells from HIV-1 infected individuals are less responsive to toll-like receptor (TLR) ligands. *Cellular Immunology* 2007; **250**(1–2): 75–84. S0008-8749(08)00023-3 [pii]10.1016/j.cellimm.2008.01.007
190. Gorden KK, Qiu XX, Binsfeld CC, Vasilakos JP, Alkan SS. Cutting edge: activation of murine TLR8 by a combination of imidazoquinoline immune response modifiers and polyT oligodeoxynucleotides. *Journal of Immunology* 2006; **177**(10): 6584–6587. 177/10/6584 [pii]
191. Gilliet M, Boonstra A, Patrel C, et al. The development of murine plasmacytoid dendritic cell precursors is differentially regulated by FLT3-ligand and granulocyte/macrophage colony-stimulating factor. *The Journal of Experimental Medicine* 2002; **195**(7): 953–958.
192. Brawand P, Fitzpatrick DR, Greenfield BW, Brasel K, Maliszewski CR, De Smedt T. Murine plasmacytoid pre-dendritic cells generated from Flt3 ligand-supplemented bone marrow cultures are immature APCs. *Journal of Immunology* 2002; **169**(12): 6711–6719.
193. Feldman SB, Ferraro M, Zheng HM, Patel N, Gould-Fogerite S, Fitzgerald-Bocarsly P, et al. Viral induction of low frequency interferon-alpha producing cells. *Virology* 1994; **204**(1): 1–7. S0042-6822(84)71504-2 [pii]10.1006/viro.1994.1504