

Review Article

The Use of Starter Cultures in Traditional Meat Products

Marta Laranjo,^{1,2} Miguel Elias,^{1,3} and Maria João Fraqueza⁴

¹*Instituto de Ciências Agrárias e Ambientais Mediterrânicas (ICAAM), Universidade de Évora, Pólo da Mitra, Ap. 94, 7002-554 Évora, Portugal*

²*Instituto de Investigação e Formação Avançada (IFA), Universidade de Évora, Évora, Portugal*

³*Departamento de Fitotecnia, Escola de Ciências e Tecnologia, Universidade de Évora, Pólo da Mitra, Ap. 94, 7002-554 Évora, Portugal*

⁴*CIISA, Faculty of Veterinary Medicine, University of Lisbon, Avenida da Universidade Técnica, Pólo Universitário do Alto da Ajuda, 1300-477 Lisbon, Portugal*

Correspondence should be addressed to Marta Laranjo; mlaranjo@uevora.pt

Received 3 July 2017; Revised 17 October 2017; Accepted 19 October 2017; Published 12 November 2017

Academic Editor: Maria Rosaria Corbo

Copyright © 2017 Marta Laranjo et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Starter cultures could play an essential role in the manufacture of traditional cured meat products. In order to achieve objectives related to meat products' quality and safety improvement, the selection of particular strains constituting a starter culture should be carried out in the context of its application, since its functionality will depend on the type of sausage and process conditions. Also, strain selection should comply with particular requirements to warrant safety. The aim of the current review is to update the knowledge on the use of starter cultures in traditional meat products, with focus on dry-fermented products. In this manuscript, we will try to give answers to some relevant questions: Which starter cultures are used and why? Why are LAB used? What are their role and their specific mode of action? Which other groups of microorganisms (bacteria and fungi) are used as starter cultures and how do they act? A particular revision of omics approach regarding starter cultures is made since the use of these techniques allows rapid screening of promising wild strains with desirable functional characteristics, enabling the development of starter cultures better adapted to the meat matrix.

1. Introduction

Starter cultures or starters are individual or mixed formulations of selected strains with a particular enzymatic activity that when added in a defined concentration to a substrate transform it into a food product with specific characteristics [1]. This concept applied to meat products could be described as viable microorganisms that are able to multiply themselves inside meat products, increasing their preservation, controlling their hygienic safety, and potentiating their acceptability by consumers, maintaining or improving their nutritional quality [1].

The preliminary use of starters in meat products resulted from adding a portion of the final meat products to their raw materials, meaning that part of the already fermented batch of sausage was thrown back into the new mix. This already fermented product contained the necessary microorganisms to start the fermentation of the new batch. This is known as back-slopping or back-inoculation [2].

Fermented meat products may be manufactured without the use of starter cultures, although their use can help to ensure safety, standardising product properties (including flavour and colour), and shorten the ripening period. Nevertheless, well-adapted and qualified presumption of safety (QPS) strains must be used and the establishment of the starter culture must be verified in order to guarantee the expected performance.

Probiotics are live microorganisms that confer a health benefit to the host when administered in adequate amounts [3]. Probiotics have been used in food products, food supplements, and pharmaceutical products. Due to increasing concerns over health, probiotic foods (e.g., probiotic dairy products) are now accepted in the world market. Recently, the possibility of developing probiotic meat products has been discussed [4]. By using probiotic starter microorganisms, potential health benefits can be introduced to meat products and it is already possible to produce probiotic meat products [5, 6]. Nevertheless, the potentially beneficial effects on

human health from eating a probiotic sausage still need confirmation [7, 8].

The starter groups used nowadays in meat industry are, by order of importance, lactic acid bacteria (LAB), Gram-positive catalase-positive cocci (GCC+) (mainly staphylococci), moulds, and yeasts.

Lactic acid bacteria (LAB) are a group of Gram-positive bacteria belonging to the Firmicutes. They are catalase-negative, either rod-shaped (bacilli) or spherical (cocci), characterised by an increased tolerance to acidity (low pH range), and have a low GC (guanine-cytosine) content. Although many genera of bacteria produce lactic acid as a primary or secondary end-product of fermentation, the term lactic acid bacteria (LAB) is conventionally reserved for genera in the order Lactobacillales, which includes *Aerococcus*, *Carnobacterium*, *Enterococcus*, *Lactococcus*, *Lactobacillus*, *Leuconostoc*, *Oenococcus*, *Pediococcus*, *Streptococcus*, *Tetragenococcus*, *Vagococcus*, and *Weissella* [9]. As food fermentation agents LAB are involved in making yogurt, cheese, cultured butter, sour cream, sausage, cucumber pickles, olives, and sauerkraut, some species may spoil beer, wine, and processed meats [10].

Gram-positive catalase-positive cocci (GCC+) are the second most important group of meat starters and are composed of nonpathogenic coagulase-negative staphylococci (CNS). The most important starters from this group are strains belonging to the genera *Staphylococcus* and *Kocuria* [11].

At the beginning of the ripening process, the surface mycobiota is mainly composed of yeasts; however, as a_w decreases, moulds outcompete yeasts and predominate in the final product [12]. Moulds colonise the surface of fermented meat products, in some cases conferring particular characteristics, however, in other cases being considered signs of spoilage.

Yeasts are characteristic components of the mycobiota growing on fermented sausages. Their origin is mainly related to the environment and to the meat used as raw material, since yeasts are naturally found on fresh meat. The most common genera are *Candida*, *Rhodotorula*, *Debaryomyces*, and *Trichosporon*. In fermented meats, the lactic acid produced by LAB changes the environment, favouring the development of yeasts, which use all of the nutrients and energy and grow fast [13].

Meat preservation by fermentation has been carried out for thousands of years, but the idea of starter cultures was first introduced for dry sausages in the 1940s with Patent US 2225783 A [14]. The first commercial starter culture was a strain of *Pediococcus acidilactici* that was made available in the US in 1957 [15]. In Europe, the first starter culture to be introduced was strain M53 from the genus *Kocuria*, isolated from a Finnish sausage, which was used to prevent colour and aroma defects [16].

Starter cultures play an essential role in the manufacture of fermented food products. Starters composed of LAB strains produce the lactic acid that acts on meat proteins modifying their water binding capacity, thus contributing to texture, moisture content, flavour, and aroma of the products, and definitively acts on its microbiological safety.

Additionally, microbial substances, namely, bacteriocins, produced by Gram-positive species of the LAB group, such as, for example, nisin and other lantibiotics or pediocin-like bacteriocins, have an antimicrobial role with an effect on preservation and safety.

Starter cultures have a number of advantages:

- (i) They are of known quantity and quality.
- (ii) They reduce the ripening time.
- (iii) They increase safety by outcompeting undesirable microorganisms.
- (iv) They enable the manufacture of a product of constant quality all year round in any climatic zone, as long as proper natural conditions or fermenting/drying chambers are available.

The aim of the current review is to update the knowledge on the use of starter cultures in traditional meat products, with focus on dry-fermented products.

In this manuscript, we will try to give answers to some relevant questions on this subject, through the analysis of published studies with some applied results. Which starter cultures are used and why? Why are LAB used? What is their role and their specific mode of action? Which other groups of microorganisms (bacteria and fungi) are also used as starter cultures and how do they act? What is their function? A revision related to omics methods applied to the screening of autochthonous strains with desirable functional characteristics, allowing the development of well adapted starter cultures to the meat matrix, will be done.

2. Starter Cultures in Dry-Fermented Meat Products

The first generation of meat starter cultures was generally based on microorganisms isolated from vegetable fermentation, such as *L. plantarum* and members of the genus *Pediococcus*. Then, a second generation of starter cultures comprising meat-borne strains, such as *L. sakei* and coagulase-negative staphylococci (CNS), was developed, harbouring phenotypic traits of technological relevance [17]. More recently, efforts have been dedicated to the study of the physiological and technological properties of LAB and CNS isolated from traditional fermented sausages, in order to develop functional starter cultures that enhance safety and nutritional advantages while maintaining industrial performance [5, 18].

The manufacturing of dry-fermented sausages involves spontaneous fermentation commanded by bacteria (LAB) and GCC+ and, less importantly, by fungi, namely, moulds and yeasts [19].

Most meat starter cultures commercially available are combined cultures of LAB (mainly *Lactobacillus* spp. and *Pediococcus* spp.) and GCC+ (primarily *Staphylococcus* spp. and *Kocuria* spp.). These bacteria are responsible for the microbial reactions that occur during meat fermentation, such as acidification, catalase activity, and bacteriocin production [11].

Several studies have addressed the importance of using starter cultures in traditional dry-fermented meat products

not only for safety or conformity reasons, but also for uniformity purposes [20–22].

Although most studies about the use of starter cultures are on dry-fermented sausages [23–25], a few works on other meat products, such as hams or fresh sausages, have also been reported [26].

Inoculation of starter cultures in dry-fermented meat products may occur either by incorporation as an ingredient in the meat batters or by surface inoculation.

Bacteria are usually incorporated in the meat batters at concentrations between 5 and 8 log colony forming units (cfu)/g [23]. Yeasts may be inoculated either on the surface of the sausage or in the meat batter at a concentration typically between 4 and 6 log cfu/g. Moulds are always surface-inoculated, due to their strictly aerobic character, frequently by dipping in an aqueous solution of spores at concentrations ranging from 3 to 4 log spores/cm².

2.1. Their Role in Quality Improvement of Sausages. The selection of starter cultures for quality improvement of sausages is based on technologically relevant traits. The autochthonous microbiota of sausages and other meat products, as well as the microbiota of the processing environment of the production units, may be a good starting point for the isolation of potential starters, because those strains are well adapted to the meat environment [19].

Bourdichon and coworkers [27] presented a list of microorganisms used in food fermentation in a wide range of food matrices (dairy products, meat, fish, vegetables, legumes, cereals, beverages, and vinegar).

2.1.1. Bacteria: LAB and GCC+. When selecting starter cultures for dry- and semidry-fermented sausages, LAB and CNS strains with useful metabolic activities and benefits during fermentation should be used.

(1) Lactic Acid Bacteria (LAB). Lactic acid bacteria (LAB) are Gram-positive, non-spore-forming cocci or bacilli with a low GC content [28]. They generally are nonrespiratory and lack catalase. They produce lactic acid as one of the main fermentation products of carbohydrates. They lack genuine catalase and do not possess cytochromes. All LAB grow anaerobically, but unlike most anaerobes, they grow in the presence of O₂ as “aerotolerant anaerobes” [9].

According to the current taxonomic classification, they belong to the phylum Firmicutes, class Bacilli, order Lactobacillales. Six different families include all genera, as shown in Table 1 (<http://www.uniprot.org/taxonomy/186826>).

Lactic acid bacteria are among the most important groups of microorganisms used in food fermentation. They contribute to the taste and texture of fermented products and inhibit food spoilage bacteria by producing growth-inhibiting substances and large amounts of lactic acid.

Based on sugar fermentation patterns, there are two broad metabolic categories of LAB: homofermentative and heterofermentative. The homofermentative pathway produces basically only lactic acid, whereas the heterofermentative pathway produces CO₂ and ethanol or acetate in addition to lactic

acid [9]. Homofermentative LAB include some lactobacilli and most enterococci, lactococci, pediococci, streptococci, tetragenococci, and vagicocci that ferment hexoses through glycolysis by the Embden-Meyerhof-Parnas pathway. On the other hand, heterofermentative LAB ferment pentoses mainly through the phosphoketolase pathway and include leuconostocs, some lactobacilli, oenococci, and *Weissella* species.

Relevant technological features for LAB starters include fast production of lactic acid; growth at different temperatures, salt concentrations, and pH values; gas production from carbohydrates; catalase activity and hydrolysis of hydrogen peroxide; nitrate and nitrite reduction; moderate proteolytic and lipolytic enzymatic activities; good performance in combined starters with other microbial components [29].

However, fermentation conditions must be controlled to avoid excessive pinholes, gas pockets, and off-flavours, resulting from gas production from carbohydrates [30]. Additionally, the production of hydrogen peroxide may result in undesirable oxidation, known as greening [30]. Furthermore, it must be taken into account that proteolytic and lipolytic activities should be moderate, to avoid undesirable sensory changes.

As for the role of LAB in the quality of dry-fermented meat products, LAB participate in the coagulation of muscle proteins by acidifying the batters, which results in increased slice stability, firmness, and cohesiveness of the final product [31, 32]. Besides, they contribute to the flavour of the final product through the formation of noticeable acidic and vinegary (acetic acid) tastes. Moreover, the existing acidic conditions may increase the activity of cathepsin D, which is again responsible for muscle proteolysis [33].

Several authors have reported the use of LAB starter cultures for the production of fermented sausages [34–38]. For example, Wang and coworkers reported the inoculation with *L. sakei* as beneficial for microbiological quality against the growth of foodborne pathogens, also improving sensory characteristics [34].

(2) Gram-Positive Catalase-Positive Cocci (GCC+). Gram-positive catalase-positive cocci GCC+, mainly nonpathogenic coagulase-negative staphylococci (CNS), are also important in the fermentation process of sausages, since they improve the quality of the final product, while standardising the production process. They enhance colour stability, contribute to flavour development, and reduce spoilage. The ones most frequently isolated from fermented sausages are summarised in Table 2.

The use of coagulase-negative staphylococci (CNS) as meat starter cultures contributes to an adequate colour development based on their nitrate reductase activity. On the other hand, their catalase activity reduces oxidative damage and their metabolism contributes to flavour. The flavour-generating potential of CNS is even more important when producing low-salt [47, 48] or low-fat [49, 50] sausages [51]. However, the full metabolic potential of CNS should be further explored, so that we may take advantage of more technological features of CNS [52].

TABLE 1: Families and genera of LAB.

Family	Genus	Cellular morphology	Sugar fermentation
Aerococcaceae	<i>Aerococcus</i>	Cocci-tetrads	Homofermentative
Carnobacteriaceae	<i>Carnobacterium</i>	Bacilli	Homofermentative
	<i>Enterococcus</i>	Cocci	Homofermentative
Enterococcaceae	<i>Tetragenococcus</i>	Cocci-tetrads	Homofermentative
	<i>Vagococcus</i>	Cocci	Homofermentative
Lactobacillaceae	<i>Lactobacillus</i>	Bacilli	Strain-dependent
	<i>Pediococcus</i>	Cocci-tetrads	Homofermentative
Leuconostocaceae	<i>Leuconostoc</i>	Cocci	Heterofermentative
	<i>Oenococcus</i>	Cocci	Heterofermentative
	<i>Weissella</i>	Cocci/bacilli	Heterofermentative
Streptococcaceae	<i>Lactococcus</i>	Cocci	Homofermentative
	<i>Streptococcus</i>	Cocci	Homofermentative

TABLE 2: Species of GCC+ isolated from fermented sausages and their role in the fermentation process.

Family	Genus	Species	Metabolic activities	References		
Staphylococcaceae	<i>Staphylococcus</i> (CNS)	<i>S. xylosus</i>	(i) Nitrate reductase (ii) Proteolytic (iii) Lipolytic (iv) Catalase	[39–42]		
		<i>S. carnosus</i>				
		<i>S. equorum</i>				
		<i>S. succinus</i>				
		<i>S. saprophyticus</i>				
Micrococcaceae	<i>Micrococcus</i>	<i>M. luteus</i>	(i) Nitrate reductase (ii) Antioxidative (iii) Catalase (iv) Lipolytic (v) Proteolytic	[27, 43]		
		<i>M. lylae</i>				
	<i>Kocuria</i>	<i>K. varians</i>			(i) Nitrate reductase (ii) Proteolytic (iii) Lipolytic	[42, 44–46]
		<i>K. kristinae</i>				

Besides contributing to flavour, *Staphylococcus* and *Kocuria* also provide nitrate-reductase and antioxidant activities [53, 54].

Numerous studies addressing the use of starter cultures in meat products have been published, with both single (either LAB or GCC+) and mixed cultures.

Several authors have reported the use of CNS starter cultures for the production of fermented sausages. According to Ravyts et al. [51], the success of CNS in flavour development seems to be determined by acidification.

Hugas and Monfort [31] highlighted the need to use selected strains of GCC+ to ensure sensory quality. Besides, other authors have described the capability of *S. xylosus* and *S. carnosus* strains to modulate aroma through the degradation of amino acids and free fatty acids (FFAs) [55–57].

Autochthonous strains of *S. xylosus* have been recommended for the production of very aromatic sausages in Southern Europe, instead of the less adapted commercial starter cultures [58].

Lusnic and colleagues have studied the effect of an added starter culture (*S. xylosus* and *S. carnosus*) to a

frankfurter-type meat emulsion in degrading polychlorinated biphenyls (PCBs) [59]. Furthermore, quite a few works have been published reporting the results obtained by the utilisation of mixed starter cultures (LAB and CNS) [25, 60–66].

Bacteriocinogenic LAB and selected strains of *S. xylosus* and *S. carnosus* are commercially available for use in improving the safety, colour, and flavour of final products. It is also important to assess positive interactions, such as growth and proteolytic activity, among the different starter cultures strains [67–70].

The effect of different starter culture combinations (*Staphylococcus carnosus*, *Pediococcus pentosaceus*, and *Lactobacillus sakei*) on the quality of Turkish type fermented sausage (*Sucuk*) has been evaluated during ripening and it was concluded that the use of lipolytic starter cultures (*S. carnosus*/*L. sakei*) would have a positive effect in accelerating ripening and enhancing the quality of dry-fermented sausages [71].

Tremonte and coworkers demonstrated that *S. xylosus* and *Kocuria varians* are able to stimulate the growth of *L.*

sakei strains, positively influencing the proteolytic activity of strains in a combined use [66].

Casquete and colleagues have emphasised the importance of autochthonous starter cultures in improving homogeneity and safety of fermented meat products, without depreciating their sensory characteristics [60–62]. Furthermore, they have highlighted the importance of choosing a starter formulation consisting of a combination of strains that is appropriate for each ripening procedure [60].

We may conclude that flavour and aroma of fermented sausages result from the combined action of different bacteria: LAB produce lactic acid and small amounts of acetic acid, ethanol, and acetoin; however, the proteolytic and lipolytic activities of both LAB and GCC+ are essential to the sensory quality of fermented sausages.

2.1.2. Fungi: Yeasts and Moulds. Fungi generally contribute to a characteristic flavour of some fermented meat products. Yeasts may be either inoculated in the meat batters or surface-inoculated, whereas moulds are always inoculated at the surface of sausages. Surface inoculation has a further physical protective role.

(1) Yeasts. The first studies with yeasts in fermented sausages were conducted in the first decades of the 20th century, when the importance of the “*fleur du saucisson*” was recognized and the use of pure yeast cultures for flavouring in fermented sausages began to be recommended. Later on, it was established that yeasts are part of the microbiota of fermented sausages and their use as starter cultures was suggested, because the addition of selected *Debaryomyces* strains could improve the curing, colour, and flavour of sausages [72].

Several studies have tried to understand the role of yeasts as secondary microbiota in fermented meat products. Yeast strains belonging to the genera *Debaryomyces*, *Yarrowia*, *Pichia*, *Rhodotorula*, *Cryptococcus*, and *Trichosporon* have been isolated from meat products [73], with clear predominance of the *Debaryomyces* genus [13].

Some yeasts have been shown to contribute to flavour and texture development throughout the curing of various products [74–76]. Moreover, some studies have shown that the characteristic flavour of dry-cured meat products may be developed through the influence of yeasts [77–79].

Furthermore, the manufacture of dry-fermented sausages with optimised concentrations of *Debaryomyces* spp. in the presence of LAB and CNS has been demonstrated to have a positive effect on the final flavour and sensory quality by inhibiting the development of rancidity and generating ethyl esters that contribute to the proper sausage aroma [78].

(2) Moulds. Surface moulding of fermented meat products is considered a desirable event in most European countries, which include Italy, Romania, Bulgaria, France, Hungary, Switzerland, Southern Germany, Spain, Austria, and Belgium [12]. In fact, the presence of mycelium at the surface of sausages has several main advantages:

- (i) It prevents excessive drying, allowing homogeneous dehydration of the product [12].

- (ii) It metabolizes peroxides, protecting fat from oxidation, thus preventing rancidity [12].

- (iii) It reduces O₂ levels on the product surface, thus avoiding oxidative processes and improving meat colour [80].

- (iv) It contributes to the flavour of the final product, by breaking up fats, proteins, and lactic acid, thus favouring pH increase [12].

The use of moulds as a seasoning for sausage can have both desirable and undesirable consequences. The desirable consequences are the creation of a successful product that appeals to consumers. The undesirable consequences are health risks associated with the growth of undesirable moulds that produce highly toxic secondary metabolites, mycotoxins, such as ochratoxin A (OTA), or penicillin produced by species of *Penicillium* [81].

Furthermore, surface moulding of fermented meat products was observed during storage and can be a quality problem, because of the undesirable effects, mainly connected to the production of off-flavours [81].

Surface mould inoculations were traditionally done with the autochthonous mycobiota, which was mainly composed of *Penicillium* spp., *Aspergillus* spp., or *Scopulariopsis* spp. The first toxicologically and technologically suitable mould starter culture for meat products, *P. nalgiovense* strain, was selected by Mintzlaff and Leistner in 1972 [82]. However, nowadays, a wide assortment of industrialised starter cultures is commercially available as an alternative to the inoculating mixtures composed of autochthonous strains.

Some studies on the use of mould starter cultures have already been performed [80, 83]. For example, quality traits of wild boar mould-ripened salami manufactured with different selections of meat and fat tissue and with and without commercial bacterial starter cultures have been investigated [84]. The use of a bacterial starter culture in the manufacture of mould-ripened wild boar salami resulted in significantly lower peroxide values, lower TBARS concentrations, and lower amounts of biogenic amines, namely, histamine, cadaverine, and putrescine, associated with better sensory evaluation scores.

Application of commercial moulds to sausage surfaces improves primarily the safety towards regarding mycotoxin production. Moreover, the production of antibiotics, namely, penicillin, also needs to be controlled [82]. Additionally, sausage producers achieve more consistent flavour, taste, and drying rate and a more uniform appearance.

Table 3 shows a list of moulds found in fermented meat products.

Among the species mentioned in Table 3, *P. nalgiovense* and *P. gladioli* are currently considered safe and are commercially available to be used as starter cultures in meat products [12].

2.2. Antimicrobial Activity of Starter Cultures. Bacteriocins, natural antimicrobial peptides, and the acid lactic produced from glucose could be used to improve the quality and safety of meat products by avoiding the presence of pathogens, such as *Listeria monocytogenes* and spoilage microorganisms, and

TABLE 3: Species of moulds usually found in dry-fermented sausages.

Common species		Uncommon species
<i>Penicillium nalgiovense</i>	<i>P. waksmanii</i>	<i>Mucor</i> spp.
<i>P. gladioli</i>	<i>Aspergillus ochraceus</i>	<i>Scopulariopsis</i> spp.
<i>P. camemberti</i>	<i>E. herbariorum</i>	<i>Cladosporium</i> spp.
<i>P. chrysogenum</i>	<i>E. repens</i>	<i>Eupenicillium</i> spp.
<i>P. aurantiogriseum</i>	<i>A. niveus</i>	<i>Eurotium</i> spp.
<i>P. brevicompactum</i>	<i>P. citrinum</i>	<i>Talaromyces</i> spp.
<i>P. nordicum</i>	<i>A. candidus</i>	<i>Geotrichum candidum</i>
<i>P. phoeniceum</i>	<i>P. crustosum</i>	<i>Talaromyces wortmannii</i>
<i>Eurotium rubrum</i>	<i>P. commune</i>	
<i>P. griseofulvum</i>	<i>A. sclerotiorum</i>	
<i>P. olsonii</i>	<i>A. versicolor</i>	
<i>P. implicatum</i>	<i>P. alii</i>	
<i>Scopulariopsis candida</i>	<i>P. fellutanum</i>	
<i>P. solitum</i>		

improving the competitiveness of their producers for survival [85].

A list of the main bacteriocins produced by LAB along with a list of bacteria they are effective against is summarised in Table 4.

Several *L. sakei* and *L. curvatus* have been reported as bacteriocin producers and have been used as protective cultures, and their activity against *L. monocytogenes* has been proved in meat products [87–90].

Lactococcus lactis and *Enterococcus* spp. strains isolated from different food matrices have been shown to produce bacteriocins [91–93].

Pediococcus acidilactici MCH14 pediocin-producing strain and the pediocin PA-1 itself have been demonstrated to inhibit the growth of the foodborne pathogens *L. monocytogenes* and *Clostridium perfringens* in Spanish dry-fermented sausages and frankfurters [94].

Bacteriocins produced by strains of *L. plantarum* isolated from Portuguese traditional pork products have been shown to have a broad spectrum of activity [95].

LAB starter cultures have been used in the production of *Nham*, which is a Thai-style fermented pork sausage, for their antilisterial activity in order to reduce the severity of postacidification and increase the shelf life of *Nham* at ambient temperature [96, 97].

Additionally, also *S. xylosus* strain SX S03/1 M/1/2 has been shown to produce a thermostable bacteriocin which could be used as starter culture or meat additive to prevent possible handling or meat processing contamination [98].

2.3. Competitiveness of Starter Cultures. One of the most important properties of meat starter cultures is the ability to colonize the meat environment, in competition with the autochthonous microbiota and dominating the microbial community of fermented products. The starter culture must compete with the natural microbiota of the raw material, which carries out the expected metabolic activities through its growth rate and survival under the prevailing conditions during sausage production. Low temperatures, high salt

concentrations, and, to a lesser extent, oxygen availability are among the most important preservative conditions during meat fermentation [17].

The main metabolic activities and their corresponding technological roles for the main microbial starter groups are shown in Table 5.

In general, CNS are poorly competitive in the presence of acidifying LAB strains [99]. On the other hand, strains of *L. sakei* have shown superior competitiveness, which could probably be explained by their specialised metabolic repertoire well adapted to the sausage environment, including the arginine deiminase (ADI) pathway [100] and the utilisation of nucleosides [101].

Genus-specific and species-specific PCR and real-time RT-PCR methods have been used to monitor and quantify the populations of the inoculated starter cultures [24]. Moreover, RT-PCR-DGGE and RNA-based pyrosequencing of the 16S rRNA gene have also been used to monitor the microbiota of fermented sausages [102].

2.4. Safety of Selected Meat Starter Cultures. Meat starter cultures or food cultures (FC) are safe live bacteria, yeasts, or moulds used in food production, and they are in themselves a characteristic food ingredient (<http://www.efca.org/content/food-culture>). Food starter cultures (microorganisms) used directly in food production are regarded as food ingredients in the European Union (EU). Starters enter in a category of food ingredients with a very long history of use in a great variety of food products. If a starter is added to a food product, the requirements established in the *General Food Law* should be accomplished by the food operator. The food cultures used as starters in the fermentation of foods are not subject to EU premarketing regulation, unless they are regarded as being novel to the EU market and their consumers. Many starters were selected from fermented foods and several microorganisms are present in spontaneously fermented foods. However, regarding safety concerns, any food cultures to be introduced in a food should be evaluated. The approaches for assessing the safety of microorganisms

TABLE 4: LAB bacteriocins, bacteriocin producers, and susceptible pathogenic bacteria.

Bacteriocin	Bacteriocin producer	Susceptible pathogenic bacteria
Sakacin	<i>Lactobacillus sakei</i>	<i>Listeria monocytogenes</i>
		<i>Staphylococcus aureus</i>
		<i>Enterococcus</i> spp.
		<i>Brochothrix thermosphacta</i>
		<i>Pseudomonas</i> spp.
Plantaricin	<i>L. plantarum</i>	<i>Campylobacter</i> spp.
		<i>Escherichia coli</i>
		<i>Klebsiella</i> spp.
		Other LAB
		<i>Listeria monocytogenes</i>
		<i>Staphylococcus aureus</i>
		<i>Clostridium perfringens</i>
		<i>Clostridium tyrobutyricum</i>
		<i>Bacillus cereus</i>
		<i>Enterococcus</i> spp.
Curvacin	<i>L. curvatus</i>	<i>Brochothrix thermosphacta</i>
		<i>Pseudomonas</i> spp.
		<i>Salmonella</i> spp.
		<i>Escherichia coli</i>
		Other LAB
Nisin	<i>Lactococcus lactis</i>	<i>Listeria monocytogenes</i>
		<i>Staphylococcus aureus</i>
		<i>Clostridium tyrobutyricum</i>
		Other LAB
Pediocins	<i>Pediococcus</i> spp.	<i>Listeria monocytogenes</i>
		<i>Enterococcus</i> spp.
		Other LAB

Adapted from Fraqueza et al. [86].

entering the human food chain differ considerably depending on the applicable legislation, if any.

Several approaches have been delineated in order to consider the starter cultures safe. The *Qualified Presumption of Safety* (QPS) list is the EFSA fast track risk assessment tool that is used by EFSA panels when evaluating products with microorganisms that require a premarket authorisation (e.g., feed additive cultures, cell factories producing enzymes/additives/vitamins, novel microorganisms, and plant protection). This approach is restricted only to the microorganisms related to regulated food and feed products and is based on history of use, body of knowledge, and the absence of adverse effects at the taxonomic unit level [103, 104].

The *Generally Recognized as Safe* (GRAS) status is open to all types of food additives, which include food cultures. The determination of GRAS status is made by the FDA and/or

external experts and is based on the history of use, body of knowledge, and the absence of adverse effects at the strain level.

Food cultures with a long history of safe use in food are considered as traditional food ingredients and are legally permitted for use in foods in the EU without premarket authorisation, as described earlier. As a consequence, EFSA panels do not evaluate microbial strains of food cultures. Nevertheless, the QPS list can be consulted when safety evaluations of food culture are made.

Microorganisms, which are not on the QPS list, are not necessarily considered to be unsafe and their assessment regarding antibioresistance, virulence, and biogenic amine characterization should be done.

The *International Dairy Federation* (IDF) and the *European Food and Feed Cultures Association* (EFFCA) have proposed additional tools and methods to evaluate the safety

TABLE 5: Requirements for starter LAB, GCC+, yeasts, and moulds.

Microbial group	Metabolic activity	Technological role
LAB	Acidification	Modulate flavour (acid/tangy)
		Inhibit pathogens
	Proteolysis	Develop texture
		Accelerate drying
		Develop flavour
Antimicrobial	Inhibit pathogens	
	Extend shelf life	
GCC+	Antioxidant	Protect colour
	Probiotic	Compete in the gastrointestinal tract
	Nitrate reductase	Develop typical red (cured) colour
Yeasts	Degradation of amino acids and FFAs	Develop flavour
	Antioxidant	Prevent rancidification
Lipolytic		
Moulds	Antioxidant	Prevent rancidification

Adapted from [17].

of food cultures with the unique target of keeping a high level of food safety and to protect human life and health. According to Laulund et al. [105], whatever the strategy applied, it is imperative to have an evaluation of the food cultures' safety at three levels: (a) at the strain level, (b) during production, and (c) in the process it is applied to and throughout the shelf life of the food.

2.4.1. Assessment of Antibioresistance. The *One Health* concept recognises that the health of people is connected to the health of animals and the environment. The food chain has been recognized as one of the main routes for the transmission of antibiotic-resistant bacteria between animal and human populations [106]. Antibiotic resistant bacterial strains may be a potential direct link between the indigenous microbiota of animals and the human gastrointestinal tract.

Bacterial strains selected as starters with technological or food protective characteristics to be introduced in food always need to be phenotypically assessed for antibiotic resistance to clinically relevant antibiotics. The phenotypic testing based on determination of a *minimum inhibitory concentration* (MIC) for a selected group of antimicrobials should be performed. The absence of phenotypic antibiotic resistance is preferred, but if a resistance profile is observed, a proper analysis of the whole genome potentially combined with information that the observed resistance is not transferable is needed; only then can the strain(s) be considered safe for use in food culture [107].

The possibility of antimicrobial resistance transfer from viable microorganisms to other microorganisms is related to the genetic basis of the resistance being considered most plausible, when the resistance is mediated by added/acquired genes. Regarding this possibility, several safety assessments have been done by several authors on the species usually selected for starters, such as CNS or LAB.

Safety hazards associated with CNS were mostly limited to the presence of antibiotic resistance [108]. CNS strains resistant to multiple antibiotics have been reported [109]. Kastner et al. [110] detected the tetracycline resistance genes *tetK* in *Staphylococcus* spp. starter cultures.

The detection of antibiotic resistant (AR) strains among LAB has resulted in their recognition as a reservoir of AR genes horizontally transmissible to pathogens through the food chain, which constitutes a problem [111, 112]. Antibiotic multiresistant strains of lactobacilli and other LAB have been isolated from dry-fermented meat products [113–120]. LAB possesses a broad spectrum of natural (intrinsic) and acquired antibiotic resistance. However, only resistance acquired by mutation or horizontal gene transfer poses a risk for public health [121].

The most common resistance genes detected in LAB isolated from dry-fermented sausages are the tetracycline resistance genes *tetM*, *tetW*, and *tetS* and the genes coding for erythromycin resistance, *ermB* and *ermC* [117, 120]. These are genes linked to mobile elements, and if the phenotypic expression of antibiotic resistance is expressed, their presence is considered a hazard.

2.4.2. Detection of Strains Producers of Biogenic Amines. Any strains to be incorporated as starters in fermented meat products should be assessed for their (in)ability to mediate the production of biogenic amines. Strategically, the use of *Lactobacillus* spp. or *Pediococcus* spp. non-BA producer strains could dominate and avoid the presence of high contents of BA in meat products. Several authors have reported the important role of starter cultures in decreasing the content in biogenic amines [47, 48, 122–126].

2.4.3. Toxigenic Potential. Among LAB, enterococci play an important role in food fermentation and may contribute to

the organoleptic uniqueness of some products, but they are also responsible for community-acquired and nosocomial infections [118]. Some of the most important virulence factors include the production of hydrolytic enzymes, namely, gelatinase, lipase, and DNase, haemolytic activity and the production of cytolysin, the presence of adhesins, and the ability to form biofilms [127].

Two studies with enterococci strains isolated from several Portuguese dry-fermented sausages revealed that although meat enterococci harbour antibiotic resistance and produce biofilms, a reduced number of virulence factors were detected [118, 128]. However, a third study with Portuguese dry-fermented products from northern Portugal has detected phenotypic and genotypic evidence of potential virulence factors among *Enterococcus* spp. isolates, which is a reason of concern [129].

Some members of the CNS group, primarily *S. epidermidis*, are common nosocomial pathogens, and the presence of regulatory elements involved in the control of virulence-factor synthesis has recently been identified. Remarkably, strains of *S. xylosus* were isolated from patients who had an underlying disease, while the same species has been reported to be involved in infections of poultry [130].

Although CNS of food origin have not been found to produce nosocomial infections, some strains that produce enterotoxins have been described. Vernozy-Rozand et al. [131] reported enterotoxin E to be the most common enterotoxin in *S. equorum* and *S. xylosus*, although it is reported that the occurrence of staphylococcal enterotoxin genes in CNS from slightly fermented sausages was rare, detecting only *entC* in *S. epidermidis* [132].

Absence of genes coding for staphylococcal enterotoxins or enterotoxin-like superantigens is a requirement for strains selected as starter cultures, and the *S. xylosus* and *S. carnosus* strains currently used as starter cultures or isolated from fermented meat products generally lack toxin genes [11].

The analysis of virulence factors in strains of *S. epidermidis*, *S. simulans*, *S. xylosus*, *S. kloosii*, and *S. caprae* revealed sometimes high percentage of incidence of the following virulence traits: production of slime, α -haemolysin, β -haemolysin, DNase, TNase, hyaluronidase, and TSST-1 and production of enterotoxins SEA, SEB, SEC, and SED [133].

2.4.4. Strains with Ability of Biofilm Formation. In food industry, biofilm formation is undesirable for hygienic and safety reasons, as it can allow the attachment of food-spoilage or pathogenic microorganisms to food or food surfaces [134]. Nevertheless, several authors believe that colonization of food surfaces by starters could be desirable, as it would inhibit colonization by pathogenic or spoilage bacteria [135].

Among CNS, biofilm formation has been studied in *S. aureus* [136], *S. epidermidis* [136], *S. hominis* [137], *S. sciuri* [135], and *S. equorum* [138]. *S. capitis*, *S. cohnii*, *S. epidermidis*, *S. lentus*, and *S. saprophyticus* have all also been reported to form biofilms [139], though due to different genetic determinants [140]. These studies concluded that, in general, biofilm formation is a strain-dependent characteristic. Furthermore, the capacity of *S. xylosus* to form biofilms may contribute to its survival of food processing [141]. On the other hand, the

inability of *S. carnosus* to form biofilms may explain why it is rarely recovered from meat processing environments [142].

LAB biofilms may be used to control the formation of biofilms by the foodborne pathogens *Listeria monocytogenes*, *Salmonella* Typhimurium, and *Escherichia coli* O157:H7 [143].

Genes potentially responsible for biofilm formation and cellular aggregation that may assist the organism to colonize meat surfaces have been identified in *L. sakei* strain 23K [144]. Moreover, the analysis of microenvironments through the scanning electron microscopy (SEM) evidenced the presence of microchannels that favour microbial flow, while the ability of *L. sakei* to form biofilm guarantees the correct colonisation of the different meat niches, throughout the fermentation process (2017).

Biofilm formation in LAB species has been reported to be a stress response and survival strategy in stressful environments [145, 146]. Some reports have also described the genes responsible for quorum sensing, adhesion, and biofilm formation [147–150].

Another possible biocontrol strategy to avoid the presence of pathogens in meat industry could be the use of bacteriocins and enzymes; this is considered important for the maintenance of biofilm-free systems and thus for the quality and safety of foods.

2.5. Functional Starter Cultures. Functional starter cultures are starters that have at least one functional property, which may contribute to food safety and/or offer one or more organoleptic, technological, nutritional, or health advantages [151]. They offer additional functionalities compared to plain starter cultures and are a way of improving the fermentation process of meat products and achieving tastier, safer, and healthier products.

2.5.1. Bioprotective Cultures. Biological preservation has gained increasing attention as a means of naturally controlling the shelf life and safety of foods. The use of protective starter cultures in the manufacture of fermented meat products is a well-established technology [86]. Bioprotective starters may contribute to the safety and increase in shelf life of fermented meat products through the release of organic acids [152], the production of bacteriocins against important food pathogens, mainly *L. monocytogenes* [153], and the control of biological hazards [86].

Potential protective starter cultures to use in fermented meat products have been identified [154] and tested [4, 155–157]. The use of bioprotective starter cultures ensures safety, while increasing shelf life, without compromising the nutritional value of fermented meat products or depreciating their sensory quality.

2.5.2. Probiotics. According to the currently adopted definition by the Food and Agriculture Organization/World Health Organization (FAO/WHO) [158], probiotics are defined as “live microorganisms which when administered in adequate amounts confer a health benefit on the host.”

Probiotics are nonpathogenic health-promoting microorganisms that when ingested in defined amounts may have a positive effect on human physiology and health [29]. In 1965,

Lilly and Stillwell proposed probiotics to be “microorganisms promoting the growth of other microorganisms.” To act as safe probiotic microorganisms, strains should be of species and genera normally present in the human gastrointestinal tract [159].

Probiotics are LAB (or bifidobacteria), mainly Gram-positive *Lactobacillus* species.

In general, health benefits of probiotic foods are based on the presence of selected strains of LAB that, having passed through the stomach and the small intestine, survive in the large intestine and confer a health benefit on the host [160].

LAB with probiotic properties may have a positive influence on product taste, flavour, and aroma, as well as on functional and physiological properties [8].

Some LAB strains are able to produce nutraceutical compounds [161]. Studies on *Lactococcus lactis* highlight the possibility of developing LAB meat starter cultures for in situ production of vitamins, by overexpression and/or disruption of relevant metabolic genes [162–164].

Although dairy products are the most commonly used food vehicles for the delivery of probiotics, several studies dealing with the use of probiotics in fermented meat products to improve their nutritional value as functional foods have been reported [5, 154, 165–167].

The commercial application of probiotics in meat products is not a current procedure, mostly because of technological issues. Although fermented meat products are processed without heating, probiotics may still be inactivated due to low pH or water activity value, as well as by the presence of native microorganisms or curing salts. The most important problem is to find a compromise between technology, safety, quality, and health-beneficial value of food [160]. For recent reviews, please refer to Neffe-Skocińska et al. [168] and Vuyst et al. [8].

Some species involved in sausage fermentation, such as *L. plantarum*, have been engineered to produce an excess of folate (vitamin B11) [162]. This gives the possibility of fortifying meat products with vitamins and other essential compounds, thus producing healthier meat products [29].

Today, the use of probiotic starters in any fermented food claiming health benefits should be scientifically demonstrated according to the legal requirements of EU for labelling [169, 170].

3. Omics of Meat Starter Cultures

The main bacterial species used in meat fermentation are LAB and CNS. *Lactobacillus sakei*, *Lactobacillus curvatus*, *Lactobacillus plantarum* (mainly in Europe), and *Pediococcus pentosaceus* and *Pediococcus acidilactici* (mainly in the US) are the starters commonly used for their fermentative role in dry-sausage production, while *Staphylococcus xylosum* and *Staphylococcus carnosus* are known for their involvement in the development and stability of colour and aroma production [171].

Using comparative genomics, transcriptomics, proteomics, and metabolomics, the diversity of strains naturally present in traditional fermented sausages is being explored. These approaches allow rapid high-throughput screening of promising wild strains with desirable functional

characteristics and a lack of negative features, enabling the development of starter cultures based on indigenous technological bacteria from traditional sausages, which are thus better adapted to the meat matrix [22, 172].

The first genome sequence of a starter to be published was the one of the LAB *L. sakei* 23K [144]. Despite the small sized genome (1,883 protein-coding genes), *L. sakei* contains seven rRNA gene clusters [144]. This redundancy may contribute to its ability to grow in complex microbial ecosystems [173]. With regard to gene products, the *L. sakei* genome shares the highest level of conservation with *Lactobacillus plantarum*, which can be used as a starter in fermented meat, dairy, and vegetable products [144, 174, 175]. Genome analysis revealed a specialized metabolic repertoire to adapt and grow on meat products. Important cellular functions are encoded by a redundancy of genes likely to enhance the organism's robustness and most probably help it to outgrow other competing bacteria. As a unique ability among lactic acid bacteria, *L. sakei* is able to use meat components, such as purine nucleosides, abundant in meat, upon glucose depletion, to grow and produce energy. Genes possibly responsible for biofilm formation and cellular aggregation, which may assist in colonising meat surfaces, were also identified [144].

The draft genome sequence of *L. sakei* subsp. *sakei* strain LS25, a commercial starter culture for fermented sausages, has been released [176]. Slightly larger than the one of *L. sakei* 23K, this genome has 1,972 predicted protein-coding genes and 7 rRNA operons [176]. Compared to the *L. sakei* 23K genome [144], 1,618 genes are orthologous, but 250 seem to be unique to LS25, including a set of genes for carbohydrate metabolism, various transporters, and dehydrogenases/oxidoreductases [176].

Complete or draft genome sequences of *Pediococcus pentosaceus* and *Pediococcus acidilactici* strains, from diverse Korean fermented food products, have been released, but none isolated from meat products [177–179].

Genomes of several strains of starter CNS, have also been published, namely, *S. xylosum* SMQ-121 [180], *S. xylosum* S04002 [181], and *S. carnosus* TM300 [182].

The draft genome sequence of *S. xylosum* SMQ-121 revealed the absence of genes coding for toxins or virulence factors. Furthermore, only four antibiotic resistance genes were found: two genes encode proteins that belong to the major facilitator superfamilies involved in phenicol and fluoroquinolone resistance; another gene encodes a putative aminoglycoside 3'-phosphotransferase for resistance to aminoglycosides; and the last one encodes trimethoprim resistance. Nevertheless, this strain was found to be sensitive to amikacin, chloramphenicol, ciprofloxacin, and trimethoprim [180].

A genome comparison of several *S. xylosum* meat starter cultures, including strain S04002, with other *S. xylosum* strains causing cow and goat mastitis, among others, has shown the presence of aroma compounds in *S. xylosum* S04002 [181].

S. carnosus TM300 genome has the highest GC content of all sequenced staphylococcal genomes [182]. It contains only one prophage and one genomic island characterised by a mosaic structure composed of species-specific genes. All starter cultures features, such as nitrate/nitrite reduction,

several sugar degradation pathways, two catalases, and nine osmoprotection systems, are present. It lacks most virulence factors, namely, the typical *S. aureus* toxins, as well as biofilm formation genes, highlighting its nonpathogenic status [182].

Following the publication of the genome sequences of several strains, global approaches based on transcriptomics and proteomics have been developed in order to better understand the adaptation of starters to the meat environment and their interactions with the ecosystem and the meat substrate.

Genes involved in safety and technologically relevant properties of food associated CNS, such as antibiotic resistance, haemolysins, toxins, amino acid decarboxylases, binding proteins to extracellular matrix (ECM), lipases, proteases, stress response factors, and nitrate dissimilation, have been detected using DNA microarrays [183].

S. xyloso C2a strain response to nitrosative [184] or nutrients and osmotic stress [185] has been investigated through DNA microarrays. *S. xyloso* has been shown to counteract nitrosative stress by developing several oxidative stress resistance mechanisms, such as modulation of the expression of genes involved in iron homeostasis, detoxifying enzymes, and DNA and protein repairs [184]. *S. xyloso* adapted its metabolism to the meat nutrients and anaerobic conditions by simultaneously using glucose and lactate as carbon sources and by using meat peptides and amino acids. *S. xyloso* responded to the osmotic stress caused by the addition of salt (NaCl) by overexpressing genes involved in transport and synthesis of osmoprotectants, particularly glycine betaine, and Na⁺ and H⁺ extrusion [185]. To overcome the damaging effects of oxidative and nitrosative stress, staphylococci have developed protection, detoxification, and repair mechanisms controlled by a network of regulators [186].

Among the overexpressed proteins in *S. xyloso* biofilm, several related to exopolysaccharide biosynthesis were reported [187]. Furthermore, with overexpression of some proteins involved in amino acids metabolism, translation, and secretion, nitrogen metabolism appeared as quite active in sessile cells of *S. xyloso*. Additionally, protein secretion systems were also upregulated in biofilms, suggesting more active protein trafficking in sessile *S. xyloso* cells [187].

L. sakei 23K strain global transcriptome response during growth on ribose [188] and *L. sakei* La22 strain transcriptomic response to meat protein environment [189] have been studied using DNA microarrays.

The ribose uptake and catabolism in *L. sakei* 23K is highly regulated at the transcriptional level, and it is closely related to the catabolism of nucleosides. A global regulation mechanism seems to allow fine tuning of the expression of enzymes, which control the efficient use of available carbon sources [188].

Whole-genome DNA microarrays were used to analyse gene expression related to growth and survival of *L. sakei* La22, when grown in a sarcoplasmic (S) or myofibrillar (M) protein-supplemented chemically defined medium (CDM). Most genes related to peptides or amino acids metabolism were overexpressed in both mediums. Still, meat proteins do not represent a stressful environment for *L. sakei* La22 because no stress response genes were induced [189].

Next generation sequencing methods will improve knowledge related to microbiota and strain characterization involved in dry-fermented meat products. Future work must be done regarding these novel approaches and certainly novel vision of starter behaviour on particular products will be given.

4. Conclusions

The increasing knowledge and exigence level of consumers have forced the search for high value traditional meat products. Consequently, the number of production units (meat transforming) has increased, sometimes in low developed regions in a bewildered way.

The production of traditional meat products, namely, dry-fermented, dry-cured sausages, is still a very traditional and laborious process subjected in several cases to uncontrolled natural environmental conditions. This poses a problem to the producers since their meat products will not be uniform throughout time. Thus, it is necessary to find solutions contributing to the reproducibility of products characteristics. The use of starter cultures based on autochthonous microbiota selection may play here an important role. In fact, the use of these starters in sausages production may improve their sensorial characteristics and contribute to their biopreservation and safety, extending their shelf life, and to increased meat products uniformity.

Selected starter cultures provide a powerful tool for driving the fermentation of meat products, allowing desired quality and safety targets to be reached. Their use in meat fermentation results in acceleration of fermentation time, an improvement of safety (by reducing undesirable microorganisms), and a better quality of the final product. The selection of a starter culture should be carried out in the context of its application, since functionality will depend on the type of sausage, the technology applied, the ripening time, and the ingredients and raw materials used. Future knowledge will be gained with omics methods approach.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was funded by National Funds through FCT-Fundação para a Ciência e a Tecnologia under the Project UID/AGR/00115/2013. M. Laranjo acknowledges a postdoc research grant from FCT (SFRH/BPD/108802/2015).

References

- [1] W. P. Hammes and C. Hertel, "New developments in meat starter cultures," *Meat Science*, vol. 49, no. 1, pp. S125–S138, 1998.
- [2] P. Kumar, M. K. Chatli, A. K. Verma et al., "Quality, functionality, and shelf life of fermented meat and meat products: A review," in *Critical Reviews in Food Science and Nutrition*, vol. 57, pp. 2844–2856, 2017.

- [3] K. Arihara, "Probiotics," in *Handbook of Fermented Meat and Poultry*, F. Toldrá, Ed., pp. 155–160, John Wiley & Sons, Ltd., 2015.
- [4] M. I. Khan, M. S. Arshad, F. M. Anjum, A. Sameen, and W. T. Gill, "Meat as a functional food with special reference to probiotic sausages," *Food Research International*, vol. 44, no. 10, pp. 3125–3133, 2011.
- [5] M. Rouhi, S. Sohrabvandi, and A. M. Mortazavian, "Probiotic Fermented Sausage: Viability of Probiotic Microorganisms and Sensory Characteristics," *Critical Reviews in Food Science and Nutrition*, vol. 53, no. 4, pp. 331–348, 2013.
- [6] A. Jofré, T. Aymerich, and M. Garriga, "Probiotic Fermented Sausages: Myth or Reality? Procedia Food," *Procedia Food Science*, vol. 5, pp. 133–136, 2015.
- [7] Y. Rivera-Espinoza and Y. Gallardo-Navarro, "Non-dairy probiotic products," *Food Microbiology*, vol. 27, no. 1, pp. 1–11, 2010.
- [8] L. Vuyst, G. Falony, and F. Leroy, "Probiotics in fermented sausages," *Meat Science*, vol. 80, no. 1, pp. 75–78, 2008.
- [9] A. V. Wright and L. Axelsson, "Lactic Acid Bacteria: An Introduction," in *Lactic Acid Bacteria: Microbiological and Functional Aspects*, S. Lahtinen, A. C. Ouwehand, S. Salminen, and A. V. Wright, Eds., pp. 1–16, CRC Press, Taylor Francis Group, New York, NY, USA, 2012.
- [10] F.-K. Lücke, "Lactic acid bacteria involved in food fermentations and their present and future uses in food industry," in *Lactic Acid Bacteria: Current Advances in Metabolism, Genetics and Applications*, T. Faruk Bozoglu and B. Ray, Eds., pp. 81–99, Springer, Berlin, Heidelberg, 1996.
- [11] P. S. Cocconcelli and C. Fontana, "Bacteria," in *Handbook of Fermented Meat and Poultry*, F. Toldrá, Ed., pp. 117–128, John Wiley & Sons, Ltd., 2015.
- [12] E. Berni, "Molds," in *Handbook of Fermented Meat and Poultry*, F. Toldrá, Ed., UK, John Wiley Sons, Ltd, 2015.
- [13] R. C. S. Mendonça, D. M. Gouvêa, H. M. Hungaro, A. D. F. Sodrê, and A. Querol-Simon, "Dynamics of the yeast flora in artisanal country style and industrial dry cured sausage (yeast in fermented sausage)," *Food Control*, vol. 29, no. 1, pp. 143–148, 2013.
- [14] L. B. Jensen and L. S. Paddock, "Sausage treatment, Patent US 2225783 A," 1940.
- [15] C. W. Everson, W. E. Danner, and P. A. Hammes, "Bacterial starter cultures in sausage products," *Journal of Agricultural and Food Chemistry*, vol. 18, no. 4, pp. 570–571, 1970.
- [16] F. P. Niinivaara, M. S. Pohja, and Se. Komulain, "Some aspects about using bacterial pure cultures in manufacture of fermented sausages," *Food Technology*, vol. 18, p. 147, 1964.
- [17] G. Vignolo, P. Castellano, and S. Fadda, "Bioprotective Cultures," in *Handbook of Fermented Meat and Poultry*, F. Toldrá, Ed., pp. 129–138, John Wiley & Sons, Ltd., 2015.
- [18] A. Galvez, R. Lucas López, H. Abriouel, E. Valdivia, and N. B. Omar, "Application of bacteriocins in the control of foodborne pathogenic and spoilage bacteria," *Critical Reviews in Biotechnology*, vol. 28, no. 2, pp. 125–152, 2008.
- [19] P. S. Cocconcelli and C. Fontana, "Characteristics and Applications of Microbial Starters in Meat Fermentations," in *Meat Biotechnology*, F. Toldrá, Ed., pp. 129–148, 2008.
- [20] T. Semedo-Lemsaddek, L. Carvalho, C. Tempera et al., "Characterization and Technological Features of Autochthonous Coagulase-Negative Staphylococci as Potential Starters for Portuguese Dry Fermented Sausages," *Journal of Food Science*, vol. 81, no. 5, pp. M1197–M1202, 2016.
- [21] R. Talon, S. Leroy, and I. Lebert, "Microbial ecosystems of traditional fermented meat products: The importance of indigenous starters," *Meat Science*, vol. 77, no. 1, pp. 55–62, 2007.
- [22] R. Talon, S. Leroy, I. Lebert et al., "Safety improvement and preservation of typical sensory qualities of traditional dry fermented sausages using autochthonous starter cultures," *International Journal of Food Microbiology*, vol. 126, no. 1–2, pp. 227–234, 2008.
- [23] M. Elias, M. E. Potes, L. C. Roseiro, C. Santos, A. Gomes, and A. C. Agulheiro-Santos, "The Effect of Starter Cultures on the Portuguese Traditional Sausage, Paio do Alentejo, in Terms of Its Sensory and Textural Characteristics and Polycyclic Aromatic Hydrocarbons Profile," *Journal of Food Research*, vol. 3, pp. 45–56, 2014.
- [24] S. Fonseca, L. I. Ivette Ouoba, I. Franco, and J. Carballo, "Use of molecular methods to characterize the bacterial community and to monitor different native starter cultures throughout the ripening of Galician chorizo," *Food Microbiology*, vol. 34, no. 1, pp. 215–226, 2013.
- [25] S. Fonseca, A. Cachaldora, M. Gómez, I. Franco, and J. Carballo, "Effect of different autochthonous starter cultures on the volatile compounds profile and sensory properties of Galician chorizo, a traditional Spanish dry fermented sausage," *Food Control*, vol. 33, no. 1, pp. 6–14, 2013.
- [26] A. G. M. Scannell, P. M. Kenneally, and E. K. Arendt, "Contribution of starter cultures to the proteolytic process of a fermented non-dried whole muscle ham product," *International Journal of Food Microbiology*, vol. 93, pp. 219–230, 2004.
- [27] F. Bourdichon, S. Casaregola, C. Farrokh et al., "Food fermentations: Microorganisms with technological beneficial use," *International Journal of Food Microbiology*, vol. 154, no. 3, pp. 87–97, 2012.
- [28] L. Morelli, M. L. Calleagri, F. K. Vogensen, and A. v. Wright, "Genetics of Lactic Acid Bacteria. In: Lactic Acid Bacteria: Microbiological and Functional Aspects," in *Lactic Acid Bacteria: Microbiological and Functional Aspects*, S. Lahtinen, A. C. Ouwehand, S. Salminen, and A. V. Wright, Eds., pp. 17–37, CRC Press, Taylor Francis Group, New York, US, 2012.
- [29] M. S. Ammor and B. Mayo, "Selection criteria for lactic acid bacteria to be used as functional starter cultures in dry sausage production: an update," *Meat Science*, vol. 76, no. 1, pp. 138–146, 2007.
- [30] S. C. Ricke, I. Z. Diaz, and J. T. Keeton, *Fermented Meat, Poultry, and Fish Products. in. Food Microbiology: Fundamentals and Frontiers*, ASM Press, Washington, wash, USA, 2007.
- [31] M. Hugas and J. M. Monfort, "Bacterial starter cultures for meat fermentation," *Food Chemistry*, vol. 59, no. 4, pp. 547–554, 1997.
- [32] J. A. Ordóñez, E. M. Hierro, J. M. Bruna, and L. de La Hoz, "Changes in the components of dry-fermented sausages during ripening," *Critical Reviews in Food Science and Nutrition*, vol. 39, no. 4, pp. 329–367, 1999.
- [33] K. Molly, D. Demeyer, G. Johansson, M. Raemaekers, M. Ghistelinck, and I. Geenen, "The importance of meat enzymes in ripening and flavour generation in dry fermented sausages. First results of a European project," *Food Chemistry*, vol. 59, no. 4, pp. 539–545, 1997.
- [34] X. H. Wang, H. Y. Ren, D. Y. Liu, W. Y. Zhu, and W. Wang, "Effects of inoculating *Lactobacillus sakei* starter cultures on the microbiological quality and nitrite depletion of Chinese fermented sausages," *Food Control*, vol. 32, no. 2, pp. 591–596, 2013.

- [35] B. T. Cenci-Goga, P. V. Rossitto, P. Sechi, S. Parmegiani, V. Cambiotti, and J. S. Cullor, "Effect of selected dairy starter cultures on microbiological, chemical and sensory characteristics of swine and venison (*Dama dama*) nitrite-free dry-cured sausages," *Meat Science*, vol. 90, no. 3, pp. 599–606, 2012.
- [36] S. Ahmad, "Sensory Quality of Fermented Sausages as Influenced by Different Combined Cultures of Lactic Acid Bacteria Fermentation during Refrigerated Storage," *Journal of Food Processing Technology*, vol. 4, 2012.
- [37] F. Ravyts and L. De Vuyst, "Prevalence and impact of single-strain starter cultures of lactic acid bacteria on metabolite formation in sourdough," *Food Microbiology*, vol. 28, no. 6, pp. 1129–1139, 2011.
- [38] J. Garcia-Diez and L. Patarata, "Influence of salt level, starter culture, fermentable carbohydrates, and temperature on the behaviour of *L. monocytogenes* in sliced chorizo during storage," *Acta Alimentaria*, vol. 46, pp. 206–213, 2017.
- [39] E. H. Drosinos, S. Paramithiotis, G. Kolovos, I. Tsikouras, and I. Metaxopoulos, "Phenotypic and technological diversity of lactic acid bacteria and staphylococci isolated from traditionally fermented sausages in Southern Greece," *Food Microbiology*, vol. 24, no. 3, pp. 260–270, 2007.
- [40] S. C. Morot-Bizot, S. Leroy, and R. Talon, "Monitoring of staphylococcal starters in two French processing plants manufacturing dry fermented sausages," *Journal of Applied Microbiology*, vol. 102, no. 1, pp. 238–244, 2007.
- [41] C. Fontana, P. S. Cocconcelli, and G. Vignolo, "Monitoring the bacterial population dynamics during fermentation of artisanal Argentinean sausages," *International Journal of Food Microbiology*, vol. 103, no. 2, pp. 131–142, 2005.
- [42] G. Mauriello, A. Casaburi, G. Blaiotta, and F. Villani, "Isolation and technological properties of coagulase negative staphylococci from fermented sausages of Southern Italy," *Meat Science*, vol. 67, no. 1, pp. 149–158, 2004.
- [43] M. C. Garcia Fontan, J. M. Lorenzo, A. Parada, I. Franco, and J. Carballo, "Microbiological characteristics of "androlla", a Spanish traditional pork sausage," *Food Microbiology*, vol. 24, no. 1, pp. 52–58, 2007.
- [44] A. Martín, B. Colín, E. Aranda, M. J. Benito, and M. G. Córdoba, "Characterization of Micrococcaceae isolated from Iberian dry-cured sausages," *Meat Science*, vol. 75, no. 4, pp. 696–708, 2007.
- [45] C. Lopez, L. M. Medina, R. Priego, and R. Jordano, "Behaviour of the constitutive biota of two types of Spanish dry-sausages ripened in a pilot-scale chamber," *Meat Science*, vol. 73, no. 1, pp. 178–180, 2006.
- [46] I. Lebert, S. Leroy, P. Giammarinaro et al., "Diversity of microorganisms in the environment and dry fermented sausages of small traditional French processing units," *Meat Science*, vol. 76, no. 1, pp. 112–122, 2007.
- [47] M. Laranjo, A. Gomes, A. C. Agulheiro-Santos et al., "Impact of salt reduction on biogenic amines, fatty acids, microbiota, texture and sensory profile in traditional blood dry-cured sausages," *Food Chemistry*, vol. 218, pp. 129–136, 2017.
- [48] M. Laranjo, A. Gomes, A. C. Agulheiro-Santos et al., "Characterisation of "Catalão" and "Salsichão" Portuguese traditional sausages with salt reduction," *Meat Science*, vol. 116, pp. 34–42, 2016.
- [49] J. M. Lorenzo, R. Montes, L. Purriños, and D. Franco, "Effect of pork fat addition on the volatile compounds of foal dry-cured sausage," *Meat Science*, vol. 91, no. 4, pp. 506–512, 2012.
- [50] A. C. Venturini, Á. D. Cavenaghi, C. J. C. Castillo, and E. M. Quiñones, "Sensory and microbiological evaluation of uncured fresh chicken sausage with reduced fat content," *Ciência e Tecnologia de Alimentos*, vol. 31, no. 3, pp. 629–634, 2011.
- [51] F. Ravyts, L. Steen, O. Goemaere, H. Paelinck, L. De Vuyst, and F. Leroy, "The application of staphylococci with flavour-generating potential is affected by acidification in fermented dry sausages," *Food Microbiology*, vol. 27, no. 7, pp. 945–954, 2010.
- [52] M. S. Mainar, D. A. Stavropoulou, and F. Leroy, "Exploring the metabolic heterogeneity of coagulase-negative staphylococci to improve the quality and safety of fermented meats: A review," *International Journal of Food Microbiology*, vol. 247, 2017.
- [53] R. Talon, D. Walter, S. Chartier, C. Barrière, and M. C. Montel, "Effect of nitrate and incubation conditions on the production of catalase and nitrate reductase by staphylococci," *International Journal of Food Microbiology*, vol. 52, no. 1-2, pp. 47–56, 1999.
- [54] R. Talon, D. Walter, and M. C. Montel, "Growth and effect of staphylococci and lactic acid bacteria on unsaturated free fatty acids," *Meat Science*, vol. 54, no. 1, pp. 41–47, 2000.
- [55] L. H. Stahnke, A. Holck, A. Jensen, A. Nilsen, and E. Zanardi, "Maturity acceleration of italian dried sausage by *Staphylococcus carnosus*-Relationship between maturity and flavor compounds," *Journal of Food Science*, vol. 67, no. 5, pp. 1914–1921, 2002.
- [56] H. C. Beck, A. M. Hansen, and F. R. Lauritsen, "Catabolism of leucine to branched-chain fatty acids in *Staphylococcus xylosus*," *Journal of Applied Microbiology*, vol. 96, no. 5, pp. 1185–1193, 2004.
- [57] P. T. Olesen, A. S. Meyer, and L. H. Stahnke, "Generation of flavour compounds in fermented sausages - The influence of curing ingredients, *Staphylococcus* starter culture and ripening time," *Meat Science*, vol. 66, no. 3, pp. 675–687, 2004.
- [58] J. Samelis, J. Metaxopoulos, M. Vlassi, and A. Pappa, "Stability and safety of traditional Greek salami—a microbiological ecology study," *International Journal of Food Microbiology*, vol. 44, no. 1-2, pp. 69–82, 1998.
- [59] M. Lusnic, T. Polak, L. Gašperlin et al., "Degradation of PCBs in a frankfurter-type meat emulsion: Effects of a meat starter, its proteins extract and thermal treatments," *Food and Chemical Toxicology*, vol. 50, no. 8, pp. 2643–2647, 2012.
- [60] R. Casquete, M. J. Benito, A. Martín, S. Ruiz-Moyano, A. Hernández, and M. G. Córdoba, "Effect of autochthonous starter cultures in the production of "salchichón", a traditional Iberian dry-fermented sausage, with different ripening processes," *LWT- Food Science and Technology*, vol. 44, no. 7, pp. 1562–1571, 2011.
- [61] R. Casquete, M. J. Benito, A. Martín, S. Ruiz-Moyano, J. J. Cordoba, and M. G. Cordoba, "Role of an autochthonous starter culture and the protease EPg222 on the sensory and safety properties of a traditional Iberian dry-fermented sausage salchichon," *Food Microbiol*, vol. 28, pp. 1432–40, 2011.
- [62] R. Casquete, M. J. Benito, A. Martín, S. Ruiz-Moyano, E. Aranda, and M. G. Córdoba, "Microbiological quality of salchichón and chorizo, traditional Iberian dry-fermented sausages from two different industries, inoculated with autochthonous starter cultures," *Food Control*, vol. 24, no. 1-2, pp. 191–198, 2012.
- [63] I. Essid and M. Hassouna, "Effect of inoculation of selected *Staphylococcus xylosus* and *Lactobacillus plantarum* strains on biochemical, microbiological and textural characteristics of a Tunisian dry fermented sausage," *Food Control*, vol. 32, no. 2, pp. 707–714, 2013.
- [64] M. Bedia, L. Méndez, and S. Bañón, "Evaluation of different starter cultures (*Staphylococci* plus Lactic Acid Bacteria) in




- semi-ripened Salami stuffed in swine gut," *Meat Science*, vol. 87, no. 4, pp. 381–386, 2011.
- [65] J. M. Aro Aro, P. Nyam-Osor, K. Tsuji, K.-I. Shimada, M. Fukushima, and M. Sekikawa, "The effect of starter cultures on proteolytic changes and amino acid content in fermented sausages," *Food Chemistry*, vol. 119, no. 1, pp. 279–285, 2010.
- [66] P. Tremonte, A. Reale, T. Di Renzo et al., "Interactions between *Lactobacillus sakei* and CNC (*Staphylococcus xylosus* and *Kocuria varians*) and their influence on proteolytic activity," *Letters in Applied Microbiology*, vol. 51, no. 5, pp. 586–594, 2010.
- [67] A. Casaburi, V. Di Martino, P. Ferranti, L. Picariello, and F. Villani, "Technological properties and bacteriocins production by *Lactobacillus curvatus* 54M16 and its use as starter culture for fermented sausage manufacture," *Food Control*, vol. 59, pp. 31–45, 2016.
- [68] M. Z. Barbosa, S. D. Todorov, I. Ivanova, J.-M. Chobert, T. Haertlé, and B. D. G. de Melo Franco, "Improving safety of salami by application of bacteriocins produced by an autochthonous *Lactobacillus curvatus* isolate," *Food Microbiology*, vol. 46, pp. 254–262, 2015.
- [69] M. Simonová, V. Stropfová, M. Marciňáková et al., "Characterization of *Staphylococcus xylosus* and *Staphylococcus carnosus* isolated from Slovak meat products," *Meat Science*, vol. 73, no. 4, pp. 559–564, 2006.
- [70] Á. M. Fiorentini, M. C. Sawitzki, T. M. Bertol, and E. S. Sant'Anna, "Viability of *Staphylococcus xylosus* isolated from artisanal sausages for application as starter cultures in meat products," *Brazilian Journal of Microbiology*, vol. 40, no. 1, pp. 129–133, 2009.
- [71] E. B. Bingol, F. Yilmaz, H. Yardibi et al., "Effect of lipolytic starter cultures on ripening and quality of Turkish type fermented sausages (sucuk)," *Current Opinion in Biotechnology*, vol. 22, p. S97, 2011.
- [72] M. D. Selgas and M. L. García, "Yeasts," in *Handbook of Fermented Meat and Poultry*, F. Toldrá, Ed., pp. 139–146, USA, Wiley Blackwell, 2015.
- [73] M. Flores, S. Corral, L. Cano-García, A. Salvador, and C. Belloch, "Yeast strains as potential aroma enhancers in dry fermented sausages," *International Journal of Food Microbiology*, vol. 212, pp. 16–24, 2015.
- [74] B. C. Viljoen, G. A. Dykes, M. Callis, and A. von Holy, "Yeasts associated with Vienna sausage packaging," *International Journal of Food Microbiology*, vol. 18, no. 1, pp. 53–62, 1993.
- [75] B. C. Viljoen and T. Greyling, "Yeasts associated with Cheddar and Gouda making," *International Journal of Food Microbiology*, vol. 28, no. 1, pp. 79–88, 1995.
- [76] E. Miteva, E. Kirova, D. Gadjeva, and M. Radeva, "Sensory aroma and taste profiles of raw-dried sausages manufactured with a lipolytically active yeast culture," *Nahrung-Food*, vol. 30, pp. 829–832, 1986.
- [77] M. A. Dura, M. Flores, and F. Toldra, "Effect of *Debaryomyces* spp. on the proteolysis of dry-fermented sausages," *Meat Science*, vol. 68, pp. 319–328, 2004.
- [78] M. Flores, M.-A. Durá, A. Marco, and F. Toldrá, "Effect of *Debaryomyces* spp. on aroma formation and sensory quality of dry-fermented sausages," *Meat Science*, vol. 68, no. 3, pp. 439–446, 2004.
- [79] A. Martin, J. J. Cordoba, E. Aranda, M. G. Cordoba, and M. A. Asensio, "Contribution of a selected fungal population to the volatile compounds on dry-cured ham," *International Journal of Food Microbiology*, vol. 110, pp. 8–18, 2006.
- [80] J. M. Bruna, E. M. Hierro, L. De La Hoz, D. S. Mottram, M. Fernández, and J. A. Ordóñez, "Changes in selected biochemical and sensory parameters as affected by the superficial inoculation of *Penicillium camemberti* on dry fermented sausages," *International Journal of Food Microbiology*, vol. 85, no. 1-2, pp. 111–125, 2003.
- [81] M. Papagianni, I. Ambrosiadis, and G. Filiouis, "Mould growth on traditional greek sausages and penicillin production by *Penicillium* isolates," *Meat Science*, vol. 76, no. 4, pp. 653–657, 2007.
- [82] L. O. Sunesen and L. H. Stahnke, "Mould starter cultures for dry sausages—selection, application and effects," *Meat Science*, vol. 65, no. 3, pp. 935–948, 2003.
- [83] V. Ludemann, M. Greco, M. P. Rodríguez, J. C. Basílico, and A. G. Pardo, "Conidial production by *Penicillium nalgioense* for use as starter cultures in dry fermented sausages by solid state fermentation," *LWT- Food Science and Technology*, vol. 43, no. 2, pp. 315–318, 2010.
- [84] P. Paulsen, S. Vali, and F. Bauer, "Quality traits of wild boar mould-ripened salami manufactured with different selections of meat and fat tissue, and with and without bacterial starter cultures," *Meat Science*, vol. 89, no. 4, pp. 486–490, 2011.
- [85] I. F. Nes and J. R. Tagg, "Novel lantibiotics and their pre-peptides," *Antonie van Leeuwenhoek-Journal of Microbiology*, vol. 69, no. 2, pp. 89–97, 1996.
- [86] M. J. Fraqueza, L. Patarata, and A. Lauková, "Protective Starter Cultures and Bacteriocins in Fermented Meats," in *Fermented Meat Products: Health Aspects*, N. Zdolec, Ed., pp. 228–269, CRC Press, New York, 2016.
- [87] M. Hugas, M. Garriga, M. T. Aymerich, and J. M. Monfort, "Inhibition of *Listeria* in dry fermented sausages by the bacteriocinogenic *Lactobacillus sake* CTC494," *Journal of Applied Bacteriology*, vol. 79, no. 3, pp. 322–330, 1995.
- [88] P. M. Foegeding, A. B. Thomas, D. H. Pilkington, and T. R. Klaenhammer, "Enhanced control of *Listeria monocytogenes* by *in situ*-produced pediocin during dry fermented sausage production," *Applied and Environmental Microbiology*, vol. 58, no. 3, pp. 884–890, 1992.
- [89] J. C. Nieto-Lozano, J. I. Reguera-Useros, M. D. C. Peláez-Martínez, and A. H. De La Torre, "Effect of a bacteriocin produced by *Pediococcus acidilactici* against *Listeria monocytogenes* and *Clostridium perfringens* on Spanish raw meat," *Meat Science*, vol. 72, no. 1, pp. 57–61, 2006.
- [90] T. Azuma, D. K. Bagenda, T. Yamamoto, Y. Kawai, and K. Yamazaki, "Inhibition of *Listeria monocytogenes* by freeze-dried piscicocin CS526 fermentate in food," *Letters in Applied Microbiology*, vol. 44, no. 2, pp. 138–144, 2007.
- [91] Á. Alegria, S. Delgado, C. Rocas, B. López, and B. Mayo, "Bacteriocins produced by wild *Lactococcus lactis* strains isolated from traditional, starter-free cheeses made of raw milk," *International Journal of Food Microbiology*, vol. 143, no. 1-2, pp. 61–66, 2010.
- [92] C. Henning, D. Gautam, and P. Muriana, "Identification of Multiple Bacteriocins in *Enterococcus* spp. Using an Enterococcus-Specific Bacteriocin PCR Array," *Microorganisms*, vol. 3, pp. 1–16, 2015.
- [93] I. F. Nes, D. B. Diep, and H. Holo, "Bacteriocin diversity in *Streptococcus* and *Enterococcus*," *Journal of Bacteriology*, vol. 189, no. 4, pp. 1189–1198, 2007.
- [94] J. C. Nieto-Lozano, J. I. Reguera-Useros, M. D. C. Peláez-Martínez, G. Sacristán-Pérez-Minayo, Á. J. Gutiérrez-Fernández, and A. H. D. la Torre, "The effect of the pediocin PA-1

- produced by *Pediococcus acidilactici* against *Listeria monocytogenes* and *Clostridium perfringens* in Spanish dry-fermented sausages and frankfurters,” *Food Control*, vol. 21, no. 5, pp. 679–685, 2010.
- [95] S. D. Todorov, P. Ho, M. Vaz-Velho, and L. M. T. Dicks, “Characterization of bacteriocins produced by two strains of *Lactobacillus plantarum* isolated from Beloura and Chouriço, traditional pork products from Portugal,” *Meat Science*, vol. 84, no. 3, pp. 334–343, 2010.
- [96] Y. Kingcha, A. Tosukhowong, T. Zendo et al., “Anti-listeria activity of *Pediococcus pentosaceus* BCC 3772 and application as starter culture for Nham, a traditional fermented pork sausage,” *Food Control*, vol. 25, no. 1, pp. 190–196, 2012.
- [97] P. Jaichumjai, R. Valyasevi, A. Assavanig, and P. Kurdi, “Isolation and characterization of acid-sensitive *Lactobacillus plantarum* with application as starter culture for Nham production,” *Food Microbiology*, vol. 27, no. 6, pp. 741–748, 2010.
- [98] A. Lauková, M. Simonová, and V. Stropfová, “*Staphylococcus xylosum* S03/1M/1/2, bacteriocin-producing meat starter culture or additive,” *Food Control*, vol. 21, no. 7, pp. 970–973, 2010.
- [99] F. Ravyts, L. D. Vuyst, and F. Leroy, “Bacterial diversity and functionalities in food fermentations,” *Engineering in Life Sciences*, vol. 12, no. 4, pp. 356–367, 2012.
- [100] T. Rimaux, G. Vrancken, V. Pothakos, D. Maes, L. De Vuyst, and F. Leroy, “The kinetics of the arginine deiminase pathway in the meat starter culture *Lactobacillus sakei* CTC 494 are pH-dependent,” *Food Microbiology*, vol. 28, no. 3, pp. 597–604, 2011.
- [101] T. Rimaux, G. Vrancken, B. Vuylsteke, L. De Vuyst, and F. Leroy, “The pentose moiety of adenosine and inosine is an important energy source for the fermented-meat starter culture *Lactobacillus sakei* CTC 494,” *Applied and Environmental Microbiology*, vol. 77, no. 18, pp. 6539–6550, 2011.
- [102] A. Greppi, I. Ferrocino, A. La Stora, K. Rantsiou, D. Ercolini, and L. Cocolin, “Monitoring of the microbiota of fermented sausages by culture independent rRNA-based approaches,” *International Journal of Food Microbiology*, vol. 212, pp. 67–75, 2015.
- [103] S. Barlow, A. Chesson, J. Collins et al., “Opinion of the Scientific Committee on a request from EFSA related to a generic approach to the safety assessment by EFSA of microorganisms used in food/feed and the production of food/feed additives,” *EFSA Journal*, vol. 3, pp. 1–12, 2005.
- [104] A. Ricci, A. Allende, D. Bolton et al., “Scientific Opinion on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA,” *EFSA Journal*, vol. 15, p. 4664, 2017.
- [105] S. Laulund, A. Wind, P. Derkx, and V. Zuliani, “Regulatory and safety requirements for food cultures,” *Microorganisms*, vol. 5, p. 28, 2017.
- [106] W. Witte, “Selective pressure by antibiotic use in livestock,” *International Journal of Antimicrobial Agents*, vol. 16, no. 1, pp. S19–S24, 2000.
- [107] G. Rychen, G. Aquilina, G. Azimonti et al., “Guidance on the characterisation of microorganisms used as feed additives or as production organisms. EFSA Journal,” *Endorsed for public consultation on 18*, 2017.
- [108] S. Even, S. Leroy, C. Charlier et al., “Low occurrence of safety hazards in coagulase negative staphylococci isolated from fermented foodstuffs,” *International Journal of Food Microbiology*, vol. 139, no. 1–2, pp. 87–95, 2010.
- [109] G. Landeta, J. A. Curiel, A. V. Carrascosa, R. Muñoz, and B. de las Rivas, “Characterization of coagulase-negative staphylococci isolated from Spanish dry cured meat products,” *Meat Science*, vol. 93, no. 3, pp. 387–396, 2013.
- [110] S. Kastner, V. Perreten, H. Bleuler, G. Hugenschmidt, C. Lacroix, and L. Meile, “Antibiotic susceptibility patterns and resistance genes of starter cultures and probiotic bacteria used in food,” *Systematic and Applied Microbiology*, vol. 29, no. 2, pp. 145–155, 2006.
- [111] B. M. Marshall, D. J. Ochieng, and S. B. Levy, “Commensals: underappreciated reservoir of antibiotic resistance,” *Microbe*, vol. 4, no. 5, pp. 231–238, 2009.
- [112] C. Devirgiliis, S. Barile, and G. Perozzi, “Antibiotic resistance determinants in the interplay between food and gut microbiota,” *Genes & Nutrition*, vol. 6, no. 3, pp. 275–284, 2011.
- [113] T. Aymerich, B. Martín, M. Garriga, M. C. Vidal-Carou, S. Bover-Cid, and M. Hugas, “Safety properties and molecular strain typing of lactic acid bacteria from slightly fermented sausages,” *Journal of Applied Microbiology*, vol. 100, no. 1, pp. 40–49, 2006.
- [114] S. Federici, F. Ciarrocchi, R. Campana, E. Ciandrini, G. Blasi, and W. Baffone, “Identification and functional traits of lactic acid bacteria isolated from Ciauscolo salami produced in Central Italy,” *Meat Science*, vol. 98, no. 4, pp. 575–584, 2014.
- [115] R. Comunian, E. Daga, I. Dupré et al., “Susceptibility to tetracycline and erythromycin of *Lactobacillus paracasei* strains isolated from traditional Italian fermented foods,” *International Journal of Food Microbiology*, vol. 138, no. 1–2, pp. 151–156, 2010.
- [116] G. Landeta, J. A. Curiel, A. V. Carrascosa, R. Muñoz, and B. De las Rivas, “Technological and safety properties of lactic acid bacteria isolated from Spanish dry-cured sausages,” *Meat Science*, vol. 95, no. 2, pp. 272–280, 2013.
- [117] D. Zonenschain, A. Rebecchi, and L. Morelli, “Erythromycin- and tetracycline-resistant lactobacilli in Italian fermented dry sausages,” *Journal of Applied Microbiology*, vol. 107, no. 5, pp. 1559–1568, 2009.
- [118] T. Ribeiro, M. Oliveira, M. J. Fraqueza et al., “Antibiotic resistance and virulence factors among Enterococci isolated from chouriço, a traditional Portuguese dry fermented sausage,” *Journal of Food Protection*, vol. 74, no. 3, pp. 465–469, 2011.
- [119] D. Gevers, G. Huys, F. Devlieghere, M. Uyttendaele, J. Debevere, and J. Swings, “Isolation and identification of tetracycline resistant lactic acid bacteria from pre-packed sliced meat products,” *Systematic and Applied Microbiology*, vol. 23, no. 2, pp. 279–284, 2000.
- [120] D. Gevers, G. Huys, and J. Swings, “In vitro conjugal transfer of tetracycline resistance from *Lactobacillus* isolates to other gram-positive bacteria,” *FEMS Microbiology Letters*, vol. 225, no. 1, pp. 125–130, 2003.
- [121] M. J. Fraqueza, “Antibiotic resistance of lactic acid bacteria isolated from dry-fermented sausages,” *International Journal of Food Microbiology*, vol. 212, pp. 76–88, 2015.
- [122] S. Lu, H. Ji, Q. Wang et al., “The effects of starter cultures and plant extracts on the biogenic amine accumulation in traditional Chinese smoked horsemeat sausages,” *Food Control*, vol. 50, pp. 869–875, 2015.
- [123] T. Komprda, D. Smělá, P. Pechová, L. Kalhotka, J. Štencl, and B. Klejduš, “Effect of starter culture, spice mix and storage time and temperature on biogenic amine content of dry fermented sausages,” *Meat Science*, vol. 67, no. 4, pp. 607–616, 2004.

- [124] M. L. Latorre-Moratalla, S. Bover-Cid, M. T. Veciana-Nogués, and M. C. Vidal-Carou, "Control of biogenic amines in fermented sausages: role of starter cultures," *Frontiers in Microbiology*, vol. 3, article 169, 2012.
- [125] M. L. Latorre-Moratalla, S. Bover-Cid, R. Talon et al., "Distribution of aminogenic activity among potential autochthonous starter cultures for dry fermented sausages," *Journal of Food Protection*, vol. 73, no. 3, pp. 524–528, 2010.
- [126] M. L. Latorre-Moratalla, S. Bover-Cid, R. Talon et al., "Strategies to reduce biogenic amine accumulation in traditional sausage manufacturing," *LWT- Food Science and Technology*, vol. 43, no. 1, pp. 20–25, 2010.
- [127] T. Semedo, M. Almeida Santos, M. F. Silva Lopes, J. J. Figueiredo Marques, M. T. Barreto Crespo, and R. Tenreiro, "Virulence factors in food, clinical and reference enterococci: A common trait in the genus?" *Systematic and Applied Microbiology*, vol. 26, no. 1, pp. 13–22, 2003.
- [128] S. C. Santos, M. J. Fraqueza, M. Elias, A. Salvador Barreto, and T. Semedo-Lemsaddek, "Traditional dry smoked fermented meat sausages: Characterization of autochthonous enterococci," *LWT- Food Science and Technology*, vol. 79, pp. 410–415, 2017.
- [129] J. Barbosa, P. A. Gibbs, and P. Teixeira, "Virulence factors among enterococci isolated from traditional fermented meat products produced in the North of Portugal," *Food Control*, vol. 21, no. 5, pp. 651–656, 2010.
- [130] F. M. Aarestrup, Y. Agersø, P. Ahrens, J. C. Ø. Jørgensen, M. Madsen, and L. B. Jensen, "Antimicrobial susceptibility and presence of resistance genes in staphylococci from poultry," *Veterinary Microbiology*, vol. 74, no. 4, pp. 353–364, 2000.
- [131] C. Vernozy-Rozand, C. Mazuy, G. Prevost et al., "Enterotoxin production by coagulase-negative staphylococci isolated from goats' milk and cheese," *International Journal of Food Microbiology*, vol. 30, no. 3, pp. 271–280, 1996.
- [132] B. Martín, M. Garriga, M. Hugas, S. Bover-Cid, M. T. Veciana-Nogués, and T. Aymerich, "Molecular, technological and safety characterization of Gram-positive catalase-positive cocci from slightly fermented sausages," *International Journal of Food Microbiology*, vol. 107, no. 2, pp. 148–158, 2006.
- [133] P. T. Fowoyo and S. T. Ogunbanwo, "Virulence and toxigenicity of coagulase-negative staphylococci in Nigerian traditional fermented foods," *Canadian Journal of Microbiology*, vol. 62, no. 7, pp. 572–578, 2016.
- [134] C. G. Kumar and S. K. Anand, "Significance of microbial biofilms in food industry: a review," *International Journal of Food Microbiology*, vol. 42, no. 1-2, pp. 9–27, 1998.
- [135] V. Leriche and B. Carpentier, "Limitation of adhesion and growth of *Listeria monocytogenes* on stainless steel surfaces by *Staphylococcus sciuri* biofilms," *Journal of Applied Microbiology*, vol. 88, no. 4, pp. 594–605, 2000.
- [136] A. Jain and A. Agarwal, "Biofilm production, a marker of pathogenic potential of colonizing and commensal staphylococci," *Journal of Microbiological Methods*, vol. 76, no. 1, pp. 88–92, 2009.
- [137] P. Kotilainen, "Association of coagulase-negative staphylococcal slime production and adherence with the development and outcome of adult septicemias," *Journal of Clinical Microbiology*, vol. 28, no. 12, pp. 2779–2785, 1990.
- [138] S. Leroy, I. Lebert, J.-P. Chacornac, P. Chavant, T. Bernardi, and R. Talon, "Genetic diversity and biofilm formation of *Staphylococcus equorum* isolated from naturally fermented sausages and their manufacturing environment," *International Journal of Food Microbiology*, vol. 134, no. 1-2, pp. 46–51, 2009.
- [139] A. Fagerlund, S. Langsrud, E. Heir, M. I. Mikkelsen, and T. Møretrø, "Biofilm matrix composition affects the susceptibility of food associated staphylococci to cleaning and disinfection agents," *Frontiers in Microbiology*, vol. 7, article no. 856, 2016.
- [140] T. Møretrø, L. Hermansen, A. L. Holck, M. S. Sidhu, K. Rudi, and S. Langsrud, "Biofilm formation and the presence of the intercellular adhesion locus *ica* among staphylococci from food and food processing environments," *Applied and Environmental Microbiology*, vol. 69, no. 9, pp. 5648–5655, 2003.
- [141] S. Planchon, B. Gaillard-Martinie, E. Dordet-Frisoni et al., "Formation of biofilm by *Staphylococcus xylosum*," *International Journal of Food Microbiology*, vol. 109, no. 1-2, pp. 88–96, 2006.
- [142] S. Planchon, B. Gaillard-Martinie, S. Leroy, M. N. Bellon-Fontaine, S. Fadda, and R. Talon, "Surface properties and behaviour on abiotic surfaces of *Staphylococcus carnosus*, a genetically homogeneous species," *Food Microbiology*, vol. 24, no. 1, pp. 44–51, 2007.
- [143] N. C. Gómez, J. M. P. Ramiro, B. X. V. Quecan, and B. D. G. de Melo Franco, "Use of potential probiotic lactic acid bacteria (LAB) biofilms for the control of *Listeria monocytogenes*, *Salmonella Typhimurium*, and *Escherichia coli* O157: H7 biofilms formation," *Frontiers in Microbiology*, vol. 7, article no. 863, 2016.
- [144] S. Chaillou, M.-C. Champomier-Vergès, M. Cornet et al., "The complete genome sequence of the meat-borne lactic acid bacterium *Lactobacillus sakei* 23K," *Nature Biotechnology*, vol. 23, no. 12, pp. 1527–1533, 2005.
- [145] P. D. Cotter and C. Hill, "Surviving the acid test: responses of gram-positive bacteria to low pH," *Microbiology and Molecular Biology Reviews*, vol. 67, no. 3, pp. 429–453, 2003.
- [146] M. Van de Guchte, P. Serror, C. Chervaux, T. Smokvina, S. D. Ehrlich, and E. Maguin, "Stress responses in lactic acid bacteria," *Antonie van Leeuwenhoek-Journal of Microbiology*, vol. 82, no. 1-4, pp. 187–216, 2002.
- [147] T. Fujii, C. Ingham, J. Nakayama et al., "Two homologous agr-like quorum-sensing systems cooperatively control adherence, cell morphology, and cell viability properties in *Lactobacillus plantarum* WCFS1," *Journal of Bacteriology*, vol. 190, no. 23, pp. 7655–7665, 2008.
- [148] S. Lebeer, S. C. J. De Keersmaecker, T. L. A. Verhoeven, A. A. Fadda, K. Marchal, and J. Vanderleyden, "Functional analysis of *luxS* in the probiotic strain *Lactobacillus rhamnosus* GG reveals a central metabolic role important for growth and biofilm formation," *Journal of Bacteriology*, vol. 189, no. 3, pp. 860–871, 2007.
- [149] M. H. J. Sturme, J. Nakayama, D. Molenaar et al., "An agr-like two-component regulatory system in *Lactobacillus plantarum* is involved in production of a novel cyclic peptide and regulation of adherence," *Journal of Bacteriology*, vol. 187, no. 15, pp. 5224–5235, 2005.
- [150] G. W. Tannock, J. B. Luchansky, L. Miller et al., "Molecular Characterization of a Plasmid-Borne (pGT633) Erythromycin Resistance Determinant (*ermGT*) from *Lactobacillus reuteri* 100-63," *Plasmid*, vol. 31, no. 1, pp. 60–71, 1994.
- [151] F. Leroy and L. de Vuyst, "Lactic acid bacteria as functional starter cultures for the food fermentation industry," *Trends in Food Science & Technology*, vol. 15, no. 2, pp. 67–78, 2004.
- [152] R. Talon, I. Lebert, S. Leroy et al., "Microbial ecosystem of traditional dry fermented sausages in Mediterranean countries and Slovakia," *Mediterranean Ecosystems: Dynamics, Management and Conservation*, pp. 115–127, 2012.

- [153] M. P. Zacharof and R. W. Lovitt, "Bacteriocins Produced by Lactic Acid Bacteria A Review Article," in *Proceedings of the 3rd International Conference on Biotechnology and Food Science* (edited by DAN, pp. 50–56, 2012.
- [154] T. D. Klingberg, L. Axelsson, K. Naterstad, D. Elsser, and B. B. Budde, "Identification of potential probiotic starter cultures for Scandinavian-type fermented sausages," *International Journal of Food Microbiology*, vol. 105, no. 3, pp. 419–431, 2005.
- [155] M. Trzaskowska, D. Kołozyn-Krajewska, K. M. Wójciak, and Z. J. Dolatowski, "Microbiological quality of raw-fermented sausages with *Lactobacillus casei* LOCK 0900 probiotic strain," *Food Control*, vol. 35, no. 1, pp. 184–191, 2014.
- [156] E. Sayas-Barberá, M. Viuda-Martos, F. Fernández-López, J. A. Pérez-Alvarez, and E. Sendra, "Combined use of a probiotic culture and citrus fiber in a traditional sausage 'Longaniza de Pascua'," *Food Control*, vol. 27, no. 2, pp. 343–350, 2012.
- [157] S. Ruiz-Moyano, A. Martín, M. J. Benito, A. Hernández, R. Casquete, and M. de Guia Córdoba, "Application of *Lactobacillus fermentum* HL57 and *Pediococcus acidilactici* SP979 as potential probiotics in the manufacture of traditional Iberian dry-fermented sausages," *Food Microbiology*, vol. 28, no. 5, pp. 839–847, 2011.
- [158] FAO/WHO, "Report of a joint FAO/WHO working group of drafting guidelines for the evaluation of probiotics in food," guidelines for the evaluation of probiotics in food, London, UK, 2002.
- [159] S. Salminen, A. von Wright, L. Morelli et al., "Demonstration of safety of probiotics—a review," *International Journal of Food Microbiology*, vol. 44, no. 1-2, pp. 93–106, 1998.
- [160] D. Kołozyn-Krajewska and Z. J. Dolatowski, "Probiotic meat products and human nutrition," *Process Biochemistry*, vol. 47, no. 12, pp. 1761–1772, 2012.
- [161] J. Hugenholtz and E. J. Smid, "Nutraceutical production with food-grade microorganisms," *Current Opinion in Biotechnology*, vol. 13, no. 5, pp. 497–507, 2002.
- [162] W. Sybesma, M. Starrenburg, L. Tijsseling, M. H. N. Hoefnagel, and J. Hugenholtz, "Effects of cultivation conditions on folate production by lactic acid bacteria," *Applied and Environmental Microbiology*, vol. 69, no. 8, pp. 4542–4548, 2003.
- [163] C. Burgess, M. O'Connell-Motherway, W. Sybesma, J. Hugenholtz, and D. Van Sinderen, "Riboflavin production in *Lactococcus lactis*: Potential for in situ production of vitamin-enriched foods," *Applied and Environmental Microbiology*, vol. 70, no. 10, pp. 5769–5777, 2004.
- [164] W. Sybesma, C. Burgess, M. Starrenburg, D. Van Sinderen, and J. Hugenholtz, "Multivitamin production in *Lactococcus lactis* using metabolic engineering," *Metabolic Engineering*, vol. 6, no. 2, pp. 109–115, 2004.
- [165] C. Pennacchia, E. E. Vaughan, and F. Villani, "Potential probiotic *Lactobacillus* strains from fermented sausages: Further investigations on their probiotic properties," *Meat Science*, vol. 73, no. 1, pp. 90–101, 2006.
- [166] R. Rubio, T. Aymerich, S. Bover-Cid, M. D. Guàrdia, J. Arnau, and M. Garriga, "Probiotic strains *Lactobacillus plantarum* 299V and *Lactobacillus rhamnosus* GG as starter cultures for fermented sausages," *LWT- Food Science and Technology*, vol. 54, no. 1, pp. 51–56, 2013.
- [167] K. Neffe-Skocińska, D. Jaworska, D. Kołozyn-Krajewska, Z. Dolatowski, and L. Jachacz-Jówko, "The effect of LAB as probiotic starter culture and green tea extract addition on dry fermented pork loins quality," *BioMed Research International*, vol. 2015, Article ID 452757, 2015.
- [168] K. Neffe-Skocińska, K. Wójciak, and D. Zielinska, "Probiotic microorganisms in dry fermented meat products," in *Probiotic Microorganisms in Dry Fermented Meat Products. in. Probiotics and Prebiotics in Human Nutrition and Health*, InTech, Rijeka, Croatia, 2016.
- [169] Regulation (EC) No. 1924/2006 of the European Parliament and of the Council of 20 December 2006 on nutrition and health claims made on foods.
- [170] EU, "Commission Regulation (EU) No 432/2012 of 16 May 2012 establishing a list of permitted health claims made on foods, other than those referring to the reduction of disease risk and to children's development and health," *Official Journal of the European Union*, vol. 136, pp. 1–40, 2012.
- [171] J. Anba-Mondoloni, M.-C. Champomier-Vergès, M. Zagorec et al., "The Genetics of Microbial Starters," in *Handbook of Fermented Meat and Poultry: Second Edition*, F. TOLDRA, Ed., pp. 161–168, Wiley Blackwell, 2015.
- [172] F. Villani, A. Casaburi, C. Pennacchia, L. Filosa, F. Russo, and D. Ercolini, "Microbial ecology of the soppressata of Vallo di Diano, a traditional dry fermented sausage from southern Italy, and *in vitro* and *in situ* selection of autochthonous starter cultures," *Applied and Environmental Microbiology*, vol. 73, no. 17, pp. 5453–5463, 2007.
- [173] J. A. Klappenbach, J. M. Dunbar, and T. M. Schmidt, "rRNA operon copy number reflects ecological strategies of bacteria," *Applied and Environmental Microbiology*, vol. 66, no. 4, pp. 1328–1333, 2000.
- [174] C.-J. Liu, R. Wang, F.-M. Gong et al., "Complete genome sequences and comparative genome analysis of *Lactobacillus plantarum* strain 5-2 isolated from fermented soybean," *Genomics*, vol. 106, no. 6, pp. 404–411, 2015.
- [175] W. P. Hammes, A. Bantleon, and S. Min, "Lactic acid bacteria in meat fermentation," *FEMS Microbiology Letters*, vol. 87, no. 1-2, pp. 165–173, 1990.
- [176] A. McLeod, D. A. Brede, I. Rud, and L. Axelsson, "Genome sequence of *Lactobacillus sakei* subsp. *sakei* LS25, a commercial starter culture strain for fermented sausage," *Genome Announcements*, vol. 1, no. 4, Article ID e00475-13, 2013.
- [177] S. H. Lee, M. Y. Jung, B. Park et al., "Complete genome sequence of *Pediococcus pentosaceus* strain wikim 20, isolated from Korean kimchi," *Genome Announcements*, vol. 4, no. 6, Article ID e01233-16, 2016.
- [178] S. H. Dantoft, E. M. Bielak, J.-G. Seo, M.-J. Chung, and P. R. Jensen, "Complete genome sequence of *Pediococcus pentosaceus* strain SL4," *Genome Announcements*, vol. 1, no. 6, Article ID e01106-13, 2013.
- [179] G.-S. Park, S.-J. Hong, B. K. Jung et al., "Whole genome sequence of lactic acid bacterium *Pediococcus acidilactici* strain Sl," *Brazilian Journal of Microbiology*, vol. 48, no. 3, 2017.
- [180] S. J. Labrie, L. El Haddad, D. M. Tremblay et al., "First complete genome sequence of *Staphylococcus xylosus*, a meat starter culture and a host to propagate *Staphylococcus aureus* phages," *Genome Announcements*, vol. 2, no. 4, Article ID e00671-14, 2014.
- [181] E. Dordet-Frisoni, G. Dorchie, C. De Araujo, R. Talon, and S. Leroy, "Genomic diversity in *Staphylococcus xylosus*," *Applied and Environmental Microbiology*, vol. 73, no. 22, pp. 7199–7209, 2007.
- [182] R. Rosenstein, C. Nerz, L. Biswas et al., "Genome analysis of the meat starter culture bacterium *Staphylococcus carnosus* TM300," *Applied and Environmental Microbiology*, vol. 75, no. 3, pp. 811–822, 2009.

- [183] M. S. Resch, C. Nerz, R. Rosenstein, F. Götz, and C. Hertel, "DNA microarray based detection of genes involved in safety and technologically relevant properties of food associated coagulase-negative staphylococci," *International Journal of Food Microbiology*, vol. 145, pp. 449–458, 2011.
- [184] A. Vermassen, A. de la Foye, V. Loux, R. Talon, and S. Leroy, "Transcriptomic analysis of *Staphylococcus xylosus* in the presence of nitrate and nitrite in meat reveals its response to nitrosative stress," *Frontiers in Microbiology*, vol. 5, article no. 691, 2014.
- [185] A. Vermassen, E. Dordet-Frisoni, A. De La Foye et al., "Adaptation of *Staphylococcus xylosus* to nutrients and osmotic stress in a salted meat model," *Frontiers in Microbiology*, vol. 7, article no. 87, 2016.
- [186] R. Gaupp, N. Ledala, and G. A. Somerville, "Staphylococcal response to oxidative stress," *Front Cell Infect Microbiol*, vol. 2, p. 33, 2012.
- [187] S. Planchon, M. Desvaux, I. Chafsey et al., "Comparative subproteome analyses of planktonic and sessile *Staphylococcus xylosus* C2a: New insight in cell physiology of a coagulase-negative staphylococcus in biofilm," *Journal of Proteome Research*, vol. 8, no. 4, pp. 1797–1809, 2009.
- [188] A. McLeod, L. Snipen, K. Naterstad, and L. Axelsson, "Global transcriptome response in *Lactobacillus sakei* during growth on ribose," *BMC Microbiology*, vol. 11, article no. 145, 2011.
- [189] H.-Q. Xu, L. Gao, Y.-S. Jiang et al., "Transcriptome response of *Lactobacillus sakei* to meat protein environment," *Journal of Basic Microbiology*, vol. 55, no. 4, pp. 490–499, 2015.



Hindawi

Submit your manuscripts at
<https://www.hindawi.com>

