

Original research

Open Access

The Utstein template for uniform reporting of data following major trauma: A joint revision by SCANTEM, TARN, DGU-TR and RITG

Kjetil G Ringdal*^{1,2}, Timothy J Coats³, Rolf Lefering⁴, Stefano Di Bartolomeo⁵, Petter Andreas Steen², Olav Røise⁶, Lauri Handolin⁷, Hans Morten Lossius¹ and Utstein TCD expert panel

Address: ¹Department of Research, Norwegian Air Ambulance Foundation, Drøbak, Norway, ²Faculty of Medicine, Faculty Division Ullevål University Hospital, University of Oslo, Norway, ³Academic Unit of Emergency Medicine, Leicester University, UK, ⁴Institute for Research in Operative Medicine, University of Witten/Herdecke, Cologne-Merheim Medical Centre, Cologne, Germany, ⁵Unit of Hygiene and Epidemiology, DPMSC, School of Medicine, University of Udine, Italy, ⁶Orthopaedic Centre, Ullevål University Hospital, Oslo, Norway and ⁷Department of Orthopaedics and Traumatology, Helsinki University Central Hospital, Finland

Email: Kjetil G Ringdal* - kjetil.ringdal@snla.no; Timothy J Coats - t.coats@virgin.net; Rolf Lefering - rolf.lefering@ifom-uni-wh.de; Stefano Di Bartolomeo - stefano.dibartolomeo@med.uniud.it; Petter Andreas Steen - p.a.steen@medisin.uio.no; Olav Røise - olav.roise@medisin.uio.no; Lauri Handolin - lauri.handolin@pp.inet.fi; Hans Morten Lossius - hans.morten.lossius@snla.no; Utstein TCD expert panel - kjetil.ringdal@snla.no

* Corresponding author

Published: 28 August 2008

Received: 19 June 2008

Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine 2008, **16**:7 doi:10.1186/1757-7241-16-7

Accepted: 28 August 2008

This article is available from: <http://www.sjtreem.com/content/16/1/7>

© 2008 Ringdal et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: In 1999, an Utstein Template for Uniform Reporting of Data following Major Trauma was published. Few papers have since been published based on that template, reflecting a lack of international consensus on its feasibility and use. The aim of the present revision was to further develop the Utstein Template, particularly with a major reduction in the number of core data variables and the addition of more precise definitions of data variables. In addition, we wanted to define a set of inclusion and exclusion criteria that will facilitate uniform comparison of trauma cases.

Methods: Over a ten-month period, selected experts from major European trauma registries and organisations carried out an Utstein consensus process based on a modified nominal group technique.

Results: The expert panel concluded that a New Injury Severity Score > 15 should be used as a single inclusion criterion, and five exclusion criteria were also selected. Thirty-five precisely defined core data variables were agreed upon, with further division into core data for Predictive models, System Characteristic Descriptors and for Process Mapping.

Conclusion: Through a structured consensus process, the Utstein Template for Uniform Reporting of Data following Major Trauma has been revised. This revision will enhance national and international comparisons of trauma systems, and will form the basis for improved prediction models in trauma care.

Background

The Utstein template for uniform reporting of data following major trauma

To permit data collection and statistics on major trauma care, in 1999 a working group from the International Trauma Anaesthesia and Critical Care Society (ITACCS) published a recommendation for the Utstein Template for Uniform Reporting of Data following Major Trauma [1]. The template extracted data for the pre-hospital phase, early in-hospital management, and for co-morbidity and outcome. In accordance with the previous Utstein templates, it was commended that data were to be classified as 'Core' (essential) or 'Optional' (supplemental). Despite the intention of facilitating studies to improve the understanding of trauma and trauma care, only a few papers have been published based on the template [2,3]. This indicates a need for further development, and in particular, a major reduction in the large number (92) of core data variables [1], as well as the addition of more precise definitions of these variables [3].

Trauma registries

Due to the practical difficulties with performing randomised controlled trials in severe trauma cases, valid scientific evidence is often lacking. Systematic prospective registry-based data collection for documenting trauma care is performed by several local, regional and national trauma registries. However, such registries cannot replace randomised clinical trials, but allow for exploration of relationships present in the collected data. The primary aims of these trauma registries are to enable comparative analyses of trauma care and outcome to provide quality improvement and optimal care of the injured patients [4]. The development of a European trauma registry may provide population-based comprehensive data on trauma incidence, epidemiology and trends. Further, it may enable development of regional outcome prediction models (taking special European factors into consideration) and thus set baseline norms for future trauma outcome studies. In Europe, there has been some reluctance to share local and national data, but it is recognised that lessons learned in one area of Europe may be useful for other European states [5]. However, when comparison is conducted, it is important to ensure that the reasons for differences in outcome are due to differences in the quality of trauma care or to differences in trauma systems, and not to variations in population characteristics [6].

TRISS methodology

Over the last two decades, the Trauma and Injury Severity Score (TRISS) method [7,8], with coefficients for prediction of outcome has been the most commonly used method for comparison of outcome in trauma patients. The TRISS coefficients were originally derived from the United States Major Trauma Outcome Study (US MTOS)

[9,10] but more recently the coefficients have been updated based on patient cases from the National Trauma Data Bank [11]. However, the TRISS method has some limitations, and it has been criticised by many authors [7,12-21]. Among other things, the TRISS model requires scoring the Revised Trauma Score (RTS) [22] components (Glasgow Coma Scale [GCS] [23], respiratory rate [RR] and systolic blood pressure [SBP]) on admission in the emergency department (ED), and does not take into account co-morbidity. Despite its limitations, TRISS continues to be the most accepted and widely-used tool for comparing trauma outcome in North America and in some parts of Europe.

Comparing and benchmarking European trauma care

In Europe, the UK Trauma Audit and Research Network (TARN) [24], along with the Trauma Registry of the German Society of Trauma Surgery (DGU-TR) [25], represent the largest trauma registries. There has also been a move towards developing a European Trauma Audit and Research Network (EuroTARN) [26], and a core dataset with inclusion and exclusion criteria has been created. Nevertheless, to date, no consensus has been reached between countries on the details and extent of the dataset. A first report from EuroTARN concluded that it is possible to collect data from established trauma registries, and the initial analysis revealed significant international variation [5]. As a continuation of this effort, a European project has been initiated by the DGU-TR, UK TARN and the Scandinavian Networking Group for Trauma and Emergency Management (SCANTEM) [27], for developing a joint European Core Dataset (EuroCoreD) for a future European Trauma Registry.

The 2007 revision of the Utstein template for uniform reporting of data following major trauma

Despite significant efforts [1,5], comparison of trauma care and outcome within Europe has not yet been carried out in a systematic way, mainly because inclusion criteria, data definitions and coding formats vary significantly between registries, and also because patient selection is not comparable [5,28]. Further efforts to establish uniform and standardised inclusion and exclusion criteria, as well as a minimum list of core data variables with precise definitions, are essential [3,5]. In addition, consistent methods of injury scoring need to be agreed upon [4,29-31]. To address this need for a European consensus, SCANTEM, TARN, DGU-TR and the Italian National Registry of Major Injuries (RITG) [32] carried out a consensus process, concluding with symposia in May and December 2007 at the Utstein Abbey [33], Norway. Selected experts met with the aim of further developing the Utstein Template for Uniform Reporting of Data following Major Trauma. At that time, they defined inclusion and exclusion criteria, and a minimum core dataset with precise

definitions. In addition, the aim of the revised template was to develop a standard for comparison of trauma data that was compatible with the large trauma registries in Europe, also adhering to EuroTARN. The template was intended to support the establishment of a European Trauma Registry, promote further development of a European model for outcome prediction and allow European and international trauma auditing and benchmarking.

Methods

This revision of the Utstein template is based on a nominal group technique (NGT) process [34,35] modified to fit the purpose. For participation in the NGT process, a European expert panel was selected.

The expert panel

The expert panel was comprised of those individuals who were central to developing and managing the largest European trauma registries; the panel included clinicians, database managers and epidemiology experts.

Data variable definition

A data variable should be unambiguously defined (with no misinterpretations) and reasonably simple to register. To meet this requirement, a data variable dictionary should contain information on 'data point number,' 'data point name,' 'descriptive field name,' 'type of data,' 'data point category/value,' 'definition of data point,' 'source of data information' and 'coding guidance.' We based recommended guidelines for data variable definitions on existing trauma registry databases, the Utstein Template for Uniform Reporting of Data following Major Trauma [1], the US National Trauma Data Standard (NTDS) [36] and the Injury Surveillance Guidelines from the World Health Organization (WHO) [37].

Core data variables

A registry should differentiate between data variables that absolutely need to be collected (core data) and the type of additional data that may be desirable (optional data) [1,37]. The current revision focuses on core data that are considered to be essential for documentation and reporting. We divided the core data into three groups ('Predictive Model,' 'System Characteristic Descriptors' and 'Process Mapping Variables') based on the role of the data variable in a registry.

Predictive model

The predictive model is composed of patient and injury severity variables that are considered to be important for outcome prediction. Predictive models are not deterministic; rather, they provide the probability of an outcome for a given patient [38]. Complex models, such as Abbreviated Injury Scale (AIS) [39] derivatives and the RTS, are often used to create such predictive models [38]. Experi-

ence from the German and UK trauma registries suggests that there may be better data variables to include in a predictive model than those traditionally used in the TRISS methodology [24,40-42].

System characteristic descriptors

Data variables in the System Characteristic Descriptor group describe trauma systems. Within Europe, there are large differences in philosophies and structures of trauma care systems, and these data variables should indicate key differences between systems and permit comparisons of the effect of system structure on outcomes.

Process mapping variables

Process mapping variables are intended to describe trauma care at an individual trauma centre (e.g., what happens to a patient after a major trauma); these are used for documentation of the patient journey, care process and care activities.

Specific premises

At present, many trauma registries have difficulty in obtaining data for patients from all involved hospitals when patients are transferred between them; therefore, the expert panel based their consensus on the premise that the core dataset was intended to cover the main hospital where a patient is treated. However, the expert group recommended that all trauma registries develop methods to track patients through the trauma system and that both the primary (local) trauma hospital and the referral trauma hospital record the same set of core data variables. The introduction of a core outcome data variable will secure that the overall effect of the entire trauma system can be measured, even if part of the patient's treatment course is not recorded in detail.

The nominal group technique

The modified NGT process consisted of four steps. First, each expert was supplied with necessary background documents (Table 1), and asked to return (by e-mail) proposals for inclusion and exclusion criteria, as well as a maximum of 30 core data variables in a prioritised order. This first proposal was summarised and structured by the coordinators (KGR, HML), and the collated results were redistributed in the second step for comments and re-prioritisation. The third step consisted of two consensus meetings in which members of the expert panel discussed their views in a structured way and then made conclusions. In the fourth step, the panellists were able to comment on the conclusions by e-mail. To complete the process, a letter of consent was signed by all experts.

Results

The expert panel concluded that a New Injury Severity Score [43] (NISS) > 15 should be used as a single inclu-

Table 1: Attachments sent to the expert panel prior to the Utstein 2007 meeting.

No.	Document name
1	Dick et al. Recommendations for uniform reporting of data following major trauma – the Utstein style [1].
2	Conclusions from the Utstein symposium on 'Improving Trauma Systems and the Role of Trauma Registries'.
3	Inclusion and exclusion criteria and data points from the European Trauma Audit & Research Network.
4	The Swedish Trauma Registry Standard (KVITTRA), Data Dictionary.
5	The Norwegian National Trauma Registry, Data Dictionary.
6	American College of Surgeons, National Trauma Data Bank; National Trauma Data Standard, Data Dictionary v. 1.2 [36].
7	ICD-10, Chapter XX. External causes of morbidity and mortality [61].

sion criterion (Table 2). Five exclusion criteria were listed (Table 2), and a total of 35 core data variables (23 in the predictive model group, eight system descriptors and four process mapping variables) were agreed upon (Tables 3, 4 and 5).

Discussion on inclusion/exclusion criteria and core data variables

Inclusion criteria

NISS is a modification of the Injury Severity Score (ISS) method [43]. ISS is calculated by summing the squares of the highest AIS severity codes in each of the three most severely injured ISS body regions [44]. Hence, ISS will ignore all but the most severe injury in a body region, and often fails to consider worse injuries in other regions of the body [43]. In contrast, NISS is defined as the sum of the square of the three most severe AIS injuries regardless of body region [43]. Several authors have argued for replacing ISS with NISS [43,45-49]. Osler et al. considered NISS to be easier to calculate and more predictive of survival than the ISS method [43], and a recent study by Lavoie et al. confirmed their findings [46]. NISS will be equal to or greater than ISS for any given patient, and it appears to be a more accurate method for rating severely injured patients [49,50]; specifically, this is true for patients with multiple head injuries [46]. The increased number of included patients by choosing NISS > 15 instead of ISS > 15 should be seen as an increase in 'sensi-

tivity' without a loss of 'specificity' of an ideal definition of major trauma. An effort should be made to secure that all patients with a NISS > 15 are included, regardless of whether or not the trauma team was activated prior to or upon the patient's arrival at the hospital, and whether or not the patient was admitted to an intensive care unit.

Exclusion criteria

Using NISS > 15 as a single inclusion criterion will include some patients that are at high risk of confounding data analysis. To remove such patients from the analysis, a set of exclusion criteria was defined. The expert panel recommended excluding first hospital admissions more than 24 hours after the injury (e.g., prolonged search and rescue missions), patients declared dead before hospital arrival, or those with no signs of life (pupillary response, spontaneous ventilation, presence of carotid pulse, measurable or palpable blood pressure, extremity movement, or cardiac electrical activity) [51] upon hospital arrival and those having no response to hospital resuscitation. In addition, it was recommended that asphyxias, drowning and burns should be excluded (Table 2).

Pre-hospital deaths should be excluded for practical reasons, since in some countries patients declared dead in the pre-hospital setting are transported directly to the morgue; whereas in other countries, they are admitted to hospital. All patients who arrive in the ED with spontane-

Table 2: Inclusion and exclusion criteria.

Inclusion criteria	NISS > 15.
Exclusion criteria	First hospital admission more than 24 hours after injury. Patients declared dead before hospital arrival, or with no signs of life on hospital arrival and no response to hospital resuscitation. Asphyxia. Drowning. Burn patients should be excluded if the burn represents the predominant injury, or if the patient is treated in a specialised burn unit.

NISS: New Injury Severity Score [43].

Signs of life: Pupillary response, spontaneous ventilation, presence of carotid pulse, measurable or palpable blood pressure, extremity movement, or cardiac electrical activity [51].

Table 3: Predictive model variables.

Data variable no.	Data variable name	Type of data	Data variable categories or values	Definition of data variable
1	Age	Continuous	Number	The patient's age at the time of injury.
2	Gender	Nominal	1 = Female 2 = Male 3 = Unknown	The patient's gender.
3	Dominating Type of Injury	Nominal	1 = Blunt 2 = Penetrating 3 = Unknown	Indication of the type of injury produced by the trauma.
4	Mechanism of Injury	Nominal	1 = Traffic: motor vehicle injury (car, pickup truck, van, heavy transport vehicle, bus) 2 = Traffic: motorcycle injury 3 = Traffic: bicycle injury 4 = Traffic: pedestrian 5 = Traffic: other (ship, airplane, railway train) 6 = Shot by handgun, shotgun, rifle, other firearm of any dimension 7 = Stabbed by knife, sword, dagger, other pointed or sharp object 8 = Struck or hit by blunt object (tree, tree branch, bar, stone, human body part, metal, other) 9 = Low energy fall (fall at the same level) 10 = High energy fall (fall from a higher level) 11 = Other 12 = Unknown	The mechanism (or external factor) that caused the injury event. The cut-off level for a fall should be defined as the person's height.
5	Intention of injury	Nominal	1 = Accident (unintentional) 2 = Self-inflicted (suspected suicide, incomplete suicide attempt, or injury attempt) 3 = Assault (suspected) 4 = Other 5 = Unknown	Information about the role of human intent in the occurrence of an injury, primarily determined by the incident and not by the resulting injury.
6	Pre-injury ASA-PS Classification System	Ordinal	1 = A normal healthy patient 2 = A patient with mild systemic disease 3 = A patient with severe systemic disease 4 = A patient with severe systemic disease that is a constant threat to life 5 = A moribund patient who is not expected to survive without the operation 6 = A declared brain-dead patient whose organs are being removed for donor purposes 7 = Unknown	The pre-injury co-morbidity existing before the incident. Derangements resulting from the injury should not be considered.
7	Pre-hospital cardiac arrest	Nominal	1 = No 2 = Yes 3 = Unknown	Did the patient suffer an injury-related pre-hospital cardiac arrest?

Table 3: Predictive model variables. (Continued)

8	Glasgow Coma Scale (GCS) upon arrival of EMS personnel at scene	Ordinal	Number	First recorded pre-interventional GCS upon arrival at scene of medical personnel trained to assess.
9	GCS motor component upon arrival of EMS personnel at scene	Ordinal	6 = Obeys commands/appropriate response to pain 5 = Localising pain 4 = Withdrawal from pain 3 = Flexion to pain (decorticate) 2 = Extension to pain (decerebrate) 1 = No motor response	First recorded pre-interventional GCS motor component upon arrival at scene of medical personnel trained to assess.
10	GCS upon arrival in ED/hospital	Ordinal	Number	First recorded GCS upon arrival in the ED/hospital.
11	GCS motor component upon arrival in ED/hospital	Ordinal	6 = Obeys commands/appropriate response to pain 5 = Localising pain 4 = Withdrawal from pain 3 = Flexion to pain (decorticate) 2 = Extension to pain (decerebrate) 1 = No motor response	First recorded GCS motor component upon arrival in the ED/hospital.
12a	Systolic Blood Pressure (SBP) upon arrival of EMS personnel at scene	Continuous	Number	First recorded SBP upon arrival at scene of medical personnel trained to assess.
12b	SBP – clinical category – upon arrival of EMS personnel at scene	Ordinal	RTS 4 = >89 ("good radial pulse") RTS 3 = 76–89 ("weak radial pulse") RTS 2 = 50–75 ("femoral pulse") RTS 1 = 1–49 ("only carotid pulse") RTS 0 = 0 ("no carotid pulse")	First recorded SBP upon arrival at scene of medical person trained to assess.
13a	SBP upon arrival in ED/hospital	Continuous	Number	First recorded SBP upon arrival in the ED/hospital.
13b	SBP – clinical category – upon arrival in ED/hospital	Ordinal	RTS 4 = >89 ("good radial pulse") RTS 3 = 76–89 ("weak radial pulse") RTS 2 = 50–75 ("femoral pulse") RTS 1 = 1–49 ("only carotid pulse") RTS 0 = 0 ("no carotid pulse")	First recorded SBP upon arrival in the ED/hospital.
14a	Respiratory Rate (RR) upon arrival of EMS personnel at scene	Continuous	Number	First recorded RR upon arrival at scene of medical personnel trained to assess.
14b	RR – clinical category – upon arrival of EMS personnel at scene	Ordinal	RTS 4 = 10–29 ("normal") RTS 3 = >29 ("fast") RTS 2 = 6–9 ("slow") RTS 1 = 1–5 ("gasp") RTS 0 = 0 ("no respiration")	First recorded RR upon arrival at scene of medical personnel trained to assess.
15a	RR upon arrival in ED/hospital	Continuous	Number	First recorded RR upon arrival in the ED/hospital.
15b	RR – clinical category – upon arrival in ED/hospital	Ordinal	RTS 4 = 10–29 ("normal") RTS 3 = >29 ("fast") RTS 2 = 6–9 ("slow") RTS 1 = 1–5 ("gasp") RTS 0 = 0 ("no respiration")	First recorded RR on arrival in the ED/hospital.

Table 3: Predictive model variables. (Continued)

16	Arterial Base Excess	Continuous	Number	First measured arterial base excess after arrival in the hospital.
17	Coagulation: INR	Continuous	Number	Use the first measured INR within the first hour after hospital arrival.
18	Number of days on ventilator	Continuous	Number	The total number of patient days spent on a mechanical ventilator (including all episodes). Record in full day increments with any partial day listed as a full day.
19	Length of stay in main hospital treating the patient	Continuous	Number	Calculate 'Date of discharge' minus 'Date of admission' from the reporting hospital.
20	Discharge destination	Nominal	1 = Home 2 = Rehabilitation 3 = Morgue 4 = Another CCU (higher treatment level) 5 = Another intermediate or low care somatic hospital ward 6 = Other 7 = Unknown	The patient's destination after end of acute care in the main hospital treating the patient. CCU = critical care unit.
21	Glasgow Outcome Scale – at discharge from main hospital	Ordinal	5 = Good Recovery 4 = Moderate Disability (Disabled but independent) 3 = Severe Disability (Conscious but disabled; depends upon others) 2 = Persistent vegetative state (unresponsive) 1 = Death 0 = Unknown	Glasgow Outcome Scale score at discharge from main hospital.
22	Survival status	Nominal	1 = Dead 2 = Alive 3 = Unknown	Alive or dead 30 days after injury.
23	Abbreviated Injury Scale (AIS)	Ordinal	Number	The AIS severity codes that reflect the patient's injuries. All injuries should be listed, even duplicated codes (e.g., bilateral femoral fractures, multiple spine fractures). The edition of the AIS coding dictionary should be indexed; AIS 2005 is recommended.

ASA-PS: American Society of Anesthesiologists Physical Status [65].

ED: Emergency Department.

EMS: Emergency Medical Services.

INR: International Normalized Ratio.

RTS: Revised Trauma Score [22].

Table 4: System characteristic descriptors.

Data variable no.	Data variable name	Type of data	Data variable categories or values	Definition of data variable
24	Time from alarm to hospital arrival	Continuous	HH:MM	The time between when the alarm call is answered (at the emergency call centre) and when the patient arrives at the reporting hospital.
25	Highest level of prehospital care provider	Ordinal	1 = Level I. No Field Care 2 = Level II. Basic Life Support 3 = Level III. Advanced Life Support, No Physician Present 4 = Level IV. Advanced Life Support On-Scene, Physician Field Care 5 = Other 6 = Unknown	The highest available level of competence of the pre-hospital care providers involved in the care of the injured patient.
26a	Pre-hospital intubation	Nominal	1 = No 2 = Yes 3 = Unknown	Was the patient intubated before arrival at the hospital?
26b	Pre-hospital intubation	Nominal	1 = A tube in the trachea (orotracheal, nasotracheal, or surgical airway) – drug assisted 2 = A supraglottic airway adjunct that prevents speech (such as esophago-tracheal combitube, the laryngeal tube, and various kinds of laryngeal masks)) – drug assisted 3 = A tube in the trachea (orotracheal, nasotracheal, or surgical airway) – not drug assisted 4 = A supraglottic airway adjunct that prevents speech (such as esophago-tracheal combitube, the laryngeal tube, and various kinds of laryngeal masks) – not drug assisted 5 = Other 6 = Unknown	Type of pre-hospital intubation. Drug assisted = anaesthesia, neuromuscular blocking drugs, and deep sedation.
27	Type of transportation	Nominal	1 = Ground ambulance 2 = Helicopter ambulance 3 = Fixed-wing ambulance 4 = Private/public vehicle 5 = Walk-in 6 = Police 7 = Other 8 = Unknown	Type of transportation delivering the patient to the hospital.

Table 4: System characteristic descriptors. (Continued)

28	Type of first key emergency intervention	Nominal	1 = Damage control thoracotomy – (any emergency or urgent thoracotomy performed for bleeding or suspected bleeding into the chest, but excluding simple thoracic tube drainage) 2 = Damage control laparotomy – (any emergency or urgent laparotomy performed for bleeding or suspected bleeding into the abdomen, including bleeding from the aorta) 3 = Extraperitoneal pelvic packing 4 = Limb revascularisation (Arterial injury necessitating vascular surgery or interventional radiology, including all interventions for pulseless limb, decreased perfusion and intimal arterial injuries) 5 = Interventional radiology (Angiographic embolisation; Stent; Stent-graft placement – excluding limb revascularisations which are classified as 4) 6 = Craniotomy 7 = Intracranial pressure (ICP) device insertion (excluding cases where the ICP device was inserted as part of a craniotomy which are classified as 6)	The first key emergency intervention performed for the treatment and stabilisation of the patient's injuries.
29	Activation of the trauma team	Nominal	1 = No 2 = Yes 3 = Unknown	Was the patient met by an activation of the trauma team prior to or upon arrival at the hospital?
30	Inter-hospital transfer	Nominal	1 = No 2 = Yes – Transferred IN to the reporting hospital 3 = Yes – Transfer OUT of the reporting hospital 4 = Yes – Transferred both IN and OUT of the reporting hospital 5 = Unknown	Was the patient transferred from/to another hospital for acute treatment?
31	Highest level of in-hospital care	Ordinal	1 = Emergency Department 2 = General Ward 3 = Operation Theatre 4 = High Dependency Unit 5 = Critical Care Unit (definition based on nurse to patient ratio) 6 = Unknown	The highest level of care in the main hospital.

ous circulation should be included, even if they have had a period of cardiac arrest before being admitted or if they die in the ED.

Asphyxia, drowning and burns are sufficiently different from blunt and penetrating injuries to require other datasets, and need to be considered separately. The UK National Burn Injury Database [52] is currently in use specifically for this purpose. Although the AIS 2005 edition has codes for asphyxia and drowning, such injuries were

not included in earlier AIS editions, making comparisons across versions more difficult. In some (but not all) countries, major burn patients are sent to dedicated burn unit hospitals, thereby confounding comparisons. Burn patients should be excluded if the burn represents the predominant injury, or if a patient is treated in a specialised burn unit. In such patients, outcome is determined by factors other than those suggested in this paper. Including burn patients will not represent a sufficient number of

Table 5: Process mapping variables.

Data variable no.	Data variable name	Type of data	Data variable categories or values	Definition of data variable
32	Time from alarm to arrival at scene	Continuous	HH:MM	The time from when the emergency call is answered (at the emergency call centre) until the first medical provider (at least the equivalent of EMT's) arrives at the patient.
33	Time until normal arterial base excess	Continuous	HH:MM	The time from first measured arterial base excess at hospital admission until first measured arterial base excess within normal range. Reference range for base excess: ± 3 mmol/l.
34	Time to first CT scan	Continuous	HH:MM	The time from hospital admission until the time marked on the first CT scan image.
35	Time until first key emergency interventions	Continuous	HH:MM	The time from hospital admission until the FIRST emergency intervention. Record the time from hospital admission until the time of FIRST knife to skin is performed. Consider only the emergency interventions listed in data variable number 28.

CT: Computed Tomography.
EMT: Emergency Medical Technician.

patients to report on; hence, burn-related injuries will add little power to the predictive model.

Predictive model variables

Age is an independent predictor of survival after trauma [53,54]. While the original TRISS model operates with only two age categories, current predictive models utilise different age groups, and we therefore recommend reporting the patient's nominal age (continuous) at the time of injury, in years without decimals, and always rounding down. Patients under one year of age should be reported with one decimal number (e.g., six months is 0.5).

Gender is recommended as a core data, since some studies have reported no association between gender and mortality after traumatic injury [55]; whereas others have found age-specific associations between male gender and outcome [56-58].

An evaluation of type of injury (blunt versus penetrating trauma) is useful for determining which patients are candidates for surgical haemostasis [59], and is essential in the TRISS model [8]. The previous Utstein document recommended that for cases involving both blunt and penetrating injuries, the predominant type of injury should be recorded [1]. The expert panel defined the dominating injury as the one with the highest AIS score. In the rare event of a patient having both blunt and penetrating traumas with the same AIS severity score, penetrating trauma is defined as the predominant injury.

The significance of the mechanism of injury (MOI) in prediction of trauma and outcome is, to a large extent, undetermined [60]. The MOI should be of value for epidemiology or subgroup analysis, and should be described in categories with reasonable prevalence rates. The International Classification of Diseases, 10th revision (ICD-10) [61], chapter XX, External causes of morbidity and mortality (V01-Y98), was initially examined for the purpose of the template; however, it was found to be too detailed, with too many injury codes. Therefore, the expert panel developed a reduced set of categories, which should make data collection easier. The set still enables the analysis of important subgroups, and since it is compatible with the ICD-10 codes, it will allow future category expansion if required.

In the ICD, most injuries can be grouped into two dimensions: intent and mechanism [62]. 'Intention of injury' provides information about the role of the human intent of an injury. The included list of categories is based on the ICD-10 codes, and is selected by the expert panel since it covers most injury intentions.

The presence of significant co-morbidity represents an independent predictor of mortality after trauma [1,53,63,64], and the expert panel recommends employing the American Society of Anaesthesiologists Physical Status (ASA-PS) classification system [65] for classifying the pre-injury co-morbidity status concretised by selected examples from the Norwegian Society of Anaesthesiology

[66] (Table 6). The ASA-PS classification system is easy to use and employed daily by anaesthesiologists world-wide. The pre-injury ASA-PS classification system should be used solely to categorise co-morbidity that exists before injury [6]. Skaga et al. found this classification system to be a significant predictor of mortality after trauma, also when adjusting for the variables in the traditional TRISS model [6]. Physiologic derangement resulting from the present injury is not reflected in the pre-injury score, only co-morbidity existing before the incident [6]. The pre-injury ASA-PS is not identical to the preoperative or the post-injury ASA score from anaesthesia records.

Pre-hospital cardiac arrest after trauma has dismal outcomes [67-70]; it was therefore chosen as a core data variable.

The panel recommends using raw values (continuous data) of the GCS, SBP, RR, where they are obtainable, and coded values (clinical categories) according to the RTS [22] (Table 7) in those cases missing raw values. In addition, the expert panel recommends recording the first pre-hospital, pre-interventional GCS, SBP and RR collected by a trained person, as well as the first GCS, SBP and RR upon arrival at the hospital.

For patients intubated before arrival at the hospital, the ED GCS is not attainable, and this could potentially result in large amounts of missing data [18]. For this reasons, the pre-hospital, pre-interventional GCS should be recorded, thus permitting its inclusion in future prediction models. Some studies have shown that the pre-hospital GCS score correlate well with the arrival GCS score [71,72], and that the pre-hospital GCS score is a strong predictor of outcome [73]. A change in the GCS from the field to hospital arrival is highly predictive of outcome [72,74]. In the RTS, GCS carries the greatest weight [22], and the motor component of GCS appears to contain most of the predictive power of the total GCS score [75]. This component needs further assessment, something that may be facilitated by reporting both the pre-hospital pre-interventional GCS motor component as well as the first recorded GCS motor component upon arrival in the ED/hospital.

Chan et al. found that even in the case of normal SBP upon arrival in the ED, out-of-hospital hypotension was a clinical predictor of severe injury [76]; it was also a strong predictor of the need for emergency surgery, according to Lipsky et al. [77]. Others have documented an increased risk of intra-abdominal injury in blunt trauma patients with hypotension in the hospital [78,79]. Franklin et al.

Table 6: American Society of Anesthesiology Physical Status (ASA-PS) Classification System.

ASA-PS 1	A normal healthy patient.
Guidelines:	No organic, physiologic, biochemical or psychiatric disturbance. Any disorder is localized, without systemic effects. Smoking <5 cigarettes/day.
Example:	Healthy non-smoker, admitted for varicose vein operation.
ASA-PS 2	A patient with mild systemic disease.
Guidelines:	Present pathology might imply specific measures or anaesthesia related precautions. The disturbance(s) might be caused by the condition to be surgically treated or by another pathologic process. Smoking >5 cigarettes/day.
Examples:	Mild organic heart disease. Uncomplicated diabetes mellitus (type 1 or 2). Benign hypertension without complications. Healthy patient with trismus.
ASA-PS 3	A patient with severe systemic disease.
Examples:	Diabetes mellitus with organ complications. Disabling heart disease. Moderate to severe respiratory disease. Angina pectoris. Myocardial infarction >6 months ago.
ASA-PS 4	A patient with severe systemic disease that is a constant threat to life.
Guidelines:	The disease is not necessarily related to the condition to be surgically treated, neither is it necessarily improved by the surgical intervention per se.
Examples:	Malignant hypertension. Myocardial infarction <6 months ago. Severe liver, kidney, respiratory, or endocrine dysfunction. Manifest cardiac failure. Unstable angina pectoris. Subarachnoid haemorrhage – patient awake or somnolent.
ASA-PS 5	A moribund patient who is not expected to survive without the operation.
Examples:	Patient in circulatory shock because of ruptured aortic aneurysm. Deeply comatose patient with intracranial haemorrhage.
ASA-PS 6	A declared brain-dead patient whose organs are being removed for donor purposes.

The six ASA-PS headings are from the American Society of Anaesthesiologists [65]. The guidelines and examples were translated from the Norwegian edition [66] by Skaga et al. [6].

For the Utstein Template, the ASA-PS classification system should solely be used to categorise pre-injury comorbidity. Derangements resulting from the injury should not be considered.

Table 7: Revised Trauma Score (RTS) categories with clinical notes.

RTS coded values	Respiratory Rate	Systolic Blood Pressure	Glasgow Coma Scale score
4	10–29 ("normal")	>89 ("good radial pulse")	13–15
3	>29 ("fast")	76–89 ("weak radial pulse")	9–12
2	6–9 ("slow")	50–75 ("femoral pulse")	6–8
1	1–5 ("gasp")	1–49 ("only carotid pulse")	4–5
0	0 ("no respiration")	0 ("no carotid pulse")	3

This table is based on (but not identical to) the RTS table in reference [22]. The parentheses represent clinical notes that were added by the expert panel.

reported 16% mortality in patients with a period of pre-hospital hypotension but with stable vital signs on ED presentation, vs. 27% mortality for patients with normal pre-hospital SBP who developed hypotension in the ED [80]. Changes in the SBP values, from the pre-hospital pre-interventional value to the SBP value on hospital arrival, could give valuable information about physiologic derangements.

Missing RR values for patients arriving in the hospital (patients intubated before arrival) is the most common cause of a lack of RTS values [81]. Collecting the pre-hospital, pre-interventional RR as well as allowing the use of coded values instead of precise values may compensate for this [18]. In addition, continued documentation of RR for later determination of its predictive power is highly recommended. Creating alternative predictive models may also necessitate the continuous collection of RR values.

Admission arterial base excess (BE) is a predictor of mortality after trauma [40,82-84]; it is also considered an indicator of haemodynamic instability, high transfusion requirement, and an indicator of metabolic and coagulatory decompensation in trauma patients [40,85]. Other results indicate that arterial BE is a predictor of intra-abdominal injury [78,79], and the European guidelines for management of bleeding following major trauma have recommended arterial base deficit as a sensitive test to estimate and monitor the extent of bleeding and shock [59]. Kroezen et al. showed that replacing RTS by base deficit as a measure of physiological disturbance could predict mortality as well as RTS in the TRISS model [86]. The expert panel recommends recording the first arterial BE after arrival in the hospital.

There is a high frequency of established coagulopathy in multiple-injured patients upon their arrival in the ED [87,88]. Coagulation abnormalities presenting early after trauma have been found to be an independent predictor

of mortality [87-90], and the prognostic value of measuring coagulation parameters upon ED admission has been documented [89]. The first measured International Normalized Ratio (INR), obtained within the first hour after hospital arrival, was chosen as a core data for measuring coagulation abnormalities.

The term 'intensive care unit' (ICU) is variously defined across the world, and in some referral hospitals, many patients are intubated upon arrival or transferred back to a local hospital while still intubated and ventilated. This makes the calculation of length of stay (LOS) in the ICU difficult. The revised template recommends recording the total number of days spent on a mechanical ventilator, rather than ICU LOS, as a measure of resource consumption. Furthermore, these data permit the use of the concept of ventilator-free days to quantify morbidity after trauma.

As outcome measures, the expert panel suggest hospital LOS, discharge destination, Glasgow Outcome Scale (GOS) [91] score at hospital discharge and 30-day mortality. The GOS contains information about morbidity, and this data collection represents an increased focus on information regarding morbidity. In the original US MTOS, "end of acute care" was used as an outcome measure [9]; however, 30-day mortality is considered to be a more fixed endpoint than "end of acute care/hospital discharge" or "end of somatic care," which will vary depending on the transfer and rehabilitation policies of an individual system [92]. Using 30-day mortality is consistent with the previous Utstein recommendations [1]. Death occurring later than 30 days after injury is more likely to be caused by other conditions, such as pre-existing disease [81]. An analysis of 69,650 patient admissions from the TARN showed that 4.8% of the patients died within 93 days of admission [41]. Of these, only 9% died later than 30 days after admission; these were mainly patients with a low ISS (< 9) and aged > 65 years. In a Scandinavian trauma registry, 4.6% of the deaths occurred

later than 30 days after injury [92], whilst data from the German trauma registry (cases with NISS ≥ 16) indicate that 4.9% of the patients that died did so later than 30 days after injury [personal communication with the DGU-TR]. Survival status of all patients at a single given endpoint is needed, and 30-day mortality was chosen to represent this information.

For injury severity description, the expert panel recommends recording all AIS codes. It has previously been shown that a comparison of survival for trauma registries that use different AIS editions is potentially possible [93]. A difference in registrars and different levels of AIS training probably represents a greater problem than problems with using different AIS versions. However, registries are recommended to use the same AIS edition as a uniform way of coding; hence, the newest available version should be used at all times. At present, this is the AIS 2005 edition.

System characteristics descriptors

The expert panel considered the time between when the alarm call is answered (at the emergency dispatch centre) and when the patient arrives at the first hospital to be an important system characteristic descriptor (this variable can also be useful for mapping the entire process of pre-hospital rescue) and recommend that this interval be reported.

Recording the highest level of competence of the pre-hospital care providers was regarded by the expert panel as an important measure for describing the pre-hospital system, and for cross-border comparisons of pre-hospital trauma care and outcome. The highest available level of the providers may vary somewhat in Europe, but the revised template's categorisation of level of provider is based on the levels proposed by McSwain [94], since these levels will encompass most pre-hospital systems.

Pre-hospital intubation is an important parameter that represents pre-hospital advanced life support (ALS) [95]. There is an ongoing debate on the use and role of ALS [96-98] measures in the out-patient management of trauma victims, and such information should be made available for possible inclusion in future survival prediction models. The original US TRISS model was derived from a dataset that excluded trauma patients who were intubated out-of-hospital [9]. However, these patients constitute a significant proportion of European trauma victims; hence, the expert panel recommends including them. In a study by Arbabi et al., early field intubation was associated with a decreased risk of fatal outcome, as compared to ED intubation [71]. In their study, intubation status was also an independent predictor of fatal outcome, after adjusting for ISS, SBP, mechanism, age and ED-GCS [71].

The type of transportation delivering the patient to the hospital is an important descriptor of the pre-hospital trauma care system, but the use of a helicopter vs. ground ambulance remains controversial [99,100], and assessment is confounded by differences in various EMS and HEMS systems. Since transportation type has been so widely debated, this is recommended as core data to be collected. The data variable does not cover what is a commonly used transport combination in some parts of Europe; ground ambulance to the local hospital and fixed-wing or rotary-wing transfer to the regional trauma centre. The data variable was developed to cover the type of transportation delivering the patient to the reporting hospital.

The expert panel recommends registering the initial key emergency intervention (EI) conducted during the hospital stay (ED, OR, critical care unit). These interventions (Table 8) represent essential emergency procedures used for the treatment and stabilisation of patients with severe injuries. Some registries will probably collect finer resolution data on interventions, but most patients will fit into one of these categories. It is recommended that the EI found in the present categories be recorded, even if there is no proof in a patient's notes that the cause of intervention was bleeding. The term "damage control", as used in some categories, implies that only the urgent (rather than later planned procedures) should be recorded. All limb

Table 8: Key emergency interventions.

No.	Emergency interventions
1	Damage control thoracotomy – (any emergency or urgent thoracotomy performed for bleeding or suspected bleeding into the chest, but excluding simple thoracic tube drainage)
2	Damage control laparotomy – (any emergency or urgent laparotomy performed for bleeding or suspected bleeding into the abdomen, including bleeding from the aorta)
3	Extraperitoneal pelvic packing
4	Limb revascularisation (Arterial injury necessitating vascular surgery or interventional radiology, including all interventions for pulseless limb, decreased perfusion and intimal arterial injuries)
5	Interventional radiology (Angiographic embolisation; Stent; Stent-graft placement – excluding limb revascularisations which are classified as No. 4)
6	Craniotomy
7	Intracranial pressure (ICP) device insertion (excluding cases where the ICP device was inserted as part of a craniotomy, which are classified as No. 6)

revascularisations should be categorised in the same group (whether surgical or radiological) since it is the fact that the patient needed revascularisation (rather than the exact method used) that is important. If radiological revascularisations are put together with all the rest of the interventional radiology, the frequency of revascularisations undertaken in a particular system will not be easily measurable. Information about the surgical revascularisations will not permit comparison of the need for revascularisation across systems. The need for revascularisation is a more interesting comparison than comparing the way in which revascularisations were achieved. Extraperitoneal pelvic packing for control of massive traumatic pelvic haemorrhage has been described by several authors [101-103], and is a separate EI subcategory. Rapid intervention is essential for patients with intracerebral haematomas that require evacuation; for patients requiring only intracranial pressure (ICP) monitoring as part of the overall intensive care management of diffuse brain injuries [104], however, this intervention would not have the same level of urgency. Nevertheless, insertion of an ICP monitoring device will provide information about neurosurgical alertness in a trauma system, and is therefore included. The suggested list of EI was developed to cover the broad majority of emergency interventions (but perhaps not all of them). For the purpose of comparative evaluation of the acute treatment process, some less common types of intervention may not be appropriate.

A formal trauma team is an essential part of an organised trauma system [104]. Activation of the trauma team is an organised initial response to a trauma [104,105] with the primary goal of securing fast and efficient treatment of severely injured patients [106-108]. Information on whether the trauma patient was met by an activation of the trauma team was recommended as vital for describing a trauma system.

The revised template recommends recording the highest level of in-hospital care for the trauma patient in the main trauma hospital (Table 4).

Process mapping variables

The "Chain of Survival" concept emphasises that all time-sensitive interventions must be optimised to maximise the chance of patient survival [109]. The expert panel recommends recording the time between when the emergency call is answered (at the emergency call centre) and when the first medical provider arrives at the patient. This core data represent parts of the first link in that chain, and is an important measure of the quality of the pre-hospital EMS system.

As an overall marker of the efficiency of patient treatment (including resuscitation, diagnostics and surgery), the

expert group suggested to consider the time required to achieve normal arterial BE. A good evidence basis does not exist for this recommendation, however, if BE worsens after arrival, prognosis worsens as well [110]. The BE should be measured regularly after hospital arrival, and the template advises to document the initial measurement of BE immediately after hospital arrival, and in cases of abnormal values, to document the time in hours until normalisation.

The efficiency of the initial in-hospital management is assessed by the time from hospital admission until the time marked on the first CT scan images. This data represents the time required to perform key in-hospital diagnostic tests.

The time to the first key EI should be recorded; this measure represents how quickly an urgent intervention fundamental to the treatment and stabilisation of a patient's specific injuries [105] is performed. This core data measures the efficiency of the trauma system in the initial phase. The time elapsed between injury and EI should be minimised.

General discussion

The present paper represents a further development of the previous Utstein Template for Uniform Reporting of Data following Major Trauma [1], and reflects a need for the creation of new prediction models that are more suitable for the type of trauma seen in Europe. Currently, both the German and UK trauma registries have developed their own predictive models, and no longer use the original TRISS system. However, uniform and standardised inclusion and exclusion criteria and a core list of data variables with precise definitions are mandatory before comparisons between trauma registries can be made. The expert panel reached a consensus on such a list of core data. This dataset does not preclude the possibility that in local trauma registries more data can legitimately be considered core information for specific purposes. The data agreed upon in the present process represent data for admissions to the first hospital within 24 hours after injury. We are aware that some countries have made further progress in developing national trauma systems and national trauma registries; whereas others only register information at local trauma hospitals. Currently, the core dataset will be difficult to use to assess the entire trauma system of some regions or countries, making direct system comparisons difficult. At this point, the expert panel recommends focusing on the hospitals of definitive treatment, and registering patient transfers (as a separate data variable) should allow for this when data are analysed. The inclusion of 30-day mortality in predictive models allows for the assessment of more of the total system performance. However, excluding pre-hospital deaths and the details of

management along the chain that precede the main treatment hospital represents a major limitation in total system assessment. The core dataset should undergo further development to allow for routine tracking of the patient through a trauma system consisting of more than one hospital. We anticipate that future development of the Utstein Template will focus more on the trauma system as European trauma registries develop, making it possible to follow transferred patients. In future trauma systems, the same core data variables should be recorded in primary, secondary and tertiary trauma centres.

The template classifies fall-related injuries as low or high (category options of the MOI data variable in Table 4) with a person's height as the cut-off value. Various definitions of falls are used internationally [61,111-115]; they range from different categories of height, different cut-off levels for height, to differing units of height measurement (feet vs. meters). This is an area in which there is a lack of uniformity; therefore, for future development, we suggest that individual registries should also record the actual estimated height of a fall in meters (as a continuous variable), so that an analysis can be performed.

We are aware that the best way of assessing the GOS as a measure of disability (morbidity), is after at least six months, but since this was not considered feasible, registration at discharge from the hospital was chosen as the endpoint. The assessment of GOS as an outcome measure at discharge from a trauma centre in trauma systems based on early discharge to specialised rehabilitation services represent a limitation, and the GOS endpoint should therefore be interpreted with caution. Nevertheless, assessment of this endpoint data can be used as a rough estimate of the amount of care needed for a patient beyond the acute hospital stay. Although GOS was developed for patients with head injuries, it represents a rough disability outcome score, and as such it should be possible to use for assessing all trauma patients.

The revised Utstein Template considers time to normalised arterial base excess as a measurement of importance for evaluating the total quality of trauma treatment. This data is not a precisely timed measurement and will probably vary from hospital to hospital. However, collecting it is considered valuable and might be important for future comparisons. The inclusion of this data also has some educational reasons; the trauma centres should be encouraged to include this value regularly in the trauma patients' charts.

The expert panel recommends reporting the first EI, and the time to first EI, disregarding the fact that some patients may have received several of the listed interventions. The first intervention is, by clinical judgement, the most

important one for comparison. This does not preclude trauma registries from documenting each of these interventions separately, or even each operation that is performed. However, the type of first EI and the time elapsed prior to its application is of significant importance; for this reason, this measure was chosen as part of the core dataset. For the purpose of comparison, focusing on the first EI performed in the current system will be adequate. However, if comparisons based on all EI are desired, the present recommendation will not be adequate. Comparison of systems based on all interventions can only be used if it is coupled with data on the time of each intervention. This will be difficult to use if the order in which the interventions occur is not known. If the first intervention is not among those listed under the EIs in the template, time to intervention is not important in our context (i.e., there will be enough time for thorough surgical preparation), and it should therefore not be part of the Utstein core dataset.

The panel acknowledges that no indicator has 100% validity and careful judgement is therefore needed. For example, in some selected cases earlier emergency surgery may imply both later CT imaging and better practice. Definition of data variables is a complex and ongoing process and in order to widen the implementation of the core dataset, facilitate participation in European trauma audit and comparisons and increase the quality of the next updates of the core dataset, we encourage all readers to ask for clarifications and point out potential improvements.

One concern with the Utstein Template is implementation, since none of the participants from the previous Utstein trauma process changed their registries to accommodate the recommendations of that meeting. The solution for the revised template was to develop a letter of consent that all members of the expert panel signed where they agreed on implementing the core data in the revised Utstein Template [see Additional file 1]. To further facilitate implementation, a complete user manual with core data variables and variable definitions will be available free of charge, at: <http://www.scantem.org>, <https://www.tarn.ac.uk>, <http://www.traumaregister.de/de/index.htm>, <http://www.pprg.infoteca.it/ritg>.

Conclusion

This paper represents a major step in perfecting the Utstein Template for Uniform Reporting of Data following Major Trauma, making the core data variables more uniform and applicable. Collecting this core dataset should be a basic component of all future studies on trauma care, and a uniform dataset such as this, will facilitate accurate description of the patient population and allow comparisons of outcome from trauma systems. It is extremely important that the data variables are collected

in a uniform manner. For this reason, each variable and response category has been specifically defined in a way that is designed to promote the collection and reporting of a comparable core dataset. A letter of consent has been signed by the expert panel, where the participants of this consensus process agreed to implement the inclusion and exclusion criteria and core data variables in their respective systems and registries.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KGR and HML designed the study and organised the consensus meetings. TJC was the chairman of the consensus process, while LH and OR were co-chairmen. The core dataset was developed by the expert panel. KGR, TJC, RL, SDB, OR, LH, PAS and HML wrote the article. The rest of the panellists read the manuscript once, made comments to it and approved the manuscript. All authors read and approved the final version of the manuscript.

Additional material

Additional file 1

Letter of Consent. The letter provided was signed by the Utstein TCD expert panel. With this letter, the expert panel members confirm that they will implement the core data agreed upon.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1757-7241-16-7-S1.pdf>]

Acknowledgements

TCD = "Trauma Core Data".

The members of the Utstein TCD expert panel were as follows:

Stefano Di Bartolomeo (Italy); Maaret Castrén (Sweden); Erika F. Christensen (Denmark); Timothy J. Coats (UK); Antoinette Edwards (UK); Torsten Eken (Norway); Ernestina Gomes (Portugal); Lauri Handolin (Finland); Morten Hestnes (Norway); Lena Klarin (Sweden); Morten Schultz Larsen (Denmark); Rolf Lefering (Germany); Ari Leppäniemi (Finland); Hans Morten Lossius (Norway); Per Örténwall (Sweden); Olav Røise (Norway); Nils Oddvar Skaga (Norway); Torben Wisborg (Norway); Maralyn Woodford (UK).

We acknowledge the Working Party of the ITACCS for developing the first Utstein Template, as well as the previous work of the EuroTARN group. We also wish to thank Jens Lauritsen MD, PhD, Accident Analysis Group, Odense University Hospital, Odense, Denmark and Professor Eldar Søreide, Departments of Emergency and Intensive Care Medicine, Division of Acute Care Medicine, Stavanger University Hospital, Stavanger, Norway for their contributions and support to this project.

Funding: The consensus process was funded by the Norwegian Air Ambulance Foundation, Drøbak, Norway (KGR, PhD scholarship, and expenses

for expert panel meetings), and the Laerdal Foundation for Acute Medicine, Stavanger, Norway (expenses for expert panel meetings). The sponsors did not influence the final decision to submit for publication, nor did they have any role in the writing process, other than facilitating the meetings.

References

- Dick WF, Baskett PJ: **Recommendations for uniform reporting of data following major trauma – the Utstein style. A report of a working party of the International Trauma Anaesthesia and Critical Care Society (ITACCS).** *Resuscitation* 1999, **42**:81-100.
- Di Bartolomeo S, Sanson G, Michelutto V, Nardi G, Burba I, Francescutti A, Lattuada L, Scian F: **Epidemiology of major injury in the population of Friuli Venezia Giulia-Italy.** *Injury* 2004, **35**:391-400.
- Lossius HM, Langhelle A, Søreide E, Pålgråm-Larsen J, Lossius TA, Laake P, Steen PA: **Reporting data following major trauma and analysing factors associated with outcome using the new Utstein style recommendations.** *Resuscitation* 2001, **50**:263-272.
- Mann NC, Guice K, Cassidy L, Wright D, Koury J: **Are statewide trauma registries comparable? Reaching for a national trauma dataset.** *Acad Emerg Med* 2006, **13**:946-953.
- The EuroTARN Writing Committee on behalf of the EuroTARN Group: **A comparison of European Trauma Registries The first report from the EuroTARN Group.** *Resuscitation* 2007, **75**:286-297.
- Skaga NO, Eken T, Sovik S, Jones JM, Steen PA: **Pre-injury ASA physical status classification is an independent predictor of mortality after trauma.** *J Trauma* 2007, **63**:972-978.
- Champion HR, Sacco WJ, Hunt TK: **Trauma severity scoring to predict mortality.** *World J Surg* 1983, **7**:4-11.
- Boyd CR, Tolson MA, Copes WS: **Evaluating trauma care: the TRISS method. Trauma Score and the Injury Severity Score.** *J Trauma* 1987, **27**:370-378.
- Champion HR, Copes WS, Sacco WJ, Lawnick MM, Keast SL, Bain LW Jr, Flanagan ME, Frey CF: **The Major Trauma Outcome Study: Establishing national norms for trauma care.** *J Trauma* 1990, **30**:1356-1365.
- Champion HR, Sacco WJ, Copes WS: **Injury severity scoring again.** *J Trauma* 1995, **38**:94-95.
- National Trauma Data Bank: **Reference Manual: Background, caveats, and resources.** [<http://www.facs.org/trauma/ntdbmanual.pdf>].
- Champion HR, Copes WS, Sacco WJ, Frey CF, Holcroft JW, Hoyt DB, Weigelt JA: **Improved predictions from a severity characterization of trauma (ASCOT) over Trauma and Injury Severity Score (TRISS): results of an independent evaluation.** *J Trauma* 1996, **40**:42-48. discussion 48-49
- Demetriades D, Chan L, Velmanos GV, Sava J, Preston C, Gruzinski G, Berne TV: **TRISS methodology: an inappropriate tool for comparing outcomes between trauma centers.** *J Am Coll Surg* 2001, **193**:250-254.
- Norris R, Woods R, Harbrecht B, Fabian T, Rhodes M, Morris J, Billiar TR, Courcoulas AP, Udekwu AO, Stinson C, Peitzman AB: **TRISS unexpected survivors: an outdated standard?** *J Trauma* 2002, **52**:229-234.
- Osler TM, Cohen M, Rogers FB, Camp L, Rutledge R, Shackford SR: **Trauma registry injury coding is superfluous: a comparison of outcome prediction based on trauma registry International Classification of Diseases-Ninth Revision (ICD-9) and hospital information system ICD-9 codes.** *J Trauma* 1997, **43**:253-256. discussion 256-257
- Rutledge R, Osler T, Emery S, Kromhout-Schiro S: **The end of the Injury Severity Score (ISS) and the Trauma and Injury Severity Score (TRISS): ICISS, an International Classification of Diseases, ninth revision-based prediction tool, outperforms both ISS and TRISS as predictors of trauma patient survival, hospital charges, and hospital length of stay.** *J Trauma* 1998, **44**:41-49.
- Demetriades D, Chan LS, Velmahos G, Berne TV, Cornwell EE 3rd, Belzberg H, Asensio JA, Murray J, Berne J, Shoemaker W: **TRISS methodology in trauma: the need for alternatives.** *Br J Surg* 1998, **85**:379-384.
- Skaga NO, Eken T, Steen PA: **Assessing quality of care in a trauma referral center: benchmarking performance by**

- TRISS-based statistics or by analysis of stratified ISS data?** *J Trauma* 2006, **60**:538-547.
19. Jones JM, Redmond AD, Templeton J: **Uses and abuses of statistical models for evaluating trauma care.** *J Trauma* 1995, **38**:89-93.
 20. Younge PA, Coats TJ, Gurney D, Kirk CJ: **Interpretation of the Ws statistic: application to an integrated trauma system.** *J Trauma* 1997, **43**:511-515.
 21. Hollis S, Yates DW, Woodford M, Foster P: **Standardized comparison of performance indicators in trauma: a new approach to case-mix variation.** *J Trauma* 1995, **38**:763-766.
 22. Champion HR, Sacco WJ, Copes WS, Gann DS, Gennarelli TA, Flanagan ME: **A Revision of the Trauma Score.** *J Trauma* 1989, **29**:623-629.
 23. Teasdale G, Jennett B: **Assessment of coma and impaired consciousness. A practical scale.** *Lancet* 1974, **2**:81-84.
 24. Trauma Audit and Research Network: **TARN.** [<https://www.tarn.ac.uk/>].
 25. Trauma Registry of the German Society of Trauma Surgery: **Traumaregister der Deutsche Gesellschaft für Unfallchirurgie.** [<http://www.traumaregister.de/de/index.htm>].
 26. European Trauma Audit & Research Network: **EuroTARN.** [<http://eurotarn.man.ac.uk/>].
 27. SCANTEM: **Scandinavian Networking Group for Trauma and Emergency Management.** [<http://www.scantem.org/>].
 28. Ringdal KG, Lossius HM, SCANTEM ad hoc group on Scandinavian MTOS and Trauma Registry: **Feasibility of comparing core data from existing trauma registries in Scandinavia. Reaching for a Scandinavian major trauma outcome study (MTOS).** *Scand J Surg* 2007, **96**:325-331.
 29. Jurkovich GJ, Mock C: **Systematic review of trauma system effectiveness based on registry comparisons.** *J Trauma* 1999, **47**:546-55.
 30. Owen JL, Bolenbaucher RM, Moore ML: **Trauma Registry Databases: A Comparison of Data Abstraction, Interpretation, and Entry at Two Level I Trauma Centers.** *J Trauma* 1999, **46**:1100-1104.
 31. Pollock DA, McClain PW: **Trauma registries. Current status and future prospects.** *JAMA* 1989, **262**:2280-2283.
 32. Italian National Registry of Major Injuries: **Registro Intraospedaliere Multiregionale Traumi Gravi.** [<http://www.pprg.infoteca.it/ritg/>].
 33. Nolan J, Soar J: **Images in Resuscitation: Utstein Abbey.** *Resuscitation* 2005, **64**:5-6.
 34. Ven A Van de, Delbecq A: **The nominal group as a research instrument for exploratory health studies.** *Am J Public Health* 1972, **62**:337-342.
 35. Delbecq A, Ven A Van de: **A group process model for problem identification and program planning.** *J Appl Behav Sci* 1971, **7**:467-492.
 36. American College of Surgeons: **National Trauma Data Bank; National Trauma Data Standard, Data Dictionary v. 1.2.** [<http://www.ntsdictionary.org>].
 37. Holder Y, Peden M, Krug E, Lund J, Gururaj G, Kobusingye O: **Injury Surveillance Guidelines.** In *Book Injury Surveillance Guidelines* (Editor ed. eds.) City: World Health Organization; 2001.
 38. Champion HR: **Trauma scoring.** *Scand J Surg* 2002, **91**:12-22.
 39. Association for the Advancement of Automotive Medicine: **The Abbreviated Injury Scale 2005** Barrington, IL: Association for the Advancement of Automotive Medicine; 2005.
 40. Rixen D, Raum M, Bouillon B, Lefering R, Neugebauer E: **Base deficit development and its prognostic significance in posttrauma critical illness: an analysis by the trauma registry of the Deutsche Gesellschaft für Unfallchirurgie.** *Shock* 2001, **15**:83-89.
 41. Bouamra O, Wrotchford A, Hollis S, Vail A, Woodford M, Lecky F: **A new approach to outcome prediction in trauma: A comparison with the TRISS model.** *J Trauma* 2006, **61**:701-710.
 42. Ruchholtz S, Lefering R, Paffrath T, Oestern HJ, Neugebauer E, Nast-Kolb D, Pape HC, Bouillon B: **Reduction in Mortality of Severely Injured Patients in Germany.** *Dtsch Arztebl Int* 2008, **105**:225-231.
 43. Osler T, Baker SP, Long W: **A modification of the injury severity score that both improves accuracy and simplifies scoring.** *J Trauma* 1997, **43**:922-925. discussion 925-926
 44. Baker SP, O'Neill B, Haddon W Jr, Long WB: **The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care.** *J Trauma* 1974, **14**:187-196.
 45. Brennenman FD, Boulanger BR, McLellan BA, Redelmeier DA: **Measuring injury severity: time for a change?** *J Trauma* 1998, **44**:580-582.
 46. Lavoie A, Moore L, LeSage N, Liberman M, Sampalis JS: **The New Injury Severity Score: a more accurate predictor of in-hospital mortality than the Injury Severity Score.** *J Trauma* 2004, **56**:1312-1320.
 47. Lavoie A, Moore L, LeSage N, Liberman M, Sampalis JS: **The Injury Severity Score or the New Injury Severity Score for predicting intensive care unit admission and hospital length of stay?** *Injury* 2005, **36**:477-483.
 48. Sacco WJ, MacKenzie EJ, Champion HR, Davis EG, Buckman RF: **Comparison of alternative methods for assessing injury severity based on anatomic descriptors.** *J Trauma* 1999, **47**:441-446. discussion 446-447
 49. Sullivan T, Haider A, DiRusso SM, Nealon P, Shaikat A, Slim M: **Prediction of mortality in pediatric trauma patients: new injury severity score outperforms injury severity score in the severely injured.** *J Trauma* 2003, **55**:1083-1087. discussion 1087-1088
 50. Tay SY, Sloan EP, Zun L, Zaret P: **Comparison of the New Injury Severity Score and the Injury Severity Score.** *J Trauma* 2004, **56**:162-164.
 51. Søreide K, Petrone P, Asensio JA: **Emergency thoracotomy in trauma: rationale, risks, and realities.** *Scand J Surg* 2007, **96**:4-10.
 52. National Burn Care Group: **National Burn Injury Database.** [<http://www.nationalburncaregroup.nhs.uk/welcome/national-burn-injury-database/>].
 53. Bergeron E, Rossignol M, Osler T, Clas D, Lavoie A: **Improving the TRISS methodology by restructuring age categories and adding comorbidities.** *J Trauma* 2004, **56**:760-767.
 54. Lefering R: **Trauma score systems for quality assessment.** *Eur J Trauma* 2002, **28**:52-63.
 55. Gannon CJ, Napolitano LM, Pasquale M, Tracy JK, McCarter RJ: **A statewide population-based study of gender differences in trauma: validation of a prior single-institution study.** *J Am Coll Surg* 2002, **195**:11-18.
 56. Knudson MM, Lieberman J, Morris JA Jr, Cushing BM, Stubbs HA: **Mortality factors in geriatric blunt trauma patients.** *Arch Surg* 1994, **129**:448-453.
 57. Morris JA Jr, MacKenzie EJ, Damiano AM, Bass SM: **Mortality in trauma patients: the interaction between host factors and severity.** *J Trauma* 1990, **30**:1476-1482.
 58. Wohltmann CD, Franklin GA, Boaz PW, Luchette FA, Kearney PA, Richardson JD, Spain DA: **A multicenter evaluation of whether gender dimorphism affects survival after trauma.** *Am J Surg* 2001, **181**:297-300.
 59. Spahn DR, Cerny V, Coats TJ, Duranseau J, Fernandez-Mondejar E, Gordini G, Stahel PF, Hunt BJ, Komadina R, Neugebauer E, et al.: **Management of bleeding following major trauma: a European guideline.** *Crit Care* 2007, **11**:R17.
 60. Boyle MJ: **Is mechanism of injury alone in the prehospital setting a predictor of major trauma – a review of the literature.** *J Trauma Manag Outcomes* 2007, **1**:4.
 61. World Health Organization: **International Statistical Classification of Diseases and Related Health Problems. 10th Revision. Version for 2007.** [<http://www.who.int/classifications/icd/en/>].
 62. Lu TH: **Changes in injury mortality by intent and mechanism in Taiwan, 1975-98.** *Inj Prev* 2002, **8**:70-73.
 63. Morris JA Jr, MacKenzie EJ, Edelman SL: **The effect of preexisting conditions on mortality in trauma patients.** *JAMA* 1990, **263**:1942-1946.
 64. Milzman DP, Boulanger BR, Rodriguez A, Soderstrom CA, Mitchell KA, Magnant CM: **Pre-existing disease in trauma patients: a predictor of fate independent of age and injury severity score.** *J Trauma* 1992, **32**:236-244.
 65. American Society of Anesthesiologists: **ASA Physical Status Classification System.** [<http://www.asahq.org/clinical/physicalstatus.htm>].
 66. Norwegian Society of Anaesthesiology: **ASA Physical Status Classification (ASA-klassifikasjonen).** [<http://nafweb.no/index.php/Standarder/asa.html>].

67. Battistella FD, Nugent W, Owings JT, Anderson JT: **Field triage of the pulseless trauma patient.** *Arch Surg* 1999, **134**:742-745. discussion 745-746
68. Shimazu S, Shatney CH: **Outcomes of trauma patients with no vital signs on hospital admission.** *J Trauma* 1983, **23**:213-216.
69. Rosemurgy AS, Norris PA, Olson SM, Hurst JM, Albrink MH: **Pre-hospital traumatic cardiac arrest: the cost of futility.** *J Trauma* 1993, **35**:468-473. discussion 473-464
70. Di Bartolomeo S, Sanson G, Nardi G, Michelutto V, Scian F: **HEMS vs. Ground-BLS care in traumatic cardiac arrest.** *Prehosp Emerg Care* 2005, **9**:79-84.
71. Arbabi S, Jurkovich GJ, Wahl WL, Franklin GA, Hemmila MR, Taheri PA, Maier RV: **A comparison of prehospital and hospital data in trauma patients.** *J Trauma* 2004, **56**:1029-1032.
72. Davis DP, Serrano JA, Vilke GM, Sise MJ, Kennedy F, Eastman AB, Velky T, Hoyt DB: **The predictive value of field versus arrival Glasgow Coma Scale score and TRISS calculations in moderate-to-severe traumatic brain injury.** *J Trauma* 2006, **60**:985-990.
73. Baxt WG, Moody P: **The impact of advanced prehospital emergency care on the mortality of severely brain-injured patients.** *J Trauma* 1987, **27**:365-369.
74. Servadei F, Nasi MT, Cremonini AM, Giuliani G, Cenni P, Nanni A: **Importance of a reliable admission Glasgow Coma Scale score for determining the need for evacuation of posttraumatic subdural hematoma: a prospective study of 65 patients.** *J Trauma* 1998, **44**:868-873.
75. Healey C, Osler TM, Rogers FB, Healey MA, Glance LG, Kilgo PD, Shackford SR, Meredith JW: **Improving the Glasgow Coma Scale score: motor score alone is a better predictor.** *J Trauma* 2003, **54**:671-678. discussion 678-680
76. Chan L, Bartfield JM, Reilly KM: **The significance of out-of-hospital hypotension in blunt trauma patients.** *Acad Emerg Med* 1997, **4**:785-788.
77. Lipsky AM, Gausche-Hill M, Henneman PL, Loffredo AJ, Eckhardt PB, Cryer HG, de Virgilio C, Klein SL, Bongard FS, Lewis RJ: **Prehospital hypotension is a predictor of the need for an emergent, therapeutic operation in trauma patients with normal systolic blood pressure in the emergency department.** *J Trauma* 2006, **61**:1228-1233.
78. Mackersie RC, Tiwary AD, Shackford SR, Hoyt DB: **Intra-abdominal injury following blunt trauma. Identifying the high-risk patient using objective risk factors.** *Arch Surg* 1989, **124**:809-813.
79. Davis JW, Mackersie RC, Holbrook TL, Hoyt DB: **Base deficit as an indicator of significant abdominal injury.** *Ann Emerg Med* 1991, **20**:842-844.
80. Franklin GA, Boaz PW, Spain DA, Lukan JK, Carrillo EH, Richardson JD: **Prehospital hypotension as a valid indicator of trauma team activation.** *J Trauma* 2000, **48**:1034-1037. discussion 1037-1039
81. Bouamra O, Wrotchford A, Hollis S, Vail A, Woodford M, Lecky F: **Outcome prediction in trauma.** *Injury* 2006, **37**:1092-1097.
82. Siegel JH, Rivkind AI, Dalal S, Goodarzi S: **Early physiologic predictors of injury severity and death in blunt multiple trauma.** *Arch Surg* 1990, **125**:498-508.
83. Davis JW, Kaups KL: **Base deficit in the elderly: a marker of severe injury and death.** *J Trauma* 1998, **45**:873-877.
84. Oestern HJ, Trentz O, Hempelmann G, Trentz OA, Sturm J: **Cardiorespiratory and metabolic patterns in multiple trauma patients.** *Resuscitation* 1979, **7**:169-183.
85. Davis JW, Parks SN, Kaups KL, Gladen HE, O'Donnell-Nicol S: **Admission base deficit predicts transfusion requirements and risk of complications.** *J Trauma* 1996, **41**:769-774.
86. Kroezen F, Bijlsma TS, Liem MS, Meeuwis JD, Leenen LP: **Base deficit-based predictive modeling of outcome in trauma patients admitted to intensive care units in Dutch trauma centers.** *J Trauma* 2007, **63**:908-913.
87. Maegele M, Lefering R, Yucel N, Tjardes T, Rixen D, Paffrath T, Simanski C, Neugebauer E, Bouillon B: **Early coagulopathy in multiple injury: an analysis from the German Trauma Registry on 8724 patients.** *Injury* 2007, **38**:298-304.
88. Brohi K, Singh J, Heron M, Coats T: **Acute traumatic coagulopathy.** *J Trauma* 2003, **54**:1127-1130.
89. Raum MR, Bouillon B, Rixen D, Lefering R, Tiling T, Neugebauer E, German Polytrauma Study Group of the German Trauma Society: **The Prognostic Value of Prothrombin Time in Predicting Survival after Major Trauma: a Prospective Analysis of 1,351 Patients from the German Trauma Registry.** *Eur J Trauma* 2001, **27**:110-116.
90. MacLeod JB, Lynn M, McKenney MG, Cohn SM, Murtha M: **Early coagulopathy predicts mortality in trauma.** *J Trauma* 2003, **55**:39-44.
91. Jennett B, Bond M: **Assessment of outcome after severe brain damage.** *Lancet* 1975, **1**:480-484.
92. Skaga NO, Eken T, Jones JM, Steen PA: **Different definitions of patient outcome: Consequences for performance analysis in trauma.** *Injury* 2008, **39**:612-622.
93. Skaga NO, Eken T, Hestnes M, Jones JM, Steen PA: **Scoring of anatomic injury after trauma: AIS 98 versus AIS 90-do the changes affect overall severity assessment?** *Injury* 2006.
94. McSwain NE Jr: **A plea for uniformity in EMS research.** *J Trauma* 2002, **52**:1220-1221.
95. Stockinger ZT, McSwain NE Jr: **Prehospital endotracheal intubation for trauma does not improve survival over bag-valve-mask ventilation.** *J Trauma* 2004, **56**:531-536.
96. Pons PT, Honigman B, Moore EE, Rosen P, Antuna B, Dernochoeur J: **Prehospital advanced trauma life support for critical penetrating wounds to the thorax and abdomen.** *J Trauma* 1985, **25**:828-832.
97. Spaite DW, Criss EA, Valenzuela TD, Meislin HW: **Prehospital advanced life support for major trauma: critical need for clinical trials.** *Ann Emerg Med* 1998, **32**:480-489.
98. Liberman M, Mulder D, Sampalis J: **Advanced or basic life support for trauma: meta-analysis and critical review of the literature.** *J Trauma* 2000, **49**:584-599.
99. Di Bartolomeo S, Sanson G, Nardi G, Michelutto V, Scian F: **HEMS vs. Ground-BLS Care in Traumatic Cardiac Arrest.** *Prehospital Emergency Care* 2005, **9**:79-84.
100. Thomas SH, Harrison TH, Buras WR, Ahmed W, Cheema F, Wedel SK: **Helicopter transport and blunt trauma mortality: a multicenter trial.** *J Trauma* 2002, **52**:136-145.
101. Pohlemann T, Gänsslen A, Bosch U, Tschern H: **The Technique of Packing for Control of Hemorrhage in Complex Pelvic Fractures.** *Tech Orthop* 1995, **4**:267-270.
102. Cothren CC, Osborn PM, Moore EE, Morgan SJ, Johnson JL, Smith WR: **Preperitoneal pelvic packing for hemodynamically unstable pelvic fractures: a paradigm shift.** *J Trauma* 2007, **62**:834-839. discussion 839-842
103. Totterman A, Madsen JE, Skaga NO, Roise O: **Extraperitoneal pelvic packing: a salvage procedure to control massive traumatic pelvic hemorrhage.** *J Trauma* 2007, **62**:843-852.
104. National Confidential Enquiry into Patient Outcome and Death: **Trauma: Who cares? A report of the National Confidential Enquiry into Patient Outcome and Death.** [<http://www.ncepod.org.uk/>].
105. American College of Surgeons Committee on Trauma: *Resources for optimal care of the injured patient 2006* Chicago, IL: American College of Surgeons; 2006.
106. Eastman AB, Lewis FR Jr, Champion HR, Mattox KL: **Regional trauma system design: critical concepts.** *Am J Surg* 1987, **154**:79-87.
107. Deane SA, Gaudry PL, Pearson I, Misra S, McNeil RJ, Read C: **The hospital trauma team: a model for trauma management.** *J Trauma* 1990, **30**:806-812.
108. Lossius HM, Langhelle A, Pillgram-Larsen J, Lossius TA, Søreide E, Laake P, Steen PA: **Efficiency of Activation of the Trauma Team in a Norwegian Trauma Referral Centre.** *European J Surg* 2000, **166**:760-764.
109. Cummins RO: **The "chain of survival" concept: how it can save lives.** *Heart Dis Stroke* 1992, **1**:43-45.
110. Rixen D, Raum M, Bouillon B, Neugebauer E: **[Base excess as prognostic indicator in patients with polytrauma].** *Anesthesiol Intensivmed Notfallmed Schmerzther* 2002, **37**:347-349.
111. ICECI Coordination and Maintenance Group: **International Classification of External Causes of Injuries (ICECI) version 1.2.** [<http://www.rivm.nl/who-fig/ICECIeng.htm>].
112. EU Injury Database: **Coding Manual V2000 for Home and Leisure Accidents. ISS Database version 2002.** [<https://webgate.ec.europa.eu/idb>].

113. Nordic Medico-Statistical Committee (NOMESCO): **NOMESCO Classification of External Causes of Injuries. Fourth revised edition.** [<http://www.nom-nos.dk/>].
114. Hannan EL, Mendeloff J, Farrell LS, Cayten CG, Murphy JG: **Multivariate models for predicting survival of patients with trauma from low falls: the impact of gender and pre-existing conditions.** *J Trauma* 1995, **38**:697-704.
115. Kennedy RL, Grant PT, Blackwell D: **Low-impact falls: demands on a system of trauma management, prediction of outcome, and influence of comorbidities.** *J Trauma* 2001, **51**:717-724.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

