

Durham Research Online

Deposited in DRO:

09 April 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Riley, Alexander H. and Fattahi, Azadeh and Pace, Andrew B. and Strigari, Louis E. and Frenk, Carlos S. and Gómez, Facundo A. and Grand, Robert J. J. and Marinacci, Federico and Navarro, Julio F. and Pakmor, Rüdiger and Simpson, Christine M. and White, Simon D.M. (2019) 'The velocity anisotropy of the Milky Way satellite system.', *Monthly notices of the Royal Astronomical Society.*, 486 (2). pp. 2679-2694.

Further information on publisher's website:

<https://doi.org/10.1093/mnras/stz973>

Publisher's copyright statement:

© 2019 The Author(s). Published by Oxford University Press on behalf of the Royal Astronomical Society.

Additional information:

Use policy


The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

The velocity anisotropy of the Milky Way satellite system

Alexander H. Riley,^{1*}  Azadeh Fattahi,² Andrew B. Pace,^{1†} Louis E. Strigari,¹
 Carlos S. Frenk,² Facundo A. Gómez,^{3,4} Robert J. J. Grand,^{5,6,7}
 Federico Marinacci,⁸ Julio F. Navarro,⁹ Rüdiger Pakmor,⁷
 Christine M. Simpson,^{10,11} and Simon D. M. White⁷

¹Department of Physics and Astronomy, Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA

²Institute for Computational Cosmology, Department of Physics, Durham University, South Road, Durham DH1 3LE, UK

³Instituto de Investigación Multidisciplinar en Ciencia y Tecnología, Universidad de La Serena, Raúl Bitrán 1305, La Serena, Chile

⁴Departamento de Física y Astronomía, Universidad de La Serena, Av. Juan Cisternas 1200 N, La Serena, Chile

⁵Heidelberger Institut für Theoretische Studien, Schloß-Wolfsbrunnenweg 35, 69118 Heidelberg, Germany

⁶Zentrum für Astronomie der Universität Heidelberg, Astronomisches Recheninstitut, Mönchhofstr. 12-14, 69120 Heidelberg, Germany

⁷Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85748, Garching, Germany

⁸Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁹Department of Physics and Astronomy, University of Victoria, Victoria, BC, V8P 1A1, Canada

¹⁰Enrico Fermi Institute, The University of Chicago, Chicago, IL 60637, USA

¹¹Department of Astronomy & Astrophysics, University of Chicago, Chicago, IL 60637, USA

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We analyse the orbital kinematics of the Milky Way (MW) satellite system utilizing the latest systemic proper motions for 38 satellites based on data from *Gaia* Data Release 2. Combining these data with distance and line-of-sight velocity measurements from the literature, we use a likelihood method to model the velocity anisotropy, β , as a function of Galactocentric distance and compare the MW satellite system with those of simulated MW-mass haloes from the APOSTLE and Auriga simulation suites. The anisotropy profile for the MW satellite system increases from $\beta \sim -2$ at $r \sim 20$ kpc to $\beta \sim 0.5$ at $r \sim 200$ kpc, indicating that satellites closer to the Galactic centre have tangentially-biased motions while those farther out have radially-biased motions. The motions of satellites around APOSTLE host galaxies are nearly isotropic at all radii, while the $\beta(r)$ profiles for satellite systems in the Auriga suite, whose host galaxies are substantially more massive in baryons than those in APOSTLE, are more consistent with that of the MW satellite system. This shape of the $\beta(r)$ profile may be attributed to the central stellar disc preferentially destroying satellites on radial orbits, or intrinsic processes from the formation of the Milky Way system.

Key words: galaxies: dwarf – Galaxy: kinematics and dynamics – dark matter

1 INTRODUCTION

Our presence within the Local Group offers it a special importance in astronomy. It is the only part of the Universe where we can detect small ($M_* \lesssim 10^5 M_\odot$) dwarf galaxies, resolve their stellar populations, and study their internal properties and kinematics. As the most dark-matter dominated galaxies in the Universe (McConnachie 2012), these dwarfs provide crucial tests of the current structure formation paradigm – cold dark matter with a cosmological constant (Λ CDM).

While several predictions of Λ CDM (e.g. large scale structure, temperature anisotropies of the cosmic microwave background) agree with observations extraordinarily well (Springel et al. 2005; Frenk & White 2012), the model faces a number of

challenges on the scales of dwarf galaxy satellites (see Bullock & Boylan-Kolchin 2017 for a recent review). Many of these challenges, including the so-called missing satellites (Klypin et al. 1999; Moore et al. 1999) and too-big-to-fail (Boylan-Kolchin et al. 2011, 2012) problems, have potential solutions through the inclusion of galaxy formation physics (Bullock et al. 2000; Benson et al. 2002; Somerville 2002) that have been reinforced by cosmological hydrodynamic simulations of galaxy formation (Okamoto & Frenk 2009; Brooks & Zolotov 2014; Sawala et al. 2016; Wetzel et al. 2016; Simpson et al. 2018; Garrison-Kimmel et al. 2018).

A growing challenge to Λ CDM is that a large fraction of satellites seem to be located on a co-rotating plane around their host galaxy (the plane-of-satellites problem; see Pawlowski 2018 for a recent review). Such planes have been observed around the Milky Way (MW; Lynden-Bell 1976; Kroupa et al. 2005; Libeskind et al. 2005; Pawlowski & Kroupa 2013; Fritz et al. 2018), Andromeda

* E-mail: alexriley@tamu.edu. Code for this work is available on Github.

† Mitchell Astronomy Fellow

PM study	N_{sats}	Satellites considered	Methodology	N_{stars}
Gaia Collaboration et al. (2018b)	12	classical satellites, Boötes I	Iterative, using only DR2 data	115/339.5/23109
Simon (2018)	17	$M_V > -8$, $d_{\odot} < 100$ kpc, w/ RV	Match RV members in DR2	2/8/68
Fritz et al. (2018) (F18)	39	$r < 420$ kpc w/ RV	Match RV members in DR2	2/18/2527
Kallivayalil et al. (2018)	13	possible satellites of LMC	Iterative, initial from RV (F18)	3/11/110
Massari & Helmi (2018)	7	$M_V < -2.5$, $d_{\odot} < 70$ kpc	Iterative, initial from RV or HB	29/53/189
Pace & Li (2018)	14	satellites in DES footprint	Probabilistic, incorporated DES DR1	5/15.5/67

Table 1. Summary of *Gaia* DR2 proper motion studies used in this analysis. N_{sats} is the number of satellites for which a proper motion was reported in the study. N_{stars} is the minimum/median/maximum number of stars for the list of satellites in the study. RV refers to spectroscopic radial velocity data, HB to photometric horizontal branch data. See Section 2 for further information.

(Ibata et al. 2013), and galaxies outside of the Local Group (Ibata et al. 2014, 2015; Müller et al. 2018). The degree to which planes of satellites pose a challenge to Λ CDM is contested; some analyses have concluded that a thin planar configuration of satellites is extremely unusual in Λ CDM (Pawlowski et al. 2012; Gillet et al. 2015), but a detailed statistical analysis (taking account of the “look elsewhere effect”) suggests that thin satellite planes like that of the MW and M31 occur in about 10% of galactic systems (Cautun et al. 2015).

Studies of the planes of satellites generally focus on two aspects of satellite kinematics: the clustering of orbital poles and the reconstruction of satellite orbits. The orbital poles of the MW satellites are more clustered than an isotropic distribution, with strong clustering measured for 8 of the 11 classical satellites (Pawlowski & Kroupa 2013). Orbit reconstruction is more challenging since the outcome is sensitive to the total and radial distribution of mass in the Milky Way, which are uncertain (Bovy 2015; McMillan 2017; see Figure 7 of Callingham et al. 2019 for a comparison of recent measurements of the total mass). This translates into large uncertainties in the reconstructed orbits, making direct comparisons to theoretical predictions more complicated.

To study the orbit structure of the satellite population in a potential-independent way, Cautun & Frenk (2017) used the velocity anisotropy, β , to characterize the orbital properties of the satellites. Introduced by Binney (1980) to quantify the orbital structure of a spherical system, β is most commonly used in spherical Jeans equation modeling to recover the mass of a system. In a Galactocentric spherical coordinate system where r corresponds to radial distance, θ the polar angle, and ϕ the azimuthal angle, β is defined as:

$$\beta(r) = 1 - \frac{\sigma_{\theta}(r)^2 + \sigma_{\phi}(r)^2}{2\sigma_r(r)^2} \quad (1)$$

where $\sigma_r, \sigma_{\theta}, \sigma_{\phi}$ are the velocity dispersions along each coordinate direction. The β parameter can take values in the range $-\infty$ to 1, where $\beta = 1$ corresponds to radial orbits plunging in and out of the Galactic centre, $\beta \rightarrow -\infty$ to circular orbits, and $\beta = 0$ to velocities being isotropically distributed at each point.

Studies of β for the MW have predominantly used either halo stars (Deason et al. 2012, 2013; Cunningham et al. 2016, 2018) or globular clusters (Sohn et al. 2018; Watkins et al. 2019; Vasiliev 2019) as tracers. While these studies have largely focused on obtaining a single value of β as an input for spherical Jeans equation modeling, the radial anisotropy profile $\beta(r)$ also contains interesting information on the accretion history of the MW. For example, Loebman et al. (2018) used high-resolution cosmological hydrodynamic simulations to find that dips in the $\beta(r)$ profile of halo stars may be associated with localized perturbations from, or remnants

of, destroyed satellites. Such perturbations have been observed in the $\beta(r)$ profile for halo stars in the Milky Way (Cunningham et al. 2018).

To date, the only measurement of β using MW satellite galaxies has been from Cautun & Frenk (2017). Using proper motions derived using the Hubble Space Telescope (HST) for the 10 brightest satellite galaxies, they obtained $\beta = -2.2 \pm 0.4$. The low number of MW satellites with measured proper motions prohibited further studies of the $\beta(r)$ profile until the second data release from *Gaia* (DR2; Gaia Collaboration et al. 2018a). Since *Gaia* DR2, proper motions for nearly every MW satellite galaxy have been measured, which now motivates further studies of β using the MW satellites as tracers.

In this paper, we compare the kinematics of the Milky Way satellites to expectations from Λ CDM using state-of-the-art cosmological hydrodynamic zoom (Katz & White 1993; Frenk et al. 1996; Oñorbe et al. 2014) simulations: APOSTLE (A Project Of Simulating The Local Environment; Fattahi et al. 2016; Sawala et al. 2016) and Auriga (Grand et al. 2017). By focusing on β , our results only depend on the present-day kinematics of the MW satellites and not the total or radial distribution of MW mass. We use the latest satellite proper motion measurements as deduced from *Gaia* DR2, increasing the number of satellites used in an anisotropy analysis from the 10 in Cautun & Frenk (2017) to 38. Furthermore, we utilize a likelihood method to determine the intrinsic σ_i 's of the MW satellite system. This more robust method, combined with the increased number of satellites spread over a wide range of Galactocentric distances ($\sim 15 - 250$ kpc), allows us to perform the first measurement of $\beta(r)$ for the satellites of the Milky Way.

This paper is organized as follows. In Section 2 we review the new *Gaia* DR2 proper motions for MW satellites included in our analysis. In Section 3 we describe the cosmological hydrodynamic zoom simulations that provide our predictions within Λ CDM. In Section 4 we detail our methodology for computing β . In Section 5 we present the main results of our analysis and in Section 6 we provide a possible interpretation of these results. In Section 7 we present our conclusions.

2 PROPER MOTIONS

The public release of *Gaia* DR2 has profoundly impacted near-field cosmology in a very short period of time. The data release contains an all-sky catalog of the five-parameter astrometric solution (position on the sky, parallax, and proper motion) for more than 1.3 billion sources (Gaia Collaboration et al. 2018a). These data have already been used in multiple studies of the kinematics of the




Figure 1. Tangential velocity excess of Milky Way satellites using proper motions from F18 (but LMC and SMC proper motions from Gaia Collaboration et al. 2018b). A ratio of radial to total kinetic energy $V_{\text{rad}}^2/V_{\text{tot}}^2 \lesssim 1/3$ indicates a tangentially biased motion.

Milky Way’s stellar halo (e.g. Deason et al. 2018), satellites (e.g. Callingham et al. 2019) and globular clusters (e.g. Vasiliev 2019).

We use results from six studies (see Table 1 for a summary) which derive *Gaia* DR2 proper motions for MW satellites with comparable precision to those derived using the Hubble Space Telescope (for a review of proper motions with HST, see van der Marel et al. 2014). *Gaia* Collaboration et al. (2018b) demonstrated *Gaia* DR2’s ability to constrain proper motions for the Magellanic Clouds, the classical (pre-SDSS) satellites, and ultra-faint dwarf Boötes I. Simon (2018) presented the first proper motions for many nearby (< 100 kpc) ultra-faint dwarf galaxies, while Fritz et al. (2018, hereafter F18) extended the limit out to 420 kpc with the largest sample of 39 satellites. Kallivayalil et al. (2018) derived proper motions for satellites located near the Magellanic Clouds, motivated by the possibility that some of them may be satellites of the LMC itself. Massari & Helmi (2018) computed proper motions for seven dwarfs, three of which do not have spectroscopic information. Pace & Li (2018) presented a probabilistic method of determining systemic proper motions that utilized the superb photometry from the first public data release of the Dark Energy Survey (Abbott et al. 2018).

The full list of satellites that we consider in this analysis is presented in Table A1, along with a summary of their properties. For this analysis we only consider satellites out to 300 kpc from the Galactic centre. We omit globular clusters and satellites whose nature is still under debate (e.g. Crater I; Kirby et al. 2015; Voggel et al. 2016). We also do not consider overdensities that are thought to be tidally disrupting dwarf galaxies: Boötes III (Carlin et al. 2009; Carlin & Sand 2018), Canis Major (Martin et al. 2004), and Hydra I (Hargis et al. 2016). Furthermore, we restrict our analysis to satellites which have published line-of-sight velocities in order to have full 6-D kinematic information.

2.1 Galactocentric coordinates

In order to convert the line-of-sight velocity and proper motion measurements into Galactocentric coordinates, we use the distance measurements from Table A1. The Galactocentric Cartesian coordi-

ates are then computed assuming a distance from the Sun to the Galactic centre of 8.2 ± 0.1 kpc, a height of the Sun relative to the Galactic plane of 25 ± 5 pc, and a solar motion relative to the Galactic centre of $(10 \pm 1, 248 \pm 3, 7 \pm 0.5)$ km s⁻¹ (Bland-Hawthorn & Gerhard 2016), in a frame where the x-axis points from the position of the Sun projected on to the Galactic plane to the Galactic centre, the y-axis points towards Galactic longitude $l = 90^\circ$ (i.e. in the direction of Galactic rotation), and the z-axis points towards the North Galactic Pole. This right-handed Cartesian system is then converted into spherical coordinates, with r the distance from the Galactic centre, polar angle θ defined from the z-axis, and azimuthal angle ϕ defined from the x-axis such that the Galactic rotation is in the $-\phi$ direction.

We perform 2,000 Monte Carlo simulations drawing satellite proper motions, line-of-sight velocities, and heliocentric distances randomly from Gaussian distributions centred on their measured values with dispersions given by their respective errors. When drawing the proper motions we account for the correlation between $\mu_{\alpha*} \equiv \mu_{\alpha} \cos \delta$ and μ_{δ} if provided in the proper motion study. The randomly drawn kinematic properties are then converted into Galactocentric spherical coordinates as described in the previous paragraph. The resulting Galactocentric positions and velocities (and corresponding uncertainties), obtained directly from the observed distance, line-of-sight velocity, and proper motion measurements, are summarized in Table A2.

To illustrate the tangential nature of the motions of the MW satellites, we show the ratio of radial to total kinetic energy $V_{\text{rad}}^2/V_{\text{tot}}^2$ for each satellite in Figure 1. A ratio $\lesssim 1/3$ indicates a tangentially-biased motion. We find that $\sim 80\%$ of MW satellites show a tangential velocity excess, comparable to Cautun & Frenk (2017) who found that 9 of the 10 brightest MW satellites had tangentially-biased motions.

2.2 Sample selections

It is important to note that the proper motions derived by both Simon (2018) and F18 were based only on matching spectroscopically confirmed member stars with *Gaia* DR2 data, in some cases depending on very few ($N \sim 2-5$) stars to derive a systemic proper motion. The small number statistics could lead to a biased result; Massari & Helmi (2018) found that the subsample of spectroscopic members in Segue 2 used by Simon (2018) and F18 is systematically shifted in proper motion space relative to the full sample recovered using their iterative method. To avoid problems from this potential bias, as well as to confirm that our results do not depend strongly on the systematics associated with a particular study, we consider three different samples of proper motion data in our analysis:

(i) 38 satellites, comprised of 36 satellites from F18 plus the LMC and SMC. This is the full list of satellites in Table A1.

(ii) a “gold” sample constructed by prioritizing studies which included steps in their analysis to increase the sample of member stars beyond the spectroscopic sample. For example, Pace & Li (2018) used a probabilistic method incorporating photometry from the first public data release of the Dark Energy Survey (Abbott et al. 2018). This “gold” sample consists of 32 satellites,¹ with proper motions taken from the five other previous studies.

¹ The satellites that are in the full sample that are excluded from the “gold” sample are: Canis Venatici I, Canis Venatici II, Hercules, Leo IV, Leo V, and Pisces II.

(iii) the same 32 satellites from the “gold” sample, but using the F18 proper motions.

Since *Gaia* DR2 proper motions for the Magellanic Clouds have only been reported by [Gaia Collaboration et al. \(2018b\)](#), we use these proper motion measurements in all samples. The exact study used for each satellite in the “gold” sample is shown in Table A1. As detailed in Section 5, we find that our results do not depend on which sample is used. We focus on the results for the full 38 satellite sample using F18 proper motions in the following sections.

3 COSMOLOGICAL SIMULATIONS

To compare our results with the expectations from the standard Λ CDM cosmology, we utilize suites of self-consistent cosmological hydrodynamic zoom simulations of Local Group analogs, APOSTLE ([Fattahi et al. 2016](#); [Sawala et al. 2016](#)), and of Milky Way analogs, Auriga ([Grand et al. 2017](#)). These two simulation suites have similar resolution and include similar baryonic processes (e.g. star formation, stellar, supernova and black hole feedback, uniform background UV field for reionization), though the numerical methods and prescriptions for subgrid physics are different (see references in following subsections for details). We also analyse dark-matter-only runs from these suites for comparison.

3.1 APOSTLE

The APOSTLE project is a suite of cosmological hydrodynamic zoom simulations of twelve volumes using the code developed for the EAGLE project ([Schaye et al. 2015](#); [Crain et al. 2015](#)). The galaxy formation model includes metallicity-dependent star formation and cooling, metal enrichment, stellar and supernova feedback, homogeneous X-ray/UV background radiation (hydrogen reionization assumed at $z_{\text{reion}} = 11.5$), supermassive black-hole formation and growth, and AGN activity ([Booth & Schaye 2009](#); [Rosas-Guevara et al. 2015](#)). The full details of the subgrid physics can be found in [Schaye et al. \(2015\)](#).

The APOSTLE volumes were selected to have similar dynamical properties as the Local Group; the full selection procedure is described in [Fattahi et al. \(2016\)](#) and a discussion of the main simulation characteristics is given in [Sawala et al. \(2016\)](#). In summary, each volume consists of a MW/M31-like pair of haloes with halo mass² ranging from $0.5 - 2.5 \times 10^{12} M_{\odot}$, separated by 800 ± 200 kpc, approaching with radial velocity $< 250 \text{ km s}^{-1}$ and tangential velocity $< 100 \text{ km s}^{-1}$. The haloes are isolated, with no additional halo larger than the smaller of the pair within 2.5 Mpc of the midpoint between the pair, and in environments where the Hubble flow is relatively unperturbed out to 4 Mpc. The simulations adopt the WMAP-7 cosmological parameters ([Komatsu et al. 2011](#)): $\Omega_M = 0.272$, $\Omega_{\Lambda} = 0.728$, $h = 0.704$, $\sigma_8 = 0.81$, and $n_s = 0.967$.

The volumes were simulated at three resolution levels, the highest of which (and the only level considered here) has primordial gas (DM) particle masses approximately $1.0(5.0) \times 10^4 M_{\odot}$, with a maximum force softening length of 134 pc. Five volumes have been simulated so far at this resolution, corresponding to AP-01, AP-04, AP-06, AP-10, and AP-11 in Table 2 of [Fattahi et al.](#)

² Defined to be the mass inside a sphere in which the mean matter density is 200 times the critical density $\rho_{\text{crit}} = 3H^2(z)/8\pi G$. Virial quantities are defined at that radius and are identified by a ‘200’ subscript.




Figure 2. Circular velocity profiles based on total mass for APOSTLE (blue) and Auriga (orange) haloes. The formation of stellar discs in Auriga is reflected in a much deeper potential near the centre of the halo. For comparison, we also show the circular velocity profile for an NFW ([Navarro et al. 1996, 1997](#)) halo with $M_{200} = 10^{12} M_{\odot}$ and $c = 10$ (black, dashed), where c is the ratio between the virial radius and the NFW scale radius.

(2016). Each halo in a pair is treated separately in this analysis, resulting in ten high-resolution APOSTLE haloes being considered in this work.

3.2 Auriga

The Auriga simulations ([Grand et al. 2017](#)) are a suite of cosmological magnetohydrodynamic zoom simulations of single MW-like haloes with halo masses in the range $1 - 2 \times 10^{12} M_{\odot}$. They were performed with the moving mesh code AREPO ([Springel 2010](#)) and a galaxy formation model that includes primordial and metal line cooling, a prescription for a uniform background UV field for reionization (completed at $z = 6$), a subgrid model for star formation and stellar feedback ([Springel & Hernquist 2003](#)), magnetic fields ([Pakmor et al. 2014, 2017](#)), and black hole seeding, accretion, and feedback.

The Auriga haloes were identified as isolated in the $z = 0$ snapshot of the parent dark-matter-only simulation with a 100 Mpc box-side length of the EAGLE project introduced in [Schaye et al. \(2015\)](#). To be considered isolated, the centre of any target halo must be located outside 9 times the halo radius of any other halo that has a mass greater than 3% that of the target. The simulations assumed the [Planck Collaboration et al. \(2014\)](#) cosmological parameters: $\Omega_M = 0.307$, $\Omega_{\Lambda} = 0.693$, $h = 0.6777$, $\sigma_8 = 0.8288$, and $n_s = 0.9611$.

The volumes were simulated at three resolution levels, the highest of which (and the only level considered here) has baryonic element (DM particle) masses approximately $0.5(4.0) \times 10^4 M_{\odot}$, with a maximum force softening length of 185 pc, comparable to the highest resolution for APOSTLE. Thus far, six haloes have been resimulated at this high resolution, corresponding to Au6, Au16, Au21, Au23, Au24, and Au27 in Table 1 of [Grand et al. \(2017\)](#).

3.3 Stellar discs

Even though the APOSTLE and Auriga haloes have similar halo masses, the difference in their baryon content at $z = 0$ affects the

Run	M_{200} [$10^{12} M_{\odot}$]	R_{200} [kpc]	M_* [$10^{10} M_{\odot}$]	$R_{1/2,*}$ [kpc]	$V_{\text{circ}}(8.2 \text{ kpc})$ [km s^{-1}]	N_{subs} ($V_{\text{max}} > 5 \text{ km s}^{-1}$)	N_{subs} ($M_* > 0$)
APOSTLE							
AP-01	1.57	238.8	2.3	8.2	173.6	1187	52
	1.02	206.8	1.1	6.2	124.3	1071	41
AP-04	1.16	216.2	1.1	5.0	155.3	1006	63
	1.12	213.7	1.6	4.7	148.8	1232	62
AP-06	2.04	260.6	2.2	5.5	172.6	1517	76
	1.07	210.3	1.2	7.4	136.7	999	27
AP-10	1.43	231.5	2.2	6.7	163.6	1105	35
	0.47	160.1	1.0	6.2	121.0	669	26
AP-11	0.90	198.5	1.0	3.3	150.4	810	36
	0.78	189.3	0.9	4.2	136.5	784	33
Auriga							
Au6	1.01	211.8	6.3	4.7	224.2	517	74
Au16	1.50	241.5	8.8	9.6	217.7	594	95
Au21	1.42	236.7	8.6	7.7	231.7	621	97
Au23	1.50	241.5	8.8	8.1	240.6	582	83
Au24	1.47	239.6	8.5	8.4	219.0	629	87
Au27	1.70	251.4	9.7	6.6	254.5	564	104
MW	1.1 ± 0.3	220.7^a	5 ± 1	–	238 ± 15	–	124^{+40}_{-27}

Table 2. Summary of simulation parameters at $z = 0$. The columns are: 1) simulation name, 2) halo mass, 3) halo radius, 4) stellar mass within $0.15 \times R_{200}$, 5) half-stellar mass radius, 6) circular velocity at $R_0 = 8.2 \text{ kpc}$, 7) number of subhaloes with maximum circular velocity $V_{\text{max}} > 5 \text{ km s}^{-1}$, and 8) number of subhaloes which form stars. Note that the APOSTLE volumes contain a pair of haloes as analogs for the MW and M31. Each halo in a pair is considered separately in this analysis. The final row provides current estimates for these quantities for the Milky Way from [Bland-Hawthorn & Gerhard \(2016\)](#) and [Newton et al. \(2018\)](#), though note that the latter considers satellites down to $M_V = 0$, too faint for APOSTLE and Auriga to resolve. ^a refers to the mean of the values for R_{200} provided in Table 8 of [Bland-Hawthorn & Gerhard \(2016\)](#), the standard deviation of which is 28.6 kpc . The value for the mass of the MW is consistent with the latest measurement from *Gaia* DR2 halo globular cluster motions ($1.41^{+1.99}_{-0.52} \times 10^{12} M_{\odot}$; [Watkins et al. 2019](#)).

shape and depth of their potentials (and hence the dynamics of their satellite systems). A main difference between the two simulation suites is the mass and morphology of the stellar discs of their main galaxies.³ The Auriga simulations are able to produce radially extended and thin discs, with sizes comparable to that of the MW ([Grand et al. 2017, 2018](#)), while their total stellar masses are slightly higher, close to $10^{11} M_{\odot}$, than that of the MW. By contrast, the APOSTLE host galaxies have morphologies that are less disky with relatively low stellar mass, $\sim 10^{10} M_{\odot}$.⁴ A summary of properties for each simulation run is shown in Table 2.

The total circular velocity profiles, $V_{\text{circ}} = \sqrt{GM(< r)}/r$, for the two simulations are shown in Figure 2. The different behaviour of the APOSTLE and Auriga haloes (blue and yellow curves) compared to the NFW circular velocity profile (black dashed curve) is due to the contraction of haloes in response to the presence of baryons. The much larger difference in the circular velocity profiles between the two simulation suites is due to the more massive stellar discs in Auriga (orange curves) combined with the enhanced dark matter contraction.

These differences are useful in quantifying the effect of a stellar disc on β . The deepening of the potential due to the large baryonic disc, combined with the non-spherical potential of the disc, can affect the tidal stripping of subhaloes. Hydro-

dynamic simulations suggest that tidal effects from a baryonic disc near the centre of a host halo can reduce the number of dark substructures by up to a factor of two ([Brooks & Zolotov 2014; Zhu et al. 2016; Sawala et al. 2017; Nadler et al. 2018](#)), an effect that is reproduced in DMO simulations with an embedded disc potential ([D’Onghia et al. 2010; Yurin & Springel 2015; Garrison-Kimmel et al. 2017b; Kelley et al. 2018](#)). This tidal disruption preferentially affects radial orbits that come close to the disc ([Garrison-Kimmel et al. 2017b](#)), implying that surviving subhaloes in the inner regions should be on circular orbits, resulting in a lower β near the centre.

3.4 Matching the radial distributions

In addition to comparing the results of simulated systems to that of the MW satellites, we would like to select samples of subhaloes that are more representative of the observed MW satellites’ radial distribution. By comparing the results of these subhalo samples to those of the full simulated systems, we will be able to understand the impact of the tracers’ radial distribution on our results. We will also provide a fairer comparison between the simulated systems and the MW satellites.

We begin by considering all subhaloes which, at $z = 0$, have maximum circular velocity $V_{\text{max}} > 5 \text{ km s}^{-1}$. This is a conservative resolution limit for both APOSTLE and Auriga, roughly corresponding to subhalo masses of $\sim 5 \times 10^6 M_{\odot}$ or containing ~ 100 DM particles. These subhaloes are a mix of dark and luminous (i.e. contain stars); typically $\sim 4\%$ contain stars in APOSTLE and $\sim 15\%$ contain stars in Auriga (see Table 2).

³ The gas content of these galaxies is sub-dominant compared to the stellar component at small radii.

⁴ We note that the host galaxies in low and medium resolution APOSTLE runs have disky morphologies and higher stellar masses compared to the high resolution runs used here.




Figure 3. The radial distribution of subhaloes with $V_{\max} > 5 \text{ km s}^{-1}$ in APOSTLE (blue) and Auriga (orange) compared to that of the MW satellites (black). The deepening of the potential in Auriga haloes results in a less centrally concentrated radial distribution compared to APOSTLE, due to more subhaloes being destroyed. The magenta contours correspond to matching the radial distribution but not number of satellites, while the green contours correspond to additionally matching the number of satellites (see Section 3.4 for details) for both suites. Solid curves indicate the median and shaded regions the total spread.

We then create two subhalo samples resulting from (1) matching the radial distribution of subhaloes to that of the MW satellites and (2) additionally matching the abundance of subhaloes to that of the MW satellites. When matching both radial distribution and abundance (case 2), we simply compare the Galactocentric distance of each MW satellite to the host-centric distance of each subhalo and select the closest match (without replacement). When only matching the radial distribution and not the abundance (case 1), we select subhaloes based on the following inverse transform sampling method:

- (i) Compute the cumulative distribution function (CDF) of MW satellite Galactocentric distances.
- (ii) Generate a random number uniformly between 0 and 1 and map that number to a distance using the CDF from step 1.
- (iii) Select the subhalo that has a host-centric distance that is closest to the randomly generated distance and add it to the sample if it is within 5 kpc of the randomly generated value. This subhalo is removed from possible selection in the future, i.e. without replacement.
- (iv) Repeat steps (ii)-(iii) until a distance is generated that does not have a subhalo match within 5 kpc.

The 5 kpc cutoff is meant to strike a balance between increasing the number of subhaloes in the sample (higher cutoff) and providing a close match between the radial distribution of subhaloes to that of the MW satellites (lower cutoff). Our results are not sensitive to the exact value of this cutoff. Using this method, we typically find subhalo radial distributions that are much closer to the MW satellite distribution than that of the original subhalo populations (see Figure 3).

4 LIKELIHOOD ANALYSIS

We seek to model the orbital kinematics of Milky Way satellites and compare the results with those of cosmological simulations using the velocity anisotropy parameter β . Two models are considered: (1) a constant value of β at all radii and (2) one in which β varies as a function of Galactocentric distance. To determine the posterior probability densities for each model, we use *emcee* (Foreman-Mackey et al. 2013), an implementation of the affine-invariant ensemble sampler for Markov chain Monte Carlo (MCMC).

4.1 Framework and constant β model

We assume that the velocity distribution of the MW satellite system in Galactocentric spherical coordinates (r, θ, ϕ) is a multivariate Gaussian with different means and dispersions in each direction. The resulting likelihood F_i for a given satellite i with velocity $\mathbf{v}_i = (v_{r,i}, v_{\theta,i}, v_{\phi,i})$ is then

$$F_i = \frac{1}{\sqrt{(2\pi)^3 |\Sigma_i|}} \exp \left[-\frac{(\mathbf{v}_i - \mathbf{v}_{\text{sys}})^T \Sigma_i^{-1} (\mathbf{v}_i - \mathbf{v}_{\text{sys}})}{2} \right], \quad (2)$$

where $\mathbf{v}_{\text{sys}} = (v_r, v_{\theta}, v_{\phi})$ are the intrinsic mean velocities of the system (i.e. the entire population of MW satellites) and the covariance matrix is

$$\Sigma_i \equiv \begin{bmatrix} \sigma_r^2 + \delta_r^2 & C_{r\theta} \delta_r \delta_{\theta} & C_{r\phi} \delta_r \delta_{\phi} \\ C_{r\theta} \delta_r \delta_{\theta} & \sigma_{\theta}^2 + \delta_{\theta}^2 & C_{\theta\phi} \delta_{\theta} \delta_{\phi} \\ C_{r\phi} \delta_r \delta_{\phi} & C_{\theta\phi} \delta_{\theta} \delta_{\phi} & \sigma_{\phi}^2 + \delta_{\phi}^2 \end{bmatrix}. \quad (3)$$

Here, $(\sigma_r, \sigma_{\theta}, \sigma_{\phi})$ are the intrinsic dispersions of the system (i.e. the entire population of MW satellites) and $(\delta_r, \delta_{\theta}, \delta_{\phi}, C_{r\theta}, C_{r\phi}, C_{\theta\phi})$ are the observed measurement errors and correlation coefficients for the velocities of the given satellite, which are obtained from the 2,000 Monte Carlo samples described in Section 2.1.

Due to the conversion from heliocentric to Galactocentric spherical coordinates, the resulting satellite velocity errors are not necessarily Gaussian in each component. We find that approximating the errors as Gaussian is reasonable in most cases, though Draco II, Tucana III, and Willman 1 show significant skewness and kurtosis in both v_{θ} and v_{ϕ} .

The combined log-likelihood for the full satellite sample is then

$$\ln \mathcal{L} = \sum_{i=1}^{N_{\text{sats}}} \ln F_i = -\frac{1}{2} \sum_{i=1}^{N_{\text{sats}}} 3 \ln 2\pi + \ln |\Sigma_i| + \mathbf{u}_i^T \Sigma_i^{-1} \mathbf{u}_i \quad (4)$$

$$\propto -\sum_{i=1}^{N_{\text{sats}}} \mathbf{u}_i^T \Sigma_i^{-1} \mathbf{u}_i + \ln |\Sigma_i|, \quad (5)$$

where $\mathbf{u}_i \equiv \mathbf{v}_i - \mathbf{v}_{\text{sys}}$. Equation 5 is the likelihood function used to probe the model parameter space with *emcee*.

The first model we consider assumes constant velocity dispersions at all radii, resulting in a constant value for β . We impose spherical symmetry by requiring $v_r = v_{\theta} = 0$ and $\sigma_r^2 = \sigma_{\theta}^2 = \sigma_{\phi}^2$, as is commonly assumed in Jeans equation modeling of the dynamics of a system. In total this model then has 3 free parameters: a mean rotational motion v_{ϕ} and dispersions σ_r and σ_{θ} . We assume uniform priors for the mean motion $-500 < v_{\phi} < 500 \text{ km s}^{-1}$ as well as




Figure 4. Posterior distributions for all 38 Milky Way satellites assuming the constant β model, using F18 proper motions. From left to right the parameters are: v_ϕ (systemic rotational motion in km s^{-1}), σ_r , and $\sigma_\theta = \sigma_\phi$ (intrinsic velocity dispersions in km s^{-1}). The contours enclose 39.4, 85.5, 98.9% of the posterior distribution corresponding to 1, 2, 3 – σ confidence intervals. The dotted lines on the 1-D histograms are the 16th, 50th, and 84th percentiles and the numerical values are quoted above.

for the dispersions $0 < \sigma_i < 300 \text{ km s}^{-1}$. We repeat the analysis with Jeffreys prior $-3 < \log_{10} \sigma_i < 3$ and find that this does not meaningfully change our results.

4.2 Variable dispersions with radius

To take advantage of the increased number of satellites with proper motions over a wide range of Galactocentric distances, we include a separate likelihood analysis in which we adopt a simple model for the velocity dispersion to vary as a function of radius in each coordinate j :

$$\sigma_j(r) = \sigma_{j,0} \left(1 + \frac{r}{r_{j,0}} \right)^{-\alpha_j} \quad (6)$$

where $\sigma_{j,0}$ and $r_{j,0}$ are the characteristic dispersion and length scales and α_j is the slope of the fall off at large radii. We then use the same likelihood function as in Section 4.1 (specifically Equation 5) with the additional parameters introduced in Equation 6. The $\beta(r)$ profile then follows from Equation 1.

As in the constant β model, we impose spherical symmetry by requiring $v_r = v_\theta = 0$ and $\sigma_\theta^2(r) = \sigma_\phi^2(r)$. In total this model then has 7 parameters: a mean rotational motion v_ϕ (which is held constant with r), the characteristic dispersion and length scales $\sigma_{i,0}$ and $r_{i,0}$, and the slope α_i , for both Galactocentric spherical coordinates r and θ . We assume the same uniform prior for the mean motion $-500 < v_\phi < 500 \text{ km s}^{-1}$ as in the constant β analysis. For the $\sigma_i(r)$ parameters we assume uniform priors $50 < \sigma_{i,0} < 1000 \text{ km s}^{-1}$, $10 < r_{i,0} < 1000 \text{ kpc}$, and $0 < \alpha_i < 10$. We repeat the analysis with Jeffreys priors $-3 < \log_{10} \sigma_{i,0} < 3$ and




Figure 5. Posterior distributions for β assuming the constant β model with F18 proper motions. The results are shown using all 38 satellites (blue), satellites within 100 kpc (orange, 23 satellites), and satellites outside of 100 kpc (green, 15 satellites). The vertical black dashed line corresponds to the isotropic case $\beta = 0$.

$-3 < \log_{10} r_{i,0} < 3$ and again find that this change of priors does not meaningfully change our results.

5 RESULTS

We now present the resulting posterior probability densities for $\beta(r)$ for the MW satellite system using the models described above. We show that satellites within $r \lesssim 100 \text{ kpc}$ have more tangentially-biased motions (lower β) than those farther away. This result is also seen in simulated MW analogs, but it is difficult to disentangle effects due to the central stellar disc from those imprinted at formation. From here onwards, we refer to dark-matter-only simulations from the APOSTLE and Auriga suites collectively as “DMO” and the haloes simulated with baryonic physics by their suite name.

5.1 Constant β model

The posterior distribution of parameters for the constant β model for the MW satellites is shown in Figure 4 and the resulting posterior for β is shown in Figure 5 (blue curve). We find that the satellites are overall on near tangential orbits, with $\beta = -1.02^{+0.37}_{-0.45}$. We do not find significant evidence for the MW satellite population exhibiting rotation parallel to the plane of the Milky Way disc ($v_\phi = -14^{+20}_{-20} \text{ km s}^{-1}$; note that a star located in the disc would have v_ϕ on the order of $\sim 100 \text{ km s}^{-1}$). We also find that the constant β results are similar when using the different samples described in Section 2.2 (“gold” sample and “gold” satellites with F18 proper motions).

To better understand the results from the variable $\beta(r)$ model, we examine two radial bins. We split the satellites into two populations, one with $r < 100 \text{ kpc}$ (23 satellites) and the other with $r > 100 \text{ kpc}$ (15 satellites), and perform the same constant β analysis on each. The inner and outer regions clearly have different posterior distributions (Figure 5, orange and green curves respectively), with the inner region having a more negative (i.e., more tangentially biased) β posterior than the outer region. These results




Figure 6. Posterior distributions for the $\beta(r)$ model for the 38 MW satellites using F18 proper motions. Top: posterior for $\beta(r)$. The horizontal black dashed line corresponds to the isotropic case, $\beta = 0$. Bottom: posterior for the systemic dispersion in σ_r (orange) and $\sigma_\theta = \sigma_\phi$ (green). The brown ticks along the middle axis mark the radial positions of the MW satellites. For both panels, solid curves correspond to the median values and the shaded region the 16 – 84% confidence interval.

do not change when considering each of the different proper motion samples described in Section 2.2. This supports the finding in the $\beta(r)$ model (discussed below) that satellites in the inner region ($r \lesssim 100$ kpc) have more tangentially-biased motions than those farther away.

5.1.1 Comparison to Cautun & Frenk 2017

Our constant β result agrees within 2σ with the result of Cautun & Frenk (2017). These authors found $\beta = -2.2 \pm 0.4$ using HST proper motions of only 10 of the brightest satellites and simply computing β from Monte Carlo realizations of the MW satellite system using observational errors. Using updated *Gaia* DR2 proper motions and our likelihood method, our result for that same subsample of 10 satellites is $\beta = -1.52^{+0.86}_{-1.23}$, which is consistent with Cautun & Frenk (2017). The small offset is likely due to different input data and analysis techniques.

5.2 Variable β model

The posterior distribution for the parametrized $\beta(r)$ model for the MW satellites is shown in Figure 6 (top panel), along with the dispersions σ_r and $\sigma_\theta = \sigma_\phi$ (bottom panel). We find that the radial profile dips in the inner (< 100 kpc) region to $\beta \sim -2$ at $r \sim 20$ kpc and flattens out to $\beta \sim 0.5$ in the outer region. This again indicates that satellites near the centre of the Milky Way have tangentially-biased motions, while satellites in the outer region have more radially-biased motions.

Using the other proper motion samples described in Section 2.2 does not impact the results; the “gold” sample and “gold” satellites with F18 proper motions have nearly the same $\beta(r)$ pro-




Figure 7. Posterior distributions for the $\beta(r)$ model using different satellite samples. The “gold” sample and “gold” satellites using F18 proper motions are as described in Section 2.2. “No LMC satellites” excludes the candidate satellites of the LMC as identified by Kallivayalil et al. (2018). We define ultra-faint dwarf galaxies as those fainter than $10^5 L_\odot$ (Bullock & Boylan-Kolchin 2017) and refer to galaxies brighter than this limit as “classical.” The results from these different input samples are all consistent (within 68% confidence) with the original full sample of 38 satellites.

file. Furthermore, this dip in β does not appear to be dependent on a particular satellite or population of satellites. We repeated the analysis removing Sagittarius (which has a well-constrained proper motion at $r \sim 18$ kpc), removing satellites with luminosities above or below the median luminosity, and removing candidate satellites of the LMC identified by Kallivayalil et al. (2018): Horologium I, Carina II, Carina III, and Hydrus I. The results from these different input samples are all consistent (within 68% confidence) with the original full sample of 38 satellites (see Figure 7).

Taken together, these results indicate that satellites closer to the Galactic centre have more tangentially-biased (near-circular) motions than those farther away. This dip in $\beta(r)$ could be a reflection of the destruction of substructure by the central stellar disc, as discussed by Sawala et al. (2017) and Garrison-Kimmel et al. (2017b). To interpret this result for the MW satellite system, we move on to analyse the simulated systems in the APOSTLE and Auriga suites using the same methods.

5.3 Simulations

The posterior distributions for the $\beta(r)$ profiles of simulated MW analogs are shown in Figure 8 for APOSTLE and Figure 9 for Auriga. When considering all subhaloes with $V_{\max} > 5$ km s $^{-1}$ (top row), it is clear that the presence of a massive stellar disc affects the radial β profile. The $\beta(r)$ profiles for DMO hosts are nearly flat at $\beta \sim 0$, indicating isotropic motions at all radii. The inclusion of baryons in APOSTLE does not have a noticeable effect on the $\beta(r)$ profiles, which are very similar to DMO. Only the Auriga haloes exhibit a dip in the β profile near the centre, resulting from the massive central disc preferentially destroying radial orbits that pass near the galaxy.

However, the $\beta(r)$ profile estimates of our simulated systems are sensitive to the radial distribution of the subhaloes. Matching the radial distribution of subhaloes with that of the MW satellites, following the procedure described in Section 3.4, results in similar




Figure 8. Posterior distribution for the simulated APOSTLE systems assuming the parametrized $\beta(r)$ model. The left column corresponds dark matter-only runs of the APOSTLE volumes, while the right column to APOSTLE with baryons. From top to bottom the rows correspond to different subhalo populations: the full sample of subhaloes ($V_{\max} > 5 \text{ km s}^{-1}$), matching the radial distribution of MW satellites, matching both the radial distribution and number (see Section 3.4 for details on matching the radial distribution/abundance), and $M_{\text{star}} > 0$ (radial matching is not applied in the final row, $M_{\text{star}} > 0$). For comparison, the result for the MW satellite system is shown in grey, with a black dashed curve, in each panel. The different simulated systems are shown with different colors, with the shaded regions corresponding to the 16 – 84% confidence intervals. Note that each halo in an APOSTLE pair is treated separately, with the more massive halo denoted as (a) and the less massive one as (b). Additionally, note that AP-10 does not have a completed high-resolution DMO run, so there are two fewer curves in the left panel than the right. The horizontal black dashed line corresponds to the isotropic case $\beta = 0$. The DMO and APOSTLE systems generally do not exhibit a dip in the $\beta(r)$ profile, but as the simulations are convolved with the observed MW radial distribution and abundance of satellites (first three rows, top to bottom), some APOSTLE systems can exhibit a dip in their $\beta(r)$ profile.

$\beta(r)$ estimates in some systems that do not contain stellar discs; the estimates for some of the APOSTLE and even DMO systems become consistent with the results for the MW satellites (second row). This similarity is even more pronounced when the subhaloes are also selected to match the total number of MW satellites in addition to the radial distribution (third row).

While these corrections bring some DMO and APOSTLE estimates of $\beta(r)$ in line with that of the MW, the Auriga systems still provide the best agreement. There are many corrected DMO and APOSTLE systems that still have $\beta \sim 0$ near the centre, but only a few corrected Auriga profiles that do not have a dip in β . These results suggest that the dip in the β profile for the MW satellite system is likely best explained by effects due to the stellar disc, but also is sensitive to the radial distribution of tracers considered.

Finally, we also consider the sample of subhaloes in APOSTLE and Auriga that contain stars at $z = 0$ ($M_* > 0$, bottom

row). Nearly every $\beta(r)$ profile matches that of the full subhalo population, albeit with increased scatter due to a smaller sample size (see Table 2 for the number in each population). The only exception is the less massive halo of AP-01, whose β profile is shifted to lower values at all radii but maintains the same shape. The agreement between the $\beta(r)$ profiles when considering all subhaloes vs. the subsample containing stars suggests that the β profile (corrected for observational distance biases) traced by the MW satellites is likely indicative of the intrinsic profile for the MW, unaffected by the complex physics that dictate which subhaloes are populated by satellites (Sawala et al. 2015; Garrison-Kimmel et al. 2017a; Nadler et al. 2019).




Figure 9. Same as Figure 8, but for simulated Auriga systems. The dip in the $\beta(r)$ profile for Auriga hydrodynamic simulations is far more pronounced than in DMO or APOSTLE hydrodynamic simulations, likely due to the formation of a central stellar disc.

6 DISCUSSION

It is clear from our results that dwarf galaxy satellites closer to the centre of the Milky Way have tangentially-biased motions while those farther from the centre have radially-biased motions. In this Section we explore some interpretations of the $\beta(r)$ profile and place our results in the context of those from other tracers.

6.1 Stellar disc

The inclusion of baryonic processes in cosmological simulations has helped resolve a number of small-scale challenges to Λ CDM. A notable effect is the destruction of substructure due to the potential of a massive stellar disc. Brooks & Zolotov (2014) found that 6 of the 8 subhaloes in a DMO simulation that did not have a baryonic simulation counterpart had pericentric passages that took them within 30 kpc of the galaxy centre. Sawala et al. (2017) found that the presence of baryons near the centre of APOSTLE haloes reduces the number of subhaloes by factors of $\sim 1/4 - 1/2$ independently of subhalo mass but increasingly towards the host halo centre. Garrison-Kimmel et al. (2017b) found similar destruction of subhaloes in the Latte simulation suite and showed that simply embedding a central disc potential in DMO simulations reproduced these radial subhalo depletion trends, arguing that the additional

tidal field from the central galaxy is the primary cause of subhalo depletion (see also D’Onghia et al. 2010; Yurin & Springel 2015; Errani et al. 2017). We also note that Zhu et al. (2016) found similar results in simulations using the AREPO code, the same code with which Auriga was performed.

A central stellar disc, whether artificially embedded in DMO simulations or formed through the inclusion of baryonic physics, preferentially destroys subhaloes on radial orbits that pass close to the disc. The surviving population then has tangentially-biased motions compared to DMO (Sawala et al. 2017; Garrison-Kimmel et al. 2017b), which is expected to be reflected in a lower value of β . However, there is also a radial dependence of β which has not yet been explored; with increasing distance from the central galaxy the destructive effects of the disc potential weaken, causing the $\beta(r)$ profile to rise to $\beta \sim 0.5$ in the outer region ($100 \text{ kpc} < r < R_{\text{vir}}$) as subhaloes are more likely to be on their first infall (Diemand et al. 2004; Navarro et al. 2010). Additionally, as surviving massive satellites pass near the stellar disc, both experience a torque and exchange angular momentum, likely inducing further circularization of the surviving satellite orbits (Gómez et al. 2017a,b).

Our simulation results are consistent with this interpretation. When considering all subhaloes with $V_{\text{max}} > 5 \text{ km s}^{-1}$ in the

APOSTLE and Auriga suites (top rows of Figures 8 and 9), the $\beta(r)$ profiles for DMO and APOSTLE haloes, which have less massive central galaxies, are relatively constant with $\beta \gtrsim 0$. In stark contrast, the $\beta(r)$ profiles for Auriga haloes have $\beta \lesssim -0.5$ near the centre of the halo and increase to $\beta \gtrsim 0$ by ~ 200 kpc. These results are similar when considering, instead, subhaloes that contain stars at $z = 0$ (bottom rows of Figures 8 and 9).

The Auriga simulations produce stellar discs that are massive, thin, and radially extended, like that of the MW, while APOSTLE forms less massive host galaxies with weaker discs. This distinction impacts the orbital distribution of subhaloes and results in the Auriga subhaloes showing a variation of β similar to that of the MW satellite system.

However, it is worth noting the possibility that not all of the subhalo disruption is due to the physical effects of the stellar disc. van den Bosch et al. (2018) and van den Bosch & Ogiya (2018) raise concerns about artificial subhalo disruption due to numerical effects, suggesting that tracking subhalo disruption requires many more particles than required for typical simulation convergence tests. For example, van den Bosch et al. (2018) find that orbits passing within 10-20% of the virial radius of a host may require $N > 10^6$ particles for an accurate treatment. More work may be required to understand the differences between these results and those from typical convergence tests. This will, in turn, inform our understanding of how much subhalo disruption is due to physical effects of the stellar disc vs. numerical effects and how this impacts the inferred $\beta(r)$ profile.

6.2 The radial distribution

This clean interpretation of a $\beta(r)$ profile caused by the tidal field of the central galaxy becomes muddier when accounting for the observed radial distribution of the Milky Way satellites. We know the current census of satellites is incomplete both radially, due to surface brightness and luminosity selection effects, and in area on the sky, as less than half of the sky has been covered by surveys capable of finding ultra-faint satellite galaxies (Kim et al. 2018; Newton et al. 2018). This results in a satellite sample that is more centrally concentrated than those found in M31 and in cosmological simulations (Yniguez et al. 2014; Graus et al. 2018; Kelley et al. 2018, however, see Li et al. 2019), giving greater weight to satellites located closer to the centre.

We attempt to account for this by matching the abundance and/or radial distribution of simulated subhaloes with $V_{\max} > 5 \text{ km s}^{-1}$ to that of the MW satellites. Applying these corrections to simulated MW-mass systems tends to lower β estimates relative to when the full population is used (see Figures 8 and 9, middle two rows). As a result, the inferred $\beta(r)$ profiles for some DMO and APOSTLE haloes, which do not contain massive central galaxies, are consistent with that of the MW satellites.

This is not to say that the impact of the central disc is not crucial to explaining the anisotropy of the MW satellite system. As shown in Figures 8 and 9, for any given selection criterion applied to the subhaloes the dip in the $\beta(r)$ profiles is most prominent for the Auriga host haloes, which have massive central discs. However, a more complete analysis of the MW disc's impact on the $\beta(r)$ profile would require understanding the true selection function for the MW satellites. Knowing this selection function, combined with a more detailed modeling procedure (e.g. using a distribution function in action/angle coordinates, as in Vasiliev 2019, would lend greater insight to orbital properties at the expense of assuming a




Figure 10. Comparison of MW $\beta(r)$ results between different tracers. The grey points and contours correspond to studies using globular clusters, while the black points correspond to studies using halo stars. The blue contours are the results from this work. The horizontal black dashed line corresponds to the isotropic case, $\beta = 0$.

MW potential), would enhance future studies of the $\beta(r)$ profile for the MW satellite galaxies.

6.3 Comparison with other tracers

Finally, it is interesting to compare our $\beta(r)$ results with those using other tracers of the MW potential (see Figure 10). Cunningham et al. (2018) used HST proper motions of $N \sim 200$ halo stars in four different fields, spherically averaging to find $\beta \sim 0.5 - 0.7$ over $19 < r < 29$ kpc. This is higher than the values found in several other studies using line-of-sight velocities alone to constrain the anisotropy of the stellar halo, which tend to prefer isotropic or tangentially-biased β values (for a summary, see Figure 6 in Cunningham et al. 2016).

Using MW globular clusters (GCs), Sohn et al. (2018) estimated $\beta = 0.609^{+0.130}_{-0.229}$ over $10.6 < r < 39.5$ kpc with proper motions from HST while Watkins et al. (2019) found $\beta = 0.48^{+0.15}_{-0.20}$ over $2.0 < r < 21.1$ kpc with proper motions from *Gaia* DR2. These two values suggest a trend for the GC orbits to become more radially-biased with increasing distance. Indeed, Vasiliev (2019) modelled $\beta(r)$ for the MW GCs using a distribution function-based method in action/angle space and found a steady increase from $\beta \sim 0.0$ at 0.5 kpc to $\beta \sim 0.6$ at 200 kpc (see Figure 7 of Vasiliev 2019), consistent with these other results.

The dip in the $\beta(r)$ profile for the MW globular cluster system detected by Vasiliev (2019) is qualitatively similar in shape to what we find for the MW dwarf satellites, but is very different both in characteristic radial scale and in overall amplitude. At $r > 100$ kpc the inferred values of β are similar. The dip in the globular cluster profile may possibly be attributed to the accretion history vs. in situ formation (Fall & Zhang 2001; Prieto & Gnedin 2008). It is also possible that both the globular clusters and stellar halo are remnants of stars previously attached to subhaloes on radial orbits, which are preferentially destroyed by the stellar disc, and maintain the anisotropy of their progenitors (see Bullock et al. 2001; Bullock & Johnston 2005; Bell et al. 2008 for halo stars; Peebles 1984; Moore et al. 2006; Boylan-Kolchin 2017 for globular clusters). This potential connection between different tracers of the MW

$\beta(r)$ profile merits further modeling, possibly with a joint analysis of Milky Way halo stars, globular clusters, and dwarf galaxies.

7 SUMMARY

In this work we have analysed the kinematics of 38 Milky Way satellites focusing on an estimate of the velocity anisotropy parameter, β , and its dynamical interpretation. Utilizing the latest satellite proper motions inferred from *Gaia* DR2 data, we modelled β using a likelihood method and, for the first time, estimated $\beta(r)$ for the MW satellite system. We then compared these results with expectations from Λ CDM using the APOSTLE and Auriga simulation suites. A summary of our main results is as follows:

- The MW satellites have overall tangentially-biased motions, with best-fitting constant $\beta = -1.02^{+0.37}_{-0.45}$. By parametrizing $\beta(r)$, we find that the anisotropy profile for the MW satellite system increases from $\beta \sim -2$ at $r \sim 20$ kpc to $\beta \sim 0.5$ at $r \sim 200$ kpc, indicating that satellites closer to the Galactic centre have tangentially-biased motions while those farther out have radially-biased motions.

- Comparing these results with the APOSTLE and Auriga galaxy formation simulations, we find that satellites surrounding the massive and radially extended stellar discs formed in Auriga have similar $\beta(r)$ profiles to that of the MW, while the weaker discs in APOSTLE produce profiles that are similar to those from DMO simulations. This suggests that the central stellar disc affects the $\beta(r)$ profile of the MW satellite system by preferentially destroying radial orbits that pass near the disc, as discussed by [Sawala et al. \(2017\)](#) and [Garrison-Kimmel et al. \(2017b\)](#).

- However, when matching the radial distributions of simulated subhaloes to that of the MW satellites, some of the inferred $\beta(r)$ profiles for APOSTLE and even DMO haloes also can match the MW data. This implies that the partial sky coverage and the increasing incompleteness with distance of the currently available satellite sample significantly impair the ability of our scheme to robustly estimate the true $\beta(r)$ profile.

The difficulty in interpreting the inferred $\beta(r)$ profile may also be alleviated by more fully exploring the Milky Way's virial volume. [Newton et al. \(2018\)](#) expect that – assuming a MW halo mass of $1.0 \times 10^{12} M_{\odot}$ – there are 46^{+12}_{-8} ultra-faint ($8 < M_V \leq 3$) satellites and 61^{+37}_{-23} hyper-faint ($-3 < M_V \leq 0$) satellites within 300 kpc that are detectable. At least half of these satellites should be found by LSST within the next decade.

Obtaining proper motions for these faint and distant objects will be challenging but clearly possible, given the results already obtained for 7 satellites fainter than $M_V = -5$ and farther than $d_{\odot} = 100$ kpc. Furthermore, since the precision in proper motion measurements grows as the 1.5 power of the time baseline, the satellite proper motions from *Gaia* should be a factor 4.5 more precise after the nominal mission and possibly a factor 12 more precise after the extended mission (F18). Artificially scaling the observed proper motion errors by a factor of 4.5 results in a $\beta(r)$ profile that has a narrower confidence interval by a factor of ~ 1.5 and provides better agreement with the Auriga $\beta(r)$ profiles. With this improved dataset, future studies will be less limited by observational selection effects and be able to study in greater depth the impact of the central stellar disc on the $\beta(r)$ profile of the Milky Way satellite system.

ACKNOWLEDGEMENTS

We thank Marius Cautun for useful discussions regarding this work and an anonymous referee for helpful suggestions regarding its presentation, as well as Emily Cunningham and Eugene Vasiliev for sharing their data. This research made use of the Python Programming Language, along with many community-developed or maintained software packages including Astropy ([Astropy Collaboration et al. 2013, 2018](#)), corner.py ([Foreman-Mackey 2016](#)), emcee ([Foreman-Mackey et al. 2013](#)), Jupyter ([Kluyver et al. 2016](#)), Matplotlib ([Hunter 2007](#)), NumPy ([van der Walt et al. 2011](#)), Pandas ([McKinney 2010](#)), and SciPy ([Jones et al. 2001](#)). This research made extensive use of [arXiv.org](#) and NASA's Astrophysics Data System for bibliographic information.

AHR acknowledges support from a Texas A&M University Merit Fellowship. AF is supported by a European Union COFUND/Durham Junior Research Fellowship (under EU grant agreement no. 609412). ABP acknowledges generous support from the Texas A&M University and the George P. and Cynthia Woods Institute for Fundamental Physics and Astronomy. LES acknowledges support from DOE Grant de-sc0010813. CSF acknowledges a European Research Council (ERC) Advanced Investigator grant (DMIDAS; GA 786910). FAG acknowledges financial support from FONDECYT Regular 1181264, and funding from the Max Planck Society through a Partner Group grant. RG acknowledges support by the Deutsche Forschungsgemeinschaft (DFG) Research Centre Sonderforschungsbereich (SFB)-881 ‘The Milky Way System’ through project A1.

Part of the simulations of this paper used the SuperMUC system at the Leibniz Computing Centre, Garching, under the project PR85JE of the Gauss Centre for Supercomputing. This work was supported in part by the UK Science and Technology Facilities Council (STFC) ST/P000541/1. This work used the DiRAC Data Centric system at Durham University, operated by the ICC on behalf of the STFC DiRAC HPC Facility ([www.dirac.ac.uk](#)). This equipment was funded by BIS National E-infrastructure capital grant ST/K00042X/1, STFC capital grant ST/H008519/1, and STFC DiRAC Operations grant ST/K003267/1 and Durham University. DiRAC is part of the National E-Infrastructure.

REFERENCES

- Abbott T. M. C., et al., 2018, *The Astrophysical Journal Supplement Series*, **239**, 18
- Adén D., et al., 2009, *A&A*, **506**, 1147
- Astropy Collaboration et al., 2013, *A&A*, **558**, A33
- Astropy Collaboration et al., 2018, *AJ*, **156**, 123
- Bechtol K., et al., 2015, *ApJ*, **807**, 50
- Bell E. F., et al., 2008, *ApJ*, **680**, 295
- Bellazzini M., Ferraro F. R., Origlia L., Pancino E., Monaco L., Oliva E., 2002, *AJ*, **124**, 3222
- Bellazzini M., Gennari N., Ferraro F. R., 2005, *MNRAS*, **360**, 185
- Belokurov V., et al., 2007, *ApJ*, **654**, 897
- Benson A. J., Lacey C. G., Baugh C. M., Cole S., Frenk C. S., 2002, *MNRAS*, **333**, 156
- Binney J., 1980, *MNRAS*, **190**, 873
- Bland-Hawthorn J., Gerhard O., 2016, *Annual Review of Astronomy and Astrophysics*, **54**, 529
- Boettcher E., et al., 2013, *AJ*, **146**, 94
- Bonanos A. Z., Stanek K. Z., Szentgyorgyi A. H., Sasselov D. D., Bakos G. Á., 2004, *AJ*, **127**, 861
- Booth C. M., Schaye J., 2009, *MNRAS*, **398**, 53

- Bovy J., 2015, *The Astrophysical Journal Supplement Series*, 216, 29
- Boylan-Kolchin M., 2017, *MNRAS*, 472, 3120
- Boylan-Kolchin M., Bullock J. S., Kaplinghat M., 2011, *MNRAS*, 415, L40
- Boylan-Kolchin M., Bullock J. S., Kaplinghat M., 2012, *MNRAS*, 422, 1203
- Brooks A. M., Zolotov A., 2014, *ApJ*, 786, 87
- Bullock J. S., Boylan-Kolchin M., 2017, *Annual Review of Astronomy and Astrophysics*, 55, 343
- Bullock J. S., Johnston K. V., 2005, *ApJ*, 635, 931
- Bullock J. S., Kravtsov A. V., Weinberg D. H., 2000, *ApJ*, 539, 517
- Bullock J. S., Kravtsov A. V., Weinberg D. H., 2001, *ApJ*, 548, 33
- Caldwell N., et al., 2017, *ApJ*, 839, 20
- Callingham T. M., et al., 2019, *MNRAS*, 484, 5453
- Carlin J. L., Sand D. J., 2018, *ApJ*, 865, 7
- Carlin J. L., Grillmair C. J., Muñoz R. R., Nidever D. L., Majewski S. R., 2009, *ApJ*, 702, L9
- Carlin J. L., et al., 2017, *AJ*, 154, 267
- Cautun M., Frenk C. S., 2017, *MNRAS*, 468, L41
- Cautun M., Bose S., Frenk C. S., Guo Q., Han J., Hellwing W. A., Sawala T., Wang W., 2015, *MNRAS*, 452, 3838
- Collins M. L. M., Tollerud E. J., Sand D. J., Bonaca A., Willman B., Strader J., 2017, *MNRAS*, 467, 573
- Crain R. A., et al., 2015, *MNRAS*, 450, 1937
- Cunningham E. C., et al., 2016, *ApJ*, 820, 18
- Cunningham E. C., et al., 2018, arXiv e-prints, p. arXiv:1810.12201
- D'Onghia E., Springel V., Hernquist L., Keres D., 2010, *ApJ*, 709, 1138
- Dall'Ora M., et al., 2006, *ApJ*, 653, L109
- Dall'Ora M., et al., 2012, *ApJ*, 752, 42
- Deason A. J., Belokurov V., Evans N. W., An J., 2012, *MNRAS*, 424, L44
- Deason A. J., Van der Marel R. P., Guhathakurta P., Sohn S. T., Brown T. M., 2013, *ApJ*, 766, 24
- Deason A. J., Belokurov V., Koposov S. E., Lancaster L., 2018, *ApJ*, 862, L1
- Diemand J., Moore B., Stadel J., 2004, *MNRAS*, 352, 535
- Drlica-Wagner A., et al., 2015, *ApJ*, 813, 109
- Errani R., Peñarrubia J., Laporte C. F. P., Gómez F. A., 2017, *MNRAS*, 465, L59
- Fall S. M., Zhang Q., 2001, *ApJ*, 561, 751
- Fattahi A., et al., 2016, *MNRAS*, 457, 844
- Foreman-Mackey D., 2016, *The Journal of Open Source Software*, 24
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *Publications of the Astronomical Society of the Pacific*, 125, 306
- Frenk C. S., White S. D. M., 2012, *Annalen der Physik*, 524, 507
- Frenk C. S., Evrard A. E., White S. D. M., Summers F. J., 1996, *ApJ*, 472, 460
- Fritz T. K., Battaglia G., Pawlowski M. S., Kallivayalil N., van der Marel R., Sohn S. T., Brook C., Besla G., 2018, *A&A*, 619, A103
- Gaia Collaboration et al., 2018a, *A&A*, 616, A1
- Gaia Collaboration et al., 2018b, *A&A*, 616, A12
- Garofalo A., et al., 2013, *ApJ*, 767, 62
- Garrison-Kimmel S., Bullock J. S., Boylan-Kolchin M., Bardwell E., 2017a, *MNRAS*, 464, 3108
- Garrison-Kimmel S., et al., 2017b, *MNRAS*, 471, 1709
- Garrison-Kimmel S., et al., 2018, preprint, p. arXiv:1806.04143 (arXiv:1806.04143)
- Gillet N., Ocvirk P., Aubert D., Knebe A., Libeskind N., Yepes G., Gottlöber S., Hoffman Y., 2015, *ApJ*, 800, 34
- Gómez F. A., White S. D. M., Grand R. J. J., Marinacci F., Springel V., Pakmor R., 2017a, *MNRAS*, 465, 3446
- Gómez F. A., et al., 2017b, *MNRAS*, 472, 3722
- Grand R. J. J., et al., 2017, *MNRAS*, 467, 179
- Grand R. J. J., et al., 2018, *MNRAS*, 481, 1726
- Graus A. S., Bullock J. S., Kelley T., Boylan-Kolchin M., Garrison-Kimmel S., Qi Y., 2018, preprint, p. arXiv:1808.03654 (arXiv:1808.03654)
- Greco C., et al., 2008, *ApJ*, 675, L73
- Hargis J. R., et al., 2016, *ApJ*, 818, 39
- Hunter J. D., 2007, *Computing In Science & Engineering*, 9, 90
- Ibata R. A., et al., 2013, *Nature*, 493, 62
- Ibata N. G., Ibata R. A., Famaey B., Lewis G. F., 2014, *Nature*, 511, 563
- Ibata R. A., Famaey B., Lewis G. F., Ibata N. G., Martin N., 2015, *ApJ*, 805, 67
- Jones E., Oliphant T., Peterson P., et al., 2001, SciPy: Open source scientific tools for Python, <http://www.scipy.org/>
- Kallivayalil N., et al., 2018, *ApJ*, 867, 19
- Karczmarek P., et al., 2015, *AJ*, 150, 90
- Katz N., White S. D. M., 1993, *ApJ*, 412, 455
- Kelley T., Bullock J. S., Garrison-Kimmel S., Boylan-Kolchin M., Pawlowski M. S., Graus A. S., 2018, arXiv e-prints, p. arXiv:1811.12413
- Kim S. Y., Peter A. H. G., Hargis J. R., 2018, *Phys. Rev. Lett.*, 121, 211302
- Kirby E. N., Boylan-Kolchin M., Cohen J. G., Geha M., Bullock J. S., Kaplinghat M., 2013, *ApJ*, 770, 16
- Kirby E. N., Simon J. D., Cohen J. G., 2015, *ApJ*, 810, 56
- Kirby E. N., Cohen J. G., Simon J. D., Guhathakurta P., Thygesen A. O., Duggan G. E., 2017, *ApJ*, 838, 83
- Kluyver T., et al., 2016, in Loizides F., Schmidt B., eds, *Positioning and Power in Academic Publishing: Players, Agents and Agendas*. pp 87 – 90
- Klypin A., Kravtsov A. V., Valenzuela O., Prada F., 1999, *ApJ*, 522, 82
- Koch A., et al., 2009, *ApJ*, 690, 453
- Komatsu E., et al., 2011, *The Astrophysical Journal Supplement Series*, 192, 18
- Koposov S. E., et al., 2011, *ApJ*, 736, 146
- Koposov S. E., Belokurov V., Torrealba G., Evans N. W., 2015a, *ApJ*, 805, 130
- Koposov S. E., et al., 2015b, *ApJ*, 811, 62
- Koposov S. E., et al., 2018, *MNRAS*, 479, 5343
- Kroupa P., Theis C., Boily C. M., 2005, *A&A*, 431, 517
- Kuehn C., et al., 2008, *ApJ*, 674, L81
- Laevens B. P. M., et al., 2015, *ApJ*, 813, 44
- Li T. S., et al., 2018, *ApJ*, 857, 145
- Li M., Gao L., Wang J., 2019, *MNRAS*, 483, 2000
- Libeskind N. I., Frenk C. S., Cole S., Helly J. C., Jenkins A., Navarro J. F., Power C., 2005, *MNRAS*, 363, 146
- Loebman S. R., et al., 2018, *ApJ*, 853, 196
- Lynden-Bell D., 1976, *MNRAS*, 174, 695
- Martin N. F., Ibata R. A., Bellazzini M., Irwin M. J., Lewis G. F., Dehnen W., 2004, *MNRAS*, 348, 12
- Martin N. F., de Jong J. T. A., Rix H.-W., 2008, *ApJ*, 684, 1075
- Martin N. F., et al., 2015, *ApJ*, 804, L5
- Martínez-Vázquez C. E., et al., 2015, *MNRAS*, 454, 1509
- Massari D., Helmi A., 2018, *A&A*, 620, A155
- Mateo M., Olszewski E. W., Walker M. G., 2008, *ApJ*, 675, 201
- McConnachie A. W., 2012, *AJ*, 144, 4
- McKinney W., 2010, in van der Walt S., Millman J., eds, *Proceedings of the 9th Python in Science Conference*. pp 51 – 56
- McMillan P. J., 2017, *MNRAS*, 465, 76
- McMonigal B., et al., 2014, *MNRAS*, 444, 3139
- Medina G. E., et al., 2017, *ApJ*, 845, L10
- Moore B., Ghigna S., Governato F., Lake G., Quinn T., Stadel J., Tozzi P., 1999, *ApJ*, 524, L19
- Moore B., Diemand J., Madau P., Zemp M., Stadel J., 2006, *MNRAS*, 368, 563
- Moretti M. I., et al., 2009, *ApJ*, 699, L125
- Muñoz R. R., Geha M., Willman B., 2010, *AJ*, 140, 138
- Müller O., Pawlowski M. S., Jerjen H., Lelli F., 2018, *Science*, 359, 534
- Musella I., et al., 2009, *ApJ*, 695, L83
- Musella I., et al., 2012, *ApJ*, 756, 121
- Mutlu-Pakdil B., et al., 2018, *ApJ*, 863, 25
- Nadler E. O., Mao Y.-Y., Wechsler R. H., Garrison-Kimmel S., Wetzel A., 2018, *ApJ*, 859, 129
- Nadler E. O., Mao Y.-Y., Green G. M., Wechsler R. H., 2019, *ApJ*, 873, 34
- Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563
- Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493
- Navarro J. F., et al., 2010, *MNRAS*, 402, 21

- Newton O., Cautun M., Jenkins A., Frenk C. S., Helly J. C., 2018, *MNRAS*, **479**, 2853
- Oñorbe J., Garrison-Kimmel S., Maller A. H., Bullock J. S., Rocha M., Hahn O., 2014, *MNRAS*, **437**, 1894
- Okamoto T., Frenk C. S., 2009, *MNRAS*, **399**, L174
- Okamoto S., Arimoto N., Tolstoy E., Jablonka P., Irwin M. J., Komiyama Y., Yamada Y., Onodera M., 2017, *MNRAS*, **467**, 208
- Pace A. B., Li T. S., 2018, preprint, p. [arXiv:1806.02345](https://arxiv.org/abs/1806.02345) ([arXiv:1806.02345](https://arxiv.org/abs/1806.02345))
- Pakmor R., Marinacci F., Springel V., 2014, *ApJ*, **783**, L20
- Pakmor R., et al., 2017, *MNRAS*, **469**, 3185
- Pawlowski M. S., 2018, *Modern Physics Letters A*, **33**, 1830004
- Pawlowski M. S., Kroupa P., 2013, *MNRAS*, **435**, 2116
- Pawlowski M. S., Pflamm-Altenburg J., Kroupa P., 2012, *MNRAS*, **423**, 1109
- Peebles P. J. E., 1984, *ApJ*, **277**, 470
- Pietrzyński G., Górski M., Gieren W., Ivanov V. D., Bresolin F., Kudritzki R.-P., 2009, *AJ*, **138**, 459
- Planck Collaboration et al., 2014, *A&A*, **571**, A16
- Prieto J. L., Gnedin O. Y., 2008, *ApJ*, **689**, 919
- Rosas-Guevara Y. M., et al., 2015, *MNRAS*, **454**, 1038
- Sand D. J., Strader J., Willman B., Zaritsky D., McLeod B., Caldwell N., Seth A., Olszewski E., 2012, *ApJ*, **756**, 79
- Sawala T., et al., 2015, *MNRAS*, **448**, 2941
- Sawala T., et al., 2016, *MNRAS*, **457**, 1931
- Sawala T., Pihajoki P., Johansson P. H., Frenk C. S., Navarro J. F., Oman K. A., White S. D. M., 2017, *MNRAS*, **467**, 4383
- Schaye J., et al., 2015, *MNRAS*, **446**, 521
- Simon J. D., 2018, *ApJ*, **863**, 89
- Simon J. D., Geha M., 2007, *ApJ*, **670**, 313
- Simon J. D., et al., 2011, *ApJ*, **733**, 46
- Simon J. D., et al., 2015, *ApJ*, **808**, 95
- Simon J. D., et al., 2017, *ApJ*, **838**, 11
- Simpson C. M., Grand R. J. J., Gómez F. A., Marinacci F., Pakmor R., Springel V., Campbell D. J. R., Frenk C. S., 2018, *MNRAS*, **478**, 548
- Sohn S. T., Watkins L. L., Fardal M. A., van der Marel R. P., Deason A. J., Besla G., Bellini A., 2018, *ApJ*, **862**, 52
- Somerville R. S., 2002, *ApJ*, **572**, L23
- Spencer M. E., Mateo M., Walker M. G., Olszewski E. W., 2017, *ApJ*, **836**, 202
- Springel V., 2010, *MNRAS*, **401**, 791
- Springel V., Hernquist L., 2003, *MNRAS*, **339**, 289
- Springel V., et al., 2005, *Nature*, **435**, 629
- Stetson P. B., Fiorentino G., Bono G., Bernard E. J., Monelli M., Iannicola G., Gallart C., Ferraro I., 2014, *Publications of the Astronomical Society of the Pacific*, **126**, 616
- Torrealba G., Koposov S. E., Belokurov V., Irwin M., 2016a, *MNRAS*, **459**, 2370
- Torrealba G., et al., 2016b, *MNRAS*, **463**, 712
- Torrealba G., et al., 2018, *MNRAS*, **475**, 5085
- Vasiliev E., 2019, *MNRAS*, **484**, 2832
- Vivas A. K., et al., 2016, *AJ*, **151**, 118
- Voggel K., Hilker M., Baumgardt H., Collins M. L. M., Grebel E. K., Husemann B., Richtler T., Frank M. J., 2016, *MNRAS*, **460**, 3384
- Walker M. G., Mateo M., Olszewski E. W., 2009, *AJ*, **137**, 3100
- Walker M. G., Olszewski E. W., Mateo M., 2015, *MNRAS*, **448**, 2717
- Walker M. G., et al., 2016, *ApJ*, **819**, 53
- Walsh S. M., Willman B., Sand D., Harris J., Seth A., Zaritsky D., Jerjen H., 2008, *ApJ*, **688**, 245
- Watkins L. L., van der Marel R. P., Sohn S. T., Evans N. W., 2019, *ApJ*, **873**, 118
- Wetzell A. R., Hopkins P. F., Kim J.-h., Faucher-Giguère C.-A., Kereš D., Quataert E., 2016, *ApJ*, **827**, L23
- Willman B., Geha M., Strader J., Strigari L. E., Simon J. D., Kirby E., Ho N., Warren A., 2011, *AJ*, **142**, 128
- Yniguez B., Garrison-Kimmel S., Boylan-Kolchin M., Bullock J. S., 2014, *MNRAS*, **439**, 73
- Yurin D., Springel V., 2015, *MNRAS*, **452**, 2367
- Zhu Q., Marinacci F., Maji M., Li Y., Springel V., Hernquist L., 2016, *MNRAS*, **458**, 1559
- van den Bosch F. C., Ogiya G., 2018, *MNRAS*, **475**, 4066
- van den Bosch F. C., Ogiya G., Hahn O., Burkert A., 2018, *MNRAS*, **474**, 3043
- van der Marel R. P., et al., 2014, in Seigar M. S., Treuthardt P., eds, Vol. 480, *Structure and Dynamics of Disk Galaxies*. p. 43 ([arXiv:1309.2014](https://arxiv.org/abs/1309.2014))
- van der Walt S., Colbert S. C., Varoquaux G., 2011, *Computing in Science Engineering*, **13**, 22

APPENDIX A: MW SATELLITE PROPERTIES

Table A1 lists the observed properties of the MW satellites used throughout this analysis. Table A2 lists the Galactocentric spherical positions and velocities, along with corresponding uncertainties, of each satellite obtained by the Monte Carlo sampling detailed in Section 2.1.

The references in the last two columns of Table A1 are as follows: [1] [Gaia Collaboration et al. \(2018b\)](#); [2] [Simon \(2018\)](#); [3] [Kallivayalil et al. \(2018\)](#); [4] [Massari & Helmi \(2018\)](#); [5] [Pace & Li \(2018\)](#); [6] [Torrealba et al. \(2016b\)](#); [7] [Dall’Ora et al. \(2006\)](#); [8] [Martin et al. \(2008\)](#); [9] [Koposov et al. \(2011\)](#); [10] [Walsh et al. \(2008\)](#); [11] [Koch et al. \(2009\)](#); [12] [Kuehn et al. \(2008\)](#); [13] [Simon & Geha \(2007\)](#); [14] [Greco et al. \(2008\)](#); [15] [Sand et al. \(2012\)](#); [16] [Karczmarek et al. \(2015\)](#); [17] [McMonigal et al. \(2014\)](#); [18] [Walker et al. \(2009\)](#); [19] [Torrealba et al. \(2018\)](#); [20] [Li et al. \(2018\)](#); [21] [Muñoz et al. \(2010\)](#); [22] [Musella et al. \(2009\)](#); [23] [Caldwell et al. \(2017\)](#); [24] [Torrealba et al. \(2016a\)](#); [25] [Walker et al. \(2015\)](#); [26] [Bonanos et al. \(2004\)](#); [27] [Laevens et al. \(2015\)](#); [28] [Pietrzyński et al. \(2009\)](#); [29] [Koposov et al. \(2015a\)](#); [30] [Walker et al. \(2016\)](#); [31] [Adén et al. \(2009\)](#); [32] [Musella et al. \(2012\)](#); [33] [Koposov et al. \(2015b\)](#); [34] [Kirby et al. \(2015\)](#); [35] [Vivas et al. \(2016\)](#); [36] [Martin et al. \(2015\)](#); [37] [Koposov et al. \(2018\)](#); [38] [Mateo et al. \(2008\)](#); [39] [Stetson et al. \(2014\)](#); [40] [Spencer et al. \(2017\)](#); [41] [Bellazzini et al. \(2005\)](#); [42] [Moretti et al. \(2009\)](#); [43] [Collins et al. \(2017\)](#); [44] [Medina et al. \(2017\)](#); [45] [McConnachie \(2012\)](#); [46] [Simon et al. \(2015\)](#); [47] [Mutlu-Pakdil et al. \(2018\)](#); [48] [Martínez-Vázquez et al. \(2015\)](#); [49] [Belokurov et al. \(2007\)](#); [50] [Simon et al. \(2011\)](#); [51] [Boettcher et al. \(2013\)](#); [52] [Kirby et al. \(2013\)](#); [53] [Okamoto et al. \(2017\)](#); [54] [Kirby et al. \(2017\)](#); [55] [Carlin et al. \(2017\)](#); [56] [Bechtol et al. \(2015\)](#); [57] [Simon et al. \(2017\)](#); [58] [Drlica-Wagner et al. \(2015\)](#); [59] [Garofalo et al. \(2013\)](#); [60] [Dall’Ora et al. \(2012\)](#); [61] [Bellazzini et al. \(2002\)](#); [62] [Willman et al. \(2011\)](#).

Table A1. Properties of the satellites used in this analysis: RA, Dec, absolute magnitude, heliocentric distance, line-of-sight velocity, proper motion from Fritz et al. (2018), proper motion for the “gold” sample described in Section 2.2, and study used for the “gold” PM.

Satellite	RA [deg]	Dec [deg]	M_V [mag]	d_\odot [kpc]	v_\odot [km s ⁻¹]	$\mu_{\alpha^*}^{\text{F18}}$ [mas yr ⁻¹]	μ_δ^{F18} [mas yr ⁻¹]	$\mu_{\alpha^*}^{\text{gold}}$ [mas yr ⁻¹]	μ_δ^{gold} [mas yr ⁻¹]	PM _{gold}	Refs.
Aquarius II	338.481	-9.327	-4.36	107.9 ± 3.3	-71.1 ± 2.5	-0.252 ± 0.53	0.011 ± 0.452	-0.491 ± 0.306	-0.049 ± 0.266	[3]	[6]
Boötes I	210.015	14.512	-6.3	66 ± 3.0	102.2 ± 0.8	-0.554 ± 0.098	-1.111 ± 0.076	-0.459 ± 0.041	-1.064 ± 0.029	[1]	[7] [8] [9]
Boötes II	209.521	12.859	-2.3	42 ± 1.6	-117.1 ± 7.6	-2.686 ± 0.393	-0.53 ± 0.292	-2.517 ± 0.325	-0.602 ± 0.235	[2]	[10] [11]
Canes Venatici I	202.016	33.559	-8.6	210 ± 6	30.9 ± 0.6	-0.159 ± 0.1	-0.067 ± 0.064	[8] [12] [13]
Canes Venatici II	194.292	34.321	-4.6	160 ± 7	-128.9 ± 1.2	-0.342 ± 0.238	-0.473 ± 0.178	[13] [14] [15]
Carina I	100.407	-50.966	-8.6	105.6 ± 5.4	222.9 ± 0.1	0.485 ± 0.038	0.131 ± 0.038	0.495 ± 0.015	0.143 ± 0.014	[1]	[16] [17] [18]
Carina II	114.107	-57.999	-4.5	37.4 ± 0.4	477.2 ± 1.2	1.867 ± 0.085	0.082 ± 0.08	1.79 ± 0.06	0.01 ± 0.05	[4]	[19] [20]
Carina III	114.630	-57.900	-2.4	27.8 ± 0.6	284.6 ± 3.25	3.046 ± 0.132	1.565 ± 0.147	3.065 ± 0.095	1.567 ± 0.104	[3]	[19] [20]
Coma Berenices I	186.746	23.908	-3.8	42 ± 1.5	98.1 ± 0.9	0.471 ± 0.113	-1.716 ± 0.11	0.546 ± 0.092	-1.726 ± 0.086	[2]	[13] [21] [22]
Crater II	177.310	-18.413	-8.2	117.5 ± 1.1	87.5 ± 0.4	-0.184 ± 0.07	-0.106 ± 0.068	-0.246 ± 0.052	-0.227 ± 0.026	[3]	[23] [24]
Draco I	260.060	57.965	-8.75	76 ± 6	-291.0 ± 0.1	-0.012 ± 0.013	-0.158 ± 0.038	-0.019 ± 0.009	-0.145 ± 0.01	[1]	[25] [26]
Draco II	238.198	64.565	-2.9	20 ± 3.0	-347.6 ± 1.75	1.242 ± 0.282	0.845 ± 0.291	1.165 ± 0.26	0.866 ± 0.27	[3]	[24] [27]
Fornax	39.962	-34.511	-13.4	147 ± 9	55.3 ± 0.1	0.374 ± 0.035	-0.401 ± 0.035	0.376 ± 0.003	-0.413 ± 0.003	[1]	[18] [28]
Grus I	344.176	-50.163	-3.4	120.2 ± 11.1	-140.5 ± 2.0	-0.261 ± 0.178	-0.437 ± 0.242	-0.25 ± 0.16	-0.47 ± 0.23	[5]	[29] [30]
Hercules	247.763	12.787	-6.6	132 ± 6	45.2 ± 1.09	-0.297 ± 0.123	-0.329 ± 0.1	[31] [32]
Horologium I	43.882	-54.119	-3.5	79 ± 7	112.8 ± 2.55	0.891 ± 0.105	-0.55 ± 0.099	0.95 ± 0.07	-0.55 ± 0.06	[5]	[33]
Hydra II	185.425	-31.985	-4.8	151 ± 8	303.1 ± 1.4	-0.416 ± 0.523	0.134 ± 0.426	-0.417 ± 0.402	0.179 ± 0.339	[3]	[34] [35] [36]
Hydrus I	37.389	-79.309	-4.71	27.6 ± 0.5	80.4 ± 0.6	3.733 ± 0.052	-1.605 ± 0.05	3.761 ± 0.029	-1.371 ± 0.027	[3]	[37]
Leo I	152.122	12.313	-12.03	258.2 ± 9.5	282.5 ± 0.1	-0.086 ± 0.069	-0.128 ± 0.071	-0.097 ± 0.056	-0.091 ± 0.047	[1]	[38] [39]
Leo II	168.370	22.152	-9.6	233 ± 15	78.5 ± 0.6	-0.025 ± 0.087	-0.173 ± 0.09	-0.064 ± 0.057	-0.21 ± 0.054	[1]	[40] [41]
Leo IV	173.233	-0.540	-4.97	154 ± 5	132.3 ± 1.4	-0.59 ± 0.534	-0.449 ± 0.362	[13] [42]
Leo V	172.784	2.222	-4.4	173 ± 5	172.1 ± 2.2	-0.097 ± 0.56	-0.628 ± 0.307	[43] [44]
LMC	80.894	-69.756	-18.1	51.0 ± 2.0	262.2 ± 3.4	1.85 ± 0.03	0.24 ± 0.03	[1]	[45]
Pisces II	344.634	5.955	-4.1	183 ± 15	-226.5 ± 2.7	-0.108 ± 0.647	-0.586 ± 0.502	[15] [34]
Reticulum II	53.949	-54.047	-3.6	31.4 ± 1.4	64.8 ± 0.5	2.398 ± 0.053	-1.319 ± 0.059	2.36 ± 0.05	-1.32 ± 0.06	[5]	[46] [47]
Sagittarius I	283.831	-30.545	-13.5	26 ± 2.0	140.0 ± 2.0	-2.736 ± 0.036	-1.357 ± 0.036	-2.692 ± 0.001	-1.359 ± 0.001	[1]	[45]
Sculptor	15.039	-33.709	-10.7	83.9 ± 1.5	111.4 ± 0.1	0.084 ± 0.036	-0.133 ± 0.0356	0.082 ± 0.005	-0.131 ± 0.004	[1]	[18] [48]
Segue 1	151.763	16.074	-1.5	23 ± 2	208.5 ± 0.9	-1.697 ± 0.198	-3.501 ± 0.178	-1.867 ± 0.11	-3.282 ± 0.102	[2]	[49] [50]
Segue 2	34.817	20.175	-2.5	36.6 ± 2.45	-40.2 ± 0.9	1.656 ± 0.161	0.135 ± 0.113	1.01 ± 0.14	-0.48 ± 0.18	[4]	[51] [52]
Sextans	153.268	-1.620	-9.3	92.5 ± 2.2	224.2 ± 0.1	-0.438 ± 0.045	0.055 ± 0.045	-0.496 ± 0.025	0.077 ± 0.02	[1]	[18] [53]
SMC	13.187	-72.829	-16.8	64.0 ± 4.0	145.6 ± 0.6	0.797 ± 0.03	-1.22 ± 0.03	[1]	[45]
Triangulum II	33.322	36.172	-1.2	28.4 ± 1.6	-381.7 ± 1.1	0.588 ± 0.194	0.554 ± 0.169	0.588 ± 0.187	0.554 ± 0.161	[3]	[54] [55]
Tucana II	343.060	-58.570	-3.9	57.5 ± 5.3	-129.1 ± 3.5	0.91 ± 0.069	-1.159 ± 0.082	0.91 ± 0.06	-1.16 ± 0.08	[5]	[30] [56]
Tucana III	359.150	-59.600	-2.4	25 ± 2	-102.3 ± 0.4	-0.025 ± 0.049	-1.661 ± 0.049	-0.03 ± 0.04	-1.65 ± 0.04	[5]	[57] [58]
Ursa Major I	158.685	51.926	-6.75	97.3 ± 5.85	-55.3 ± 1.4	-0.683 ± 0.1	-0.72 ± 0.135	-0.659 ± 0.093	-0.635 ± 0.131	[2]	[13] [59]
Ursa Major II	132.874	63.133	-3.9	34.7 ± 2.1	-116.5 ± 1.9	1.691 ± 0.064	-1.902 ± 0.075	1.661 ± 0.053	-1.87 ± 0.065	[2]	[13] [60]
Ursa Minor	227.242	67.222	-8.4	76 ± 4	-246.9 ± 0.1	-0.184 ± 0.044	0.082 ± 0.042	-0.182 ± 0.01	0.074 ± 0.008	[1]	[61]
Willman 1	162.341	51.053	-2.7	38 ± 7	-12.8 ± 1.0	0.199 ± 0.194	-1.342 ± 0.37	0.382 ± 0.119	-1.152 ± 0.216	[2]	[8] [62]

Table A2. Galactocentric positions and velocities for each coordinate using [Fritz et al. \(2018\)](#) proper motions (LMC and SMC proper motions from [Gaia Collaboration et al. 2018b](#)). Quoted values are the 16th, 50th, and 84th percentiles.

Satellite	r [kpc]	θ [deg]	ϕ [deg]	v_r [km s ⁻¹]	v_θ [km s ⁻¹]	v_ϕ [km s ⁻¹]
Aquarius II	105.2 ^{+3.3} _{-2.2}	145.0 ^{+0.1} _{-0.1}	61.7 ^{+0.2} _{-0.2}	43.2 ^{+19.5} _{-20.2}	-282.3 ^{+222.8} _{-239.9}	29.0 ^{+268.5} _{-252.0}
Boötes I	63.6 ^{+2.9} _{-2.8}	13.6 ^{+0.3} _{-0.3}	-2.9 ^{+0.1} _{-0.1}	97.5 ^{+3.3} _{-3.4}	110.7 ^{+26.2} _{-25.4}	-126.0 ^{+32.4} _{-33.9}
Boötes II	39.8 ^{+1.6} _{-1.5}	10.4 ^{+0.4} _{-0.5}	-13.3 ^{+0.6} _{-0.7}	-48.1 ^{+16.5} _{-16.2}	-311.1 ^{+70.7} _{-68.3}	-246.9 ^{+70.5} _{-67.4}
Canes Venatici I	209.8 ^{+5.9} _{-5.8}	9.8 ^{+0.0} _{-0.0}	86.0 ^{+0.5} _{-0.4}	83.5 ^{+3.4} _{-3.4}	93.1 ^{+80.0} _{-85.8}	83.8 ^{+79.6} _{-86.1}
Canes Venatici II	160.9 ^{+6.7} _{-7.0}	8.8 ^{+0.1} _{-0.1}	130.4 ^{+0.7} _{-0.6}	-93.2 ^{+8.4} _{-8.9}	-144.2 ^{+127.7} _{-137.4}	135.0 ^{+171.6} _{-181.1}
Carina I	106.9 ^{+5.8} _{-5.2}	111.8 ^{+0.0} _{-0.0}	-104.6 ^{+0.3} _{-0.2}	-4.6 ^{+3.0} _{-2.9}	-165.0 ^{+22.0} _{-24.4}	-22.5 ^{+17.4} _{-18.2}
Carina II	38.3 ^{+0.3} _{-0.4}	106.7 ^{+0.0} _{-0.0}	-103.0 ^{+0.2} _{-0.2}	204.4 ^{+4.3} _{-4.3}	-228.6 ^{+15.5} _{-15.6}	195.6 ^{+14.3} _{-13.6}
Carina III	29.0 ^{+0.6} _{-0.9}	106.1 ^{+0.0} _{-0.0}	-107.2 ^{+0.4} _{-0.4}	46.6 ^{+6.4} _{-6.6}	-383.6 ^{+18.9} _{-20.1}	36.1 ^{+16.9} _{-18.1}
Coma Berenices I	43.2 ^{+1.3} _{-1.4}	14.9 ^{+0.4} _{-0.4}	-158.2 ^{+0.6} _{-0.6}	28.9 ^{+7.7} _{-5.1}	-252.6 ^{+21.9} _{-24.0}	104.5 ^{+25.3} _{-25.3}
Crater II	116.4 ^{+1.1} _{-1.1}	47.5 ^{+0.0} _{-0.0}	-82.3 ^{+0.1} _{-0.1}	-83.7 ^{+3.4} _{-3.6}	-77.0 ^{+38.5} _{-37.3}	-24.2 ^{+40.5} _{-39.0}
Draco I	75.9 ^{+6.0} _{-5.6}	55.3 ^{+0.0} _{-0.0}	93.8 ^{+0.6} _{-0.6}	-88.5 ^{+2.8} _{-2.9}	124.1 ^{+5.7} _{-5.7}	-50.7 ^{+14.2} _{-13.5}
Draco II	22.4 ^{+2.8} _{-2.7}	52.5 ^{+1.2} _{-1.0}	125.4 ^{+3.5} _{-3.0}	-154.4 ^{+12.4} _{-12.1}	300.5 ^{+26.0} _{-23.8}	-68.8 ^{+31.6} _{-32.6}
Fornax	149.5 ^{+8.6} _{-9.0}	153.9 ^{+0.1} _{-0.1}	-129.1 ^{+0.3} _{-0.4}	-40.9 ^{+1.5} _{-1.5}	-104.5 ^{+30.8} _{-32.9}	112.5 ^{+28.4} _{-28.3}
Grus I	116.3 ^{+11.5} _{-10.6}	151.6 ^{+0.3} _{-0.3}	-24.5 ^{+0.3} _{-0.4}	-203.2 ^{+7.0} _{-7.0}	-187.5 ^{+133.9} _{-135.2}	123.7 ^{+110.8} _{-113.6}
Hercules	126.3 ^{+6.1} _{-6.0}	51.3 ^{+0.1} _{-0.1}	30.9 ^{+0.1} _{-0.1}	150.5 ^{+3.3} _{-3.2}	-10.0 ^{+73.4} _{-70.7}	-54.8 ^{+73.4} _{-68.2}
Horologium I	79.4 ^{+7.0} _{-7.0}	144.4 ^{+0.1} _{-0.1}	-99.1 ^{+0.9} _{-1.0}	-33.7 ^{+5.2} _{-4.9}	-193.4 ^{+46.6} _{-49.0}	0.6 ^{+42.0} _{-40.1}
Hydra II	148.3 ^{+7.9} _{-8.3}	58.9 ^{+0.0} _{-0.0}	-67.6 ^{+0.2} _{-0.2}	129.3 ^{+21.1} _{-21.2}	-164.7 ^{+281.8} _{-287.8}	-187.8 ^{+392.9} _{-396.2}
Hydrus I	25.7 ^{+0.5} _{-0.5}	129.9 ^{+0.0} _{-0.0}	-84.4 ^{+0.5} _{-0.5}	-57.2 ^{+3.2} _{-3.3}	-328.9 ^{+9.6} _{-9.9}	-161.9 ^{+9.0} _{-8.6}
LMC	50.3 ^{+2.0} _{-1.9}	123.3 ^{+0.0} _{-0.0}	-90.7 ^{+0.4} _{-0.5}	63.1 ^{+4.3} _{-4.3}	-310.3 ^{+18.4} _{-18.0}	-40.9 ^{+8.3} _{-9.3}
Leo I	261.9 ^{+9.3} _{-9.3}	41.7 ^{+0.0} _{-0.0}	-135.8 ^{+0.1} _{-0.1}	168.6 ^{+3.1} _{-3.1}	24.4 ^{+66.4} _{-74.3}	-71.4 ^{+101.2} _{-102.0}
Leo II	235.2 ^{+15.2} _{-14.5}	24.1 ^{+0.1} _{-0.1}	-142.7 ^{+0.2} _{-0.2}	18.5 ^{+3.8} _{-4.0}	-72.0 ^{+86.1} _{-88.7}	-14.0 ^{+112.0} _{-101.7}
Leo IV	154.7 ^{+5.1} _{-4.9}	33.8 ^{+0.0} _{-0.0}	-99.8 ^{+0.2} _{-0.2}	13.8 ^{+20.8} _{-21.5}	321.4 ^{+265.2} _{-270.9}	-183.6 ^{+372.5} _{-393.5}
Leo V	174.0 ^{+4.6} _{-5.0}	31.9 ^{+0.0} _{-0.0}	-102.9 ^{+0.2} _{-0.2}	40.5 ^{+19.9} _{-18.9}	225.1 ^{+373.5} _{-351.4}	236.3 ^{+369.4} _{-381.6}
Pisces II	181.8 ^{+14.6} _{-14.5}	137.4 ^{+0.0} _{-0.0}	83.1 ^{+0.3} _{-0.3}	-79.7 ^{+24.4} _{-24.1}	173.6 ^{+475.1} _{-471.3}	-356.8 ^{+537.6} _{-533.7}
Reticulum II	32.8 ^{+1.4} _{-1.3}	136.9 ^{+0.2} _{-0.2}	-115.3 ^{+0.8} _{-0.9}	-99.8 ^{+3.0} _{-3.1}	-215.8 ^{+18.7} _{-19.1}	56.4 ^{+10.7} _{-9.8}
SMC	61.3 ^{+4.2} _{-3.8}	136.9 ^{+0.1} _{-0.1}	-66.8 ^{+0.6} _{-0.7}	-5.6 ^{+2.3} _{-2.4}	-245.3 ^{+26.3} _{-27.0}	-67.5 ^{+16.2} _{-17.1}
Sagittarius I	18.3 ^{+2.0} _{-2.0}	110.6 ^{+0.8} _{-0.8}	8.2 ^{+0.3} _{-0.3}	140.0 ^{+2.3} _{-2.6}	-275.2 ^{+17.2} _{-17.0}	-53.3 ^{+21.6} _{-21.4}
Sculptor	84.0 ^{+1.5} _{-1.5}	172.5 ^{+0.1} _{-0.1}	-119.7 ^{+0.9} _{-0.8}	75.0 ^{+1.6} _{-1.6}	169.6 ^{+13.9} _{-14.4}	-72.8 ^{+16.2} _{-15.1}
Segue 1	27.9 ^{+1.9} _{-1.9}	50.4 ^{+0.9} _{-0.8}	-153.8 ^{+0.8} _{-0.9}	116.8 ^{+5.9} _{-5.7}	142.1 ^{+34.5} _{-31.3}	142.0 ^{+34.0} _{-30.2}
Segue 2	42.4 ^{+2.4} _{-2.3}	121.9 ^{+0.3} _{-0.3}	156.1 ^{+0.4} _{-0.3}	72.8 ^{+4.6} _{-4.5}	-214.7 ^{+26.2} _{-25.2}	9.5 ^{+28.7} _{-30.1}
Sextans	95.5 ^{+2.3} _{-2.2}	49.3 ^{+0.0} _{-0.0}	-122.2 ^{+0.1} _{-0.1}	79.2 ^{+2.6} _{-2.6}	-12.2 ^{+17.6} _{-18.3}	-239.5 ^{+24.9} _{-24.9}
Triangulum II	34.7 ^{+1.6} _{-1.6}	109.2 ^{+0.2} _{-0.3}	150.0 ^{+0.5} _{-0.4}	-255.2 ^{+4.9} _{-5.0}	-175.7 ^{+23.6} _{-24.0}	-122.6 ^{+25.5} _{-25.5}
Tucana II	54.0 ^{+5.2} _{-5.2}	148.1 ^{+0.6} _{-0.5}	-40.9 ^{+0.9} _{-1.1}	-187.6 ^{+4.6} _{-4.1}	48.6 ^{+18.3} _{-18.1}	-208.0 ^{+40.5} _{-43.1}
Tucana III	23.0 ^{+1.9} _{-1.9}	154.5 ^{+0.2} _{-0.3}	-80.4 ^{+3.4} _{-4.2}	-228.1 ^{+2.3} _{-2.2}	28.2 ^{+19.3} _{-21.2}	48.3 ^{+11.0} _{-13.6}
Ursa Major I	102.1 ^{+3.7} _{-3.9}	39.0 ^{+0.2} _{-0.2}	161.9 ^{+0.1} _{-0.1}	11.5 ^{+3.7} _{-3.8}	165.7 ^{+54.0} _{-54.7}	206.1 ^{+62.6} _{-60.8}
Ursa Major II	40.9 ^{+2.2} _{-2.0}	58.8 ^{+0.3} _{-0.3}	158.6 ^{+0.3} _{-0.3}	-57.7 ^{+2.8} _{-2.8}	-280.3 ^{+23.7} _{-24.1}	32.6 ^{+19.9} _{-17.0}
Ursa Minor	78.2 ^{+4.0} _{-4.0}	46.5 ^{+0.1} _{-0.1}	112.9 ^{+0.4} _{-0.4}	-71.4 ^{+2.7} _{-2.8}	136.9 ^{+12.8} _{-12.3}	-11.5 ^{+18.3} _{-18.4}
Willman 1	42.5 ^{+6.8} _{-6.5}	41.9 ^{+1.6} _{-1.3}	164.5 ^{+0.9} _{-0.7}	17.8 ^{+6.3} _{-6.4}	-106.4 ^{+47.6} _{-59.3}	-55.5 ^{+75.0} _{-59.6}