

 Open access • Journal Article • DOI:10.1080/07352689.2017.1360112

## The Worldwide Spread, Success, and Impact of Ragweed (*Ambrosia* spp.)

— [Source link](#) 

Chiara Montagnani, Rodolfo Gentili, Matt Smith, Maria Francesca Guarino ...+1 more authors

**Institutions:** University of Milano-Bicocca, University of Worcester

**Published on:** 21 Sep 2017 - Critical Reviews in Plant Sciences (Taylor & Francis)

**Topics:** Ambrosia artemisiifolia

Related papers:

- [Biological Flora of the British Isles: \*Ambrosia artemisiifolia\*.](#)
- [Effects of climate change and seed dispersal on airborne ragweed pollen loads in Europe](#)
- [Let native species take their course: \*Ambrosia artemisiifolia\* replacement during natural or “artificial” succession](#)
- [The historical spread of \*Ambrosia artemisiifolia\* L. in France from herbarium records](#)
- [Ophraella communa, the ragweed leaf beetle, has successfully landed in Europe: fortunate coincidence or threat?](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/the-worldwide-spread-success-and-impact-of-ragweed-ambrosia-whlz5a8chb>

***The worldwide spread, success and impact of ragweed (*Ambrosia spp.*)***

Montagnani C.<sup>a\*</sup>, Gentili R.<sup>a</sup>, Smith M.<sup>b</sup>, Guarino M. F.<sup>a</sup>, Citterio S.<sup>a</sup>

<sup>a</sup> Department of Earth and Environmental Sciences, University of Milano-Bicocca, Piazza della Scienza 1, 20126

Milano, Italy

chiara.montagnani@unimib.it, rodolfo.gentili@unimib.it, m.guarino2@campus.unimib.it,

sandra.citterio@unimib.it

<sup>b</sup> Institute of Science and the Environment, University of Worcester, Worcester, United Kingdom

aeromattsmith@gmail.com

\*Corresponding author: Chiara Montagnani, Department of Earth and Environmental Sciences,  
University of Milano-Bicocca, Piazza della Scienza 1, 20126 Milano, Italy, email:  
chiara.montagnani@unimib.it

## **Abstract**

*Ambrosia* species represent one of the most problematic groups of invasive weeds around the world. The ease they are introduced and spread in new countries, their generalist ecological requirements and functional traits facilitate their invasion and subsequent naturalization in new areas. All of these aspects contribute to increasing their global social and economic impact, which is mostly related to pollen allergy. Here we analyse available scientific publications about *Ambrosia artemisiifolia*, *A. psilostachya*, *A. tenuifolia* and *A. trifida*, with the aim of defining the current level of knowledge and summarizing important data that is currently scattered throughout the literature. Specifically, we analysed the following: (1) their current global distribution and current stage of invasion; (2) traits and requirements promoting their introduction, reproductive success and adaptation to climate and environment in the non-native range; as well as (3) current knowledge about allergens and elements increasing their impact.

**Key words:** *Ambrosia artemisiifolia*, *A. psilostachya*, *A. tenuifolia*, *A. trifida*, invasive alien plants, pollen allergy.

---

## Contents

|   |    |
|---|----|
| I. Introduction.....  | 4  |
| II. Literature screening .....  | 6  |
| III. Species considered and their description.....                                    | 6  |
| IV. Geographical barriers .....   | 7  |
| <i>A. Native range</i> .....  | 7  |
| <i>B. From native to invasive range: pathways of introduction</i> .....               | 8  |
| <i>C. Global distribution: current status and invasion history</i> .....              | 11 |
| V. Environmental barriers .....   | 15 |
| <i>A. Habitat types and environmental matching</i> .....                              | 16 |
| <i>B. Climate matching and temperature tolerance</i> .....                            | 20 |
| <i>C. Moisture and soil types tolerance</i> .....                                     | 22 |
| <i>D. Light requirements and tolerance</i> .....                                      | 26 |
| <i>E. Plant traits involved in resistance and resilience</i> .....                    | 27 |
| 1. Resistance.....  | 28 |
| 2. Resilience .....   | 30 |
| VI. Reproductive and dispersal barriers .....   | 37 |
| <i>A. Pollination</i> .....   | 38 |
| <i>B. Seed and propagule pressure</i> .....   | 38 |
| <i>C. Dispersal</i> .....   | 40 |
| <i>D. Human-mediated dispersal pathways (medium- and long-distance vectors)</i> ..... | 42 |
| VII. Allergenic impact and environment.....   | 44 |
| <i>A. Pollen characteristics related to allergy</i> .....                             | 44 |
| <i>B. Pollen Allergens</i> .....  | 45 |
| <i>C. Allergenic impact and environment</i> .....                                     | 47 |
| VIII. Conclusion.....   | 49 |
| REFERENCES.....   | 50 |

## **I. Introduction**

There are over 40 species in the genus *Ambrosia* L. (Asteraceae) (Rich, 1994; Makra *et al.*, 2015; [www.theplantlist.org](http://www.theplantlist.org)), most of which are native to the Americas. Over the last 200 years, human impacts on land use (i.e. urbanization, the intensification of farming practices and increased transportation networks) have had serious effects on the distribution and ecology of several *Ambrosia* species. In particular, the introduction of the following species in non-native continents: *A. artemisiifolia* L. (common or short ragweed), *A. trifida* L. (giant ragweed), *A. tenuifolia* Spreng. (slender or slim-leaf burr ragweed) and *A. psilostachya* DC. (Western or perennial ragweed).

These *Ambrosia* (ragweed) species have been introduced into new countries since the 19<sup>th</sup> century, especially *A. artemisiifolia* which has quickly become an invasive species (Smith *et al.* (2013) and references therein) and is presently a species of concern for public health in both its native and invasive ranges because of its highly allergenic pollen. In North America, *Ambrosia* pollen is the second most important cause of seasonal allergic rhinitis and asthma, affecting more than 15 million people (about 20-25% of the United States population), with a prevalence of about 45% in atopic individuals (Wopfner *et al.*, 2005; Kats and Carey 2014). In Europe, *Ambrosia* has become a serious problem in the past decades, contributing to an evident increase in respiratory allergic reactions in areas where it is distributed (D'Amato, 1992; D'Amato, 2007). It is therefore recognized globally that *Ambrosia* species represent one of the most problematic groups of invasive weeds.

Biological invasions can be represented as a chronological series of decisive stages that can allow or halt the entry and establishment of an organism in a new range (Blackburn *et al.*, 2011; Richardson and Pysek, 2012). A non-native organism needs to overcome geographical, environmental and reproductive barriers to establish in a new area, and some factors and traits

can be more meaningful than others in predicting or explaining its success or failure during the process of introduction, establishment and spread (Van Kleunen *et al.*, 2015). Expressing the invasion process formulaically as "introduction - naturalization - invasion *continuum*", Richardson and Pysek (2012) stressed the importance of focusing on "naturalization", which is the fundamental preliminary step before invasion. According to their review, this "naturalization" is understudied, but its predictors/mediators can be more robust than those formulated for the invasion phase as they depend on factors less unpredictable and highly context-dependent.

On this basis, our review examines the main predictors of introduction-naturalization of the most widespread *Ambrosia* species at a global level, considering the environmental requirements and traits discussed and reported in scientific literature. After starting with a brief description of the taxa considered, this review then follows the logical sequence of the *continuum* in order to examine factors and traits involved in overcoming:

- *Geographical barriers*. Mediators of the introduction of species into new areas: native extent of occurrence, pathways of introduction and their effectiveness in terms of space (extent of the invasive range) and time (protraction of invasion).
- *Environmental barriers*. Predictors of the likelihood of persistence of *Ambrosia* species: environmental and climatic (broad-scale) matching between native and invasive range, requirements and tolerance to the main environmental abiotic factors (temperature, soil, light) in sensitive phases of the life cycle, resilience (strategies) to disturbances and competition.
- *Reproductive and dispersal barriers*. Predictors of successful propagation and spread of species, such as: specific pollinators, reproduction and dispersal strategies, propagule pressure.

The review ends with an overview of the allergenic impact of ragweed and how environmental factors and plant traits influence the magnitude of pollinosis.

## **II. Literature screening**

Google Scholar, Web of Science and Scopus databases were consulted to identify scientific literature about the *Ambrosia* genus. However, given the availability of interesting information also in additional web repositories, technical reports and online databases (national/regional floras, CABI, IUCN, EPPO, DAISIE, HEAR) were examined.

## **III. Species considered and their description**

The *Ambrosia* species considered in this review are presently introduced in more than ten countries: *A. artemisiifolia*, *A. trifida*, *A. psilostachya* and *A. tenuifolia*. All *taxa* are herbaceous or slightly suffruticose species. *Ambrosia artemisiifolia* and *A. trifida* are annual plants, while *A. psilostachya* and *A. tenuifolia* are perennials. Besides the presence of different belowground organs, diagnostic elements are mainly leaves, whose shape and clefts, as well as the presence of petioles, are useful in identifying different species. Leaves are variously pubescent or glabrous. They are monoic species with inflorescences of unisexual heads. All *taxa* produce 1-seeded cypselae and its size and coat ornaments may differ: *A. trifida* produces the biggest seeds and, among others species, spines may differ in number and bluntness.

*A. trifida* is an easily identifiable *taxon*, whose height (up to 4 m) and shape of leaves are unmistakable. However, the determination of other *taxa* can often be hard owing to a high variability in leaf or seed shape. Furthermore, the taxonomy is complicated by the possible presence of hybrids between *A. artemisiifolia* and *A. psilostachya* (*A. x intergradiens*; Wagner and Beals, 1958) and *A. artemisiifolia* and *A. trifida* (*Ambrosia x helenae*; Wagner, 1958; Strother, 2006).

Recently, the SMARTER project (COST Action FA1203) brought together a European team of botanists from different countries who reviewed the discriminating characters among species to provide a proper key of identification; as the most recent and reliable source, their findings relevant for this section are summarized in Table 1 and they are also available at <http://internationalragweedsociety.org/>

#### **IV. Geographical barriers**

Geographical barriers are the first obstacle an organism has to overcome before reaching a new range (Richardson *et al.*, 2000; Blackburn *et al.*, 2011). All the *taxa* considered here are native to the Americas, where the highest number of species of the genus have occurred: the genus appears to have originated and diversified from arid and semi-arid regions in Southwestern North America (Payne, 1970; Gerber *et al.*, 2011).

From the current distribution of ragweed, it is clear that these species have often successfully crossed the natural borders of their broad native ranges (Fig. 1). Pathways of introduction are linked to involuntary human actions and the flux of propagules has been in operation for at least a century. Here the pathways of introduction from native to invasive ranges are explained and the present global distribution is discussed, starting from a description of the native environments of the single species.

##### *A. Native range*

*Ambrosia artemisiifolia* is native to North America and is presently widespread in the United States and Canada except for Yukon and Nunavut (Strother, 2006; Essl *et al.*, 2015). In its native area, it was first recorded before 1838 in the United States (Kazinczi *et al.*, 2008) and in 1822 in Canada (Bassett and Crompton, 1975; Mitich, 1996; Lavoie *et al.*, 2007). It is, however, difficult to understand its primitive range of distribution because its spread has been human-mediated for a long time (McAndrews, 1988). According to Bassett and Crompton (1982) the



species spread widely with the increase of settlements of white men in North America. As a consequence of this, and due to taxonomic disagreements, there are several incongruities among floras about the native or introduced status of *A. artemisiifolia* in a number of countries, above all regarding Central and South America (Fig. 1).

Like *A. artemisiifolia*, *A. trifida* is native to North America. According to Bazzaz (1979), it was found in repeatedly disturbed ground only in the Midwestern and Eastern United States. However, based on more recent sources, *A. trifida* occurs in a wider range, covering almost all the USA (except for Nevada) and Canada (except for the Northwest Territories, Nunavut, and Yukon (Strother, 2006); it moved into Canada from the South, following the retreat of the last glacial ice (Fig. 1) (Bassett and Crompton, 1982).

*Ambrosia psilostachya* is also native to Western North America. As with *A. trifida*, it migrated to Canada after the glacial retreat, colonizing the Eastern Canada, where it has been present for a considerable time (Fig. 1) (Mitich, 1996). Bovey (1966) confirmed that *A. psilostachya* is widely distributed from California, Texas, Mexico, Idaho and Saskatchewan eastward to Illinois and Louisiana, on pasture land in Nebraska but especially in the Rocky Mountain States (Bovey, 1966). According to Strother (2006), its native range in the United States includes almost all States (except for Maryland, Delaware, and New Jersey) and the Southern part of Canada (from Columbia to Nova Scotia and Newfoundland).

In contrast to the other three species, *A. tenuifolia* is native to temperate South American countries, particularly Brazil, Paraguay, Uruguay, Argentina, and probably Perù (Fig. 1) (Randall, 2012).

#### *B. From native to invasive range: pathways of introduction*

Due to their ethnobotanical value, *Ambrosia* spp. have been used in traditional medicine since ancient times. In North and Central America, *A. artemisiifolia*, *A. psilostachya* and *A. trifida* were medicinal plants for Native Americans (Chamberlin, 1911; Shemluck, 1982; Mamedov *et*

*al.*, 2015) and today some of them are still used (i.e. *A. psilostachya*, see Gioanetto *et al.*, 2010). Moreover, *A. trifida* was domesticated by Indigenous North Americans, who collected and planted seeds mainly for alimentary purposes (Simon, 2009; Journey, 2012), In South America, predominantly in the Southern Cone, *A. tenuifolia* has also been a traditional medicinal plant (Mongelli *et al.*, 1996; Del Vitto, 1997; Trillo *et al.*, 2014). Thus, owing to the long-lasting utilization of ragweed in traditional medicine and uses, it is likely that European colonizers carried seeds to their countries by growing plants in botanical gardens (Essl. *et al.*, 2015). However, the scientific or ethnobotanical interest was probably not the main root of introduction for these species, as the amounts of seeds or plants moved were small. Ragweed plants are not suitable for flower market trading or collection, and so it is likely that their massive expansion followed involuntary human pathways. Today, owing to the frequency and abundance of ragweeds in anthropic environments in their native range, it is widely accepted that seeds and/or propagules of plants have been unintentionally transported outside the Americas by human activities along trade routes.

The introduction of *A. artemisiifolia* into different parts of the world has been ascribed to contaminated seed lots of grain, vegetables (e.g. potatoes), seed for forage or oil-seeds (e.g. sunflower (Genton *et al.*, 2005; Chauvel *et al.*, 2006; Smith *et al.*, 2013; Essl *et al.*, 2015) and also in bird food (Brandes and Nitzsche, 2006; Frick *et al.*, 2011), fodder, ship ballasts, and military movements (Kazinczi *et al.*, 2008; Gaudeul *et al.*, 2011). During the 20th century, particularly during the World Wars, *A. artemisiifolia* was introduced into Europe principally through agricultural products from several North American sources. Based on molecular studies, repeated introductions occurred during the invasion in different parts of the new range (Genton *et al.*, 2005; Kiss and Béres, 2006; Gaudeul *et al.*, 2011; Hodgins and Rieseberg, 2011; Smith *et al.*, 2013; Ciappetta *et al.*, 2016).

The seeds of *A. artemisiifolia* were not the only ragweed species to follow such routes. Frick *et al.* (2011) showed that *Ambrosia* sp. seeds, including *A. artemisiifolia* as well as other

ragweed species, occur in 21 to 75% of the bird feed products available on the German, Hungarian and Danish markets. For instance, *A. trifida* was introduced by imports of commercial grain and oil-seed, e.g. between North America and European countries, which repeatedly ensured the reinforcement of populations into new areas (Follak *et al.*, 2013). Military movements during World War II were also vectors of introduction for *A. trifida* in several sites in Northern Italy (Ardenghi and Polani, 2016).

The mechanism of introduction into the invasive range of *A. psilostachya* and *A. tenuifolia* was likely the same as those mentioned previously (Makra *et al.*, 2015). Data from Moskalenko (2001) and CABI (2017) revealed that Russia cereals coming from Canada were contaminated by seeds of *A. psilostachya*. Verloove (2016b) ascribed the arrival of *A. psilostachya* in Belgium to the American forces during the First World War and Parsons and Cuthbertson (2001) underlined the contribution of United States military movements during World War II to the spread of *A. psilostachya* in Australia and other parts of the world. These latter assumptions were mainly based on the fact that the species was unknown before the arrival of US troops. On the other hand, *A. tenuifolia*, was mentioned (Nelson, 1917) as a "ballast-plant", a species involuntary transported in solid sailing ballasts (currently replaced by water ballasts) and then released in new countries during the de-ballasting phase. This pathway was inferred by the author after the finding of *A. tenuifolia* and other non-native species in dumping areas near harbours in Oregon; nevertheless, this route is also plausible if past transoceanic travels to Europe are considered (Thellung, 1912). In any case, at present, the pathways of introduction of *A. psilostachya* and *A. tenuifolia* are of less certain than those of *A. artemisiifolia*, given the few studies available. Moreover, considering that the species have different dispersal strategies, additional vectors should be taken into consideration. Specifically, the reproduction strategy of *A. psilostachya* is mainly vegetative and the amount of seeds produced is quite small. Thus, the role of alternative propagules (i.e. rhizomes) and different

vectors in the global spread of the *taxon* should be considered in defining reliable pathways of introduction.

### *C. Global distribution: current status and invasion history*

Vectors for the spread of ragweed work in very effective ways. The percentage of countries where the species are native or alien and their status (casual, naturalized or invasive) are shown in Fig. 2. Fig. 2 was developed according to the database reported in Table 2.

At the moment three species out of four (*A. artemisiifolia*, *A. psilostachya*, *A. tenuifolia*) are present in every continent and the fourth (*A. trifida*) has colonized all continents except for Africa and Oceania. In its non-native range, *A. artemisiifolia* occurs in eighty countries (including those where its native status is uncertain) and it is classified as invasive in 32% of them. *Ambrosia psilostachya* is alien in almost forty countries, but, although it is naturalized in at least fourteen countries representing the 36% of its non-native range, only in seven (19%) does it show invasive behaviour. Also *A. trifida* has colonized almost forty countries, but it is included among invasive plants only in three of these (one doubtful), representing 7% of the total. To date, *A. tenuifolia* appears not to be an invasive *taxon* in its non-native range, consisting of fewer than fifteen countries, but it is naturalized in over half of the range of introduction (64%).

Without considering its cultivation in botanical gardens, which dates back to the 18<sup>th</sup> century, the first record of *A. artemisiifolia* outside its native range comes from the United Kingdom, where the species was recorded as casual in 1836 (Essl *et al.*, 2015). Then, in 1854, it was found in the Hawaiian Islands (Wagner *et al.*, 1990) and in the same decade it was recorded again in Europe, in Switzerland (Bullock *et al.*, 2012). Later, in 1860 and 1863, it was found respectively in Germany (Brandes and Nitzsche, 2006) and France (Chauvel *et al.*, 2006) (Western Europe). In all these countries the species is still present. The native or alien status of *A. artemisiifolia* is uncertain in many countries in the Americas. Essl *et al.* (2015) asserted that

*A. artemisiifolia* is surely alien to Argentina, Chile, Brazil, Bahamas and the island of Hispaniola. Villaseñor and Espinosa-Garcia (2004) listed *A. artemisiifolia* among the alien species in Mexico, while for Cuban populations there are clearly uncertainties (Acevedo-Rodríguez and Strong, 2012). Going back to Europe, from the second half of the 19<sup>th</sup> century onwards, the species has rapidly spread over all the continent. Records of *A. artemisiifolia* in Northern countries are later than Germany and France: Denmark in 1865 (Bullock *et al.*, 2012) and Sweden in 1866 (Anderberg, 2000a). In Eastern Europe, it appeared in 1873 in Poland (Tokarska-Guzik *et al.*, 2011) and ten years later in the Czech Republic (Bullock *et al.*, 2012). In Southern Europe the first signs of introduction probably occurred 1879 in Croatia (Galzina *et al.*, 2010) and then in Italy (1902; Gentili *et al.*, 2016). Newly colonized countries have continued to be recorded also in recent times (e.g. Greece) (Greuter and Raus, 2008).

Concerning Asia, the first record dates back to 1877 and is from Japan, where *A. artemisiifolia* was found as a casual (naturalization stage was recorded around 1925; <https://www.nies.go.jp/biodiversity/invasive/DB/detail/80400e.html>), while in China the species was found later (1930s) (Qin *et al.*, 2014). The invasive behaviour of *A. artemisiifolia* was reported by Washitani (2004) as beginning in the 1960s. In Taiwan, South Korea and Turkey, *A. artemisiifolia* was found later in the 1990s, whereas in other Asian countries (e.g. Armenia, Kazakhstan, Iran and India) information about the introduction time is not available.

In other parts of the world, Quézel and Santa (1963) reported the presence of *A. artemisiifolia* in Africa in Algerian flora and Lawlree (1947) asserted that the species was found there later than 1890. The species was recorded in other African countries more recently and, at the moment, it is also naturalized in Egypt (Boulos, 2002). Moreover it appears that the species is expanding toward Southern Africa (Botswana, South Africa, and Swaziland) (Setshogo, 2005; Henderson, 2007; Randall, 2012; Swaziland's Alien Plants Database at: <http://www.sntc.org.sz/alienplants/index.asp>). Skalova *et al.* (2015) also reported the presence of *A. artemisiifolia* in Madagascar, while Kull *et al.* (2012) mention only *A. maritima* as an

introduced and naturalized *taxon* on the island. Finally, in Oceania the species officially appeared in 1908 in Australia, spreading rapidly only after 1940s (Parsons and Cuthbertson, 2001), and in 1911 in New Zealand (Webb *et al.*, 1988).

Europe was the first continent that *A. trifida*, was introduced. It is recorded in the 17<sup>th</sup> century in botanical gardens (e.g. 1699 in the United Kingdom - see Online Atlas British and Irish Flora <http://www.brc.ac.uk/plantatlas/index.php?q=plant/ambrosia-trifida>). However, the oldest collection in the wild was reported from Western Europe where *A. trifida* was recorded in Belgium in 1829 and has been found with more continuity from 1896 onwards (Verloove, 2016a). It was later collected in Germany in 1877 (Follak *et al.*, 2013). In Northern Europe, the species was found first in Ireland (1894) and, in 1897, in the United Kingdom (Rich, 1994; Sell and Murrell, 2006), Finland (Lampinen and Lahti, 2016), Switzerland (Follak *et al.*, 2013) and Latvia (Gudzinskas *et al.*, 1993). In Southern Europe, the species was first recorded in South Tyrol in 1899 (Chauvel *et al.*, 2015) and recorded in 1909 in Northwest Italy (Vignolo Lutati, 1935); in other countries it has been found in more recent times (1982 in Serbia and 1983 in Spain) (Amor Morales *et al.*, 2012; Follak *et al.*, 2013). In Eastern Europe *A. trifida* was mentioned much later than in other parts of the continent, starting from 1960 (e.g. Czech Republic) (Pyšek *et al.*, 2012) up to the 1980s and beyond.

In comparison to *A. artemisiifolia*, *A. trifida* appeared later in Asia. In 1935 it was observed in China (Qin *et al.*, 2014), and it was noted in Japan almost twenty years later (1952; Invasive species of Japan: <https://www.nies.go.jp/biodiversity/invasive/DB/detail/80410e.html>). In other countries in Asia, records have been mostly since 1970 (South Korea) and the following decades (Israel, India). In Central America, *A. trifida* has been introduced into Mexico (Villaseñor and Espinosa-Garcia, 2004). Unlike common ragweed, giant ragweed rarely shows invasive behaviour (e.g. China and Japan) and it is often casual (e.g. Austria and British Isles) (Essl and Rabitsch, 2002; Reynolds, 2002; EPPO, 2016), which suggests that its persistence in some areas is only possible due to repeated introductions (Follak *et al.*, 2013).

*Ambrosia psilostachya* has a wider invasive range than *A. trifida*, but it is not listed alongside giant and common ragweed as invasive species by EPPO (EPPO, 2016). The species occurs in all continents and is naturalized in a large number of countries, although it is not as aggressive as *A. artemisiifolia* (Table 2). In any case, it should be remembered that *A. psilostachya* has not been as deeply studied and is often confused with *A. artemisiifolia* or other *taxa*. For these reasons, its distribution may be affected by misidentification, and dates of introduction are often uncertain. On the American continent, outside its recognized native range, the position of *A. psilostachya* as an independent *taxon* is discussed (Pruski, 2017). On the Hawaiian Islands it is listed as a quarantine weed (Randall, 2012), while it is reported as both wild and cultivated in Guadeloupe (Hibon, 1942). In Europe, it has been found in the United Kingdom since the 1880s (Rich, 1994), where today it is naturalized. However, it is not as widespread in the UK as it is in Germany, where it appeared a few years later in 1897 (NetPhyD: Deutschlandflora WebGIS). At the beginning of the 20th century, *A. psilostachya* was found in Hungary (Puc, 2004) and then in the Netherlands (1905; Odé and Beringen, 2017b), where according to Van Denderen *et al.* (2010) it is the only well established *taxon*. In Southern Europe, it first appeared in Italy at the end of the 1920s (Vignolo Lutati, 1935). In France, colonization probably dates back to the same time, although this may be influenced by misidentification of the specimens (Hibon, 1942; Queney, 1942). New European records of the species have also been collected recently, since the 1980s for instance in Spain, Finland, Czech Republic and Greece (Anderberg, 2005; Amor Morales *et al.*, 2012; Pyšek *et al.*, 2012; Von Raab Straube and Raus, 2016). Concerning Africa, information about the arrival of the *taxon* is quite fragmented. The oldest date of collection (1916) relates to Algeria (Maire, 1928), although an earlier record by Battandier (1888) discusses the presence of a perennial plant growing on maritime sands "with *Ambrosia* leaves" different from *A. maritima* (the only ragweed identified in other Algerian localities, probably native to the Old World (Montagnani *et al.*, 2017). *Ambrosia psilostachya* has also been alien to Moroccan flora since the 1990s (Tanji, 2005). In

Southern Africa, it is a weed of sugar cane fields in Mauritius (Macdonald *et al.*, 2003) and it is naturalized in different areas (Germishuizen and Meyer, 2003; SANBI, 2015a). In Australia, the plant was found in 1922 (Parsons and Cuthbertson, 2001). Records from Asia are quite recent, from the 1990s in India (Ramachandra Prasad *et al.*, 2013) to 2000 in Taiwan (Tseng and Peng, 2004), and the plant is often naturalized and shows an invasive behaviour in several countries (e.g. Japan and India).

*Ambrosia tenuifolia* is widespread at global level, but is usually much more localized than the other species outside its native range. Fairly close to its native area, *A. tenuifolia* has been introduced into Louisiana (North America), Puerto Rico and Chile, where it was identified for the first time in 1923 (Ugarte *et al.*, 2011; USDA - NRCS, 2017; Acevedo-Rodriguez and Strong, 2012). In Europe, according to available data, the oldest record (1839) of *A. tenuifolia* is from France (Thellung, 1912). Successive records date one century later - 1935 in Italy (Vignolo Lutati, 1935) and 1954 in Spain (Amor Morales *et al.*, 2012). According to Randall (2012), *A. tenuifolia* also occurs in the United Kingdom, but there is no bibliographic evidence of this. In Asia, the collections of *A. tenuifolia* are few and quite recent: in Israel in 1991 (Waisel *et al.*, 2008) and Turkey in 2000 (Behçet, 2004; Özhatay and Kültür, 2006). In Australia and New Zealand, the presence of the species has been confirmed since 1932 and 1950 (Parsons and Cuthbertson, 2001; Howell and Sawyer, 2006) while in Africa, the species is only present in South Africa (Germishuizen and Meyer, 2003; SANBI, 2015b).

## **V. Environmental barriers**

In this section, the main environmental requirements and plant traits, predictor of the likelihood of persistence of the species in new ranges, are reported and discussed. Slatyer *et al.* (2013) found a positive relationship between niche breadth and range size, suggesting that a wide tolerance to abiotic conditions facilitates occupancy of a larger area, and that habitat breadth is a good predictor of a wide distribution. Environmental matching is important along the



naturalization-invasion continuum (Richardson and Pysek, 2012). Such generalist, common-habitat colonizer, species are highly competitive and tolerant to disturbance and have great potential to become invasive (Volta *et al.*, 2013). Looking at the global distribution of ragweed species (Fig.1), they have generally found suitable conditions to persist and spread, often becoming naturalised (Fig.2). Thus, overcoming environmental barriers appears to be a solvable issue for ragweed species.

#### *A. Habitat types and environmental matching*

Before describing and discussing the habitats elected by the ragweed species in both their native and invasive ranges (Table 3), it is necessary to point out that it is currently quite difficult to define the native habitat of *A. artemisiifolia* with any certainty, because, as already mentioned, its distribution has been human-mediated for such a long time (McAndrews, 1988).

It has been suggested that *A. artemisiifolia* should be native to the Great Plains (Hodgins and Rieseberg 2011; Hodgins *et al.*, 2013). However, both in the United States and in its invasive range *A. artemisiifolia* has rarely been found in natural habitats, such as prairies, while it is abundant and often invasive in ruderal ones (roadside verges, wastelands, railway embankments, construction sites, quarries etc.), at the edge of croplands or in arable fields and also on riverbanks (Bassett and Crompton, 1975; Fumanal *et al.*, 2008a; Milakovic *et al.*, 2014; Essl *et al.*, 2015; Gentili *et al.*, 2016). Consistent with this, floras and specialized sources indicate synantropic environments as the main habitats for common ragweed (Basset and Crompton, 1975; Smith *et al.* 2013).

It is likely that common ragweed has shifted gradually from its primary habitat to ruderal areas and croplands, following the advance of human settlements into unexploited American lands, when synantropic environments became more frequent (Basset and Crompton, 1975; Smith *et al.* 2013). In support of this hypothesis, the analysis of herbarium records by Lavoie *et al.* (2007) showed that in Quebec common ragweed spread along rivers at first, and only later entered agricultural fields.

In the last thirty years in the USA (native range), *A. trifida* has also shown the tendency to colonize cultivated fields (e.g. corn, soybean and cotton), probably as local adaptation of the species that originally lived in riparian (riverbanks, floodplains) or non-riparian edge habitats near cultivated areas or railroads and wastelands (Basset and Crompton, 1982; Regnier *et al.*, 2016). On floodplains, where floods occasionally occur, it does not grow at the lowest elevation but dominates in communities located 60 cm above the water level (Menges and Waller, 1983). Outside of North America, *A. trifida* lives along rivers and generally shares the ruderal behaviour of *A. artemisiifolia*, occurring especially in cultivated fields, along railways (Gudzinskis, 1993; Chauvel *et al.*, 2015), on maritime docks (e.g. Britain) (Rich, 1994) and fluvial ports (e.g. along Rhine and Elbe rivers in Europe) (Chauvel *et al.*, 2015). Nevertheless, in France, Follak *et al.* (2013) and Chauvel *et al.* (2015) found that *A. trifida* is more often recorded in ruderal places and along railways than in native elective riverine habitats.

Regarding *A. psilostachya* and *A. tenuifolia*, elective habitats are more recognizable than the ragweed species previously described. According to Albertson (1937) *A. psilostachya* "invades the short grasses from the disturbed places along the slopes". In its native range in North America, *A. psilostachya* shares ruderal habits with *A. artemisiifolia*, as it is common in open disturbed habitats such as abandoned fields, vacant lots and along transportation corridors (Basset and Crompton, 1975). In contrast to *A. artemisiifolia*, however, it is also common in semi-natural/natural environments: it is a typical forb of tallgrass prairies (temperate grasslands) of the American Great Plains (Reece *et al.*, 2004) and a weed in pastures and rangelands where it is favoured by overgrazing and fire (Baker and Guthery, 1990; Abrams, 1988; Vermeire and Gillen, 2000; Funderburg *et al.*, 2014). It is a sand-loving species and colonizes dry sand prairies on sand hills (Hart and Gleason, 1907; Hulett *et al.*, 1988; Ebinger *et al.*, 2006; Uresk, 2012). It also colonizes coastal dunes, mainly secondary dunes and vegetated flats behind them (Carls *et al.*, 1991) mostly where there is a certain grade of human impact (e.g. vehicle passage) (Stephenson, 1999). It is also listed among halophytic plants living

in harsh environments such as inland saline plains (Flores-Olvera *et al.*, 2016) and is referred to as a colonizer of riparian habitats, where it is considered a "mesoriparian plant", i.e. not typical of the wettest areas of riverbeds (Stromberg, 2013). Contrary to the other ragweed species, *A. psilostachya* also lives in the understory of non-dense woodland (e.g. *Pinus ponderosa* forests) (Bojorquez Tapia *et al.*, 1990). There is little information in the literature concerning the habitats of *A. psilostachya* outside of its native range. However, it suggests that this species, as in its native range, frequently colonizes coastal areas, dunes, sandy soils usually exposed to human impacts (Rich, 1994; Mandrioli, 1998; Weeda, 2010; Del Vecchio *et al.*, 2015; Fried *et al.*, 2015), rivers and ruderal habitats (Amor Morales *et al.*, 2012; Ardenghi and Polani, 2016). In a recently colonized area of India, *A. psilostachya* has also been observed in croplands and pastures (Ramachandra Prasad *et al.*, 2013).

As with the congeners examined previously, in South America, *A. tenuifolia* is a plant typical of open habitats and, like *A. psilostachya*, it is native to grasslands traditionally subjected to disturbing factors that in this case are ascribable to grazing and periodic flooding (as the result of heavy rainfall, flat topography and poor drainage). In Argentina, it represents a characterizing element of the flooding Pampa grasslands, as one of the co-dominant taxa of one of the most widespread plant communities of the area (e.g. communities characterized by *Piptochaetium montevidense*, *Ambrosia tenuifolia*, *Eclipta bellidioides* and *Mentha pulegium*) (Burkart *et al.*, 1990; Insausti *et al.*, 1995; Insausti and Grimoldi, 2006). The typical community of *A. tenuifolia* usually lives in raised flat lowlands, less subjected to inundation. However, *A. tenuifolia* can also dominate communities typical of more humid conditions, localized along river valleys, drainage basins, and coastal salty lagoons (Burkart *et al.*, 1990). It is also a characteristic element of coastal dune vegetation (Fontana, 2005; Marcomini and López, 2013). In Argentina, Marcomini *et al.* (2016) found it in stable dune systems, occurring between dunes where the vegetation cover is more relevant and dominated by *Cortaderia selloana* (*cortaderal* community). As in the Pampean Plains, these environments are subjected to periodic floods,

although they are also subjected to drought (Marcomini *et al.*, 2016). Furthermore, *A. tenuifolia* has been listed in Paraguay as an agricultural weed (De Egea *et al.*, 2016). Outside its native range, specifically Spain, *A. tenuifolia* lives in habitats very similar to those colonized by *A. psilostachya* (Amor Morales *et al.* (2012) and in other European countries the two species can be found in the same sites, for instance sharing an halophytic behaviour (e.g. in some areas of Italy) (Mandrioli *et al.*, 1998). Additional findings in Turkey revealed that *A. tenuifolia* can be also found in orchards and cultivated fields (tomatoes, cucumbers, wheat) and that it could prefer humid ruderal places (Behçet, 2004; Özhatay and Kültür, 2006).

Overall, analysis of the spectrum of colonized environments shows that a limited shift of habitat types between native and invasive ranges can be observed. All species are in fact typical of naturally or artificially disturbed open areas, both in America and in the rest of their acquired range. However, all species are more frequently present in natural environments in their native ranges, likely representing their original habitats. It is difficult to have an idea of the primary environments of *A. artemisiifolia*, whereas *A. trifida* spreads from riparian habitats, and the perennial *A. tenuifolia* and *A. psilostachya* originated from temperate grasslands and colonize inland and coastal sand dunes. On the whole, it is clear that all these species spread gradually from their primary habitat to synanthropic environments. It follows that although the spreading of each of these species requires specific conditions, they are capable of quickly shifting their habitat in changing circumstances and taking advantage of environmental disturbance. Being tolerant to disturbance, these common-habitat colonizer species have a great potentiality as invader. They can be considered pioneer species naturally colonizing harsh environments and ready to spread in ruderal habitats, where conditions are maintained suitable mainly by human action.

### *B. Climate matching and temperature tolerance*

Climate matching is a basic requirement for persistence in a new area. According to their global distribution, all these species come from temperate areas (Fig.1). Although with several exceptions, they mainly "move" from/to warm temperate climate regions, generally avoiding equatorial, arid and snow climates (Fig.1). Petitpierre *et al.* (2012) observed a limited shift of climatic niche between the native and adventive range of *A. artemisiifolia*. Analogously, a limited shift also appears valid for the other ragweed species on the basis of our preliminary and basic comparison (Fig.1). In keeping with these observations, many studies demonstrate the influence of climate on ragweed germination, growth and reproduction. Regarding this, *A. artemisiifolia* has been the most widely studied species. About germination, it has been reported that *A. artemisiifolia* seeds can germinate in a wide range of temperatures; the minimum temperature of germination ranges from 3.4 to 3.6° C (Essl *et al.*, 2015) while germination decreases up to 40°C (Bullock *et al.*, 2012). Nevertheless, Leiblein-Wild *et al.* (2014) observed differences between native and introduced populations: in the invasive range, seeds generally have a larger mass and can germinate faster under a wider range of conditions. The authors attributed this better performance to favourable biotic and abiotic factors occurring in the invasive range and speculated about a possible case of Evolution in Increased Competitive Ability (EICA). In any case, *A. artemisiifolia* seeds follow a quite complicated cycle of primary/secondary dormancy and they need an exposure to winter temperature or stratification in lighting conditions to break primary dormancy (Baskin and Baskin, 1980). In sites where the growing season is too short for seed maturation (e.g. Northern Europe) or seasonal temperatures are too high for vernalization (e.g. some areas of the Mediterranean basin), the species cannot become naturalized and occurs just in few small ephemeral populations (Dahl *et al.*, 1999; Kazinczi *et al.*, 2008; Van Denderen *et al.*, 2010; Makra *et al.*, 2014; Smith *et al.*, 2013).

Seasonal temperature variations also play an important role for *A. trifida* (Davis, 1930; Bazzaz, 1979) and *A. tenuifolia* (Insausti *et al.*, 1995), whose seeds need low temperatures to

germinate (primary/secondary dormancy cycle). The range of germination temperatures for *A. trifida* is wide (from 4 to 41°C, with an optimum between 10 to 24°C), but only if soil moisture conditions are suitable (17% to 55% soil moisture, with an optimum at 20 to 30%) (Abul-Fatih and Bazzaz, 1979a; Ballard *et al.*, 1996). For *A. tenuifolia*, the lack of alternating temperatures prevents seed germination (Insausti *et al.*, 1995).

The climatic requirements for *A. psilostachya* are not clear (Basset and Crompton, 1975). Martison *et al.* (2011) identified, among climatic and soil variables, mean annual temperature as the factor that mostly contributes to explaining the distribution of *A. psilostachya* in the United States, whereas in Europe Rasmussen *et al.* (2017) found that minimum temperature to be highly influential. The life cycle of the species in relation to seasonality in its native habitat (e.g. snowing, freezing winter and summer drought), indicates that the seeds of *A. psilostachya* should also be characterized by dormancy and they may need a vernalization phase to germinate (Baskin and Baskin, 2014). Nevertheless, the propagation of *A. psilostachya* is mainly vegetative, so if the root system can survive, environmental limits on germination are less important. According to Rich (1994) and Bassett and Crompton (1975), roots are generally cold-resistant, able to survive in the extremely cold of Canadian winters, and can continue growing the following spring for over 30 years. It has been supposed that independence from germination requirements may allow *A. psilostachya* to colonize those countries where *A. artemisiifolia* cannot successfully conclude its life cycle (e.g. The Netherlands; Van Denderen *et al.*, 2010).

Concerning plant development, data are available mainly for *A. artemisiifolia*. Cunze *et al.* (2013) estimated that it requires an accumulated temperature sum of 1400°C to produce mature seeds. In their report, Bullock *et al.* (2012) found that the maximum photosynthetic rate is at 20°C (halved at 30°C). Nevertheless, they also highlighted that *A. artemisiifolia* persists where the climate is hotter, deducing that high temperatures are likely to have a lesser impact on its performance than low temperatures. Bazzaz (1974) attributed its tolerance to high

temperatures to high transpiration rates, which allow a transfer of latent heat in leaves at temperatures below ambient temperature. Frost in late spring or early autumn can be fatal for seedlings and adult plants (Essl *et al.*, 2015), although Leiblein Wild *et al.* (2014) observed that seedlings are more frost tolerant in Europe than in native countries, thus supporting the assumption of local adaptation. In general, Rasmussen *et al.* (2017) recently found that common and giant ragweed perform better in relatively wet conditions, while perennial ragweed in drier ones. Growing degree days are generally cited as the most influential climatic factor explaining the distribution of short - day plants *A. artemisiifolia*, *A. trifida* and *A. psilostachya* in Europe. The main climate requirements of the four ragweed species considered are summarized in Table 3.

#### *C. Moisture and soil types tolerance*

Soil is another factor determining the colonization and successful survival of plants. Concerning soil pH, few studies were undertaken and the results are not totally consistent. Fumanal *et al.* (2008a) demonstrated that *A. artemisiifolia* can grow both on acid and alkaline soils (extreme values of pH KCl: 4.1–8.6), even if preferentially occupies sites with a pH range between 7 and 8. Coherently, Essl *et al.* (2015) reported that *A. artemisiifolia* grows best under moderately basic condition (pH = 8). On the other hand, Pinke *et al.* (2011) found the highest common ragweed cover at the edge of Hungarian sunflower fields when the soil pH was acid (< 5); this in agreement with the information reported in Hungarian literature, namely that this weed thrives best on acidic sandy soils (Ujvárosi, 1973; Szigetvári and Benkő, 2008). Regarding germination, Sang and collaborators (2011) demonstrated that *A. artemisiifolia* germination success exceeded 48% in solutions with pH values between 4 and 12, with maximum rates occurring in distilled water at pH 5.57. However, under laboratory conditions, germination occurs in a wider range of pH (Bullock *et al.*, 2012). Silt loam and silt clay loam soils are elected as the optimum by Basset and Crompton (1975) and, in France, Fumanal *et al.* (2008a) found

the plant on sand to clay or silty loam, but mostly on sandy soils. Regarding soil water content, *A. artemisiifolia* can be very resilient to short-term drought (Bullock *et al.*, 2012). Nevertheless, Hodgins and Rieseberg (2011) demonstrated the poor survivorship under drought conditions of the European populations in comparison to the American ones, probably due to the evolution of a life-history that has favoured a more rapid growth and reproduction than drought tolerance in the invasive range. Leiblein and Lössch (2011) observed a major growth of *A. artemisiifolia* in moist soil conditions, but also the capacity to survive in dry, moist and waterlogged soils (5%, 22% and 39% of water). In the latter situation, the plants are far smaller, but able to reach maturity and produce seeds, although in small quantities. In keeping with Essl *et al.* (2015), *A. artemisiifolia* is not typical of wet areas, but its seeds can potentially tolerate and remain viable in soils with high water content. Concerning salinity (Table 3), Di Tommaso (2004) showed that the seeds of *A. artemisiifolia* can also germinate at high levels of sodium chloride (5 to 12% of germinated seed at 400 mmol L<sup>-1</sup>). However, he highlighted that the percentage of germination in his experiment was negatively correlated to the increase in salt, but the recovery in distilled water of viable non-germinated seeds was rapid. This study also suggested an adaptation of *A. artemisiifolia* to local conditions, as the seeds collected from plants living along roadsides showed a higher percentage of germination than those collected in cultivated fields. Another soil parameter that *A. artemisiifolia* appears to manage quite well is the presence of metals. Bae *et al.* (2016) proved that under metal stresses (Zn, Pb, Ni, Cd, Cu), *A. artemisiifolia* performs better in germination and seedling growth in comparison with native flora. This experiment simulated roadside conditions and delineated a potential empty niche where the species would have almost no competitors.

Information about soil requirements for the other ragweed species is not as exhaustive as for *A. artemisiifolia*, although it is possible to understand several differences by reviewing available data. *Ambrosia trifida* is typical of more mesophytic conditions than *A. artemisiifolia* (Abul-Fatih and Bazzaz, 1979a; Basset and Crompton, 1982), consistent with its native habitat (e.g.



floodplains temporarily including presence of standing water; Menges and Waller, 1983). Wortman *et al.* (2012) and Follak *et al.* (2013) showed how the distribution of *A. trifida* is more closely related to the seasonality of precipitation and summer precipitation than other variables such as land use and landscape structure. Low rainfall is a limiting factor in its native range (Basset and Crompton, 1982); germination occurs in a wide range of soil moisture with an optimum of 20-33% (Abul-Fatih and Bazzaz, 1979a). Nonetheless, Schutte *et al.* (2008a) found that giant ragweed seedling emergence is insensitive to dry conditions of the top layers of soil (1 cm of soil) and that emergence usually occurs during relatively dry periods. Soil texture is not specified, but based on habitat (floodplains, drainage ditches, open stream banks), it can be deduced that *A. trifida* colonizes incoherent soils; in cultivated fields, it is usually found in low silty substratum (Basset and Crompton, 1982). There are no data regarding preferences of soil pH and salinity tolerance, while Cui *et al.* (2007) showed that *A. trifida* can live where the concentration of metals (Pb, Zn, Cu, Cd) is quite high and can be considered a good accumulator at root level.

Concerning *A. psilostachya*, as previously stated, it is a sand-loving species and prefers well drained, alkaline soils (Basset and Crompton, 1975). It lives in soils characterized by high salinity both in the native and invasive ranges, although salt can strongly limit the growth of plants (Salzman, 1985). Salzman and Parker (1985) experimentally demonstrated that the wide root system can balance stress through the connection between ramets; taking advantage of local salinity variations, ramets living in lower salinity conditions contribute to the survival of those living in high salinity conditions. This physiological integration (exchange of resources among connected ramets) helps *A. psilostachya* persist in stressful conditions. It boosts its efficiency of colonization and habitat exploration, thus promoting a greater dispersal ability (through rhizomes) in adverse conditions, to increase the probability of finding favourable microsites (Salzman, 1985). Furthermore, *A. psilostachya* persists even when high concentrations of metals in soil are lethal for other plants (e.g. Zn, Cu, Mn, etc.) (Basset and

Crompton, 1975). The successful survival of *A. psilostachya* in metal-rich soils appears to be positively mediated by mycorrhizal symbiosis (Rivera-Becerril *et al.*, 2013). Habitats colonized by Western ragweed are typically subjected to seasonal drought. It is a xeric adapted species (Corbett and Anderson, 2006), with a very long root able to draw water from deep sources and persist without desiccating at more humid deep soil levels (1.83 m in depth according to Stromberg, 2013). In the grasslands of the Nebraska Sandhills, *A. psilostachya* is relatively stable through drought (and grazing), while other resistant perennial grasses are damaged (Reece *et al.*, 2004; Stubbendieck and Tunnell, 2008). Although severe events leading to a serious desiccation of rhizome (- 60% of weight), can have negative effects on shoot emergence (Karnkowsky, 2001). Regarding tolerance to waterlogged conditions, *A. psilostachya* prefers drained soils, but it lives in the riparian habitat and according to Stromberg (2013) it is a mesoriparian species. Finally, from data by Towne (2000), who monitored the impact of large ungulate carcasses (e.g. bison, cattle and deer) on grassland dynamics, it can be argued that *A. psilostachya* dominates in nutrient-rich soils and easily tolerates high levels of organic compounds.

Information about soil preferences of *A. tenuifolia* is quite dispersed. It prefers fertile, well aerated (Insausti and Soriano, 1987), not very deep hydro halomorphic soils (Burkart *et al.*, 1990; Anton *et al.*, 2012). Soil water content can be determinant in the life cycle of the plant. *Ambrosia tenuifolia* co-/dominates plant communities frequently exposed to floods of varying intensity and duration, usually brief (1-2 months), of reduced magnitude (water cover does not exceed the depth of 7 cm in spring), with only occasionally prolonged events with heavy impact occurring (water cover 10–30 cm deep for 3–5 months) (Insausti and Grimoldi, 2006). Insausti and Soriano (1987) observed that *A. tenuifolia* frequently grows on anthills, and argued that those sites are suitable as they are not affected by prolonged waterlogged soil conditions and the subsequent anoxia is not tolerated by roots for more than 1-2 months (Insausti and Grimoldi, 2006). On the other hand, seeds can tolerate immersion in water and

low temperature even for long periods, without negative effects on dormancy release (Insausti *et al.*, 1995). Resilience to drought has not been specifically investigated but a certain degree of tolerance can be inferred as environments where *A. tenuifolia* lives are involved in periods of drought, both on the Pampean Plains (summer drought) and coastal areas (Marcomini *et al.*, 2016). A summary of the aforementioned soil requirements is shown in Table 3.

#### *D. Light requirements and tolerance*

The tolerance to light/shade has been briefly discussed in the description of habitats and the main requirements for ragweed species are reported in Table 3. In general, ragweed species are pioneer plants living in open sunny environments. However, based on habitat preferences, tolerance of moderate shade has been shown for *A. psilostachya*, and Essl *et al.* (2015) affirmed that *A. artemisiifolia* is also medium shade-tolerant. Generally, shade suppresses *A. artemisiifolia* (Bullock *et al.*, 2012) and the lack of adequate light intensity strongly contributes to a progressive decrease in plant performance and recruitment (Gentili *et al.*, 2015; 2017). Conversely, *A. artemisiifolia*, both at mature and seedling stage, is extremely tolerant to high light intensities, which are characteristic of open environments (Bazzaz, 1974). *Ambrosia trifida* is not only tolerant to high light intensities but Menges and Waller (1983) indicated it as a high light specialist or light-loving species. The authors distinguished this species from the low light specialists and light generalist herbs in floodplain communities; the early emergence, development of seedlings and the great growth of plants would indicate that *A. trifida* is strongly projected to light exploitation. However, several studies show that the species is tolerant also to shady conditions likely to occur along both the natural vegetation dynamic and cultivated fields. It appears that *A. trifida* can allocate resources differently, based on light situations (Hartnett *et al.*, 1987; Abul-Fatih *et al.*, 1979; Webster *et al.*, 1994; Jurik, 1991).

Although a multi-factorial process, *A. artemisiifolia* germination is also light-induced. For instance, in this species the secondary dormancy (Baskin and Baskin, 1980; 1985) is induced

by the lack of light in combination with low temperature fluctuations, high CO<sub>2</sub> concentration in the soil and hot dry summer periods (Bazzaz, 1979; Essl *et al.*, 2015). Nevertheless, in lab experiments, seeds germinated also in the dark in a range of temperature corresponding to late spring and summer (Bullock *et al.*, 2012; Baskin and Baskin, 1980).

Along with alternating temperatures, light and in particular the R:FR ratio also promotes *A. tenuifolia* seed germination and plant growth, which benefits from vegetation gaps (Insausti *et al.*, 1995; Insausti and Grimoldi, 2006). Concerning germination, the same can probably be said for *A. psilostachya* (CABI, 2017). Conversely, light is not one of the main factors promoting the germination of *A. trifida* seeds (Davis, 1930; Schutte *et al.*, 2012), which are bigger in comparison to those of the other species and have different mechanisms of quiescence release mediated by pericarp and/or embryo-covering structures (Harrison *et al.*, 2007; Schutte *et al.*, 2012).

#### *E. Plant traits involved in resistance and resilience*

To support and explain environmental requirements, changes in several plant traits have been highlighted as key strategies in enhancing resistance or resilience to abiotic stresses (e.g. dormancy, rhizome features, early emergence of seedlings, etc.). However, stress can also derive from unfavourable biotic interactions with parasites and predators, as well as from competition with the local plant community. Furthermore, on observing colonized environments, there is a series of potential human-mediated disturbances such as fire, grazing, mowing, agricultural practices and human settlement development. All these factors can affect the reproductive and vegetative fitness of individuals or prevent their persistence through direct suppression or a dramatic change in environmental parameters. Ragweed plant traits involved in stress resistance and resilience are analyzed and discussed below.

## 1. Resistance

Starting from strategies and traits that enhance the resistance of individuals to natural or human perturbation (intending resistance as the capacity of an individual to resist the displacement of its biomass (Grime, 2001), it is found that morphological adaptations can avoid or limit the effect of several perturbations. Spines, pubescent or sclerophyllous leaves, or incorporation of granular minerals into plant tissues are typical defensive structures against grazing (Hanley *et al.*, 2007). In a xeric habitat, waxy leaves or a particular form of plants (e.g. cushion plants) can prevent desiccation and damage from wind, drought or frost, while other morphological structures can promote the persistence of individuals in flooded and waterlogged riparian areas (Catford and Jansson, 2014).

Ragweed plants (Table 4), do not show any evident morphological adaptation to overcome stressing factors that damage aerial parts. Only perennials (*A. psilostachya*, *A. tenuifolia*) have quite densely short-haired leaves that play a relatively protective role in avoiding desiccation; *A. psilostachya* may assume a prostrate habit that enhances resistance to several stresses. The presence of phytoliths (mineral deposits in epidermal cell walls) in aerial parts has been reported for *A. psilostachya* and *A. trifida* (Bozarth, 1992) and this increases resistance to leaf-eating invertebrates (Hanley *et al.*, 2007).

Rather than morphological adaptations, chemical protections, such as leaf-coating resins, are important in ragweeds: *Ambrosia* species characteristically possess glandular trichomes, especially on the lower leaf surfaces but also on stems, thus producing resinous excreta rich in secondary metabolites such as sesquiterpenes and flavonoids (Mitchell *et al.*, 1971; Wollenweber *et al.* 1987; 1995). In general, like many other Asteraceae (Heinrich *et al.*, 1998), *Ambrosia* species can biosynthesize many types of secondary metabolites (Hodgins *et al.*, 2013; Wan *et al.*, 2002; Wang *et al.*, 2005; Kong, 2010; Sülsen *et al.*, 2008, 2013) that contribute to protecting plants from abiotic and biotic perturbations (Table 4) Parts of plants and seeds can be included in the diet of several wild mammals, birds and insects, and due to

secondary metabolites ragweed species are unpalatable for cattle that only resort to eating the plants when there is no alternative forage (Marten and Andersen, 1975; Reece *et al.*, 2004; Bullock *et al.*, 2012). It is worth noting that many authors (Gerber *et al.* 2011; Essl *et al.*, 2015; Goeden and Ricker, 1976) have pointed out that *Ambrosia* species are attacked by specialized parasites that affect their life cycles in their native range, rather than in their invasive range where the parasites are less specialized and the damage inflicted is often not relevant.

The allelopathic effects of ragweeds on other plants are also well documented (Table 4). Root exudates, leaf leachate and decaying leaves produce allochemical compounds that inhibit germination and growth of other species, both in natural and agricultural environments. Most studies relate to *A. artemisiifolia* (Rice, 1965; Bullock *et al.*, 2012; Vidotto *et al.*, 2013) and *A. trifida* (Wang *et al.*, 2005; Kong *et al.*, 2007), which have been taken into account as a potential bioherbicide (Kong, 2010; Molinaro *et al.*, 2016). However, allelopathy is known also for *A. psilostachya* (Neill and Rice, 1971; Dalrymple and Rogers, 1983) and *A. tenuifolia* (Mongelli *et al.*, 1997).

Another important factor involved in resistance of ragweed species to stress and disturbance is the presence of mycorrhizal fungi (Table 4). Mycorrhiza improve plant growth and health by enhancing mineral nutrition and increasing tolerance to abiotic and biotic stresses (Lenoir *et al.*, 2016). *Ambrosia artemisiifolia* is considered in obligatory symbiosis with mycorrhizal fungi (arbuscular mycorrhizal fungi, AMF) and studies have demonstrated that fungal colonization is positively correlated to environment disturbance (Essl *et al.*, 2015). Similarly, *A. psilostachya* has also been found associated with mycorrhizal fungi in disturbed and polluted environments (Busby *et al.*, 2011; Pendleton and Smith, 1983; Rivera-Becerril *et al.*, 2013). For *A. trifida*, evidence of root colonization by mycorrhizal fungi are few (MacDougall and Glasgow, 1929; Bassett and Crompton, 1982) and no data are available for *A. tenuifolia*. Both the species live naturally in environments subjected to seasonal flooding and

wet areas are not suitable to mycorrhizal colonization, probably due to the lack of well-aerated soils (Entry *et al.*, 2002; Escudero and Mendoza, 2005).

The re-allocation of resources is considered a further important mechanism allowing ragweed plants to tolerate and respond to environmental stresses (Table 4). For instance, *A. trifida* allocates resources differently as a reaction to light variations (see above). Following defoliation due to herbivory attacks, *A. artemisiifolia* can efficiently re-allocate resources from root to shoot biomass and avoid evident costs for fitness (Gard *et al.*, 2013); it can also enhance ramification when the stem apex has been removed (Brandes and Nitzsche, 2006).

Interestingly, when environmental stress is lower or absent, as can occur in introduced ranges, alien plants can reallocate resources and thereby improve their growth and competitive ability. This is at the foundation of the Evolution of Increased Competitive Ability hypothesis (EICA) which has been associated with *A. artemisiifolia* in relation to changes in climate (Leiblein-Wild *et al.*, 2014), environmental conditions (Hodgins and Rieseberg, 2011) and parasites (Fukano and Yahara, 2012). Nevertheless, further studies have demonstrated that the hypothesis is not always valid for *A. artemisiifolia* (Genton *et al.*, 2005; MacKay and Kotanen, 2008).

## 2. Resilience

Traits and strategies related to resilience ensure rapid recovery from disturbance and stress and a return to control levels (Grime, 2001). In the case of highly disruptive natural and human-related perturbations, species need to rely on a series of regenerative strategies to have a speedy and complete return to the earlier status. Attributing traits and strategies of plants to resistance or resilience can often be tricky, and this paragraph focuses on the main strategies involved in recovery from severe events that definitely lead to suppression of individuals or their aerial part in the case of geophytes.

Resprouting capacity (a resistance/resilience trait) can be considered one of the main functional traits related to successfully overcoming fire, mowing, intensive grazing and some severe atmospheric events leading to suppression of the aerial parts of plants (Table 4; Keeley *et al.*, 2011). *Ambrosia psilostachya* and *A. tenuifolia* live on plains traditionally subjected to these types of severe perturbations and are not highly affected by them (Wolfe, 1973; Menghi *et al.*, 1993; Hartnett *et al.*, 1996; Madanes *et al.*, 2007); on the contrary, they are often favoured and thus show their weediness (Abrams, 1988; Hartnett *et al.*, 1996; Vermeire and Gillen, 2000; Vermeire *et al.*, 2005; Insausti and Grimoldi, 2006). This is related to their rhizome, a "resistance" structure that confers resilience on these species. As already underlined, the life strategy of *A. psilostachya* is mainly based on its belowground system which allows it to overcome adverse moments and unpredictability deriving from human action or climate: through the rhizome, *A. psilostachya* can form clones of plants occupying areas larger than 100 m<sup>2</sup> (Karnkowski, 2001). In suitable situations, the presence of the weed can be very massive: 1132 kg ha<sup>-1</sup> dry weight according to Bovey *et al.* (1966). It has been estimated that the establishment of a "competitive" root system takes one year; after the emergence of a seedling from one of the few mature seeds, a shoot emerges from the root during the second year and in only one season it can colonize an area of 2 m<sup>2</sup> (Basset and Crompton, 1975; Mitich, 1996). Reece *et al.* (2004) demonstrated that *A. psilostachya* can maintain primordia for several years even with limited plant growth, and Wan *et al.* (2002) showed that clipping stimulates the growth of new stems. Limits to rhizome viability/resprouting derive from several climatic conditions and burial depth (Miziniak and Praczyk, 2002). Several authors affirm that sprouting from buds is possible when soil thickness is up to 5 cm (Miziniak and Praczyk, 2002; Vermeire *et al.*, 2005). In comparison to *A. psilostachya*, the life strategy of *A. tenuifolia* is based not only on rhizome sprouting, but also on seeds as explained below. Although fewer investigations have been carried out on the *A. tenuifolia* rhizome, it is clear that resprouting from belowground buds permits the recolonization of vegetation gaps after disturbance (Insausti and Grimoldi, 2006),



thus ensuring the persistence of the species (Semmartin *et al.*, 2010). *Ambrosia tenuifolia* is a highly productive species, 1,330 g m<sup>-2</sup> of total biomass according to Semmartin *et al.*, (2010) and it can advance quite rapidly thanks to rhizomes, with an estimated rate of 1.72- 0.2 m<sup>2</sup>/month (Insausti and Grimoldi, 2006). As already mentioned, rhizome persistence in this species is limited by anaerobic condition. Resprouting is also important for *A. artemisiifolia* resilience: removal of stem, as can occur in the mowing or grazing regime, induces the resprouting of plants from buds at the base (Brandes and Nitzsche, 2006; Patracchini *et al.*, 2011; Milakovic and Karrer, 2016). By contrast, resprouting capacity has never been reported for *A. trifida* species.

In addition to their reproductive capacity, seeds also play a crucial role as "survival structures" deputized to respond to environmental unpredictability and adversity and so confer resilience to species. Through dispersal, seeds can allow the species to strategically escape from unsuitable conditions. Moreover, seed dormancy leads to a "delayed germination" that prevents the germination of fresh seeds when the environmental parameters are unsuitable and promotes the establishment of a soil seed bank (Finch-Savage and Leubner-Metzger, 2006; Gioria *et al.*, 2016), which buffer plant populations against environmental variability and increase the time of (local) extinction (Thompson, 2000).

Apart from *A. psilostachya*, ragweed species produce large amounts of seeds, which establishes conspicuous soil seed banks (Table 3). For instance, the density of seeds of *A. artemisiifolia* ranged from 4.5 to 536 units per m<sup>2</sup> in the upper 20 cm of soil depending on the habitat type (Fumanal *et al.*, 2008b). As seeds can remain viable in the soil for decades, even more than 40 years (Toole and Browne, 1946), they can be considered as forming long-lasting soil seed banks. However, it must be taken into account that burial depth is crucial for the viability of seeds, decreasing to 4 years on the soil surface (Essl *et al.*, 2015). Experiments and observations have been conducted at a depth between 0 and 25 cm, which is considered the living seed bank limit (Fumanal *et al.*, 2008b; Essl *et al.*, 2015; Karrer *et al.*, 2016). Fumanal

*et al.* (2008b) recorded a lower viability of seeds between 0-5 cm than at 5-15 cm. Karrer *et al.* (2016) confirmed that the deep soil condition (down to 25 cm) is more suitable for lengthening seed viability, but differences between seeds buried at 5 and 25 cm are not so pronounced as in the study by Fumanal *et al.* (2008b). The authors also speculated that viability is more influenced by seed origin and habitat than burial depth. In any case, beyond viability, germination of seeds is strongly influenced by burial depth: if seed germination is quite high on the soil surface, it decreases with increasing depth (below 8 cm), where parameters dramatically change and dormancy cannot be interrupted (Essl *et al.*, 2015). Guillemin and Chauvel (2011) observed a decrease in germination for seeds buried from 2 to 8 cm and null germination between 10 and 12 cm of depth. It is likely that with the decrease of soil depth most seeds tend to germinate and leave a greater amount of non-viable seeds in the upper soil. Moreover, seed mass also appears to influence the percentage of germination, and the lightest seeds are more sensitive to burial (Guillemin and Chauvel 2011). On the other hand, the time of germination is also important as demonstrated by Ortman *et al.* (2016) who showed that *A. artemisiifolia* seed traits have a minimal effect, while foliage cover and aboveground biomass are more relevant.

Burial depth also has a determinant effect on seed germination and seedling emergency for *A. trifida*; germination decreases with depth after the first winter burial period (vernalization) (Harrison *et al.*, 2007). According to Harrison *et al.* (2007), the lowest depth from which giant ragweed can emerge is probably between 16 and 20 cm and no seedling emerges beyond 20 cm of depth. Soil seed bank viability in *A. trifida* is lower than that observed for *A. artemisiifolia*, as the total percentage of germination strongly decreases after 4 years (Harrison *et al.*, 2007), and the combination of low viability and high post-dispersal predation of seeds leads to a limited effectiveness of soil seed banks in this species (Harrison *et al.*, 2001, 2003). In any case, in suitable conditions, *A. trifida* enriches its soil seed bank by repeated dispersal of seeds year after year. Moreover, this species produces polymorphic seeds and, in

contrast to *A. artemisiifolia*, their size is relevant for their persistence in soil. Even with some exceptions, small seeds appear to be viable for a longer than larger ones (Schutte; 2008b). Nevertheless, the likelihood of the emergence of seedlings from large seeds is higher at 5 and 10 cm of soil depth (Harrison *et al.*, 2007).

A viable soil seed bank also allows *A. tenuifolia* to be more resilient to critical phases such as floods. As reported above, the root system of this perennial species does not survive in anaerobic conditions due to prolonged water coverage, while seeds remain viable. According to Insausti and Grimoldi (2006), seeds are released in large quantities and they can remain viable for several years in soil seed banks. Unfortunately, information such as germination percentage and critical depth are not available for this species.

Seed production of *A. psilostachya* is very low and, in some studies, the species was even classified as a non-seeded forb (Table 4; Grygiel *et al.*, 2012). Moreover, studies performed in tallgrass prairies and coastal areas showed no seed of this species in the soil seed bank (McNicoll and Augspurger, 2010; Barton *et al.*, 2016). Nevertheless, from samples of soils collected in North American mixed prairies and then stored in artificial conditions, Lippert and Hopkins (1950) observed emergence of seedlings of *A. psilostachya* but in very low numbers. Information about seed viability is very scarce but is likely to be around 3-5 years, independent of persistence in the soil (Barton *et al.*, 2016). Thus, resilience strategies in *A. psilostachya* do not include soil seed banks, but the rhizome is likely to be the target organ in overcoming stressful conditions and disturbances.

#### *F. Competition with local plant communities*

From previous sections, it is clear that ragweed species colonize habitats quite far from *equilibrium*, characterized by the influence of several stresses and disturbances. They also occur in "stable" situations, but usually when the *equilibrium* is determined and "arrested" by particular conditions, such as a high concentration of salt or metal in soil. Essl *et al.* (2015)

underlined that germination and early seedling establishment are mostly related to disturbance and lack of competition from local communities. This would explain the rarity of plants in natural habitats. As pioneer species, their life strategies are shaped to harsh situations mostly in early successional stages of vegetation. Accordingly, almost all biological and functional traits of ragweeds can be related to these low evolved environments. For instance, the need of light intensity to break seed dormancy and the need for rather shallow and compact soils, suitable to the growth of seedlings from soil seed banks and resprouting from rhizomes, are all requirements related to poorly evolved or disturbed vegetation. The main characteristics of the four-ragweed species, related to their ability in competing with local plants, are compared in Table 4.

Considering *A. artemisiifolia*, environmental conditions are only suitable for its growth when human action or natural events influence the natural evolution of vegetation by removing competitors or operating on soil components (agricultural practices, excavations, floods, etc.). Gentili *et al.* (2015) demonstrated that the plant is a weak competitor in evolved stages of vegetation. These authors showed that germination and recruitment, as well as plant growth, are mainly inhibited by the presence of perennial and/or winter annual grassland species characteristic of more advanced stages (Gentili *et al.* 2015). In keeping with this, Fenesi *et al.* (2014) reported that in a competitive regime, *A. artemisiifolia* seed germination is delayed and seedling development is restrained by the presence of heterospecific neighbours (e.g. *Erigeron* spp.). They reported that on average the cost in *A. artemisiifolia* biomass associated with only a 3-day emergence delay is very high (-97%).

In contrast to *A. artemisiifolia*, *A. trifida* is usually a good and vigorous competitor. Its life strategies are mostly based on very rapid and relevant growth: early germination, followed by very rapid growth, allows the plant to reach a height and biomass superior to other plants. Accordingly, the growth of later-growing plants is strongly inhibited by the canopy shadow determined by its large leaves (Abul-Fatih and Bazzaz (1979b). Therefore, once giant ragweed

finds good conditions in which to persist, it inhibits species diversity, biomass and density of the local community. In its native community, *A. trifida* is in fact a dominant species, one that strongly inhibits the colonization and growth of other annual plants (Abul-Fatih and Bazzaz, 1979b). Early emergence also ensures a timely capitalization of resources that avoid the competition of dominant perennials as well (Hartnett *et al.*, 1987; Schutte *et al.*, 2012). As *A. trifida* is one of the most problematic crop weeds in the United States, its impact on other plants is extremely evident as shown also by the analysis of crop yield losses (Regnier *et al.*, 2016). Interestingly, in agricultural environments, but not in rarely disturbed natural successional areas, *A. trifida* has been reported as being able to modulate its emergence time to adapt to different, "scheduled" selective pressures (Hartnett *et al.*, 1987; Schutte *et al.*, 2012; Regnier *et al.* 2016). In general, environmental or human-mediated disturbance must surely contribute to *A. trifida* persistence but, contrary to *A. artemisiifolia*, it does not disappear when the natural vegetation dynamic evolves and perennials become dominant. Hartnett *et al.* (1987) reported that *A. trifida* can persist for years, even penetrating dense vegetation.

Similarly, perennial ragweed species, being pioneer plants, take advantage from vegetation gaps and also persist in more evolved environments when suitable conditions are maintained. Nevertheless, unlike *A. trifida*, *A. psilostachya* is not a superior competitor of grasses under "normal circumstances", and is in fact present in undisturbed, healthy pastures, but in low quantities (Vermeire *et al.*, 2005). Vermeire and Gillen (2000) demonstrated that its abundance does not affect the presence of other grasses in mixed prairies, and there is a positive correlation between them. However, they speculate that Western ragweed is less abundant where there are grasses with roots forming a dense mat in the upper soil level; in this way the vegetative propagation of the plant is inhibited, as upper soil roots compete with the quite superficial creeping rhizome of *A. psilostachya*. Thus, the species only endures competition with some plants. This is inferred by observing its allelopathic effect, which only negatively affects some species.

Again, *A. tenuifolia* takes advantage of gaps in vegetation covers (caused by floods, grazing, etc.) which make light, nutrients and water more available for seed germination and recruitment (Insausti *et al.*, 1995; Insausti and Grimoldi, 2006). As a result, this species is a good competitor in poorly evolved environments, although it usually persists in the following stage of vegetation succession where it can become co-dominant (see above). After disturbance, gaps constitute focal points where recolonization of the grassland by *A. tenuifolia* originates from seedlings, which initially have a slow growth phase. It then grows rapidly and continues outside the original gaps by lateral clonal expansion, thus allowing the species to occupy new areas. In any case, independent of the starting population's abundance, the species multiplies many times the surface of colonization in the quite short time of about 4 months (Insausti and Grimoldi, 2006).

## **VI. Reproductive and dispersal barriers**

Survival without reproduction reduces an exotic species to a casual alien *taxon* as it cannot reach the naturalization phase (Blackburn *et al.*, 2011). In this section, reproductive and dispersal strategies (also human-mediated) of ragweed species are analysed. Dependence on specialized pollinators or particular requirements within reproduction phases (e.g. obligatory outcrossing) are often indicated as traits that prevent the establishment of plants in new ranges (Van Kleunen *et al.*, 2015). In general, when a species does not encounter unfavourable factors that strongly limit reproduction, the road to establishment is much less difficult. As already seen, the life cycle of ragweed species includes a series of adaptations useful in avoiding adversity. However, the chance of easily shifting its range through reproductive structure dispersal can be a positive trait determinant in facing unpredictability and escaping unfavourable conditions (Estrada *et al.*, 2016).

### *A. Pollination*

In ragweed species, flowers are organized in heads containing either male or female flowers. The pollen-producing male raceme grows at the tips of the principal stem and lateral branches; seed producing female heads containing one or a few pistillate flowers are sessile and situated in the axils of the leaves immediately below the staminate spikes (Smith *et al.*, 2013). According to Smith *et al.* (2013), within the Asteraceae family, ragweeds possess a strongly modified inflorescence, highly adapted to wind pollination (Table 5). As highlighted by Franz Essl and coauthors (2015), *A. artemisiifolia* is strongly self-incompatible and has high outcrossing rates, both in its invasive and native ranges. This may limit its reproductive efficiency, but due to the large production of airborne pollen, genetic flux is also maintained between distant or isolated populations. Unfortunately, there is no information about the other ragweed species concerning this.

### *B. Seed and propagule pressure*

In the literature, the term "seed" usually indicates the whole diaspore unit of ragweed, which is a one-seeded cypsela for the species considered. The characteristics of ragweed seeds are shown in Table 5. It can be observed that *A. trifida* produces the largest seeds (more than 6 mm long) with a consistent outer coat (Bassett and Crompton, 1982). The other ragweed species produce smaller seeds: seeds of *A. tenuifolia* are slightly larger, 3-5 mm long (Parsons and Cuthbertson, 2001; Behçet, 2004), than those of *A. artemisiifolia* (~3.5 mm long) (Bassett and Crompton, 1975) and *A. psilostachya* (3-4.5 mm long) (Table 5) (Barton *et al.*, 2016).

Propagule pressure of common ragweed can be very high because the species produces between 3,000 and 100,000 seeds per plant, depending on the size of individuals and thus on growth conditions (Dickerson and Sweet, 1971; Bassett and Crompton, 1975; <https://gd.eppo.int/reporting/article-3032>; Fumanal *et al.*, 2007). Seed production is also quite

noteworthy for giant ragweed and varies between a few hundred and 5000 ca. units per plant, depending on plant density and environmental conditions (Abul-Fatih and Bazzaz, 1979a; Baysinger and Sims, 1991; Harrison *et al.*, 2001; MacDonald and Kotanen, 2010). Nevertheless, unlike *A. artemisiifolia*, the potential dissemination of this species in its native range can be strongly reduced by a relevant post-dispersal predation by rodents and invertebrates (Harrison *et al.*, 2003; Regnier *et al.*, 2008). Moreover, the viability of its seeds is not very high, 50 - 66% (Goplen *et al.*, 2016; Harrison *et al.*, 2001).

In contrast to common and giant ragweed which, being annual species, have seeds as their main dispersal units, Western and slender ragweed are perennial and show additional dispersal structures. Wagner and Beals (1958) observed that only 66 out of 118 flowering heads developed to maturity in one plant and speculated that the reproductive potential by seeds of *A. psilostachya* is six times less than that of *A. artemisiifolia*. Similarly, Basset and Crompton (1975) showed that *A. psilostachya* produces just one seed per flowering head, thus indicating that vegetative reproduction is predominant for the species. In agreement with this, the main dispersal structure of the species is its highly vigorous creeping root system, capable of sprouting from pieces of rhizome, which make this ragweed species an even harder weed to fight than common ragweed (Table 5). Vegetative reproduction through rhizomes is also relevant for *A. tenuifolia*, even though it produces a large number of seeds (Table 5; Insausti and Grimoldi, 2006). Indeed, in this species both these strategies are important for its success in different stages of its life, as they react to dramatic environmental events, such as floods that are frequent on the Pampean Plains or human pressures such as agriculture and grazing (Insausti and Soriano, 1987; Insausti *et al.*, 1995; Insausti and Grimoldi, 2006; Soriano, 1982). As already discussed in previous sections, the rhizome of *A. tenuifolia* is less resistant to extreme conditions (i.e. anoxia due to floods) than seeds that showed a longer viability (Insausti and Grimoldi, 2006). Furthermore, there is no evidence of enemies in the native range that affect the persistence and productivity of this plant, given its toxicity or unpalatability to cattle and



the low dietary interest for other vertebrates such as rodents (Freire *et al.*, 2005; Semmartin, 2010; Ellis *et al.*, 1998).

### *C. Dispersal*

Regarding ways of dispersal relevant to local movements (medium-short-range dispersal in native and invasion range), ragweed seeds show no morphological structure strictly representing a specific dispersal vector. According to Basset and Crompton (1975), the primary way of dispersal of *Ambrosia artemisiifolia* seeds is barochory. Anemochory is often cited as a potential dispersal vector, but owing to the absence of suitable structures and the weight of seeds, wind may represent only a "facilitator" of spread rather than a driver of diffusion (Table 5; Bullock *et al.*, 2012). Zoochory and hydrochory (Table 5) were considered by Essl *et al.* (2015), who reviewed all vectors contributing to *A. artemisiifolia* diffusion. Regarding the first, epizoochory by bison was proven in the native range of common ragweed, and endozoochory was also reported as an additional plausible mechanism of dispersion (Rosas *et al.*, 2008; Bullock *et al.*, 2012). Viable seeds of common ragweed resulting from feed intake have been found in cattle manure both in the United States and Europe (Pleasant and Schlather, 1994; Vitalos and Karrer, 2008). In addition, Wright, (1941) and Vitalos and Karrer (2008) proved that seeds are part of the diet of some birds (e.g. sparrows, pheasants, and quails), and Essl *et al.* (2015) reported that these vectors, along with rodents, play a role in dispersal. Hydrochory (intended both as streams, flowing water, and as runoff), has been indicated as a way of dispersal in the native range of the species (Table 5; Payne, 1970). Recently, Fumanal *et al.* (2007) underlined how the polymorphism of seeds of *A. artemisiifolia* contributes to dispersal by flowing water and indicated this mechanism as fundamental in the spread of the *taxon* within France, and also as a long-distance means of diffusion.

Compared to *A. artemisiifolia* and the other ragweed species considered, *A. trifida* produces the largest seeds, reaching almost 1 cm in length (Bassett and Crompton, 1982). Seed size and weight point to barochory as the main way of dispersal also for this species (Table 5).

However, Osawa *et al.* (2013) indicated hydrochory and zoochory as additional important mechanisms for *A. trifida* diffusion, mainly supporting the former in accordance with the shape and size of the seeds. Yoshikawa *et al.* (2013) demonstrated that owing to its weight, a giant ragweed seed transported by a stream settles quite rapidly when the current velocity decreases, thus suggesting that these seeds are transported by water flow rather than floating. Concerning zoochory, although giant ragweed seeds have a quite developed crown of large spines, their size and weight represent a limiting factor for epizoochory. Hejný and Jelík (1972) mentioned the presence of *A. trifida* seeds in wool scraps, but as an exceptional circumstance in the former Czechoslovakia. In addition, Verloove (2016a) reported that it is rarely seen as a wool alien in Belgium and Pyšek (2005) excludes the possibility that *A. trifida* dispersal could be associated to wool processing in Czech Republic. As already mentioned, owing to their palatability, rodents and invertebrates eat seeds (Harrison *et al.*, 2003; 2007), but if they cached instead of being immediately eaten then post-dispersal predation allows an estrangement of viable seeds from the mother plant. An association between *A. trifida* and earthworms has only very recently been observed: in the native range of *A. trifida*, non-native earthworms (*Lumbricus terrestris*) cache its seeds in burrows. Beyond the benefits for the plants (reduction of fast seed predation), giant ragweed appears to increase dispersal opportunities through this acquired form of diplochory (Regnier *et al.*, 2008, 2016; Schutte *et al.*, 2010). However, it allows a very short-range translocation, probably less than one meter from the mother plant, considering the homing capability and movements of earthworms (Nuutinen and Butt, 2005).

Little information is available regarding *A. psilostachya*. However, considering that its seeds are slightly larger than those of *A. artemisiifolia* (Basset and Crompton, 1975), it is likely that this species also mainly disperses through barochory (Table 5). Moreover, the crown of rudimental spines is less developed or even absent in Western ragweed seeds compared to those of common ragweed, which suggests that epizoochory is probably not as important a dispersal mechanism in this species (Wagner and Beals, 1958). However, Amor Morales *et al.* (2012)

and Parsons and Cuthbertson (2001) cited epizoochory for *A. psilostachya* seed dispersal in Spain and Australia, respectively. Finally, although the species is not highly palatable to cattle or bison, endozoochory was also inferred by Rosas *et al.* (2008), who found a percentage of *Ambrosia* spp. seeds in dung of bison grazing in prairies where *A. psilostachya* was a common forb. In any case, although studies about the role of Western ragweed seed intake in bird diets have been published (Campbell-Kissock *et al.*, 1985), clear evidence of seed dispersal by animals is yet to be collected, as well as other means of diffusion. For instance, CABI (2017) reported that in springtime *A. psilostachya* seed can be "transferred by water in ditches, canals and rivers" (hydrochory). Moreover, as discussed above, seeds are probably not the main dispersal unit of Western ragweed, owing to their paucity, but the role of rhizome fragmentation is yet to be investigated.

Like *A. psilostachya*, *A. tenuifolia* is a perennial, but its life strategy is not mainly based on vegetative propagation and seed production is not low, although data on the precise quantity are not available. No real evidence for epizoochory is present in literature, but several sources stress the evidence that seeds are caught on sheep wool ([http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon\\_id=17510#](http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=17510#)). Available studies suggest that seeds can be dispersed by water flow without losing their viability (Insausti and Soriano, 1982; Insausti *et al.*, 2006).

#### *D. Human-mediated dispersal pathways (medium- and long-distance vectors)*

Trade routes and all connected elements, previously discussed as important pathways of introduction and global diffusion, are relevant for short- and medium-range movements (Table 5; Ferus *et al.*, 2015). Nevertheless, on a regional scale, further important vectors of spread linked to human activity stand out: movement of soils, spread of seeds through mowing or agricultural machinery and car and train passage (Table 5). All these vectors are associated with the spread of *A. artemisiifolia* particularly within the European area as extensively reviewed by

Bullock *et al.* (2012) and Essl *et al.* (2015). The presence of common ragweed in a new area, after the setup of a construction site, can easily be attributed to the transport and dumping of contaminated soils from different sites. Moreover, the abundance of common ragweed along railways and road networks indicates transportation corridors as one of the main drivers of introduction. The importance of which is even more stressed by the explanatory power of this variable in spatial distribution models and other studies (e.g. Dullinger *et al.*, 2009 and Joly *et al.*, 2011). However, few experimental data about the dispersion mechanism of propagules are available. Two studies (Vitalos and Karrer, 2009; Von der Lippe *et al.*, 2013) tested the effect of vehicles on dispersion of seeds and both demonstrated that the sole car slipstream or seed attachment cannot completely explain a long-distance dispersal. Even the action of mowing machines along roads, which strongly boosts the process, does not fully explain the distance of dispersal (Vitalos and Karrer, 2009). As a result, the spread of *A. artemisiifolia* along corridors of transportation appears to be a multifactorial phenomenon not yet completely understood. Regarding the impact of agricultural machinery, it is effective at the local and more extended level. Karrer (2014) demonstrated that harvesters and other machines can transport several thousands of viable seeds, colonizing new fields or reinforcing already present metapopulations of common ragweed. Contaminated machines from already colonized French regions are even thought to be responsible for the introduction of *A. artemisiifolia* into some virgin Swiss areas (Buttenschøn *et al.*, 2010).

All these vectors may also be relevant for the other species of ragweeds, and especially for *A. tenuifolia* and *A. psilostachya* as light-seed producers (Table 5). Conversely, *A. trifida* produces large seeds and may exploit these vectors less (Table 5). The transport of viable rhizomes through soil movements may also be an effective vector for the diffusion of *A. psilostachya*, but unfortunately, no evidence for this is available.

## VII. Allergenic impact and environment

The *Ambrosia* genus represents a global risk to public health owing to the allergic reactions induced by pollen allergens in atopic subjects. In Europe, ragweed pollen affects more than 36 million people each year and the prevalence of sensitization is growing mainly due to the plant's spread (Mihajlovic, 2015; Bordas-Le Floch *et al.*, 2015). Among all ragweed species, pollen of *A. artemisiifolia*, *A. psilostachya* and *A. trifida* have long been acknowledged as a significant cause of allergic disease (Ziska *et al.*, 2011). Similarly, *A. tenuifolia* is reported to be severely allergenic in its native range and produces a huge amount of pollen (Giscafre and Ragonese, 1942; Vaz Ferreira, 1946; Tejera and Beri, 2005; Del Vitto *et al.*, 2015; pollenlibrary.com). However, more specific and up-to-date medical evidence needs to be collected to better define the allergenic impact of this species (Tejera and Beri, 2005; Marco and Pirovani, 2009).

### *A. Pollen characteristics related to allergy*

The major source of allergenic proteins in ragweed plants is pollen. Ragweed pollen grains are small particles containing air chambers between the layers of the outer wall. These characteristics allow them to become easily airborne under favourable conditions and transported by wind for very long distances at a continental scale, even reaching areas not colonized by *Ambrosia* spp. plants (Smith *et al.*, 2013; Mahmoudi, 2016; Šikoparija *et al.*, 2013). It is striking that these pollen grains maintain their allergenic power over such a long distances, even after spending days in the atmosphere (Makra *et al.*, 2016; Grewling *et al.*, 2016). This means that exposed individuals may become sensitized to ragweed pollen allergens and develop symptoms even in areas where the plant is not widely distributed (Grewling *et al.*, 2016). Furthermore, sub-pollen particles (SPPs) of respirable size (0.5 to 4.5  $\mu\text{m}$ ) contain allergens that can be released by *A. artemisiifolia* pollen after hydration (i.e. after thunderstorms). These particles, along with pollen fragments, can also be transported for long distances, thus contributing to allergen exposure even when no airborne pollen grains are identifiable. This

generates out-of season pollinosis in highly ragweed-sensitive subjects (Table 6) (Busse *et al.*, 1972; Bacsi *et al.*, 2006; Pazmandi *et al.*, 2012).

The species' morphological characteristics are similar (Table 6) and identifying pollen grains to a single species by optical microscopy is not usually feasible. A few authors have investigated the pollen structure of ragweed species other than *A. artemisiifolia*. They show that *A. psilostachya* pollen grains are likely to be larger than the other species (Table 6) (Jacobson and Morris, 1976; Robbins *et al.*, 1979; Wan *et al.*, 2002) and that *A. artemisiifolia* pollen is distinguishable from that of *A. trifida* through the analysis of exine characters. Specifically, Bassett and Crompton (1982) reported that *A. trifida* has 60-65 spines on one-half of the grain surface, whereas *A. artemisiifolia* has 70-75 spines. No distinctive traits are reported for *A. tenuifolia* pollen.

#### *B. Pollen Allergens*

The allergens of the four ragweed species officially recorded by the International Union of Immunological Societies (IUIS) are shown in Table 7. The pectate lyase Amb a 1 is currently considered the most important allergenic group for the *Ambrosia* genus (Gadermaier *et al.*, 2014), although it has been identified only in *A. artemisiifolia* and not in the other species. Concerning *A. artemisiifolia*, ten different allergen groups are reported in the IUIS database and they were extensively reviewed by Gadermaier *et al.* (2014) and Bordas-Le Floch *et al.* (2015). Briefly, the list includes: Amb a 1 (comprising the formerly called Amb a 2, recently designated as Amb a 1 isoallergen 5), to which more than 95% of ragweed pollen allergic patients are sensitized; Amb a 3, classified as a minor allergen (sensitization prevalence of 30–50%); Amb a 4 (defensin), to which sensitization frequencies are variable; Amb a 5 (unknown function) a minor allergen, affecting only 10% to 15% of ragweed pollen-allergic individuals; the panallergen Amb a 6 (non-specific lipid transfer protein) which is considered a minor allergen (sensitization prevalence of 21% among ragweed sensitized patients); Amb a 7 (plastocyanins), another minor allergen (reaction in 15% to 20% of ragweed pollen-allergic

patients); Amb a 8 (profilin), highly cross-reactive with mugwort (*Artemisia*) profilin (Art v 4); Amb a 9/Amb a 10 (polcalcins), minor panallergens (reaction in 10% to 15% of ragweed pollen-allergic patients). Finally, the list includes the very recently discovered allergen Amb a 11 (cysteine protease), which has been classified as one of the major allergens along with Amb a 1 for this ragweed species (Bouley *et al.*, 2015). Additional IgE reactive pollen proteins, identified by transcriptomic and proteomic approaches, have been indicated as “*bona fide* allergens”, probably extending the list of *A. artemisiifolia* allergens (Bordas-Le Floch *et al.*, 2015).

In contrast to *A. artemisiifolia*, *A. trifida* and *A. psilostachya* allergens are less characterized. Only two (Amb t 5, Amb t 8) and one allergenic proteins (Amb p 5) have been identified respectively in *A. trifida* and *A. psilostachya*, and reported in IUIS and/or allergome databases. Amb t 5 and Amb p 5 belong to the 5 group of allergens of the *Ambrosia* genus and cross-react with Amb a 5. In *A. psilostachya*, two isoforms of Amb p 5, Amb p 5.0101 and Amb p 5.0201 have been characterized. Ghosh *et al.* (1994) investigated the variants of Amb p 5 from *A. psilostachya* pollen and suggested that these forms are part of the natural variation within the *A. psilostachya* species, which exhibits polyploidy and can form hybrids with related ragweed species. Amb t 8 is a profilin, an actin binding proteins (Girodet, 2013). However, these data are not present in the IUIS database and only scanty information about it is available.

*A. artemisiifolia*, *A. trifida*, and *A. psilostachya* pollen allergens have long been considered largely cross-reactive (Weber *et al.* 2007), and it is generally believed that one species is sufficient for skin testing and immunotherapy. However, in the Northern area of Milan (widely invaded only by *A. artemisiifolia*), about 50% of patients submitted to injection of specific immunotherapy with *A. trifida* showed little or no clinical response, although an excellent outcome was obtained if they were shifted to *A. artemisiifolia* specific immunotherapy (Asero *et al.*, 2005). By comparing the proteome of *A. artemisiifolia* with those of *A. trifida* and *A. psilostachya*, Barton and Schomacker (2017) recently found that only *A. psilostachya* pollen

contains all five Amb a 1 isoallergens identified in *A. artemisiifolia* and reported in the IUIS database. In contrast, they found only three Amb a 1 isoallergens (Amb a 1.2, Amb a 1.4, and Amb a 1.5) in *A. trifida*, the lesser IgE reactive isoforms. Although more specific analyses are needed to characterize the allergenic profile of these species, this information suggests that *A. artemisiifolia* is more similar to *A. psilostachya* than to *A. trifida*, thus explaining the results reported by Asero and collaborators (2005). The allergenicity of *A. tenuifolia* is still poorly known (no allergens are reported in allergen databases), but it shows little cross-reactivity with the other *Ambrosia* species (Girodet, 2013).

### *C. Allergenic impact and environment*

Regarding quantities of pollen, ragweed has the potential to release billions of pollen grains: for *A. artemisiifolia* it is well known that  $1.19 \pm 0.14$  billion pollen grains can be released per plant (Fumanal *et al.*, 2007; Smith *et al.*, 2013). However, pollen production is closely related to size, growth, phenology and fitness of plants. For *A. artemisiifolia*, there is a positive correlation between dry plant biomass and reproductive success, as bigger individuals produce more pollen grains (Fumanal *et al.*, 2007), although decreasing plant size is generally associated with increasing maleness and decreasing femaleness (Paquin and Aarssen, 2004). Biomass and flowering phenology can follow a latitudinal gradient (Allard, 1945; Dickerson, 1968; Leiblein Wild, 2014): both in Europe and North America, plants from southern populations grow larger and flower later than northern populations (e.g., Gudzinaskas, 1993; Li *et al.*, 2015). The time of flowering greatly depends on germination time and the average springtime temperature (April, May and June) (Kazinczi *et al.*, 2008); for instance, it has been shown that earlier germination during spring leads to higher biomass allocation and higher pollen and seed production. Consequently, environmental conditions can alter plant fitness and result in pollen production change (Smith *et al.*, 2013). It is worth noting that adaptations to newly invaded environments (e.g. Europe) often have a positive effect on the fitness of plants, reproduction,



and biomass allocation as well as influencing the length of flowering time (Hodgins and Rieseberg, 2011; Leiblein Wild, 2014).

Furthermore, pollen production is influenced directly or indirectly by human practices. Sensitization of the population to *A. artemisiifolia* is constantly increasing and is probably correlated with the increased civilization, urbanization and pollution of the last decades (D'Amato, 2007; Ghiani, 2012). Several studies warn that global changes are going to worsen the situation in the next few decades. Effects will include changes in ragweed distribution, plant growth and life cycle as well as pollen allergenicity itself (e.g. Ziska and Caulfield, 2000; Rogers *et al.*, 2006). Species Distribution Models (SDMs) for *A. artemisiifolia* predict that its potential distribution will increase globally (Essl *et al.*, 2015; Chapman *et al.*, 2016). In Europe, warmer summers and later autumn frosts will allow a spread of *A. artemisiifolia* northward and uphill, leading by the mid-21st century to the inclusion of northern areas (e.g. southern Scandinavia and the British Isles) in its climatically suitable regions, and southward a regression of the species' range. Regarding the *Ambrosia* genus in North America, Ziska *et al.* (2011) showed an increase in recent decades (since 1995) of the duration of the ragweed (*Ambrosia* spp.) pollen season as a function of latitude (latitudinal effects are primarily associated with a delay in first frost of the fall season and lengthening of the frost free period). *Ambrosia trifida* in China is predicted to slightly increase its range (<1%) although it had the potential to spread northward (Qin *et al.*, 2014). Recently Rasmussen *et al.* (2017) found that, by the year 2100, the distribution range of *A. artemisiifolia*, *A. trifida* and *A. psilostachya* will increase towards Northern and Eastern Europe under all climate scenarios and consequently the high allergy risk areas will expand in Europe. Effects of the increase in temperature influence the flowering season length, but also the growth of plants and pollen production. Wan *et al.* (2002) tested the effects of warming and mowing on *A. psilostachya*, and showed that both can increase above ground biomass of plants (AGB), and the ratio of ragweed AGB to total AGB. With warming, total pollen production increased by 84% because ragweed stems were more abundant.

Moreover, experimental warming significantly increased pollen diameter (13% increase). El Kelish *et al.* (2014) demonstrated that both an elevated level of CO<sub>2</sub> and drought stress have an effect on *A. artemisiifolia* pollen allergenicity because expressed sequence tags (ESTs) encoding allergenic ragweed proteins increased under those conditions. Zhao *et al.* (2016) showed the direct influence of elevated NO<sub>2</sub> on the increased allergenicity of ragweed pollen and Ghiani *et al.* (2012) demonstrated that traffic-related pollution enhanced ragweed pollen allergenicity, showing that pollen collected along high-traffic roads had a higher whole allergenicity than pollen from low-traffic roads and vegetated areas. Conversely, several studies have shown no effect on the content of the major ragweed allergen Amb a 1 due to high concentrations of ozone or extended exposure of the plant to this pollutant (Sénéchal *et al.*, 2015; Kanter *et al.*, 2013).

## **VIII. Conclusion**

The successful invasion of the ragweed species considered can be ascribed to a synergy of anthropogenic and bio-ecological factors. The globalization of commerce and changes in land use have dramatically favoured their spread into new areas. Firstly, the species were used as medicinal plants in the Americas and were transported to Europe and cultivated in botanical gardens. They then spread as a contaminant of crop and forage seeds, and in a wide variety of goods, by means of transportation, to become noxious pests. Climatic changes are predicted to worsen the impact of these species by increasing both their colonisation range and allergenic potential. Thus, the setting up of effective measures to prevent and stop their spread is essential. Until now, researchers have mainly focused on common ragweed, the most widespread species, and the results have often been automatically associated with the other three species, although their ecology, biology and allergenic and ecological impact can differ significantly. Although other ragweeds are less widespread globally than *A. artemisiifolia*, their impact could differ in terms of type and magnitude. For instance, *A. psilostachya*, *A. tenuifolia*, and *A. trifida* are able

to colonize environments different from *A. artemisiifolia*, thus potentially expanding their range of impacts as one of the most "black-listed" genera in the world. For this reason, further research efforts and data collection about factors that have allowed ragweed species to overcome geographical and environmental barriers are needed. Specifically, more in-depth research is necessary about:

- The impact, biology and ecology of ragweeds other than *A. artemisiifolia*, which may represent a severe threat to local plant communities, given their ability to colonize semi-natural habitats;
- dispersal vectors and introduction pathways and their role in the spreading of *taxa*, with a particular focus for the rhizomatous "low-seed producer" *A. psilostachya*;
- allergenic impact of ragweed species other than *A. artemisiifolia*;
- taxonomy of ragweeds to clarify their distribution and relations among them;
- competition mechanisms and strategies with local plant communities (all ragweed species).

In conclusion, "for a fistful of ragweeds" a great deal of work has been done, but it is mandatory to remain alert and not underestimate the role of basic research in elaborating consistent strategies and models (e.g. SDMs) for better understanding and controlling of the ragweed invasion.

## REFERENCES

- Abrams, M. D. 1988. Effects of burning regime on buried seed banks and canopy coverage in a Kansas. *Southwest. Nat.* **33**: 65–70.
- Abul-Fatih, H. A. and Bazzaz, F. A. 1979a. The biology of *Ambrosia trifida* L. II Germination, emergence, growth and survival. *New Phytol.* **83**: 817–827.
- Abul-Fatih, H. A. and Bazzaz, F. A. (1979b). The biology of *Ambrosia trifida* L. I. Influence of species removal on the organization of the plant community. *New Phytol.* **83**: 813–816.
- Abul-Fatih, H., A. Bazzaz, F. A. and Hunt, R. 1979. The biology of *Ambrosia trifida* L. III Growth and biomass allocation. *New Phytol.* **83**: 829-838.

- Acevedo-Rodríguez, P. and Strong, M. T. 2012. *Flora of the West Indies: catalogue of the seed plants of the West Indies*. Washington, DC: Smithsonian Institution, National Museum of Natural History.
- Albertson, F. W. 1937. Ecology of mixed prairie in West Central Kansas. *Ecol. Monogr.* **7**: 481-547.
- Allard, H. A. 1945. Flowering behavior and natural distribution of the Eastern ragweeds (*Ambrosia*) as affected by length of day. *Ecology*, **26**: 387–394.
- Alves, M. and Rocha, N. 2016. Flora of Bahia: Asteraceae – Tribe Heliantheae. *Sitientibus Sér. Ci. Biol.* **16**: 1-63.
- Amor Morales, Á., Navarro Andrés, F. and Sánchez Anta, M. A. 2012. Datos corológicos y morfológicos de las especies del género *Ambrosia* L. (Compositae) presentes en la Península Ibérica. *Bot. Complut.* **36**: 85–96.
- Anderberg, A. 2000a. *Ambrosia artemisiifolia* L. Den Virtuella Flora, Naturhistoriska riksmuseet 2000. <http://linnaeus.nrm.se/flora/di/astera/ambro/ambrrart.html> (24.01.2017)
- Anderberg A. 2000b. *Ambrosia trifida* L. Den Virtuella Flora, Naturhistoriska riksmuseet 2000 . <http://linnaeus.nrm.se/flora/di/astera/ambro/ambrrtri.html> (24.01.2017).
- Anderberg, A. 2005. *Ambrosia psilostachya* DC. Den Virtuella Flora, Naturhistoriska riksmuseet 2000. <http://linnaeus.nrm.se/flora/di/astera/ambro/ambrrpsi.html> (24.01.2017)
- Anton, A., Zuloaga, F.O. and Rúgolo de Agrasar, Z. 2012. *Flora vascular de la República Argentina*. INTA, IMBIV CONICET, IBODA CONICET available at <http://www.floraargentina.edu.ar/>
- Ardenghi, N. M. G. and Polani, F. 2016. La flora della provincia di Pavia (Lombardia, Italia settentrionale). 1. L’Oltrepò Pavese. *Nat. Hist. Sci.* **3**: 51-79.
- Arianoutsou, M., Bazos, I., Delipetrou, P. and Kokkoris, Y. 2010. The alien flora of Greece: taxonomy, life traits and habitat preferences. *Biol. Invasions* **12**: 3525-3549.
- Arslan, Z. F., Uludag, A. and Uremis, I. 2015. Status of invasive alien plants included in EPPO Lists in Turkey. *EPPO Bull.* **45**: 66-72.
- Asero, R., Weber, B., Mistrello, G., Amato, S., Madonini, E. and Cromwell, O. 2005. Giant ragweed specific immunotherapy is not effective in a proportion of patients sensitized to short ragweed: Analysis of the allergenic differences between short and giant ragweed. *J. Allergy Clin. Immunol.* **116**: 1036–1041.
- Auld, B., Morita, H., Nishida, T., Ito, M. and Michael, P. 2003. Shared exotica: Plant invasions of Japan and South Eastern Australia. *Cunninghamia* **8**: 147–152.
- Bacsi, A., Choudhury, B. K., Dharajiya, N. M. D., Sur, S. M. D. and Boldogh, I. 2006. Subpollen particles: Carriers of allergenic proteins and oxidases. *J. Allergy Clin. Immunol.* **118**: 844–850.
- Bae, J., Benoit, D. L. and Watson, A. K. 2016. Effect of heavy metals on seed germination and seedling growth of common ragweed and roadside ground cover legumes. *Environ. Pollut.* **213**: 112–118.
- Baker, D. L. and Guthery, F. S. 1990. Effects of continuous grazing on habitat and density of ground-foraging birds in south Texas. *J. Range Manag.* **43**: 2–5.
- Ballard, T. O., Foley, M. E. and Bauman, T. T. 1996. Germination, viability, and protein changes during cold stratification of giant ragweed (*Ambrosia trifida* L.) seed. *J. Plant Physiol.* **149**: 229–232.
- Bararpour, M.T. 2014. *Weed flora of Mazandaran*. Iran. URL: <https://agritechresearch.files.wordpress.com/2014/06/weed-flora-of-mazandaran.pdf>
- Barina, Z., Rakaj, M. and Pifkó, D. 2013. Contributions to the flora of Albania, 4. *Willdenowia* **43**: 165–184.
- Barina, Z., Rakaj, M., Somogyi, G., Erős-Honti, Z. and Pifkó, D. 2014. The alien flora of Albania: history, current status and future trends. *Weed Res.* **54**: 196–215.

- Barton, M. L., Medel, I. D., Johnston, K. K. and Whitcraft, C. R. 2016. Seed collection and germination strategies for common wetland and coastal sage scrub species in Southern California. *Bull. - South. Calif. Acad. Sci.* **115**: 41-71.
- Barton, J. S., and Schomacker, R. 2017. Comparative protein profiles of the *Ambrosia* plants. *Biochim. Biophys. Acta* **1865**: 633-639.
- Baskin, J. M. and Baskin, C. C. 1980. Ecophysiology of Secondary Dormancy in Seeds of *Ambrosia artemisiifolia*. *Ecology* **61**: 475-480.
- Baskin, J. M. and Baskin, C. C. 1985. The Annual Dormancy Cycle in Buried Weed Seeds: A Continuum. *BioScience* **35**: 492-498.
- Baskin, C. C. and Baskin, J. M. 2014. *Seeds: ecology, biogeography, and, evolution of dormancy and germination*. 2nd ed., San Diego, CA, USA: Academic/Elsevier.
- Bassett, I. J. and Crompton, C. W. 1975. The biology of Canadian Weeds - *Ambrosia artemisiifolia* L. and *Ambrosia psilostachya* DC. *Can. J. Plant Sci.* **55**: 463-476.
- Bassett, I. J. and Crompton, C. W. 1982. The biology of Canadian weeds. 55.: *Ambrosia trifida* L. *Can. J. Plant Sci.* **62**: 1003-1010.
- Battandier, J. A. 1888. *Flore de l'Algérie. Dicotylédones*. 1<sup>er</sup> Fascicule. Thalamiflores. J. A. Battandier & L. Trabut, Eds. Alger: Adolphe Jourdan; Paris: F. Savy
- Baysinger, J. A. and Sims, B. D. 2016. Giant ragweed (*Ambrosia trifida*) interference in soybeans (*Glycine max*). *Weed Sci.* **39**: 358-362.
- Bazzaz, F. A. 1974. Ecophysiology of *Ambrosia artemisiifolia*: a successional dominant. *Ecology* **55**: 112-119.
- Bazzaz, F. A. 1979. The physiological ecology of plant succession. *Annu. Rev. Ecol. Syst.* **10**: 351-371.
- Behçet, L. 2004. A new record for the flora of Turkey: *Ambrosia tenuifolia* Spreng. (Compositae). *Turk. J. Botany* **28**: 201-203.
- Blackburn, T. M., Pyšek, P., Bacher, S., Carlton, J. T., Duncan, R. P., Jarošík, V., Wilson J.R.U. and Richardson, D. M. 2011. A proposed unified framework for Biol. Invasions. *Trends Ecol. Evol.* **26**: 333-339.
- Bojorquez Tapia, L. A. B., Ffolliott, P. F. and Guertin, D. P. 1990. Hbage production-forest overstory relationships in two Arizona ponderosa pine forests. *J. Range Manag.* **43**: 25-28.
- Bordas-le Floch, V., Groeme, R., Chabre, H., Baron-Bodo, V., Nony, E., Mascarell, L. and Moingeon, P. 2015. New insights into ragweed pollen allergens. *Curr. Allergy Asthma Rep.* **15**: 1-7.
- Borges, P. A., Abreu, C., Aguiar, A. M. F., Carvalho, P., Jardim, R., Melo, I. Oliveira, P., Sérgio, C., Serrano, A. R. M. and Vieira P. 2008. *Listagem dos fungos, flora e fauna terrestres dos arquipélagos da Madeira e Selvagens - A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos*. Secretaria Regional do Ambiente e dos Recursos Naturais do Governo Regional da Madeira, Funchal, Madeira.
- Bouley, J., Groeme, R., Le Mignon, M., Jain, K., Chabre, H., Bordas-Le Floch, V., Couret M.N., Bussièrès, L., Lautrette, A., Naveau, M., Baron-Bodo, V., Lombardi, V., Mascarell, L., Batard, T., Nony, E. and Moingeon, P. 201). Identification of the cysteine protease Amb a 11 as a novel major allergen from short ragweed. *J. Allergy Clin. Immunol.* **136**: 1055-1064.
- Boulos, L. 2002. Flora of Egypt, Volume 3 (Vebenaceae-Compositae). Al Hadara Publishing, Cairo, Egypt.
- Bovey, R. W., McCarty, M. K. and Davis, F. S. 1966. Control of Perennial Ragweed on Western Nebraska Rangeland. *J. Range Manag.* **19**: 220-222.
- Bozarth, S. R. 1992. Classification of opal phytoliths formed in selected Dicotyledons native to the Great Plains. **In**: *Phytolith systematics*. pp. 193-214. George Rapp Jr., G. and Mulholland, S. C., Eds., Springer, US.

- Buttler, K.P. 2016. *Florenliste von Deutschland – Gefäßpflanzen*. Version 8 <http://www.kp-buttler.de/florenliste/index.htm>
- Buttler, K. P. and Harms, K. H. 1999. *Florenliste von Baden-Württemberg, Liste der Farn- und Samenpflanzen (Pteridophyta et Spermatophyta)*. Naturschutz-Praxis, Artenschutz Karlsruhe 1.
- Brandes, D. and Nitzsche, J. 2006. Biology, introduction, dispersal, and distribution of common ragweed (*Ambrosia artemisiifolia* L.) with special regard to Germany. *Nachrichtenblatt-Deutschen Pflanzenschutzdienstes Braunschweig* **58**: 286–291.
- Bullock, J. M., Chapman, D., Schafer, S., Roy, D., Girardello, M., Haynes, T., *et al.* 2012. *Assessing and controlling the spread and the effects of common ragweed in Europe (ENV.B2/ETU/2010/0037)*. European Commission, Final Report.
- Burkart, S. E., Leon, R. J. and Movia, C. P. 1990. Inventario fitosociológico del pastizal de la Depresión del Salado (Prov. Bs. As.) en un área representativa de sus principales ambientes. *Darwiniana* **30**: 27-69.
- Busby, R. R., Gebhart, D. L., Stromberger, M. E., Meiman, P. J. and Paschke, M. W. 2011. Early seral plant species' interactions with an arbuscular mycorrhizal fungi community are highly variable. *Applied soil ecology* **48**: 257-262.
- Busse, W., Reed, C. E. and Hoehne, J. H. 1972. Where is the allergic reaction in ragweed asthma? *J. Allergy Clin. Immunol.* **50**: 289–293.
- Buttenschøn, R. M., Waldispühl, S. and Bohren, C. 2010. *Guidelines for management of common ragweed, Ambrosia artemisiifolia*. University of Copenhagen.
- Byfield, A. J. and Baytop, A. 1998. Three alien species new to the flora of Turkey. *Turk. J. Botany* **22**: 205-208.
- CABI, 2017. *Invasive Species Compendium*. Wallingford, UK: CAB International. [www.cabi.org/isc](http://www.cabi.org/isc).
- Campbell-Kissock, L., Blankenship, L. H. and Stewart, J. W. 1985. Plant and animal foods of bobwhite and scaled quail in Southwest Texas. *Southwest. Nat.* **30**:543–553.
- Carls, E. G., Lonard, R. I. and Dennis, B. 1991. Notes on the Vegetation and Flora of North Padre Island, Texas. *Southwest. Nat.* **36**:121–125.
- Catford, J. A. and Jansson, R. 2014. Drowned, buried and carried away: effects of plant traits on the distribution of native and alien species in riparian ecosystems. *New Phytol.* **204**: 19-36.
- Celesti-Grapow, L., Alessandrini, A., Arrigoni, P.V., Banfi, E., Bernardo, L., Bovio, M., Brundu, G., Cagiotti, M. R., Camarda, I., Carli, E., Conti, F., Fascetti, S., Galasso, G., Gubellini, L., La Valva, V., Lucchese, F., Marchiori, S., Mazzola, P., Peccenini, S., Poldini, L., Pretto, F., Prosser, F., Siniscalco, C., Villani, M. C., Viegi, L., Wilhalm, T., Blasi, C., 2009. Inventory of the non-native flora of Italy. *Plant Biosyst.* **143**: 386-430.
- Chamberlin, R. V. 1911. The Ethno-Botany of the Gosiute Indians. *Proc. Acad. Nat. Sci. Philadelphia* **63**: 24–99.
- Chapman, D.S., Makra, L., Albertini, R., Bonini, M., Páldy, A., Rodinkova, V., Šikoparija, B., Weryszko-Chmielewska, E. and Bullock, J.M. 2016. Modelling the introduction and spread of non-native species: International trade and climate change drive ragweed invasion. *Glob. Chang. Biol.* **22**: 3067-3079.
- Chauvel, B., Dessaint, F., Cardinal-Legrand, C. and Bretagnolle, F. 2006. The historical spread of *Ambrosia artemisiifolia* L. in France from herbarium records. *J. Biogeogr.* **33**: 665–673.
- Chauvel, B., Rodriguez, A., Moreau, C., Martinez, Q., Bilon, R. and Fried, G. 2015. Développement d'*Ambrosia trifida* L. en France: connaissances historiques et écologiques en vue d'une éradication de l'espèce. *J. Bot. Soc. Bot. France* **71**: 25–38.

- Chen, Y. S. and Hind, D. J. N. 2011. Heliantheae. **In:** Wu, Z. Y., Raven, P. H. & Hong, D. Y., eds., *Flora of China Volume 20–21 (Asteraceae)*. pp. 852–878. Science Press (Beijing) & Missouri Botanical Garden Press (St. Louis).
- Cheraghian, A. 2016a. *A guide for detection and diagnosis of quarantine pests: Perennial ragweed *Ambrosia psilostachya* DC. Asterales:Asteraceae*. Islamic Republic of Iran Ministry of Jihad -e- Agriculture Plant Protection Organization.
- Cheraghian, A. 2016b. *A guide for detection and diagnosis of quarantine pests: Giant ragweed *Ambrosia trifida* L. Asterales:Asteraceae*. Islamic Republic of Iran Ministry of Jihad -e- Agriculture Plant Protection Organization.
- Ciappetta, S., Ghiani, A., Gilardelli, F., M., B., Citterio, S. and Gentili, R. 2016. Invasion of *Ambrosia artemisiifolia* in Italy: Assessment via analysis of genetic variability and herbarium data. *Flora-Morphology, Distrib. Funct. Ecol. Plants* **223**:173–178.
- Conti, F., Abbate, G. , Alessandrini, A. , Blasi, C. 2005. *An annotated checklist of the Italian vascular flora*. Fratelli Palombi Roma.
- Corbett, E. and Anderson, R. C. 2006. Landscape analysis of Illinois and Wisconsin remnant prairies. *J. Torrey Bot. Soc.* **133**: 267-279.
- Csontos, P., Vitalos, M., Barina, Z. and Kiss, L. 2010. Early distribution and spread of *Ambrosia artemisiifolia* in Central and Eastern Europe. *Bot. Helv.* **120**: 75–78
- Cui, S., Zhou, Q. and Chao, L. 2007. Potential hyperaccumulation of Pb, Zn, Cu and Cd in enduring plants distributed in an old smeltery, Northeast China. *Environ. Geol.* **51**: 1043-1048.
- Cunze, S., Leiblein, M. C. and Tackenberg, O. 2013. Range Expansion of *Ambrosia artemisiifolia* in Europe is promoted by climate change. *ISRN Ecol.* 2013, 1–9.
- Dahl, Å., Strandhede, S. O. and Wihl, J. Å. 1999. Ragweed - An allergy risk in Sweden? *Aerobiologia*, **15**: 293–297.
- Dalrymple, R. L. and Rogers, J. L. 1983. Allelopathic effects of Western ragweed on seed germination and seedling growth of selected plants. *J. Chem. Ecol.*, **9**: 1073–1078.
- Danin, A. 2000. The nomenclature news of Flora Palaestina. *Flora Mediterranea* **10**: 109-172.
- Danin, A. 2016. Analytical Flora of Isarel: *Ambrosia trifida* L. available at <http://flora.org.il/en/plants/AMBTRI/>
- Davis, W. E. 1930. Primary dormancy, after-ripening , and the development of secondary dormancy in embryos of *Ambrosia trifida*. *Am. J. Bot.* **17**: 58–76.
- D’Amato, G., Cecchi, L., Bonini, S., Nunes, C., Annesi-Maesano, I., Behrendt, H., Liccardi, G., Popov, T. and Van Cauwenberge, P. (2007). Allergenic pollen and pollen allergy in Europe. *Allergy Eur. J. Allergy Clin. Immunol.* **62**: 976–990.
- D’Amato, G. and Spiekma, F. T. M. 1992. European allergenic pollen types. *Aerobiologia*, **8**: 447–450.
- De Egea, J., Mereles, F., Del Carmen Pena-Chocarro, M. and Céspedes, G. 2016. Checklist for the crop weeds of Paraguay. *PhytoKeys*, **73**: 13-92.
- Del Vecchio, S., Pizzo, L. and Buffa, G. 2015. The response of plant community diversity to alien invasion: evidence from a sand dune time series. *Biodivers. Conserv.* **24**: 371–392.
- Del Vitto, L. A., Petenatti, E. M. and Petenatti, M. E. 1997. Recursos herbolarios de San Luis (República Argentina) Primera parte: plantas nativas. *Multequina*, **6**: 49–66.
- Del Vitto, L. A. and Petenatti, E. M. 2015. Asteráceas de importancia económica y ambiental. Segunda parte: Otras plantas útiles y nocivas. *Multequina*, **24**: 47-74.
- Dickerson, C. T. 1968. Studies on the germination, growth, development and control of common ragweed, *Ambrosia artemisiifolia* L. PhD thesis, Cornell University.
- Dickerson, C. T. and Sweet, R. D. 1971. Common Ragweed Ecotypes. *Weed Sci.* **19**: 64–66.
- DiTommaso, A. 2004. Germination behavior of common ragweed (*Ambrosia artemisiifolia*) populations across a range of salinities. *Weed Sci.* **52**: 1002-1009.

- Dullinger, S., Kleinbauer, I., Peterseil, J., Smolik, M. and Essl, F. 2009. Niche based distribution modelling of an invasive alien plant: effects of population status, propagule pressure and invasion history. *Biol. Invasions* **11**: 2401–2414.
- Ebinger, J. E., Phillippe, L. R., Nyboer, R. W., McClain, W. E., Busemeyer, D. T., Robertson, K. and Levin, G. A. 2006. Vegetation and flora of the sand deposits of the Mississippi River Valley in North Western Illinois. *Bull. Ill. Nat. Hist. Surv.* **37**: 1-56.
- El Kelish, A., Zhao, F., Werner, H., Durner, J., Winkler, J. B., Behrendt, H., Traidl-Hoffmann, C., Horres, R., Pfeifer, M., Frank, U. and Ernst, D. 2014. Ragweed (*Ambrosia artemisiifolia*) pollen allergenicity: SuperSAGE transcriptomic analysis upon elevated CO<sub>2</sub> and drought stress. *BMC Plant Biol.* **14**: 1–16.
- Ellis, B. A., Mills, J. N., Glass, G. E., Mckee, K. T., Delia, A. and Childs, J. E. 1998. Dietary habits of the common rodents in an agroecosystem in Argentina. *J. Mammal.* **79**: 1203–1220.
- Entry, J. A., Rygielwicz, P. T., Watrud, L. S. and Donnelly, P. K. 2002. Influence of adverse soil conditions on the formation and function of arbuscular mycorrhizas. *Adv. Environ. Res.* **7**: 123-138.
- EPPO, 2016. PQR - EPPO database on quarantine pests (available online). <http://www.eppo.int>
- Escudero, V. and Mendoza, R. 2005. Seasonal variation of arbuscular mycorrhizal fungi in temperate grasslands along a wide hydrologic gradient. *Mycorrhiza*, **15**: 291-299.
- Essl, F. and Rabitsch, W. 2002. Neobiota in Österreich. Umweltbundesamt, Wien.
- Essl, F., Dullinger, S., and Kleinbauer, I. 2009. Changes in the spatio-temporal patterns and habitat preferences of *Ambrosia artemisiifolia* during its invasion of Austria. *Preslia* **81**: 119–133
- Essl, F., Biró, K., Brandes, D., Broennimann, O., Bullock, J. M., Chapman, D. S., Chauvel, B., Dullinger, S., Fumanal, B., Guisan, A., Karrer, G., Kazinczi, G., Kueffer, C., Laitung, B., Lavoie, C., Leitner, M., Mang, T., Moser, D., Müller-Schärer, H., Petitpierre, B., Richter, R., Schaffner, U., Smith, M., Starfinger, U., Vautard, R., Vogl, G., Von Der Lippe, M. & Follak, S. 2015. Biological Flora of the British Isles: *Ambrosia artemisiifolia*. *J. Ecol.* **103**: 1069–1098.
- Estrada, A., Morales-Castilla, I., Caplat, P. and Early, R. 2016. Usefulness of species traits in predicting range shifts. *Trends Ecol. Evol.* **31**: 190-203
- Fenesi, A., Albert, Á. J. and Ruprecht, E. 2014. Fine-tuned ability to predict future competitive environment in *Ambrosia artemisiifolia* seeds. *Weed Res.* **54**: 58-69.
- Ferus, P., Sîrbu, C., Eliáš, P., Konôpková, J., Ďurišová, L., Samuil, C. and Oprea, A. 2015. Reciprocal contamination by invasive plants: Analysis of trade exchange between Slovakia and Romania. *Biologia*, **70**: 893–904.
- Finch-Savage, W. E. and Leubner-Metzger, G. 2006. Seed dormancy and the control of germination. *New Phytol.* **171**: 501-523.
- Finnish Ministry of Agriculture and Forestry, 2012. *Finland's National Strategy on Invasive Alien Species*. Helsinki, Finland.
- Flores-Olvera, H., Czaja, A., Estrada-Rodríguez, J. L. and Méndez, U. R. 2016. Floristic diversity of halophytic plants of Mexico. **In**: *Sabkha Ecosystems Volume V: The Americas*. pp. 299-327. Springer International Publishing.
- Follak, S., Dullinger, S., Kleinbauer, I., Moser, D. and Essl, F. 2013. Invasion dynamics of three allergenic invasive asteraceae (*Ambrosia trifida*, *Artemisia annua*, *Iva xanthiifolia*) in Central and Eastern Europe. *Preslia*, **85**: 41–61.
- Fontana, S. L. 2005. Coastal dune vegetation and pollen representation in South Buenos Aires Province, Argentina. *J. Biogeogr.* **32**: 719–735.
- Fraga, P. and García, Ó. 2004. Notes i contribucions al coneixement de la flora de Menorca (VI). *Bol. Soc. Hist. Nat. Balears* **47**: 143-152.



- Freire, S. E., Arambarri, A. N. A. M., Bayón, N. D., Sancho, G., Urtubey, E., Monti, C., Novoa, M. and Colares, M. N. 2005. Epidermal characteristics of toxic plants for cattle from the Salado river basin (Buenos Aires, Argentina). *Boletín la Soc. Argentina Botánica* **40**: 241–281.
- Fremstad, E. and Elven, R. 1997. Alien plants in Norway and dynamics in the flora: a review. *Nor. Geogr. Tidsskr* **51**:199–218.
- Frick, G., Boschung, H., Schulz-schroeder, G., Russ, G., Ujčič-vrhovnik, I., Jakovac-strajn, B., Angetter, D., John, I. and Jørgensen, J. S. 2011. Ragweed (*Ambrosia* sp.) seeds in bird feed. *Biotechnol. Agron. Société Environ.* **15**: 39–44.
- Fried, G., Belaud, A. and Chauvel, B. 2015. Ecology and impact of an emerging invasive species in France: Western ragweed (*Ambrosia psilostachya* DC.). *Rev. Ecol.-Terre Vie* **70**: 53-67.
- Fuentes, N., Pauchard, A., Sánchez, P., Esquivel, J. and Marticorena, A. 2013. A new comprehensive database of alien plant species in Chile based on herbarium records. *Biol. Invasions* **15**: 847-858
- Fukano Y. and Yahara T 2012. Changes in defense of an alien plant *Ambrosia artemisiifolia* before and after the invasion of a native specialist enemy *Ophraella communa*. *PLoS One* **7**: 10.1371/annotation/868d00f2-375e-421f-8435-0e628c0567bd.
- Fumanal, B., Girod, C., Fried, G., Bretagnolle, F. and Chauvel, B. 2008a. Can the large ecological amplitude of *Ambrosia artemisiifolia* explain its invasive success in France? *Weed Res.* **48**: 349–359.
- Fumanal, B., Gaudot, I. and Bretagnolle, F. 2008b. Seed-bank dynamics in the invasive plant, *Ambrosia artemisiifolia* L. *Seed Sci. Res.* **18**: 101-114.
- Fumanal, B., Chauvel, B., Sabatier, A. and Bretagnolle, F. 2007. Variability and cryptic heteromorphism of *Ambrosia artemisiifolia* seeds: what consequences for its invasion in France? *Ann. Bot.* **100**: 305–313.
- Funderburg, E. R., Locke, J. M. and Biermacher, J. T. 2014. Evaluation of Aminopyralid Applied PRE to Control Western Ragweed (*Ambrosia psilostachya*) in Oklahoma Pastureland. *Weed Technol.* **28**: 395–400.
- Gadermaier, G., Hauser, M. and Ferreira, F. 2014. Allergens of weed pollen: An overview on recombinant and natural molecules. *Methods*, 66: 55–66.
- Galzina, N., Barić, K., Šćepanović, M., Goršić, M. and Ostojić, Z. 2010. Distribution of invasive weed *Ambrosia artemisiifolia* L. in Croatia. *Agric. Conspec. Sci.* **75**: 75-81.
- Gann, G.D., J.C. Trejo-Torres and C.G. Stocking. 2015-2017. *Plantas de la Isla de Puerto Rico/Plants of the Island of Puerto Rico*. The Institute for Regional Conservation. Delray Beach, Florida USA.
- Gard, B., Bretagnolle, F., Dessaint, F. and Laitung, B. 2013. Invasive and native populations of common ragweed exhibit strong tolerance to foliar damage. *Basic Appl. Ecol.* **14**: 28-35.
- Gaudeul, M., Giraud, T., Kiss, L. and Shykoff, J. A. 2011. Nuclear and chloroplast microsatellites show multiple introductions in the worldwide invasion history of common ragweed, *Ambrosia artemisiifolia*. *PLoS One*, **6**: e17658. doi:10.1371/journal.pone.0017658
- Gentili, R., Gilardelli, F., Ciappetta, S., Ghiani, A. and Citterio, S. (2015). Inducing competition: Intensive grassland seeding to control *Ambrosia artemisiifolia*. *Weed Res.* **55**: 278–288.
- Gentili, R., Gilardelli, F., Bona, E., Prosser, F., Selvaggi, A., et al. (2016). Distribution map of *Ambrosia artemisiifolia* L. (Asteraceae) in Italy. *Plant Biosyst.* **151**: 381-386.
- Gentili R., Montagnani C., Gilardelli F., Guarino M. F., Citterio S. 2017. Let native species take their course: *Ambrosia artemisiifolia* replacement during natural or “artificial” succession, *Acta Oecologica*, 82, 32-40.

- Genton, B. J., Shykoff, J. A. and Giraud, T. 2005. High genetic diversity in French invasive populations of common ragweed, *Ambrosia artemisiifolia*, as a result of multiple sources of introduction. *Mol. Ecol.* **14**: 4275–4285.
- Gerber, E., Schaffner, U., Gassmann, A., Hinz, H. L., Seier, M. and Müller-Schärer, H. 2011. Prospects for biological control of *Ambrosia artemisiifolia* in Europe: learning from the past. *Weed Res.* **51**: 559–573.
- Germishuizen, G. and Meyer, N.L. 2003. Plants of Southern Africa: an annotated checklist. *Strelitzia*, 14. National Botanical Institute, Pretoria. Download from POSA (<http://posa.sanbi.org>) on January 25, 2017.
- Ghiani, A., Aina, R., Asero, R., Bellotto, E. and Citterio, S. 2012. Ragweed pollen collected along high-traffic roads shows a higher allergenicity than pollen sampled in vegetated areas. *Allergy*, **67**: 887–94.
- Ghosh, B., Perry, M. P., Bassolino-klimas, D., Rafnar, T., Klapper, D. G. and Marsh, D. G. 1994. Immunologic and molecular characterization of Amb p V allergens from *Ambrosia psilostachya* (Western Ragweed) pollen. *J. Immunol.* **152**:2882–2889.
- Gioanetto, F., Díaz Vilchis, J. T. and Quintero Sánchez, R. 2010. *Manual de utilización de las malezas silvestres de Michoacán. Usos allelopáticos, agroecológicos, medicinales, alimentarios, veterinarios y rituales de las adventicias silvestres y arvenses*. Graphopolis. Morelia.
- Gioria, M., Pyšek, P. and Osborne, B. A. 2016. Timing is everything: does early and late germination favor invasions by herbaceous alien plants? *J Plant Ecol.* doi:10.1093/jpe/rtw105
- Girodet, B. 2013. Les allergènes de l' ambrosie. *Rev. Fr. Allergol.* **53**: 473–476.
- Giscafre, L. and Ragonese, A. E. 1942. Importancia del género *Ambrosia* como factor responsable de polinosis en la provincia de Santa Fe. *Darwiniana*, **6**: 31–44.
- Goeden, R. D. and Ricker, D. W. 1976. The phytophagous insect fauna of the ragweed, *Ambrosia psilostachya*, in Southern California. *Environ. Entomol.* **5**: 1169-1177.
- Goplen, J. J., Sheaffer, C. C., Becker, R. L., Coulter, J. A., Breitenbach, F. R., Behnken, L. M., Johnson, G. A. and Gunsolus, J. L. 2016. Giant Ragweed (*Ambrosia trifida*) seed production and retention in soybean and field margins. *Weed Technol.* **30**: 246–253.
- Greuter, W., 2006: Compositae (pro parte majore). In: Greuter, W. & Raab-Straube, E. von (ed.): *Compositae. Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity*. Published on the Internet: <http://ww2.bgbm.org/EuroPlusMed/> [24.01.2017].
- Greuter, W. and Raus, T. 2008. Med-checklist Notulae 27. *Willdenowia* **38**: 465-474.
- Greuter, W. and Raus, T. 1995. Med checklist Notulae 16. *Willdenowia* **25**: 171-176.
- Grime, J. P. 2001. *Plant strategies, vegetation processes, and ecosystem properties*. 2nd ed., John Wiley & Sons, New York.
- Grygiel, C. E., Norland, J. E. and Biondini, M. E. 2012. Can Carbon and Phosphorous amendments increase native forbs in a restoration process? A case study in the Northern tall-grass prairie (USA). *Restor. Ecol.* **20**: 122-130.
- Gudzinskas, Z. 1993. Genus *Ambrosia* L.(Asteraceae) in Lithuania. *Thaiszia* **3**: 89–96.
- Gutte, P. 1978. Beitrag zur Kenntnis zentralperuanischer Pflanzengesellschaften I Ruderalpflanzengesellschaften von Lima und Huanuco. *Feddes Repert.* **89**: 75-97.
- Guillemin, J. P. and Chauvel, B. 2011. Effects of the seed weight and burial depth on the seed behavior of common ragweed (*Ambrosia artemisiifolia*). *Weed Biol. Manag.* **11**: 217-223.
- Hanley, M. E., Lamont, B. B., Fairbanks, M. M. and Rafferty, C. M. 2007. Plant structural traits and their role in anti-herbivore defence. *Perspect. Plant Ecol. Evol. Syst.* **8**: 157-178.
- Harrison, K. S., Regnier, E. E., Schmoll, J. T. and Webb, J. E. 2001. Competition and fecundity of giant ragweed in corn. *Weed Sci.* **49**: 224–229.

- Harrison, S. K., Regnier, E. E. and Schmoll, J. T. 2003. Postdispersal predation of giant ragweed (*Ambrosia trifida*) seed in no-tillage corn. *Weed Sci.* **51**: 955–964.
- Harrison, S. K., Regnier, E. E., Schmoll, J. T. and Harrison, J. M. 2007. Seed size and burial effects on giant ragweed (*Ambrosia trifida*) emergence and seed demise. *Weed Sci.* **55**: 16–22.
- Hart, C. A. and Gleason, H. A. 1907. On the biology of the sand areas of Illinois. *Bull. Ill. Nat. Hist. Surv.* **7**: 137-272.
- Hartnett, D. C., Hickman, K. R. and Walter, L. E. F. 1996. Effects of bison grazing, fire, and topography on floristic diversity in tallgrass prairie. *J. Range Manag.* **49**: 413-420.
- Hartnett, D. C., Hartnett, B. B. and Bazzaz, F. A. 1987. Persistence of *Ambrosia trifida* populations in old fields and responses to successional changes. *Am. J. Bot.* **74**: 1239–1248.
- Heinrich, M., Robles, M., West, J. E., Ortiz de Montellano, B. R. and Rodriguez, E. 1998. Ethnopharmacology of Mexican Asteraceae (Compositae). *Ann. Rev. Pharm. Toxicol.* **38**: 539-565.
- Hejný, S. and Jelík, V. 1972. Hemerochorous dispersal of adventitious plants from the viewpoint of frequency of different ways of introduction - a proposal of terminology. *Folia Geobot. Phytotaxon.* **7**: 91–93.
- Henderson, L. 2007. Invasive, naturalized and casual alien plants in Southern Africa: a summary based on the Southern African Plant Invaders Atlas (SAPIA). *Bothalia*, **37**: 215–248.
- Hess, H. E., Landolt, E., Müller-Hirzel, R. and Baltisberger, M. 2006. *Bestimmungsschlüssel zur Flora der Schweiz und angrenzender Gebiete*. Birkhauser Verlag.
- Hibon, G. 1942. *Ambrosia psilostachya* adventice parisienne. *Bull. Soc. Bot. France* **89**: 233-234,
- Hodgins, K. A. and Rieseberg, L. 2011. Genetic differentiation in life-history traits of introduced and native common ragweed (*Ambrosia artemisiifolia*) populations. *J. Evol. Biol.* **24**: 2731–2749.
- Hodgins, K. A., Lai, Z., Nurkowski, K., Huang, J. and Rieseberg, L. H. 2013. The molecular basis of invasiveness: differences in gene expression of native and introduced common ragweed (*Ambrosia artemisiifolia*) in stressful and benign environments. *Mol. Ecol.* **22**: 2496-2510.
- Howell, C. J. and Sawyer, J. W. D. 2006. *New Zealand naturalised vascular plant checklist*. New Zealand Plant Conservation Network, Wellington.
- Hulett, G. K., Tomelleri, J. R. and Hampton, C. O. 1988. Vegetation and flora of a sandsage prairie site in Finney County, Southwestern Kansas. *Trans. Kans. Acad. Sci.* **91**: 83-95.
- Insausti, P. and Grimoldi, A. A. 2006. Gap disturbance triggers the recolonization of the clonal plant *Ambrosia tenuifolia* in a flooding grassland of Argentina. *Austral Ecol.* **31**: 828–836.
- Insausti, P. and Soriano, A. 1982. Comportamiento de las semillas de *Ambrosia tenuifolia* (altamisa) en un pastizal de la Depresion del Salado (provincia de Buenos Aires). *Revista Fac. Agron.* **3**: 75-80
- Insausti, P. and Soriano, A. 1987. Efecto del anegamiento prolongado en un pastizal de la Depresion del Salado (provincia de Buenos Aires — Argentina): dinamica del pastizal en conjunto y de *Ambrosia tenuifolia* (Asteraceae). *Darwiniana*, **28**: 397–403.
- Insausti, P., Soriano, A. and Sanchez, R. A. 1995. Effects of flood-influenced factors on seed germination of *Ambrosia tenuifolia*. *Oecologia*, **103**: 127–132.
- Jacobson, A. R. and Morris, S. C. 1976. The Primary Air Pollutants—Viable Particulates Their Occurrence, Sources, and Effects. In: Stern A.C. (eds.) *Air Pollution VI: Air Pollutants, Their Transformation and Transport*, V.1, Academic Press, New York.
- Jaramillo Díaz, P. and Guézou, A. 2013. CDF Checklist of Galapagos Vascular Plants - FCD Lista de especies de Plantas Vasculares de Galápagos. In: Bungartz, F., Herrera, H.,

- Jaramillo, P., Tirado, N., Jiménez-Uzcátegui, G., Ruiz, D., Guézou, A. & Ziemmeck, F. (eds.). *Charles Darwin Foundation Galapagos Species Checklist - Lista de Especies de Galápagos de la Fundación Charles Darwin*. Charles Darwin Foundation / Fundación Charles Darwin, Puerto Ayora, Galapagos.
- Joly, M., Bertrand, P., Gbangou, R. Y., White, M. C., Dubé, J. and Lavoie, C. 2011. Paving the way for invasive species: road type and the spread of common ragweed (*Ambrosia artemisiifolia*). *Environ. Manage.* **48**: 514-522.
- Jørgensen, P. M., and S. León-Yáñez (eds.) 1999. Catalogue of the vascular plants of Ecuador. *Monogr. Syst. Bot. Missouri Bot. Gard.* **75**: 1–1182. Missouri Botanical Garden Press, St. Louis.
- Jørgensen, P. M., M. H. Nee and S. G. Beck. (eds.) 2014. Catálogo de las plantas vasculares de Bolivia. *Monogr. Syst. Bot. Missouri Bot. Gard.* **127**: 1–1744. Missouri Botanical Garden Press, St. Louis.
- Jurik, T. W. 1991. Population distributions of plant size and light environment of giant ragweed (*Ambrosia trifida* L.) at three densities. *Oecologia*, **87**: 539-550.
- Jurney, D. H. 2012. Anthropology of fire in the Ozark Highland Region. **In**: D. C. Dey, M. C. Stambaugh, S. L. Clark and C. J. Schweitzer (Eds.), *Proceedings of the 4th Fire in Eastern Oak Forests Conference*. pp. 12–33. Northern Research Station, Newtown Square, Pennsylvania, USA: USDA Forest Service General Technical Report GTR-NRS-P-102.
- Kanter, U., Heller, W., Durner, J., Winkler, J. B., Engel, M., Behrendt, H., Holzinger, A., Braun, P., Hauser, M., Ferreira, F., Mayer, K., Pfeifer, M. and Ernst, D. 2013. Molecular and immunological characterization of ragweed (*Ambrosia artemisiifolia* L.) pollen after exposure of the plants to elevated ozone over a whole growing season. *PLoS One*, **8**: e61518. doi:10.1371/journal.pone.0061518
- Karnkowski W. 2001. Can the weeds be recognized as quarantine pests? - Polish experiences with *Ambrosia* spp. **In**: *Proceedings of the 5th Slovenian Conference on Plant Protection, Catez ob Savi, Slovenia, 6-8 March 2001*. pp. 396-402. Društvo za varstvo rastlin Slovenije.
- Karrer G. 2014. Das österreichische Ragweed Projekt – übertragbare Erfahrungen? *Julius-Kühn-Archiv* **445**: 27–33. doi: 10.5073/jka.2013.445.003.
- Karrer, G. 2016. *Morphological and synecological deterioration of European Ambrosia species in the Northern Mediterranean*. Report COST-STSM-ECOST-STSM-FA1203-070415-059092
- Karrer, G., Hall, R., Lener, F., Waldhäuser, N., Kazinczi, G., Kerepesi, I. Máté, S., Sölter, U., Starfinger, U., Verschwele, A., Mathiassen, S. K., Kudsk, P., Leskovšek, R. and Simončič A. 2016. Field experiment on longevity of the seeds in the soil seed bank (joint experiment). *Halt Ambrosia-Final Project Report and General Publication of Project Findings*.
- Katz, D. S. W. and Carey, T. S. 2014. Heterogeneity in ragweed pollen exposure is determined by plant composition at small spatial scales. *Sci. Total Environ.* **485**: 435–440. <http://doi.org/10.1016/j.scitotenv.2014.03.099>
- Kazinczi, G., Béres, I., Novák, R., Bíró, K. and Pathy, Z. 2008. Common ragweed (*Ambrosia artemisiifolia*): a review with special regards to the results in Hungary. I. Taxonomy, origin and distribution, morphology, life cycle and reproduction strategy. *Herbologia*, **9**: 55–91.
- Keeley, J. E., Pausas, J. G., Rundel, P. W., Bond, W. J. and Bradstock, R. A. 2011. Fire as an evolutionary pressure shaping plant traits. *Trends Plant Sci.* **16**: 406-411.
- Kohli, R.K., Batishi, D.R., Singh, J.S., Singh, H.P., Bhatt J.R. 2012. Plant Invasion in India: an overview. **In**: *Invasive alien plants an ecological appraisal for the Indian subcontinent*. pp: 1-9. Eds. Bhatt JR, Singh JS, Singh SP, Tripathi RS and Kohli RK

- Khuroo, A. A., Reshi, Z. A., Malik, A. H., Weber, E., Rashid, I. and Dar, G. H. 2012. Alien flora of India: taxonomic composition, invasion status and biogeographic affiliations. *Biol. Invasions* **14**: 99-113.
- Kikodze, D., Memiadze, N., Kharazishvili, D., Manvelidze, Z. and Mueller-Schaerer, H. 2010. *The Alien Flora of Georgia*. 2nd ed. Swiss National Science Foundation, Swiss Agency for Development and Cooperation and SCOPES (project number IB73A0-110830).
- Kim, C. G. and Kil, J. 2016. Alien flora of the Korean Peninsula. *Biol. Invasions* **18**: 1843-1852.
- Kiss, L. and Béres, I. 2006. Anthropogenic factors behind the recent population expansion of common ragweed (*Ambrosia artemisiifolia* L.) in Eastern Europe: is there a correlation with political transitions? *J. Biogeogr.* **33**: 2156–2157.
- Kong, C. H. 2010. Ecological pest management and control by using allelopathic weeds (*Ageratum conyzoides*, *Ambrosia trifida*, and *Lantana camara*) and their allelochemicals in China. *Weed Biol. Manag.* **10**: 73-80.
- Kong, C. H., Wang, P. and Xu, X. H. 2007. Allelopathic interference of *Ambrosia trifida* with wheat (*Triticum aestivum*). *Agric. Ecosyst. Environ.* **119**: 416-420.
- Kull, C. A., Tassin, J., Moreau, S., Ramiarantsoa, H. R., Blanc-Pamard, C. and Carrière, S. M. 2012. The introduced flora of Madagascar. *Biol. Invasions* **14**: 875–888.
- Kumar, P., Singh, P.K. and Dubey, R.K. 2009. *National conference on Invasive Alien Species – A threat to native biodiversity*. Uttar Pradesh State Biodiversity Board, Lucknow, U.P.
- Lampinen, R. & Lahti, T. 2016. Kasviatlas -- Helsingin Yliopisto, Luonnontieteellinen keskusmuseo, Helsinki. Levinneisyyskartat osoitteessa <http://www.luomus.fi/kasviatlas>.
- Lavoie, C., Jodoin, Y. and Goursaud de Merlis, A. 2007. How did common ragweed (*Ambrosia artemisiifolia* L.) spread in Québec? A historical analysis using herbarium records. *J. Biogeogr.* **34**: 1751–1761.
- Lawalree A. 1947. Les Ambrosia adventices en Europe Occidentale. *Bull. Jard. Bot. État Bruxelles* **18**: 305-315.
- Lee, C. S., Cho, Y. C., Shin, H. C., Kim, G. S., and Pi, J. H. 2010. Control of an invasive alien species, *Ambrosia trifida* with restoration by introducing willows as a typical riparian vegetation. *J.Ecol. Environ.* **33**: 157-164.
- Leiblein, M. C. and Lösch, R. 2011. Biomass development and CO<sub>2</sub> gas exchange of *Ambrosia artemisiifolia* L. under different soil moisture conditions. *Flora-Morphology, Distrib. Funct. Ecol. Plants* **206**: 511-516.
- Leiblein-Wild, M. C., Kaviani, R. and Tackenberg, O. 2014. Germination and seedling frost tolerance differ between the native and invasive range in common ragweed. *Oecologia*, **174**: 739–750.
- Lenoir, I., Fontaine, J. and Sahraoui, A. L. H. 2016. Arbuscular mycorrhizal fungal responses to abiotic stresses: A review. *Phytochemistry*, **123**: 4-15.
- Li, X. M., She, D. Y., Zhang, D. Y. & Liao, W. J. 2015. Life history trait differentiation and local adaptation in invasive populations of *Ambrosia artemisiifolia* in China. *Oecologia* **177**: 669–677.
- Liogier, A. H. 1997. *Flora of Puerto Rico and adjacent islands*. Editorial Universidad de Puerto Rico.
- Lippert, R. D. and Hopkins, H. H. 1950. Study of viable seeds in various habitats in mixed prairie. *Trans. Kans. Acad. Sci.* **53**: 355-364.
- MacDonald, A. A. M. and Kotanen, P. M. 2010. The effects of disturbance and enemy exclusion on performance of an invasive species, common ragweed, in its native range. *Oecologia*, **162**: 977–986.
- Macdonald, I.A.W., J.K. Reaser, Bright, C., Neville, L.E., Howard, G.W., Murphy S.J. and Preston G. (eds.) 2003. *Invasive alien species in Southern Africa: national reports & directory of resources*. Global Invasive Species Programme, Cape Town, South Africa.

- MacDougall, W. B., and Glasgow, O. E. 1929. Mycorrhizas of the Compositae. *Am. J. Bot.* **16**: 225-228.
- MacKay, J. and Kotanen, P. M. 2008. Local escape of an invasive plant, common ragweed (*Ambrosia artemisiifolia* L.), from above-ground and below-ground enemies in its native area. *J. Ecol.* **96**: 1152-1161.
- Madanes, N., Fischer, S., and Vicari, R. 2007. Fire effects on a *Spartina densiflora* salt marsh in the floodplain of the Paraná River, Argentina. *Rev. Chil. Hist. Nat.* **80**: 187-199.
- Mahmoudi M. 2016. Prevalence of pollens in the United States and elsewhere. **In:** *Allergy and Asthma*. 2nd ed. pp. 53-62. Massoud Mahmoudi Editor. Springer International Publishing, San Francisco.
- Maire, R. 1928. Contribution à l'étude de la flore de l'Afrique du Nord. *Bull. Soc. Hist. Nat. Afrique N.* **19**: 29-66
- Makra, L., Matyasovszky, I. and Deák, Á. J. 2014. Ragweed in Eastern Europe. In: *CAB International 2014. Invasive Species and Global Climate Change*. p.p. 117-128. Ziska, L.H. and Dukes J.S., Eds. Wallingford.
- Makra, L., Matyasovszky, I., Hufnagel, L. and Tusnády, G. 2015. The history of ragweed in the world. *Appl. Ecol. Environ. Res.*, **13**: 489–512.
- Makra, L., Matyasovszky, I., Tusnády, G., Wang, Y., Csépe, Z., Bozóki, Z., Nyúl, L. G., Erostyák, J., Bodnár, K., Sümeghy, Z., Vogel, H., Pauling, A., Páldy, A., Magyar, D., Mányoki, G., Bergmann, K.C., Bonini, M., Šikoparija, B., Radišić, P., Gehrig, R., Kofol Seliger, A., Stjepanović, B., Rodinkova, V., Prikhodko, A., Maleeva, A., Severova, E., Ščevková, J., Ianovici, N., Peternel, R., Thibaudon, M. 2016. Biogeographical estimates of allergenic pollen transport over regional scales: Common ragweed and Szeged, Hungary as a test case. *Agric. For. Meteorol.* **221**: 94-110.
- Mamedov, N., Mehdiyeva, N. P. and Craker, L. E. 2015. Medicinal plants used in traditional medicine of the Caucasus and North America. *J. Med. Act. Plants* **4**: 42-66
- Mandrioli, P., Di Cecco, M. and Andina, G. 1998. Ragweed pollen: The aeroallergen is spreading in Italy. *Aerobiologia*, **14**: 13–20.
- Marco, L. N. and Pirovani, M. 2009. Relevamiento de flora alergógena en Concepción del Uruguay. *Arch. Alerg. e Inmunol. Clin.* **40**: 44–50.
- Marcomini, S. C. and López, R. A. 2013. Erosion and management in coastal dunes. **In:** *Coastal Hazards*. pp. 511-553. Springer Netherlands.
- Marcomini, S., López, R., Picca, P., Madanes, N. and Bertolín, L. 2016. Natural Coastal Dune-Field Landforms, Plant Communities, and Human Intervention along Buenos Aires Northern Aeolian Barrier. *J. Coast. Res.* In-Press.
- Marten, G. C. and Andersen, R. N. 1975. Forage nutritive value and palatability of 12 common annual weeds. *Crop Sci.* **15**: 821-827.
- Martinson, E. J., Zachary B. Eddy, Z. B, Julie L. Commerford, J. L. , Emilie Blevins E., Susan J. Rolfsmeier, S. J. and McLauchlan, K. K. 2011. Biogeographic Distributions of Selected North American Grassland Plant Species. *Phys. Geogr.* **32**: 583-602
- Mcandrews, J. H. (1988). Human disturbance of North American forests and grasslands: the fossil pollen record. **In:** Huntley, B. B. and Webb III, T. Eds., *Vegetation history*. pp. 673–697. Springer Netherlands.
- McNicoll, M. B. and Augspurger, C. K. 2010. A comparison of vegetation and seed bank community structure in a sand prairie in Illinois, USA. *Am. Midl. Nat.* **164**: 136-150.
- Medvecká, J., Kliment, J., Majeková, J., Halada, L., Zaliberova, M., Gojdičová, E., Feráková, V. and Jarolímeckl, I. 2012. Inventory of the alien flora of Slovakia. *Preslia* **84**: 257-309.
- Menghi, M., Cabido, M., Acosta, A., Peco, B. and Pineda, F. D. 1993. Changes in pasture communities subject to burning in the Córdoba Mountains, Argentina. *Coenoses*, **8**: 1-10.
- Menges, E. S. and Waller, D. M. 1983. Plant strategies in relation to elevation and light in floodplain herbs. *Am. Nat.* **122**: 454-473.

- Mihajlovic, L., Radosavljevic, J., Burazer, L., Smiljanic, K. and Cirkovic Velickovic, T. 2015. Phytochemistry composition of polyphenol and polyamide compounds in common ragweed (*Ambrosia artemisiifolia* L.) pollen and sub-pollen particles. *Phytochemistry*, **109**: 125–132.
- Milakovic, I., Fiedler, K. and Karrer, G. 2014. Management of roadside populations of invasive *Ambrosia artemisiifolia* by mowing. *Weed Res.* **54**: 256–264.
- Milakovic, I. and Karrer, G. 2016. The influence of mowing regime on the soil seed bank of the invasive plant *Ambrosia artemisiifolia* L. *NeoBiota*, **28**: 39-49.
- Mitchell, J. C., Roy, A. K., Dupuis, G. and Towers, G. N. (1971). Allergic contact dermatitis from ragweeds (*Ambrosia* species): the role of sesquiterpene lactones. *Arch. Dermatol.* **104**: 73-76.
- Mitich, L. W. 1996. Ragweeds (*Ambrosia* spp.): The Hay Fever Weeds. *Weed Technol.* **10**: 236–240.
- Mito, T., and Uesugi, T. 2004. Invasive alien species in Japan: the status quo and the new regulation for prevention of their adverse effects. *Global Environmental Research*, **8**: 171-193.
- Miziniak, W. and Praczyk, T. 2002. Regrowth of *Ambrosia psilostachya* D.C. from rhizomes on different type of soils. *Prog. Plant Prot.* **42**: 547–550.
- Molinaro, F., Monterumici, C. M., Ferrero, A., Tabasso, S. and Negre, M. 2016. Bioherbicidal activity of a germacranolide sesquiterpene dilactone from *Ambrosia artemisiifolia* L. *J. Environ. Sci. Health B* **51**: 847-852.
- Mondin, C.A. and Nakajima, J. N. 2015. *Ambrosia*. In: *Lista de Espécies da Flora do Brasil*. Jardim Botânico do Rio de Janeiro. Disponível em: <<http://reflora.jbrj.gov.br/jabot/floradobrasil/FB103255>>
- Mongelli, E., Martino, V., Coussio, J. and Ciccia, G. 1996. Screening of Argentine medicinal plants using the brine shrimp microwell cytotoxicity assay. *Int. J. Pharmacogn.* **34**: 249–254.
- Montagnani, C., Gentili, R., Citterio, S., Fenu, G., Nicolella, G., Karrer, G. 2017 *Ambrosia maritima* L. In: Orsenigo, S., Astuti, G., Bartolucci, F., Citterio, S., Conti, F., Garrido-Becerra, J. A., Gentili, R., del Galdo, G. G., Jiménez-Martínez, J.F., Karrer, G., Lahora, A., Martínez-Hernández, F., Mendoza-Fernández, A. J., Merlo, M. E., Montagnani, C., Mota, J., Nicolella, G., Pérez-García, F. J., Peruzzi, L., Robles, J., Roma-Marzio, F., Salmerón-Sánchez, E., Sánchez-Gómez, P., Serra, L., Stinca, A., Fenu, G. 2017. Global and Regional IUCN Red List Assessments: 3. *Italian Botanist* **3**: 83-98.
- Moskalenko G.P. 2001. *Quarantine Weeds for Russia*. Moscow, Russia: Plant Quarantine Inspectorate.
- Neill, R. L. and Rice, E. L. 1971. Possible role of *Ambrosia psilostachya* on pattern and succession in old-fields. *Am. Midl. Nat.* **86**: 344–357.
- Nelson, J. C. 1917. The introduction of foreign weeds in ballast as illustrated by ballast- plants at Linnton, Oregon. *Torreyia*, **17**: 151–160.
- NetPhyD: Deutschlandflora WebGIS. Floristische Verbreitungskarten in Deutschland. Accessed in January 2017.
- Novara, L.J. and Gutierrez, D.G. 2010. Asteraceae- Tribu Heliantheae. In: Novara, L.J. (Dir), *Flora del Valle de Lerma*. Aportes Bot. Salta, sér. Flora 9: 1-201.
- Nuutinen, V. and Butt, K. R. 2005. Homing ability widens the sphere of influence of the earthworm *Lumbricus terrestris* L. *Soil Biol. Biochem.* **37**: 805–807.
- Odé, B. and Beringen, R. 2017a. Alsemamambrosia *Ambrosia artemisiifolia*. Nederlands Soortenregister, [www.nederlandsesoorten.nl](http://www.nederlandsesoorten.nl). Geraadpleegd op (Accessed 24.01.2017).
- Odé, and Beringen, R. 2017b. Zandambrosia *Ambrosia psilostachya*. Nederlands Soortenregister, [www.nederlandsesoorten.nl](http://www.nederlandsesoorten.nl). Geraadpleegd op (Accessed 24.01.2017).

- Odé, B. and Beringen, R. 2017c. Driedelige ambrosia *Ambrosia trifida*. Nederlands Soortenregister, [www.nederlandsesoorten.nl](http://www.nederlandsesoorten.nl). Geraadpleegd op (Accessed 24.01.2017).
- Onen H., Gunal H. and Ozcan S. 2014. The Black Sea highway: The Route of Common Ragweed (*Ambrosia artemisiifolia* L.) Invasion in Turkey. **In:** Uludag et al. (eds.) *Proceedings of 8th International Conference on Biological Invasions from understanding to action*. Antalya, Turkey.
- Ortmans, W., Mahy, G. and Monty, A. 2016. Effects of seed traits variation on seedling performance of the invasive weed, *Ambrosia artemisiifolia* L. *Acta Oecologica*, **71**: 39-46.
- Osawa, T., Mitsuhashi, H. and Niwa, H. 2013. Many alien invasive plants disperse against the direction of stream flow in riparian areas. *Ecol. Complex.* **15**: 26–32.
- Otto, C. 2006. *Ambrosia artemisiifolia*. Bundesamt für Naturschutz - Neobiota.de: <http://neobiota.bfn.de/12655.html> (access: 1.02.2017);
- Özhatay, N. and Kültür, Ş. 2006. Check-list of additional *taxa* to the Supplement Flora of Turkey III. *Turk. J. Botany* **30**: 281-316.
- Pacific Island Ecosystems at Risk (PIER), 2013a. Species list – *Ambrosia artemisiifolia* L. – Asteraceae: [http://www.hear.org/pier/species/ambrosia\\_artemisiifolia.htm](http://www.hear.org/pier/species/ambrosia_artemisiifolia.htm) (access 19.01.2017)
- Pacific Island Ecosystems at Risk (PIER), 2013b. species list – *Ambrosia psilostachya* DC. – Asteraceae: [http://www.hear.org/pier/species/ambrosia\\_psilostachya.htm](http://www.hear.org/pier/species/ambrosia_psilostachya.htm) (access 19.01.2017)
- Paquin, V. and Aarssen, L. W. 2004. Allometric gender allocation in *Ambrosia artemisiifolia* (Asteraceae). *J. Bot.* **91**: 430–438.
- Parsons W.T & Cuthbertson E.G. 2001. *Noxious weeds of Australia*. 2nd ed. CSIRO, Collingwood.
- Patracchini, C., Vidotto, F. and Ferrero, A. 2011. Common ragweed (*Ambrosia artemisiifolia*) growth as affected by plant density and clipping. *Weed Technol.* **25**: 268-276.
- Payne, W. W. 1970. Preliminary reports on the flora of Wisconsin. No. 62 Compositae VI. (Composite Family VI). The genus *Ambrosia*—the ragweeds. *Trans. Wisconsin Acad. Sci. Arts, Lett.* **58**: 353–371.
- Pazmandi, K., Kumar, B. V, Szabo, K., Boldogh, I., Szoor, A., Vereb, G., Veres, A., Lanyi, A., Rajnavolgyi, E. and Bacsi, A. 2012. Ragweed subpollen particles of respirable size activate human dendritic cells. *PloS One*, **7**: e52085. doi:10.1371/journal.pone.0052085.
- Pendleton, R. L. and Smith, B. N. 1983. Vesicular-arbuscular mycorrhizae of weedy and colonizer plant species at disturbed sites in Utah. *Oecologia*, **59**: 296-301.
- Peng, C.I. 2013. Digital Flora of Taiwan. [http://www.efloras.org/florataxon.aspx?flora\\_id=100&taxon\\_id=200023072](http://www.efloras.org/florataxon.aspx?flora_id=100&taxon_id=200023072) (access 27.01.2017).
- Petitpierre, B., Kueffer, C., Broennimann, O., Randin, C., Daehler, C. and Guisan, A. 2012. Climatic niche shifts are rare among terrestrial plant invaders. *Science* **335**: 1344-1348.
- Pinke, G., Karácsony, P., Czúcz, B. and Botta-Dukát, Z. 2011. Environmental and land-use variables determining the abundance of *Ambrosia artemisiifolia* in arable fields in Hungary. *Preslia* **83**: 219-235.
- Plank, L., Zak, D., Getzner, M., Follak, S., Essl, F., Dullinger, S., Kleinbauere, I., Moserc, D., Gattringer, A. 2016. Benefits and costs of controlling three allergenic alien species under climate change and dispersal scenarios in Central Europe. *Environ. Sci. Policy* **56**: 9–21.
- Pleasant, J. M. and Schlather, K. J. 1994. Incidence of weed seed in cow (*Bos* sp.) manure and its importance as a weed source for cropland. *Weed Technol.* **8**: 304–310.
- Pruski, J. F. 2017. 1. *Ambrosia* L. **In:** G. Davidse, M. Sousa Sánchez, S. Knapp & F. Chiang Cabrera, Eds., *Flora Mesoamericana, Volume 5 (2), Asteraceae*. pp. 632-636. Missouri Botanical Garden, St. Louis.



- Puc, M. 2004. Ragweed pollen in the air of Szczecin. *Ann. Agric. Environ. Med.* **11**: 53-57.
- Pyšek, P. 2005. Survival rates in the Czech Republic of introduced plants known as wool aliens. *Biol. Invasions* **7**: 567–576
- Pyšek, P., Danihelka, J., Sádlo, J., Chrtek, J., Chytrý, M., Jarošík, V., Kaplan, Z., Krahulec, F., Moravcová, L., Pergl, J., Štajerová, K. and Tichý, L. 2012. Catalogue of alien plants of the Czech Republic: checklist update, taxonomic diversity and invasion patterns. *Preslia* **84**: 155–255.
- Quéney, A. 1942. *Ambrosia psilostachya* D.C. : espèce américaine nouvelle dans la banlieue de Lyon; ses rapports avec *Ambrosia artemisiaefolia* L.. *Bull. Mens. Soc. Linn.* **11**: 66-70.
- Quézel, P. and Santa, S. 1963. *Nouvelle flore de l'Algérie et des régions désertiques méridionales*. Tome 2. Paris.
- Qin, Z., DiTommaso, A., Wu, R. S. and Huang, H. Y. 2014. Potential distribution of two *Ambrosia* species in China under projected climate change. *Weed Res.* **54**: 520–531.
- Ramachandra Prasad, T. V, Rao, R. R., Sanjay, M. T. and Sharma, R. A. 2013. *Ambrosia psilostachya* DC (Asteraceae) - a new record but a potential threat to Indian flora. *Current Science*, **104**: 294–296.
- Rasmussen, K., Thyrring, J., Muscarella, R. and Borchsenius, F. 2017. Climate-change-induced range shifts of three allergenic ragweeds (*Ambrosia* L.) in Europe and their potential impact on human health. *PeerJ*, **5**: e3104 <https://doi.org/10.7717/peerj.3104>.
- Randall, R.P. 2012. *A Global Compendium of Weeds*. 2nd eds. Department of Agriculture and Food, Western Australia.
- Reece, P. E., Brummer, J. E., Northup, B. K., Koehler, A. E. and Moser, L. 2004. Interactions among Western ragweed and other sandhills species after drought. *J. Range Manag.* **57**: 583–589.
- Regnier, E., Harrison, S. K., Liu, J., Schmoll, J. T., Edwards, C. A., Arancon, N. and Holloman, C. 2008. Impact of an exotic earthworm on seed dispersal of an indigenous US weed. *J. Appl. Ecol.* **45**: 1621–1629. <http://doi.org/10.1111/j.1365-2664.2008.01489.x>
- Regnier, E. E., Harrison, S. K., Loux, M. M., Holloman, C., Venkatesh, R., Diekmann, F., Taylor R., Ford, R. A., Stoltenberg D. E., Hartzler, R. G., Davis A. S., Schutte B. J., Cadina, J., Mahoney K. J. and Johnson, W. G. 2016. Certified Crop Advisors' perceptions of giant ragweed (*Ambrosia trifida*) distribution, herbicide resistance, and management in the Corn Belt. *Weed Sci.* **64**: 361–377.
- Reynolds, S. C. P. 2002. *A catalogue of alien plants in Ireland*. National Botanic Gardens, Glasnevin, Dublin.
- Rice, E. L. 1965. Inhibition of nitrogen-fixing and nitrifying bacteria by seed plants IV. The inhibitors produced by *Ambrosia elatior* and *A. psilostachya*. *Southwest. Nat.* **10**: 248-255.
- Rich, T. C. G. 1994. Ragweeds (*Ambrosia* L.) in Britain. *Grana* **33**: 38–43.
- Richardson, D. M., and Pyšek, P. 2012. Naturalization of introduced plants: Ecological drivers of biogeographical patterns. *New Phytol.* **196**: 383–396.
- Richardson, D. M., Pyšek, P., Rejmánek, M., Barbour, M. G., Dane Panetta, F. and West, C. J. 2000. Naturalization and invasion of alien plants: Concepts and definitions. *Divers. Distrib.* **6**: 93–107.
- Ries, C. (eds.) 2017. *Ambrosia artemisiifolia* L. **In**: *neobiota.lu - Invasive Alien Species in Luxembourg*. URL: <http://neobiota.lu/ambrosia-artemisiifolia/> [23.01.2017].
- Rivera-Becerril, F., Juarez-Vazquez, L. V., Hernandez-Cervantes, S. C., Acevedo-Sandoval, O. A., Vela-Correa, G., Cruz-Chavez, E., Moreno-Espíndola, I. P., Esquivel-Herrera, A. and de León-González 2013. Impacts of manganese mining activity on the environment : interactions among soil, plants, and arbuscular mycorrhiza. *Arch. Environ. Contam. Toxicol.* **64**: 219–227.

- Robbins, R. R., Dickinson, D. B. and Rhodes, A. M. 1979. Morphometric analysis of pollen from four species of *Ambrosia* (Compositae). *Am. J. Bot.* **66**: 538–545.
- Rogers, C., Wayne, P.M., Macklin, E.A., Muilenberg, M.L., Wagner, C.J., Epstein, P.R. and Bazzaz, F. 2006. Interaction of the onset of spring and elevated atmospheric CO<sub>2</sub> on ragweed (*Ambrosia artemisiifolia* L.) pollen production. *Environ. Health persp.* **114**: 865–869.
- Roldán, F.P. and Vibrans, H., 2009. Malezas de México, Ficha - *Ambrosia psilostachya* DC. 24.01.2017. <http://www.conabio.gob.mx/malezasdemexico/asteraceae/ambrosia-psilostachya/fichas/ficha.htm#2.%20Origen%20y%20distribuci%C3%B3n%20geogr%C3%A1fica>
- Rosas, C. A., Engle, D. M., Shaw, J. H. and Palmer, M. W. 2008. Seed dispersal by *Bison bison* in a tallgrass prairie. *J. Veg. Sci.* **19**: 769–778.
- Sáenz, A. A. and Gutiérrez, D. G. 2008. Tribu Heliantheae (Asteraceae). In: Freire, S. E. and Molina, A. M. (eds.), *Flora Chaqueña -Argentina- (Formosa, Chaco y Santiago del Estero)*. Colecc. Ci. Inst. Nac. Tecnol. Agropecu. pp.: 422–533.
- Salzman, A. G. 1985. Habitat selection in a clonal plant. *Science*, **228**: 603–604.
- Salzman, A. G. and Parker, M. A. 1985. Neighbours ameliorate local salinity stress for a rhizomatous plant in a heterogeneous environment. *Oecologia*, **65**: 273–277.
- SANBI. 2015a. *Ambrosia psilostachya* DC. National Assessment: Red List of South African Plants version 2015.1. Accessed on 2017/01/24.
- SANBI. 2015b. *Ambrosia tenuifolia* Spreng. National Assessment: Red List of South African Plants version 2015.1. Accessed on 2017/01/24.
- Sang, W., Liu, X. and Axmacher, J.C. 2011. Germination and emergence of *Ambrosia artemisiifolia* L. under changing environmental conditions in China. *Plant Species Biol.* **26**: 125–133.
- Sauvalle Chanceaulme, F.A. 1873. *Flora Cubana. Enumeratio Nova Plantarum Havanae*. La Antilla, de Cacho-Negrete.
- Schutte, B. J., Regnier, E. E., Harrison, S. K., Schmoll, J. T., Spokas, K. and Forcella, F. 2008a. A hydrothermal seedling emergence model for giant ragweed (*Ambrosia trifida*). *Weed Sci.* **56**: 555–560.
- Schutte, B. J., Regnier, E. E. and Harrison, S. K. 2008b. The association between seed size and seed longevity among maternal families in *Ambrosia trifida* L. populations. *Seed Sci. Res.* **18**: 201–211.
- Schutte, B. J., Liu, J., Davis, A. S., Harrison, S. K. and Regnier, E. E. (2010). Environmental factors that influence the association of an earthworm (*Lumbricus terrestris* L.) and an annual weed (*Ambrosia trifida* L.) in no-till agricultural fields across the eastern U. S. Corn Belt. *Agric. Ecosyst. Environ.* **138**: 197–205.
- Schutte, B. J., Regnier, E. E. and Harrison, S. K. (2012). Seed Dormancy and Adaptive Seedling Emergence Timing in Giant Ragweed (*Ambrosia trifida*). *Weed Sci.* **60**: 19–26.
- Sell, P. and Murrell, G. 2006. *Flora of Great Britain and Ireland: Volume 4, Campanulaceae-Asteraceae*. Cambridge University Press.
- Semmartin, M., Di Bella, C. and Garcia de Salamone, I. 2010. Grazing-induced changes in plant species composition affect plant and soil properties of grassland mesocosms. *Plant Soil* **328**: 471–481.
- Sénéchal, H., Visez, N., Charpin, D., Shahali, Y., Peltre, G., Biolley, J., Lhuissier, F., Courdec, R., Yamada, O., Malrat-Domenge, A., Pham-Thi, N., Poncet, P. and Sutra, J. P. 2015. A review of the effects of major atmospheric pollutants on pollen grains, pollen content, and allergenicity. *Sci. World J.* **2015**: Article ID 940243.
- Setshogo, M.P. 2005. *Preliminary Checklist of the Plants of Botswana*. Southern African Botanical Diversity Network, Report No. 37. SABONET, Pretoria and Gaborone.
- Shaltout, K. H. 2004. An updated flora of Egypt. *Divers. Distrib.* **10**: 77–78.

- Shemluck, M. 1982. Medicinal and Other Uses of the Compositae by Indians. *J. Ethnopharmacol.* **5**: 303–358.
- Šikoparija, B., Galan, C., Smith, M. and Group, E. Q. W. 2016. Pollen-monitoring: between analyst proficiency testing. *Aerobiologia*. <http://doi.org/10.1007/s10453-016-9461-3>
- Šikoparija, B., Skjøth, C. A., Alm Kübler, K., Dahl, A., Sommer, J., Grewling, Ł., Radišić, P. and Smith, M. 2013. A mechanism for long distance transport of *Ambrosia* pollen from the Pannonian Plain. *Agric. For. Meteorol.* **180**: 112–117.
- Simon, M. L. 2009. *A Regional and Chronological Synthesis of Archaic Period Plant Use in the Midcontinent*. Emerson, T. E., McElrath, D. L. and Fortier A. C. Eds. Suny Press, Albany, New York.
- Sîrbu, C. 2012. *Plante de carantină invazive în România*. Universitatea de Științe Agricole și Medicină Veterinară Ion Ionescu de la Brad Iași.
- Skálová, H., Moravcová, L., Dixon, A. F., Kindlmann, P. and Pyšek, P. 2015. Effect of temperature and nutrients on the growth and development of seedlings of an invasive plant. *AoB Plants*, **7**: plv044 <http://doi.org/10.1093/aobpla/plv044>.
- Skarpe, C., du Toit, J.T. and Moe, S.R. (eds.) 2014. *Elephants and savanna woodland ecosystems : a study from Chobe National Park, Botswana*. Wiley Blackwell.
- Slatyer, R. A., Hirst, M., and Sexton, J. P. 2013. Niche breadth predicts geographical range size: a general ecological pattern. *Ecol. Lett.* **16**: 1104-1114.
- Smith, M., Skjøth, C. A., Myszkowska, D., Uruska, A., Puc, M., Stach, A., Balwierz, Z., Chlopek, K., Piotrowska, K., Kasprzyk, K., Brandt, J., 2008. Long-range transport of *Ambrosia* pollen to Poland. *Agric. For. Meteorol.* **148**: 1402-1411.
- Smith, M., Cecchi, L., Skjøth, C. A., Karrer, G. and Šikoparija, B. 2013. Common ragweed: a threat to environmental health in Europe. *Environ. Int.* **61**: 115–126.
- Song, U., Mun, S. and Lee, E. J. 2012. Responses of two invasive plants under various microclimate conditions in the Seoul metropolitan region. *Environ. Manage.* **49**: 1238–1246.
- Soriano, A. (1982). La germinacion y el establecimiento de las plantas como episodios ecologicos. **In:** *Jornada Académica (9 y 10 de diciembre, Universidad Nacional de Río Cuarto, Córdoba*. pp. 58–65.
- Stace, C. 2010. *New flora of the British Isles*. Cambridge University Press.
- Stephenson, G. 1999. *Vehicle impacts on the biota of sandy beaches and coastal dunes: a review from a New Zealand perspective*. Departement of Conservation. Wellington, N.Z.
- Stešević, D., and Petrović, D. 2010. Preliminary list of plant invaders in Montenegro. *Biologica Nyssana*, **1**: 35-42.
- Stoyanov, S., Vladimirov, V. and Milanova, S. 2014. *Ambrosia trifida* (Asteraceae), a new non-native species for the Bulgarian flora. *C. R. Acad. Bulg. Sci.* **67**: 1653–1656.
- Stromberg, J. C. 2013. Root patterns and hydrogeomorphic niches of riparian plants in the American Southwest. *J. Arid Environ.* **94**: 1-9.
- Strother J.L. 2006. *Flora of North America, North of Mexico*, Vol. 21. Magnoliophyta: Asteridae, part 8: Asteraceae, part 3:10–18. Oxford University Press. New York.
- Stubbendieck, J. and Tunnell, S. J. 2008. Seventy-eight years of vegetation dynamics in a Sandhills Grassland. *Nat. Area. J.* **28**: 58-65.
- Sülsen, V. P., Frank, F. M., Cazorla, S. I., Anesini, C. A., Malchiodi, E. L., Freixa, B., Vila, R., Muschiatti, L. V. and Martino V. S. 2008. Trypanocidal and leishmanicidal activities of sesquiterpene lactones from *Ambrosia tenuifolia* Sprengel (Asteraceae). *Antimicrob. Agents Chemother.* **52**: 2415-2419.
- Sülsen V.P., Cazorla S.I., Frank F.M., Laurella L.C., Muschiatti L.V., Catalán, C. A., Martino, V. S. and Malchiodi, E. L. 2013. Natural Terpenoids from *Ambrosia* species are active in vitro and in vivo against human pathogenic trypanosomatids. *PLoS Negl. Trop. Dis.* **7**: e2494.

- Szigetvári, C. and Benkő, Z. 2008. Common ragweed (*Ambrosia elatior* L.). **In:** Botta-Dukát Z. & Balogh L. (eds.) *The most important invasive plants in Hungary*. pp.189–201, Institute of Ecology and Botany, Hungarian Academy of Sciences, Vácrátót.
- Tamanyan, K. and Fayvush, G. 2010. Invasive alien plants in Armenia. **In** Brunel, S., Uludag, A.U., Fernandez-Galiano, E. and Brundu, G. (Eds.) *Proceedings of the 2nd International Workshop on Invasive Plants in the Mediterranean Type Regions of the World*. Trabzon, Turkey.
- Tamarcaz, P., Lambelet, C., Clot, B., Keimer, C. and Hauser, C. 2005. Ragweed (*Ambrosia*) progression and its health risks: Will Switzerland resist this invasion? *Swiss Med Wkly* **135**: 538–548.
- Tanji, A. 2005. *Adventices du blé et de l'orge au Maroc*. Editions INRA Maroc, Rabat.
- Tejera, L. and Beri, A. 2005. First volumetric airborne pollen sampling in Montevideo City, Uruguay. *Aerobiologia*, **21**: 33–41.
- Thellung, A. 1912. La flore adventice de Montpellier. *M. Soc. natl. sci. nat. math. Cherb.* **38**: 57–728.
- Tokarska-Guzik, B., Bzdęga, K., Koszela, K., Żabińska, I., Krzuś, B., Sajan, M. and Sendek, A. 2011. Allergenic invasive plant *Ambrosia artemisiifolia* L. in Poland: threat and selected aspects of biology. *Biodivers. Res. Conserv.* **21**: 39–48.
- Thompson K. 2000. The functional ecology of soil seed banks. **In:** *Seeds: the ecology of regeneration in plant communities*. pp.: 215–235. Fenner M., Eds., Wallingford, UK: CAB International.
- Toole, H. E. and Brown, E. 1946. Final results of the Duvel buried seed experiment. *J. Agricult. Res.* **72**: 201–210
- Towne, E. G. 2000. Prairie vegetation and soil nutrient responses to ungulate carcasses. *Oecologia*, **122**: 232-239.
- Trillo, C., Colantonio, S. and Galetto, L. 2014. Perceptions and use of native forests in the arid Chaco of Córdoba, Argentina. *Ethnobot. Res. Appl.* **12**: 497–510.
- Tseng, Y.-H. and Peng, C.-I. 2004. *Ambrosia psilostachya* DC. (Asteraceae), a newly naturalized plant in Taiwan. *Endangered Sp. Res.* **6**: 71-74.
- Tye, A. 2001. *Invasive Plant Problems and Requirements for Weed Risk Assessment in the Galapagos Islands*. Proceedings of the Weed Risk Assessment Workshop. CSIRO Publishing;
- Ugarte, E., Lira, F., Fuentes, N. and Klotz, S. 2011. Vascular alien flora. *Chile Check List* **7**: 365-382.
- Uresk, D. W. 2012. Monitoring standing herbage of the sands and choppy sands ecological vegetation types in the Nebraska Sandhills. *Great Plains Research* **22**: 181-186.
- USDA, NRCS. 2017. The PLANTS Database (<https://plants.usda.gov/core/profile?symbol=AMTE5>) National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- Ujvárosi, M. 1973. *Gyomnövények* [Weeds]. Mezőgazdasági Kiadó, Budapest.
- Van Denderen, P. D., Tamis, W. L. M. and Van Valken, J. L. C. H. 2010. Risico's van introductie van exotische plantensoorten, in het bijzonder uit het geslacht *Ambrosia* L., via import van zaden voor met name veevoer en vogelvoer. *Gorteria* **34**: 65–85.
- Van Kleunen, M., Dawson, W. and Maurel, N. 2015. Characteristics of successful alien plants. *Mol. Ecol.* **24**: 1954–1968.
- Vaz Ferreira R. 1946. Los agentes de la polinosis. **In:** *Alergia en la practica clinica*. pp. 314–348. Varela, B., Recarte, R. and Graña, A. Eds., Espasa-Calpe-Argentina-S.A, Buenos Aires.
- Verloove F. 2016a. *Ambrosia trifida*. **In:** *Manual of the Alien Plants of Belgium*. Botanic Garden of Meise, Belgium. At: alienplantsbelgium.be, accessed 20/11/2016.

- Verloove F. 2016b. *Ambrosia psilostachya*. In: *Manual of the Alien Plants of Belgium*. Botanic Garden of Meise, Belgium. At: alienplantsbelgium.be, accessed 20/11/2016.
- Verloove F., 2016c. *Ambrosia artemisiifolia*. In: *Manual of the Alien Plants of Belgium*. Botanic Garden of Meise, Belgium. At: alienplantsbelgium.be, accessed 20/11/2016.
- Vermeire, L. T. and Gillen, R. L. 2000. Western Ragweed Effects on Herbaceous Standing Crop in Great Plains Grasslands. *J. Range Manag.* **53**: 335–341.
- Vermeire, L. T., Gillen, R. L. and Bidwell, T. G. 2005. *Ecology and management of Western ragweed on rangeland*. Division of Agricultural Sciences and Natural Resources, Oklahoma State University.
- Vibrans, H. 1998. Urban weeds of México City. Floristic composition and important families. *Anales Inst. Biol. Univ. Nac. Auton. Mexico, Bot.* **69**: 37-69
- Vidotto, F., Tesio, F. and Ferrero, A. 2013. Allelopathic effects of *Ambrosia artemisiifolia* L. in the invasive process. *Crop Prot.* **54**: 161-167.
- Vignolo Lutati, F. 1935. Il genere "*Ambrosia*" in Italia. *Nuovo Giornale Botanico Italiano*, **42**: 364–378.
- Villaseñor, J. L. and Espinosa-Garcia, F. J. 2004. The alien flowering plants of Mexico. *Divers. Distrib.* **10**: 113-123.
- Vinogradova, Y.K., Maiorov, S.R. and Khorun, L.V. 2010. *Black book of the flora of Central Russia: alien plant species in Central Russian ecosystems*. GEOS Moscow (RU). EPPO Reporting Service - NO. 2, PARIS, 2011-02-01 2011/045.
- Vitalos, M. and Karrer, G. 2008. Distribution of *Ambrosia artemisiifolia* L. - Is birdseed a relevant vector? *J. Plant Dis. Protection Spec. Issue* **21**: 345–348.
- Vitalos, M. and Karrer, G. 2009. Dispersal of *Ambrosia artemisiifolia* seeds along roads: the contribution of traffic and mowing machines. *Neobiota* **8**: 53-60
- Volta P., Jeppesen E., Leoni B., Campi B., Sala P., Garibaldi L., Lauridsen T.L., Winfield I.J., 2013. Recent invasion by a non-native cyprinid (*Abramis brama*) is followed by major changes in the ecological quality of a shallow lake in southern Europe. *Biol. Invasions* **15**: 2065-2079.
- Von der Lippe, M., Bullock, J. M., Kowarik, I., Knopp, T. and Wichmann, M. 2013. Human-Mediated Dispersal of Seeds by the Airflow of Vehicles. *PLoS One* **8**: e52733.
- Von Raab-Straube, E. and Raus, T. 2016. Euro+Med-Checklist Notulæ, 6. *Willdenowia*, **46**: 423–442.
- Vrbničanin, S., Karadžić, B. and Dajić-Stevanović, Z. 2004. Advantivne i invazivne korovske vrste na području Srbije. *Acta Herbologica*, **13**: 1-12
- Wagner, W. H. 1958. The hybrid ragweed, *Ambrosia artemisiifolia* × *trifida*. *Rhodora*, **60**: 309-316.
- Wagner, W. H. and Beals, T. F. 1958. Perennial ragweeds (*Ambrosia*) in Michigan, with the description of a new, intermediate *taxon*. *Rhodora*, **60**:177–204.
- Wagner, W. L., Herbst, D. R., Sohmer, S. H. 1990. *Manual of the flowering plants of Hawaii*. Bishop Museum Press, Honolulu, Hawaii, USA.
- Waisel, Y., Eshel, A., Keynan, N. and Langgut, D. 2008. Ambrosia: A new impending disaster for the Israeli allergic population. *Isr. Med. Assoc. J.* **10**: 856–857.
- Waldburger, E. and Staub, R. 2006. Neophyten im Fürstentum Liechtenstein. *Berichte der Botanisch-Zoologischen Gesellschaft Liechtenstein-Sargans-Werdenberg Schaan*, **32**: 95-112.
- Wan, S., Yuan, T., Bowdish, S., Wallace, L., Russell, S. D. and Luo, Y. 2002. Response of an allergenic species, *Ambrosia psilostachya* (Asteraceae), to experimental warming and clipping: implications for public health. *Am. J. Bot.* **89**: 1843–1846.
- Wang, P., Liang W.J., Kong C.H. and Jiang Y. 2005. Allelopathic potentials of volatile allelochemicals from *Ambrosia trifida* L. on other plants. *Allelopathy J.*, **15**: 131–136.

- Washitani, I. 2004. Invasive alien species problems in Japan: an introductory ecological essay. *Glob. Environ. Res.* **8**: 1-12.
- Wasowicz, P., Przedpelska-Wasowicz, E. M. and Kristinsson, H. 2013. Alien vascular plants in Iceland: diversity, spatial patterns, temporal trends, and the impact of climate change. *Flora-Morphology, Distrib. Funct. Ecol. Plants* **208**: 648-673.
- Webb, C.J., Sykes, W.R., Garnock-Jones, P.J. 1988. *Flora of New Zealand*. Vol. IV. Naturalised Pteridophytes, Gymnosperms, Dicotyledons. Botany Division DSIR, Christchurch.
- Weber, R. W. 2007. Cross-reactivity of pollen allergens: impact on allergen immunotherapy. *Annals of Ann. Allergy, Asthma Immunol.* **99**: 203–212.
- Webster, T. M., Loux, M. M., Regnier, E. E. and Harrison, S. K. 1994. Giant ragweed (*Ambrosia trifida*) canopy architecture and interference studies in soybean (*Glycine max*). *Weed Technol.* **8**: 559-564.
- Weeda, E. J. 2010. The role of archaeophytes and neophytes in the Dutch coastal dunes. *J. Coast. Conserv.* **14**: 75–79.
- Wells, M.J., Balsinhas, V.M., Joffe, H., Engelbrecht, V.M., Harding, G. and Stirton, C.H. 1986. *A Catalogue of Problem Plants in Southern Africa, incorporating The National Weed List of South Africa*. Memoirs of the Botanical Survey of South Africa, No. 53. Botanical Research Institute, Pretoria, South Africa
- Wolfe, C. W. 1973. Effects of fire on a sand hills grassland environment. **In: Proceedings of the Tall Timbers Fire Ecology Conference; 8–9 June 1972.** pp. 241-255. Tall Timbers Research Station. Tallahassee, FL, USA.
- Wollenweber, E., Hradetzky, D., Mann, K., Roitman, J. N., Yatskievych, G., Proksch, M. and Proksch, P. 1987. Exudate flavonoids from aerial parts of five *Ambrosia* species. *J. Plant Physiol.* **131**:7-43.
- Wollenweber, E., Mann, K., Dörr, M., Fritz, H., Roitman, J. N. and Yatskievych, G. 1995. Exudate flavonoids in three *Ambrosia* species. *Nat. Prod. Lett.* **7**: 109-116.
- Wopfner, N., Gadermaier, G., Egger, M., Asero, R., Ebner, C., Jahn-Schmid, B. and Ferreira, F. 2005. The spectrum of allergens in ragweed and mugwort pollen. *Int. Arch. Allergy Immunol.* **138**: 337–346.
- Wortman, S. E., Davis, A. S., Schutte, B. J., Lindquist, J. L., Cardina, J., Felix, J., Sprague, C. L., Dille, J. A., Ramirez, A. H. M., Reicks, G. and Clay, S. A. 2012. Local conditions, not regional gradients, drive demographic variation of giant ragweed (*Ambrosia trifida*) and common sunflower (*Helianthus annuus*) across northern US maize belt. *Weed Sci.* **60**: 440-450.
- Wright, T. J. 1941. A study of the Fall food supply of the ring-necked pheasant and the bob-white quail in Washington county , Rhode Island. *J. Wildl. Manage.* **5**: 279–296.
- Wu, S. H., Aleck Yang, T. Y., Teng, Y. C., Chang, C. Y., Yang, K. C. and Hsieh, C. F. 2010. Insights of the latest naturalized flora of Taiwan: Change in the past eight years. *Taiwania* **55**: 139–159.
- Wu, S-H., Hsieh C.-F. and Rejmánek M. 2004. Catalogue of the naturalized flora of Taiwan. *Taiwania* **49**: 16-31.
- Yair, Y., Sibony, M. and Rubin, B. 2017. Four *Ambrosia* species in Israel: invasive, naturalized and casual alien plants. *Isr. J. Plant Sci.* DOI: 10.1080/07929978.2017.1288399
- Yavorska, O. 2009. The North American species of the non-native flora of the Kyiv urban area (Ukraine): a checklist and analysis. *Biodivers. Conserv.* **13**: 25-30.
- Yoshikawa, M., Hoshino, Y. and Iwata, N. 2013. Role of seed settleability and settling velocity in water for plant colonization of river gravel bars. *J. Veg. Sci.* **24**: 712–723.
- Zárate, R., Macedo, N. L., Gallardo, G. P., Flores, M., Martínez-Dávila, P., Ramírez, F. F. and Torres, L. A. 2015. Contribución al conocimiento de la composición florística del departamento de Huánuco, Perú. *Folia Amazónica* **24**: 91-100.

- Zelnik, I. 2012. The presence of invasive alien plant species in different habitats: case study from Slovenia. *Acta Biol Slov.* **55**: 25-38.
- Zemmer, F., Karaca, F. and Ozkaragoz, F. 2012. Ragweed pollen observed in Turkey: detection of sources using back trajectory models. *Sci. Total Environ.* **430**: 101-108.
- Zhao, F., El Kelish, A., Durner, J., Lindermayr, C., Winkler, J. B., Ruëff, F., Behrendt, H., Traidl-Hoffmann, C., Holzinger, A., Kofler, W., Braun, P., von Toerne, C., Hauck, S. M., Ernst, D. and Frank, U. 2016. Common ragweed (*Ambrosia artemisiifolia* L.): allergenicity and molecular characterization of pollen after plant exposure to elevated NO<sub>2</sub>. *Plant Cell Environ.* **39**: 147–164.
- Ziska, L.H. and Caulfield, F.A. 2000. Rising CO<sub>2</sub> and pollen production of common ragweed (*Ambrosia artemisiifolia*), a known allergy-inducing species: implications for public health. *Australian J. Plant Physiol.* **27**: 893-898.
- Ziska, L., Knowlton, K., Rogers, C., Dalan, D., Tierney, N., Elder, M. A., Filley, W., Shropshire, J., Ford, L. B., Hedberg, C., Fleetwood, P., Hovanky, K. T., Kavanaugh, T., Fulford, G., Vrtis, R. F., Patz, J. A., Portnoy, J., Coates, F., Bielory, L., and Frenz, D. 2011. Recent warming by latitude associated with increased length of ragweed pollen season in central North America. *Proc. Natl. Acad. Sci. U. S. A.* **108**: 4248–51.
- Zuloaga, F. O., Morrone, O. and Belgrano, M. J. 2008. *Catalogue of the vascular plants of the Southern Cone (Argentina, southern Brazil, Chile, Paraguay and Uruguay)*. Volume 2: Dicotyledoneae: Acanthaceae-Fabaceae (Abarema-Schizolobium). Missouri Botanical Garden.

## Tables

Table 1 – *Ambrosia* species characteristics: morphological data allowing the identification of the four ragweeds.

| Species         | <i>Ambrosia artemisiifolia</i> L.   | <i>Ambrosia trifida</i> L.  | <i>Ambrosia psilostachya</i> DC.   | <i>Ambrosia tenuifolia</i> Spreng.  |
|-----------------|---|---|--|---|
| Life form       | Annual  | Annual  | Perennial  | Perennial   |
| Plant size (cm) | 10 to 250   | 40 to 400   | 10 to 90   | 20-100  |
| Belowground     | Taproot   | Taproot   | Root sprouter  | Root sprouter   |
| Stem            | +/- intensively branched, branches with wide angles   | +/- intensively branched  | Few branches, with narrow angles   | Few branches, with narrow angles  |
| Leaves          | Pinnatifid to bipinnate, rarely entire; leaf segments broadened and separated, rarely narrow; lower leaves with distinct narrow petiole; upper leaves alternate; long and short hairs mixed | Palmate, 1 to 5 lobes; glabrous or few short hairs; all leaves opposite   | Pinnatifid, rarely entire; leaf segments lineal and connected, often sharpened towards the tip; +/- sessile; upper leaves alternate; dense short hairs | Bipinnate to pinnatifid; leaf segments as narrow as the rachis, lineal, connected; lower leaves with distinct narrow petiole; upper leaves alternate; dense short hairs |
| Diaspore coat   | Few hairs and glands; 2-5 short lateral spines with sharpened tips; dark brown  | Glabrous or few hairs; 2-4 indistinct lateral spines; dark brown to black | Few glands and short hairs; blunt, short lateral spines few or none; dark brown  | Short hairs and glands, 2-5 lateral short blunt spines; olive to dark brown   |



Table 2 - Geography of *Ambrosia* (ragweed) species: distribution, time of arrival (first record in the wild), current status of ragweeds and references supporting reported data.

| Continent   | Country            | Species                           | First record | Status               | References  |
|-------------|--------------------|-----------------------------------|--------------|----------------------|---|
| Europe      | Albany             | <i>Ambrosia artemisiifolia</i> L. | /            | Doubtful occurrence  | Barina <i>et al.</i> , 2013; Barina <i>et al.</i> , 2014                            |
| Africa      | Algeria            | <i>A. artemisiifolia</i>          | > 1890       | Alien Casual         | Lawalree, 1947; Quézel and Santa, 1963; Greuter, 2006                               |
| Africa      | Algeria            | <i>A. psilostachya</i> DC.        | 1916         | Alien Naturalized    | Maire, 1928; Quézel and Santa, 1963; Greuter, 2006                                  |
| America (S) | Argentina          | <i>A. artemisiifolia</i>          | /            | Species occurring    | Freire <i>et al.</i> , 2008; Gerber <i>et al.</i> , 2011; Essl <i>et al.</i> , 2015 |
| America (S) | Argentina          | <i>A. tenuifolia</i> Spreng.      | /            | Native               | Freire <i>et al.</i> , 2008; Novara and Gutiérrez, 2010                             |
| Asia        | Armenia            | <i>A. artemisiifolia</i>          | /            | Alien Naturalized    | Tamanyan and Fayvush, 2010; Randall, 2012   |
| Oceania     | Australia          | <i>A. artemisiifolia</i>          | 1908         | Alien Invasive       | Parsons and Cuthbertson , 2001; Essl, <i>et al.</i> , 2015                          |
| Oceania     | Australia          | <i>A. psilostachya</i>            | 1922         | Alien Invasive       | Parsons and Cuthbertson , 2001  |
| Oceania     | Australia          | <i>A. tenuifolia</i>              | 1932         | Alien Naturalized    | Parsons and Cuthbertson , 2001  |
| Europe      | Austria            | <i>A. artemisiifolia</i>          | 1883         | Alien Invasive       | Essl <i>et al.</i> , 2009; Smith <i>et al.</i> , 2013                               |
| Europe      | Austria            | <i>A. trifida</i> L.              | 1948         | Alien Casual         | Essl and Rabitsch, 2002; Follak <i>et al.</i> , 2013                                |
| Europe      | Austria            | <i>A. psilostachya</i>            | /            | Alien Casual         | Essl and Rabitsch, 2002   |
| Asia        | Azerbaijan         | <i>A. artemisiifolia</i>          | /            | Alien Invasive       | Greuter, 2006; Gerber <i>et al.</i> , 2011  |
| America (C) | Bahamas            | <i>A. artemisiifolia</i>          | /            | Species occurring    | Acevedo-Rodriguez and Strong, 2012; Essl <i>et al.</i> , 2015                       |
| Europe      | Belarus            | <i>A. artemisiifolia</i>          | /            | Alien Naturalized    | Greuter, 2006   |
| Europe      | Belarus            | <i>A. psilostachya</i>            | /            | Alien status unknown | EPPO, 2016  |
| Europe      | Belarus            | <i>A. trifida</i>                 | /            | Alien Casual         | EPPO, 2016  |
| Europe      | Belgium            | <i>A. trifida</i>                 | 1829         | Alien Casual         | Verloove, 2016a   |
| Europe      | Belgium            | <i>A. psilostachya</i>            | 1917         | Alien Naturalized    | Verloove, 2016b   |
| Europe      | Belgium            | <i>A. artemisiifolia</i>          | 1883         | Alien Naturalized?   | Bullock <i>et al.</i> , 2012; Verloove, 2016c                                       |
| America (S) | Bolivia            | <i>A. artemisiifolia</i>          | /            | Alien status unknown | Jørgensen <i>et al.</i> , 2014  |
| America (S) | Bolivia            | <i>A. tenuifolia</i>              | /            | Native               | Jørgensen <i>et al.</i> , 2014  |
| Europe      | Bosnia Herzegovina | <i>A. artemisiifolia</i>          | /            | Alien status unknown | Kazinczi <i>et al.</i> , 2008; Smith <i>et al.</i> , 2008                           |
| Africa      | Botswana           | <i>A. artemisiifolia</i>          | /            | Alien Naturalized    | Setshogo, 2005; Randall, 2012; Skarpe <i>et al.</i> , 2014                          |

|                    |                     |                          |        |                      |   |
|--------------------|---------------------|--------------------------|--------|----------------------|---|
| <b>America (S)</b> | Brazil              | <i>A. artemisiifolia</i> | /      | Species occurring    | Mondin and Nakajima, 2015; Essl <i>et al.</i> , 2015; Alves and Rocha, 2016               |
| <b>America (S)</b> | Brazil              | <i>A. tenuifolia</i>     | /      | Native               | Sáenz and Gutiérrez, 2008   |
| <b>Europe</b>      | Bulgaria            | <i>A. artemisiifolia</i> | 1975   | Alien Naturalized    | Kazinczi <i>et al.</i> , 2008; Bullock <i>et al.</i> , 2012                               |
| <b>Europe</b>      | Bulgaria            | <i>A. trifida</i>        | 2014   | Alien status unknown | Stoyanov <i>et al.</i> , 2014   |
| <b>America (S)</b> | Canada              | <i>A. artemisiifolia</i> | /      | Native               | Bassett and Crompton, 1975; Kazinczi <i>et al.</i> , 2008                                 |
| <b>America (N)</b> | Canada              | <i>A. psilostachya</i>   | /      | Native               | Bassett and Crompton, 1975  |
| <b>America (N)</b> | Canada              | <i>A. trifida</i>        | /      | Native               | Bassett and Crompton, 1982  |
| <b>America (S)</b> | Chile               | <i>A. artemisiifolia</i> | 1959   | Alien Naturalized    | Essl <i>et al.</i> , 2015; Ugarte <i>et al.</i> , 2011; Fuentes <i>et al.</i> , 2013      |
| <b>America (S)</b> | Chile               | <i>A. tenuifolia</i>     | 1923   | Alien Naturalized?   | Ugarte <i>et al.</i> , 2011   |
| <b>Asia</b>        | China               | <i>A. artemisiifolia</i> | 1930s  | Alien Invasive       | Qin et a., 2014   |
| <b>Asia</b>        | China               | <i>A. trifida</i>        | 1935   | Alien Invasive       | Qin et a., 2014   |
| <b>Asia</b>        | China               | <i>A. psilostachya</i>   | /      | Alien status unknown | Chen and Hind 2011  |
| <b>America (S)</b> | Colombia            | <i>A. artemisiifolia</i> | /      | Species occurring    | Gerber <i>et al.</i> , 2011; CABI, 2017   |
| <b>Europe</b>      | Croatia             | <i>A. artemisiifolia</i> | 1879   | Alien Invasive       | Galzina <i>et al.</i> , 2010; Csontos <i>et al.</i> , 2010; Kazinczi <i>et al.</i> , 2008 |
| <b>America (C)</b> | Cuba                | <i>A. artemisiifolia</i> | < 1873 | Species occurring    | Sauvalle Chanceaulme, 1873; Acevedo-Rodriguez and Strong, 2012                            |
| <b>Europe</b>      | Czech Republic      | <i>A. artemisiifolia</i> | 1883   | Alien Invasive       | Kazinczi <i>et al.</i> , 2008; Smith <i>et al.</i> , 2008; Bullock <i>et al.</i> , 2012   |
| <b>Europe</b>      | Czech Republic      | <i>A. trifida</i>        | 1960   | Alien Casual         | Pyšek <i>et al.</i> , 2012  |
| <b>Europe</b>      | Czech Republic      | <i>A. psilostachya</i>   | 1999   | Alien Casual         | Pyšek <i>et al.</i> , 2012  |
| <b>Europe</b>      | Denmark             | <i>A. psilostachya</i>   | /      | Alien Casual         | Greuter, 2006   |
| <b>Europe</b>      | Denmark             | <i>A. trifida</i>        | /      | Alien Casual         | EPPO, 2016  |
| <b>Europe</b>      | Denmark             | <i>A. artemisiifolia</i> | 1865   | Alien Casual         | Bullock <i>et al.</i> , 2012  |
| <b>America (S)</b> | Ecuador             | <i>A. artemisiifolia</i> | /      | Species occurring    | Jørgensen and León-Yáñez, 1999  |
| <b>America (S)</b> | Ecuador (Galapagos) | <i>A. artemisiifolia</i> | /      | Alien Casual         | Tye, 2001; Jaramillo Díaz and Guézou, 2013.   |
| <b>Africa</b>      | Egypt               | <i>A. artemisiifolia</i> | 2002?  | Alien Naturalized    | Greuter, 2006; Shaltout, 2004.  |
| <b>Europe</b>      | Estonia             | <i>A. artemisiifolia</i> | 1954   | Alien Casual         | Gudzinskas, 1993  |
| <b>Europe</b>      | Estonia             | <i>A. psilostachya</i>   | /      | Alien Casual         | Greuter, 2006   |
| <b>Europe</b>      | Estonia             | <i>A. trifida</i>        | /      | Alien Casual         | EPPO, 2016  |

|                    |            |                          |           |                                      |  |
|--------------------|------------|--------------------------|-----------|--------------------------------------|--|
| <b>Europe</b>      | Finland    | <i>A. artemisiifolia</i> | 1900<1950 | Alien Naturalized                    | Finnish Ministry of Agriculture and Forestry, 2012; Lampinen and Lahti, 2016   |
| <b>Europe</b>      | Finland    | <i>A. trifida</i>        | 1900<1950 | Alien status unknown                 | Lampinen and Lahti, 2016   |
| <b>Europe</b>      | Finland    | <i>A. psilostachya</i>   | 1990s     | Alien status unknown                 | Lampinen and Lahti, 2016   |
| <b>Europe</b>      | France     | <i>A. tenuifolia</i>     | 1839      | Alien Naturalized                    | Thellung 1912; Chauvel <i>et al.</i> , 2015  |
| <b>Europe</b>      | France     | <i>A. artemisiifolia</i> | 1863      | Alien Invasive [Corse- Alien Casual] | Chauvel <i>et al.</i> , 2006; Csontos <i>et al.</i> , 2010   |
| <b>Europe</b>      | France     | <i>A. trifida</i>        | 1901      | Alien Naturalized                    | Chauvel <i>et al.</i> , 2015   |
| <b>Europe</b>      | France     | <i>A. psilostachya</i>   | 1931      | Alien Naturalized (Invasive?)        | Hibon, 1942; Fried <i>et al.</i> , 2015  |
| <b>Asia</b>        | Georgia    | <i>A. artemisiifolia</i> | /         | Alien Invasive                       | Kikodze <i>et al.</i> , 2010; EPPO, 2016   |
| <b>Asia</b>        | Georgia    | <i>A. trifida</i>        | /         | Alien status unknown                 | Kikodze <i>et al.</i> , 2010   |
| <b>Europe</b>      | Germany    | <i>A. artemisiifolia</i> | 1860      | Alien Naturalized                    | Buttler and Harms, 1999; Brandes and Nitzsche, 2006; Otto, 2006; Kazinczi <i>et al.</i> , 2008; Bullock <i>et al.</i> , 2012; Buttler, 2016.   |
| <b>Europe</b>      | Germany    | <i>A. trifida</i>        | 1877      | Alien Naturalized                    | Buttler and Harms, 1999; Follak <i>et al.</i> , 2013; Buttler, 2016; DAISIE, Species Factsheet: <i>A. trifida</i> . available at <a href="http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#">http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#</a> (Accessed in January 2017); Deutschlandflora WebGIS. Floristische Verbreitungskarten in Deutschland: <a href="https://deutschlandflora.de">https://deutschlandflora.de</a> (Accessed in January 2017) |
| <b>Europe</b>      | Germany    | <i>A. psilostachya</i>   | 1897      | Alien Naturalized                    | Buttler and Harms, 1999; Buttler, 2016; Bundesamt für Naturschutz - Floraweb, 2017: <a href="http://www.floraweb.de/pflanzenarten/artenhome.xsql?suchnr=20068&amp;">http://www.floraweb.de/pflanzenarten/artenhome.xsql?suchnr=20068&amp;</a> (Accessed in January 2017); Deutschlandflora WebGIS. Floristische Verbreitungskarten in Deutschland: <a href="https://deutschlandflora.de">https://deutschlandflora.de</a> (Accessed in January 2017)                                      |
| <b>Europe</b>      | Germany    | <i>A. tenuifolia</i>     | /         | Alien Casual                         | Buttler and Harms, 1999; Buttler, 2016   |
| <b>Europe</b>      | Greece     | <i>A. psilostachya</i>   | 2016      | Alien Naturalized?                   | Von Raab-Straube and Raus, 2016  |
| <b>Europe</b>      | Greece     | <i>A. artemisiifolia</i> | 2008?     | Alien status unknown                 | Arianoutsou <i>et al.</i> , 2010; Greuter and Raus, 2008   |
| <b>America (C)</b> | Guadeloupe | <i>A. artemisiifolia</i> | /         | Alien status unknown                 | Gerber <i>et al.</i> , 2011  |
| <b>America (S)</b> | Guatemala  | <i>A. artemisiifolia</i> | /         | Alien status unknown                 | Gerber <i>et al.</i> , 2011  |

|                    |                                 |                          |           |                               |   |
|--------------------|---------------------------------|--------------------------|-----------|-------------------------------|---|
| <b>America (C)</b> | Hawaiian Islands                | <i>A. artemisiifolia</i> | 1854      | Alien Invasive                | Wagner <i>et al.</i> , 1990; Pacific Island Ecosystems at Risk (PIER), 2013a  |
| <b>America (N)</b> | Hawaiian Islands                | <i>A. psilostachya</i>   | /         | Alien status unknown          | Randall, 2012; Pacific Island Ecosystems at Risk (PIER), 2013b  |
| <b>America (C)</b> | Hispaniola (Dominican Republic) | <i>A. artemisiifolia</i> | /         | Species occurring             | Acevedo-Rodriguez and Strong, 2012; Essl <i>et al.</i> , 2015   |
| <b>Europe</b>      | Hungary                         | <i>A. psilostachya</i>   | 1900 ca.  | Alien Invasive                | Puc, 2004; CABI, 2017   |
| <b>Europe</b>      | Hungary                         | <i>A. trifida</i>        | /         | Alien Invasive?               | Plank <i>et al.</i> , 2016  |
| <b>Europe</b>      | Hungary                         | <i>A. artemisiifolia</i> | 1922      | Alien Invasive                | Csontos <i>et al.</i> , 2010  |
| <b>Europe</b>      | Iceland                         | <i>A. artemisiifolia</i> | 1948      | Alien Casual                  | Wasowicz <i>et al.</i> , 2013   |
| <b>Asia</b>        | India                           | <i>A. artemisiifolia</i> | /         | Alien Invasive                | Khuroo <i>et al.</i> , 2012; Kohli <i>et al.</i> , 2012   |
| <b>Asia</b>        | India                           | <i>A. psilostachya</i>   | 1990s     | Alien Invasive                | Ramachandra Prasad <i>et al.</i> , 2013   |
| <b>Asia</b>        | India                           | <i>A. trifida</i>        | 2004-2009 | Alien status unknown          | Kumar <i>et al.</i> , 2009; Randall, 2012   |
| <b>Asia</b>        | Iran                            | <i>A. artemisiifolia</i> | /         | Alien status unknown          | Gerber <i>et al.</i> , 2011; Randall, 2012; Bararpour, 2014.  |
| <b>Asia</b>        | Iran                            | <i>A. psilostachya</i>   | /         | Alien status unknown          | Cheraghian, 2016a   |
| <b>Asia</b>        | Iran                            | <i>A. trifida</i>        | /         | Alien status unknown          | Randall, 2012; Bararpour, 2014; Cheraghian, 2016b   |
| <b>Europe</b>      | Ireland                         | <i>A. trifida</i>        | 1894      | Alien Casual                  | Reynolds, 2002  |
| <b>Europe</b>      | Ireland                         | <i>A. artemisiifolia</i> | 1900      | Alien Casual                  | Rich, 1994; Reynolds, 2002; Bullock <i>et al.</i> , 2012; Essl <i>et al.</i> , 2015                                     |
| <b>Asia</b>        | Israel                          | <i>A. artemisiifolia</i> | 1925      | Alien Casual                  | Waisel <i>et al.</i> , 2008; Yair <i>et al.</i> , 2017  |
| <b>Asia</b>        | Israel                          | <i>A. tenuifolia</i>     | 1984      | Alien Naturalized             | Greuter and Raus, 1995; Danin, 2000; Waisel <i>et al.</i> , 2008; Yair <i>et al.</i> , 2017                             |
| <b>Asia</b>        | Israel                          | <i>A. trifida</i>        | 1987      | Alien Casual (still present?) | Danin, 2000; Waisel <i>et al.</i> , 2008; Danin, 2016; Yair <i>et al.</i> , 2017  |
| <b>Asia</b>        | Israel                          | <i>A. psilostachya</i>   | 2006 ca.  | Alien Naturalized             | Yair <i>et al.</i> , 2017   |
| <b>Europe</b>      | Italy                           | <i>A. trifida</i>        | 1899      | Alien Naturalized             | Vignolo-Lutati, 1935; Mandrioli <i>et al.</i> , 1998; Celesti-Grappo <i>et al.</i> , 2009; Chauvel <i>et al.</i> , 2015 |
| <b>Europe</b>      | Italy                           | <i>A. psilostachya</i>   | 1927      | Alien Invasive                | Vignolo-Lutati, 1935; Mandrioli <i>et al.</i> , 1998; Conti <i>et al.</i> , 2005  |
| <b>Europe</b>      | Italy                           | <i>A. tenuifolia</i>     | 1935      | Alien Naturalized             | Vignolo-Lutati, 1935; Mandrioli <i>et al.</i> , 1998; Conti <i>et al.</i> , 2005  |
| <b>Europe</b>      | Italy                           | <i>A. artemisiifolia</i> | 1902      | Alien Invasive                | Gentili <i>et al.</i> , 2016  |

|                    |                  |                          |      |                      |  |
|--------------------|------------------|--------------------------|------|----------------------|--|
| <b>America (S)</b> | Jamaica          | <i>A. artemisiifolia</i> | /    | Alien status unknown | Gerber <i>et al.</i> , 2011  |
| <b>Asia</b>        | Japan            | <i>A. artemisiifolia</i> | 1877 | Alien Invasive       | Auld <i>et al.</i> , 2003; Kazinczi <i>et al.</i> , 2008; Fukano and Yahara, 2012; Essl <i>et al.</i> , 2015; Invasive Species of Japan (NIES). <i>A. artemisiifolia</i> . Available at: <a href="https://www.nies.go.jp/biodiversity/invasive/DB/detail/80400e.html">https://www.nies.go.jp/biodiversity/invasive/DB/detail/80400e.html</a> (Accessed January 2017) |
| <b>Asia</b>        | Japan            | <i>A. trifida</i>        | 1952 | Alien Invasive       | Auld <i>et al.</i> , 2003  |
| <b>Asia</b>        | Japan            | <i>A. psilostachya</i>   | /    | Alien Invasive       | Auld <i>et al.</i> , 2003; Mito and Uesugi, 2004; Ramachandra Prasad <i>et al.</i> , 2013.   |
| <b>Asia</b>        | Kazakhstan       | <i>A. artemisiifolia</i> | /    | Alien status unknown | Gerber <i>et al.</i> , 2011  |
| <b>Asia</b>        | Kazakhstan       | <i>A. psilostachya</i>   | /    | Alien Naturalized    | Von Raab-Straube and Raus, 2016.   |
| <b>Asia</b>        | Korean Peninsula | <i>A. trifida</i>        | 1964 | Alien Invasive       | Lee <i>et al.</i> , 2010; Kim and Kil, 2016  |
| <b>Asia</b>        | Korean Peninsula | <i>A. artemisiifolia</i> | 1955 | Alien Invasive       | Song <i>et al.</i> , 2012; Kim and Kil, 2016.  |
| <b>Europe</b>      | Latvia           | <i>A. trifida</i>        | 1900 | Alien Casual?        | Gudzinskas, 1993   |
| <b>Europe</b>      | Latvia           | <i>A. artemisiifolia</i> | 1936 | Alien Casual         | Gudzinskas, 1993   |
| <b>Europe</b>      | Latvia           | <i>A. psilostachya</i>   | /    | Alien Casual         | DAISIE, Species Factsheet: <i>A. coronopifolia</i> available at <a href="http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21701#">http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21701#</a> (Accessed in January 2017)   |
| <b>Europe</b>      | Liechtenstein    | <i>A. artemisiifolia</i> | 1995 | Alien Casual         | Greuter, 2006; Waldburger and Staub 2006   |
| <b>Europe</b>      | Lithuania        | <i>A. trifida</i>        | 1987 | Alien Casual         | Gudzinskas, 1993   |
| <b>Europe</b>      | Lithuania        | <i>A. artemisiifolia</i> | 1884 | Alien Casual         | Gudzinskas, 1993   |
| <b>Europe</b>      | Luxembourg       | <i>A. artemisiifolia</i> | /    | Alien Naturalized    | Ries, 2017   |
| <b>Africa</b>      | Lybia            | <i>A. artemisiifolia</i> | /    | Doubtful occurrence  | Greuter, 2006  |
| <b>Africa</b>      | Madagascar       | <i>A. artemisiifolia</i> | /    | Doubtful occurrence  | Kull <i>et al.</i> , 2012; Skálová <i>et al.</i> , 2015  |
| <b>America (C)</b> | Martinique       | <i>A. artemisiifolia</i> | /    | Alien status unknown | Gerber <i>et al.</i> , 2011  |
| <b>Africa</b>      | Mauritius        | <i>A. psilostachya</i>   | /    | Alien Invasive       | Macdonald <i>et al.</i> , 2003   |
| <b>America (C)</b> | Mexico           | <i>A. artemisiifolia</i> | /    | Species occurring    | Villaseñor and Espinosa-Garcia, 2004; Gerber <i>et al.</i> , 2011.   |
| <b>America (C)</b> | Mexico           | <i>A. psilostachya</i>   | /    | Native               | Vibrans, 1998; Roldán & Vibrans, 2009  |
| <b>America (C)</b> | Mexico           | <i>A. trifida</i>        | /    | Species occurring    | Villaseñor and Espinosa-Garcia, 2004; EPPO, 2016; CABI, 2017.  |

|                    |             |                          |                          |                      |   |
|--------------------|-------------|--------------------------|--------------------------|----------------------|---|
| <b>Europe</b>      | Moldova     | <i>A. artemisiifolia</i> | 1975?                    | Alien Naturalized    | Greuter, 2006; Bullock <i>et al.</i> , 2012   |
| <b>Europe</b>      | Moldova     | <i>A. trifida</i>        | /                        | Alien Casual         | EPPO, 2016  |
| <b>Asia</b>        | Mongolia    | <i>A. trifida</i>        | /                        | Alien status unknown | EPPO, 2016  |
| <b>Europe</b>      | Montenegro  | <i>A. artemisiifolia</i> | ?                        | Alien status unknown | Stešević and Petrović 2010; Karrer, 2016  |
| <b>Europe</b>      | Montenegro  | <i>A. psilostachya</i>   | /                        | Alien status unknown | Greuter, 2006   |
| <b>Africa</b>      | Morocco     | <i>A. psilostachya</i>   | 1994                     | Alien status unknown | Tanji, 2005   |
| <b>Europe</b>      | Netherlands | <i>A. artemisiifolia</i> | 1875                     | Alien Naturalized    | Van Denderen <i>et al.</i> , 2010; Odé and Beringen, 2017a  |
| <b>Europe</b>      | Netherlands | <i>A. psilostachya</i>   | 1905                     | Alien Naturalized    | Van Denderen <i>et al.</i> , 2010; Odé and Beringen, 2017b  |
| <b>Europe</b>      | Netherlands | <i>A. trifida</i>        | more frequent from 1960s | Alien Casual         | Van Denderen <i>et al.</i> , 2010; Odé and Beringen 2017c   |
| <b>Oceania</b>     | New Zealand | <i>A. tenuifolia</i>     | 1950                     | Alien Casual         | Howell and Sawyer, 2006   |
| <b>Oceania</b>     | New Zealand | <i>A. artemisiifolia</i> | 1911                     | Alien Casual         | Webb <i>et al.</i> , 1988; Essl <i>et al.</i> , 2015  |
| <b>Europe</b>      | Norway      | <i>A. artemisiifolia</i> | 1930                     | Alien Casual         | Fremstad and Elven 1997; Kazinczi <i>et al.</i> , 2008  |
| <b>Europe</b>      | Norway      | <i>A. psilostachya</i>   | /                        | Alien Casual         | Greuter, 2006   |
| <b>Europe</b>      | Norway      | <i>A. trifida</i>        | /                        | Alien Casual         | Randall, 2012; EPPO, 2016; CABI, 2017; DAISIE, Species Factsheet: <i>A. trifida</i> . available at <a href="http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#">http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#</a> (Accessed in January 2017) |
| <b>America (S)</b> | Paraguay    | <i>A. artemisiifolia</i> | /                        | Species occurring    | Zuloaga <i>et al.</i> , 2008; Essl <i>et al.</i> , 2015; CABI, 2017; Tropicos.org. Missouri Botanical Garden. 01 Feb 2017 < <a href="http://www.tropicos.org/Name/2701648">http://www.tropicos.org/Name/2701648</a>   |
| <b>America (S)</b> | Paraguay    | <i>A. tenuifolia</i>     | /                        | Native               | Sáenz and Gutiérrez, 2008   |
| <b>America (S)</b> | Perù        | <i>A. artemisiifolia</i> | /                        | Species occurring    | Gutte, 1978; Gerber <i>et al.</i> , 2011; Zárate <i>et al.</i> , 2015; Tropicos.org. Missouri Botanical Garden. 20 Jan 2017 < <a href="http://www.tropicos.org/Name/2701648">http://www.tropicos.org/Name/2701648</a> >   |
| <b>America (S)</b> | Perù        | <i>A. tenuifolia</i>     | /                        | Native               | Randall, 2012   |
| <b>Europe</b>      | Poland      | <i>A. artemisiifolia</i> | 1873                     | Alien Naturalized    | Gudzinkas, 1993; Kazinczi <i>et al.</i> , 2008; Tokarska-Guzik <i>et al.</i> , 2011   |
| <b>Europe</b>      | Poland      | <i>A. psilostachya</i>   | /                        | Alien Naturalized    | Kazinczi <i>et al.</i> , 2008; Tokarska-Guzik <i>et al.</i> , 2011  |
| <b>Europe</b>      | Poland      | <i>A. trifida</i>        | /                        | Alien Casual         | Kazinczi <i>et al.</i> , 2008; Tokarska-Guzik <i>et al.</i> , 2011  |

|                    |                      |                          |                        |   |  |
|--------------------|----------------------|--------------------------|------------------------|---|--|
| <b>Europe</b>      | Portugal             | <i>A. artemisiifolia</i> | 1972                   | Alien Invasive<br>(Isle of Madeira<br>Alien Casual) | Borges <i>et al.</i> , 2008; Amor Morales <i>et al.</i> , 2012   |
| <b>America (C)</b> | Puerto Rico          | <i>A. tenuifolia</i>     | /                      | Alien Naturalized                                   | Liogier, 1997; Acevedo-Rodriguez and Strong, 2012; Gann <i>et al.</i> , 2015-2017                      |
| <b>Europe</b>      | Romania              | <i>A. artemisiifolia</i> | 1907                   | Alien Invasive                                      | Kazinczi <i>et al.</i> , 2008; Csontos <i>et al.</i> , 2010; Bullock <i>et al.</i> , 2012; Sîrbu, 2012 |
| <b>Europe</b>      | Romania              | <i>A. trifida</i>        | 1970-1980              | Alien Naturalized                                   | Sîrbu, 2012; Stoyanov <i>et al.</i> , 2014   |
| <b>Europe</b>      | Romania              | <i>A. psilostachya</i>   | /                      | Alien Naturalized                                   | Sîrbu, 2012  |
| <b>Europe</b>      | Russia               | <i>A. psilostachya</i>   | /                      | Alien Naturalized                                   | EPPO, 2016   |
| <b>Europe</b>      | Russia<br>(European) | <i>A. artemisiifolia</i> | 1918                   | Alien Invasive                                      | Gudzinkas, 1993; Csontos <i>et al.</i> , 2010; Vinogradova <i>et al.</i> , 2010; Randall, 2012         |
| <b>Europe</b>      | Russia<br>(European) | <i>A. trifida</i>        | /                      | Alien Naturalized                                   | Randall, 2012; EPPO, 2016  |
| <b>Europe</b>      | Russia<br>(European) | <i>A. psilostachya</i>   | /                      | Alien Naturalized                                   | EPPO, 2016   |
| <b>Europe</b>      | Serbia               | <i>A. artemisiifolia</i> | 1935                   | Alien Invasive                                      | Vrbničanin <i>et al.</i> , 2004; Kazinczi <i>et al.</i> , 2008; Bullock <i>et al.</i> , 2012           |
| <b>Europe</b>      | Serbia               | <i>A. trifida</i>        | 1982                   | Alien Naturalized                                   | Vrbničanin <i>et al.</i> , 2004; Follak <i>et al.</i> , 2013; EPPO, 2016                               |
| <b>Europe</b>      | Serbia               | <i>A. tenuifolia</i>     | /                      | Alien Naturalized                                   | Vrbničanin <i>et al.</i> , 2004  |
| <b>Europe</b>      | Slovakia             | <i>A. artemisiifolia</i> | 1949                   | Alien Invasive                                      | Medvecká <i>et al.</i> 2012  |
| <b>Europe</b>      | Slovakia             | <i>A. trifida</i>        | 1980                   | Alien Casual  | Medvecká <i>et al.</i> 2012  |
| <b>Europe</b>      | Slovenia             | <i>A. artemisiifolia</i> | 1993 (after<br>WW II?) | Alien Invasive                                      | Kazinczi <i>et al.</i> , 2008.; Galzina <i>et al.</i> , 2010; Zelnik, 2012                             |
| <b>Europe</b>      | Slovenia             | <i>A. trifida</i>        | late 1980s             | Alien Casual  | Follak <i>et al.</i> , 2013; EPPO, 2016  |
| <b>Africa</b>      | South Africa         | <i>A. artemisiifolia</i> | /                      | Alien Naturalized                                   | Germishuizen and Meyer, 2003; Henderson, 2007; Essl <i>et al.</i> , 2015                               |
| <b>Africa</b>      | South Africa         | <i>A. psilostachya</i>   | /                      | Alien Naturalized                                   | Wells <i>et al.</i> , 1986; Germishuizen and Meyer, 2003; Randall, 2012; SANBI, 2015a                  |
| <b>Africa</b>      | South Africa         | <i>A. tenuifolia</i>     | /                      | Alien Naturalized                                   | Germishuizen and Meyer, 2003; SANBI, 2015b   |
| <b>Europe</b>      | Spain                | <i>A. tenuifolia</i>     | 1954                   | Alien Naturalized                                   | Amor Morales <i>et al.</i> , 2012  |
| <b>Europe</b>      | Spain                | <i>A. psilostachya</i>   | 1981                   | Alien Invasive                                      | Amor Morales <i>et al.</i> , 2012  |
| <b>Europe</b>      | Spain                | <i>A. artemisiifolia</i> | 1983                   | Alien Invasive                                      | Amor Morales <i>et al.</i> , 2012  |
| <b>Europe</b>      | Spain                | <i>A. trifida</i>        | 1983                   | Alien Naturalized                                   | Amor Morales <i>et al.</i> , 2012  |
| <b>Europe</b>      | Spain-<br>Balears    | <i>A. tenuifolia</i>     | 2004                   | Alien Naturalized                                   | Fraga and García, 2004   |

|                    |                          |                          |       |                      |   |
|--------------------|--------------------------|--------------------------|-------|----------------------|---|
| <b>Africa</b>      | Swaziland                | <i>A. artemisiifolia</i> | /     | Alien Naturalized    | Randall, 2012; Swaziland's Alien Plants Database. <a href="http://www.sntc.org.sz/alienplants/index.asp">http://www.sntc.org.sz/alienplants/index.asp</a>   |
| <b>Europe</b>      | Sweden                   | <i>A. trifida</i>        | 1909  | Alien Casual         | Anderberg, 2000; Gerber <i>et al.</i> , 2011; Randall, 2012; DAISIE, Species Factsheet: <i>A. trifida</i> . available at <a href="http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#">http://www.europe-aliens.org/speciesFactsheet.do?speciesId=21722#</a> (Accessed in January 2017) |
| <b>Europe</b>      | Sweden                   | <i>A. psilostachya</i>   | 1928  | Alien Naturalized    | Dahl <i>et al.</i> , 1999; Anderberg, 2005  |
| <b>Europe</b>      | Sweden                   | <i>A. artemisiifolia</i> | 1866  | Alien Casual         | Dahl <i>et al.</i> , 1999; Anderberg, 2000; Smith <i>et al.</i> , 2013; Smith <i>et al.</i> , 2008; Dahl, 1999; Smith <i>et al.</i> , 2013  |
| <b>Europe</b>      | Switzerland              | <i>A. trifida</i>        | 1900  | Alien status unknown | Follak <i>et al.</i> , 2013; EPPO, 2016.  |
| <b>Europe</b>      | Switzerland              | <i>A. psilostachya</i>   | /     | Alien status unknown | Greuter, 2006; Hess <i>et al.</i> , 2006  |
| <b>Europe</b>      | Switzerland              | <i>A. artemisiifolia</i> | 1850s | Alien Invasive       | Tamarcaz <i>et al.</i> , 2005; Kazinczi <i>et al.</i> , 2008; Bullock <i>et al.</i> , 2012  |
| <b>Asia</b>        | Taiwan                   | <i>A. artemisiifolia</i> | 1971  | Alien Naturalized    | Wu <i>et al.</i> , 2004; Wu <i>et al.</i> , 2010; Peng, 2013  |
| <b>Asia</b>        | Taiwan                   | <i>A. psilostachya</i>   | 2000  | Alien Naturalized    | Tseng <i>et al.</i> , 2004; Ramachandra Prasad <i>et al.</i> , 2012; Wu, <i>et al.</i> , 2010; Chen and Hind, 2011  |
| <b>Asia</b>        | Turkey                   | <i>A. artemisiifolia</i> | 1995  | Alien Invasive       | Byfield and Baytop 1998; Zemmer <i>et al.</i> 2012; Behçet 2004; Onen <i>et al.</i> , 2014; Arslan <i>et al.</i> , 2015   |
| <b>Asia</b>        | Turkey                   | <i>A. tenuifolia</i>     | 2000  | Alien Naturalized?   | Behçet 2004; Özhatay and Kültür, 2006   |
| <b>Europe</b>      | Ukraine                  | <i>A. artemisiifolia</i> | 1925  | Alien Invasive       | Smith <i>et al.</i> , 2013; Bullock <i>et al.</i> , 2012; EPPO, 2016.   |
| <b>Europe</b>      | Ukraine                  | <i>A. psilostachya</i>   | /     | Alien status unknown | Greuter, 2006   |
| <b>Europe</b>      | Ukraine                  | <i>A. trifida</i>        | /     | Alien Casual         | Yavorska, 2009  |
| <b>Europe</b>      | United Kingdom           | <i>A. artemisiifolia</i> | 1836  | Alien Invasive       | Rich, 1994; Bullock <i>et al.</i> , 2012; Essl <i>et al.</i> , 2015   |
| <b>Europe</b>      | United Kingdom           | <i>A. trifida</i>        | 1897  | Alien Casual         | Rich, 1994; Sell and Murrell, 2006; EPPO, 2016; Online Atlas British and Irish Flora: <a href="http://www.brc.ac.uk/plantatlas/index.php?q=plant/A.-trifida">http://www.brc.ac.uk/plantatlas/index.php?q=plant/A.-trifida</a>   |
| <b>Europe</b>      | United Kingdom           | <i>A. psilostachya</i>   | 1880s | Alien Naturalized    | Rich, 1994; Sell and Murrell, 2006; On line Atlas of the British and Irish Flora: <a href="http://www.brc.ac.uk/plantatlas/index.php?q=plant/A.-psilostachya">http://www.brc.ac.uk/plantatlas/index.php?q=plant/A.-psilostachya</a>   |
| <b>Europe</b>      | United Kingdom           | <i>A. tenuifolia</i>     | /     | Doubtful occurrence  | Stace, 2010; Randall, 2012  |
| <b>America (N)</b> | United States of America | <i>A. psilostachya</i>   | /     | Native               | Bassett and Crompton, 1975  |



|                    |                          |                          |   |                      |   |
|--------------------|--------------------------|--------------------------|---|----------------------|---|
| <b>America (N)</b> | United States of America | <i>A. tenuifolia</i>     | / | Alien status unknown | Liogier, 1997; USDA - NRCS, 2017  |
| <b>America (N)</b> | United States of America | <i>A. trifida</i>        | / | Native               | Bassett and Crompton, 1982  |
| <b>America (N)</b> | United States of America | <i>A. artemisiifolia</i> | / | Native               | Bassett and Crompton, 1975  |
| <b>America (S)</b> | Uruguay                  | <i>A. artemisiifolia</i> | / | Species occurring    | Tejera and Beri 2005; Zuloaga <i>et al.</i> , 2008; Essl <i>et al.</i> , 2015 |
| <b>America (S)</b> | Uruguay                  | <i>A. tenuifolia</i>     | / | Native               | Sáenz and Gutiérrez, 2008   |

**Table 3 - Environmental requirements of *Ambrosia* species (ragweeds): main data related to colonized habitat types (native and invasive range), suitable climatic, soil and light conditions. Requirements of each species are highlighted in grey and doubtful attributions are signaled by "(?)".**

| <b>Species</b> | <i>Ambrosia artemisiifolia</i> L.        | <i>Ambrosia trifida</i> L.     | <i>Ambrosia psilostachya</i> DC.         | <i>Ambrosia tenuifolia</i> Spreng. |
|----------------|--|--------------------------------|--|------------------------------------|
|                | Disturbed open habitat                   | Disturbed open habitat         | Disturbed open habitat                   | Disturbed open habitat             |
|                | Semi-natural grasslands                  | Semi-natural grasslands        | Semi-natural grasslands                  | Semi-natural grasslands            |
|                | Croplands                                | Croplands                      | Croplands                                | Croplands                          |
|                | Along transportation corridors           | Along transportation corridors | Along transportation corridors           | Along transportation corridors     |
|                | Wastelands                               | Wastelands                     | Wastelands                               | Wastelands                         |
|                | Riparian habitat                         | Riparian habitat               | Riparian habitat                         | Riparian habitat                   |
|                | Dunes                                    | Dunes                          | Dunes                                    | Dunes                              |
|                | Non dense wood                           | Non dense wood                 | Non dense wood                           | Non dense wood                     |
|                | Disturbed open habitat                   | Disturbed open habitat         | Disturbed open habitat                   | Disturbed open habitat             |
|                | Semi-natural grasslands                  | Semi-natural grasslands        | Semi-natural grasslands                  | Semi-natural grasslands            |
|                | Croplands                                | Croplands                      | Croplands                                | Croplands                          |
|                | Along transportation corridors           | Along transportation corridors | Along transportation corridors           | Along transportation corridors (?) |
|                | Wastelands                               | Wastelands                     | Wastelands                               | Wastelands                         |
|                | Riparian habitat                         | Riparian habitat               | Riparian habitat                         | Riparian habitat (?)               |
|                | Dunes                                    | Dunes                          | Dunes                                    | Dunes                              |
|                | Non dense wood                           | Non dense wood                 | Non dense wood                           | Non dense wood                     |
|                | Warm temperate climate (with exceptions) | Warm temperate climate         | Warm temperate climate (with exceptions) | Warm temperate climate             |
|                | Drought tolerant                         | Drought tolerant               | Drought tolerant                         | Drought tolerant                   |
|                | Freeze tolerant                          | Freeze tolerant                | Freeze tolerant                          | Freeze tolerant                    |
|                | Alkaline                                 | Alkaline?                      | Alkaline                                 | Alkaline?                          |
|                | Acid                                     | Acid                           | Acid                                     | Acid                               |
|                | Silty                                    | Silty                          | Silty                                    | Silty                              |
|                | Sandy                                    | Sandy                          | Sandy                                    | Sandy                              |
|                | Well drained/ Dry                        | Well drained/ Dry              | Well drained/ Dry                        | Well drained/ Dry                  |
|                | Moist/Wet                                | Moist/Wet                      | Moist/Wet                                | Moist/Wet                          |
|                | Saline                                   | Saline                         | Saline                                   | Saline                             |
|                | Metal                                    | Metal                          | Metal                                    | Metal                              |
|                | Heliophyllous                            | Heliophyllous                  | Heliophyllous                            | Heliophyllous                      |
|                | Shady-tolerant                           | Shady-tolerant                 | Shady-tolerant                           | Shady-tolerant                     |




**Table 4 - Life strategies of *Ambrosia* species (ragweeds): relevant traits contributing to strengthen resistance, resilience and competition of ragweeds in the wild. Traits of each species are highlighted in gray.**

| <b>Species</b> | <i>Ambrosia artemisiifolia</i> L.                 | <i>Ambrosia trifida</i> L.                        | <i>Ambrosia psilostachya</i> DC.                  | <i>Ambrosia tenuifolia</i> Spreng.                |
|----------------|---|---|---|---|
|                | Morphologic structures                            | Morphologic structures                            | Morphologic structures                            | Morphologic structures                            |
|                | Chemical defence against stress and predators     | Chemical defence against stress and predators     | Chemical defence against stress and predators     | Chemical defence against stress and predators     |
|                | Allelopathy                                       | Allelopathy                                       | Allelopathy                                       | Allelopathy                                       |
|                | Mychorrhiza                                       | Mychorrhiza                                       | Mychorrhiza                                       | Mychorrhiza                                       |
|                | Reallocation biomass                              | Reallocation biomass                              | Reallocation biomass                              | Reallocation biomass                              |
|                | Resprouting                                       | Resprouting                                       | Resprouting                                       | Resprouting                                       |
|                | Rhizome   | Rhizome   | Rhizome   | Rhizome   |
|                | Secondary dormancy                                | Secondary dormancy                                | Secondary dormancy                                | Secondary dormancy                                |
|                | Soil seed bank                                    | Soil seed bank                                    | Soil seed bank                                    | Soil seed bank                                    |
|                | Long lasting soil seed bank                       | Long lasting soil seed bank                       | Long lasting soil seed bank                       | Long lasting soil seed bank                       |
|                | Advantages from vegetation gaps                   | Advantages from vegetation gaps                   | Advantages from vegetation gaps                   | Advantages from vegetation gaps                   |
|                | Weak competitor in more evolved vegetation stages | Weak competitor in more evolved vegetation stages | Weak competitor in more evolved vegetation stages | Weak competitor in more evolved vegetation stages |
|                | Persistence in more evolved vegetation stages     | Persistence in more evolved vegetation stages     | Persistence in more evolved vegetation stages     | Persistence in more evolved vegetation stages     |

**Table 5 - Reproduction and dispersal of *Ambrosia* species (ragweeds): relevant data in understanding the reproductive and dispersal potential of ragweeds. The characteristic of each species is highlighted in grey and doubtful attributions are signaled by "(?)".**

| <b>Species</b>                | <i>Ambrosia artemisiifolia</i> L.                                       | <i>Ambrosia trifida</i> L.  | <i>Ambrosia psilostachya</i> DC.  | <i>Ambrosia tenuifolia</i> Spreng.                                      |
|-------------------------------|---|---|---|---|
| <b>POLLINATION</b>            | Anemophylous  | Anemophylous  | Anemophylous  | Anemophylous  |
|                               | Sexual  | Sexual  | Sexual  | Sexual  |
|                               | Vegetative  | Vegetative  | Vegetative  | Vegetative  |
| <b>SEED DIMENSION</b>         | 3.5 mm long   | more than 6 mm long   | 3-4.5 mm long   | 3-5 mm long   |
|                               | Very high   | Very high   | Very high   | Very high   |
|                               | High  | High  | High  | High (?)  |
|                               | Scarce  | Scarce  | Scarce  | Scarce  |
|                               | Existing  | Existing  | Existing  | Existing?   |
|                               | Highly relevant   | Highly relevant   | Highly relevant   | Highly relevant   |
|                               | No evidences  | No evidences  | No evidences  | No evidences  |
|                               | Vernalization   | Vernalization   | Vernalization   | Vernalization   |
|                               | Light   | Light   | Light   | Light   |
|                               | Soil moisture   | Soil moisture   | Soil moisture   | Soil moisture   |
| <b>PRIMARY SEED DISPERSAL</b> | Barochory   | Barochory   | Barochory   | Barochory   |
|                               | Anemochory  | Anemochory  | Anemochory (?)  | Anemochory (?)  |
|                               | Epizoochory   | Epizoochory   | Epizoochory   | Epizoochory   |
|                               | Endozoochory  | Endozoochory  | Endozoochory (?)  | Endozoochory  |
|                               | Hydrochory  | Hydrochory  | Hydrochory  | Hydrochory  |
|                               | Movement of soils   | Movement of soils (?)   | Movement of soils (?)   | Movement of soils (?)   |
|                               | Mowing or agricultural machinery  | Mowing or agricultural machinery (?)                                    | Mowing or agricultural machinery (?)                                    | Mowing or agricultural machinery (?)                                    |
|                               | Car and train passage   | Car and train passage (?)   | Car and train passage (?)   | Car and train passage   |
|                               | Grain, vegetables, fodder, bird food and oil-seeds commercial exchanges | Grain, vegetables, fodder, bird food and oil-seeds commercial exchanges | Grain, vegetables, fodder, bird food and oil-seeds commercial exchanges | Grain, vegetables, fodder, bird food and oil-seeds commercial exchanges |

**Table 6 - Allergenic potential of *Ambrosia* species (ragweeds): relevant elements of plants contributing to determine and increase allergy reaction. Elements of each species are highlighted in gray.**

| <b>Species</b>   | <b>Ambrosia artemisiifolia L.</b>   | <b>Ambrosia trifida L.</b>  | <b>Ambrosia psilostachya DC.</b>   | <b>Ambrosia tenuifolia Spreng.</b>    |
|--|---|---|--|---------------------------------------|
|  | Pollen  | Pollen  | Pollen   | Pollen                                |
|  | Plant debris  | Plant debris  | Plant debris   | Plant debris                          |
| <b>POLLEN DIMENSION</b>                                    | 18–22 µm  | 19.25 µm  | 21 - 23 µm up to 26.4 µm   | ?                                     |
| <b>CROSS-REACTIVITY</b>                                    | Amb a 5 and Amb t 5   | Amb a 5 and Amb t 5   | Amb a 5 and Amb p 5  | poorly known                          |
|  | Long-distance transport of pollen   | Long-distance transport of pollen (?)   | Long-distance transport of pollen (?)  | Long-distance transport of pollen (?) |
|  | Atmospheric Pollution   | Atmospheric Pollution(?)  | Atmospheric Pollution (?)  | Atmospheric Pollution (?)             |
|  | Warming climate   | Warming climate   | Warming climate  | Warming climate (?)                   |
|  | Disturbance   | Disturbance (?)   | Disturbance  | Disturbance (?)                       |
| <b>POLLEN GRAIN</b><br>(from Robbins <i>et al.</i> , 1979) |  |  |  | -                                     |

**Table 7 - Allergens in *Ambrosia* species (ragweeds): characteristics of identified allergens of each analyzed species (Source: \* = IUIS, \*\* = ExPASy).**

| Allergen*                                | Isoform*      | MW(SDS-PAGE) (kDa)*   | Theoretical pI ** | Biological function*         | Allergenic Potential (%)* |
|--|---------------|---|-------------------|------------------------------|---------------------------|
| <b><i>Ambrosia artemisiifolia</i> L.</b> |               |   |                   |                              |                           |
|  | Amb a 1.0101  |   | 5.58              |                              |                           |
|  | Amb a 1.0201  |   | 6.63              |                              |                           |
|  | Amb a 1.0202  |   | 6.63              |                              |                           |
|  | Amb a 1.0301  |   | 5.72              |                              |                           |
|  | Amb a 1.0302  |   | 5.72              |                              |                           |
|  | Amb a 1.0303  |   | 5.79              |                              |                           |
|  | Amb a 1.0304  |   | 5.79              |                              |                           |
|  | Amb a 1.0305  |   | 5.79              |                              |                           |
|  | Amb a 1.0401  |   | 5.61              |                              |                           |
|  | Amb a 1.0402  |   | 5.22              |                              |                           |
|  | Amb a 1.0501  |   | 6.00              |                              |                           |
|  | Amb a 1.0502  |   | 5.79              |                              |                           |
| Amb a 3                                  | Amb a 3.0101  | 11  | 6.11              | Plastocyanin                 | 51                        |
| Amb a 4                                  | Amb a 4.0101  | 30  | 4.88              | Defense-like protein         | Unknown                   |
| Amb a 5                                  | Amb a 5.0101  | 5   | 8.19              | unknown                      | 10-20                     |
| Amb a 6                                  | Amb a 6.0101  | 10  | 8.93              | Lipid Transfer protein (LTP) | 21                        |
| Amb a 7                                  | Amb a 7.0101  | 12  | -                 | Plastocyanin                 | 15-20                     |
|  | Amb a 8.0101  | 14  | 4.79              |                              |                           |
|  | Amb a 8.0102  |   | 4.88              |                              |                           |
|  | Amb a 9.0101  | 10  | 4.17              |                              |                           |
|  | Amb a 9.0102  |   | 4.15              |                              |                           |
| Amb a 10                                 | Amb a 10.0101 | 18  | 4.25              | Polcalcine-like protein      | 10-15                     |
| Amb a 11                                 | Amb a 11.0101 | 37 kDa (natural purified mature protein), 52 kDa (natural purified zymogen) | 6.43              | Cysteine protease            | 54                        |
| <b><i>Ambrosia psilostachya</i> DC.</b>  |               |   |                   |                              |                           |
|  | Amb p 5.0101  |   |                   |                              |                           |
|  | Amb p 5.0201  |   |                   |                              |                           |
| <b><i>Ambrosia trifida</i> L.</b>        |               |   |                   |                              |                           |
| Amb t 5                                  | Amb t 5.0101  | 5   |                   | Unknown                      | 5                         |

## Figures Legend

**Figure 1** - Global distribution of *Ambrosia* species (ragweeds). **Alien:** the species is not native to a country. Status (*invasive, naturalized, and casual*) is attributed when the condition is confirmed at country and/or local level; “?” indicate uncertainty due to lack of confirmations. **Alien status unknown:** the species is alien to a country, but its status is indefinite. **Species occurring:** the species occurs in one country, but there are uncertainties/ inconsistencies about its origin (alien/native). **Native:** the species is not introduced from other countries; it is part of local flora. **Doubtful occurrence:** the occurrence of the species is not confirmed.

**Figure 2** - Status of *Ambrosia* species (ragweeds) at world level. Each pie chart describes the percentage of countries where the species is native or alien, their status and the table below the numbers for each category.