

## The XMM-Newton extended survey of the Taurus molecular cloud (XEST)<sup>★,★★</sup>

M. Güdel<sup>1</sup>, K. R. Briggs<sup>1</sup>, K. Arzner<sup>1</sup>, M. Audard<sup>2,★★★</sup>, J. Bouvier<sup>3</sup>, E. D. Feigelson<sup>4</sup>, E. Franciosini<sup>5</sup>, A. Glauser<sup>1</sup>, N. Grosso<sup>3</sup>, G. Micela<sup>5</sup>, J.-L. Monin<sup>3</sup>, T. Montmerle<sup>3</sup>, D. L. Padgett<sup>6</sup>, F. Palla<sup>7</sup>, I. Pillitteri<sup>8</sup>, L. Rebull<sup>6</sup>, L. Scelsi<sup>8</sup>, B. Silva<sup>9,10</sup>, S. L. Skinner<sup>11</sup>, B. Stelzer<sup>5</sup>, and A. Telleschi<sup>1</sup>

<sup>1</sup> Paul Scherrer Institut, Würenlingen and Villigen, 5232 Villigen PSI, Switzerland  
e-mail: guedel@astro.phys.ethz.ch

<sup>2</sup> Columbia Astrophysics Laboratory, Mail Code 5247, 550 West 120th Street, New York, NY 10027, USA

<sup>3</sup> Laboratoire d'Astrophysique de Grenoble, Université Joseph Fourier - CNRS, BP 53, 38041 Grenoble Cedex, France

<sup>4</sup> Department of Astronomy & Astrophysics, Penn State University, 525 Davey Lab, University Park, PA 16802, USA

<sup>5</sup> INAF - Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90134 Palermo, Italy

<sup>6</sup> Spitzer Science Center, California Institute of Technology, Mail Code 220-6, Pasadena, CA 91125, USA

<sup>7</sup> INAF - Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi, 5, 50125 Firenze, Italy

<sup>8</sup> Dipartimento di Scienze Fisiche ed Astronomiche, Università di Palermo, Piazza del Parlamento 1, 90134 Palermo, Italy

<sup>9</sup> Centro de Astrofísica da Universidade do Porto, Rua das Estrelas, 4150 Porto, Portugal

<sup>10</sup> Departamento de Matemática Aplicada, Faculdade de Ciências da Universidade do Porto, 4169 Porto, Portugal

<sup>11</sup> CASA, 389, University of Colorado, Boulder, CO 80309-0389, USA

Received 31 May 2006 / Accepted 5 August 2006

### ABSTRACT

**Context.** The Taurus Molecular Cloud (TMC) is the nearest large star-forming region, prototypical for the distributed mode of low-mass star formation. Pre-main sequence stars are luminous X-ray sources, probably mostly owing to magnetic energy release.

**Aims.** The *XMM-Newton Extended Survey of the Taurus Molecular Cloud* (XEST) presented in this paper surveys the most populated  $\approx 5$  square degrees of the TMC, using the *XMM-Newton* X-ray observatory to study the thermal structure, variability, and long-term evolution of hot plasma, to investigate the magnetic dynamo, and to search for new potential members of the association. Many targets are also studied in the optical, and high-resolution X-ray grating spectroscopy has been obtained for selected bright sources.

**Methods.** The X-ray spectra have been coherently analyzed with two different thermal models (2-component thermal model, and a continuous emission measure distribution model). We present overall correlations with fundamental stellar parameters that were derived from the previous literature. A few detections from *Chandra* observations have been added.

**Results.** The present overview paper introduces the project and provides the basic results from the X-ray analysis of all sources detected in the XEST survey. Comprehensive tables summarize the stellar properties of all targets surveyed. The survey goes deeper than previous X-ray surveys of Taurus by about an order of magnitude and for the first time systematically accesses very faint and strongly absorbed TMC objects. We find a detection rate of 85% and 98% for classical and weak-line T Tau stars (CTTS resp. WTTS), and identify about half of the surveyed protostars and brown dwarfs. Overall, 136 out of 169 surveyed stellar systems are detected. We describe an X-ray luminosity vs. mass correlation, discuss the distribution of X-ray-to-bolometric luminosity ratios, and show evidence for lower X-ray luminosities in CTTS compared to WTTS. Detailed analysis (e.g., variability, rotation-activity relations, influence of accretion on X-rays) will be discussed in a series of accompanying papers.

**Key words.** stars: formation – stars: pre-main sequence – X-rays: stars – gravitational waves

## 1. Introduction

Studies of star-forming regions have drawn a picture in which cool, molecular gas, contracting to protostars with surrounding accretion disks, co-exists with high-energy radiation. The

latter is emitted by plasma in the stellar environment that is continuously heated to temperatures beyond one million degrees. This radiation is most prominently seen in the soft (0.1–10 keV) X-ray range and is conventionally attributed to the presence of magnetically trapped plasma in the outer stellar atmosphere. This plasma is reminiscent of, in the broadest sense, the solar coronal plasma. There is little doubt that both lower-energy (extreme ultraviolet) and higher-energy (hard X-ray and gamma-ray) radiation is amply present, but strong photoelectric absorption or, respectively, extremely low flux precludes direct detection. Further, non-thermal radio emission from relativistic electrons trapped in the magnetic fields gives unambiguous evidence for the presence of an accelerated population of electrons

\* Table 3, Full Tables 4–11 and Appendices A and B are only available in electronic form at <http://www.aanda.org>

\*\* Full Table A.1 is only available in electronic form at the CDS via anonymous ftp to [cdsweb.u-strasbg.fr](http://cdsweb.u-strasbg.fr) (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/468/353>

\*\*\* New address: Integral Science Data Centre, Ch. d'Ecogia 16, 1290 Versoix, and Geneva Observatory, University of Geneva, Ch. des Maillettes 51, 1290 Sauverny, Switzerland.

similar to populations seen in the solar and in stellar coronae during flares (Güdel 2002). Again, by implication from the solar analogy, we expect that high-energy, non-thermal particles (electrons and ions) play a crucial role also outside the confines of the magnetic stellar atmosphere.

But in contrast to the solar case, high-energy radiation from forming stars has a profound influence on its environment as dense accretion disks and the larger-scale molecular envelope are irradiated, heated, and ionized by high-energy photons and particles. There are numerous consequences of these processes. Ultraviolet and X-ray irradiation of molecular disks heats their upper layers (Alexander et al. 2004; Glassgold et al. 2004), making them accessible to magnetic fields, thus inducing, for example, the magnetorotational instability (Balbus & Hawley 1991). The latter is thought to be a prime source for increased disk viscosity, which in turn is the main driver of the accretion process and thus the stellar formation process in the first place. Complex chemical networks are also put in operation by heating and ionizing the molecular environment (Glassgold et al. 2004), thus fundamentally altering composition, opacity, and cooling behavior of the material. Magnetic fields are thought to attach to the inner border of the lightly ionized accretion disk, thus transporting material to the stellar surface while at the same time applying torques to the star that may regulate its rotation period (Montmerle et al. 2000). If the stellar magnetic fields are generated in a solar-like dynamo in which rotation is a key parameter, then the long-term evolution of the magnetic field production and thus of the high-energy radiation is indeed directly controlled by the accretion process in the feedback loop sketched above. Further, magnetic fields and ionization of the disk may be relevant for the launch of bipolar jets and molecular outflows (Pudritz & Norman 1986), a widely observed but poorly understood mechanism that seems to regulate the mass buildup of the forming star. Alternatively, these jets may originate from the magnetic fields spanning between the star and the disk (Shu et al. 1994), or from plasma clouds ejected in the process of magnetic reconnection of these same fields (Hayashi et al. 1996).

The origin of the X-ray emission is still not entirely clear. While the bulk of the X-rays from T Tau stars is thought to originate in a magnetically trapped corona consisting of complex magnetic loops anchored in photospheric magnetic regions like in the Sun, features unknown to the Sun may exist. Speculation has arisen that some very soft X-rays may be formed in accretion shocks near the stellar surface (e.g., Kastner et al. 2002). Herbig-Haro objects far from the forming stars have been detected to be very soft X-ray emitters due to shock interactions with the interstellar medium (e.g., Pravdo et al. 2001; Bally et al. 2003; Güdel et al. 2005), but new spectral phenomenology suggests that appreciable X-ray emission may also be formed close to the stars, at the (spatially unresolved) base of the jets otherwise seen in radio and in the optical (Güdel et al. 2005). Whatever the analogy to the Sun, we also note that the level of the X-ray luminosity (between a few times  $10^{28}$  erg s<sup>-1</sup> and  $10^{31}$  erg s<sup>-1</sup> for low-mass pre-main sequence stars) and the measured plasma temperatures of up to 100 MK are extreme and may imply heating mechanisms related to continuous flaring (Güdel et al. 2003; Wolk et al. 2005).

Studies of X-ray radiation in pre-main sequence stars are thus crucial for our further understanding of star formation, accretion processes, angular momentum transport, and mass outflow mechanisms in the earliest stages of a star's life. For recent reviews of X-rays from pre-main sequence stars, we refer the reader to Feigelson & Montmerle (1999), Güdel (2004), and Feigelson et al. (2007). *XMM-Newton* and *Chandra*

penetrate dense molecular envelopes providing unequalled information on high-energy processes. A recent, large survey of the Orion Nebula Cluster (ONC) with *Chandra* (the COUP project, Getman et al. 2005) has monitored approximately 1400 young stars in a  $17' \times 17'$  field around the Trapezium. This study has provided unprecedented information on the ONC stellar population and the evolution of X-ray production. For example, Preibisch et al. (2005) concluded that a significant correlation between rotation and X-ray luminosity consistently found in main-sequence stars is absent in the ONC sample, arguing for a different magnetic dynamo process or a very different internal structure of these youngest stars. X-ray radiation was found to be largely variable, with numerous flares contributing significantly to the overall X-ray production (Wolk et al. 2005). We refer the interested reader to a special volume of papers from this study (Getman et al. 2005 and associated papers), or to a summary provided by Feigelson et al. (2007).

The ONC represents the so-called “clustered mode” of star formation in which hundreds to thousands of low-mass stars form around massive O-type stars such as those in the Trapezium. Such environments produce extreme stellar densities, and the presence of hot stars has a decisive impact on star formation. UV radiation and the winds of the central O stars have a deteriorating effect on the molecular environment, the pre-stellar cores and the circumstellar disks by ionizing and dispersing the molecular material, prematurely altering the accretion environment (O'Dell 1998) and eventually halting the star formation process altogether (Dolan & Mathieu 2001). The ionized H II Orion Nebula is the most vivid expression of this environment.

Star formation occurs, however, also in relatively unspectacular low-mass star forming regions devoid of the influence of radiation and outflow-related effects from luminous O stars. These so-called “dark clouds” typically form a smaller number of stars and often do so in a “dispersed” or “isolated mode” in which stars or very small groups of stars form in relative isolation. Stars form in a similar manner also in high-mass star formation regions, outside the clusters and at larger distances from the central OB stars (Carpenter 2000).

The closest and best-studied large, low-mass star formation region is the Taurus Molecular Cloud (TMC) complex that extends to the adjacent Auriga Clouds. TMC has served as a testbed for low-mass star-formation theory and has provided some of the best examples of phenomenology relevant to star formation. There are only a few Herbig Ae/Be stars in this vast region (V892 Tau and AB Aur in our survey). The rest are predominantly sub-solar in mass although there have been claims that a substantial population of non-accreting B stars may be members of the Taurus population (Walter & Boyd 1991), in agreement with expectations from the initial mass function. Owing to the near-absence of massive stars and their mechanical/radiative effects, the molecular gas is still present in large amounts ( $M_{\text{cloud}} \approx 3 \times 10^4 M_{\odot}$ , Ungerechts & Thaddeus 1987) much in excess of the total mass in stars already formed ( $M_{\text{stars}}$  of order  $2 \times 10^2 M_{\odot}$ ).

The Taurus region is quite large (some 10–15 degrees in diameter, corresponding to about 25–35 pc at a distance of 140 pc), making comprehensive studies of the entire population difficult. We have initiated a large X-ray study of the TMC with the *XMM-Newton* X-ray observatory, concentrating on the denser cloud areas that contain the majority of the TMC stellar population. The emphasis of this *XMM-Newton Extended Survey of the Taurus Molecular Cloud* (XEST henceforth) is on a wide-field sampling rather than on long exposures. It goes

systematically deeper by about an order of magnitude than previous Taurus X-ray surveys. While it is less sensitive than the COUP survey, it surveys an area about 60 times larger ( $\approx 5$  square degrees), which is required because of the low surface density of Taurus members. The survey is accompanied by a deep, large-field optical survey with the Canada-France-Hawaii Telescope (CFHT) and a mid-infrared survey of the entire cloud complex with the *Spitzer Space Telescope* (Padgett et al. 2007). These latter surveys will be presented separately (see summary of all three surveys in Güdel et al. 2007a). All studies combined provide an unsurpassed database for the nearest major star-forming cloud complex.

The purpose of the present paper is to give an introduction to the XEST project, to present an overview of the surveyed stellar sample, and to discuss the strategies followed during the data reduction and analysis. The detailed results are described in a series of associated papers.

The outline of this paper is as follows. We discuss some pivotal properties of the TMC in Sect. 2, including results from previous studies that support the key role of this region for further star-formation studies. Section 3 introduces the TMC X-ray survey, and Sect. 4 presents the data reduction and analysis strategies. Section 5 contains an overview of the basic results, presenting detected and undetected stellar populations, accompanied by comprehensive tabulations of fundamental properties of all surveyed objects and spectral X-ray results from the detections. We summarize overall statistical properties of XEST in Sect. 6.

The subsequent series of papers related to this survey discusses X-ray properties in the context of stellar class and accretion (Telleschi et al. 2007a), rotation-activity relations (Briggs et al. 2007), a dedicated study of the L1495E subsample (Silva et al., in preparation), correlated behavior of X-rays and optical/ultraviolet emission (Audard et al. 2007), X-rays from jet-driving stars in TMC (Güdel et al. 2007b), high-resolution X-ray spectroscopy of classical and weak-line T Tau stars (Telleschi et al. 2007b), X-rays from brown dwarfs (Grosso et al. 2007a), a *U*-band survey of brown dwarfs with the *XMM-Newton* Optical Monitor (Grosso et al. 2007b), light-curve variability studies (Stelzer et al. 2007), interpretation of X-ray flares (Franciosini et al. 2007), an investigation of statistical fluctuations in X-ray light curves (Arzner et al. 2007a), an investigation of spectral parameters in extremely faint sources (Arzner et al. 2007b), an analysis of the gas-to-dust ratio in IRAS 04158+2805 (Glaser et al. 2007), a study of the gas-to-dust ratios in the TMC in general (Glaser et al., in preparation), a case study of the accreting, prototypical CTTS T Tau (Güdel et al. 2007c), a study of the supposedly single Herbig star in TMC, AB Aurigae (Telleschi et al. 2007c), and a search for new TMC members based on near-infrared and X-ray properties of field sources detected in XEST (Scelsi et al. 2007).

## 2. The Taurus molecular cloud complex

At a distance around 140 pc (e.g., Loinard et al. 2005; Kenyon et al. 1994), the TMC is the nearest large star formation region and reveals characteristics that make it ideal for detailed physical studies. One of the most notable properties of TMC in this regard is its structure in which several loosely associated but otherwise rather isolated molecular cores each produce one or only a few low-mass stars, different from the much denser cores in  $\rho$  Oph or in Orion. TMC features a low stellar density of only 1–10 stars  $\text{pc}^{-3}$  (e.g., Luhman 2000). Strong mutual influence due to outflows, jets, or gravitational effects are therefore minimized. Strong stellar winds and ionizing UV radiation are mostly absent

in TMC because there are no O stars and only very few B and A stars (Walter & Boyd 1991). Further, most stars in TMC are subject to relatively modest extinction, providing access to a broad spectrum of stars at all evolutionary stages from Class 0 sources to near-zero age main-sequence T Tau stars. TMC has also become of central interest for the study of substellar objects, in particular brown dwarfs, with regard to their evolutionary history and their spatial distribution and dispersal (Briceño et al. 2002; Guieu et al. 2006).

TMC has figured prominently in star-formation studies at all wavelengths. It has provided the best-characterized sample of classical and weak-line T Tau stars (CTTS and WTTS, respectively, or “Class II” and “Class III” objects in the infrared classification - Kenyon & Hartmann 1995); most of our current picture of low-density star formation is indeed based on IRAS studies of Taurus (Strom et al. 1989; Weaver & Jones 1992). Among the key results from TMC studies as listed in Kenyon & Hartmann (1995) figure the following: i) more than 50% of the TMC objects have IR excess beyond the photospheric contribution, correlating with other activity indicators ( $H\alpha$ , UV excess etc.) and indicating the presence of warm circumstellar material predominantly in the form of envelopes for Class I protostars and circumstellar disks for Class II stars. ii) Class III sources (mostly to be identified with WTTS) are distinctly different from Class I-II objects by not revealing optically thick disks or signatures of accretion. iii) Star formation has been ongoing at a similar level during the past 1–2 Myr, with the Class-I protostars having ages of typically 0.1–0.2 Myr. iv) There is clear support for an evolutionary sequence Class I→II→III, although there is little luminosity evolution along this sequence, indicating different evolutionary speeds for different objects (see also Hartmann 2002). The infall time scale is a few times  $10^5$  yrs, while the disk phase amounts to a few times  $10^6$  yrs. An evolutionary scenario is also suggested from the different spatial distribution of Class 0/I stars vs. CTTS/WTTS with respect to the gas distribution: the former classes are still within the boundaries of the high density gas, while the latter are found in regions of lower density (Palla & Stahler 2001).

TMC has also been well-studied at millimeter wavelengths. This region has better high-resolution molecular line maps than any other star-forming region (Onishi et al. 2002). Most of the higher-mass CTTS and Class I protostars have been surveyed by millimeter interferometers for molecular line emission from disks (Dutrey et al. 1996; Ohashi et al. 1996, etc.) and many detailed studies of individual sources have been published (e.g., Qi et al. 2003; Duchêne et al. 2003). A variety of millimeter continuum observations of these sources have enabled studies of cold disk frequency into substellar mass ranges and raised the possibility of grain growth within circumstellar disks (Scholz et al. 2006; Rodman et al. 2006; Wolf et al. 2003; Kitamura et al. 2002, etc.). Due to its proximity and easy accessibility to northern hemisphere radio telescopes, the TMC has served as a template for millimeter studies of young stellar objects.

Although TMC has been regarded, together with the  $\rho$  Oph dark cloud, as the prototypical low-mass star-forming region, a few apparent peculiarities deserve to be mentioned. TMC contains an anomalous number of binaries (Ghez et al. 1993; Duchêne 1999), compared with other SFRs (e.g., Orion) or with field stars. In TMC, about two thirds of all members are bound in multiple systems, with an average separation of about  $0.3''$  (e.g., Leinert et al. 1993; Mathieu 1994; Simon et al. 1995; Duchêne et al. 1999; White & Ghez 2001; Hartigan & Kenyon 2003). Petr et al. (1998) reported, at the 96% confidence level, a three times higher binary occurrence in TMC compared to Orion for

component separations in the range of 63–225 AU. Also, TMC cloud cores are comparatively small and of low mass, at least when compared with cores in Orion or Perseus (Kun 1998).

Initially, TMC was also found to be deficient of lowest-mass stars and brown dwarfs, with a mass distribution significantly enriched in 0.5–1  $M_{\odot}$  stars, compared to Orion samples (Luhman 2000; Briceño et al. 2002). It was speculated that the formation of brown dwarfs could be different in the low-density environment of TMC compared to the dense packing of stars in Orion. However, a new wide-field search for low-mass TMC members now indicates that there is no BD deficit in Taurus (Guiou et al. 2006).

In X-rays, Taurus has again played a key role in our understanding of high-energy processes and circumstellar magnetic fields around pre-main sequence stars. Taurus X-ray studies provided the first detailed view of the X-ray behavior of T Tauri stars (Feigelson & Kriss 1981; Feigelson & DeCampli 1981; Walter & Kuhi 1981). Among the key surveys of the entire region are those by Feigelson et al. (1987), Walter et al. (1988), Bouvier (1990), Strom et al. (1990), Damiani & Micela (1995), Damiani et al. (1995), based on *Einstein Observatory* observations, and the work by Strom & Strom (1994), Neuhäuser et al. (1995), and Stelzer & Neuhäuser (2001) based on *ROSAT*. These surveys have characterized the overall luminosity behavior of TTS and studied the dependence of X-ray activity on rotation. Bally et al. (2003) and Favata et al. (2003) reported, respectively, *Chandra* and *XMM-Newton* studies of the L1551 cloud region (the longer exposure from *XMM-Newton* will be included in our survey; the *Chandra* exposure contains no additional sources).

But again, for reasons very poorly understood, TMC differs from other SFRs significantly also with regard to X-ray properties. Whereas no X-ray activity-rotation correlation (analogous to that in main-sequence stars) is found for samples in the Orion star-forming regions, perhaps suggesting that all stars are in a saturated state (Flaccomio et al. 2003a; Preibisch et al. 2005), the X-ray activity in TMC stars has been reported to decrease for increasing rotation period (e.g., Neuhäuser et al. 1995; Damiani & Micela 1995; Stelzer & Neuhäuser 2001). Also, claims have been made that the X-ray behavior of TMC CTTS and WTTS is significantly different, CTTS being less luminous than WTTS (Strom & Strom 1994; Damiani et al. 1995; Neuhäuser et al. 1995; Stelzer & Neuhäuser 2001). This contrasts with other star-forming regions (Flaccomio et al. 2000; Preibisch & Zinnecker 2001), but recent reports reveal a similar segregation also for Orion and some other SFRs (Flaccomio et al. 2003b; Preibisch et al. 2005). Some of these discrepancies may be due to selection and detection bias (e.g., WTTS are predominantly identified in X-ray studies, in contrast to CTTS), but also to incomplete samples due to high detection limits. Issues related to rotation and selection biases will be discussed in detail in one of the companion papers, see Briggs et al. (2007).

### 3. The XEST project

#### 3.1. Scientific goals

The scientific goals of XEST are:

- To collect X-ray spectra and light curves from a statistically meaningful sample of TMC objects, and to characterize them in terms of X-ray emission measure distributions, temperatures, X-ray luminosities, and variability.
- To interpret X-ray emission in the context of other stellar properties such as rotation, mass, and radius.

- To investigate changes in the X-ray behavior as a young stellar object evolves.
- To obtain a census of X-ray emitting objects at the stellar mass limit and in the substellar regime (brown dwarfs = BDs).
- To study in what sense the stellar environment (circumstellar disks, jets, accretion) influences the X-ray production, and vice versa.
- To study the gas-to-dust ratio in the circumstellar environment, making use of extinction measurements obtained in the optical or near-infrared.
- To assess the role of flares in coronal heating, and to study flare characteristics in their own right.
- To search for new, hitherto unrecognized, TMC members.

The outstanding characteristics of the survey are its sensitivity and its energy resolution, covering a large fraction of the most densely populated regions of the Taurus clouds. It is by far the largest-area Taurus survey at this sensitivity. It reaches sensitivities about ten times better than previous surveys such as those conducted with *ROSAT* (Neuhäuser et al. 1995; Stelzer & Neuhäuser 2001) and therefore permits a systematic study of the lowest-mass Taurus members. Because of the harder band used for the XEST project (compared to *ROSAT*'s 0.1–2.4 keV band), XEST also detects several deeply embedded Taurus protostars systematically for the first time, while they remained undetected in softer-band surveys. Given the achieved sensitivity, the XEST project accesses the known TMC population in the surveyed fields nearly completely, thus suppressing potential bias that previous surveys may have been subject to. Energy resolution permits a detailed description of the plasma properties together with the measurement of the absorbing gas columns that are located predominantly in the TMC clouds themselves, and even in the immediate circumstellar environment in the case of strongly absorbed objects.

#### 3.2. Instruments and exposures

XEST is a wide-field X-ray survey obtained with the *XMM-Newton* X-ray observatory, principally based on combined CCD camera exposures, but complemented by exposures of a few bright targets with the reflection grating spectrometers and fields observed with the optical monitor. A few complementary observations obtained with the *Chandra X-Ray Observatory* have also been included.

*XMM-Newton* (Jansen et al. 2001) orbits the Earth in a 48 hr orbit, permitting long, uninterrupted exposures. It carries three high-throughput telescopes feeding two suites of X-ray instruments, and also features an optical telescope. A short description of the instruments and the chosen setups follows.<sup>1</sup>

i) The *European Photon Imaging Cameras* (EPICs) are three CCD-based X-ray cameras, one per telescope, that operate entirely independently. Two cameras are of the MOS type (Turner et al. 2001), and one is of the PN type (Strüder et al. 2001). The circular field of view of each camera has a diameter of 30', and the three telescopes are nearly co-aligned. The PN camera provides most counts because the beams to the MOS cameras are intersected by the reflection gratings that consume approximately half of the X-ray flux. The MOS PSF has a full-width-at-half-maximum (FWHM) of about 4–5'', while the half-energy width (HEW) is larger, amounting to some 13–15''. The cameras provide pixel sizes of 1.1'' (MOS) or 4'' (PN). The energy

<sup>1</sup> For further details, see *XMM-Newton* User's Handbook (<http://xmm.vilspa.esa.es/>)

resolution is approximately  $E/\Delta E \approx 45$  at 6.7 keV, scaling as  $E^{-1/2}$ . To prevent the spectra from being altered by optical load, various filters can be inserted.

We usually applied the medium filter to all EPIC cameras, providing a combined on-axis effective area for all three cameras of  $\approx 1800 \text{ cm}^2$  at 1.5 keV. Exceptions were exposure XEST-26 for which the thick filter was used for the two MOS cameras to suppress optical load from the Herbig star AB Aur (the PN camera was not recording), and exposures XEST-27 and XEST-28 for which the thick filter was used for each EPIC camera. All cameras were operated in full window mode, i.e., the entire field of view was exposed, confining the time resolution to 2.6 s for the MOS and to 73 ms for the PN camera, entirely appropriate for the moderate brightness of our sources. The only exceptions here were the fields around V773 Tau (XEST-20) where MOS2 was operated in the small-window mode, and the field of T Tau (XEST-01) for which MOS1 was operated in small-window mode, and MOS2 was operated in large-window mode. These modes confine the central CCD to  $100 \times 100$  and  $300 \times 300$  pixels, respectively, compared to  $600 \times 600$  pixels for the full window mode, but avoiding pile-up during potential large flares.

ii) Two *Reflection Grating Spectrometers* (RGS, den Herder et al. 2001) are each fed by about 50% of the light of one of the telescopes. The dispersed spectra comprise the wavelength range from 5 to 35 Å, although in each spectrometer, one of the nine chips has failed earlier in the mission, leaving a gap in the 10.6–13.8 Å range for RGS1 and in the 20.0–24.1 Å range for RGS2. The first-order spectral resolution is approximately 60–70 mÅ (*FWHM*) at any wavelength, corresponding to  $\lambda/\Delta\lambda \approx 300$  at  $\lambda = 20 \text{ Å}$ . The combined first-order effective area is  $\approx 115 \text{ cm}^2$  at 15 Å.

iii) The *Optical Monitor* (OM, Mason et al. 2001) is a microchannel-plate enhanced CCD camera fed by a 30 cm mirror co-aligned with the EPIC cameras. Its square field of view is, however, somewhat smaller than the latter's, amounting to 17' side-length, i.e., a radius of 12' along the diagonal. The OM can be used in combination with various filters. Filter use may be dictated by the brightness of the stars in the field of view. We obtained most of the exposures with the *U* band filter inserted, given the diagnostic importance of this band for both chromospheric flare processes and accretion mechanisms. In a few cases, another near-UV filter was used (*UVW1*: 2500–3500 Å; *UVW2*: 1700–2500 Å) to suppress excessive optical load. The window mode we applied essentially observed the entire field of view. The detector area is split up into a series of windows, some of which are sequentially exposed and read out. The exposure time for one of these frames determines the time resolution for any source within the window. This time resolution is of order 1000–2000 s but varies within any given XEST OM field, and is also not identical between different XEST fields. A detailed presentation is given by Audard et al. (2007). Additionally, if a bright source was present on-axis, the central  $10.5'' \times 10.5''$  of the OM field were exposed within a high-time resolution window that collects photons with a time resolution of 0.5 s.

Our initial project collected 19 fields in the TMC coherently with exposure times of about 30 ks each. The effective exposure time varied between 31.2 ks and 41.9 ks. Relevant information is given in Table 1. The start and end times of the observations as listed in Table 1 are the times of the earliest and latest recording of any of the three EPIC cameras, and the quoted exposure time is the difference between end and start times. The pointing coordinates are the nominal boresight coordinates, which are not identical with the coordinates of the center of the EPIC fields

of view, the latter being slightly misaligned (e.g., to avoid the PN CCD edges crossing the exact field center). The last column gives the filter used for the OM (*U* band, or one of the ultraviolet filters, *UVW1* or *UVW2*). A letter *F* indicates that the central star was recorded in fast mode (see Audard et al. 2007).

The fields of view were selected such that they cover the densest concentrations of CO gas, which also show the strongest accumulations of TMC stellar and substellar members. We complemented this sample with exposures that were obtained as part of separate projects, partly retrieved from the *XMM-Newton* archive (marked in Table 1); in all of these observations very similar instrument setups were used, although in most cases the exposure times were longer and the strategies for the use of the OM were different. The eight additional fields are: 1) L1495 centered at V410 Tau (XEST exposures 23 and 24, see Table 1; two exposures totaling 117.9 ks; PI F. Walter); 2) L1551, centered near L1551 IRS-5 (XEST exposure 22, 56.9 ks, PI F. Favata); 3) a field centered at AA Tau (eight exposures of 16–19.5 ks each of which we use only one [XEST exposure 25] that reveals the strongly variable AA Tau X-ray source in an average state – details in Grosso et al., in preparation; PI J. Bouvier); 4) a field centered at HD 283572 (XEST exposure 21, 46.2 ks, PI R. Pallavicini); 5) a field around BP Tau (XEST exposure 28, 131.3 ks, PI J. Schmitt); 6) a field around SU Aur (XEST exposure 26, 129.6 ks, PI R. Pallavicini); 7) a field around  $\zeta$  Per (XEST exposure 27, 43.4 ks, PI W. Waldron); 8) a field around T Tau (XEST exposure 1, 83.0 ks – a special discussion of this observation is given by Güdel et al. (2007c); PI M. Güdel). The entire survey includes approximately 5 sq. degrees in total, with a total exposure time of about 1.3 Ms. The spatial coverage is illustrated in Fig. 1.

We have included a few exposures obtained by *Chandra* that contain important objects outside the *XMM-Newton* coverage, or faint objects not detected by *XMM-Newton* but detected by *Chandra*. Objects that were covered by the XEST program (regardless of their detection status) and were *not* detected in these *Chandra* exposures will not be reported in our survey tables. The additional fields are defined in Table 2. Note that different detectors were used for these observations. The entire fields of view were scanned for TMC objects that add to the project, but no systematic survey of all X-ray sources in the fields was performed. The exposure around GV Tau does not contain new sources beyond those detected in the *XMM-Newton* field but provides better angular resolution for the study of the origin of the X-rays in this binary system (Güdel et al. 2007b). We also add the strong X-ray detection coincident with HD 28867 = HR 1442 that has been suspected to be a late-B (non-Herbig) binary member of the TMC (e.g., Walter & Boyd 1991). This object has been extensively studied by Walter et al. (2003) who found evidence for a G-type companion to one of the B-type components.

The XEST work presented here and in the accompanying papers addresses X-ray properties of the commonly known stellar and substellar population. A considerable number of hitherto unrecognized Taurus members may be present in the survey, e.g., deeply embedded sources or extincted stars on the far side of the clouds. An attempt to identify some objects of this population has been undertaken by Scelsi et al. (2007).

#### 4. Strategy and analysis of the TMC survey

Our *XMM-Newton* data analysis procedure predominantly uses tasks provided in the *XMM-Newton* Science Analysis Software (SAS) v6.1.0, augmented by procedures in FTOOLS, and is tied together in Perl scripts developed at the Paul Scherrer

**Table 1.** Observing log of XEST program.

Exposure #	ObsID <sup>a</sup>	RA(J2000.0) <sup>b</sup> h m s	$\delta$ (J2000.0) <sup>b</sup> deg ' "	Start time <sup>c</sup> y-m-d h:m:s	Stop time <sup>c</sup> y-m-d h:m:s	Exposure time (s)	OM mode, OM filter <sup>d</sup>
1 <sup>e</sup>	0301500101	04 21 59.4	19 32 06	2005-08-15 13:52:13	2005-08-16 12:55:22	82989	F, UVW1
2	0203540201	04 27 19.6	26 09 25	2004-08-17 06:08:10	2004-08-17 17:32:46	41076	U
3	0203540301	04 32 18.9	24 22 28	2004-08-22 06:45:22	2004-08-22 16:37:04	35502	F, U
4	0203540401	04 33 34.4	24 21 08	2005-02-21 01:18:21	2005-02-21 10:31:17	33176	F, U
5	0203540501	04 39 34.9	25 41 46	2005-02-21 11:25:51	2005-02-21 20:13:47	31676	U
6	0203540601	04 04 42.9	26 18 56	2004-08-25 20:53:54	2004-08-26 05:41:50	31676	U
7	0203540701	04 41 12.5	25 46 37	2005-02-24 20:28:00	2005-02-25 05:15:58	31678	U
8	0203540801	04 35 52.9	22 54 23	2004-08-26 06:36:23	2004-08-26 18:10:59	41676	F, U
9	0203540901	04 35 55.1	22 39 24	2005-02-25 08:38:53	2005-02-25 17:26:50	31677	U
10	0203542201	04 42 20.9	25 20 35	2005-03-05 05:56:38	2005-03-05 14:44:30	31672	F, U
11	0203541101	04 21 51.1	26 57 33	2004-08-18 06:44:13	2004-08-18 18:21:54	41861	U
12	0203542101	04 35 17.4	24 15 00	2005-03-04 20:22:29	2005-03-05 05:01:54	31165	F, U
13	0203541301	04 29 52.0	24 36 47	2004-08-25 11:11:23	2004-08-25 19:59:19	31676	U
14	0203541401	04 30 30.6	26 02 14	2005-02-09 03:01:02	2005-02-09 12:30:38	34176	U
15	0203541501	04 29 42.4	26 32 51	2005-02-09 13:12:40	2005-02-09 22:38:18	33938	F, U
16	0203541601	04 19 43.0	27 13 34	2004-08-21 05:53:24	2004-08-21 17:11:22	40678	–
17	0203541701	04 33 21.2	22 52 41	2005-02-11 01:19:56	2005-02-11 10:07:50	31674	U
18	0203541801	04 33 54.7	26 13 28	2004-08-13 21:48:05	2004-08-14 06:36:03	31678	F, U
19	0203541901	04 32 43.0	25 52 32	2004-08-14 07:18:05	2004-08-14 17:50:41	37956	F, U
20	0203542001	04 14 12.9	28 12 12	2004-09-12 07:04:43	2004-09-12 15:52:37	31674	F, U
21 <sup>e</sup>	0101440701	04 21 59.0	28 18 08	2000-09-05 02:57:44	2000-09-05 15:47:55	46211	–
22 <sup>e</sup>	0109060301	04 31 39.0	18 10 00	2000-09-09 18:29:39	2000-09-10 10:18:12	56913	UVW2
23 <sup>e</sup>	0086360301	04 18 31.2	28 27 16	2001-03-11 12:46:45	2001-03-12 09:13:24	73599	(UVW2) <sup>f</sup>
24 <sup>e</sup>	0086360401	04 18 31.2	28 27 16	2001-03-12 09:29:38	2001-03-12 21:47:51	44293	(UVW2) <sup>f</sup>
25 <sup>e</sup>	0152680201 <sup>g</sup>	04 34 55.5	24 28 54	2003-02-14 02:18:48	2003-02-14 07:01:32	16964	UVW2
26 <sup>e</sup>	0101440801	04 55 59.0	30 34 02	2001-09-21 01:34:17	2001-09-22 13:34:31	129614	–
27 <sup>e</sup>	0201550201	03 54 07.9	31 53 01	2004-02-13 21:45:10	2004-02-14 09:48:07	43377	–
28 <sup>e</sup>	0200370101	04 19 15.8	29 06 27	2004-08-15 06:14:30	2004-08-16 18:42:57	131307	UVW1

<sup>a</sup> XMM-Newton observation identification number.<sup>b</sup> Nominal boresight coordinates.<sup>c</sup> Earliest respectively latest time of recording for any EPIC camera is reported.<sup>d</sup> F = fast + imaging mode (only imaging otherwise); for filter name, see XMM-Newton User's Handbook (<http://xmm.vilspa.esa.es/>).<sup>e</sup> Observation from separate projects.<sup>f</sup> The OM was recording only for short time intervals.<sup>g</sup> Eight different observations of  $\approx 16$ – $19$  ks (PN) each around AA Tau, 2003-02-14 – 2003-02-28, ObsID = 0152680201-901.**Table 2.** Complementary *Chandra* observations.

Exposure #	Main Target	Obs ID <sup>a</sup>	RA(J2000.0) <sup>b</sup> h m s	$\delta$ (J2000.0) <sup>b</sup> deg ' "	Start time y-m-d h:m:s	Stop time y-m-d h:m:s	Exposure time (s)	Instr.
C1	V410 Tau	3364	04 18 34.6	28 22 47	2002-03-07 06:16:32	2002-03-07 11:45:24	17734	ACIS-S
C2	FS Tau	4488	04 22 00.2	26 58 07	2003-11-08 12:57:58	2003-11-08 21:56:34	29674	ACIS-S
C3	DG Tau	4487	04 27 02.3	26 04 56	2004-01-11 02:58:51	2004-01-11 11:52:21	29717	ACIS-S
C4	GV Tau	4498	04 29 23.3	24 32 44	2003-12-28 08:56:57	2003-12-28 16:37:59	24650	ACIS-I
C5	HR 1442	612	04 33 33.1	18 01 15	2000-09-11 17:53:33	2000-09-11 19:41:57	4679	HRC-I
C6	L1527	2563	04 39 52.7	26 03 05	2002-12-06 08:30:12	2002-12-06 14:21:15	19317	ACIS-I

<sup>a</sup> *Chandra* observation identification number. <sup>b</sup> Nominal boresight coordinates.


Institut. Source identification was based on procedures involving maximum-likelihood algorithms and wavelet analysis. In order to optimize detection of faint sources, periods of high background radiation levels due to local particles were cut out. All programs named below can be assumed to be SAS tasks unless otherwise noted.

#### 4.1. Creation of event lists and event selection

An event list was produced for each active EPIC instrument from the original data files using the standard SAS procedures EPCHAIN and EMCHAIN. MOS event lists were filtered to include only events flagged as ‘good’ and with patterns 0–12 (Kirsch 2006). PN event lists were filtered using the selection

expression (FLAG & 0xfb0825) = 0, which rejects events with patterns >12, close to CCD windows or on-board bad pixels, in spoiled frames, or outside the field of view, but retains events on or next to offset columns, close to bright pixels or dead pixels. Remaining bright pixels, columns and partial columns observed in images at energies >300 eV were further added to the bad pixel list and rejected.

Source detection was performed in three energy bands: soft (500–2000 eV), hard (2001–7300 eV) and full (500–7300 eV). These were chosen with consideration of the energy-dependent sensitivity of the EPIC instruments, the expected spectra of our sources of interest, the energies of strong fluorescent spectral features from the detectors (particularly Ni, Cu and Zn  $K\alpha$  at energies 7.3–9 keV), and compatibility of count-rates and fluxes



**Fig. 1.** Map of the TMC region (oriented in RA - dec, north is up, east to the left). The grayscale background map is an extinction ( $A_V$ ) map from Dobashi et al. (2005). Contours show the CO emission (Dame et al. 1987). The large (0.5 degrees diameter) circles show the fields of view of the *XMM-Newton* survey (the dashed circles marking fields from separate projects also used for the survey). Small white dots mark the positions of individual young TMC members. The labels correspond to the XEST exposure numbers in Table 1. Note the outlying *XMM-Newton* fields XEST-26 around SU Aur (NE corner), XEST-27 around  $\zeta$  Per (NW corner), XEST-01 around T Tau, and XEST-22 around L1551 (the latter two at the southern border of the figure).

derived therefrom with those quoted using previous and current instruments for X-ray astronomy. We give an outline of the source detection procedure here and describe each stage in more detail below.

For each EPIC instrument and energy band:

- Define good time intervals that reject times of high background rate.

- Extract an image using good time intervals.

- Generate an exposure map.

- Create a background map.

For each band:

- Mosaic the images of the (up to) three instruments to use the full sensitivity of the combined EPIC detectors.

- Mosaic the background maps.
- Scale the exposure maps to account for the differences in sensitivity and mosaic.
- Locate candidate sources using a wavelet transform algorithm.

Finally:

- Match candidate sources from all bands and concatenate into a single list.
- Parameterize candidate sources in all bands simultaneously via maximum-likelihood fitting of the point-spread function and reject those with low detection significance.
- Verify by eye and reject obvious spurious detections.
- Calculate upper limits in bands where each source is undetected.
- Calculate and correct any systematic offset of the X-ray source positions from the positions of near-infrared counterparts.

#### 4.2. Definition of good time intervals

In almost every observation the background count-rate was highly variable, by factors up to several hundred<sup>2</sup>. To optimize sensitivity to the detection of faint sources, it was necessary to exclude intervals of highest background count-rate. Good time intervals (GTIs) were defined independently for each instrument and energy band as follows.

A light curve was extracted of the whole active detector area (excluding bright sources, as described in the creation of background maps, below) with a binning of 52 s, scaled to a 1 sq. arcsec area, and ordered by increasing count-rate. Its cumulative distribution was made and the source count-rate required to give a  $4\sigma$  detection in a circular region of  $15 \text{ arcsec}^3$  was calculated for each bin, and the minimum was found. This defined the maximum background count-rate accepted, and hence the good time intervals (see Fig. 2). In some observations, the source count-rate required for a  $4\sigma$  detection is a factor three higher than would be expected for a 30-ks observation performed with background at the quiescent level. The total accepted good exposure times for the full energy band for each instrument in each observation are listed in Table 3.

#### 4.3. Extraction of images


An image was created for each instrument and each energy band, in sky coordinates with pixel size  $2'' \times 2''$ , using data in good time intervals only. The intrinsic pixel sizes of the PN and MOS detectors are  $4''$  and  $1.1''$ , respectively. In processing, a randomization of the event position within a pixel is performed to avoid fixed pattern noise.  $2''$  was simply found to provide much better separation of close sources than  $4''$ .

For each energy band, the images from the three detectors were mosaicked to use the full sensitivity of the combined EPIC cameras. The relative astrometry of the EPIC detectors is less than  $1.5''$ , so we applied no offset between the images.

An exposure map was generated for each image using EEXPMAP. A summed exposure map was also produced for each energy band, but first MOS exposure maps were scaled

<sup>2</sup> The EPIC background is described at [http://xmm.vilspa.esa.es/external/xmm\\_sw\\_cal/background/index.shtml](http://xmm.vilspa.esa.es/external/xmm_sw_cal/background/index.shtml); see also Read & Ponman (2003).

<sup>3</sup> This encloses approximately 68 per cent of source counts and is the area used by the SAS source parameterization task EMLDETECT.



**Fig. 2.** The selection of good time intervals to reject times affected by high background and so optimize sensitivity to the detection of faint sources. The background count rate per unit area in the 0.5–7.3 keV energy band is shown in the top panel. The expected quiescent background level in this band is marked by the green dot-dashed lines. The bottom panel shows the background count-rate sorted in ascending order. The middle panel shows as a function of accumulated good exposure time the minimum source count-rate that will give a  $4\sigma$  detection against the accumulated background in a circular source extraction region of radius  $15''$ , containing 67 per cent of the source counts. The minimum defines the amount of accepted good time, noted in the bottom panel in green in seconds and as a percentage of the total exposure time, and the maximum acceptable background rate, marked in the upper panel by a green solid line. The time bins marked in red in the upper panel are rejected. XEST-16, shown here, was one of the observations worst-affected by high background, but time intervals with background levels more than 100 times the quiescent level occurred in most observations.

to account for the lower sensitivity compared with the PN. The scaling factors used were 0.370, 0.345 and 0.385 in the full, soft and hard bands, respectively: these factors are strictly dependent on the individual source spectrum; the values used are median values found for all sources detected in the first 12 analyzed observations. The count rates resulting from our source detection are therefore approximate PN-equivalent count rates.

#### 4.4. Creation of background maps

A background map was created for each image: each instrument and energy band. The basic method was to mask out regions of the image affected by source counts and apply a 2D Gaussian smoothing filter of  $\sigma = 60''$ . The background map was produced in three iterations which successively improved



the source parameterization and hence the mask. We first used the SAS wavelet transform algorithm EWAVELET to locate and approximately parameterize sources. The resultant source list and background map were input into the maximum-likelihood fitting algorithm EMLDETECT to produce an improved background which was input into a second run of EMLDETECT to produce the final background map. The mask, produced using REGION, masked out each source with an elliptical area, whose eccentricity reflected the shape of the point-spread function at the source position and whose size reflected the number of source counts. Smoothing was performed using ASMOOTH. EMLDETECT outputs an image of the parameterized sources, which was masked and smoothed in the same way as the image and subtracted from the background to remove the effect of remaining source counts in the background map. Out-of-time events (OOTEs) in PN images were modeled into the background map. An image of OOTEs was created<sup>4</sup> and smoothed only along the CCD readout direction, subtracted from the true image before source detection and smoothing, and added to the resulting background map. Where a central MOS CCD observed a bright source in a window mode such that no reliable background level could be assessed from this CCD, the background in that CCD was estimated as 1.15 times the mean background in the other CCDs, as measured from MOS observations in full frame mode lacking a bright central source. For each energy band, we mosaicked the background maps from all EPIC detectors.

#### 4.5. Location of source candidates

Source candidates in the mosaicked EPIC image in each energy band were located using two wavelet transform algorithms: the SAS task EWAVELET, and PWXDETECT developed at the Osservatorio Astronomico di Palermo (Damiani et al. 1997). In PWXDETECT the wavelet transform (WT) is obtained from the rate image (image/exposure map)  $r(x, y)$  as:

$$w(x, y; a) = \int \int g\left(\frac{x-x'}{a}, \frac{y-y'}{a}\right) r(x', y') dx' dy'. \quad (1)$$

The WT is evaluated at different values of the scale parameter,  $a$ , of the generating wavelet<sup>5</sup>, going from 2.8'' to 16'' with a logarithmic step of  $\sqrt{2}$ . This property makes this multiscale WT particularly suited when the PSF varies across the detector image, as well as very effective in detecting extended sources: for the case of the EPIC images, point-like sources are detected typically at  $a = 8''$  (although faint sources with undersampled PSF may be detected either at low and at high detection scale), while extended sources typically result with  $a = 16''$ .

Since sources are defined as local maxima in the WT space, thresholds for detection at a given confidence level must be assessed for each field and each band. This is obtained by means of extensive simulations on summed PN and MOS images containing only background photons. The number of background photons in each simulated PN or MOS image is evaluated from the observed image relevant to that band, instrument and field. To avoid rejecting possible source candidates, unconservative thresholds were adopted so as to give about 150 spurious detections due to background fluctuations, per band and per field.

<sup>4</sup> See [http://xmm.vilspa.esa.es/sas/new/documentation/threads/EPIC\\_0oT.html](http://xmm.vilspa.esa.es/sas/new/documentation/threads/EPIC_0oT.html)

<sup>5</sup> The wavelet function  $g$  adopted in this method is a two-dimensional ‘‘Mexican hat’’:  $g(x/a, y/a) = (2 - r^2/a^2) e^{-r^2/2a^2}$  ( $r^2 = x^2 + y^2$ ).

These simulations were also used to assess the number of false detections expected after maximum likelihood fitting of the candidate sources (see below). EWAVELET uses a similar methodology and we used a similarly low detection significance threshold of  $3.5\sigma$ . For each algorithm the source lists in the three bands were merged to create a single list input to EMLDETECT.

#### 4.6. Parameterization of detected sources

Source candidates were parameterized in the three energy bands simultaneously via maximum likelihood fitting of the spatially-varying point spread function (PSF) using EMLDETECT. The free fit parameters were the source location, fixed to be the same in all energy bands, and the count rate in each energy band.

We retained detections with maximum likelihood  $ML > 8$  in at least one band. Extensive tests of simulated blank-field EPIC images have demonstrated that we expect approximately 1.5 false detections per full-band image due to background fluctuations alone.

Detections were overplotted on the image and examined by eye. Obvious false detections, e.g. in the wings of the point spread function of bright sources, were removed and the maximum likelihood fitting repeated.

This procedure was run using the EWAVELET candidate list and the PWXDETECT candidate list. The two source lists were merged, adding 0–5 extra sources per observation, and the maximum likelihood fitting repeated.

For each source, the parameters fitted by EMLDETECT were retained in each band where the source was detected with  $ML > 5$ , else upper limits were calculated at the 95 per cent confidence level using the prescription of Kraft et al. (1991).

#### 4.7. Boresight correction

2MASS counterpart positions were used to correct X-ray source positions for systematic shifts; this boresight correction was made iteratively. First, we cross-correlated the XEST catalog and the 2MASS All-Sky Catalogue of Point Sources (Cutri et al. 2003), using a 5'' correlation radius and keeping only the minimum-distance match to compute the XEST-2MASS position offsets. The XEST positions were corrected by subtracting the median position offset. Then, we repeated this correction process, using a 3.5'' correlation radius, until the median position offset became zero, which was obtained in a few steps. The boresight corrections are small (0.4''–3.8''), and the final residual registration error between 2MASS and XEST sources is 0.9''–2.2''.

The resultant boresight correction was applied to the X-ray source positions. The RMS error on the boresight shift of each field was calculated and added in quadrature to the statistical positional error calculated for each source in that field by EMLDETECT to give a  $1\sigma$  positional error circle.

We applied the same method to OM sources using 10'' and 3'' radii. We found larger boresight corrections (1''–8'') but with residual registration errors lower than 1.6''.

#### 4.8. Candidate TMC members

To assess membership in TMC of the detected X-ray sources, we compiled a comprehensive membership catalog mostly based on the following references: Cohen & Kuhl (1979), Kenyon & Hartmann (1995), Briceño et al. (1998), White & Ghez (2001), Briceño et al. (2002), Hartmann (2002),

Hartigan & Kenyon (2003), Luhman et al. (2003), Luhman (2004), Luhman (2006), White & Hillenbrand (2004), Andrews & Williams (2005), and Guieu et al. (2006). The catalog was subsequently used to decide which X-ray sources to use for the physical interpretation, and to assess which TMC members remained undetected. The adopted coordinates (mostly from 2MASS) are described in more detail below (Sect. 5.2.1).

#### 4.9. Extraction of source spectra and light curves

Source events were extracted from circular regions centered on the EMLDETECT source positions. Although the off-axis PSF is obviously elongated, there is currently no available parameterization of an elliptical form of the PSF and the enclosed energy correction calculated by the SAS task ARFGEN in creating the ancillary response file is not energy dependent unless a circular source extraction region is used. We calculated the source extraction radius that maximized the source signal-to-noise ratio in the full energy band using: the total number of source counts derived by EMLDETECT for the full-band mosaicked EPIC image, the expected number of background counts per unit area in the source region, calculated from the full-band EPIC background map, and the circularly-symmetric EPIC PSF formulation of Ghizzardi et al. (2002). The SAS task REGION was used to calculate smaller extraction radii in cases where nearby contaminating sources existed. The source extraction radius was constrained to be between  $8''$  and  $60''$ .

Annular background extraction regions were calculated by the REGION task, with inner radii calculated such that the source count density (counts per sq. arcsec) had fallen to 30 per cent of the local background count density. The task excluded neighboring sources using the same criterion. The outer radius was simply 3 times the inner radius. For each case where a source of interest was clearly contaminated by counts from a neighboring source, the background region was individually defined to account for these, as an annulus around the contaminating source with inner and outer radii exactly enclosing the extraction region of the source of interest (with the source itself excluded with a larger circle), or as a number of circles within this annulus of identical size to the extraction region of the source of interest, chosen to avoid other nearby sources.

The same source and background regions were used for the extraction of PN and MOS products. Background products were scaled by the ratio of active detector area in the source region to that in the background region.

Light curves and spectra were extracted for all sources identified as members of Taurus–Auriga, for all other sources with more than 150 EPIC counts (see e.g. Scelsi et al. 2007), and all other sources detected by the Optical Monitor (Audard et al. 2007).

##### 4.9.1. Light curves

Light curves were extracted for each instrument in the full-band only and used all available exposure time. Time intervals affected by high background were not excluded as intrinsic source variability can still be detected in these times. Also, the background flares provide a sensitive test of the reliability of the background subtraction.

The active exposure time in each bin was calculated, accounting for intervals when the instrument (or CCD) was not live, which are frequent when the instrument experiences high

event rates during background flares. The background light curve was corrected for differences in active exposure

The time bin size was calculated such that the average number of EPIC source counts in each bin was 20. However, the minimum bin size was fixed to 100 s to produce easily-viewed plots and the maximum bin size was fixed to 1000 s, the timescale on which flares are typically observed, to avoid short bright flares on faint sources being smoothed out. The same bin size was used for PN and MOS light curves to enable simple summation to form an EPIC light curve.

The light curves often have few counts per bin, in which case the traditional approach – calculating the source counts as  $T - B$ , where  $T$  is the total number of counts in the source extraction region and  $B$  is the number of expected background counts, with a symmetric  $1\sigma$  uncertainty calculated by propagating  $\sqrt{N}$  errors – breaks down. We used the Bayesian approach of Kraft et al. (1991) to calculate the most-likely source count-rate and the 68 percent confidence interval in each bin with fewer than 20 net source counts or where the traditional  $1\sigma$  uncertainty extended below zero.

We also calculated a summed EPIC light curve. The counts in the PN, MOS1 and MOS2 source light curves were summed, as were those in the background light curves, and the net source light curve was calculated as described above. A PN-effective count rate was calculated for each bin by scaling the MOS active exposure time by the factor 0.370 (see above) and dividing the summed net counts by the sum of active exposure times in each bin. Two example light curves are shown in Fig. 3, one containing a strong flare (IQ Tau = XEST-14-006), and one containing some low-level activity (V807 Tau = XEST-04-012) seen in most stellar light curves in XEST.

##### 4.9.2. Spectra


Source and background spectra were produced for each instrument using data in the GTIs used for source detection. This may be too conservative for some bright sources, but in trying to extract products automatically for sources with a wide range of count-rates, this is most straightforward. For sources whose light curves showed clear flares we also excluded times of flaring emission where possible, as we aimed to avoid measuring properties of a source in a temporary peculiar state.

PN spectra were extracted using events with patterns 0–4, as the spectral response is not calibrated for higher patterns. Ancillary responses files (ARFs) were produced for each source spectrum using the SAS task ARFGEN. We used the canned response matrix files (RMFs) appropriate for SAS v6.1.0. For each source spectrum we used the PN RMF appropriate for patterns 0–4, the observing mode and the source position on the CCD, and MOS RMFs appropriate for Imaging mode, patterns 0–12, and the orbit (date) of observation.

Three examples of EPIC PN spectra are shown in Fig. 4 together with the best DEM fits (see below).

#### 4.10. Overall sensitivity


An example of a rich EPIC field in L1495, centered at V410 Tau, is displayed in Figure 5, showing a particularly deep observation within our survey. A typical exposure of  $\approx 30$  ks duration with an average background contamination level reached a detection threshold of  $\approx 9 \times 10^{27}$  erg s $^{-1}$  on-axis and  $\approx 1.3 \times 10^{28}$  erg s $^{-1}$  at  $10'$  off-axis for an X-ray source with a thermal spectrum characteristic of T Tau stars (see below) subject to a hydrogen




**Fig. 3.** Two X-ray light curves, summed from all three EPIC detectors in the energy range 0.5–7.3 keV. The total count rate has been re-normalized to the effective area of the PN. The upper panel shows IQ Tau = XEST-14-006 that flared strongly in the second half of the observation. The low-level episode used for spectral fitting was defined to cover the first 18 ks (bin size: 250 s). The lower panel shows V807 Tau = XEST-04-012, exhibiting low-level flaring variability seen in most XEST objects (bin size: 150 s). Here, all data were used for the spectral fits.

absorption column density of  $N_{\text{H}} = 3 \times 10^{21} \text{ cm}^{-2}$ . This threshold turns out to be appropriate to detect essentially every T Tau star in the surveyed fields. The precise detection limits depend on the level of particle background contamination (hence the exposure time obtained during low background radiation), and the off-axis angle. The effective area drops to about 50% at 10' distance from the center and to 35% at the border of the detectors (15' distance).

A typical XEST exposure contained between 50 and 100 X-ray point sources (Table 3) of which, however, a minority (of order 10) are known TMC stellar members. The remainder are probably mostly extragalactic sources, but a non-negligible contribution may be defined by as yet unidentified new TMC members that may be embedded or be located behind the molecular cloud complex (see Scelsi et al. 2007 for a detailed study).



**Fig. 4.** Three example CCD spectra from the EPIC PN camera. All spectra extracted from the source position were binned to a minimum of 15 cts per bin. The histograms show the best fits derived from DEM fits. *Top:* Hubble 4 = XEST-23-056, an X-ray strong WTTS with modest absorption (total of 27648 cts in EPIC,  $N_{\text{H}} = 3.1 \times 10^{21} \text{ cm}^{-3}$ ). *Middle:* FS Tau = XEST-11-057, an intermediately X-ray strong CTTS with strong absorption (total of 7398 cts in EPIC,  $N_{\text{H}} = 1.4 \times 10^{22} \text{ cm}^{-3}$ ). *Bottom:* L1489 IRS = IRAS 04016+2610 = XEST-06-059, an X-ray bright protostar with very strong absorption (total of 3336 cts in EPIC,  $N_{\text{H}} = 6.6 \times 10^{22} \text{ cm}^{-3}$ ).



**Fig. 5.** Example of an XMM EPIC image (the field around V410 Tau = L1495). The field diameter is  $30'$ , and the point-spread function has an *FWHM* of about  $5''$ . North is up. The colors indicate X-ray hardness, the red-orange sources being softest and the blue sources being hardest. This exposure was longer than average ( $74$  ks compared to the more typical  $30$ – $40$  ks). It contains  $76$  X-ray detections, of which  $20$  are TMC members.

#### 4.11. Chandra data

The few exposures obtained with *Chandra* were reduced using standard strategies for ACIS, while we used the standard, reduced events file for the HRC-I observation around HR 1442. The ACIS-S observations of the fields around DG Tau and FS Tau used the “Very Faint Mode” of ACIS, while those around the fields of GV Tau, V410 Tau, L1527 used the “Faint Mode”. The data were reduced in CIAO vers. 3.3.0.1 following the standard analysis threads<sup>6</sup>. These procedures included, for ACIS, corrections for charge transfer inefficiency and afterglow, and selection of good time intervals. Spectra were extracted from level 2 event files with DMEXTRACT, and response and ancillary response files were generated with the MKRMF and MKARF tasks, respectively. We re-emphasize that these data are added here as they provide information on some objects undetected by *XMM-Newton*, but the observations have been obtained with rather different exposure times, and the detectors in use vary (ACIS-S, ACIS-I, HRC-I). Some of the objects were at large off-axis angles and were thus subject to severe distortion (PSF *FWHM* up to several arcseconds). The data quality, e.g., positional accuracy or spectral information, is therefore very inhomogeneous for this sample.

#### 4.12. Spectral analysis

##### 4.12.1. Multi-thermal fits

For the spectral analysis, the MOS spectra provide little or no additional constraints for the many fainter sources, with a total count rate smaller than the count rate of the PN. To keep our analysis as uniform as possible, we therefore used only the PN

spectrum for our spectral fits. Exceptions are those sources for which the PN data were not available, e.g., for sources whose image falls into a PN CCD gap, or for all sources in XEST exposure 26 for which the PN was not operational. To derive basic X-ray properties from the spectra, we followed two strategies. First, conventional one- or two-component spectral fits ( $1-T$  or  $2-T$  henceforth) were performed. The two components define two plasmas with different temperatures  $T_{1,2}$  and emission measures  $EM_{1,2}$ . The latter parameters were fitted in XSPEC (Arnaud 1996) using the *vapec* thermal collisional ionization equilibrium model that includes emission lines and continua. The abundances were fixed at values typical for pre-main sequence or extremely active zero-age main-sequence stars (based on results from Telleschi et al. 2005; Argiroffi et al. 2004; Garcia-Alvarez et al. 2005; and Scelsi et al. 2005)<sup>7</sup>. The spectral components were subject to photoelectric absorption based on the Morrison & McCammon (1983) cross sections. The equivalent hydrogen column density,  $N_H$ , was treated as a further fit parameter. For the faintest sources, a  $2-T$  fit provided too many free parameters for the information contained in the spectrum. A  $1-T$  fit was performed in these cases.

##### 4.12.2. Emission-measure distributions

For highly absorbed spectra, the multi-thermal fit method can become problematic because the softer component is strongly suppressed or may not be detected at all, biasing the fits toward high temperatures. This may further bias the total X-ray luminosity and  $N_H$ . We therefore, in an alternative approach, combined several thermal components in such a way that they describe the shape of the differential emission measure distribution (DEM) in a similar way as previously found for nearby pre-main sequence stars or extremely active zero-age main sequence stars (Telleschi et al. 2005; Argiroffi et al. 2004; Garcia-Alvarez et al. 2005; Scelsi et al. 2005). In short, we have adopted a model in which the DEM shows one peak located at the temperature  $T_0$ , and two power-law distributions toward lower and higher temperatures, characterized by their power-law indices  $\alpha$  and  $\beta$ , respectively. Given the poor temperature discrimination of CCD spectra in the softer X-ray range (in particular in the presence of significant photoelectric absorption), the power-law index on the low-temperature side was kept fixed at a value often found for magnetically active main-sequence and pre-main sequence stars, namely  $\alpha = 2$ , while  $\beta$  was treated as a free parameter. The DEM was cut off at  $\log T = 6.0$  below which EPIC is not sensitive to photons, and at  $\log T = 7.95$  above which EPIC is insensitive to temperature. Within this range, the DEM was binned to  $d \log T = 0.1$ , i.e., 20 bins (centered at  $\log T = 6.0, 6.1, \dots, 7.9$ ) were used. The element abundances were fixed at the same values as those adopted for the  $1$ - or  $2-T$  fits. Again, the absorbing hydrogen column density  $N_H$  was also fitted to the spectrum. The final fit parameter was the normalization, defined as the EM in the temperature bin at  $T_0$ . This method thus assumes that the DEM can be described by four fit parameters, namely  $T_0$ ,  $\beta$ ,  $EM_0$ , and  $N_H$ . We summed the EM over all EM bins used for the calculation to obtain the total  $EM_t$ .

We set the following hard limits to the parameters:  $-3 \leq \beta \leq 1$ ,  $2 \text{ MK} \leq T_0 \leq 32 \text{ MK}$ . In exceptional cases (very hard, absorbed spectra), we extended the range for  $T_0$  to slightly

<sup>7</sup> The adopted abundances are, with respect to the solar photospheric abundances given by Anders & Grevesse (1989) = AG89: C = 0.45, N = 0.788, O = 0.426, Ne = 0.832, Mg = 0.263, Al = 0.5, Si = 0.309, S = 0.417, Ar = 0.55, Ca = 0.195, Fe = 0.195, Ni = 0.195.

<sup>6</sup> <http://asc.harvard.edu/ciao/threads/>

higher values, although we emphasize that at such temperatures, the dominant contribution to the spectrum is a featureless bremsstrahlung continuum, apart from the Fe K complex at 6.7 keV.

For very faint sources, such fits provide too many degrees of freedom. We thus fixed one or more of the following parameters:  $\beta$ , usually fixed at a value of  $-1$  which is often found for brighter sources, or  $N_H$ , fixed at a value corresponding to the tabulated  $A_V$  assuming a standard gas-to-dust ratio (after Vuong et al. 2003), or both. In a few cases, it was necessary to fix  $T_0$  as well, usually at  $\log T_0 = 7$ . Finally, for non-detections with reasonably well-known  $A_V$ , we derived upper limits to  $L_X$  as follows: We used the 95% confidence upper limits to the count rates derived at the optical or infrared position of the star, and then adopted an  $N_H$  derived from  $A_V$ . We thus estimated the upper limit using an average thermal model. For brown dwarfs and very-low mass stars, we adopted  $\beta = -3$  and  $\log T_0 = 6.9$ , characteristic values found for the detected low-mass objects. For more massive T Tau stars, we adopted  $\beta = -1$  and  $\log T_0 = 7$ , again characteristic values found for the detected stars. These upper limits should only be used as rough estimates. For embedded sources, uncertain visual or infrared extinctions and the strong influence of  $N_H$  on the measured count rate make such estimates unreliable, which will therefore not be given. As we will predominantly use the DEM-fit results, we have not derived upper limits based on 1- $T$  plasmas.

For both models, the total X-ray luminosity was computed for the energy range 0.3-10 keV based on the integration of the best-fit model. We adopted a distance of 140 pc for all objects.

The same spectral-fit procedures were applied to the *Chandra* ACIS data if feasible (most of the faint sources were interpreted only with a 1- $T$  model). For the HRC data, we assumed a standard electron temperature of 10 MK and estimated  $N_H$  from  $A_V$  using standard interstellar conversion ratios (Vuong et al. 2003 and references therein). The unabsorbed source flux and the corresponding luminosity were then estimated based on the PIMMS software<sup>8</sup>.

## 5. Results

Tables 4 – 11 summarize basic X-ray parameters and fundamental properties of all observed TMC members. We reproduce the first ten entries per table for illustration. The entire tables with comments and references are available in the electronic version of this paper.

All tables are of identical length, each line number referring to the same observation of the same target across all tables. The tables are sorted in increasing right ascension. The first column of each table gives the XEST catalog entry. This number is composed of the two-digit code corresponding to the exposure number listed in Table 1, followed by the X-ray source number for this exposure (including every X-ray source regardless of TMC membership). This source number is ‘000’ for X-ray non-detections of known TMC members. The second column gives a conventional name frequently used in the literature. Preference was mostly given to variable-star names if available. Note that several objects were observed twice within the principal XEST project (in particular in exposures 23 and 24). They are listed as separate entries. XEST-25 is one of a series of exposures containing the TMC member AA Tau. This star is highly X-ray variable probably due to variable absorption by its disk. We use only one observation to report the typical X-ray properties of this star.

A more detailed analysis of the temporal behavior of AA Tau is given by Grosso et al. (in preparation). There are no other known TMC stellar members in this field.

The final portion of each table refers to results from the complementary *Chandra* observations. Here, we list TMC objects that were either not covered by *XMM-Newton*, regardless of their detection status with *Chandra*, or objects that were detected by *Chandra* but not by *XMM-Newton*. We also add FS Tau because *XMM-Newton* observed this star during a slow flux decrease, presumably from a flare, and Haro 6-5 B which is slightly contaminated by the wings of FS Tau A in the *XMM-Newton* observation. GV Tau is added because of the higher resolution of *Chandra* that attributes the flux to GV Tau A (Güdel et al. 2007b), while the embedded object Haro 6-10 B at 1.3" remains undetected.

### 5.1. The XEST X-ray results

The individual X-ray results from our survey will be discussed in depth in the series of accompanying papers. We collect here the basic, statistical X-ray results for the stellar TMC sources. Our results refer to episodes during the observations that were not affected by outstanding stellar flares (i.e., flares in excess of the typical slow modulation within factors of  $\approx 2$  of the average level), although low-level variability is common to pre-main sequence X-ray sources. Exceptions are CFHT-BD-Tau 1 = XEST-17-068 that showed a flare with a slow decay, dominating most of the exposure time, and FS Tau = XEST-11-057, DH Tau = XEST-15-040 and V830 Tau = XEST-04-016 that showed slowly decreasing light curves, probably following a strong flare starting before our exposures. Specific discussions of the time-dependent behavior of all observed sources will be given by Stelzer et al. (2007), Franciosini et al. (2007), Arzner et al. (2007a), and Audard et al. (2007).

Table 4 provides a summary of detection results for each XEST TMC member. The table lists the detection coordinates with statistical errors and the offsets from the expected position as given later in Table 7. For non-detections, the expected coordinates of the source is given (see below), which were used to calculate upper limits to the count rates. Then, the maximum likelihood (ML) is given for each detection, referring to the full (0.5–7.3 keV) band. Non-detections show  $ML = 0$ . For the *Chandra* observations in Table 4, we list the “significance” of the detection as provided by the CIAO WAVDETECT algorithm instead of  $ML_{det}$ . Then, we list the total number of detected EPIC counts, the effective PN-equivalent on-axis exposure time that corrects for vignetting and also takes the (lower-sensitivity) MOS data into account, as described above. For *Chandra*, these numbers refer to the respective detector used for the observations. For *XMM-Newton* non-detections, the 95% upper limits are reported. Next, the table gives the average count rate (on-axis equivalent rate for PN). Finally, we report a variability flag from inspection of the light curves; the codes have the following meaning: 0 = no or only low-level variability, no flares visible that may dominate the X-ray spectrum; 1 = clear flaring observed and flare intervals removed in spectral fit; 2 = clear flaring observed but flare intervals not removed; 3 = slow decay of flare throughout the observation, all data used. Examples for light curves with flag 1 (IQ Tau) and flag 0 (V807 Tau) are shown in Fig. 3. In the former case, spectral fits were obtained using only data during the low-level episode, defined by the first 18 ks of the light curve. A more quantitative analysis of light curve variability and an interpretation of the role of flares in coronal heating will be given by Stelzer et al. (2007).

<sup>8</sup> <http://heasarc.gsfc.nasa.gov/Tools/w3pimms.html>

**Table 4.** X-ray parameters of targets in XEST (1): Positions and count rates [first ten entries].

XEST	Name	RA <sub>X</sub>			Dec <sub>X</sub>		Poserr ( $''$ )	Offset ( $''$ )	ML <sub>det</sub> <sup>a</sup>	Scts	T <sub>exp</sub> (s)	Rate (ct s <sup>-1</sup> )	Var <sup>b</sup>	
		h	m	s	deg	'								''
27-115	HBC 352	3	54	29.54	32	03	02.2	1.04	0.89	9395	3832	15 776	0.2429	0
27-000	HBC 353	3	54	30.17	32	03	04.3	0.00	0.00	0	<183	14 469	<0.0127	0
06-005	HBC 358 AB	4	03	49.27	26	10	53.1	1.48	1.22	1073	879	7361	0.1194	0
06-007	HBC 359	4	03	50.82	26	10	53.0	1.46	0.34	2921	1605	7664	0.2094	0
06-059	L1489 IRS	4	04	43.07	26	18	56.3	1.46	0.10	7632	3336	46 853	0.0712	0
20-001	LkCa 1	4	13	14.03	28	19	09.9	1.58	1.71	455	336	7813	0.0430	0
20-005	Anon 1	4	13	27.28	28	16	23.3	1.53	1.64	24658	8299	21 634	0.3836	0
20-000	IRAS 04108+2803 A	4	13	53.29	28	11	23.4	0.00	0.00	0	<26	29 957	<0.0009	0
20-022	IRAS 04108+2803 B	4	13	54.72	28	11	32.2	1.55	0.70	1145	696	30 660	0.0227	1
20-000	2M J04141188+28	4	14	11.88	28	11	53.5	0.00	0.00	0	<54	48 645	<0.0011	0

<sup>a</sup> Maximum likelihood for detection for *XMM-Newton* data, CIAO WAVDETECT “Significance” for *Chandra* data.

<sup>b</sup> Variability flag: 0 = no or only low-level variability; 1 = clear flaring, flare intervals removed in spectral fit; 2 = clear flaring observed but flare intervals not removed; 3 = slow decay of flare throughout observation, all data used.

**Table 5.** X-ray parameters of targets in XEST (2): Plasma parameters from the DEM fits [first ten entries].

XEST	Name	N <sub>H</sub> (1σ range) (10 <sup>22</sup> cm <sup>-2</sup> )	T <sub>0</sub> (1σ range) (MK)	β (1σ range)	EM <sub>t</sub> <sup>a</sup> (10 <sup>52</sup> )	L <sub>X</sub> <sup>b</sup> (range) (10 <sup>30</sup> erg s <sup>-1</sup> )	log L <sub>X</sub> /L <sub>*</sub>	T <sub>av</sub> (MK)	χ <sub>red</sub> <sup>2</sup>	d.o.f.
27-115	HBC 352	0.22 (0.19,0.25)	6.5 (5.0, 8.2)	-0.60 (-0.79, -0.47)	25.21	2.657 (2.43, 2.87)	-3.03	12.00	0.85	129
27-000	HBC 353	0.17	10.0	-1.00	...	<0.176	<-4.04	...	...	...
06-005	HBC 358 AB	0.01 (0.00,0.04)	4.7 (3.0, 7.5)	-0.60 (-1.24, -0.34)	3.79	0.383 (0.37, 0.44)	-3.45	9.42	0.74	21
06-007	HBC 359	0.01 (0.00,0.02)	7.2 (5.8, 8.9)	-1.25 (-1.84, -0.91)	6.65	0.663 (0.64, 0.69)	-3.18	9.20	1.05	57
06-059	L1489 IRS	6.63 (6.18,7.09)	50.9 (30.9,60.5)	-3.00 (-3.00, 0.02)	30.39	4.471 (4.07, 4.91)	-3.63	40.42	0.94	99
20-001	LkCa 1	0.07 (0.04,0.15)	4.9 (3.7, 5.7)	-3.00 (-3.00, -1.94)	2.66	0.232 (0.20, 0.37)	-3.80	4.35	0.57	18
20-005	Anon 1	0.33 (0.30,0.35)	7.6 (6.4, 9.1)	-1.01 (-1.26, -0.88)	40.27	4.139 (3.84, 4.42)	-3.38	10.72	0.92	228
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	5.68 (2.99,8.19)	4.5 (2.0,31.6)	-1.00	4.39	0.417 (0.07, 0.57)	-3.57	6.93	0.68	8
20-000	2M J04141188+28	0.18	7.9	-3.00	...	<0.028	<-3.31	...	...	...

<sup>a</sup> EM<sub>t</sub> is sum of EM over all DEM bins from log *T* = 6.0 to log *T* = 7.9; given in units of 10<sup>52</sup> cm<sup>-3</sup>.

<sup>b</sup> L<sub>X</sub> for [0.3,10] keV, in units of 10<sup>30</sup> erg s<sup>-1</sup>.

**Table 6.** X-ray parameters of targets in XEST (3): Plasma parameters from the 1-*T* and 2-*T* fits [first ten entries].

XEST	Name	N <sub>H</sub> (1σ range) (10 <sup>22</sup> cm <sup>-2</sup> )	T <sub>1</sub> <sup>a</sup> (MK)	T <sub>2</sub> (MK)	EM <sub>1</sub> <sup>b</sup> (10 <sup>52</sup> )	EM <sub>2</sub> <sup>b</sup> (10 <sup>52</sup> )	L <sub>X</sub> <sup>c</sup> (10 <sup>30</sup> )	log L <sub>X</sub> /L <sub>*</sub>	T <sub>av</sub> (MK)	χ <sub>red</sub> <sup>d</sup>	d.o.f.
27-115	HBC 352	0.19 (0.17,0.21)	7.54	23.77	10.35	11.52	2.307	-3.09	13.81	0.87	128
27-000	HBC 353	...	...	...	...	...	...	...	...	...	...
06-005	HBC 358 AB	0.00 (0.00,0.03)	4.29	14.26	1.41	2.12	0.346	-3.49	8.82	0.73	20
06-007	HBC 359	0.00 (0.00,0.02)	5.33	15.19	3.06	3.53	0.628	-3.20	9.34	1.09	56
06-059	L1489 IRS	6.54 (6.08,6.88)	...	50.20	...	26.57	4.003	-3.67	50.20	1.02	100
20-001	LkCa 1	0.07 (0.00,0.21)	1.39	8.35	1.76	0.82	0.205	-3.85	2.46	1.27	15
20-005	Anon 1	0.28 (0.27,0.29)	8.58	24.12	19.05	14.11	3.473	-3.46	13.32	1.02	228
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	7.86 (3.40,28.0)	...	11.59	...	5.95	0.588	-3.42	11.59	0.61	8
20-000	2M J04141188+28	...	...	...	...	...	...	...	...	...	...

<sup>a</sup> “=” sign before number indicates that parameter was fixed. <sup>b</sup> EM in units of 10<sup>52</sup> cm<sup>-3</sup>. <sup>c</sup> L<sub>X</sub> for [0.3,10] keV, in units of 10<sup>30</sup> erg s<sup>-1</sup>. <sup>d</sup> Numbers followed by “C” denote C statistic (for low-background *Chandra* data).

Table 5 summarizes the basic spectral-fit results from our DEM method. The abundances were kept fixed at characteristic values as described in Sect. 4.12. The columns provide N<sub>H</sub>, the temperature of the DEM peak, T<sub>0</sub>, the power-law index β of the high-temperature DEM slope, the total EM integrated over all bins centered at values of log *T* = 6.0 to log *T* = 7.9 in steps of *d* log *T* = 0.1, and the inferred L<sub>X</sub> in the 0.3–10 keV range. We list 68% (1σ) confidence ranges for N<sub>H</sub>, β, and T<sub>0</sub> in parentheses. For fixed parameters, no error range is given. For non-detections, the spectral parameters used to estimate upper limits to L<sub>X</sub> are also given as fixed parameters. The next columns list the logarithms of the normalized luminosities L<sub>X</sub>/L<sub>\*</sub> where the stellar bolometric luminosity L<sub>\*</sub> is adopted from Table 9, and the

average electron temperature that is obtained as a mean of log *T* over the entire DEM, where DEM(*T*) is used as a weight function, calculated analytically as follows:

$$A = \frac{T^\alpha}{\alpha T_0^\alpha} \left( \log T - \frac{1}{c\alpha} \right) \Big|_{T_l}^{T_0} + \frac{T^\beta}{\beta T_0^\beta} \left( \log T - \frac{1}{c\beta} \right) \Big|_{T_0}^{T_h} \quad (2)$$

$$B = \frac{T^\alpha}{\alpha T_0^\alpha} \Big|_{T_l}^{T_0} + \frac{T^\beta}{\beta T_0^\beta} \Big|_{T_0}^{T_h} \quad (3)$$

$$\log T_{\text{av}} = \frac{A}{B} \quad (4)$$

**Table 7.** Fundamental parameters of targets in XEST (1): Names and coordinates [first ten entries].

XEST	Name	2MASS <sup>a</sup>	IRAS <sup>b</sup>	Alternative names	RA(J2000.0) <sup>c</sup>			Dec(J2000.0) <sup>c</sup>		
					h	m	s	deg	'	''
27-115	HBC 352	03542950+3203013	...	NTTS 035120+3154SW	3	54	29.51	32	03	01.4
27-000	HBC 353	03543017+3203043	...	NTTS 035120+3154NE	3	54	30.17	32	03	04.3
06-005	HBC 358 AB	04034930+2610520	...	NTTS 040047+2603W	4	03	49.31	26	10	52.0
06-007	HBC 359	04035084+2610531	...	TTS 040047+2603	4	03	50.84	26	10	53.2
06-059	L1489 IRS	04044307+2618563	04016+2610	...	4	04	43.07	26	18	56.4
20-001	LkCa 1	04131414+2819108	...	HBC 365, V1095 Tau, JH 141	4	13	14.14	28	19	10.8
20-005	Anon 1	04132722+2816247	...	HBC 366, V1096 Tau	4	13	27.23	28	16	24.8
20-000	IRAS 04108+2803 A	04135328+2811233	04108+2803A	L1495N IRS	4	13	53.29	28	11	23.4
20-022	IRAS 04108+2803 B	04135471+2811328	04108+2803	...	4	13	54.72	28	11	32.9
20-000	2M J04141188+28	04141188+2811535	...	...	4	14	11.88	28	11	53.5

<sup>a</sup> Nearest 2MASS entry within 5'' to coordinates given by references 5, 18, 33, or SIMBAD. Unlikely identifications in parentheses.

<sup>b</sup> Nearest IRAS catalog entry, within 10''.

<sup>c</sup> 2MASS coordinates. For unlikely identifications, SIMBAD (S).

**Table 8.** Fundamental parameters of targets in XEST (2): Multiplicity [first ten entries].

XEST	Name	Comp	Separations ('' )		Refs.
27-115	HBC 352	1	...	...	...
27-000	HBC 353	1	...	...	...
06-005	HBC 358 AB	3	0.15, 1.55	11, 20, 31, 40	...
06-007	HBC 359	1	...	...	...
06-059	L1489 IRS	1	...	...	...
20-001	LkCa 1	1	...	...	...
20-005	Anon 1	1	...	...	...
20-000	IRAS 04108+2803 A	1	...	...	...
20-022	IRAS 04108+2803 B	1	...	...	...
20-000	2M J04141188+28	1	...	...	...

**Table 9.** Fundamental parameters of targets in XEST (3): Photometry and spectroscopy [first ten entries].

XEST	Name	Spec <sup>a</sup>	Refs.	$A_V^a$ (mag)	$A_J$ (mag)	$T_{\text{eff}}^a$ (K)	$L_*^b$ ( $L_\odot$ )	Refs.
27-115	HBC 352	G0	27	0.87	0.25	6030	0.740	27
27-000	HBC 353	G5	27	0.97	0.28	5770	0.500	27
06-005	HBC 358 AB	M2	27	0.21	0.06	3580	0.280	27
06-007	HBC 359	M2	27	0.49	0.14	3580	0.260	27
06-059	L1489 IRS	K4	59	10.20	...	4500	4.900	10, 59
20-001	LkCa 1	M4	5	0.00	0.00	3270	0.380	5, 27
20-005	Anon 1	M0	5	1.32	1.03	3850	2.600	5, 27
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	...	...	...	...	3500	0.400	10
20-000	2M J04141188+28	M6.25	33	...	0.28	2962	0.015	33

<sup>a</sup> For multiples, first number or spectral type refers to primary, second to secondary component.

<sup>b</sup> For multiples, three numbers give primary/secondary/total system luminosity.

<sup>c</sup> Referring to  $L_{\text{bol}}$  as derived from integration of the optical and infrared spectrum.

where  $c = \ln 10 \approx 2.3$  (note that we use decadic logarithms throughout), and we define the limits of the integration as  $T_l = 10^6$  K and  $T_h = 10^8$  K. Finally, we list the reduced  $\chi_{\text{red}}^2$  achieved by the fit together with the number of degrees of freedom (d.o.f.).

Table 6 equivalently reports the results from 2- $T$  or, for faint sources, 1- $T$  fits. The columns give  $N_H$  (with errors if fitted),  $T_1$ ,  $T_2$ , and the corresponding  $EM_1$ ,  $EM_2$ . If only one component was fitted, its temperature and EM values are listed in the columns for  $T_2$  and  $EM_2$  if the temperature exceeds 10 MK, and in the columns for  $T_1$  and  $EM_1$  otherwise. If a temperature was held fixed, its value is preceded by a “=” sign. Further, the table gives  $L_X$ ,  $\log(L_X/L_*)$ , and  $T_{\text{av}}$ , the latter defined as the

EM-weighted logarithmic average of the temperature in the case of two non-vanishing components:

$$\log T_{\text{av}} = \frac{EM_1 \log T_1 + EM_2 \log T_2}{EM_1 + EM_2}. \quad (5)$$

Finally, the table lists the  $\chi_{\text{red}}^2$  values and the number of degrees of freedom (d.o.f.).

Errors of EM and  $L_X$  are difficult to compute, and their meaning is not unequivocal. Our EM models are rather crude approximations to true emission measure distributions. Formal error ranges for the EM values may therefore bear little relevance. Further,  $L_X$  is a complicated function of the thermal emission model. Although resolved spectral features can usually be fitted

**Table 10.** Fundamental parameters of targets in XEST (4): Age, mass, radius, rotation [first ten entries].

XEST	Name	Age <sup>a,b</sup> (Myr)	Mass <sup>a,c</sup> ( $M_{\odot}$ )	Refs.	Radius <sup>d</sup> ( $R_{\odot}$ )	$P$ (d)	Refs.	$v \sin i$ ( $\text{km s}^{-1}$ )	Refs.
27-115	HBC 352	...	1.05	2	0.79	<0.53	C	>75.00	48
27-000	HBC 353	...	...	...	0.71	<4.08	C	8.80	2
06-005	HBC 358 AB	3.26	0.41	27	1.38	...	C	<10.00	2
06-007	HBC 359	3.54	0.41	27	1.33	...	C	<10.00	48
06-059	L1489 IRS	0.80	1.45	10	3.65	<4.02	C	46.00	10
20-001	LkCa 1	0.87	0.27	5	1.93	<4.18	C	23.30	48
20-005	Anon 1	0.50	0.56	5	3.63	...	...	...	...
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	1.60	0.36	10	1.72	<6.23	C	14.00	10
20-000	2M J04141188+28	...	0.08	46	0.47	...	...	...	...

<sup>a</sup> For binaries, first number refers to primary, second to secondary component (calculated from  $L_*$  and  $T_{\text{eff}}$ ).

<sup>b</sup> Ages derived after Siess et al. (2000) using the same principal parameters as for masses, quoted in Table 9.

<sup>c</sup> Masses derived after Siess et al. (2000) using principal parameters quoted in Table 9.

<sup>d</sup> For multiples, radius is given only for primary if luminosity of primary is explicitly known

**Table 11.** Fundamental parameters of targets in XEST (5): Accretion and evolution [first ten entries].

XEST	Name	$\dot{M}$ (min/max) <sup>a</sup> ( $M_{\odot}\text{yr}^{-1}$ )	Refs.	$EW(\text{H}\alpha)$ <sup>b</sup> ( $\text{\AA}$ )	TTS type	Refs. <sup>c</sup>	IR <sup>d</sup> class	Refs. <sup>d</sup>	Type
27-115	HBC 352	...	...	0	W	29	III	1, 27	3
27-000	HBC 353	...	...	0	W	29	III	27	3
06-005	HBC 358 AB	<8.97	20	4- 10	W/W	37, 29	III	1, 27	3
06-007	HBC 359	...	...	2- 9	W	56, 37	III	27	3
06-059	L1489 IRS	-7.15	59	41- 56	C*	59, 29	I	1, 59, 27	1
20-001	LkCa 1	<-9.72	58	3- 4	W	29, 45	III	1, 27	3
20-005	Anon 1	<-8.94	58	1- 3	W	37, 29	III	1, 27	3
20-000	IRAS 04108+2803 A	...	...	37	C	29	II	27	2
20-022	IRAS 04108+2803 B	...	...	...	...	...	I	59, 27	1
20-000	2M J04141188+28	-10.00	46	250	C	46	...	...	4

<sup>a</sup> Range of  $\dot{M}$  reported in literature given. For multiple systems, numbers refer to primary or integrated system.

<sup>b</sup> Range of  $EW$  reported in literature given. For multiple systems, numbers refer to primary or integrated system.

<sup>c</sup> For  $EW$  range, first reference for minimum, second for maximum reported.

<sup>d</sup> Infrared classification; double entries: '/' for transition objects, '+' for components, ';' for different types, ',' for different references. FS = flat-spectrum source.


sufficiently well in a  $\chi^2$  sense using a number of thermal components that describe the absorbed, observed spectral flux density, the major uncertainty in  $L_X$  is introduced by fitting (and correcting for)  $N_H$ . An instability discussed below makes determinations of  $L_X$  rather uncertain once cool plasma components are subject to large absorption while the spectrum is of modest quality. To quantify this latter effect that dominates the uncertainty of  $L_X$  in many cases, we proceeded as follows. We adopted the upper and lower  $1\sigma$  bounds of  $N_H$  (if  $N_H$  was a fit parameter), kept these values fixed and refitted the spectrum. The  $L_X$  values thus determined,  $L_{X,1}$  and  $L_{X,2}$  for the lower and the upper bound, are taken to bracket the error range of the tabulated  $L_X$ . We performed this error analysis for the DEM fit method. We graphically show the magnitude of the ratio  $L_{X,2}/L_{X,1}$  in Fig. 6 as a function of the best-fit  $N_H$  (set to  $0.01 \times 10^{22} \text{ cm}^{-2}$  for sources with  $N_H$  smaller than this value) and the number of counts used in the spectral fit. Expectedly, the error range is largest for faint and strongly absorbed sources, and smallest for bright and weakly absorbed sources. For 83 sources,  $L_{X,2}/L_{X,1} \leq 2$  and for 48 sources,  $L_{X,2}/L_{X,1} > 2$ . For 99 sources,  $L_{X,2}/L_{X,1} \leq 3$  and for 32 sources,  $L_{X,2}/L_{X,1} > 3$ . Gagné et al. (2004) described a similar analysis performed for errors in  $L_X$  for *Chandra* observations of the  $\rho$  Ophiuchus Cloud, with similar conclusions.

We note, however, that T Tau stars are manifestly variable on timescales of hours, typically within a factor of at least two, while most of the EM errors are formally much smaller. A detailed variability analysis of all XEST sources will be presented by Stelzer et al. (2007). Therefore, for the majority of our sources, the formal uncertainties from the spectral fits underestimate the true uncertainty. Realistically, thus, standard error bars of (at least) 0.3 dex should be adopted for all values of  $L_X$  (and EM). Larger statistical errors from spectral fits are typically found for  $N_H \gtrsim 2 \times 10^{21} \text{ cm}^{-2}$  and  $\lesssim 600$  cts in the spectrum, or in regions where  $\text{cts}/(N_H/10^{22} \text{ cm}^{-2}) \lesssim 500$ .

## 5.2. The XEST catalog

Tables 7–11 identify the X-ray sources and provide a summary of the fundamental properties of the stellar systems. The parameters were extracted from the existing literature, in particular from Cohen & Kuhi (1979), Kenyon & Hartmann (1995), Briceño et al. (1998), White & Ghez (2001), Briceño et al. (2002), Hartmann (2002), Hartigan & Kenyon (2003), Luhman et al. (2003), Luhman (2004), White & Hillenbrand (2004), Andrews & Williams (2005), and Guieu et al. (2006). Several further references are listed in the bibliography summary in the table





**Fig. 6.** Magnitude of the error ranges for  $L_X$ , estimated from maximum variations of  $N_H$  (see text for details), as a function of the best-fit  $N_H$  and the number of counts used in the spectral fit. Results from the DEM fit method are shown. The circle size scales with the logarithm of the ratio between the upper and the lower bounds of  $L_X$ . Red circles denote sources for which the  $\beta$  parameter was held fixed.

section. For the suspected late-B member of TMC, HD 28867, all basic information except for the 2MASS identification comes from Walter et al. (2003) and references therein. A short description of the table entries follows.

### 5.2.1. Names, coordinates, and multiplicity

Table 7 lists the 2MASS catalog entry likely to be identified with the X-ray source. For this identification, we first adopted object coordinates mostly from Briceño et al. (2002), Luhman et al. (2003), Luhman (2004), and Guieu et al. (2006) if available, or from SIMBAD in other cases, and then identified the closest object in the most recent 2MASS catalog version (Cutri et al. 2003) within a circle of radius  $5''$  around this position. In three cases, no 2MASS object was found within  $5''$ . The next columns of Table 7 give the nearest IRAS catalog entry, confined to a search radius of  $10''$  or directly identified from SIMBAD, and a selection of commonly used alternative names. Finally, the J2000.0 coordinates refer to the adopted 2MASS identifications. If no 2MASS identification is given, the coordinates were taken from SIMBAD, marked with '(S)' after the coordinates.

Table 8 provides binary information, i.e., the number of components, the approximate component separations, and references. Note that binary components with a separation of  $\lesssim 5''$  are unresolved in *XMM-Newton* observations, and pairs at  $5$ – $10''$  are difficult to separate. Typically, binaries with separations up to about  $10''$  have been recorded as a single X-ray source (e.g., V819 Tau, but note the case of HP Tau/G2 and G3, two stars with a separation of  $10''$  for which a joint spectral model was derived, while the count rates and thus the luminosities were estimated separately based on PSF fitting).

### 5.2.2. Photometry and spectroscopy

Table 9 lists spectral types, visual and  $J$ -band extinctions ( $A_V$  and  $A_J$ , respectively, the latter only for the primary in multiple systems), effective temperatures ( $T_{\text{eff}}$ ; for binaries, separately given for primary and secondary if available), and bolometric luminosities ( $L_*$ ) of the stars. The latter are bolometric luminosities of the stellar photospheres derived from the optical or near-IR except in cases where only the IR bolometric luminosity, possibly including contributions from the disk and/or the envelope, was available (these cases are marked with “Lb” in the reference column). In the case of binaries,  $L_*$  is given for the primary and the secondary if available, and the entire system. The component and the system values may originate from different references, which accounts for some discrepancies with regard to the sum of the component  $L_*$ . Because spectral types,  $A_V$ ,  $A_J$ , and  $L_*$  sensitively depend on the interpretation of the measured  $T_{\text{eff}}$  and the optical and near-infrared photometry, we attempted to use the same references for these parameters for a given star, and opted predominantly for Briceño et al. (2002), Luhman et al. (2003) and Luhman (2004), and Kenyon & Hartmann (1995) (in this sequence) for single stars or integrated multiple systems, and for White & Ghez (2001) and Hartigan & Kenyon (2003) for binary components. Significant information for deeply embedded sources was adopted from White & Hillenbrand (2004), and for brown dwarfs from Guieu et al. (2006), Luhman et al. (2003), and Luhman (2004). Grosso et al. (2007a) rederived  $T_{\text{eff}}$  of some BDs from the spectral types, while we report here values as originally published. Strict consistency was not possible, in particular for  $A_V$  and  $A_J$  that had to be adopted from different authors and that are inconsistent in several cases (the second references in the last column mostly refer to  $A_V$ ).

### 5.2.3. Ages, masses, and rotation

Table 10 lists ages and masses,  $M$ , that we derived from  $T_{\text{eff}}$  and  $L_*$  as listed in Table 9 using the Siess et al. (2000) isochrones. For binaries, component ages and masses are given if the fundamental parameters were available separately. In a few cases, we did not find a solution from the Siess et al. (2000) tracks; if available, some masses (and ages) were therefore taken directly from the literature (see footnotes to Table 10). The references specifically used for  $T_{\text{eff}}$  and  $L_*$  are listed. Assessing the uncertainties of these parameters is difficult because the uncertainties in  $T_{\text{eff}}$  and  $L_*$  are usually not reported in the literature. If multiple measurements have been reported for an object, then the agreement was (for the TMC) within 100 K for  $T_{\text{eff}}$  for about half of the objects, and within 150 K for two thirds. We conservatively adopt an error of  $\pm 150$  K for  $T_{\text{eff}}$ . Using the Siess et al. calculations for the regions of densest population in the HRD ( $T_{\text{eff}} = 4700, 4000$ , and  $3500$  K with  $L_* = 2, 1$ , and  $0.1 L_\odot$ , respectively), we find mass uncertainties of  $\pm(0.07\text{--}0.16)M_\odot$  or a fractional mass uncertainty of about  $\pm(10\text{--}20)\%$  (higher for cooler stars). An uncertainty in  $T_{\text{eff}}$  also affects the age estimate, resulting in typical deviations of  $\pm(0.1\text{--}0.18)$  dex (factors of 1.3–1.5) from the best-fit value. The age estimate is further affected by uncertainties in  $L_*$  (while  $L_*$  has little influence on the mass estimate in the region where the evolutionary tracks are nearly vertical). We found that in the region of  $T_{\text{eff}} \approx 3500\text{--}4000$  K, a shift of  $d \log L_*$  in  $\log L_*$  results in a shift of approximately  $-d \log L_*$  in age. For  $\approx 80\%$  of the TMC objects with multiple reports of  $L_*$  in the literature, deviations are up to a factor of 2 (0.3 dex), but often much smaller. We thus adopt a conservative, characteristic uncertainty of factors of 2.6–3.0 for the ages, noting that this

is a gross overestimate for many objects, while for a minority of objects, in particular for embedded protostars with poor  $L_*$  determinations, ages may become entirely unreliable.

The radius,  $R$ , was calculated from  $L_* = 4\pi R^2 \sigma T_{\text{eff}}^4$ , where  $\sigma = 5.67 \times 10^{-5} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ K}^{-4}$  is the Stefan-Boltzmann constant. Rotation periods  $P$  are given as measured by various authors. The values were taken from Rebull et al. (2004) who give a comprehensive list of references (not reproduced here). That work includes periods of the lower-mass sample. We have added further rotation periods (Rebull, private communication) extracted from the same bibliography. If not available from the literature, rotation periods were estimated as upper limits if spectroscopic  $v \sin i$  measurements were available (listed in Col. 9, with references in column 10, many again referring to the tabulations and the bibliography given by Rebull et al. 2004). In those cases,  $P \leq 2\pi R / (v \sin i)$ , and the reference column for these calculated periods contains a “C”.

#### 5.2.4. Accretion and evolutionary stage

Table 11 provides parameters relevant for the interpretation of the evolutionary status. Column 3 lists mass accretion rates  $\dot{M}$ . Accretion rates may be variable, and different indirect methods have been used by previous studies to estimate  $\dot{M}$ , hence significantly discrepant values may be found in the published literature. In those cases, we provide ranges, giving a reference for each the minimum and maximum values reported. The same applies to the equivalent width of the  $H\alpha$  line [ $EW(H\alpha)$ ], reported in Col. 5. In multiple systems, both of the above values refer to the primary star even if, in a few cases, a separate measurement was available for a companion. We adopted this strategy because the primary star is most likely also to be the dominant X-ray source.

We assessed the “T Tauri” type (classical or weak-line) based on  $EW$  as follows. For spectral types G and K, stars with  $EW(H\alpha) \geq 5 \text{ \AA}$  are defined as classical T Tauri stars (C), all other stars are weak-line T Tauri stars (W). For early-M stars, the separation was adopted at  $EW(H\alpha) = 10 \text{ \AA}$ , and for mid-M type stars, we adopted  $EW(H\alpha) = 20 \text{ \AA}$  for discrimination. For late-M stars, i.e., typically BDs, the low continuum makes a sensible definition difficult. We will therefore not include BDs when studying differences between CTTS and WTTS, although we have adopted the criterion described by Barrado y Navascués & Martín (2003) that defines accreting stars and BDs as those that show  $L(H\alpha)/L_*$  greater than the chromospheric saturation limit. This criterion also supports our classifications at earlier spectral types. Borderline cases are marked with a question mark. In such cases, we also included other supporting evidence. For example, KPNO-Tau 14 (XEST-18-004) was classified as “W?” (and subsequently as “type 3”, see below) despite one published measurement of  $EW(H\alpha)$  somewhat higher than the saturation limit, because there is one report with much lower  $EW(H\alpha)$ , a very low upper limit to  $\dot{M}$ , and no  $K$  band excess (Luhman et al. 2003; Muzerolle et al. 2005). “C\*” denotes sources for which accretion signatures are evident, but that are embedded and share characteristics with protostars. We have added some “C” and “W” designations from Luhman (2004) even if the explicit  $EW$  values were not given.

Column 8 lists the “Young Stellar Object” (YSO) infrared class as derived from the infrared spectral energy distribution, which is thought to provide information on the presence of disks and molecular envelopes. Class 0 and I objects are protostars, Class II objects are disk-surrounded T Tau stars, whereas disk signatures are weak or absent in Class III stars. “FS” designates

“flat spectrum” sources, intermediate between Class I and II. Most entries refer to IRAS measurements (and were mostly adopted from Kenyon & Hartmann 1995), but some recent information from *Spitzer* is also available (Hartmann et al. 2005). If discrepant entries are given, they are separated by a semi-colon, and so are the corresponding references in the subsequent column. References referring to the same classification are separated by a comma. Transition objects are designated by two types separated by a “/”, while for multiple systems with different component classifications, we give the two classes separated by “+”.


There is a rather good correlation between TTS type indicating presence or absence of accretion (CTTS and WTTS) and YSO infrared class (Classes II and III, respectively) indicating the presence or absence of circumstellar disks. There are, however, exceptions that may be borderline cases or in which IR companions may bias the YSO classification. We therefore adopted a final classification scheme that is predominantly based on the accretion signatures (WTTS vs. CTTS) and takes the IR classification as supporting evidence in borderline cases, except for protostars for which the IR classification is the only relevant parameter. Hence, in col. 10, “0” and “1” stand, respectively, for a protostar of Class 0 or Class I, “2” for an accreting (classical) T Tau star that usually shows a Class-II IR spectrum; “3” corresponds to a weak-line or Class-III object; “4” designates a brown dwarf (spectral class equal to or later than M6.25) irrespective of its accretion signatures; and “5” marks the Herbig Ae/Be stars in TMC. For uncertain classifications or other object types, we use the code “9”.

## 6. Discussion and conclusions

### 6.1. Spectral interpretation: quality of results

To assess the quality of our spectral interpretations, we now compare results derived from the two fit methods (DEM vs 1- or 2- $T$  plasma). Figure 7 compares  $N_{\text{H}}$  values derived from the DEM fits (shown along the abscissa) and from the 1- or 2- $T$  fits (along the ordinate). We show results for bright spectra ( $>600$  cts for the source counts parameter in Table 4 of which typically about half are found in the PN camera) and for faint spectra ( $<600$  cts) separately. The agreement is excellent. The brighter sources show a trend toward  $N_{\text{H}}$  being slightly lower for the 1- $T$  or 2- $T$  fit than for the DEM fit. This can be understood as follows. The DEM model assumes the presence of cool material down to  $\log T = 6$ . This plasma accounts for some of the soft emission where the 1- or 2- $T$  model may be subject to an EM deficit. To account for the same spectral flux, the trend is to lower  $N_{\text{H}}$  for the spectrum with deficient cool EM. The scatter obviously becomes larger for fainter sources, but the error bars grow accordingly so that the agreement is of similar quality (lower plot).


Figure 8 analogously compares  $L_{\text{X}}$ . For most of the brighter stars, the agreement is excellent although there are a few cases for which the 1- or 2- $T$  fit produces significantly higher  $L_{\text{X}}$ . The reason for this deviation is the following. Soft components can be added with relatively little constraints if they are sufficiently absorbed. The temperature then becomes uncertain. A cooler plasma subject to photoelectric absorption will, however, require a much larger EM to produce the same observed flux. If the cool EM is a free fit parameter, an instability may occur with  $T$  converging to low values while  $N_{\text{H}}$  and EM grow excessively. This becomes more evident for fainter sources (lower plot). In several cases, we therefore fixed  $N_{\text{H}}$  at the value expected from  $A_{\text{V}}$  for



**Fig. 7.** Comparison of  $N_{\text{H}}$  values derived from the DEM method ( $x$  axis) and from 1- $T$  or 2- $T$  fits ( $y$  axis; Tables 5 and 6). Only results from *XMM-Newton* data are reported. Upper plot is for sources with more than 600 counts in total (after Table 4), lower plot for sources with fewer than 600 source counts. The solid lines indicate equal  $N_{\text{H}}$ .

the DEM fit, whereas the 1- $T$  fits, requiring one parameter less, were performed without this constraint. The sources that likely suffer from this numerical instability in the 1- $T$  fit are CIDA 7 (XEST-10-034), HO Tau (XEST-09-010), V410 X6 (XEST-23-061), CFHT-Tau 7 (XEST-03-017), MHO 9 (XEST-22-013), and V410 A25 (XEST-23-029) that all show  $L_{\text{X}}$  from the 1- $T$  or 2- $T$  fits at least 3 times higher compared to the DEM fit. In the former, they show  $N_{\text{H}}$  considerably higher than in the latter, or much higher than expected from  $A_{\text{V}}$ , while the dominant electron temperature was very low, between 1 MK and 3.3 MK. The first four of those six stars are faint, while MHO 9 is at the limit between bright and faint as defined for the plots in Fig. 8.


Given that the DEM model is more rigid and avoids biases toward strongly absorbed soft emission from large amounts of cool EM, we will base most of our statistical investigations in the accompanying papers on the DEM results. This model is also more physically meaningful because work on bright spectra of nearby stars has clearly shown continuous emission measure distributions of this kind (e.g., Telleschi et al. 2005). As shown here and below, however, the results for  $L_{\text{X}}$ ,  $N_{\text{H}}$ , and  $T_{\text{av}}$  from the two methods agree well for the brighter spectra. Caution is in order essentially only for spectra defined as “faint” here.



**Fig. 8.** Comparison of  $L_{\text{X}}$  values derived from the DEM method ( $x$  axis) and from 1- $T$  or 2- $T$  fits ( $y$  axis; Tables 5 and 6). Only results from *XMM-Newton* data are reported. Upper plot is for sources with more than 600 counts in total (after Table 4), lower plot for sources with fewer than 600 source counts. The solid lines indicate equal  $L_{\text{X}}$ .

Figure 9 shows the same comparison for weakly absorbed sources (upper plot, for  $N_{\text{H}} < 5 \times 10^{21} \text{ cm}^{-2}$ ) and for more strongly absorbed sources (lower plot, for  $N_{\text{H}} > 5 \times 10^{21} \text{ cm}^{-2}$ ). It is evident that higher absorption introduces more uncertainty to the precise determination of  $L_{\text{X}}$ . Also, these sources obviously tend to be fainter (in terms of count rates).

Figure 10 compares the average electron temperature,  $T_{\text{av}}$ , derived from the DEM and the 2- $T$  or 1- $T$  fits. Given the rather different methodology, the agreement is satisfactory for the brighter sources, while the scatter increases for faint sources, making accurate statements on electron temperatures difficult. The few outliers among the bright sources show very low average temperatures in 1- $T$  or 2- $T$  fits probably due to the same numerical instability described above. The worst agreement is found for the very faint spectrum of LR 1 (XEST-23-048) for which  $T_0$  of the DEM fit is unconstrained despite fixed  $\beta = -1$ . In the presence of severe absorption, the DEM fit finds a low best-fit  $T_0$ , while the 1- $T$  fit converges to a very high temperature. Both  $L_{\text{X}}$  are reasonable, while the DEM fit converges to a higher  $N_{\text{H}}$  which is preferred in the light of the very high  $A_{\text{J}}$  of this star.




**Fig. 9.** Comparison of  $L_X$  values derived from the DEM method ( $x$  axis) and from 1- $T$  or 2- $T$  fits ( $y$  axis; Tables 5 and 6). Only results from *XMM-Newton* data are reported. Upper plot is for sources with  $N_H < 5 \times 10^{21} \text{ cm}^{-2}$ , lower plot for  $N_H > 5 \times 10^{21} \text{ cm}^{-2}$  (based on the DEM fits in Table 5). The solid lines indicate equal  $L_X$ .

Concluding from this comparison, we are confident that  $N_H$  values are meaningful within factors of about 1.5 for the brighter sample and mostly within factors of 2 for the fainter sample. X-ray luminosities appear to be reasonably constrained within a factor of two (although with exceptions), which is a typical range of variability for most sources in any case.

## 6.2. Population statistics

Figure 11 shows an Hertzsprung-Russell Diagram (HRD) of all sources observed in the framework of this survey with isochrones and evolutionary tracks (after Siess et al. 2000) overplotted. The source symbols indicate the source type, and the symbol size corresponds to the X-ray luminosity.

The following characteristics are noteworthy: i) The surveyed sources cover a broad range of masses, from the substellar range up to  $2.5\text{--}3 M_\odot$ , the three most massive stars being HD 28867 and the Herbig stars AB Aur and V928 Tau (the latter being a multiple system). Most objects are concentrated at ages between  $3 \times 10^5\text{--}3 \times 10^6$  yr, and very few at ages around  $10^7$  yr. ii) As pointed out by Kenyon & Hartmann (1995), CTTS and WTTS (Class II and III, shown in gray and white,



**Fig. 10.** Comparison of  $T_{av}$  values derived from the DEM method ( $x$  axis) and from 1- $T$  or 2- $T$  fits ( $y$  axis; Tables 5 and 6). Only results from *XMM-Newton* data are reported. Upper plot is for sources with more than 600 counts in total (after Table 4), lower plot for sources with fewer than 600 source counts. The solid lines indicate equal  $T_{av}$ .

respectively) occupy the same region in the HRD. In fact, this is also true for the detected Class-I protostars. The low-mass and substellar HRD will be more specifically discussed by Grosso et al. (2007a). iii) Most undetected sources are in the very low-mass/substellar regimes and/or have the largest isochronal ages, although for some of the latter objects, the fundamental parameters are poorly determined, as detailed below.

A few objects are apparently located below the zero-age main sequence (ZAMS) which is likely to be a consequence of poor photometry. This is particularly true for embedded sources where scattered light distorts photometric measurements. The stars below the ZAMS are, in order of increasing  $T_{\text{eff}}$ : HH 30 (XEST-22-000, an edge-on star-disk system, Burrows et al. 1996), IRAS S04301+261 (XEST-18-000, a highly reddened object probably primarily detected in scattered light, Briceño et al. 2002), Haro 6-5 B = FS Tau B (XEST-11-054 = C2-1, a near-edge-on star-disk system, Padgett et al. 1999, with  $A_V \approx 10$  mag), HBC 353 (XEST-27-000), and HBC 352 (XEST-27-115). These stars will not be considered in statistical studies whenever  $L_*$  is required (e.g., in  $L_X/L_*$ ). A few stars are located between the 10 Myr isochrone and the ZAMS, making them much older than the bulk population in Taurus. Several

of these objects are relatively poorly studied and may suffer from inaccurate photometry. For most of them, no  $A_V$  has been given, or the extinction is large, and indeed several of them have not been detected (crosses in Fig. 11). These objects are, in order of increasing  $T_{\text{eff}}$  above the  $0.1 M_{\odot}$  track: ITG 33A (XEST-07-000, suggested to be an edge-on star-disk system, Martín 2000:  $A_V = 3.5$  mag), FS Tau (XEST-11-057, a strongly extinguished close binary of CTTS,  $A_V \approx 5$  mag), CoKu Tau 1 (XEST-23-000, an edge-on star-disk system, Padgett et al. 1999,  $A_V = 6.8$  mag), HN Tau (C5-2), LR 1 (XEST-23-048, with a large  $A_J = 6.4$  mag), V955 Tau (XEST-10-020,  $A_V = 3.7$  mag), V410 A20 (XEST-23-000, with a large  $A_J = 6.57$  mag), and V410 A24 (XEST-23-000, with a large  $A_J = 6.73$  mag – see Table 9 for  $A_V$  and  $A_J$  values).

The detection statistics of our X-ray survey is summarized in Table 12 (considering only the *XMM-Newton* observations), based on our final classification scheme in Table 11. Here, the protostar L1551 IRS5 has been treated as a non-detection as the source closely associated with this binary has been interpreted as X-ray emission from the jets (Favata et al. 2002; Bally et al. 2003). We also list the statistics (in parentheses) if *Chandra* detections and non-detections are counted (note that DG Tau B and FV Tau/c have been counted as non-detections in the *XMM-Newton* sample, while they are detections with *Chandra*). An important point for further statistical studies is that the X-ray sample of detected CTTS and WTTS is nearly complete for the surveyed fields (as far as the population is known). Most of the few remaining, undetected objects are considerably extinguished and by implication X-ray absorbed: IRAS S04301+261 = XEST-18-000 with  $A_J \approx 1.8$ , and CoKu Tau 1 = XEST-23-000, CFHT-Tau 19 = XEST-11-000, FV Tau/c = XEST-02-000, CFHT-Tau 20 = XEST-13-000, ITG33 A = XEST-07-000, CFHT-Tau 8 = XEST-07-000, and HH 30 = XEST-22-000 with  $A_V = 6.8, 7.3, 3.25, 3.6, 3.5, 1.8,$  and  $\approx 3$  mag and CFHT-Tau 12 = XEST-17-000 with  $A_J = 3.44$  mag, where the  $A_V$  of IRAS S04301+261, CoKu Tau 1, HH 30, ITG 33A may have been severely underestimated owing to scattered light and strong extinction by near-edge-on disks (see above). FV Tau/c was, however, detected in a *Chandra* observation (C3-1) reported here, with an unexpectedly high photoelectric absorption corresponding to  $N_{\text{H}} \approx 10^{23} \text{ cm}^{-2}$ . Some of the non-detections are very-low mass stars with late spectral types (CFHT-Tau 19 = XEST-11-000, CFHT-Tau 20 = XEST-13-000, CFHT-Tau 12 = XEST-17-000, and CFHT-Tau 8 = XEST-07-000 with spectral types of M5.25, M5.5, M6, and M5.5, respectively). Very little is known about IRAS 04108+2803 A. Its companion, a Class I object, was detected, its light curve showing a strong flare and its spectrum revealing very strong absorption. Perhaps, both objects are subject to similar absorption. The only formally undetected WTTS in the *XMM-Newton* sample is HBC 353, but it is located on the wings of HBC 352 = XEST-27-115, falls into a CCD gap of the PN detector, and is located far off-axis, elevating the corresponding detection limit to  $\approx 180$  cts. From Table 8, we derive that only 3 out of 45 multiple systems remain undetected (2/47 including *Chandra* results), while 29 out of 114 single stars are undetected (31/122 including *Chandra* results). In total, 45 of the 159 systems surveyed by *XMM-Newton* are multiple (47/169 including *Chandra* results).

Previous X-ray surveys of TMC did not detect the intrinsically fainter TTS population. Those previous surveys covered a wider field in Taurus (e.g., the ROSAT surveys by Neuhäuser et al. 1995; Stelzer & Neuhäuser 2001) but were about ten times less sensitive and were confined to soft sources. Stellar classes that were thus inaccessible by these earlier surveys but are

**Table 12.** XEST X-ray detection statistics.

Object type	Members surveyed	Detections	Detection fraction
0, 1 Protostars	20 (21)	8 (10)	40% (48%)
2 CTTS	65 (70)	55 (60)	85% (86%)
3 WTTS	50 (52)	49 (50)	98% (96%)
4 BDs	16 (17)	8 (9)	50% (53%)
5 Herbig	2	2	100%
9 others/unident.	6 (7)	4 (5)	67% (71%)
Total	159 (169)	126 (136)	79% (80%)

Notes:


Numbers in parentheses include *Chandra* observations. Source near L1551 IRS5 not considered (non-detection for XEST-22-040)

systematically (albeit not completely) detected in XEST are: i) strongly absorbed, embedded protostars or T Tau stars absorbed by their own disks seen nearly edge-on; their non-detection in previous surveys is little surprising given the strong photoelectric absorption. Most of the detected protostars show no X-ray counts below 2 keV. ii) Low- $L_X$  brown dwarfs; in XEST, the detection rate of BDs (50% for the *XMM-Newton* sample, 53% if one detection from *Chandra* is added) is high; the remaining objects of this class are likely to be intrinsically fainter than our detection limit rather than being excessively absorbed by gas ( $A_V$  of those objects typically being no more than a few magnitudes; see Grosso et al. 2007a for further discussion). iii) Several T Tau stars with double-peaked X-ray spectra in which a strongly absorbed hard component is accompanied by a soft excess (Güdel et al. 2007b).

The incomplete statistics in previous surveys potentially introduces bias into statistical correlations and population studies. On the other hand, we emphasize that the so far cataloged population of Taurus members may not be complete either. In particular, WTTS are predominantly detected by their X-rays (e.g., Neuhäuser et al. 1995). The WTTS statistics in Taurus may thus be incomplete in particular with regard to faint X-ray sources or WTTS located behind the TMC. Several dozen new TMC member candidates are indeed reported from XEST (Scelsi et al. 2007) but will require spectroscopic follow-up observations to confirm membership.

### 6.3. Statistical properties and correlations

Table 13 summarizes statistical properties of the surveyed sources, providing median, mean, and standard deviation of the sample for various parameters. The sample sizes vary (given in the last column, N) because not all parameters are available for all sources (see details in the footnotes of the table). The salient features of this compilation are the following: i) The agreement between the sample X-ray parameters derived from the DEM model (“DEM” in column 2) and the 1- $T$  or 2- $T$  model (“2- $T$ ” in column 2) is excellent. ii) Characteristic electron temperatures are around 6-8 MK as characterized by  $T_0$  in the DEM model and by  $T_1$  in the 1- $T$  or 2- $T$  model. The hotter component shows a mean and median around 23 MK. iii) The large majority of the sample shows  $-4 \lesssim \log L_X/L_* \lesssim -3$ . iv) Protostars are, as expected, significantly more extinguished in the visual band ( $A_V$ ) in our sample than T Tau stars; among the latter, objects of type 2 (CTTS, Class-II objects) are more strongly extinguished than objects of type 3 (WTTS, Class III). The median extinction of the detected sources is  $\approx 1.4$  mag, corresponding to a hydrogen column density of  $\log N_{\text{H}} \approx 21.44$ , close to the measured median of  $\log N_{\text{H}} \approx 21.5$ –21.6. iv) As pointed out on the



**Fig. 11.** HRD of all observed TMC objects for which  $L_*$  and  $T_{\text{eff}}$  are known. Symbol indicates object type (see panel in lower left corner for the key), and symbol size scales with  $\log L_X$ , crosses marking undetected sources. Additional sources from *Chandra* have been included, but note that the L1551 IRS5 protostar is marked here as a non-detection. Solid lines show evolutionary tracks toward the zero-age main sequence (ZAMS) for masses as given, in units of the solar mass (ZAMS). Dotted curves indicate isochrones at ages given (after Siess et al. 2000). The lowest of these curves marks the ZAMS. The spectral class ranges given at the bottom of the figure refer to main-sequence stars.

HRD, the  $L_*$  distributions of objects of type 1, 2, and 3 strongly overlap, although a marginal trend is seen toward a decreasing sample *mean* of  $L_*$  with increasing type, perhaps reflecting some overall evolutionary trend. v) A wide range of accretion rates is covered (for protostars and CTTS), with a mean and median around  $10^{-8} M_{\odot} \text{ yr}^{-1}$ .

In Fig. 12 we characterize the detected stellar population by plotting  $L_X$  (from the DEM models) as a function of mass  $M$  for all observed stars and substellar objects with known masses, although we exclude the following objects: DG Tau A (XEST-02-022), GV Tau (XEST-13-004), DP Tau (XEST-10-045), and CW Tau (XEST-20-046) have composite X-ray spectra that may originate from two unrelated sources (Güdel et al. 2007b). L1551 IRS5 is a deeply embedded protostar, while the X-ray source seen close to it is only lightly absorbed and may be related to the jet (Favata et al. 2002; Bally et al. 2003). Finally, the light curves of DH Tau (XEST-15-040), FS Tau (XEST-11-057), and V830 Tau (XEST-04-016) showed a decay presumably from large flares. For objects that were observed and detected twice, the logarithmic average of  $L_X$  has been adopted. Although for pre-main sequence stars there is no strict correlation between  $L_*$  and  $M$ , it is interesting that we find a rather well-developed correlation between  $L_X$  and  $M$ . The correlation coefficient for the  $\log L_X$  vs.  $\log M$  sample is  $C = 0.79$  for 99 data points. For the subsample of type 2 (“CTTS”) and type 3 objects (“WTTS”), we find  $C = 0.75$  and  $C = 0.85$  for 45 and 43 data points, respectively. All values point to a very significant correlation. We find a

linear regression,  $\log L_X = (1.54 \pm 0.12) \log M + 30.31 \pm 0.06$ . The slopes for type 2 and type 3 objects are, respectively,  $1.52 \pm 0.21$  and  $1.78 \pm 0.17$ . While these regressions are not significantly different, we note the larger errors for type 2 objects, and also their lower correlation coefficient, which is due to a larger scatter of  $L_X$  at a given mass. Similar trends have been noted in Orion (Preibisch et al. 2005), with  $\log L_X = (1.44 \pm 0.10) \log M + 30.37 \pm 0.06$  for the entire sample. This coincidence between XEST and COUP reveals that in TMC and ONC, *the basic X-ray production mechanism scales precisely the same way with the most basic property of the stellar objects, namely their mass*, regardless of any differences in age distribution, star-formation mode, or the presence or (near-) absence of high-mass stars. Part of this correlation might be explained by higher-mass stars being larger, i.e., providing more surface area for coronal active regions. Assuming similar average internal stellar density, the correlation between surface area and  $L_X$  is, however, considerably weaker than the trend shown in Fig. 12, so that additional effects may play a role (convection zone depth, convective mass, etc).

The figure shows another important feature of our survey. Almost all sources plotted are at  $L_X$  considerably higher than our approximate detection limit of  $\approx 10^{28} \text{ erg s}^{-1}$ . There are hardly any objects in the range of  $(1-5) \times 10^{28} \text{ erg s}^{-1}$  although they would have been detected. Exceptions are the brown dwarfs. This again testifies to the near-completeness of our survey within the observed fields.

**Table 13.** XEST parameter statistics.

Parameter	Sample <sup>a</sup>	Median	Mean	Std. dev. <sup>b</sup>	<i>N</i>
X-ray parameters					
$\beta^c$	DEM	-1.25	-1.49	0.95	99
$\log T_0^c$ [K]	DEM	6.90	6.90	0.28	119
$\log T_1^{c,d}$ [K]	2- <i>T</i>	6.79	6.76	0.21	94
$\log T_2^{c,d}$ [K]	2- <i>T</i>	7.37	7.38	0.25	105
$\log T_{av}^e$ [K]	DEM	7.04	7.03	0.24	123
	2- <i>T</i>	7.05	7.08	0.32	125
$\log N_H^d$ [cm <sup>-2</sup> ]	DEM	21.57	21.52	0.67	123
	2- <i>T</i>	21.53	21.50	0.65	123
$\log L_X^e$ [erg s <sup>-1</sup> ]	DEM	29.82	29.76	0.77	123
	2- <i>T</i>	29.80	29.72	0.78	125
$\log L_X/L_*^{e,f}$	DEM	-3.56	-3.55	0.50	116
	2- <i>T</i>	-3.56	-3.58	0.56	118
Fundamental parameters					
$A_V$ [cm <sup>-2</sup> ]	all	1.46	2.80	3.38	116
	1	10.20	9.52	1.93	6
	2	1.95	3.30	3.61	57
	3	0.87	1.36	1.56	38
	det	1.35	2.60	3.36	97
$\log L_*^{f,g}$ [L <sub>⊙</sub> ]	all	-0.34	-0.46	0.82	146
	1	0.08	-0.01	0.49	16
	2	-0.25	-0.33	0.61	60
	3	-0.43	-0.44	0.59	46
	det	-0.30	-0.33	0.71	119
$\log(\text{age})^{f,h}$ [Myr]	all	0.39	0.38	0.46	112
	1	0.05	0.23	0.54	9
	2	0.43	0.44	0.42	52
	3	0.39	0.32	0.35	43
$\log(\text{mass})^{f,g}$ [M <sub>⊙</sub> ]	all	-0.29	-0.37	0.43	123
	1	-0.17	-0.23	0.22	10
	2	-0.25	-0.27	0.31	54
	3	-0.39	-0.40	0.34	43
$\log(P)^i$ [d]	all	0.69	0.64	0.28	69
	1	0.63	0.54	0.32	10
	2	0.77	0.73	0.21	35
	3	0.58	0.56	0.33	22
$\log(\dot{M})^j$ [M <sub>⊙</sub> yr <sup>-1</sup> ]	all	-8.08	-8.20	1.21	57

Notes: Only results from *XMM-Newton* have been considered.

For multiple detections, parameters were averaged.

<sup>a</sup> DEM: from DEM fit; 2-*T*: from 1-*T* or 2-*T* fit; all; all objects; 1, 2, 3; object types; det = X-ray detected objects.

<sup>b</sup> Standard deviation of distribution.

<sup>c</sup> Only non-fixed fit parameters for detections considered.

<sup>d</sup> Only non-zero values for detections considered.

<sup>e</sup> Only detections considered.

<sup>f</sup> Only for known  $L_*$  above the ZAMS.


<sup>g</sup> For multiples, value for primary if available.

<sup>h</sup> For multiples, logarithmic average of components if available.


<sup>i</sup> Upper limits are adopted as measured values.

<sup>j</sup> Average of reported range if at least one non-upper limit given.

Figure 13 shows the distribution of the  $L_X$  (derived from the DEM models) as a function of the (stellar, photospheric) bolometric luminosity  $L_*$  for all spectrally modeled TTS and protostars, and also including BDs. We again exclude the peculiar spectral sources with two absorbers, the protostar L1551 IRS5, the three stars with gradually decaying light curves (DH Tau = XEST-15-040, FS Tau = XEST-11-057, V830 Tau = XEST-04-016), and also all sources located below the ZAMS in the HRD. For stars observed and detected twice, we again plot the logarithmic average of  $L_X$ . Most stars cluster between  $L_X/L_* = 10^{-4}$ – $10^{-3}$  as is often found in star-forming regions




**Fig. 12.** X-ray luminosity  $L_X$  (based on DEM fits) vs. stellar mass  $M$  for all detected XEST sources (*Chandra* sources and Herbig stars are not included), or upper limits (arrows). For multiple systems, the primary mass has been used if available. Symbols defining the object type are given in the lower right corner. The horizontal bars for some objects show the ranges of masses derived from literature  $L_*$  and  $T_{\text{eff}}$ , while the circle is centered at the adopted  $M$ . The straight line gives a linear regression using the logarithmic values for the X-ray detections, and the  $1\sigma$  ranges for the slopes:  $\log L_X = (1.54 \pm 0.12) \log M + 30.31 \pm 0.06$ .



**Fig. 13.** X-ray luminosity  $L_X$  (based on DEM fits) vs. stellar luminosity  $L_*$  for all X-ray detected XEST sources (*Chandra* sources, Herbig stars, and objects “below” the ZAMS not included), or upper limits (arrows). For multiple systems, total (“system”) luminosity  $L_*$  has been used. Symbols defining the object type are given in the lower right corner. The horizontal bars for some objects show the ranges of literature values for  $L_*$ , while the circle marks the adopted  $L_*$ .

(see Preibisch et al. 2005 for a similar analysis for the Orion sample). The value  $L_X/L_* = 10^{-3}$  corresponds to the saturation value for rapidly rotating main-sequence stars. One key parameter that drives the X-ray output is obviously  $L_*$ .

We finally present the X-ray luminosity function (XLF) of our Taurus sample in Fig. 14. The XLF has been calculated using the Kaplan-Meier estimator in the ASURV software package (Lavalley et al. 1992) that considers upper limits to  $L_X$  also. Non-detections for which no reasonable upper limit can be given were



**Fig. 14.** X-ray luminosity function for the entire sample (black), type 2 objects (red), and type 3 objects (blue).

not included. We also dropped the same non-standard X-ray sources excluded above (double-absorber spectra, L1551 IRS 5, and sources with gradual flare decays), but include BDs and Herbig stars, thus using 136 X-ray luminosity values, including 17 upper limits. We also plot separately the XLF for type 2 (CTTS, 56 values, including 6 upper limits) and type 3 (WTTS, 49 values, 1 upper limit). We note the shift of the CTTS distribution toward lower luminosities compared to WTTS, by a factor of about two. This is also evident for  $L_X/L_{bol}$  (Telleschi et al. 2007a). A two-sample test based on the Wilcoxon test and the logrank test performed in ASURV indicates a probability of only 6–9% that the two distributions are drawn from the same parent population. A more detailed discussion on distinctions between CTTS and WTTS is given by Telleschi et al. (2007a).

## 7. Outlook

The XEST project gives near-complete access to the entire pre-main sequence population of the Taurus Molecular Clouds within the  $\approx 5$  sq. degrees covered by this survey. Essentially all WTTS and nearly all CTTS have been detected, exceptions being a few low-mass, strongly absorbed sources. About half of the embedded protostellar sample is detected, and a similar fraction of the BDs. This survey thus goes deeper than previous X-ray surveys of Taurus (e.g., Neuhäuser et al. 1995; Stelzer & Neuhäuser 2001) by about an order of magnitude for low-extinction objects and provides systematic access to the protostellar and the brown-dwarf samples for the first time. The survey forms the basis for more specialized studies as discussed in the accompanying series of papers.

Furthermore, the survey will provide a useful database for correlation studies at other wavelengths. In particular, gas-to-dust ratio studies and investigations of the effects of accretion disks on magnetic fields in pre-main sequence systems will be undertaken in conjunction with a wide-field survey available from the *Spitzer Space Telescope* (Padgett et al. 2007).

**Acknowledgements.** The XEST project was made possible by the co-operation of many individuals. In particular, we thank the *XMM-Newton* SOC team in Vilsba for its excellent support before, during, and after the observations. We warmly thank the International Space Science Institute (ISSI) in Bern for their financial support of the project and their hospitality during several XEST team meetings at ISSI. Thanks go in particular to Vittorio Manno, Brigitte Fasler, and

Saliba F. Saliba for their efforts to make these events enjoyable. In the course of the scientific analysis, we obtained advice from numerous colleagues, of which we mention in particular Claude Catala, Laurence DeWarf, Ed Fitzpatrick, Sylvain Guieu, Antonio Maggio, and Karl Stapelfeldt. Francesco Damiani is thanked for giving us access to his private version of PWXDETECT for *XMM-Newton* data. We are grateful to the referee, Marc Gagné, for his competent review of this paper and his constructive comments that helped improve the paper. This research is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA member states and the USA (NASA). This publication makes use of data products from the Two Micron All Sky Survey (2MASS), which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. Further, our research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. We have made use of the ASURV statistical software package maintained by Penn State. X-ray astronomy research at PSI has been supported by the Swiss National Science Foundation (grants 20-66875.01 and 20-109255/1). MA acknowledges support by NASA grants NNG05GF92G. The Palermo group acknowledges financial contributions from contract ASI-INAF I/023/05/0.

## References

- Alexander, R. D., Clarke, C. J., & Pringle, J. E. 2004, *MNRAS*, 354, 71  
 Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197 (AG89)  
 Andrews, S. M., & Williams, J. P. 2005, *ApJ*, 631, 1134  
 Argiroffi, C., Drake, J. J., Maggio, A., et al. 2004, *ApJ*, 609, 925  
 Arnaud, K. A. 1996, in *ASP Conf. Ser.*, 101, *Astronomical Data Analysis Software and Systems V*, ed. G. Jacoby, & J. Barnes (San Francisco: ASP), 17  
 Arzner, K., Güdel, M., Briggs, K. R., Telleschi, A., & Audard, M. 2007a, *A&A*, 468, 477  
 Arzner, K., Güdel, M., Briggs, K. R., et al. 2007b, *A&A*, 468, 501  
 Audard, M., Briggs, K. R., Grosso, N., et al. 2007, *A&A*, 468, 379  
 Balbus, S. A., & Hawley J. F. 1991, *ApJ*, 376, 214  
 Bally, J., Feigelson, E., & Reipurth, B. 2003, *ApJ*, 584, 843  
 Barrado y Navascués, D., & Martín, E. L. 2003, *AJ*, 126, 2997  
 Berghoefer, T. W., Schmitt, J. H. M. M., & Cassinelli, J. P. 1996, *A&AS*, 118, 481  
 Böhm, T., & Catala, C. 1993, *A&AS*, 101, 629  
 Bouvier, J. 1990, *AJ*, 99, 946  
 Bouvier, J., 2006, private communication  
 Briceño, C., Calvet, N., Gomez, M., et al. 1993, *PASP*, 105, 686  
 Briceño, C., Hartmann, L., Stauffer, J., & Martín, E. 1998, *AJ*, 115, 2074  
 Briceño, C., Luhman, K. L., Hartmann, L., Stauffer, J. R., & Kirkpatrick, J. D. 2002, *ApJ*, 580, 317  
 Briggs, K. R., Güdel, M., Telleschi, A., et al. 2007, *A&A*, 468, 413  
 Burrows, C. J., Stapelfeldt, K. R., Watson, A. M., et al. 1996, *ApJ*, 473, 437  
 Calvet, N., Muzerolle, J., Briceño, C., et al. 2004, *AJ*, 128, 1294  
 Carpenter, J. M. 2000, *AJ*, 120, 3139  
 Cohen, M., & Kuhl, L. V. 1979, *ApJS*, 41, 743  
 Cutri, R. M., et al. 2003, *The IRSA 2MASS All-Sky Point Source Catalog*, NASA/IPAC Infrared Science Archive.  
<http://irsa.ipac.caltech.edu/applications/Gator/>  
 Dame, T. M., Ungerechts, H., Cohen, R. S., et al. 1987, *ApJ*, 322, 706  
 Damiani, F., & Micela, G. 1995, *ApJ*, 446, 341  
 Damiani, F., Micela, G., Sciortino, S., & Harden, F. R. Jr. 1995, *ApJ*, 446, 331  
 Damiani, F., Maggio, A., Micela, G., & Sciortino, S. 1997, *ApJ*, 483, 350  
 den Herder, J. W., Brinkman, A. C., Kahn, S. M., et al. 2001, *A&A*, 365, L7  
 DeWarf, L. E., & Fitzpatrick, E. 2006, private communication  
 DeWarf, L. E., Sepinsky, J. F., Guinan, E. F., Ribas, I., & Nadalin, I. 2003, *ApJ*, 590, 357  
 Dobashi, K., Uehara, H., Kandori, R., et al. 2005, *PASJ*, 57, 1  
 Dolan, C. J., & Mathieu, R. D. 2001, *AJ*, 121, 2124  
 Doppmann, G. W., Greene, T. P., Covey, K. R., & Lada, C. J. 2005, *AJ*, 130, 1145  
 Duchêne, G. 1999, *A&A*, 341, 547  
 Duchêne, G., Monin, J.-L., Bouvier, J., & Ménard, F. 1999, *A&A*, 351, 954  
 Duchêne, G., Ghez, A. M., & McCabe, C. 2002, *ApJ*, 568, 771  
 Duchêne, G., Ménard, F., Stapelfeldt, K., & Duvert, G. 2003, *A&A*, 400, 559  
 Duchêne, G., Bouvier, J., Bontemps, S., André, P., & Motte, F. 2004, *A&A*, 427, 651  
 Dutrey, A., Guilloteau, S., Duvert, G., et al. 1996, *A&A*, 309, 493  
 Favata, F., Fridlund, C. V. M., Micela, G., Sciortino, S., & Kaas, A. A. 2002, *A&A*, 386, 204  
 Favata, F., Giardino, G., Micela, G., Sciortino, S., & Damiani, F. 2003, *A&A*, 403, 187  
 Feigelson, E. D., & DeCampli, W. M. 1981, *ApJ*, 243, L89



- Feigelson, E. D., & Kriss, G. A. 1981, *Ap*, 248, L35
- Feigelson, E. D., & Montmerle, T. 1999, *ARA&A*, 37, 363
- Feigelson, E. D., Jackson, J. M., Mathieu, R. D., Myers, P. C., & Walter, F. M. 1987, *AJ*, 94, 1251
- Feigelson, E. D., Townsley, L., Güdel, M., & Stassun, K. 2007, *Protostars & Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil (Tucson: University of Arizona Press), 313
- Fernández, M., Ortiz, E., Eiroa, C., & Miranda, L. F. 1995, *A&AS*, 114, 439
- Finkenzeller, U., & Mundt, R. 1984, *A&AS*, 55, 109
- Flaccomio, E., Micela, G., Sciortino, S., et al. 2000, *A&A*, 355, 651
- Flaccomio, E., Micela, G., & Sciortino S. 2003a, *A&A*, 402, 277
- Flaccomio, E., Damiani, F., Micela, G., et al. 2003b, *ApJ*, 582, 398
- Franciosini, E., Pillitteri, I., Stelzer, B., et al. 2007, *A&A*, 468, 485
- Gagné, M., Skinner, S. L., & Daniel, K. J. 2004, *ApJ*, 613, 393
- García-Alvarez, D., Drake, J. J., Lin, L., Kashyap, V. L., & Ball, B. 2005, *ApJ*, 621, 1009
- Getman, K. V., Flaccomio, E., Broos, P. S., et al. 2005, *ApJS*, 160, 319
- Ghez, A. M., Neugebauer, G., & Matthews, K. 1993, *AJ*, 106, 2005
- Ghez, A. M., White, R. J., & Simon, M. 1997, *ApJ*, 490, 353
- Ghizzardi, S., et al. 2002, In-flight calibration of the PSF for the PN camera, XMM-SOC-CAL-TN-0029
- Glassgold, A. E., Najita, J., & Igea, J. 2004, *ApJ*, 615, 972
- Glauser, A. M., Ménard, F., Pinte, C., et al. 2007, *A&A*, submitted
- Grosso, N., Briggs, K. R., Güdel, M., et al. 2007a, *A&A*, 468, 391
- Grosso, N., Audard, M., Bouvier, J., et al. 2007b, *A&A*, 468, 557
- Güdel, M. 2002, *ARA&A*, 40, 217
- Güdel, M. 2004, *A&AR*, 12, 71
- Güdel, M., Audard, M., Kashyap, V. L., Drake, J. J., & Guinan, E. F. 2003, *ApJ*, 582, 483
- Güdel, M., Skinner, S. L., Briggs, K. R., et al. 2005, *ApJ*, 626, L53
- Güdel, M., Padgett, D. L., & Dougados, C. 2007a, *Protostars & Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil (Tucson: University of Arizona Press), 329
- Güdel, M., Telleschi, A., Audard, M., et al. 2007b, *A&A*, 468, 515
- Güdel, M., Skinner, S. L., Mel'nikov, S. Y., et al. 2007c, *A&A*, 468, 529
- Guiou, S. 2006, private communication
- Guiou, S., Dougados, C., Monin, J.-L., Magnier, E., & Martín, E. L. 2006, *A&A*, 446, 485
- Hartigan, P., & Kenyon, S. J. 2003, *ApJ*, 583, 334
- Hartigan, P., Strom, K. M., & Strom, S. E. 1994, *ApJ*, 427, 961
- Hartmann, L. 2002, *ApJ*, 578, 914
- Hartmann, L., Megeath, S. T., Allen, L., et al. 2005, *ApJ*, 629, 881
- Hayashi, M. R., Shibata, K., & Matsumoto, R. 1996, *ApJ*, 468, L37
- Herbig, G. H., Vrba, F. V., & Rydgren, A. E. 1986, *AJ*, 91, 575
- Itoh, Y., Hayashi, M., Tamura, M., et al. 2005, *ApJ*, 620, 984
- Jansen, F., Lumb, D., Altieri, B., et al. 2001, *A&A*, 365, L1
- Jensen, E. L., & Akeson, R. L. 2003, *ApJ*, 584, 875
- Kastner, J. H., Huenemoerder, D. P., Schulz, N. S., Canizares, C. R., & Weintraub, D. A. 2002, *ApJ*, 567, 434
- Kenyon, S. J., & Hartmann, L. 1995, *ApJS*, 101, 117
- Kenyon, S. J., Calvet, N., & Hartmann, L. 1993, *ApJ*, 414, 676
- Kenyon S. J., Dobrzycka D., & Hartmann L. 1994, *AJ*, 108, 1872
- Kenyon S. J., Brown, D. I., Tout, C. A., & Berlind, P. 1998, *AJ*, 115, 2491
- Kirsch, M. 2006, EPIC status of calibration and data analysis, XMM-SOC-CAL-TN-0018, <http://xmm.esac.esa.int/docs/documents/CAL-TN-0018.pdf>
- Kitamura, Y., Momose, M., Yokogawa, S., et al. 2002, *ApJ*, 581, 357
- Kraft, R. P., Burrows, D. N., & Nousek, J. A. 1991, *ApJ*, 374, 344
- Kun, M. 1998, *ApJS*, 115, 59
- Lavalley, M., Isobe, T., & Feigelson, E. 1992, in *Astronomical Data Analysis Software and Systems I*, ed. D. M. Worrall, C. Biemesderfer, & J. Barnes (San Francisco: ASP), 245
- Leinert, Ch., & Haas, M. 1989, *ApJ*, 342, L39
- Leinert, Ch., Zinnecker, H., Weitzel, N., et al. 1993, *A&A*, 278, 129
- Leinert, C., Richichi, A., & Haas, M. 1997, *A&A*, 318, 472
- Loinard, L., Mioduszewski, A. J., Rodríguez, L. F., et al. 2005, *ApJ*, 619, L179
- Luhman, K. L. 2000, *ApJ*, 544, 1044
- Luhman, K. L. 2004, *ApJ*, 617, 1216
- Luhman, K. L. 2006, *ApJ*, 645, 676
- Luhman, K. L., Briceño C., Stauffer, J. R., Hartmann, L., et al. 2003, *ApJ*, 590, 348
- Martín, E. L. 2000, *AJ*, 120, 2114
- Martín E. L., & Magazzù, A. 1999, *A&A*, 342, 173
- Martín, E. L., Rebolo, R., Magazzù, A., & Pavlenko, Ya. V. 1994, *A&A*, 282, 503
- Martín, E. L., Basri, G., & Zapatero Osorio, R. 1999, *AJ*, 118, 1005
- Martín, E. L., Dougados, C., Magnier, E., Ménard, F., Magazzù, A., Cuillandre, J.-C., & Delfosse, X. 2001, *ApJ*, 561, L195
- Mason, K. O., Breeveld, A., Much, R., et al. 2001, *A&A*, 365, L36
- McCabe, C., Ghez, A. M., Prato, L., et al. 2006, *ApJ*, 636, 932
- Mathieu, R. D. 1994, *ARA&A*, 32, 465
- Mohanty, S., Jayawardhana, R., & Basri, G. 2005, *ApJ*, 626, 498
- Moneti, A., & Zinnecker, H. 1991, *A&A*, 242, 438
- Monin, J.-L., Ménard, F., & Duchêne, G. 1998, *A&A*, 339, 113
- Montmerle, T., Grosso, N., Tsuboi, Y., & Koyama, K. 2000, *ApJ*, 532, 1097
- Morrison, R., & McCammon, D. 1983, *ApJ*, 270, 119
- Mundt, R., Walter, F. M., Feigelson, E. D., et al. 1983, *ApJ*, 269, 229
- Muzerolle, J., Hartmann, L., & Calvet, N. 1998, *AJ*, 116, 2965
- Muzerolle, J., Hillenbrand, L., Calvet, N., Briceño, C., & Hartmann, L. 2003, *ApJ*, 592, 266
- Muzerolle, J., Luhman, K. L., Briceño, C., Hartmann, L., & Calvet, N. 2005, *ApJ*, 625, 906
- Neuhäuser, R., Sterzik, M. F., Schmitt, J. H. M. M., Wichmann, R., & Krautter, J. 1995, *A&A*, 297, 391
- Ohashi, N., Hayashi, M., Kawabe, R., & Ishiguro, M. 1996, *ApJ*, 466, 317
- O'Dell, C. R. 1998, *AJ*, 115, 263
- Onishi, T., Mizuno, A., Kawamura, A., Tachihara, K., & Fukui, Y. 2002, *ApJ*, 575, 950
- Padgett, D. L., Brandner, W., Stapelfeldt, K. R., et al. 1999, *AJ*, 117, 1490
- Padgett, D. L., et al. 2007, in preparation
- Palla, F., & Stahler, S. W. 2002, *ApJ*, 581, 1194
- Petr, M. G., Coude Du Foresto, V., Beckwith, S. V. W., Richichi, A., & McCaughrean, M. J. 1998, *ApJ*, 500, 825
- Prato, L., & Simon, M. 1997, *ApJ*, 474, 455
- Pravdo, S. H., Feigelson, E. D., Garmire, G., et al. 2001, *Nature*, 413, 708
- Preibisch, T., & Zinnecker, H. 2001, *AJ*, 122, 866
- Preibisch, T., Kim, Y.-C., Favata, F., et al. 2005, *ApJS*, 160, 401
- Pudritz, R. E., & Norman, C. A. 1986, *ApJ*, 301, 571
- Qi, C., Kessler, J. E. Koerner, D. W., Sargent, A. I., & Blake, G. A. 2003, *ApJ*, 597, 986
- Read, A. M., & Ponman, T. J. 2003, *A&A*, 409, 395
- Rebull, L. M., Wolff, S. C., & Strom, S. E. 2004, *AJ*, 127, 1029
- Reipurth, B., & Zinnecker, H. 1993, *A&A*, 278, 81
- Reipurth, B., Rodríguez, L. F., Anglada, G., & Bally, J. 2002, *AJ*, 124, 1045
- Richichi, A., Leinert, Ch., Jameson, R., & Zinnecker, H. 1994, *A&A*, 287, 145
- Rodmann, J., Henning, Th., Chandler, C. J., Mundy, L. G., & Wilner, D. J. 2006, *A&A*, 466, 211
- Rodríguez, L. F., D'Alessio, P., Wilner, D. J., et al. 1998, *Nature*, 395, 355
- Scelsi, L., Maggio, A., Peres, G., & Pallavicini, R. 2005, *A&A*, 432, 671
- Scelsi, L., Maggio, A., Micela, G., et al. 2007, *A&A*, 468, 405
- Scholz, A., Jayawardhana, R., & Wood, K. 2006, *ApJ*, 645, 1498
- Shu, F., Najita, J., Ostriker, E., et al. 1994, *ApJ*, 429, 781
- Siess, L., Dufour, E., & Forestini, M. 2000, *A&A*, 358, 593
- Simon, M., Chen, W. P., Howell, R. R., & Slovick, D. 1992, *ApJ*, 384, 212
- Simon, M., Ghez, A. M., Leinert, Ch., et al. 1995, *ApJ*, 443, 625
- Simon, M., Holfeltz, S. T., & Taff, L. G. 1996, *ApJ*, 469, 890
- Smith, K. W., Balega, Y. Y., Duschl, W. J., et al. 2005, *A&A*, 431, 307
- Stelzer, B., & Neuhäuser, R. 2001, *A&A*, 377, 538
- Stelzer, B., Flaccomio, E., Micela, G., et al. 2007, *A&A*, 468, 463
- Strom, K. M., & Strom, S. E. 1994, *ApJ*, 424, 237
- Strom, K. M. Strom, S. E., Edwards, S., Cabrit, S., & Skrutskie, M. 1989, *AJ*, 97, 1451
- Strom, K. M., Strom, S. E., Wilkin, F. P., et al. 1990, *ApJ*, 362, 168
- Strüder, L., Briel, U., Dennerl, K., et al. 2001, *A&A*, 365, L18
- Telleschi, A., Güdel, M., Briggs, K., et al. 2005, *ApJ*, 622, 653
- Telleschi, A., Güdel, M., Briggs, K. R., Audard, M., & Palla, F. 2007a, *A&A*, 468, 425
- Telleschi, A., Güdel, M., Briggs, K. R., Audard, M., & Scelsi, L. 2007b, *A&A*, 468, 443
- Telleschi, A., Güdel, M., Briggs, K. R., et al. 2007c, *A&A*, 468, 541
- Turner, M. J. L., Abbey, A., Arnaud, M., et al. 2001, *A&A*, 365, L27
- Ungerechts, H., & Thaddeus, P. 1987, *ApJS*, 63, 645
- Vuong, M. H., Montmerle, T., Grosso, N., et al. 2003, *A&A*, 408, 581
- Walter, F. M., & Boyd, W. T. 1991, *ApJ*, 370, 318
- Walter, F. M., & Kuhl, L. V. 1981, *ApJ*, 250, 254
- Walter, F. M., Brown, A., Mathieu, R. D., Myers, P. C., & Vrba, F. J. 1988, *AJ*, 96, 297
- Walter, F. M., Beck, T. L., Morse, J. A., & Wolk, S. 2003, *AJ*, 125, 2123
- Weaver, W. B., & Jones, G. 1992, *ApJS*, 78, 239
- Welty, A. D. 1995, *5AJ*, 110, 776
- White, R. J., & Basri, G. 2003, *ApJ*, 582, 1109
- White, R. J., & Ghez, A. M. 2001, *ApJ*, 556, 265
- White, R. J., & Hillenbrand, L. A. 2004, 616, 998
- Woitak, J. 2003, *A&A*, 406, 685
- Woitak, J., Leinert, Ch., & Köhler, R. 2001, *A&A*, 376, 982
- Wolf, S., Padgett, D., & Stapelfeldt, K. 2003, *ApJ*, 588, 373
- Wolk, S. J., Harnden, F. R., Jr., Flaccomio, E., et al. 2005, *ApJS*, 160, 423

# Online Material

**References cited in the tables:**

- 1 = Andrews & Williams (2005)
- 2 = Bouvier (2006)
- 3 = Briceño et al. (1993)
- 4 = Briceño et al. (1998)
- 5 = Briceño et al. (2002)
- 6 = Böhm & Catala (1993)
- 7 = Cohen & Kuhl (1979)
- 8 = DeWarf et al. (2003)
- 9 = DeWarf & Fitzpatrick (2006)
- 10 = Doppmann et al. (2005)
- 11 = Duchêne et al. (1999)
- 12 = Duchêne et al. (2002)
- 13 = Duchêne et al. (2004)
- 14 = Fernández et al. (1995)
- 15 = Finkenzeller & Mundt (1984)
- 16 = Ghez et al. (1993)
- 17 = Ghez et al. (1997)
- 18 = Guieu et al. (2006)
- 19 = Guieu (2006)
- 20 = Hartigan & Kenyon (2003)
- 21 = Hartigan et al. (1994)
- 22 = Hartmann (2002)
- 23 = Hartmann et al. (2005)
- 24 = Herbig et al. (1986)
- 25 = Itoh et al. (2005)
- 26 = Jensen & Akeson (2003)
- 27 = Kenyon & Hartmann (1995)
- 28 = Kenyon et al. (1993)
- 29 = Kenyon et al. (1998)
- 30 = Leinert & Haas (1989)
- 31 = Leinert et al. (1993)
- 32 = Leinert et al. (1997)
- 33 = Luhman (2004)
- 34 = Luhman et al. (2003)
- 35 = Martín (2000)
- 36 = Martín & Magazzù (1999)
- 37 = Martín et al. (1994)
- 38 = Martín et al. (1999)
- 39 = Martín et al. (2001)
- 40 = McCabe et al. (2006)
- 41 = Mohanty et al. (2005)
- 42 = Moneti & Zinnecker (1991)
- 43 = Monin et al. (1998)
- 44 = Muzerolle et al. (1998)
- 45 = Muzerolle et al. (2003)
- 46 = Muzerolle et al. (2005)
- 47 = Prato & Simon (1997)
- 48 = Rebull et al. (2004) and references therein
- 49 = Reipurth & Zinnecker (1993)
- 50 = Richichi et al. (1994)
- 51 = Rodríguez et al. (1998)
- 52 = Simon et al. (1992) and references therein
- 53 = Simon et al. (1995)
- 54 = Smith et al. (2005)
- 55 = Strom & Strom (1994)
- 56 = Walter et al. (1988)
- 57 = White & Basri (2003)
- 58 = White & Ghez (2001)
- 59 = White & Hillenbrand (2004)
- 60 = Mundt et al. (1983)
- 61 = Welty (1995)
- 62 = Woitas (2003)
- 63 = Calvet et al. (2004)
- 64 = Reipurth et al. (2002)
- 65 = Walter et al. (2003) and references therein
- 66 = Luhman (2006)
- 67 = Simon et al. (1996)

**Table 3.** Accepted good exposure time and fraction of total exposure time remaining after filtering out intervals of high background in the full energy band (0.5–7.3 keV). Calculated for each EPIC instrument in each *XMM-Newton* observation used in XEST. The penultimate column lists the approximate lowest detectable count-rate for a  $4\sigma$  detection in the PN; for a 30 ks exposure in purely quiescent background, this would be  $1.1 \text{ c ks}^{-1}$ . For XEST-26 only the MOS cameras were used and the MOS1 value has been multiplied by a factor 2.7 to calculate a PN-equivalent source count-rate. The final column lists the total number of detections made in the summed EPIC images.

Field	M1		M2		PN		$C_{4\sigma}$ $\text{c ks}^{-1}$	EPIC $N_{\text{det}}$
	$t_{\text{good}}$ s	Per cent	$t_{\text{good}}$ s	Per cent	$t_{\text{good}}$ s	Per cent		
01	50 215	62.9	50 944	63.9	41 756	57.2	1.06	104
02	39 078	95.4	39 638	96.8	27 895	71.0	1.63	76
03	22 814	65.2	23 908	68.3	15 647	47.1	1.81	37
04	29 722	89.9	29 710	89.9	29 131	95.0	1.19	70
05	24 125	76.3	24 229	76.6	15 963	53.9	1.86	36
06	30 777	97.4	30 894	97.7	27 858	92.9	1.30	103
07	30 013	95.1	30 183	95.6	25 338	84.2	1.29	52
08	35 577	85.6	35 841	86.2	26 699	71.7	1.66	82
09	25 009	79.0	25 170	79.5	19 272	64.3	1.82	81
10	28 597	90.3	28 649	90.5	25 476	84.8	1.31	66
11	34 245	82.5	35 075	84.2	30 972	80.7	1.14	104
12	30 398	97.9	30 451	98.1	28 673	97.4	1.17	90
13	24 561	77.8	24 979	79.1	20 918	70.5	3.11	37
14	28 832	84.6	28 942	84.9	24 173	74.9	1.65	78
15	23 372	75.5	23 475	75.9	19 166	63.1	1.67	84
16	23 864	58.9	25 683	63.3	13 705	40.2	2.54	46
17	26 750	86.7	26 804	86.9	22 898	78.3	1.59	68
18	28 540	90.1	28 654	90.5	26 500	89.8	1.24	66
19	30 235	80.0	30 544	80.8	26 086	75.4	1.29	90
20	31 494	100.0	31 479	100.0	29 877	100.0	1.18	76
21	29 060	74.9	29 094	75.4	18 386	40.4	2.03	75
22	55 031	96.8	54 931	96.6	50 484	92.7	0.76	116
23	57 519	79.9	57 496	79.8	41 190	59.5	1.11	76
24	32 069	72.6	31 991	72.4	24 253	60.2	1.57	63
25	16 411	97.8	16 466	98.1	14 399	94.9	1.84	54
26	124 995	97.4	125 270	97.6	–	–	0.81	165
27	37 770	87.3	27 864	87.5	35 416	85.2	1.16	153
28	124 833	95.4	124 951	95.5	120 693	93.1	0.52	199

**Table 4.** X-ray parameters of targets in XEST (1): Positions and count rates.

XEST	Name	RA <sub>X</sub>			Dec <sub>X</sub>		Poserr ( $''$ )	Offset ( $''$ )	ML <sub>det</sub> <sup>a</sup>	Scts	T <sub>exp</sub> (s)	Rate (ct s <sup>-1</sup> )	Var <sup>b</sup>	
		h	m	s	deg	'								''
27-115	HBC 352	3	54	29.54	32	03	02.2	1.04	0.89	9395	3832	15 780	0.2429	0
27-000	HBC 353	3	54	30.17	32	03	04.3	0.00	0.00	0	<183	14 469	<0.0127	0
06-005	HBC 358 AB	4	03	49.27	26	10	53.1	1.48	1.22	1073	879	7371	0.1194	0
06-007	HBC 359	4	03	50.82	26	10	53.0	1.46	0.34	2921	1605	7668	0.2094	0
06-059	L1489 IRS	4	04	43.07	26	18	56.3	1.46	0.10	7632	3336	46 889	0.0712	0
20-001	LkCa 1	4	13	14.03	28	19	09.9	1.58	1.71	455	336	7820	0.0430	0
20-005	Anon 1	4	13	27.28	28	16	23.3	1.53	1.64	24 658	8299	21 632	0.3836	0
20-000	IRAS 04108+2803 A	4	13	53.29	28	11	23.4	0.00	0.00	0	<26	29 957	<0.0009	0
20-022	IRAS 04108+2803 B	4	13	54.72	28	11	32.2	1.55	0.70	1145	696	30 670	0.0227	1
20-000	2M J04141188+28	4	14	11.88	28	11	53.5	0.00	0.00	0	<54	48 645	<0.0011	0
20-042	V773 Tau ABC	4	14	12.92	28	12	11.8	1.53	0.60	213 164	59 321	48 974	1.2113	0
20-043	FM Tau	4	14	13.57	28	12	48.6	1.54	0.61	2627	2988	48 901	0.0611	0
20-046	CW Tau	4	14	16.99	28	11	00.6	1.84	2.80	11	52	35 943	0.0015	0
20-047	CIDA 1	4	14	17.72	28	06	09.8	1.69	1.46	63	105	33 985	0.0031	0
20-056	MHO 2/1	4	14	26.41	28	06	00.9	1.54	1.21	6260	2735	32 359	0.0845	0
20-058	MHO 3	4	14	30.58	28	05	14.7	1.55	0.40	1234	766	26 856	0.0285	0
20-069	FO Tau AB	4	14	49.23	28	12	31.3	1.60	1.06	238	240	33 352	0.0072	0
20-073	CIDA 2	4	15	05.16	28	08	46.5	1.56	0.30	814	584	22 342	0.0261	0
23-002	CY Tau	4	17	33.61	28	20	47.7	1.08	1.77	683	457	10 256	0.0446	1
24-002	CY Tau	4	17	33.73	28	20	47.9	1.01	1.00	407	284	8366	0.0340	2
23-004	LkCa 5	4	17	38.91	28	32	59.6	1.04	0.98	5286	2855	32 431	0.0880	0
24-004	LkCa 5	4	17	38.94	28	33	00.2	0.94	0.30	4767	2221	18 626	0.1192	2
23-008	CIDA 3	4	17	49.67	28	29	35.9	1.10	0.48	292	358	45 960	0.0078	0
24-008	CIDA 3	4	17	49.59	28	29	35.3	0.99	1.28	338	332	26 423	0.0126	0
23-015	V410 X3	4	18	07.98	28	26	02.8	1.09	0.94	576	561	64 938	0.0086	0
24-015	V410 X3	4	18	07.97	28	26	02.1	0.99	1.61	388	341	37 758	0.0090	0
23-018	V410 A13	4	18	17.12	28	28	40.5	1.45	1.41	20	72	66 746	0.0011	0
24-000	V410 A13	4	18	17.11	28	28	41.9	0.00	0.00	0	<24	34 966	<0.0007	0
23-000	V410 A24	4	18	22.39	28	24	37.6	0.00	0.00	0	<26	68 500	<0.0004	0
24-000	V410 A24	4	18	22.39	28	24	37.6	0.00	0.00	0	<30	42 210	<0.0007	0
23-029	V410 A25	4	18	29.15	28	26	18.9	1.04	0.69	4542	3586	76 359	0.0470	0
24-027	V410 A25	4	18	29.12	28	26	19.1	0.94	0.26	2447	1812	43 216	0.0419	0
23-032	V410 Tau ABC	4	18	31.08	28	27	16.7	1.03	0.57	365 212	101 132	78 334	1.2910	1
24-028	V410 Tau ABC	4	18	31.09	28	27	16.8	0.93	0.61	137 848	38 215	44 779	0.8534	3
23-033	DD Tau AB	4	18	31.11	28	16	28.7	1.08	0.40	1006	885	39 614	0.0223	1
24-029	DD Tau AB	4	18	31.10	28	16	29.3	1.04	0.50	295	320	22 841	0.0140	0
23-035	CZ Tau AB	4	18	31.60	28	16	59.2	1.05	0.71	1756	1271	41 620	0.0305	0
24-030	CZ Tau AB	4	18	31.57	28	16	59.2	0.97	0.75	1509	995	23 898	0.0417	0
23-036	IRAS 04154+2823	4	18	31.94	28	31	15.6	1.11	1.20	211	308	64 773	0.0048	1
24-031	IRAS 04154+2823	4	18	31.88	28	31	14.9	1.59	2.04	18	61	36 863	0.0017	0
23-037	V410 X2	4	18	34.40	28	30	30.1	1.04	0.67	5641	3007	65 004	0.0463	0
24-032	V410 X2	4	18	34.38	28	30	30.6	0.94	1.01	3959	2015	36 402	0.0554	1
23-045	V410 X4	4	18	40.24	28	24	25.1	1.06	0.61	859	760	69 092	0.0110	0
24-038	V410 X4	4	18	40.26	28	24	25.2	0.95	0.80	1153	778	39 074	0.0199	2
23-047	V892 Tau	4	18	40.64	28	19	15.9	1.03	0.48	58 194	20 108	48 862	0.4115	0
24-040	V892 Tau	4	18	40.62	28	19	15.6	0.93	0.10	111 462	31 125	28 058	1.1093	1
23-048	LR 1	4	18	41.29	28	27	26.9	1.22	1.97	45	134	69 240	0.0019	1
24-000	LR 1	4	18	41.33	28	27	25.0	0.00	0.00	0	<67	41 278	<0.0016	0
23-050	V410 X7	4	18	42.53	28	18	49.9	1.04	0.41	2808	2319	46 487	0.0499	1
24-042	V410 X7	4	18	42.49	28	18	50.0	0.93	0.24	10 724	6309	26 634	0.2369	1
23-000	V410 A20	4	18	45.06	28	20	52.8	0.00	0.00	0	<23	32 774	<0.0007	0
24-000	V410 A20	4	18	45.06	28	20	52.8	0.00	0.00	0	<15	17 241	<0.0009	0
23-056	Hubble 4	4	18	47.03	28	20	07.7	1.03	0.42	95 259	27 648	45 560	0.6069	0
24-047	Hubble 4	4	18	47.02	28	20	07.7	0.93	0.48	43 119	14 030	26 926	0.5211	0
23-000	KPNO-Tau 2	4	18	51.16	28	14	33.2	0.00	0.00	0	<28	15 428	<0.0018	0
24-000	KPNO-Tau 2	4	18	51.16	28	14	33.2	0.00	0.00	0	<15	8642	<0.0018	0
23-000	CoKu Tau 1	4	18	51.48	28	20	26.5	0.00	0.00	0	<128	48 084	<0.0027	0
24-000	CoKu Tau 1	4	18	51.48	28	20	26.5	0.00	0.00	0	<57	27 535	<0.0021	0
23-061	V410 X6	4	19	01.07	28	19	42.1	1.10	0.54	298	369	38 775	0.0095	0

Table 4. continued.

XEST	Name	RA <sub>X</sub>			Dec <sub>X</sub>		Poserr ( $''$ )	Offset ( $''$ )	ML <sub>det</sub> <sup>a</sup>	Scts	T <sub>exp</sub> (s)	Rate (ct s <sup>-1</sup> )	Var <sup>b</sup>
		h	m	s	deg	'							
24-054	V410 X6	4 19 01.08	28 19 42.2	0.98	0.44	514	442	22 047	0.0201	2			
23-063	V410 X5	4 19 01.94	28 22 34.4	1.05	1.31	1688	1179	43 645	0.0270	0			
24-055	V410 X5	4 19 01.94	28 22 33.8	0.95	0.80	2422	1339	24 869	0.0539	2			
23-067	FQ Tau AB	4 19 12.77	28 29 33.3	1.21	0.56	100	178	34 347	0.0052	0			
24-058	FQ Tau AB	4 19 12.77	28 29 34.3	1.20	1.31	64	101	18 919	0.0054	0			
28-100	BP Tau	4 19 15.90	29 06 26.6	2.24	0.84	142 902	46 278	197 539	0.2343	1			
23-074	V819 Tau AB	4 19 26.26	28 26 14.5	1.04	0.20	19 437	7785	28 888	0.2695	0			
24-061	V819 Tau AB	4 19 26.26	28 26 14.7	0.94	0.40	9399	3967	16 434	0.2414	0			
16-000	IRAS 04166+2706	4 19 43.00	27 13 33.7	0.00	0.00	0	<30	30 078	<0.0010	0			
16-000	IRAS 04169+2702	4 19 58.45	27 09 57.1	0.00	0.00	0	<38	23 883	<0.0016	0			
11-000	CFHT-Tau 19	4 21 07.95	27 02 20.4	0.00	0.00	0	<18	22 031	<0.0008	0			
11-000	IRAS 04181+2655	4 21 10.90	27 02 06.0	0.00	0.00	0	<18	22 755	<0.0008	0			
11-000	IRAS 04181+2654AB	4 21 11.47	27 01 09.4	0.00	0.00	0	<34	24 662	<0.0014	0			
11-023	2M J04213459	4 21 34.51	27 01 38.7	1.88	1.22	140	171	38 717	0.0044	0			
01-028	IRAS 04187+1927	4 21 43.23	19 34 12.5	1.55	0.81	2842	1460	30 378	0.0481	0			
11-037	CFHT-Tau 10	4 21 46.39	26 59 28.4	2.25	1.61	9	35	48 347	0.0007	0			
11-000	2M J04215450+2652	4 21 54.51	26 52 31.5	0.00	0.00	0	<17	37 210	<0.0004	0			
21-038	RY Tau	4 21 57.40	28 26 35.4	1.36	0.10	17 357	6495	23 482	0.2766	0			
21-039	HD 283572	4 21 58.87	28 18 06.1	1.36	0.64	282 825	76 762	37 198	2.0636	0			
01-045	T Tau N(+Sab)	4 21 59.44	19 32 05.8	1.54	0.62	211 562	59 050	72 108	0.8189	0			
11-054	Haro 6-5 B	4 22 00.72	26 57 32.4	1.88	0.29	28	188	51 812	0.0036	0			
11-057	FS Tau AC	4 22 02.19	26 57 30.9	1.78	0.42	20 767	7398	51 050	0.1449	3			
21-044	LkCa 21	4 22 03.12	28 25 39.1	1.36	0.28	6256	2932	25 588	0.1146	0			
01-054	RX J0422.1+1934	4 22 04.91	19 34 48.8	1.54	0.87	54 106	17 801	52 504	0.3390	0			
01-062	2M J04221332+1934	4 22 13.24	19 34 40.2	1.72	1.51	69	117	33 836	0.0035	0			
11-079	CFHT-Tau 21	4 22 16.80	26 54 58.2	1.82	1.22	286	274	39 533	0.0070	0			
02-013	FV Tau AB	4 26 53.48	26 06 54.9	1.60	0.84	781	662	31 329	0.0212	0			
02-000	FV Tau/c AB	4 26 54.41	26 06 51.0	0.00	0.00	0	<94	35 140	<0.0027	0			
02-016	KPNO-Tau 13	4 26 57.34	26 06 27.2	1.63	1.21	217	288	22 911	0.0126	0			
02-000	DG Tau B	4 27 02.66	26 05 30.5	0.00	0.00	0	<27	38 087	<0.0007	0			
02-022	DG Tau A	4 27 04.70	26 06 15.5	1.59	0.80	1669	1185	40 865	0.0290	2			
02-000	KPNO-Tau 4	4 27 28.00	26 12 05.3	0.00	0.00	0	<41	50 463	<0.0008	0			
02-000	IRAS 04248+2612AB	4 27 57.31	26 19 18.3	0.00	0.00	0	<25	17 943	<0.0014	0			
15-020	JH 507	4 29 20.67	26 33 40.2	1.56	0.73	2000	1102	28 851	0.0382	0			
13-004	GV Tau AB	4 29 23.69	24 33 00.2	1.63	0.55	518	615	20 976	0.0293	1			
13-000	IRAS 04264+2433	4 29 30.08	24 39 55.1	0.00	0.00	0	<68	26 868	<0.0025	0			
15-040	DH Tau AB	4 29 41.56	26 32 58.5	1.55	0.20	130 666	36 137	33 680	1.0729	3			
15-042	DI Tau AB	4 29 42.51	26 32 49.3	1.56	0.40	926	3305	33 773	0.0979	0			
15-044	KPNO-Tau 5	4 29 45.81	26 30 47.9	1.82	2.06	40	86	32 576	0.0026	0			
14-006	IQ Tau A	4 29 51.56	26 06 44.9	2.02	0.00	3811	1972	22 748	0.0867	1			
13-000	CFHT-Tau 20	4 29 59.51	24 33 07.9	0.00	0.00	0	<22	30 935	<0.0007	0			
14-000	KPNO-Tau 6	4 30 07.24	26 08 20.8	0.00	0.00	0	<27	26 206	<0.0010	0			
13-035	FX Tau AB	4 30 29.59	24 26 47.2	1.65	2.22	313	378	7123	0.0532	0			
14-057	DK Tau AB	4 30 44.25	26 01 25.5	2.01	1.00	8459	3697	37 386	0.0989	0			
14-000	KPNO-Tau 7	4 30 57.19	25 56 39.5	0.00	0.00	0	<31	25 489	<0.0012	0			
22-013	MHO 9	4 31 15.83	18 20 05.9	1.62	1.48	777	611	37 684	0.0162	0			
22-021	MHO 4	4 31 24.18	18 00 21.6	1.63	1.71	700	569	37 887	0.0150	0			
22-040	L1551 IRS5	4 31 34.02	18 08 04.2	1.68	1.11	139	191	74 201	0.0026	0			
22-042	LkHa 358	4 31 36.13	18 13 43.3	1.66	0.00	115	273	76 104	0.0036	0			
22-000	HH 30	4 31 37.47	18 12 24.5	0.00	0.00	0	<21	81 198	<0.0003	0			
22-043	HL Tau	4 31 38.38	18 13 57.1	1.60	1.04	6052	5618	75 594	0.0743	0			
22-047	XZ Tau AB	4 31 40.05	18 13 56.6	1.60	0.66	60 086	18 391	75 388	0.2440	1			
22-056	L1551 NE	4 31 44.53	18 08 32.3	1.98	1.39	13	47	80 720	0.0006	0			
03-005	HK Tau AB	4 31 50.57	24 24 16.1	1.77	2.00	60	86	19 411	0.0045	0			
22-070	V710 Tau BA	4 31 57.72	18 21 37.2	1.60	1.34	11 966	4639	35 159	0.1320	0			
19-009	JH 665	4 31 58.50	25 43 30.6	2.27	1.07	108	123	8791	0.0141	0			
22-089	L1551 51	4 32 09.45	17 57 22.9	1.60	2.57	11 539	4287	13 801	0.3107	2			
22-097	V827 Tau	4 32 14.50	18 20 14.5	1.60	1.02	36 001	10 530	18 303	0.5753	0			
03-016	Haro 6-13	4 32 15.40	24 28 59.9	1.59	0.24	338	288	20 435	0.0141	0			

**Table 4.** continued.

XEST	Name	RA <sub>X</sub>			Dec <sub>X</sub>		Poserr ( $''$ )	Offset ( $''$ )	ML <sub>det</sub> <sup>a</sup>	Scts	T <sub>exp</sub> (s)	Rate (ct s <sup>-1</sup> )	Var <sup>b</sup>
		h	m	s	deg	'							
22-100	V826 Tau	4 32	15.91	18 01	39.1	1.60	1.08	94 853	27 095	36 121	0.7501	0	
22-101	MHO 5	4 32	16.07	18 12	46.5	1.62	0.10	1457	963	51 887	0.0186	0	
03-017	CFHT-Tau 7	4 32	17.83	24 22	13.1	1.71	1.94	29	121	30 227	0.0040	0	
03-019	V928 Tau AB	4 32	18.90	24 22	26.1	1.54	1.14	4972	2243	30 636	0.0732	0	
03-022	FY Tau	4 32	30.68	24 19	58.1	1.55	1.58	4931	2110	25 663	0.0822	0	
03-023	FZ Tau	4 32	31.83	24 20	04.1	1.59	1.46	213	440	19 003	0.0232	0	
17-002	IRAS 04295+2251	4 32	32.07	22 57	26.4	1.36	0.41	322	303	17 292	0.0176	2	
19-049	UZ Tau E+W(AB)	4 32	42.89	25 52	32.6	2.19	2.52	4110	2085	44 868	0.0465	0	
17-009	JH 112	4 32	49.09	22 53	01.9	1.33	0.94	1517	867	24 276	0.0357	0	
03-031	CFHT-Tau 5	4 32	50.28	24 22	11.4	1.73	0.24	54	82	15 890	0.0052	0	
04-003	CFHT-Tau 5	4 32	50.31	24 22	11.1	1.73	0.74	367	312	26 369	0.0119	0	
03-035	MHO 8	4 33	01.97	24 21	02.7	1.66	2.70	105	130	17 386	0.0075	0	
04-009	MHO 8	4 33	02.02	24 20	59.0	1.77	1.14	181	172	28 001	0.0062	0	
04-010	GH Tau AB	4 33	06.33	24 09	32.5	1.78	2.13	96	169	11 180	0.0151	0	
04-012	V807 Tau SNab	4 33	06.75	24 09	54.6	1.68	1.56	10 380	4068	20 068	0.2027	0	
18-004	KPNO-Tau 14	4 33	07.79	26 16	05.7	2.13	0.94	1424	840	19 063	0.0441	0	
17-000	CFHT-Tau 12	4 33	09.46	22 46	48.7	0.00	0.00	0	<10	30 963	<0.0003	0	
04-016	V830 Tau	4 33	09.98	24 33	43.4	1.68	0.68	44 530	13 508	15 863	0.8516	3	
18-000	IRAS S04301+261	4 33	14.36	26 14	23.5	0.00	0.00	0	<17	22 545	<0.0008	0	
17-000	IRAS 04302+2247	4 33	16.50	22 53	20.4	0.00	0.00	0	<28	29 406	<0.0010	0	
17-027	IRAS 04303+2240	4 33	19.10	22 46	34.1	1.31	0.43	9238	3723	30 569	0.1218	0	
04-034	GI Tau	4 33	34.06	24 21	17.9	1.69	0.90	792	1707	46 957	0.0364	0	
04-035	GK Tau AB	4 33	34.52	24 21	08.0	1.68	2.17	17 201	6635	47 269	0.1404	0	
18-019	IS Tau AB	4 33	36.83	26 09	48.3	2.13	1.05	3999	1835	31 294	0.0586	0	
17-058	CI Tau	4 33	52.02	22 50	30.8	1.34	0.66	647	485	23 775	0.0204	1	
18-030	IT Tau AB	4 33	54.72	26 13	26.7	2.12	0.84	58 042	17 849	44 176	0.4041	0	
17-066	JH 108	4 34	11.01	22 51	44.7	1.32	0.34	9235	3875	14 379	0.2695	1	
17-068	CFHT-BD Tau 1	4 34	15.31	22 50	33.1	1.45	2.17	163	204	14 010	0.0146	2	
25-026	AA Tau	4 34	55.39	24 28	53.6	1.82	0.57	3334	1471	24 628	0.0597	0	
09-010	HO Tau AB	4 35	20.31	22 32	11.1	1.58	3.82	82	122	18 311	0.0067	0	
08-019	FF Tau AB	4 35	20.90	22 54	24.3	1.60	0.10	3812	2076	30 501	0.0681	0	
12-040	DN Tau	4 35	27.34	24 14	58.9	1.57	0.41	28 146	9246	47 293	0.1955	0	
12-000	IRAS 04325+2402AB	4 35	35.39	24 08	19.4	0.00	0.00	0	<44	34 141	<0.0013	0	
12-059	CoKu Tau 3 AB	4 35	40.95	24 11	09.0	1.57	0.24	69 838	20 015	38 317	0.5224	0	
09-022	KPNO-Tau 8	4 35	41.91	22 34	10.9	1.44	1.20	3919	1858	27 822	0.0668	0	
08-037	HQ Tau AB	4 35	47.39	22 50	22.0	1.59	0.75	25 465	9258	30 355	0.3050	0	
09-026	HQ Tau AB	4 35	47.33	22 50	20.9	1.43	0.81	20 604	7171	14 030	0.5111	3	
08-043	KPNO-Tau 15	4 35	51.13	22 52	40.1	1.59	0.41	25 254	10 329	46 282	0.2232	0	
09-031	KPNO-Tau 15	4 35	51.01	22 52	40.3	1.49	1.26	633	529	11 974	0.0443	0	
08-000	KPNO-Tau 9	4 35	51.43	22 49	11.9	0.00	0.00	0	<28	37 232	<0.0008	0	
09-000	KPNO-Tau 9	4 35	51.43	22 49	11.9	0.00	0.00	0	<26	17 141	<0.0016	0	
08-048	HP Tau AB	4 35	52.78	22 54	22.9	1.59	0.20	3573	5706	49 366	0.1156	0	
08-051a	HP Tau/G3 AB	4 35	53.50	22 54	09.0	0.00	0.00	134 269	3006	48 905	0.0648	0	
08-051	HP Tau/G2	4 35	54.14	22 54	12.8	1.59	0.71	134 269	39 787	48 905	0.8136	0	
08-058	Haro 6-28 AB	4 35	56.80	22 54	37.1	1.63	1.23	290	614	49 680	0.0124	0	
08-000	CFHT-BD Tau 2	4 36	10.39	22 59	56.0	0.00	0.00	0	<29	36 650	<0.0008	0	
08-080	CFHT-BD Tau 3	4 36	38.91	22 58	13.2	1.86	1.36	20	64	25 098	0.0026	0	
05-005	CFHT-Tau 6	4 39	04.10	25 44	26.4	1.90	1.89	48	84	21 111	0.0040	0	
05-000	IRAS 04 361+2547	4 39	13.89	25 53	20.9	0.00	0.00	0	<35	11 917	<0.0030	0	
05-013	GN Tau AB	4 39	20.86	25 45	00.9	1.77	1.38	301	248	22 952	0.0108	0	
05-017	IRAS 04365+2535	4 39	35.25	25 41	45.4	1.93	1.07	34	75	31 546	0.0024	0	
05-024	IRAS 04369+2539	4 39	55.70	25 45	01.7	1.75	0.74	435	335	20 227	0.0166	0	
07-011	JH 223	4 40	49.54	25 51	20.0	1.87	0.90	374	318	24 335	0.0131	0	
07-022	Haro 6-32	4 41	04.32	25 57	55.5	1.89	1.23	351	311	17 222	0.0181	0	
07-000	ITG 33 A	4 41	08.26	25 56	07.5	0.00	0.00	0	<21	22 254	<0.0010	0	
07-000	CFHT-Tau 8	4 41	10.78	25 55	11.7	0.00	0.00	0	<16	23 759	<0.0007	0	
07-000	IRAS 04381+2540	4 41	12.68	25 46	35.4	0.00	0.00	0	<32	44 259	<0.0007	0	
07-041	IRAS 04385+2550AB	4 41	38.80	25 56	27.4	1.86	0.66	503	397	16 994	0.0234	2	
10-017	CoKuLk332/G2 AB	4 42	05.47	25 22	56.2	1.68	0.29	17 061	6651	36 423	0.1826	0	

Table 4. continued.

XEST	Name	RA <sub>X</sub>			Dec <sub>X</sub> deg ' ''	Poserr ('')	Offset ('')	ML <sub>det</sub> <sup>a</sup>	Scts	T <sub>exp</sub> (s)	Rate (ct s <sup>-1</sup> )	Var <sup>b</sup>
		h	m	s								
10-018	CoKuLk332/G1 AB	4 42 07.32	25 23 03.2	1.69	0.14	243	1448	30 835	0.0470	0		
10-020	V955 Tau AB	4 42 07.76	25 23 11.6	1.70	0.24	308	1360	23 616	0.0576	0		
10-034	CIDA 7	4 42 21.02	25 20 35.5	1.83	1.10	71	119	43 153	0.0028	2		
10-045	DP Tau	4 42 37.72	25 15 36.8	1.83	0.75	64	93	30 465	0.0031	2		
10-060	GO Tau	4 43 03.12	25 20 19.8	1.72	1.08	611	479	21 286	0.0225	0		
26-012	2M J04552333+30	4 55 23.12	30 27 38.2	1.98	3.15	17	54	28 154	0.0019	0		
26-034	2M J04554046+30	4 55 40.34	30 39 07.1	2.19	2.09	4	45	35 843	0.0013	0		
26-043	AB Aur	4 55 45.83	30 33 03.3	1.73	1.10	7917	3472	69 171	0.0502	0		
26-050	2MJ04554757/801	4 55 47.83	30 28 05.3	1.76	4.13	426	418	43 514	0.0096	2		
26-067	SU Aur	4 55 59.34	30 34 00.9	1.73	0.87	245 679	59 141	78 106	0.7572	1		
26-072	HBC 427	4 56 02.05	30 21 03.9	1.73	0.40	91 805	25 437	28 045	0.9070	1		
Additional sources from Chandra							Sig. <sup>a</sup>					
C1-0	KPNO-Tau 10	4 17 49.55	28 13 31.9	0.00	0.00	0.0	0	17 734	0.0000	0		
C1-1	IRAS 04158+2805	4 18 58.15	28 12 23.3	0.93	0.24	6.0	100	17 734	0.0056	0		
C2-1	Haro 6-5 B	4 22 00.71	26 57 32.2	0.07	0.33	15.4	32	29 674	0.0011	0		
C2-2	FS Tau AC	4 22 02.20	26 57 30.4	0.04	0.29	101.0	254	29 674	0.0086	0		
C3-1	FV Tau/c AB	4 26 54.34	26 06 51.3	0.21	0.99	6.2	13	29 717	0.0005	0		
C3-2	DG Tau B	4 27 02.58	26 05 30.8	0.22	1.12	4.3	9	29 717	0.0003	0		
C4-1	GV Tau AB	4 29 23.74	24 33 00.5	0.05	0.24	26.7	57	24 650	0.0023	0		
C5-2	HN Tau AB	4 33 39.34	17 51 51.3	1.40	1.11	3.6	22	4679	0.0047	0		
C5-1	L1551 55	4 32 43.70	18 02 54.4	1.17	1.95	6.7	74	4679	0.0160	0		
C5-4	HD 28867	4 33 33.06	18 01 00.1	0.01	1.18	39.2	1582	4679	0.3390	0		
C5-3	DM Tau	4 33 48.63	18 10 11.5	1.28	1.97	5.1	33	4679	0.0071	0		
C6-1	CFHT-BD Tau 4	4 39 47.50	26 01 40.8	0.14	0.27	16.8	30	19 317	0.0016	0		
C6-0	L1527 IRS	4 39 53.59	26 03 05.5	0.00	0.00	0.0	0	19 317	0.0000	0		
C6-0	CFHT-Tau 17	4 40 01.75	25 56 29.2	0.00	0.00	0.0	0	19 317	0.0000	0		
C6-2	IRAS 04370+2559	4 40 08.02	26 05 25.5	0.09	0.29	74.6	229	19 317	0.0119	0		

<sup>a</sup> Maximum likelihood for detection for *XMM-Newton* data, CIAO WAVDETECT "Significance" for *Chandra* data.

<sup>b</sup> Variability flag: 0 = no or only low-level variability; 1 = clear flaring, flare intervals removed in spectral fit; 2 = clear flaring observed but flare intervals not removed; 3 = slow decay of flare throughout observation, all data used.

Notes on individual objects:

- CoKu Tau 1 = XEST-23-000: *Chandra* observation C1 may contain a marginal off-axis detection ( $\approx 6$  counts).
- V892 Tau = XEST-23-047 = XEST-24-040: Companion at 4'' is visible as a faint source in *Chandra* observation C1.
- HP Tau/G2 and G3 = XEST-08-051 (separation: 10'') were treated as one source. The ratios of the counts and rates were derived from PSF fitting in the image.



**Table 5.** X-ray parameters of targets in XEST (2): Plasma parameters from the DEM fits.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_0$ ( $1\sigma$ range) (MK)	$\beta$ ( $1\sigma$ range)	EM $^a$ ( $10^{52}$ )	$L_{\text{X}}^b$ (range) ( $10^{30}$ erg s $^{-1}$ )	log $L_{\text{X}}/L_{\odot}$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^2$	d.o.f.
27-115	HBC 352	0.22 (0.19,0.25)	6.5 (5.0, 8.2)	-0.60 (-0.79,-0.47)	25.21	2.657 (2.43, 2.87)	-3.03	12.00	0.85	129
27-000	HBC 353	0.17	10.0	-1.00	...	<0.176	<-4.04	...	...	...
06-005	HBC 358 AB	0.01 (0.00,0.04)	4.7 (3.0, 7.5)	-0.60 (-1.24,-0.34)	3.79	0.383 (0.37, 0.44)	-3.45	9.42	0.74	21
06-007	HBC 359	0.01 (0.00,0.02)	7.2 (5.8, 8.9)	-1.25 (-1.84,-0.91)	6.65	0.663 (0.64, 0.69)	-3.18	9.20	1.05	57
06-059	L1489 IRS	6.63 (6.18,7.09)	50.9 (30.9,60.5)	-3.00 (-3.00, 0.02)	30.39	4.471 (4.07, 4.91)	-3.63	40.42	0.94	99
20-001	LkCa 1	0.07 (0.04,0.15)	4.9 (3.7, 5.7)	-3.00 (-3.00,-1.94)	2.66	0.232 (0.20, 0.37)	-3.80	4.35	0.57	18
20-005	Anon 1	0.33 (0.30,0.35)	7.6 (6.4, 9.1)	-1.01 (-1.26,-0.88)	40.27	4.139 (3.84, 4.42)	-3.38	10.72	0.92	228
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	5.68 (2.99,8.19)	4.5 (2.0,31.6)	-1.00	4.39	0.417 (0.07, 0.57)	-3.57	6.93	0.68	8
20-000	2M J04141188+28	0.18	7.9	-3.00	...	<0.028	<-3.31	...	...	...
20-042	V773 Tau ABC	0.17 (0.17,0.17)	8.7 (8.4, 9.1)	-0.87 (-0.93,-0.82)	89.77	9.488 (9.39, 9.54)	-3.36	12.96	1.06	526
20-043	FM Tau	0.15 (0.13,0.17)	6.1 (3.9, 9.6)	-0.02 (-0.29, 0.11)	4.59	0.532 (0.51, 0.56)	-3.52	18.47	1.12	78
20-046	CW Tau	6.49 (4.02,8.12)	4.5 (2.0,12.4)	-3.00 (-3.00,-2.21)	33.14	2.844 (0.28, 4.00)	-3.17	4.07	1.76	2
20-047	CIDA 1	0.22 (0.03,0.47)	5.0 (2.5,10.7)	-1.00	0.26	0.025 (0.01, 0.07)	...	7.57	0.68	3
20-056	MHO 2/1	0.92 (0.83,1.15)	17.7 (10.4,28.9)	-1.20 (-3.00,-0.35)	14.44	1.636 (1.48, 2.39)	-3.12	20.53	0.98	85
20-058	MHO 3	0.61 (0.48,1.02)	10.0 (2.0,15.4)	-1.40 (-3.00,-0.74)	4.46	0.459 (0.36, 2.18)	-3.88	11.76	0.86	20
20-069	FO Tau AB	0.25 (0.17,0.77)	12.5 (2.0,18.7)	-2.01 (-3.00,-0.37)	0.64	0.065 (0.05, 0.52)	-4.66	12.33	1.29	4
20-073	CIDA 2	0.13 (0.06,0.27)	6.0 (3.3, 7.8)	-2.87 (-3.00,-1.49)	1.96	0.178 (0.14, 0.36)	-3.84	5.34	0.80	18
23-002	CY Tau	0.00 (0.00,0.17)	6.7 (2.2,10.3)	-1.05 (-3.00,-0.36)	1.32	0.133 (0.13, 0.29)	-4.16	9.42	1.71	10
24-002	CY Tau	0.02	9.5 (6.6,13.7)	-0.79 (-2.00,-0.29)	1.79	0.194 (0.19, 0.30)	-4.00	14.46	0.80	12
23-004	LkCa 5	0.02 (0.01,0.03)	7.9 (6.4, 9.1)	-1.33 (-1.86,-0.91)	4.32	0.432 (0.41, 0.45)	-3.52	9.67	0.59	85
24-004	LkCa 5	0.04 (0.02,0.06)	5.6 (3.7, 8.3)	-0.50 (-0.96,-0.28)	6.46	0.681 (0.64, 0.73)	-3.32	11.62	0.99	72
23-008	CIDA 3	1.38 (0.78,2.09)	5.7 (2.0,31.6)	-0.67 (-1.67,-0.45)	2.67	0.277 (0.12, 1.62)	-3.29	10.45	1.15	11
24-008	CIDA 3	0.85 (0.67,1.09)	31.6 (22.3,31.6)	-1.00	1.95	0.254 (0.22, 0.31)	-3.33	33.60	0.87	7
23-015	V410 X3	0.13 (0.07,0.25)	4.9 (3.1, 8.7)	-1.22 (-2.75,-0.71)	0.63	0.059 (0.05, 0.10)	-3.74	6.67	0.76	13
24-015	V410 X3	0.24 (0.00,0.52)	3.5 (2.0, 9.9)	-1.98 (-3.00,-1.04)	1.10	0.092 (0.03, 0.47)	-3.55	3.81	0.79	5
23-018	V410 A13	0.42 (0.00,0.78)	4.0 (2.0,20.6)	-1.00	0.10	0.010 (0.00, 0.03)	-4.18	6.21	1.11	2
24-000	V410 A13	0.50	10.0	-1.00	...	<0.011	<-4.14	...	...	...
23-000	V410 A24	4.37	10.0	-1.00	...	<0.042	<-5.41	...	...	...
24-000	V410 A24	4.37	10.0	-1.00	...	<0.078	<-5.14	...	...	...
23-029	V410 A25	0.59 (0.53,0.65)	25.9 (23.1,29.1)	-3.00 (-3.00,-2.17)	5.98	0.676 (0.63, 0.72)	-3.96	21.75	1.27	90
24-027	V410 A25	0.57 (0.49,0.66)	33.6 (26.4,40.0)	-3.00 (-3.00,-1.10)	4.79	0.587 (0.55, 0.64)	-4.02	27.79	1.11	47
23-032	V410 Tau ABC	0.01 (0.01,0.02)	9.0 (7.7,10.1)	-1.14 (-1.43,-0.85)	36.33	3.762 (3.68, 3.84)	-3.35	11.74	0.95	115
24-028	V410 Tau ABC	0.02 (0.02,0.03)	10.4 (9.4,10.8)	-1.17 (-1.28,-0.96)	44.47	4.663 (4.58, 4.76)	-3.26	13.20	1.10	414
23-033	DD Tau AB	0.23 (0.16,0.37)	3.8 (2.0,31.6)	0.93 (0.19, 1.00)	0.69	0.093 (0.08, 0.11)	-4.15	37.84	1.37	6
24-029	DD Tau AB	0.29 (0.21,0.45)	31.6 (2.0,31.6)	0.13 (-3.00, 1.00)	1.00	0.139 (0.12, 0.16)	-3.97	41.71	1.37	5
23-035	CZ Tau AB	0.34 (0.24,0.43)	5.0 (3.9, 6.5)	-3.00 (-3.00,-2.57)	4.86	0.423 (0.28, 0.64)	-3.39	4.41	1.09	28
24-030	CZ Tau AB	0.33 (0.24,0.37)	5.0 (4.6, 6.4)	-3.00 (-3.00,-2.73)	6.22	0.547 (0.37, 0.64)	-3.28	4.45	1.86	22
23-036	IRAS 04154+2823	6.80 (4.36,13.1)	31.6 (2.0,31.6)	-1.00	1.44	0.188 (0.13, 0.88)	-3.43	33.60	0.89	8
24-031	IRAS 04154+2823	5.43 (1.89,10.6)	9.1 (2.0,31.6)	-1.00	1.56	0.164 (0.03, 0.72)	-3.48	12.58	1.16	1
23-037	V410 X2	1.47 (1.32,1.59)	2.9 (2.0, 5.2)	-1.65 (-1.82,-1.48)	76.83	6.192 (3.06,11.49)	-3.27	3.48	0.81	84
24-032	V410 X2	1.46 (0.95,1.60)	2.0 (2.0,14.1)	-1.47 (-1.75,-1.15)	109.71	7.853 (1.18,11.25)	-3.17	2.79	0.58	24
23-045	V410 X4	0.94 (0.79,1.16)	15.0 (9.9,19.1)	-3.00 (-3.00,-1.34)	2.10	0.211 (0.17, 0.30)	-4.38	12.73	0.74	19
24-038	V410 X4	0.74 (0.32,1.43)	17.6 (2.0,31.6)	-2.84 (-3.00, 0.42)	2.06	0.213 (0.13, 1.69)	-4.37	15.15	0.12	4
23-047	V892 Tau	0.94 (0.89,0.98)	12.0 (10.5,14.5)	-1.18 (-1.50,-1.04)	86.14	9.207 (8.51, 9.90)	-4.51	14.86	1.17	377
24-040	V892 Tau	1.04 (0.90,1.31)	9.0 (3.3,13.7)	-1.39 (-1.97,-1.04)	78.05	7.944 (6.02,18.12)	-4.57	10.74	0.90	86
23-048	LR 1	2.77 (2.11,3.66)	2.0 (2.0,31.6)	-1.00	5.33	0.417 (0.29, 0.55)	-3.64	3.63	0.59	5
24-000	LR 1	4.16	10.0	-1.00	...	<0.169	<-4.03	...	...	...
23-050	V410 X7	0.80 (0.70,0.92)	12.6 (8.5,17.9)	-1.34 (-2.48,-0.88)	8.60	0.913 (0.78, 1.16)	-3.32	14.68	0.98	44
24-042	V410 X7	0.72 (0.67,0.78)	29.4 (19.4,32.5)	-2.99 (-3.00,-1.06)	25.28	2.975 (2.82, 3.14)	-2.81	24.57	0.93	82
23-000	V410 A20	4.27	10.0	-1.00	...	<0.075	<-4.42	...	...	...
24-000	V410 A20	4.27	10.0	-1.00	...	<0.094	<-4.32	...	...	...
23-056	Hubble 4	0.25 (0.24,0.25)	9.1 (8.6, 9.6)	-1.08 (-1.17,-0.99)	51.25	5.342 (5.24, 5.38)	-3.29	12.14	1.02	383
24-047	Hubble 4	0.24 (0.23,0.25)	9.2 (8.6,10.0)	-1.16 (-1.29,-1.03)	44.83	4.668 (4.52, 4.76)	-3.35	11.96	0.98	280
23-000	KPNO-Tau 2	0.07	7.9	-3.00	...	<0.012	<-3.38	...	...	...
24-000	KPNO-Tau 2	0.07	7.9	-3.00	...	<0.012	<-3.38	...	...	...
23-000	CoKu Tau 1	...	...	...	...	...	...	...	...	...
24-000	CoKu Tau 1	...	...	...	...	...	...	...	...	...
23-061	V410 X6	0.36 (0.26,0.43)	3.2 (2.1, 4.6)	-1.00	1.39	0.124 (0.08, 0.17)	-3.79	5.15	1.26	13

Table 5. continued.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_0$ ( $1\sigma$ range) (MK)	$\beta$ ( $1\sigma$ range)	$EM_r^a$ ( $10^{52}$ )	$L_X^b$ (range) ( $10^{30}$ erg s $^{-1}$ )	$\log L_X/L_e$	$T_{\text{av}}$ (MK)	$\chi^2_{\text{red}}$	d.o.f.
24-054	V410 X6	0.45 (0.32,0.63)	5.0 (2.4, 9.1)	-1.19 (-2.55,-0.71)	2.93	0.278 (0.18, 0.57)	-3.44	6.86	1.21	9
23-063	V410 X5	0.45 (0.37,0.56)	7.0 (4.6,11.6)	-1.16 (-1.90,-0.77)	3.87	0.387 (0.31, 0.52)	-2.92	9.30	1.17	32
24-055	V410 X5	0.61 (0.52,0.72)	4.0 (2.2, 7.0)	-0.50 (-0.76,-0.30)	11.07	1.115 (0.87, 1.59)	-2.46	9.16	0.94	41
23-067	FQ Tau AB	0.50 (0.22,0.93)	5.0 (2.0,10.1)	-3.00 (-3.00,-1.51)	1.36	0.120 (0.05, 0.83)	-3.83	4.46	1.79	6
24-058	FQ Tau AB	0.35	5.9 (2.0,10.0)	-3.00	5.59	0.049 (0.04, 0.16)	-4.22	5.17	0.87	1
28-100	BP Tau	0.06 (0.06,0.07)	7.1 (6.7, 7.6)	-0.67 (-0.73,-0.60)	12.84	1.365 (1.35, 1.40)	-3.43	12.34	1.10	430
23-074	V819 Tau AB	0.21 (0.20,0.23)	4.9 (4.3, 5.4)	-0.99 (-1.10,-0.90)	25.41	2.445 (2.33, 2.61)	-3.16	7.45	1.04	199
24-061	V819 Tau AB	0.22 (0.18,0.26)	4.5 (3.5, 5.7)	-1.00 (-1.13,-0.87)	23.24	2.205 (1.91, 2.66)	-3.20	6.87	0.90	110
16-000	IRAS 04166+2706	...	...	...	...	...	...	...	...	...
16-000	IRAS 04169+2702	...	...	...	...	...	...	...	...	...
11-000	CFHT-Tau 19	1.31	7.9	-3.00	...	<0.047	<-3.77	...	...	...
11-000	IRAS 04181+2655	...	...	...	...	...	...	...	...	...
11-000	IRAS 04181+2654AB	...	...	...	...	...	...	...	...	...
11-023	2M J04213459	0.31	5.4 (3.8,10.0)	-1.37 (-3.00,-0.91)	0.49	0.043 (0.04, 0.05)	-3.76	6.79	0.90	4
01-028	IRAS 04187+1927	0.62 (0.55,0.70)	6.3 (4.7,12.1)	-1.02 (-1.28,-0.84)	9.09	0.909 (0.75, 1.15)	...	9.15	0.94	79
11-037	CFHT-Tau 10	0.65	10.0	-3.00	0.16	0.015	-3.73	8.59	...	...
11-000	2M J04215450+2652	0.54	7.9	-3.00	...	<0.008	<-3.17	...	...	...
21-038	RY Tau	0.77 (0.70,0.86)	2.0 (2.0, 4.2)	0.20 (-0.03, 0.41)	49.07	5.520 (4.82, 6.38)	-3.72	15.70	1.18	158
21-039	HD 283572	0.08 (0.07,0.08)	10.4 (10.0,10.7)	-0.94 (-1.02,-0.87)	114.33	13.003 (2.68,13.05)	-3.28	14.43	1.12	522
01-045	T Tau N(+Sab)	0.27 (0.27,0.28)	11.3 (10.8,11.8)	-0.65 (-0.72,-0.58)	71.84	8.048 (7.99, 8.15)	-3.63	17.62	1.21	627
11-054	Haro 6-5 B	19.3 (12.9,23.8)	6.6 (2.0,16.7)	-3.00 (-3.00,-2.30)	18.95	17.551 (1.49,22.38)	-1.01	5.74	1.26	7
11-057	FS Tau AC	1.42 (1.34,1.48)	39.8 (34.6,44.8)	-2.90 (-3.00,-1.53)	24.99	3.224 (3.09, 3.36)	-2.58	32.76	1.13	155
21-044	LkCa 21	0.08 (0.06,0.10)	8.5 (6.0, 9.7)	-2.21 (-3.00,-1.31)	6.65	0.646 (0.60, 0.71)	-3.57	8.18	1.07	82
01-054	RX J0422.1+1934	0.28 (0.27,0.28)	13.6 (12.8,14.5)	-1.64 (-1.93,-1.22)	29.59	3.110 (3.05, 3.12)	...	14.46	1.05	411
01-062	2M J04221332+1934	0.37 (0.19,0.60)	4.4 (2.5, 7.7)	-3.00 (-3.00,-1.63)	0.67	0.057 (0.03, 0.22)	-3.07	3.92	1.06	6
11-079	CFHT-Tau 21	1.19	31.6 (23.4,31.6)	-1.00	1.16	0.151 (0.15, 0.15)	-3.99	33.60	0.73	8
02-013	FV Tau AB	1.20 (0.87,1.50)	2.0 (2.0,31.6)	0.32 (-0.41, 1.00)	4.51	0.532 (0.37, 0.72)	-3.94	18.76	1.06	13
02-000	FV Tau/c AB	0.59	10.0	-1.00	...	<0.047	<-4.24	...	...	...
02-016	KPNO-Tau 13	0.32 (0.23,0.45)	5.0 (2.2, 8.1)	-0.97 (-1.93,-0.53)	1.42	0.138 (0.10, 0.24)	-3.62	7.71	0.79	11
02-000	DG Tau B	...	...	...	...	...	...	...	...	...
02-022	DG Tau A	...	...	...	...	...	...	...	...	...
02-000	KPNO-Tau 4	0.44	7.9	-3.00	...	<0.013	<-3.09	...	...	...
02-000	IRAS 04248+2612AB	...	...	...	...	...	...	...	...	...
15-020	JH 507	0.26 (0.19,0.36)	3.8 (2.7, 5.2)	-1.56 (-1.97,-1.25)	5.20	0.455 (0.34, 0.75)	-3.55	4.62	0.47	31
13-004	GV Tau AB	...	...	...	...	...	...	...	...	...
13-000	IRAS 04264+2433	...	...	...	...	...	...	...	...	...
15-040	DH Tau AB	0.20 (0.19,0.21)	11.5 (11.0,12.1)	-1.38 (-1.49,-1.26)	80.60	8.458 (8.23, 8.64)	-2.41	13.42	1.06	381
15-042	DI Tau AB	0.14 (0.13,0.16)	9.5 (8.4,11.3)	-1.43 (-2.01,-1.18)	15.39	1.568 (1.51, 1.65)	-3.39	11.07	0.92	111
15-044	KPNO-Tau 5	0.01	7.9 (4.6,12.6)	-3.00 (-3.00,-0.66)	0.10	0.010 (0.01, 0.01)	-3.95	6.88	1.11	2
14-006	IQ Tau A	0.49 (0.36,0.80)	10.1 (2.0,19.6)	-0.95 (-3.00,-0.06)	3.89	0.416 (0.33, 1.17)	-3.91	14.06	1.57	9
13-000	CFHT-Tau 20	0.65	7.9	-3.00	...	<0.017	<-4.49	...	...	...
14-000	KPNO-Tau 6	0.16	7.9	-3.00	...	<0.008	<-3.17	...	...	...
13-035	FX Tau AB	0.26 (0.18,0.55)	7.9 (2.0,10.1)	-3.00 (-3.00,-0.94)	5.26	0.502 (0.39, 2.36)	-3.89	6.86	0.98	5
14-057	DK Tau AB	0.25 (0.23,0.27)	9.2 (6.7,13.3)	-0.81 (-1.57,-0.52)	8.54	0.916 (0.87, 0.96)	-3.74	13.90	0.86	106
14-000	KPNO-Tau 7	0.01	7.9	-3.00	...	<0.006	<-3.33	...	...	...
22-013	MHO 9	0.05 (0.02,0.08)	6.8 (5.6, 7.9)	-3.00 (-3.00,-1.92)	0.86	0.080 (0.07, 0.09)	-4.02	5.93	1.05	20
22-021	MHO 4	0.16 (0.11,0.22)	6.1 (3.8,11.2)	-1.15 (-3.00,-0.60)	1.25	0.123 (0.10, 0.16)	-3.18	8.26	0.73	16
22-040	L1551 IRS5	0.17 (0.11,0.29)	31.6 (21.4,31.6)	-1.00	0.17	0.021 (0.02, 0.02)	-5.68	33.60	1.40	6
22-042	LkHa 358	3.01 (2.26,4.05)	3.0 (2.0,30.9)	-1.00	4.58	0.404 (0.29, 0.53)	-3.75	4.99	0.40	11
22-000	HH 30	...	...	...	...	...	...	...	...	...
22-043	HL Tau	2.81 (2.59,3.04)	20.1 (13.6,29.9)	-1.40 (-3.00,-0.80)	26.84	3.838 (3.22, 4.73)	-3.19	21.80	1.14	87
22-047	XZ Tau AB	0.28 (0.23,0.33)	8.5 (6.0,11.5)	-0.90 (-1.45,-0.62)	9.14	0.962 (0.86, 1.12)	-3.12	12.49	1.07	38
22-056	L1551 NE	7.66 (3.96,13.8)	31.6	-1.00	0.46	0.059 (0.03, 0.09)	...	33.60	0.10	1
03-005	HK Tau AB	0.92 (0.49,1.89)	63.1 (14.1,63.1)	-1.00	0.53	0.079 (0.06, 0.12)	-4.44	51.83	3.19	1
22-070	V710 Tau BA	0.28 (0.26,0.30)	7.9 (6.7, 8.8)	-1.44 (-1.71,-1.22)	13.81	1.378 (1.32, 1.49)	-3.49	9.33	0.94	158
19-009	JH 665	0.14 (0.05,0.37)	5.0 (2.0,11.8)	-0.94 (-3.00,-0.26)	0.81	0.079 (0.06, 0.23)	-4.10	7.78	0.58	3
22-089	L1551 51	0.11 (0.10,0.12)	7.7 (7.0, 8.4)	-1.63 (-1.88,-1.42)	18.67	1.841 (1.78, 1.91)	-2.99	8.61	1.13	204
22-097	V827 Tau	0.06 (0.05,0.06)	9.4 (8.7,10.7)	-0.74 (-0.96,-0.65)	37.02	4.010 (3.88, 4.01)	-3.02	14.64	1.11	273
03-016	Haro 6-13	1.26 (0.44,1.41)	2.5 (2.0,31.6)	-1.00	9.49	0.799 (0.14, 0.91)	-4.01	4.29	1.04	6

Table 5. continued.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_0$ ( $1\sigma$ range) (MK)	$\beta$ ( $1\sigma$ range)	$EM_{\text{H}}^0$ ( $10^{52}$ )	$L_{\text{X}}^b$ (range) ( $10^{30}$ erg s $^{-1}$ )	log $L_{\text{X}}/L_{\odot}$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^2$	d.o.f.
22-100	V826 Tau	0.07 (0.06,0.07)	9.1 (8.6, 9.4)	-1.26 (-1.36,-1.16)	44.06	4.523 (4.42, 4.57)	-2.90	11.27	1.12	359
22-101	MHO 5	0.09 (0.06,0.14)	5.8 (4.2, 8.6)	-1.62 (-3.00,-1.09)	1.18	0.111 (0.10, 0.14)	-3.58	6.63	0.95	31
03-017	CFHT-Tau 7	0.01	8.1 (5.9,12.0)	-3.00 (-3.00,-0.90)	0.21	0.020 (0.02, 0.02)	-4.07	7.03	1.82	3
03-019	V928 Tau AB	0.38 (0.31,0.44)	5.0 (3.9, 7.0)	-1.40 (-1.78,-1.23)	11.21	1.046 (0.81, 1.33)	-3.71	6.27	0.93	46
03-022	FY Tau	0.31 (0.28,0.36)	13.7 (8.7,19.6)	-0.99 (-2.31,-0.40)	7.30	0.807 (0.76, 0.90)	-3.68	17.73	0.54	34
03-023	FZ Tau	0.53 (0.33,0.82)	4.8 (2.0,12.0)	-1.55 (-2.44,-1.05)	7.05	0.644 (0.35, 2.91)	-3.77	5.65	0.70	16
17-002	IRAS 04295+2251	3.75 (2.74,5.26)	16.8 (5.2,27.0)	-3.00 (-3.00,-1.21)	14.56	1.489 (0.82, 4.80)	-3.49	14.28	0.46	7
19-049	UZ Tau E+W(AB)	0.44 (0.29,0.53)	2.7 (2.0, 7.4)	-0.75 (-0.95,-0.58)	9.98	0.890 (0.51, 1.35)	-3.59	5.44	0.74	61
17-009	JH 112	0.72 (0.57,0.86)	3.9 (2.0, 7.8)	-0.90 (-1.20,-0.61)	8.71	0.820 (0.52, 1.48)	-3.54	6.58	0.64	22
03-031	CFHT-Tau 5	0.84	8.1 (3.5,11.5)	-3.00 (-3.00,-1.27)	1.91	0.181 (0.14, 0.26)	-3.20	7.03	1.26	4
04-003	CFHT-Tau 5	0.84 (0.51,1.48)	11.9 (2.0,17.8)	-3.00 (-3.00,-1.07)	2.51	0.247 (0.14, 2.28)	-3.07	10.14	0.63	7
03-035	MHO 8	0.21 (0.13,0.34)	4.0 (2.1, 6.1)	-1.00	0.70	0.065 (0.05, 0.12)	-4.00	6.21	2.00	3
04-009	MHO 8	0.68 (0.46,0.82)	2.5 (2.0, 4.4)	-3.00 (-3.00,-2.51)	6.18	0.444 (0.13, 0.93)	-3.17	2.43	0.97	5
04-010	GH Tau AB	0.09	12.6 (9.2,18.8)	-1.00	1.00	0.109 (0.10, 0.12)	-4.46	16.59	0.71	7
04-012	V807 Tau SNab	0.10 (0.08,0.12)	5.4 (4.4, 6.4)	-1.18 (-1.40,-0.99)	10.92	1.049 (0.97, 1.15)	-3.89	7.35	1.06	84
18-004	KPNO-Tau 14	0.85 (0.69,1.31)	12.6 (2.0,21.3)	-1.00 (-2.30,-0.40)	8.44	0.923 (0.76, 3.08)	-2.66	16.50	1.05	27
17-000	CFHT-Tau 12	0.61	7.9	-3.00	...	<0.007	<-4.30	...	...	...
04-016	V830 Tau	0.05 (0.04,0.05)	9.2 (8.4, 9.6)	-0.53 (-0.64,-0.42)	46.62	5.181 (5.07, 5.23)	-2.76	16.23	1.22	309
18-000	IRAS S04301+261	1.13	10.0	-1.00	...	<0.023	<-3.62	...	...	...
17-000	IRAS 04302+2247	...	...	...	...	...	...	...	...	...
17-027	IRAS 04303+2240	1.66 (1.44,1.79)	2.0 (2.0, 6.8)	0.07 (-0.10, 0.48)	41.37	5.006 (3.50, 5.97)	-3.23	12.95	0.95	114
04-034	GI Tau	0.38 (0.33,0.45)	4.0 (2.0, 6.8)	-0.19 (-0.33,-0.05)	7.76	0.833 (0.73, 1.06)	-3.66	12.52	0.76	49
04-035	GK Tau AB	0.45 (0.41,0.71)	7.8 (2.0, 9.5)	-1.03 (-1.31,-0.72)	14.24	1.471 (1.33, 4.41)	-3.56	10.92	1.02	80
18-019	IS Tau AB	0.40 (0.35,0.50)	14.2 (8.4,16.6)	-2.49 (-3.00,-1.17)	6.48	0.658 (0.60, 0.85)	-3.59	12.83	1.07	56
17-058	CI Tau	0.55 (0.38,1.18)	17.7 (2.0,31.6)	-1.00	1.69	0.195 (0.16, 0.89)	-4.23	21.78	0.68	9
18-030	IT Tau AB	0.69 (0.66,0.72)	22.6 (17.1,34.5)	-0.86 (-3.00,-0.38)	53.13	6.492 (6.28, 6.73)	-3.15	27.22	0.92	427
17-066	JH 108	0.20 (0.17,0.24)	9.4 (6.5,12.5)	-0.85 (-1.49,-0.44)	11.48	1.233 (1.14, 1.36)	-2.97	13.93	1.07	38
17-068	CFHT-BD Tau 1	0.56	17.0 (12.6,25.7)	-1.00	1.48	0.169 (0.17, 0.17)	-2.59	21.04	1.10	4
25-026	AA Tau	1.09 (0.96,1.20)	25.7 (20.6,31.0)	-3.00 (-3.00,-1.79)	11.00	1.241 (1.11, 1.36)	-3.39	21.55	0.91	42
09-010	HO Tau AB	0.20	5.3 (3.7, 8.0)	-1.00	0.48	0.047 (0.05, 0.05)	-4.14	8.01	1.06	5
08-019	FF Tau AB	0.31 (0.26,0.41)	8.0 (4.7,10.6)	-1.50 (-2.47,-1.00)	7.99	0.796 (0.69, 1.12)	-3.52	9.21	1.07	59
12-040	DN Tau	0.07 (0.07,0.08)	10.5 (9.5,11.5)	-1.25 (-1.51,-1.02)	11.03	1.155 (1.14, 1.17)	-3.52	12.93	1.12	227
12-000	IRAS 04325+2402AB	...	...	...	...	...	...	...	...	...
12-059	CoKu Tau 3 AB	0.38 (0.37,0.39)	14.2 (13.3,15.0)	-2.07 (-2.36,-1.80)	56.63	5.851 (5.71, 5.94)	-2.81	13.80	1.06	388
09-022	KPNO-Tau 8	0.11 (0.08,0.16)	3.4 (2.0, 5.8)	-0.11 (-0.23, 0.01)	4.66	0.504 (0.45, 0.60)	-2.21	12.62	0.75	59
08-037	HQ Tau AB	0.40 (0.37,0.45)	8.0 (6.4, 9.6)	-1.16 (-1.43,-0.99)	24.25	2.476 (2.29, 2.84)	...	10.57	1.07	135
09-026	HQ Tau AB	0.56 (0.52,0.61)	6.3 (4.7, 7.5)	-0.84 (-0.95,-0.71)	79.94	8.161 (7.43, 9.56)	...	10.05	0.84	166
08-043	KPNO-Tau 15	0.38 (0.36,0.43)	13.7 (7.1,15.0)	-0.33 (-0.45, 0.10)	21.46	2.624 (2.49, 2.78)	-2.31	23.46	0.91	248
09-031	KPNO-Tau 15	0.37 (0.26,0.52)	6.3 (3.6,10.1)	-1.07 (-2.05,-0.59)	4.85	0.483 (0.37, 0.77)	-3.05	8.92	1.22	11
08-000	KPNO-Tau 9	0.01	7.9	-3.00	...	<0.004	<-3.13	...	...	...
09-000	KPNO-Tau 9	0.01	7.9	-3.00	...	<0.007	<-2.89	...	...	...
08-048	HP Tau AB	0.53 (0.48,0.60)	8.8 (6.6,11.4)	-1.09 (-1.44,-0.89)	24.55	2.550 (2.28, 3.03)	-3.33	11.81	1.07	129
08-051a	HP Tau/G3 AB	0.41 (0.39,0.42)	9.2 (8.6, 9.9)	-1.37 (-1.46,-1.28)	12.85	1.293 (1.09, 1.39)	-3.33	11.03	1.04	382
08-051	HP Tau/G2	0.41 (0.39,0.42)	9.2 (8.6, 9.9)	-1.37 (-1.46,-1.28)	94.61	9.653 (9.15, 9.90)	-3.41	11.03	1.04	382
08-058	Haro 6-28 AB	0.38 (0.33,0.47)	10.0 (6.5,12.2)	-2.10 (-3.00,-1.16)	2.49	0.248 (0.22, 0.34)	-3.27	9.80	0.85	32
08-000	CFHT-BD Tau 2	0.01	7.9	-3.00	...	<0.004	<-4.02	...	...	...
08-080	CFHT-BD Tau 3	0.18	7.9 (4.0,17.8)	-3.00	0.13	0.012 (0.01, 0.01)	-3.36	6.88	0.16	1
05-005	CFHT-Tau 6	0.47 (0.33,0.64)	7.9	-3.00	0.75	0.071 (0.04, 0.09)	-3.11	6.88	1.53	3
05-000	IRAS 04361+2547	...	...	...	...	...	...	...	...	...
05-013	GN Tau AB	2.34 (1.48,2.80)	4.2 (2.0,17.7)	-1.00	8.35	0.785 (0.21, 1.00)	-3.55	6.52	1.33	6
05-017	IRAS 04365+2535	30.7 (20.0,51.8)	10.0	-1.00	18.70	1.986 (1.23, 3.52)	-3.63	13.65	0.28	2
05-024	IRAS 04369+2539	7.69 (5.40,8.71)	6.4 (2.0,31.6)	-1.00	46.50	4.671 (1.18, 5.55)	-4.23	9.32	1.12	7
07-011	JH 223	0.09	9.3 (7.6,10.6)	-3.00 (-3.00,-1.74)	0.66	0.064 (0.06, 0.07)	-4.03	8.00	1.22	10
07-022	Haro 6-32	0.08 (0.03,0.14)	12.0 (5.8,14.7)	-3.00 (-3.00,-0.76)	1.02	0.101 (0.09, 0.12)	-3.66	10.28	0.50	5
07-000	ITG 33 A	...	...	...	...	...	...	...	...	...
07-000	CFHT-Tau 8	0.32	7.9	-3.00	...	<0.008	<-4.06	...	...	...
07-000	IRAS 04381+2540	...	...	...	...	...	...	...	...	...
07-041	IRAS 04385+2550AB	0.80 (0.69,1.02)	31.6 (17.9,31.6)	-1.00	3.08	0.401 (0.37, 0.50)	-3.24	33.60	0.74	10
10-017	CoKuLk332/G2 AB	0.53 (0.49,0.59)	6.3 (4.8, 7.3)	-1.14 (-1.26,-1.02)	32.92	3.257 (2.91, 3.94)	-3.11	8.59	0.83	163

Table 5. continued.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_0$ ( $1\sigma$ range) (MK)	$\beta$ ( $1\sigma$ range)	$EM_{\text{r}}^a$ ( $10^{52}$ )	$L_{\text{X}}^b$ (range) ( $10^{30}$ erg s $^{-1}$ )	log $L_{\text{X}}/L_{\odot}$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^2$	d.o.f.
10-018	CoKuLk332/G1 AB	0.33 (0.24,0.58)	10.0 (2.0,15.5)	-0.70 (-1.73,-0.26)	4.50	0.493 (0.42, 1.11)	-4.12	15.78	0.64	21
10-020	V955 Tau AB	0.89 (0.70,1.30)	9.5 (2.0,21.1)	-0.83 (-1.90,-0.41)	15.07	1.622 (1.23, 4.55)	-3.38	14.25	0.91	36
10-034	CIDA 7	0.32	4.3 (2.4, 7.7)	-1.88 (-3.00,-0.90)	0.46	0.040 (0.04, 0.05)	-3.68	4.70	0.76	6
10-045	DP Tau	...	...	...	...	...	...	...	...	...
10-060	GO Tau	0.31 (0.25,0.44)	6.2 (2.0,10.2)	-0.23 (-0.71, 0.13)	2.24	0.249 (0.22, 0.36)	-3.76	15.54	1.56	13
26-012	2M J04552333+30	0.01	12.6 (8.3,18.6)	-3.00	0.13	0.013 (0.01, 0.01)	-3.65	10.75	1.55	2
26-034	2M J04554046+30	0.05	10.2 (6.5,17.0)	-3.00	0.11	0.011 (0.01, 0.01)	-3.87	8.79	0.52	1
26-043	AB Aur	0.05 (0.03,0.07)	4.8 (4.3, 5.5)	-1.55 (-1.78,-1.41)	3.80	0.349 (0.32, 0.38)	-5.73	5.70	1.25	125
26-050	2MJ04554757/801	0.10 (0.06,0.15)	10.8 (8.3,12.3)	-3.00 (-3.00,-1.76)	0.71	0.070 (0.06, 0.08)	-3.78	9.29	1.01	21
26-067	SU Aur	0.47 (0.43,0.48)	6.4 (6.2, 7.6)	-1.11 (-1.21,-1.06)	95.36	9.464 (8.42, 9.70)	-3.61	8.89	1.56	397
26-072	HBC 427	0.04 (0.03,0.05)	9.3 (7.8,10.8)	-0.81 (-1.08,-0.57)	33.01	3.558 (3.39, 3.57)	-3.08	14.11	0.90	107
Additional sources from Chandra										
C1-0	KPNO-Tau 10	...	...	...	...	...	...	...	...	...
C1-1	IRAS 04158+2805	4.36 (2.58,6.12)	2.3 (2.0,31.6)	0.04 (-3.00, 1.00)	8.10	0.882	-2.34	13.10	0.88	6
C2-1	Haro 6-5 B	...	...	...	...	...	...	...	...	...
C2-2	FS Tau AC	1.10 (0.95,1.38)	3.0 (2.0,31.6)	0.99 (0.26, 1.00)	2.84	0.391	-3.50	39.18	1.13	11
C3-1	FV Tau/c AB	...	...	...	...	...	...	...	...	...
C3-2	DG Tau B	...	...	...	...	...	...	...	...	...
C4-1	GV Tau AB	...	...	...	...	...	...	...	...	...
C5-2	HN Tau AB	...	...	...	...	...	...	...	...	...
C5-1	L1551 55	...	...	...	...	...	...	...	...	...
C5-4	HD 28867	...	...	...	...	...	...	...	...	...
C5-3	DM Tau	...	...	...	...	...	...	...	...	...
C6-1	CFHT-BD Tau 4	...	...	...	...	...	...	...	...	...
C6-0	L1527 IRS	...	...	...	...	...	...	...	...	...
C6-0	CFHT-Tau 17	...	...	...	...	...	...	...	...	...
C6-2	IRAS 04370+2559	1.31 (1.19,1.51)	31.6 (23.4,31.6)	-1.00	5.94	0.773	-3.02	33.60	1.03	12

<sup>a</sup>  $EM_{\text{r}}$  is sum of EM over all DEM bins from  $\log T = 6.0$  to  $\log T = 7.9$ ; given in units of  $10^{52}$  cm $^{-3}$

<sup>b</sup>  $L_{\text{X}}$  for [0.3,10] keV, in units of  $10^{30}$  erg s $^{-1}$

Notes on individual objects:

- For several detected, faint *Chandra* sources, only 1- $T$  fits were derived; see Table 6.
- Fit performed with MOS spectra for: HBC 352 (XEST-27-115), CY Tau (23-002), CY Tau (24-002), FQ Tau (23-067), 2M-J04213459 (11-023), FX Tau (13-035), V827 Tau (22-097), FZ Tau (03-023), CFHT-Tau 5 (03-031), GH Tau (04-010), V955 Tau (10-020), 2M J04552333+30, 2M J04554046+30, 2M J04554757/801, AB Aur, SU Aur, 2M J04552333+30 (26-012), 2M J04554046+30 (26-034), 2M J04554757/801 (26-050), AB Aur (26-043), SU Aur (26-067).
- HP Tau/G2 and G3 = XEST-08-051 (separation: 10'') were fitted as one source. The ratio of the normalizations was derived from PSF fitting in the image.
- CW Tau = XEST-20-046: fit is of low quality. CW Tau may be a two-absorber X-ray source (Güdel et al. 2007b).
- DG Tau A (XEST-02-022), GV Tau (XEST-13-004), and DP Tau (XEST-10-045) require 2- $T$  fits with two absorbers; see Table 6.
- L1489 IRS = XEST-06-049 required Fe = 0.46 (AG89) for acceptable fit.
- HD 283572 = XEST-21-039 required Fe = 0.27 (AG89) for acceptable fit.
- IRAS 04303+2240 = XEST-17-027 required Fe = 0.43 (AG89) for acceptable fit.
- HL Tau = XEST-22-043 required Fe = 0.77 (AG89) for acceptable fit.
- L1551 IRS 5 = XEST-22-040: the lightly absorbed source cannot originate from heavily absorbed protostar.
- CFHT-Tau 10 = XEST-11-037: Too few counts in spectrum for reliable spectral fit. Adopted model parameters are characteristic for low-mass stars, and  $N_{\text{H}}$  has been derived from  $A_{\text{V}}$ .  $L_{\text{X}}$  has been determined from count rate-to-flux conversion using spectral model in XSPEC.

**Table 6.** X-ray parameters of targets in XEST (3): Plasma parameters from the 1-*T* and 2-*T* fits.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_1^a$ (MK)	$T_2$ (MK)	$\text{EM}_1^b$ ( $10^{52}$ )	$\text{EM}_2^b$ ( $10^{52}$ )	$L_X^c$ ( $10^{30}$ )	$\log L_X/L_*$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^{2/d}$	d.o.f.
27-115	HBC 352	0.19 (0.17,0.21)	7.54	23.77	10.35	11.52	2.307	-3.09	13.81	0.87	128
27-000	HBC 353	...	...	...	...	...	...	...	...	...	...
06-005	HBC 358 AB	0.00 (0.00,0.03)	4.29	14.26	1.41	2.12	0.346	-3.49	8.82	0.73	20
06-007	HBC 359	0.00 (0.00,0.02)	5.33	15.19	3.06	3.53	0.628	-3.20	9.34	1.09	56
06-059	L1489 IRS	6.54 (6.08,6.88)	...	50.20	...	26.57	4.003	-3.67	50.20	1.02	100
20-001	LkCa 1	0.07 (0.00,0.21)	1.39	8.35	1.76	0.82	0.205	-3.85	2.46	1.27	15
20-005	Anon 1	0.28 (0.27,0.29)	8.58	24.12	19.05	14.11	3.473	-3.46	13.32	1.02	228
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	7.86 (3.40,28.0)	...	11.59	...	5.95	0.588	-3.42	11.59	0.61	8
20-000	2M J04141188+28	...	...	...	...	...	...	...	...	...	...
20-042	V773 Tau ABC	0.14 (0.14,0.15)	8.81	28.17	41.39	37.63	8.588	-3.40	15.32	1.28	525
20-043	FM Tau	0.14 (0.12,0.15)	7.19	29.45	1.41	3.06	0.494	-3.55	18.87	1.17	77
20-046	CW Tau	...	...	...	...	...	...	...	...	...	...
20-047	CIDA 1	0.34 (0.04,1.10)	6.38	...	0.24	...	0.031	...	6.38	0.02	2
20-056	MHO 2/1	1.01 (0.84,1.15)	8.00	28.87	6.35	10.35	1.853	-3.07	17.72	0.99	84
20-058	MHO 3	0.50 (0.40,0.69)	11.01	35.94	2.12	1.18	0.357	-3.99	16.80	0.88	19
20-069	FO Tau AB	0.18 (0.10,0.50)	...	13.80	...	0.47	0.052	-4.76	13.80	1.41	4
20-073	CIDA 2	0.03 (0.00,0.06)	7.77	...	1.18	...	0.115	-4.03	7.77	0.79	19
23-002	CY Tau	0.01 (0.00,0.16)	7.77	...	1.18	...	0.108	-4.25	7.77	1.85	9
24-002	CY Tau	0.12 (0.03,0.22)	8.81	...	2.12	...	0.202	-3.98	8.81	1.19	12
23-004	LkCa 5	0.03 (0.01,0.04)	4.75	17.16	2.35	2.35	0.440	-3.51	9.03	0.75	84
24-004	LkCa 5	0.01 (0.00,0.02)	8.35	28.29	3.06	2.35	0.588	-3.38	14.19	1.10	71
23-008	CIDA 3	0.71 (0.52,0.84)	...	38.38	...	0.94	0.115	-3.67	38.38	0.89	9
24-008	CIDA 3	0.43 (0.19,0.68)	...	126.38	...	1.18	0.226	-3.38	126.38	0.62	7
23-015	V410 X3	0.14 (0.09,0.24)	4.41	63.07	0.49	0.16	0.071	-3.66	8.58	0.65	12
24-015	V410 X3	0.00 (0.00,0.07)	7.88	...	0.31	...	0.031	-4.02	7.88	0.78	6
23-018	V410 A13	0.29 (0.02,0.62)	=10.00	...	0.05	...	0.006	-4.40	10.00	0.50	3
24-000	V410 A13	...	...	...	...	...	...	...	...	...	...
23-000	V410 A24	...	...	...	...	...	...	...	...	...	...
24-000	V410 A24	...	...	...	...	...	...	...	...	...	...
23-029	V410 A25	0.71 (0.60,0.80)	1.28	22.14	103.48	6.35	3.807	-3.21	1.51	1.11	89
24-027	V410 A25	0.65 (0.51,0.73)	1.62	27.83	11.05	4.94	1.174	-3.72	3.90	1.04	46
23-032	V410 Tau ABC	0.00 (0.00,0.01)	8.81	24.81	19.52	14.11	3.523	-3.38	13.60	1.03	115
24-028	V410 Tau ABC	0.02 (0.01,0.02)	8.70	22.03	19.28	23.05	4.402	-3.28	14.43	1.30	413
23-033	DD Tau AB	0.28 (0.22,0.54)	5.10	139.48	0.26	0.52	0.122	-4.03	46.29	1.41	5
24-029	DD Tau AB	0.27 (0.18,0.40)	...	38.72	...	0.94	0.132	-4.00	38.72	1.09	6
23-035	CZ Tau AB	0.17 (0.14,0.21)	7.42	...	2.12	...	0.205	-3.71	7.42	0.86	29
24-030	CZ Tau AB	0.19 (0.15,0.23)	7.07	...	3.06	...	0.292	-3.55	7.07	1.43	23
23-036	IRAS 04154+2823	4.10 (2.49,6.38)	...	67.83	...	1.18	0.176	-3.45	67.83	0.98	7
24-031	IRAS 04154+2823	7.67 (1.15,25.8)	...	14.14	...	2.89	0.282	-3.25	14.14	0.95	1
23-037	V410 X2	1.23 (1.14,1.31)	7.54	32.81	16.93	2.12	1.973	-3.77	8.88	0.81	83
24-032	V410 X2	1.43 (1.12,1.67)	3.48	15.19	46.33	9.88	4.953	-3.37	4.51	0.57	23
23-045	V410 X4	0.86 (0.71,1.05)	...	14.72	...	1.76	0.172	-4.46	14.72	0.59	16
24-038	V410 X4	0.56 (0.28,1.12)	...	19.13	...	1.58	0.160	-4.50	19.13	0.11	5
23-047	V892 Tau	0.86 (0.83,0.89)	10.55	28.87	32.69	39.04	7.923	-4.57	18.25	1.15	372
24-040	V892 Tau	1.25 (1.12,1.40)	4.87	22.49	108.65	35.98	13.779	-4.33	7.13	0.87	85
23-048	LR 1	0.63 (0.31,1.60)	...	277.10	...	0.24	0.045	-4.60	277.10	0.30	3
24-000	LR 1	...	...	...	...	...	...	...	...	...	...
23-050	V410 X7	0.77 (0.64,0.91)	9.97	18.09	3.06	3.29	0.626	-3.49	13.58	0.73	28
24-042	V410 X7	0.86 (0.73,1.01)	7.88	30.72	10.58	20.70	3.584	-2.73	19.39	0.90	81
23-000	V410 A20	...	...	...	...	...	...	...	...	...	...
24-000	V410 A20	...	...	...	...	...	...	...	...	...	...
23-056	Hubble 4	0.31 (0.30,0.33)	4.29	18.55	33.39	35.04	6.491	-3.20	9.08	1.10	382
24-047	Hubble 4	0.21 (0.20,0.22)	8.70	22.96	20.46	19.52	4.167	-3.40	13.97	1.04	279
23-000	KPNO-Tau 2	...	...	...	...	...	...	...	...	...	...
24-000	KPNO-Tau 2	...	...	...	...	...	...	...	...	...	...
23-000	CoKu Tau 1	...	...	...	...	...	...	...	...	...	...
24-000	CoKu Tau 1	...	...	...	...	...	...	...	...	...	...
23-061	V410 X6	0.73 (0.51,0.99)	3.25	...	6.35	...	0.541	-3.15	3.25	0.64	7

Table 6. continued.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_1^a$ (MK)	$T_2$ (MK)	$\text{EM}_1^b$ ( $10^{52}$ )	$\text{EM}_2^b$ ( $10^{52}$ )	$L_X^c$ ( $10^{30}$ )	$\log L_X/L_*$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^{2,d}$	d.o.f.
24-054	V410 X6	0.37 (0.28,0.64)	7.19	50.20	1.65	0.47	0.214	-3.56	11.07	1.21	8
23-063	V410 X5	0.40 (0.34,0.48)	6.96	18.20	1.65	1.65	0.322	-3.00	11.25	1.08	31
24-055	V410 X5	0.81 (0.75,0.85)	4.64	55.19	15.05	3.06	1.823	-2.24	7.05	0.88	40
23-067	FQ Tau AB	0.45 (0.26,0.80)	6.26	...	0.86	...	0.085	-3.98	6.26	1.48	7
24-058	FQ Tau AB	0.35	6.84	...	0.41	...	0.040	-4.31	6.84	0.88	1
28-100	BP Tau	0.09 (0.08,0.09)	4.75	22.26	9.64	7.53	1.482	-3.39	9.35	1.10	427
23-074	V819 Tau AB	0.21 (0.20,0.23)	4.64	17.51	15.52	9.64	2.356	-3.17	7.72	1.05	198
24-061	V819 Tau AB	0.20 (0.17,0.22)	4.17	14.72	11.29	9.64	1.945	-3.26	7.46	0.85	109
16-000	IRAS 04166+2706	...	...	...	...	...	...	...	...	...	...
16-000	IRAS 04169+2702	...	...	...	...	...	...	...	...	...	...
11-000	CFHT-Tau 19	...	...	...	...	...	...	...	...	...	...
11-000	IRAS 04181+2655	...	...	...	...	...	...	...	...	...	...
11-000	IRAS 04181+2654AB	...	...	...	...	...	...	...	...	...	...
11-023	2M J04213459	0.12 (0.01,1.10)	9.97	...	0.24	...	0.024	-4.02	9.97	0.63	3
01-028	IRAS 04187+1927	0.68 (0.64,0.80)	5.10	21.68	7.48	3.53	1.072	...	8.11	0.87	78
11-037	CFHT-Tau 10	0.65	=10.00	...	0.07	...	0.007	-4.06	10.00	0.36	2
11-000	2M J04215450+2652	...	...	...	...	...	...	...	...	...	...
21-038	RY Tau	0.76 (0.66,0.85)	5.91	37.57	15.29	27.28	5.242	-3.75	19.34	1.15	157
21-039	HD 283572	0.08 (0.08,0.08)	8.46	23.30	59.50	61.61	12.687	-3.30	14.16	1.16	522
01-045	T Tau N(+Sab)	0.34 (0.33,0.35)	4.52	23.65	33.86	57.85	9.395	-3.56	12.84	1.36	626
11-054	Haro 6-5 B	22.0 (14.4,33.6)	...	12.17	...	69.61	6.848	-1.42	12.17	0.90	8
11-057	FS Tau AC	1.70 (1.53,1.87)	3.83	36.17	36.92	24.69	6.434	-2.28	9.42	1.06	154
21-044	LkCa 21	0.10 (0.08,0.12)	4.29	13.22	4.00	3.53	0.698	-3.53	7.27	1.13	81
01-054	RX J0422.1+1934	0.27 (0.27,0.28)	8.46	21.91	10.82	18.11	3.010	...	15.35	1.04	410
01-062	2M J04221332+1934	0.40 (0.17,0.65)	4.64	9.97	0.63	0.05	0.059	-3.06	4.89	1.11	5
11-079	CFHT-Tau 21	1.48 (1.09,1.96)	...	30.26	...	1.41	0.169	-3.94	30.26	0.69	7
02-013	FV Tau AB	1.51 (0.97,2.21)	3.94	39.65	8.94	2.82	1.176	-3.59	6.86	1.04	12
02-000	FV Tau/c AB	...	...	...	...	...	...	...	...	...	...
02-016	KPNO-Tau 13	0.30 (0.22,0.42)	6.03	21.33	0.82	0.42	0.122	-3.68	9.26	0.71	10
02-000	DG Tau B	...	...	...	...	...	...	...	...	...	...
02-022	DG Tau A	0.11 (0.08,0.14)	3.71	22.96	0.80	1.69	0.252	-4.41	12.80	0.74	33
02-000	KPNO-Tau 4	...	...	...	...	...	...	...	...	...	...
02-000	IRAS 04248+2612AB	...	...	...	...	...	...	...	...	...	...
15-020	JH 507	0.27 (0.21,0.34)	3.94	20.75	4.23	0.94	0.456	-3.55	5.33	0.46	30
13-004	GV Tau AB	0.12 (0.01,0.79)	5.80	47.77	0.45	5.48	0.647	-4.03	40.75	1.08	5
13-000	IRAS 04264+2433	...	...	...	...	...	...	...	...	...	...
15-040	DH Tau AB	0.18 (0.18,0.19)	8.58	23.54	33.16	42.80	8.003	-2.43	15.15	0.94	380
15-042	DI Tau AB	0.12 (0.11,0.14)	7.65	18.20	6.58	7.76	1.437	-3.42	12.23	0.96	110
15-044	KPNO-Tau 5	0.01	6.61	...	0.09	...	0.009	-3.99	6.61	0.44	3
14-006	IQ Tau A	0.41 (0.33,0.52)	...	14.26	...	3.29	0.313	-4.03	14.26	1.57	10
13-000	CFHT-Tau 20	...	...	...	...	...	...	...	...	...	...
14-000	KPNO-Tau 6	...	...	...	...	...	...	...	...	...	...
13-035	FX Tau AB	0.20 (0.14,0.29)	8.46	...	0.39	...	0.393	-4.00	8.46	0.68	6
14-057	DK Tau AB	0.31 (0.28,0.34)	4.17	18.67	4.70	6.58	1.061	-3.67	10.00	0.83	105
14-000	KPNO-Tau 7	...	...	...	...	...	...	...	...	...	...
22-013	MHO 9	0.20 (0.12,0.28)	1.04	5.45	6.11	1.18	0.235	-3.56	1.36	0.77	19
22-021	MHO 4	0.13 (0.09,0.16)	5.10	13.45	0.52	0.54	0.101	-3.26	8.37	0.75	14
22-040	L1551 IRS5	0.18 (0.10,0.27)	...	27.83	...	0.16	0.019	-5.72	27.83	1.29	6
22-042	LkHa 358	1.78 (0.85,3.24)	...	36.64	...	0.49	0.066	-4.54	36.64	0.44	11
22-000	HH 30	...	...	...	...	...	...	...	...	...	...
22-043	HL Tau	2.79 (2.55,3.21)	22.38	159.42	21.64	2.12	3.285	-3.25	26.66	1.14	85
22-047	XZ Tau AB	0.24 (0.21,0.27)	8.70	26.43	4.00	3.76	0.849	-3.18	14.91	1.07	37
22-056	L1551 NE	5.51 (1.64,13.5)	...	43.13	...	0.33	0.045	...	43.13	0.03	2
03-005	HK Tau AB	0.89 (0.40,1.90)	...	79.54	...	0.47	0.080	-4.43	79.54	2.46	2
22-070	V710 Tau BA	0.23 (0.22,0.25)	7.77	18.32	6.82	4.94	1.166	-3.56	11.14	0.94	157
19-009	JH 665	0.13 (0.02,0.27)	6.84	...	0.71	...	0.059	-4.23	6.84	0.99	4
22-089	L1551 51	0.08 (0.07,0.09)	7.30	16.81	9.88	6.35	1.611	-3.05	10.12	1.14	203
22-097	V827 Tau	0.05 (0.04,0.05)	8.35	22.38	13.64	21.64	3.702	-3.06	15.29	1.16	272
03-016	Haro 6-13	0.54 (0.41,0.72)	...	27.25	...	1.29	0.151	-4.73	27.25	0.78	8

Table 6. continued.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_1^a$ (MK)	$T_2$ (MK)	$\text{EM}_1^b$ ( $10^{52}$ )	$\text{EM}_2^b$ ( $10^{52}$ )	$L_{\text{X}}^c$ ( $10^{30}$ )	$\log L_{\text{X}}/L_{\odot}$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^{2/d}$	d.o.f.
22-100	V826 Tau	0.06 (0.06,0.06)	7.54	17.97	19.52	22.34	4.210	-2.93	11.99	1.37	373
22-101	MHO 5	0.09 (0.06,0.13)	2.55	8.46	0.28	0.80	0.108	-3.59	6.19	1.06	30
03-017	CFHT-Tau 7	0.47 (0.24,1.46)	3.13	...	2.12	...	0.176	-3.13	3.13	0.80	3
03-019	V928 Tau AB	0.40 (0.33,0.46)	4.64	20.99	9.17	2.59	1.096	-3.69	6.47	0.92	45
03-022	FY Tau	0.30 (0.27,0.34)	9.28	27.94	2.35	4.47	0.769	-3.70	19.11	0.99	33
03-023	FZ Tau	0.28 (0.22,0.47)	...	11.48	...	2.70	0.268	-4.15	11.48	0.83	17
17-002	IRAS 04295+2251	3.43 (2.67,4.59)	...	20.17	...	9.88	1.025	-3.65	20.17	0.33	8
19-049	UZ Tau E+W(AB)	0.41 (0.33,0.44)	3.83	19.01	4.94	3.06	0.736	-3.67	7.07	0.77	60
17-009	JH 112	0.90 (0.68,1.02)	4.06	30.26	13.17	2.12	1.399	-3.31	5.36	0.55	21
03-031	CFHT-Tau 5	1.51 (0.88,2.29)	4.41	...	10.82	...	0.962	-2.48	4.41	0.88	4
04-003	CFHT-Tau 5	0.33 (0.21,0.55)	...	19.83	...	1.18	0.115	-3.40	19.83	0.51	6
03-035	MHO 8	0.29 (0.18,0.58)	4.99	...	0.87	...	0.080	-3.91	4.99	0.54	3
04-009	MHO 8	0.72 (0.50,0.97)	3.25	...	4.42	...	0.376	-3.24	3.25	0.50	6
04-010	GH Tau AB	0.11 (0.06,0.19)	...	12.29	...	1.20	0.120	-4.41	12.29	0.78	6
04-012	V807 Tau SNab	0.10 (0.09,0.12)	4.52	17.04	6.82	4.23	1.030	-3.89	7.51	1.22	83
18-004	KPNO-Tau 14	0.63 (0.59,0.70)	...	22.84	...	5.88	0.647	-2.82	22.84	1.15	28
17-000	CFHT-Tau 12	...	...	...	...	...	...	...	...	...	...
04-016	V830 Tau	0.04 (0.04,0.04)	8.12	22.49	15.05	30.57	4.807	-2.80	16.07	1.24	308
18-000	IRAS S04301+261	...	...	...	...	...	...	...	...	...	...
17-000	IRAS 04302+2247	...	...	...	...	...	...	...	...	...	...
17-027	IRAS 04303+2240	1.59 (1.41,1.78)	4.06	39.19	32.22	19.99	5.548	-3.18	9.67	0.89	114
04-034	GI Tau	0.41 (0.36,0.47)	4.87	32.58	4.23	3.76	0.872	-3.64	11.91	0.75	48
04-035	GK Tau AB	0.40 (0.37,0.55)	8.81	28.87	7.29	4.47	1.244	-3.64	13.83	1.04	79
18-019	IS Tau AB	0.65 (0.50,0.80)	4.41	17.86	9.41	4.70	1.310	-3.29	7.03	1.08	55
17-058	CI Tau	0.62 (0.31,1.10)	7.88	43.71	0.94	1.65	0.329	-4.01	23.44	0.52	10
18-030	IT Tau AB	0.68 (0.65,0.71)	11.71	37.33	11.76	38.57	6.307	-3.17	28.47	0.90	426
17-066	JH 108	0.17 (0.15,0.21)	8.81	24.93	4.70	5.64	1.115	-3.02	15.54	1.07	36
17-068	CFHT-BD Tau 1	0.65 (0.44,0.80)	...	15.77	...	1.88	0.174	-2.58	15.77	1.38	2
25-026	AA Tau	0.94 (0.84,1.03)	...	27.13	...	8.94	1.039	-3.47	27.13	0.82	43
09-010	HO Tau AB	0.64 (0.02,1.48)	3.25	...	0.47	...	0.289	-3.36	3.25	0.36	3
08-019	FF Tau AB	0.27 (0.21,0.28)	7.42	15.07	3.53	3.29	0.658	-3.61	10.45	1.13	58
12-040	DN Tau	0.06 (0.05,0.06)	8.81	23.42	5.17	5.17	1.072	-3.56	14.36	1.14	226
12-000	IRAS 04325+2402AB	...	...	...	...	...	...	...	...	...	...
12-059	CoKu Tau 3 AB	0.50 (0.48,0.52)	4.41	18.67	41.86	43.98	8.207	-2.66	9.24	1.01	387
09-022	KPNO-Tau 8	0.11 (0.09,0.13)	4.41	20.99	1.65	3.06	0.459	-2.25	12.16	0.80	58
08-037	HQ Tau AB	0.35 (0.32,0.37)	8.70	24.58	11.99	8.00	2.074	...	13.18	1.11	134
09-026	HQ Tau AB	0.66 (0.62,0.70)	4.75	24.23	75.72	34.81	10.724	...	7.93	0.83	165
08-043	KPNO-Tau 15	0.38 (0.37,0.41)	8.81	33.51	5.64	15.52	2.526	-2.33	23.47	0.92	260
09-031	KPNO-Tau 15	0.39 (0.25,0.55)	5.68	24.58	3.29	2.12	0.553	-2.99	10.08	1.22	10
08-000	KPNO-Tau 9	...	...	...	...	...	...	...	...	...	...
09-000	KPNO-Tau 9	...	...	...	...	...	...	...	...	...	...
08-048	HP Tau AB	0.48 (0.45,0.51)	8.58	23.88	10.11	10.58	2.187	-3.39	14.48	1.08	128
08-051a	HP Tau/G3 AB	0.53 (0.52,0.56)	4.64	20.64	12.46	6.82	1.839	-3.17	7.87	1.08	381
08-051	HP Tau/G2	0.53 (0.52,0.56)	4.64	20.64	93.83	51.27	13.830	-3.26	7.86	1.08	381
08-058	Haro 6-28 AB	0.34 (0.29,0.45)	...	10.55	...	2.12	0.202	-3.36	10.55	0.87	33
08-000	CFHT-BD Tau 2	...	...	...	...	...	...	...	...	...	...
08-080	CFHT-BD Tau 3	0.11 (0.00,0.41)	=10.00	...	0.09	...	0.009	-3.49	10.00	0.12	1
05-005	CFHT-Tau 6	0.32 (0.18,0.46)	9.97	...	0.40	...	0.040	-3.36	9.97	1.59	2
05-000	IRAS 04361+2547	...	...	...	...	...	...	...	...	...	...
05-013	GN Tau AB	2.42 (1.28,3.33)	...	14.38	...	4.23	0.423	-3.82	14.38	1.60	5
05-017	IRAS 04365+2535	24.3 (9.76,63.0)	...	49.16	...	3.29	0.499	-4.23	49.16	0.48	1
05-024	IRAS 04369+2539	6.15 (4.93,7.81)	...	32.70	...	10.35	1.315	-4.78	32.70	1.17	7
07-011	JH 223	0.09 (0.05,0.14)	8.70	...	0.71	...	0.061	-4.06	8.70	1.04	10
07-022	Haro 6-32	0.04 (0.00,0.08)	...	11.25	...	0.94	0.085	-3.74	11.25	0.47	6
07-000	ITG 33 A	...	...	...	...	...	...	...	...	...	...
07-000	CFHT-Tau 8	...	...	...	...	...	...	...	...	...	...
07-000	IRAS 04381+2540	...	...	...	...	...	...	...	...	...	...
07-041	IRAS 04385+2550AB	0.76 (0.60,0.97)	...	32.12	...	2.82	0.367	-3.28	32.12	0.70	9
10-017	CoKuLk332/G2 AB	0.63 (0.59,0.67)	4.75	22.96	33.63	11.52	4.313	-2.99	7.10	0.82	162

Table 6. continued.

XEST	Name	$N_{\text{H}}$ ( $1\sigma$ range) ( $10^{22}$ cm $^{-2}$ )	$T_1^a$ (MK)	$T_2$ (MK)	$\text{EM}_1^b$ ( $10^{52}$ )	$\text{EM}_2^b$ ( $10^{52}$ )	$L_{\text{X}}^c$ ( $10^{30}$ )	$\log L_{\text{X}}/L_{\odot}$	$T_{\text{av}}$ (MK)	$\chi_{\text{red}}^{2,d}$	d.o.f.
10-018	CoKuLk332/G1 AB	0.23 (0.15,0.31)	11.48	43.01	1.88	1.65	0.414	-4.20	21.26	0.64	20
10-020	V955 Tau AB	1.09 (0.83,1.34)	4.17	25.86	27.99	10.11	3.617	-3.03	6.77	1.00	33
10-034	CIDA 7	0.88 (0.58,1.11)	2.20	...	11.52	...	0.884	-2.34	2.20	0.58	6
10-045	DP Tau	0.00 (0.00,1.05)	3.25	43.83	0.05	0.61	0.101	-3.88	36.40	1.27	7
10-060	GO Tau	0.35 (0.26,0.55)	6.26	36.29	1.18	1.18	0.308	-3.67	15.07	1.53	12
26-012	2M J04552333+30	0.07 (0.00,0.24)	=10.00	...	0.15	...	0.015	-3.59	10.00	1.07	2
26-034	2M J04554046+30	0.04 (0.00,0.18)	=10.00	...	0.10	...	0.010	-3.91	10.00	0.17	1
26-043	AB Aur	0.06 (0.04,0.07)	2.43	7.54	1.18	2.59	0.353	-5.73	5.29	1.36	124
26-050	2MJ04554757/801	0.06 (0.02,0.12)	8.58	12.17	0.24	0.24	0.056	-3.88	10.22	1.03	20
26-067	SU Aur	0.56 (0.54,0.60)	5.22	23.30	87.01	32.45	11.641	-3.52	7.84	1.68	396
26-072	HBC 427	0.02 (0.01,0.03)	9.04	28.06	14.98	15.07	3.276	-3.11	15.96	0.92	106
Additional sources from Chandra											
C1-0	KPNO-Tau 10	...	...	...	...	...	...	...	...	...	...
C1-1	IRAS 04158+2805	3.24 (1.95,5.47)	...	70.72	...	2.59	0.414	-2.67	70.72	0.70	7
C2-1	Haro 6-5 B	26.4 (10.4,37.2)	...	15.54	...	43.27	3.774	-1.68	15.54	142.3C	512
C2-2	FS Tau AC	0.82 (0.58,1.05)	...	70.96	...	1.95	0.329	-3.57	70.96	1.02	12
C3-1	FV Tau/c AB	10.8 (5.70,16.4)	...	16.46	...	4.94	0.430	-3.27	16.46	96.3C	512
C3-2	DG Tau B	51.3 (11.0,63.0)	...	11.59	...	111.94	9.689	-3.34	11.59	70.20C	512
C4-1	GV Tau AB	...	...	...	...	...	...	...	...	...	...
C5-2	HN Tau AB	0.20	=10.00	...	...	...	0.153	-3.80	10.00	...	...
C5-1	L1551 55	0.14	=10.00	...	...	...	0.597	-3.38	10.00	...	...
C5-4	HD 28867	0.05	=10.00	...	...	...	6.867	-4.85	10.00	...	...
C5-3	DM Tau	0.12	=10.00	...	...	...	0.181	-3.80	10.00	...	...
C6-1	CFHT-BD Tau 4	0.52	...	14.26	...	0.94	0.096	-3.40	14.26	0.087	3
C6-0	L1527 IRS	...	...	...	...	...	...	...	...	...	...
C6-0	CFHT-Tau 17	...	...	...	...	...	...	...	...	...	...
C6-2	IRAS 04370+2559	1.09 (0.90,1.33)	...	50.55	...	4.47	0.666	-3.08	50.55	0.97	12

<sup>a</sup> “=” sign before number indicates that parameter was fixed

<sup>b</sup> EM in units of  $10^{52}$  cm $^{-3}$

<sup>c</sup>  $L_{\text{X}}$  for [0.3,10] keV, in units of  $10^{30}$  erg s $^{-1}$

<sup>d</sup> Numbers followed by ‘C’ denote C statistic (for low-background *Chandra* data)

Notes on individual objects:

- Fit performed for MOS1 + MOS2 spectra for: HBC 352 (XEST-27-115), CY Tau (23-002), CY Tau (24-002), FQ Tau (23-067), 2M-J04213459 (11-023), FX Tau (13-035), V827 Tau (22-097), FZ Tau (03-023), CFHT-Tau 5 (03-031), GH Tau (04-010), V955 Tau (10-020), 2M J04552333+30 (26-012), 2M J04554046+30 (26-034), 2M J04554757/801 (26-050), AB Aur (26-043), and SU Aur (26-067).
- HN Tau A (C5-2), L1551 55 (C5-1), DM Tau (C5-3), and HD 28867 (C5-4):  $T = 10$  MK was adopted for *Chandra* HRC observations, and  $N_{\text{H}}$  was derived from  $A_{\text{V}}$  assuming standard gas-to-dust ratios.
- DG Tau A (XEST-02-022), GV Tau (XEST-13-004), and DP Tau (XEST-10-045) require different  $N_{\text{H}}$  for the two components (also CW Tau; see Güdel et al. 2007b).  $N_{\text{H}}$  for cooler component is given. For hotter component:  $N_{\text{H}} = 1.8, 4.1,$  and  $3.8,$  respectively.
- CW Tau = XEST-20-046: spectrum cannot be reliably fitted. CW Tau may be two-absorber X-ray source (Güdel et al. 2007b).
- HP Tau/G2 and G3 = XEST-08-051 (separation:  $10''$ ) were fitted as one source. The ratio of the normalizations was derived from PSF fitting in the image.
- BP Tau = XEST-28-100: for an acceptable fit, two abundances were adjusted: O = 0.267, Fe = 0.10 (AG89).
- LkCa 1 = XEST-20-001: for an acceptable fit, Fe was adjusted: Fe = 0.57 (AG89).
- HL Tau = XEST-22-043: for an acceptable fit, Fe was adjusted: Fe = 1.06 (AG89).
- L1551 IRS 5 = XEST-22-040: the lightly absorbed source cannot originate from heavily absorbed protostar.



**Table 7.** Fundamental parameters of targets in XEST (1): Names and coordinates.

XEST	Name	2MASS <sup>a</sup>	IRAS <sup>b</sup>	Alternative names	RA(J2000.0) <sup>c</sup>			Dec(J2000.0) <sup>c</sup>		
					h	m	s	deg	'	"
27-115	HBC 352	03542950+3203013	...	NTTS 035120+3154SW	3	54	29.51	32	03	01.4
27-000	HBC 353	03543017+3203043	...	NTTS 035120+3154NE	3	54	30.17	32	03	04.3
06-005	HBC 358 AB	04034930+2610520	...	NTTS 040047+2603W	4	03	49.31	26	10	52.0
06-007	HBC 359	04035084+2610531	...	TTS 040047+2603	4	03	50.84	26	10	53.2
06-059	L1489 IRS	04044307+2618563	04016+2610	...	4	04	43.07	26	18	56.4
20-001	LkCa 1	04131414+2819108	...	HBC 365, V1095 Tau, JH 141	4	13	14.14	28	19	10.8
20-005	Anon 1	04132722+2816247	...	HBC 366, V1096 Tau	4	13	27.23	28	16	24.8
20-000	IRAS 04108+2803 A	04135328+2811233	04108+2803A	L1495N IRS	4	13	53.29	28	11	23.4
20-022	IRAS 04108+2803 B	04135471+2811328	04108+2803	...	4	13	54.72	28	11	32.9
20-000	2M J04141188+28	04141188+2811535	...	...	4	14	11.88	28	11	53.5
20-042	V773 Tau ABC	04141291+2812124	04111+2804	HBC 367, HD 283447	4	14	12.92	28	12	12.4
20-043	FM Tau	04141358+2812492	...	HBC 23, Haro 6-1	4	14	13.58	28	12	49.2
20-046	CW Tau	04141700+2810578	04112+2803	HBC 25	4	14	17.00	28	10	57.8
20-047	CIDA 1	04141760+2806096	...	C4101	4	14	17.61	28	06	09.7
20-056	MHO 2/1	04142639+2805597	04113+2758	...	4	14	26.40	28	05	59.7
20-058	MHO 3	04143054+2805147	04114+2757	...	4	14	30.55	28	05	14.7
20-069	FO Tau AB	04144928+2812305	04117+2804	HBC 369	4	14	49.29	28	12	30.6
20-073	CIDA 2	04150515+2808462	...	C3601	4	15	05.16	28	08	46.2
23-002	CY Tau	04173372+2820468	C04144+2813	HBC 28	4	17	33.73	28	20	46.9
24-002	CY Tau	04173372+2820468	C04144+2813	HBC 28	4	17	33.73	28	20	46.9
23-004	LkCa 5	04173893+2833005	...	HBC 371, JH 153	4	17	38.94	28	33	00.5
24-004	LkCa 5	04173893+2833005	...	HBC 371, JH 153	4	17	38.94	28	33	00.5
23-008	CIDA 3	04174965+2829362	...	C2701, V410 X1, Kim3-76	4	17	49.65	28	29	36.3
24-008	CIDA 3	04174965+2829362	...	C2701, V410 X1, Kim3-76	4	17	49.65	28	29	36.3
23-015	V410 X3	04180796+2826036	...	...	4	18	07.96	28	26	03.7
24-015	V410 X3	04180796+2826036	...	...	4	18	07.96	28	26	03.7
23-018	V410 A13	04181710+2828419	...	MHO 10	4	18	17.11	28	28	41.9
24-000	V410 A13	04181710+2828419	...	MHO 10	4	18	17.11	28	28	41.9
23-000	V410 A24	04182239+2824375	...	...	4	18	22.39	28	24	37.6
24-000	V410 A24	04182239+2824375	...	...	4	18	22.39	28	24	37.6
23-029	V410 A25	04182909+2826191	...	...	4	18	29.10	28	26	19.1
24-027	V410 A25	04182909+2826191	...	...	4	18	29.10	28	26	19.1
23-032	V410 Tau ABC	04183110+2827162	C04152+2820	HBC 29, HD 283518	4	18	31.10	28	27	16.2
24-028	V410 Tau ABC	04183110+2827162	C04152+2820	HBC 29, HD 283518	4	18	31.10	28	27	16.2
23-033	DD Tau AB	04183112+2816290	04154+2809	HBC 30	4	18	31.13	28	16	29.0
24-029	DD Tau AB	04183112+2816290	04154+2809	HBC 30	4	18	31.13	28	16	29.0
23-035	CZ Tau AB	04183158+2816585	...	HBC 31	4	18	31.59	28	16	58.5
24-030	CZ Tau AB	04183158+2816585	...	HBC 31	4	18	31.59	28	16	58.5
23-036	IRAS 04154+2823	04183203+2831153	04154+2823	...	4	18	32.03	28	31	15.4
24-031	IRAS 04154+2823	04183203+2831153	04154+2823	...	4	18	32.03	28	31	15.4
23-037	V410 X2	04183444+2830302	...	...	4	18	34.45	28	30	30.2
24-032	V410 X2	04183444+2830302	...	...	4	18	34.45	28	30	30.2
23-045	V410 X4	04184023+2824245	...	...	4	18	40.23	28	24	24.5
24-038	V410 X4	04184023+2824245	...	...	4	18	40.23	28	24	24.5
23-047	V892 Tau	04184061+2819155	04155+2812	HBC 373, Elias 1	4	18	40.62	28	19	15.5
24-040	V892 Tau	04184061+2819155	04155+2812	HBC 373, Elias 1	4	18	40.62	28	19	15.5
23-048	LR 1	04184133+2827250	...	...	4	18	41.33	28	27	25.0
24-000	LR 1	04184133+2827250	...	...	4	18	41.33	28	27	25.0
23-050	V410 X7	04184250+2818498	...	MHO-11	4	18	42.50	28	18	49.8
24-042	V410 X7	04184250+2818498	...	MHO-11	4	18	42.50	28	18	49.8
23-000	V410 A20	04184505+2820528	...	...	4	18	45.06	28	20	52.8
24-000	V410 A20	04184505+2820528	...	...	4	18	45.06	28	20	52.8
23-056	Hubble 4	04184703+2820073	04157+2813	V1023 Tau, HBC 374	4	18	47.04	28	20	07.3
24-047	Hubble 4	04184703+2820073	04157+2813	V1023 Tau, HBC 374	4	18	47.04	28	20	07.3
23-000	KPNO-Tau 2	04185115+2814332	...	...	4	18	51.16	28	14	33.2
24-000	KPNO-Tau 2	04185115+2814332	...	...	4	18	51.16	28	14	33.2
23-000	CoKu Tau 1	04185147+2820264	...	HBC 375	4	18	51.48	28	20	26.5
24-000	CoKu Tau 1	04185147+2820264	...	HBC 375	4	18	51.48	28	20	26.5
23-061	V410 X6	04190110+2819420	...	Kim3-89	4	19	01.11	28	19	42.0

Table 7. continued.

XEST	Name	2MASS <sup>a</sup>	IRAS <sup>b</sup>	Alternative names	RA(J2000.0) <sup>c</sup>			Dec(J2000.0) <sup>c</sup>		
					h	m	s	deg	'	''
24-054	V410 X6	04190110+2819420	...	Kim3-89	4	19	01.11	28	19	42.0
23-063	V410 X5	04190197+2822332	...	MHO-12	4	19	01.98	28	22	33.2
24-055	V410 X5	04190197+2822332	...	MHO-12	4	19	01.98	28	22	33.2
23-067	FQ Tau AB	04191281+2829330	C04161+2822	HBC 377, Haro 6-3	4	19	12.81	28	29	33.1
24-058	FQ Tau AB	04191281+2829330	C04161+2822	HBC 377, Haro 6-3	4	19	12.81	28	29	33.1
28-100	BP Tau	04191583+2906269	04161+2859	HD 281934, HBC 32	4	19	15.84	29	06	26.9
23-074	V819 Tau AB	04192625+2826142	C04162+2819	HBC 378, TAP 27	4	19	26.26	28	26	14.3
24-061	V819 Tau AB	04192625+2826142	C04162+2819	HBC 378, TAP 27	4	19	26.26	28	26	14.3
16-000	IRAS 04166+2706	...	04166+2706	...	4	19	43.00	27	13	33.7 (S)
16-000	IRAS 04169+2702	04195844+2709570	04169+2702	...	4	19	58.45	27	09	57.1
11-000	CFHT-Tau 19	04210795+2702204	...	...	4	21	07.95	27	02	20.4
11-000	IRAS 04181+2655	...	04181+2655	...	4	21	10.90	27	02	06.0 (S)
11-000	IRAS 04181+2654AB	04211146+2701094	04181+2654	...	4	21	11.47	27	01	09.4
11-023	2M J04213459	04213459+2701388	...	...	4	21	34.60	27	01	38.9
01-028	IRAS 04187+1927	04214323+1934133	04187+1927	...	4	21	43.24	19	34	13.3
11-037	CFHT-Tau 10	04214631+2659296	...	...	4	21	46.31	26	59	29.6
11-000	2M J04215450+2652	04215450+2652315	...	...	4	21	54.51	26	52	31.5
21-038	RY Tau	04215740+2826355	04188+2819	HBC 34, HD 283571	4	21	57.40	28	26	35.5
21-039	HD 283572	04215884+2818066	C04185+2811	V987 Tau, HBC 380	4	21	58.84	28	18	06.6
01-045	T Tau N(+Sab)	04215943+1932063	04190+1924	HD 284419, HBC 35	4	21	59.43	19	32	06.4
11-054	Haro 6-5 B	04220069+2657324	...	Haro 381, FS Tau B	4	22	00.70	26	57	32.5
11-057	FS Tau AC	04220217+2657304	04189+2650	HBC 383, Haro 6-5 A	4	22	02.18	26	57	30.5
21-044	LkCa 21	04220313+2825389	C04185+2818	V1071 Tau, HBC 382	4	22	03.14	28	25	39.0
01-054	RX J0422.1+1934	04220496+1934483	...	...	4	22	04.96	19	34	48.3
01-062	2M J04221332+1934	04221332+1934392	...	...	4	22	13.32	19	34	39.2
11-079	CFHT-Tau 21	04221675+2654570	...	...	4	22	16.76	26	54	57.1
02-013	FV Tau AB	04265352+2606543	C04238+2600	HBC 386, Haro 6-8	4	26	53.53	26	06	54.4
02-000	FV Tau/c AB	04265440+2606510	...	HBC 387	4	26	54.41	26	06	51.0
02-016	KPNO-Tau 13	04265732+2606284	...	...	4	26	57.33	26	06	28.4
02-000	DG Tau B	04270266+2605304	...	...	4	27	02.66	26	05	30.5
02-022	DG Tau A	04270469+2606163	04240+2559	HBC 37	4	27	04.70	26	06	16.3
02-000	KPNO-Tau 4	04272799+2612052	...	...	4	27	28.00	26	12	05.3
02-000	IRAS 04248+2612AB	04275730+2619183	04248+2612	HH 31 IRS 2	4	27	57.31	26	19	18.3
15-020	JH 507	04292071+2633406	...	J1-507	4	29	20.71	26	33	40.7
13-004	GV Tau AB	04292373+2433002	04263+2426	HBC 389, Haro 6-10 AB	4	29	23.73	24	33	00.3
13-000	IRAS 04264+2433	04293008+2439550	04264+2433	...	4	29	30.08	24	39	55.1
15-040	DH Tau AB	04294155+2632582	04267+2626	HBC 38	4	29	41.56	26	32	58.3
15-042	DI Tau AB	04294247+2632493	...	HBC 39	4	29	42.48	26	32	49.3
15-044	KPNO-Tau 5	04294568+2630468	...	...	4	29	45.68	26	30	46.8
14-006	IQ Tau A	04295156+2606448	04267+2600	HBC 41, LkHa 265	4	29	51.56	26	06	44.9
13-000	CFHT-Tau 20	04295950+2433078	...	...	4	29	59.51	24	33	07.9
14-000	KPNO-Tau 6	04300724+2608207	...	...	4	30	07.24	26	08	20.8
13-035	FX Tau AB	04302961+2426450	04267+2420	HBC 44, Haro 6-11	4	30	29.61	24	26	45.0
14-057	DK Tau AB	04304425+2601244	04276+2554	HBC 45	4	30	44.25	26	01	24.5
14-000	KPNO-Tau 7	04305718+2556394	...	...	4	30	57.19	25	56	39.5
22-013	MHO 9	04311578+1820072	...	...	4	31	15.78	18	20	07.2
22-021	MHO 4	04312405+1800215	...	RXJ0431.4+1800	4	31	24.06	18	00	21.5
22-040	L1551 IRS5	04313407+1808049	04287+1801	...	4	31	34.08	18	08	04.9
22-042	LkHa 358	04313613+1813432	...	CoKu Tau 2, HBC 394	4	31	36.13	18	13	43.3
22-000	HH 30	04313747+1812244	04287+1806	V1213 Tau	4	31	37.47	18	12	24.5
22-043	HL Tau	04313843+1813576	04287+1807	HBC 49, Haro 6-14	4	31	38.44	18	13	57.7
22-047	XZ Tau AB	04314007+1813571	...	HBC 50, Haro 6-15	4	31	40.07	18	13	57.2
22-056	L1551 NE	04314444+1808315	...	...	4	31	44.45	18	08	31.5
03-005	HK Tau AB	04315056+2424180	04288+2417	HBC 48, Haro 6-12	4	31	50.57	24	24	18.1
22-070	V710 Tau BA	04315779+1821380	04290+1815	HBC 395+51, LkHa266S	4	31	57.79	18	21	38.1
19-009	JH 665	04315844+2543299	...	J1-665	4	31	58.44	25	43	29.9
22-089	L1551 51	04320926+1757227	042916+1751	V1075 Tau, HBC 397	4	32	09.27	17	57	22.8
22-097	V827 Tau	04321456+1820147	...	HBC 399, TAP 42	4	32	14.57	18	20	14.7
03-016	Haro 6-13	04321540+2428597	04292+2422	V806 Tau, HBC 396	4	32	15.41	24	28	59.7

**Table 7.** continued.

XEST	Name	2MASS <sup>a</sup>	IRAS <sup>b</sup>	Alternative names	RA(J2000.0) <sup>c</sup>			Dec(J2000.0) <sup>c</sup>		
					h	m	s	deg	'	''
22-100	V826 Tau	04321583+1801387	...	HBC 400, TAP 43	4 32	15.84	18 01	38.7		
22-101	MHO 5	04321606+1812464	...	...	4 32	16.07	18 12	46.4		
03-017	CFHT-Tau 7	04321786+2422149	...	...	4 32	17.86	24 22	15.0		
03-019	V928 Tau AB	04321885+2422271	...	HK Tau/G2, JH 91, HBC 398	4 32	18.86	24 22	27.1		
03-022	FY Tau	04323058+2419572	04294+2413	HBC 401, Haro 6-17	4 32	30.58	24 19	57.3		
03-023	FZ Tau	04323176+2420029	...	HBC 402, Haro 6-18	4 32	31.76	24 20	03.0		
17-002	IRAS 04295+2251	04323205+2257266	04295+2251	L1536 IRS	4 32	32.05	22 57	26.7		
19-049	UZ Tau E+W(AB)	04324303+2552311	04296+2546	HBC 52	4 32	43.04	25 52	31.1		
17-009	JH 112	04324911+2253027	04298+2246	...	4 32	49.11	22 53	02.8		
03-031	CFHT-Tau 5	04325026+2422115	...	...	4 32	50.27	24 22	11.6		
04-003	CFHT-Tau 5	04325026+2422115	...	...	4 32	50.27	24 22	11.6		
03-035	MHO 8	04330197+2421000	...	...	4 33	01.98	24 21	00.0		
04-009	MHO 8	04330197+2421000	...	...	4 33	01.98	24 21	00.0		
04-010	GH Tau AB	04330622+2409339	04300+2403	HBC 55, Haro 6-20	4 33	06.22	24 09	34.0		
04-012	V807 Tau SNab	04330664+2409549	...	HBC 404, Elias 12	4 33	06.64	24 09	55.0		
18-004	KPNO-Tau 14	04330781+2616066	...	...	4 33	07.81	26 16	06.6		
17-000	CFHT-Tau 12	04330945+2246487	...	...	4 33	09.46	22 46	48.7		
04-016	V830 Tau	04331003+2433433	C04301+2427	HBC 405, TAP 46	4 33	10.03	24 33	43.4		
18-000	IRAS S04301+261	04331435+2614235	...	...	4 33	14.36	26 14	23.5		
17-000	IRAS 04302+2247	04331650+2253204	04302+2247	Butterfly Star	4 33	16.50	22 53	20.4		
17-027	IRAS 04303+2240	04331907+2246342	04303+2240	L1536S	4 33	19.07	22 46	34.2		
04-034	GI Tau	04333405+2421170	04305+2414	HBC 56, Haro 6-21	4 33	34.06	24 21	17.0		
04-035	GK Tau AB	04333456+2421058	...	HBC 57, Haro 6-22	4 33	34.56	24 21	05.9		
18-019	IS Tau AB	04333678+2609492	04308+2607	HBC 59, Haro 6-23	4 33	36.79	26 09	49.2		
17-058	CI Tau	04335200+2250301	04308+2244	HBC 61, Haro 6-25	4 33	52.00	22 50	30.2		
18-030	IT Tau AB	04335470+2613275	...	Haro 6-26	4 33	54.70	26 13	27.5		
17-066	JH 108	04341099+2251445	...	...	4 34	10.99	22 51	44.5		
17-068	CFHT-BD Tau 1	04341527+2250309	...	...	4 34	15.27	22 50	31.0		
25-026	AA Tau	04345542+2428531	04318+2422	HBC 63	4 34	55.42	24 28	53.2		
09-010	HO Tau AB	04352020+2232146	C04323+2226	HBC 64, Haro 6-27	4 35	20.20	22 32	14.6		
08-019	FF Tau AB	04352089+2254242	...	HBC 409	4 35	20.90	22 54	24.2		
12-040	DN Tau	04352737+2414589	04324+2408	HBC 65	4 35	27.37	24 14	58.9		
12-000	IRAS 04325+2402AB	04353539+2408194	04325+2402	L1535 IRS	4 35	35.39	24 08	19.4		
12-059	CoKu Tau 3 AB	04354093+2411087	...	HBC 411	4 35	40.94	24 11	08.8		
09-022	KPNO-Tau 8	04354183+2234115	...	...	4 35	41.84	22 34	11.6		
08-037	HQ Tau AB	04354733+2250216	04327+2244	...	4 35	47.34	22 50	21.7		
09-026	HQ Tau AB	04354733+2250216	04327+2244	...	4 35	47.34	22 50	21.7		
08-043	KPNO-Tau 15	04355109+2252401	...	...	4 35	51.10	22 52	40.1		
09-031	KPNO-Tau 15	04355109+2252401	...	...	4 35	51.10	22 52	40.1		
08-000	KPNO-Tau 9	04355143+2249119	...	...	4 35	51.43	22 49	11.9		
09-000	KPNO-Tau 9	04355143+2249119	...	...	4 35	51.43	22 49	11.9		
08-048	HP Tau AB	04355277+2254231	04328+2248	HBC 66, LkHa 258	4 35	52.78	22 54	23.1		
08-051a	HP Tau/G3 AB	04355349+2254089	...	HBC 414	4 35	53.50	22 54	09.0		
08-051	HP Tau/G2	04355415+2254134	...	V1025 Tau, HBC 415	4 35	54.15	22 54	13.5		
08-058	Haro 6-28 AB	04355684+2254360	...	V1026 Tau, HBC 416	4 35	56.84	22 54	36.0		
08-000	CFHT-BD Tau 2	04361038+2259560	...	...	4 36	10.39	22 59	56.0		
08-080	CFHT-BD Tau 3	04363893+2258119	...	...	4 36	38.94	22 58	11.9		
05-005	CFHT-Tau 6	04390396+2544264	...	...	4 39	03.96	25 44	26.4		
05-000	IRAS 04361+2547	04391389+2553208	04361+2547	TMR 1, ITG 12	4 39	13.89	25 53	20.9		
05-013	GN Tau AB	04392090+2545021	04362+2539	Haro 6-31	4 39	20.91	25 45	02.1		
05-017	IRAS 04365+2535	04393519+2541447	04365+2535	TMC-1A	4 39	35.19	25 41	44.7		
05-024	IRAS 04369+2539	04395574+2545020	04369+2539	Elias 18, IC 2087/IR	4 39	55.75	25 45	02.0		
07-011	JH 223	04404950+2551191	...	...	4 40	49.51	25 51	19.2		
07-022	Haro 6-32	04410424+2557561	...	...	4 41	04.24	25 57	56.1		
07-000	ITG 33 A	04410826+2556074	...	...	4 41	08.26	25 56	07.5		
07-000	CFHT-Tau 8	04411078+2555116	...	...	4 41	10.78	25 55	11.7		
07-000	IRAS 04381+2540	04411267+2546354	04381+2540	TMC-1, ITG 35	4 41	12.68	25 46	35.4		
07-041	IRAS 04385+2550AB	04413882+2556267	04385+2550	Haro 6-33	4 41	38.82	25 56	26.8		
10-017	CoKuLk332/G2 AB	04420548+2522562	...	V999 Tau, HBC 422	4 42	05.49	25 22	56.3		

**Table 7.** continued.

XEST	Name	2MASS <sup>a</sup>	IRAS <sup>b</sup>	Alternative names	RA(J2000.0) <sup>c</sup>			Dec(J2000.0) <sup>c</sup>		
					h	m	s	deg	'	''
10-018	CoKuLk332/G1 AB	04420732+2523032	...	HBC 423	4	42	07.33	25	23	03.2
10-020	V955 Tau AB	04420777+2523118	04390+2517	HBC 69, JH 226, LkHa 332	4	42	07.77	25	23	11.8
10-034	CIDA 7	04422101+2520343	...	C31	4	42	21.02	25	20	34.4
10-045	DP Tau	04423769+2515374	04395+2509	HBC 70	4	42	37.70	25	15	37.5
10-060	GO Tau	04430309+2520187	C04400+2514	HBC 71	4	43	03.09	25	20	18.8
26-012	2M J04552333+30	04552333+3027366	...	...	4	55	23.33	30	27	36.6
26-034	2M J04554046+30	04554046+3039057	...	...	4	55	40.46	30	39	05.7
26-043	AB Aur	04554582+3033043	04525+3028	HD 31293, HBC 78	4	55	45.83	30	33	04.4
26-050	2MJ04554757/801	04554757+3028077	...	...	4	55	47.57	30	28	07.7
26-067	SU Aur	04555938+3034015	04528+3029	HBC 79, HD 282624	4	55	59.38	30	34	01.6
26-072	HBC 427	04560201+3021037	...	V397 Aur	4	56	02.02	30	21	03.8
Additional sources from Chandra										
C1-0	KPNO-Tau 10	04174955+2813318	...	...	4	17	49.55	28	13	31.9
C1-1	IRAS 04158+2805	04185813+2812234	04158+2805	...	4	18	58.14	28	12	23.5
C2-1	Haro 6-5 B	04220069+2657324	...	Haro 381, FS Tau B	4	22	00.70	26	57	32.5
C2-2	FS Tau AC	04220217+2657304	04189+2650	HBC 383, Haro 6-5 A	4	22	02.18	26	57	30.5
C3-1	FV Tau/c AB	04265440+2606510	...	HBC 387	4	26	54.41	26	06	51.0
C3-2	DG Tau B	04270266+2605304	...	...	4	27	02.66	26	05	30.5
C4-1	GV Tau AB	04292373+2433002	04263+2426	HBC 389, Haro 6-10 AB	4	29	23.73	24	33	00.3
C5-2	HN Tau AB	04333935+1751523	...	...	4	33	39.35	17	51	52.4
C5-1	L1551 55	04324373+1802563	042950+1757	...	4	32	43.73	18	02	56.3
C5-4	HD 28867	04333297+1801004	04306+1754	HR 1442, L1551 53	4	33	32.98	18	01	00.4
C5-3	DM Tau	04334871+1810099	04309+1803	HBC 62	4	33	48.72	18	10	10.0
C6-1	CFHT-BD Tau 4	04394748+2601407	04368+2557N3	...	4	39	47.48	26	01	40.8
C6-0	L1527 IRS	...	04368+2557	...	4	39	53.59	26	03	05.5 (S)
C6-0	CFHT-Tau 17	04400174+2556292	...	...	4	40	01.75	25	56	29.2
C6-2	IRAS 04370+2559	04400800+2605253	04370+2559	...	4	40	08.00	26	05	25.4

<sup>a</sup> Nearest 2MASS entry within 5'' to coordinates given by references 5, 18, 33, or SIMBAD. Unlikely identifications in parentheses.<sup>b</sup> Nearest IRAS catalog entry, within 10''.<sup>c</sup> 2MASS coordinates. For unlikely identifications, SIMBAD (S).

**Table 8.** Fundamental parameters of targets in XEST (2): Multiplicity.

XEST	Name	Comp	Separations (")	Refs
27-115	HBC 352	1	...	...
27-000	HBC 353	1	...	...
06-005	HBC 358 AB	3	0.15, 1.55	11, 20, 31, 40
06-007	HBC 359	1	...	...
06-059	L1489 IRS	1	...	...
20-001	LkCa 1	1	...	...
20-005	Anon 1	1	...	...
20-000	IRAS 04108+2803 A	1	...	...
20-022	IRAS 04108+2803 B	1	...	...
20-000	2M J04141188+28	1	...	...
20-042	V773 Tau ABC	4	0.002,0.11,0.2	16, 17, 31, 58, 61, 62
20-043	FM Tau	1	...	...
20-046	CW Tau	1	...	...
20-047	CIDA 1	1	...	...
20-056	MHO 2/1	2	4.0	13
20-058	MHO 3	1	...	...
20-069	FO Tau AB	2	0.16	16, 20, 31, 58
20-073	CIDA 2	1	...	...
23-002	CY Tau	1	...	...
24-002	CY Tau	1	...	...
23-004	LkCa 5	1	...	...
24-004	LkCa 5	1	...	...
23-008	CIDA 3	1	...	...
24-008	CIDA 3	1	...	...
23-015	V410 X3	1	...	...
24-015	V410 X3	1	...	...
23-018	V410 A13	1	...	...
24-000	V410 A13	1	...	...
23-000	V410 A24	1	...	...
24-000	V410 A24	1	...	...
23-029	V410 A25	1	...	...
24-027	V410 A25	1	...	...
23-032	V410 Tau ABC	3	0.12, 0.29	16, 17, 58
24-028	V410 Tau ABC	3	0.12, 0.29	16, 17, 58
23-033	DD Tau AB	2	0.54	16, 20, 31, 40, 47, 58
24-029	DD Tau AB	2	0.54	16, 20, 31, 40, 47, 58
23-035	CZ Tau AB	2	0.33	31, 40, 58
24-030	CZ Tau AB	2	0.33	31, 40, 58
23-036	IRAS 04154+2823	1	...	...
24-031	IRAS 04154+2823	1	...	...
23-037	V410 X2	1	...	...
24-032	V410 X2	1	...	...
23-045	V410 X4	1	...	...
24-038	V410 X4	1	...	...
23-047	V892 Tau	3	0.05, 4	32, 54
24-040	V892 Tau	3	0.05, 4	32, 54
23-048	LR 1	1	...	...
24-000	LR 1	1	...	...
23-050	V410 X7	1	...	...
24-042	V410 X7	1	...	...
23-000	V410 A20	1	...	...
24-000	V410 A20	1	...	...
23-056	Hubble 4	1	...	...
24-047	Hubble 4	1	...	...
23-000	KPNO-Tau 2	1	...	...
24-000	KPNO-Tau 2	1	...	...
23-000	CoKu Tau 1	1	...	...
24-000	CoKu Tau 1	1	...	...
23-061	V410 X6	1	...	...

**Table 8.** continued.

XEST	Name	Comp	Separations (")	Refs
24-054	V410 X6	1	...	...
23-063	V410 X5	1	...	...
24-055	V410 X5	1	...	...
23-067	FQ Tau AB	2	0.73	20, 31, 40, 58
24-058	FQ Tau AB	2	0.73	20, 31, 40, 58
28-100	BP Tau	1	...	...
23-074	V819 Tau AB	2	10.5	31 <sup>a</sup>
24-061	V819 Tau AB	2	10.5	31 <sup>a</sup>
16-000	IRAS 04166+2706	1	...	...
16-000	IRAS 04169+2702	1	...	...
11-000	CFHT-Tau 19	1	...	...
11-000	IRAS 04181+2655	1	...	...
11-000	IRAS 04181+2654AB	2	...	...
11-023	2M J04213459	1	...	...
01-028	IRAS 04187+1927	1	...	...
11-037	CFHT-Tau 10	1	...	...
11-000	2M J04215450+2652	1	...	...
21-038	RY Tau	1	...	...
21-039	HD 283572	1	...	...
01-045	T Tau N(+Sab)	3	0.67, 0.13	12, 47, 58
11-054	Haro 6-5 B	1	...	...
11-057	FS Tau AC	2	0.30	20, 31, 40, 52, 53, 58
21-044	LkCa 21	1	...	...
01-054	RX J0422.1+1934	1	...	...
01-062	2M J04221332+1934	1	...	...
11-079	CFHT-Tau 21	1	...	...
02-013	FV Tau AB	2	0.71	16, 31, 40, 47, 53, 58
02-000	FV Tau/c AB	2	0.67	31, 40, 52, 53, 58
02-016	KPNO-Tau 13	1	...	...
02-000	DG Tau B	1	...	...
02-022	DG Tau A	1	...	...
02-000	KPNO-Tau 4	1	...	...
02-000	IRAS 04248+2612AB	2	4.55	13
15-020	JH 507	1	...	...
13-004	GV Tau AB	2	1.2	30, 31, 47, 53, 58
13-000	IRAS 04264+2433	1	...	...
15-040	DH Tau AB	2	2.3	25
15-042	DI Tau AB	2	0.12	16, 53
15-044	KPNO-Tau 5	1	...	...
14-006	IQ Tau A	1	...	<sup>b</sup>
13-000	CFHT-Tau 20	1	...	...
14-000	KPNO-Tau 6	1	...	...
13-035	FX Tau AB	2	0.85	11, 16, 31, 40, 53, 58
14-057	DK Tau AB	2	2.27	11, 31, 40, 43, 47, 52, 53, 58
14-000	KPNO-Tau 7	1	...	...
22-013	MHO 9	1	...	...
22-021	MHO 4	1	...	...
22-040	L1551 IRS5	2	0.3	51
22-042	LkHa 358	1	...	...
22-000	HH 30	1	...	...
22-043	HL Tau	1	...	...
22-047	XZ Tau AB	2	0.3	16, 20, 31, 47, 58
22-056	L1551 NE	2	0.5	64
03-005	HK Tau AB	2	2.4	11, 31, 40, 42, 43, 53, 58
22-070	V710 Tau BA	2	3.1	21, 31, 40, 43, 58
19-009	JH 665	1	...	...
22-089	L1551 51	1	...	...
22-097	V827 Tau	1	...	...
03-016	Haro 6-13	1	...	...

**Table 8.** continued.

XEST	Name	Comp	Separations (")		Refs
22-100	V826 Tau	1	...	...	...
22-101	MHO 5	1	...	...	...
03-017	CFHT-Tau 7	1	...	...	...
03-019	V928 Tau AB	2	0.17	...	16, 31, 53, 58
03-022	FY Tau	1	...	...	<sup>c</sup>
03-023	FZ Tau	1	...	...	...
17-002	IRAS 04295+2251	1	...	...	...
19-049	UZ Tau E+W(AB)	3	3.5, 0.36	...	16, 17, 20, 21, 31, 40, 52, 53, 58
17-009	JH 112	1	...	...	...
03-031	CFHT-Tau 5	1	...	...	...
04-003	CFHT-Tau 5	1	...	...	...
03-035	MHO 8	1	...	...	...
04-009	MHO 8	1	...	...	...
04-010	GH Tau AB	2	0.31	...	16, 31, 40, 58
04-012	V807 Tau SNab	3	0.3, 0.023	...	16, 20, 31, 53, 58
18-004	KPNO-Tau 14	1	...	...	...
17-000	CFHT-Tau 12	1	...	...	...
04-016	V830 Tau	1	...	...	...
18-000	IRAS S04301+261	1	...	...	...
17-000	IRAS 04302+2247	1	...	...	...
17-027	IRAS 04303+2240	1	...	...	...
04-034	GI Tau	1	...	...	...
04-035	GK Tau AB	2	2.5	...	21, 49
18-019	IS Tau AB	2	0.22	...	16, 20, 53, 58
17-058	CI Tau	1	...	...	...
18-030	IT Tau AB	2	2.4	...	11, 40, 52, 53, 58
17-066	JH 108	1	...	...	...
17-068	CFHT-BD Tau 1	1	...	...	...
25-026	AA Tau	1	...	...	...
09-010	HO Tau AB	2	6.9	...	21 <sup>d</sup>
08-019	FF Tau AB	2	0.03	...	52, 53
12-040	DN Tau	1	...	...	...
12-000	IRAS 04325+2402AB	1	8.15	...	13
12-059	CoKu Tau 3 AB	2	2.06	...	31, 53, 58
09-022	KPNO-Tau 8	1	...	...	...
08-037	HQ Tau AB	2	0.009	...	52 <sup>e</sup>
09-026	HQ Tau AB	2	0.009	...	52 <sup>e</sup>
08-043	KPNO-Tau 15	1	...	...	...
09-031	KPNO-Tau 15	1	...	...	...
08-000	KPNO-Tau 9	1	...	...	...
09-000	KPNO-Tau 9	1	...	...	...
08-048	HP Tau AB	2	0.017	...	50, 53
08-051a	HP Tau/G3 AB	2	0.022	...	50, 53
08-051	HP Tau/G2	1	...	...	...
08-058	Haro 6-28 AB	2	0.65	...	20, 31, 40, 53, 58
08-000	CFHT-BD Tau 2	1	...	...	...
08-080	CFHT-BD Tau 3	1	...	...	...
05-005	CFHT-Tau 6	1	...	...	...
05-000	IRAS 04361+2547	1	...	...	...
05-013	GN Tau AB	2	0.36	...	52, 53, 58, 67 <sup>f</sup>
05-017	IRAS 04365+2535	1	...	...	...
05-024	IRAS 04369+2539	1	...	...	...
07-011	JH 223	1	...	...	...
07-022	Haro 6-32	1	...	...	...
07-000	ITG 33 A	1	...	...	<sup>g</sup>
07-000	CFHT-Tau 8	1	...	...	...
07-000	IRAS 04381+2540	1	...	...	...
07-041	IRAS 04385+2550AB	2	18.9	...	13
10-017	CoKuLk332/G2 AB	2	0.27	...	20, 31, 58

**Table 8.** continued.

XEST	Name	Comp	Separations ( $''$ )	Refs
10-018	CoKuLk332/G1 AB	2	0.23	16, 31, 58
10-020	V955 Tau AB	2	0.34	20, 31, 40, 58
10-034	CIDA 7	1	...	...
10-045	DP Tau	1	...	...
10-060	GO Tau	1	...	...
26-012	2M J04552333+30	1	...	...
26-034	2M J04554046+30	1	...	...
26-043	AB Aur	1	...	...
26-050	2MJ04554757/801	1	...	...
26-067	SU Aur	1	...	...
26-072	HBC 427	1	...	...
Additional sources from Chandra				
C1-0	KPNO-Tau 10	1	...	...
C1-1	IRAS 04158+2805	1	...	...
C2-1	Haro 6-5 B	1	...	...
C2-2	FS Tau AC	2	0.25	20, 31, 52, 53, 58
C3-1	FV Tau/c AB	2	0.67	31, 40, 52, 53, 58
C3-2	DG Tau B	1	...	...
C4-1	GV Tau AB	2	1.2	30, 31, 47, 53, 58
C5-2	HN Tau AB	2	3.1	11, 21, 31, 42, 43, 58
C5-1	L1551 55	1	...	...
C5-4	HD 28867	3	3.08	65
C5-3	DM Tau	1	...	...
C6-1	CFHT-BD Tau 4	1	...	...
C6-0	L1527 IRS	1	...	...
C6-0	CFHT-Tau 17	1	...	...
C6-2	IRAS 04370+2559	1	...	...

<sup>a</sup> V819 Tau B suspected by Woitas (2001) to be a background star.

<sup>b</sup> IQ Tau B, 10.2 $''$  from IQ Tau A = XEST-14-006, has been interpreted as an extinguished background G star by Hartigan et al. (1994).

<sup>c</sup> FY Tau is a suspected binary with separation of 0.153 $''$  (Richichi et al. 1994).

<sup>d</sup> HO Tau B suspected by Hartigan et al. (1994) to be a background star.

<sup>e</sup> HQ Tau B not confirmed by Simon et al. (1995).

<sup>f</sup> GN Tau AB: A component detected from lunar occultation by Simon et al. (1995) at a projected separation of 0.041 $''$  may be identical to GN Tau B at  $\approx 0.36''$  (Simon et al. 1996).

<sup>g</sup> ITG 33 B, 5.2 $''$  from ITG 33 A, has been interpreted as a reddened background object by Martín (2000).



**Table 9.** Fundamental parameters of targets in XEST (3): Photometry and spectroscopy.

XEST	Name	Spec <sup>a</sup>	Refs.	$A_V^a$ (mag)	$A_J$ (mag)	$T_{\text{eff}}^a$ (K)	$L_*^b$ ( $L_{\odot}$ )	Refs
27-115	HBC 352	G0	27	0.87	0.25	6030	0.740	27
27-000	HBC 353	G5	27	0.97	0.28	5770	0.500	27
06-005	HBC 358 AB	M2	27	0.21	0.06	3580	0.280	27
06-007	HBC 359	M2	27	0.49	0.14	3580	0.260	27
06-059	L1489 IRS	K4	59	10.20	...	4500	4.900	10, 59
20-001	LkCa 1	M4	5	0.00	0.00	3270	0.380	5, 27
20-005	Anon 1	M0	5	1.32	1.03	3850	2.600	5, 27
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	...	...	...	...	3500	0.400	10
20-000	2M J04141188+28	M6.25	33	...	0.28	2962	0.015	33
20-042	V773 Tau ABC	K2/M0	58	1.39	0.31	4898/3873	1.89/ 1.17/ 5.60	5, 58
20-043	FM Tau	M0	5	0.69	0.59	3850	0.460	5, 27
20-046	CW Tau	K3	5	2.29	0.55	4730	1.100	5, 27
20-047	CIDA 1	M5.5	57	...	...	...	...	...
20-056	MHO 2/1	M2.5/2.5	5	4.58	1.85	3488/3488	0.42/ 0.14/ 0.56	5, 4
20-058	MHO 3	K7	5	6.01	2.15	4060	0.910	5, 4
20-069	FO Tau AB	M2	5	1.87	0.63	3556/3556	0.48/ 0.45/ 0.77	5, 58, 27
20-073	CIDA 2	M5.5	5	0.83	0.28	3058	0.320	5, 27
23-002	CY Tau	M1.5	5	0.10	0.23	3632	0.500	5, 27
24-002	CY Tau	M1.5	5	0.10	0.23	3632	0.500	5, 27
23-004	LkCa 5	M2	27	0.10	0.11	3560	0.370	5, 27
24-004	LkCa 5	M2	27	0.10	0.11	3560	0.370	5, 27
23-008	CIDA 3	M4	5	...	0.26	3270	0.140	5
24-008	CIDA 3	M4	5	...	0.26	3270	0.140	5
23-015	V410 X3	M6	5	0.54	0.17	2990	0.084	5, 58
24-015	V410 X3	M6	5	0.54	0.17	2990	0.084	5, 58
23-018	V410 A13	M5.75	5	2.78	0.79	3024	0.039	5, 4
24-000	V410 A13	M5.75	5	2.78	0.79	3024	0.039	5, 4
23-000	V410 A24	G1	5	...	6.73	5945	2.800	5
24-000	V410 A24	G1	5	...	6.73	5945	2.800	5
23-029	V410 A25	M1	5	...	6.55	3705	1.600	5
24-027	V410 A25	M1	5	...	6.55	3705	1.600	5
23-032	V410 Tau ABC	K4	58	0.67	0.00	4602/3076	2.15/ 0.05/ 2.20	58
24-028	V410 Tau ABC	K4	58	0.67	0.00	4602/3076	2.15/ 0.05/ 2.20	58
23-033	DD Tau AB	M3	5	0.39/ 0.39	0.05	3412/3412	0.13/ 0.13/ 0.34	5, 58
24-029	DD Tau AB	M3	5	0.39/ 0.39	0.05	3412/3412	0.13/ 0.13/ 0.34	5, 58
23-035	CZ Tau AB	M3	5	1.32	0.48	3415	0.270	5, 27
24-030	CZ Tau AB	M3	5	1.32	0.48	3415	0.270	5, 27
23-036	IRAS 04154+2823	M2.5	5	...	4.26	3488	0.130	5
24-031	IRAS 04154+2823	M2.5	5	...	4.26	3488	0.130	5
23-037	V410 X2	M0	5	...	6.03	3850	3.000	5
24-032	V410 X2	M0	5	...	6.03	3850	3.000	5
23-045	V410 X4	M4	5	...	5.29	3270	1.300	5
24-038	V410 X4	M4	5	...	5.29	3270	1.300	5
23-047	V892 Tau	B9	5	5.93	2.39	10500	77.000	5, 27
24-040	V892 Tau	B9	5	5.93	2.39	10500	77.000	5, 27
23-048	LR 1	K4.5	5	...	6.40	4470	0.470	5
24-000	LR 1	K4.5	5	...	6.40	4470	0.470	5
23-050	V410 X7	M0.75	5	8.18	2.34	3741	0.500	5, 58
24-042	V410 X7	M0.75	5	8.18	2.34	3741	0.500	5, 58
23-000	V410 A20	K3	5	...	6.57	4730	0.510	5
24-000	V410 A20	K3	5	...	6.57	4730	0.510	5
23-056	Hubble 4	K7	5	0.76	0.68	4060	2.700	5, 27
24-047	Hubble 4	K7	5	0.76	0.68	4060	2.700	5, 27
23-000	KPNO-Tau 2	M6.75	18	0.37	0.00	2889	0.007	18, 5
24-000	KPNO-Tau 2	M6.75	18	0.37	0.00	2889	0.007	18, 5
23-000	CoKu Tau 1	K7	59	6.80	...	4000	0.150	59
24-000	CoKu Tau 1	K7	59	6.80	...	4000	0.150	59
23-061	V410 X6	M5.5	5	...	0.17	3058	0.200	5

Table 9. continued.

XEST	Name	Spec <sup>a</sup>	Refs.	$A_V^a$ (mag)	$A_I$ (mag)	$T_{\text{eff}}^a$ (K)	$L_*^b$ ( $L_\odot$ )	Refs.
24-054	V410 X6	M5.5	5	...	0.17	3058	0.200	5
23-063	V410 X5	M5.5	5	3.36	0.72	3058	0.083	5, 58
24-055	V410 X5	M5.5	5	3.36	0.72	3058	0.083	5, 58
23-067	FQ Tau AB	M3/M3.5	20	1.95/ 1.80	0.16	3416/3345	0.20/ 0.28/ 0.21	5, 20
24-058	FQ Tau AB	M3/M3.5	20	1.95/ 1.80	0.16	3416/3345	0.20/ 0.28/ 0.21	5, 20
28-100	BP Tau	K7	27	0.49	0.14	4060	0.950	27
23-074	V819 Tau AB	K7	5	1.35	0.48	4060	0.910	5, 27
24-061	V819 Tau AB	K7	5	1.35	0.48	4060	0.910	5, 27
16-000	IRAS 04166+2706	<M0	66	...	...	...	...	...
16-000	IRAS 04169+2702	...	...	...	...	...	0.800	13
11-000	CFHT-Tau 19	M5.25	18	7.30	...	3100	0.072	18
11-000	IRAS 04181+2655	...	...	...	...	4000	1.800	10
11-000	IRAS 04181+2654AB	...	...	...	...	...	0.25/ 0.25/ 0.50	13
11-023	2M J04213459	M5.5	33	...	0.49	3058	0.065	33
01-028	IRAS 04187+1927	M0	27	...	...	3850	...	27
11-037	CFHT-Tau 10	M5.75	66	3.59	...	3030	0.021	18, 19
11-000	2M J04215450+2652	M8.5	66	2.97	...	2642	0.003	19
21-038	RY Tau	K1	27	1.84	0.53	5080	7.600	27
21-039	HD 283572	G5	27	0.38	0.11	5770	6.500	27
01-045	T Tau N(+Sab)	K0	27	1.39	0.40	5250	8.910	27
11-054	Haro 6-5 B	K5	59	9.96	...	4395	0.047	59
11-057	FS Tau AC	M0/M3.5	20	4.95/ 5.15	0.53	3876/3345	0.15/ 0.17/ 0.32	27, 20
21-044	LkCa 21	M3	27	0.73	0.21	3470	0.620	27
01-054	RX J0422.1+1934	...	...	...	...	...	...	...
01-062	2M J04221332+1934	M8	66	1.02	...	2713	0.017	19
11-079	CFHT-Tau 21	M1.25	18	6.60	...	3665	0.381	18
02-013	FV Tau AB	K5/K6	58	5.33/ 5.33	1.15	4395/4130	0.98/ 0.44/ 1.20	5, 58
02-000	FV Tau/c AB	M2.5/3.5	20	3.25/ 7.00	0.49	3412/3155	0.18/ 0.06/ 0.21	5, 20, 58
02-016	KPNO-Tau 13	M5	34	...	0.70	3125	0.150	34
02-000	DG Tau B	...	...	...	...	4000	5.500	10
02-022	DG Tau A	K6	5	1.41	0.36	4205	1.700	5, 58
02-000	KPNO-Tau 4	M9.5	5, 18	2.45	0.00	2500	0.004	5, 18
02-000	IRAS 04248+2612AB	M2-5.5	5, 59	7.02	1.51	2845	0.270	5, 59
15-020	JH 507	M4	5	0.76	0.29	3270	0.420	5, 27
13-004	GV Tau AB	K3-7	27, 59	12.10	...	4000	1.820	59
13-000	IRAS 04264+2433	M1	59	10.40	...	3605	0.140	59
15-040	DH Tau AB	M1	5	1.25	0.32	3705	0.560	5, 27
15-042	DI Tau AB	M0	5	0.76	0.43	3850	0.990	5, 27
15-044	KPNO-Tau 5	M7.5	5, 19	0.00	0.00	2783	0.023	19
14-006	IQ Tau A	M0.5	5	1.25	0.45	3778	0.880	5, 27
13-000	CFHT-Tau 20	M5.5	18	3.60	...	3065	0.138	18
14-000	KPNO-Tau 6	M9.0	18	0.88	0.00	2571	0.003	5, 18
13-035	FX Tau AB	M1	27	1.08	0.31	3720	1.020	27
14-057	DK Tau AB	K7	5	0.76	0.28	4060	1.300	5, 27
14-000	KPNO-Tau 7	M8.25	5	...	0.00	2632	0.003	5
22-013	MHO 9	M4.25	5	1.73	0.56	3234	0.220	5, 4
22-021	MHO 4	M7.1	5, 18	1.10	0.14	2880	0.048	5, 58
22-040	L1551 IRS5	...	...	...	...	4800	2.600	10
22-042	LkHa 358	M5.5	5	13.60	3.84	3058	0.590	5, 45
22-000	HH 30	M0	59	2.96	...	3800	0.006	59
22-043	HL Tau	K5	59	7.43	...	4395	1.530	59
22-047	XZ Tau AB	M2/M3.5	20	1.40/ 1.35	0.11	3561/3345	0.17/ 0.31/ 0.33	5, 20
22-056	L1551 NE	...	...	...	...	...	...	...
03-005	HK Tau AB	M0.5/M2	5	2.32	0.97	3778/3560	0.560	5, 27
22-070	V710 Tau BA	M0.5/M2	5	1.80/ 1.82	0.62	3778/3560	0.58/ 0.52/ 1.10	5, 58
19-009	JH 665	M5.5	5	0.97	0.54	3058	0.260	5, 27
22-089	L1551 51	K7	5	0.00	0.00	4060	0.470	5, 27
22-097	V827 Tau	K7	5	0.28	0.31	4060	1.100	5, 27
03-016	Haro 6-13	M0	59	11.90	...	3800	2.110	59

Table 9. continued.

XEST	Name	Spec <sup>a</sup>	Refs.	$A_V^a$ (mag)	$A_J$ (mag)	$T_{\text{eff}}^a$ (K)	$L_*^b$ ( $L_\odot$ )	Refs.
22-100	V826 Tau	K7	5	0.28	0.19	4060	0.940	5, 27
22-101	MHO 5	M6	5	0.23	0.17	2990	0.110	5, 58
03-017	CFHT-Tau 7	M5.75	66	0.00	...	3030	0.061	18, 19
03-019	V928 Tau AB	M0.5	5	1.87	1.04	3778	1.400	5, 27
03-022	FY Tau	K5	5	3.47	1.07	4350	1.000	5, 27
03-023	FZ Tau	M0	5	2.72	0.99	3850	0.980	5, 58
17-002	IRAS 04295+2251	...	...	...	...	3400	1.200	10
19-049	UZ Tau E+W(AB)	M1/2/3	27, 20	1.49/ 0.83	0.26	3705/3560	0.40/ 0.49/ 0.89	5, 27
17-009	JH 112	K6	27	3.23	0.93	4205	0.740	27
03-031	CFHT-Tau 5	M7.5	18	9.22	...	2783	0.075	18
04-003	CFHT-Tau 5	M7.5	18	9.22	...	2783	0.075	18
03-035	MHO 8	M6	5	0.19	0.28	2990	0.170	5, 4
04-009	MHO 8	M6	5	0.19	0.28	2990	0.170	5, 4
04-010	GH Tau AB	M1.5/M2	58	0.69/ 0.64	0.11	3631/3556	0.38/ 0.34/ 0.81	5, 58
04-012	V807 Tau SNab	K7/M3	58	0.36/ 0.36	0.04	3999/3388	1.07/ 0.32/ 2.10	5, 58
18-004	KPNO-Tau 14	M6	34	...	0.85	2990	0.110	34
17-000	CFHT-Tau 12	M6	66	3.44	...	2995	0.036	18, 19
04-016	V830 Tau	K7	5	0.28	0.13	4060	0.780	5, 27
18-000	IRAS S04301+261	M0	5	...	1.76	3850	0.025	5
17-000	IRAS 04302+2247	...	...	...	...	...	0.300	13
17-027	IRAS 04303+2240	M0.5	59	11.70	...	3700	2.200	59
04-034	GI Tau	K7	5	0.87	0.43	4060	1.000	5, 27
04-035	GK Tau AB	K7	5	0.87	0.46	4060	1.400	5, 27
18-019	IS Tau AB	K7/M4.5	58	3.67/ 3.67	0.95	3999/3221	0.50/ 0.19/ 0.66	5, 58
17-058	CI Tau	K7	27	1.77	0.51	4060	0.870	27
18-030	IT Tau AB	K2	5	4.09	1.86	4900	2.400	5, 27
17-066	JH 108	M1	27	1.46	0.42	3720	0.300	27
17-068	CFHT-BD Tau 1	M7.1	39, 19	3.10	...	2853	0.017	39, 19
25-026	AA Tau	K7	5	0.49	0.27	4060	0.800	5, 27
09-010	HO Tau AB	M0.5	5	1.11	0.46	3778	0.170	5, 27
08-019	FF Tau AB	K7	5	2.22	0.51	4060	0.690	5, 27
12-040	DN Tau	M0	5	0.49	0.34	3850	1.000	5, 27
12-000	IRAS 04325+2402AB	...	...	...	...	...	0.720	28
12-059	CoKu Tau 3 AB	M1	5	3.26	2.15	3705	0.980	5, 27
09-022	KPNO-Tau 8	M5.75	5	...	0.14	3024	0.021	5
08-037	HQ Tau AB	...	...	...	...	...	...	...
09-026	HQ Tau AB	...	...	...	...	...	...	...
08-043	KPNO-Tau 15	M2.75	34	...	0.56	3451	0.140	34
09-031	KPNO-Tau 15	M2.75	34	...	0.56	3451	0.140	34
08-000	KPNO-Tau 9	M8.5	5	...	0.00	2555	0.001	5
09-000	KPNO-Tau 9	M8.5	5	...	0.00	2555	0.001	5
08-048	HP Tau AB	K3	5	2.26	0.91	4730	1.400	5, 27
08-051a	HP Tau/G3 AB	K7	5	2.32	0.77	4060	0.710	5, 27
08-051	HP Tau/G2	G0	5	2.08	0.66	6030	6.500	5, 27
08-058	Haro 6-28 AB	M2/M3.5	20	2.30/ 1.90	1.34	3556/3342	0.09/ 0.03/ 0.12	5, 20
08-000	CFHT-BD Tau 2	M7.5	5	0.00	0.56	2795	0.011	5, 39
08-080	CFHT-BD Tau 3	M7.75	5	0.00	0.28	2747	0.007	5, 39
05-005	CFHT-Tau 6	M7.25	33, 18	0.41	0.07	2818	0.024	33, 18
05-000	IRAS 04361+2547	...	...	...	...	...	3.700	13
05-013	GN Tau AB	M2.5	33	...	1.17	3488	0.720	33
05-017	IRAS 04365+2535	...	...	...	...	...	2.200	27(Lb) <sup>c</sup>
05-024	IRAS 04369+2539	K4	59	18.10	...	4580	20.700	59
07-011	JH 223	M2	27	...	0.14	3560	0.180	33
07-022	Haro 6-32	M5	33	...	0.17	3125	0.120	33
07-000	ITG 33 A	M3	33	3.50	1.89	3415	0.051	33, 35
07-000	CFHT-Tau 8	M5.5	33	1.77	0.70	3058	0.024	33, 18, 19
07-000	IRAS 04381+2540	...	...	...	...	...	0.700	13
07-041	IRAS 04385+2550AB	M0	33	10.20	1.13	3850	0.180	33, 59
10-017	CoKuLk332/G2 AB	M0.5/2.5	20	2.00/ 3.30	0.99	3778/3485	0.24/ 0.15/ 1.10	33, 20

**Table 9.** continued.

XEST	Name	Spec <sup>a</sup>	Refs.	$A_V^a$ (mag)	$A_J$ (mag)	$T_{\text{eff}}^a$ (K)	$L_*^b$ ( $L_\odot$ )	Refs.
10-018	CoKuLk332/G1 AB	K7/M1	58	4.13/ 4.13	1.33	3707/3945	0.85/ 0.58/ 1.70	33, 58
10-020	V955 Tau AB	K5/M1	58	3.72/ 3.72	0.90	4395/3715	0.32/ 0.18/ 1.00	33, 58
10-034	CIDA 7	M4.75	66	...	...	...	0.050	27(Lb) <sup>c</sup>
10-045	DP Tau	M0.5	27	1.46	0.41	3778	0.200	33, 27
10-060	GO Tau	M0	27	1.18	0.77	3850	0.370	33, 27
26-012	2M J04552333+30	M6.25	33, 19	0.00	0.00	2959	0.015	33, 19
26-034	2M J04554046+30	M5.25	33	...	0.07	3091	0.021	33
26-043	AB Aur	B9.5-A0	27, 7	0.25	0.24	10050	49.000	33, 27, 8, 9
26-050	2MJ04554757/801	M4.75/5.6	33	...	0.00	3161/3044	0.10/ 0.01/ 0.11	33
26-067	SU Aur	G2	27	0.90	0.21	5860	9.900	33, 27
26-072	HBC 427	K7	27	0.00	0.17	4350	1.100	33, 27
Additional sources from Chandra								
C1-0	KPNO-Tau 10	M5	34	...	0.14	3125	0.052	34
C1-1	IRAS 04158+2805	M3	5	8.63	...	2760	0.050	59
C2-1	Haro 6-5 B	K5	59	9.96	...	4395	0.047	59
C2-2	FS Tau AC	M0/M3.5	20	4.95/ 5.15	0.53	3876/3345	0.15/ 0.17/ 0.32	27, 20
C3-1	FV Tau/c AB	M2.5/3.5	20	3.25/ 7.00	0.49	3412/3155	0.18/ 0.06/ 0.21	5, 20, 58
C3-2	DG Tau B	...	...	...	...	4000	5.500	10
C4-1	GV Tau AB	K3-7	27, 59	12.10	...	4000	1.820	59
C5-2	HN Tau AB	K5/M4	5	1.18/ 0.91	0.12	4395/3273	0.22/ 0.03/ 0.25	5, 58
C5-1	L1551 55	K7	5	0.69	0.20	4060	0.370	5, 27
C5-4	HD 28867	B9IVn	65	0.25	...	10500	64.90/62.10/127.00	65
C5-3	DM Tau	M1	5	0.59	0.31	3705	0.300	5, 58
C6-1	CFHT-BD Tau 4	M7	33	2.60	0.70	2853	0.062	33, 39, 19
C6-0	L1527 IRS	...	...	...	...	...	1.300	27(Lb) <sup>c</sup>
C6-0	CFHT-Tau 17	M5.5	66	6.50	...	3030	0.068	18
C6-2	IRAS 04370+2559	...	...	...	...	...	0.210	27(Lb) <sup>c</sup>

<sup>a</sup> For multiples, first number or spectral type refers to primary, second to secondary component.

<sup>b</sup> For multiples, three numbers give primary/secondary/total system luminosity.

<sup>c</sup> Referring to  $L_{\text{bol}}$  as derived from integration of the optical and infrared spectrum.

Notes on individual objects:

- V773 Tau = XEST-20-042: Multiple entries refer to A and C, respectively
- HD 28867 = C5-4: total  $L_*$  (Berghoefer et al. 1996) scaled to  $d = 140$  pc, split to components according to  $V$  mag (Walter et al. 2003).
- FQ Tau = XEST-23-067 = XEST-24-058, FS Tau = XEST-11-057, XZ Tau = XEST-22-047, CoKuLk332/G2 = XEST-10-017:  $T_{\text{eff}}$  of components read from Fig. 10 in Ref. 20.

**Table 10.** Fundamental parameters of targets in XEST (4): Age, mass, radius, rotation.

XEST	Name	Age <sup>a,b</sup> (Myr)	Mass <sup>a,c</sup> ( $M_{\odot}$ )	Refs.	Radius <sup>d</sup> ( $R_{\odot}$ )	$P$ (d)	Refs.	$v \sin i$ ( $\text{km s}^{-1}$ )	Refs.
27-115	HBC 352	...	1.05	2	0.79	<0.53	C	>75.00	48
27-000	HBC 353	...	...	...	0.71	<4.08	C	8.80	2
06-005	HBC 358 AB	3.26	0.41	27	1.38	...	C	<10.00	2
06-007	HBC 359	3.54	0.41	27	1.33	...	C	<10.00	48
06-059	L1489 IRS	0.80	1.45	10	3.65	<4.02	C	46.00	10
20-001	LkCa 1	0.87	0.27	5	1.93	<4.18	C	23.30	48
20-005	Anon 1	0.50	0.56	5	3.63	...	...	...	...
20-000	IRAS 04108+2803 A	...	...	...	...	...	...	...	...
20-022	IRAS 04108+2803 B	1.60	0.36	10	1.72	<6.23	C	14.00	10
20-000	2M J04141188+28	...	0.08	46	0.47	...	...	...	...
20-042	V773 Tau ABC	6.35/0.93	1.53/ 0.57	58	1.91	3.43	48	55.00	48
20-043	FM Tau	2.76	0.57	5	1.53	...	...	...	...
20-046	CW Tau	6.97	1.40	27	1.57	8.25	48	27.40	48
20-047	CIDA 1	...	...	...	...	...	...	5.30	48
20-056	MHO 2/1	1.54/4.85	0.36/ 0.34	5	1.78	<4.19	C	21.50	59
20-058	MHO 3	2.0	0.75	5	1.93	...	...	...	...
20-069	FO Tau AB	1.45/1.54	0.38/ 0.38	58	1.83	...	...	...	...
20-073	CIDA 2	0.39	0.20	5	2.02	...	...	...	...
23-002	CY Tau	1.53	0.42	5	1.79	7.50	48	10.00	48
24-002	CY Tau	1.53	0.42	5	1.79	7.50	48	10.00	48
23-004	LkCa 5	2.31	0.40	5	1.60	<2.19	C	37.00	48
24-004	LkCa 5	2.31	0.40	5	1.60	<2.19	C	37.00	48
23-008	CIDA 3	3.34	0.24	5	1.17	...	...	...	...
24-008	CIDA 3	3.34	0.24	5	1.17	...	...	...	...
23-015	V410 X3	2.68	0.12	5	1.08	<3.79	C	14.10	48
24-015	V410 X3	2.68	0.12	5	1.08	<3.79	C	14.10	48
23-018	V410 A13	...	0.10	58	0.72	<3.72	C	9.80	45
24-000	V410 A13	...	0.10	58	0.72	<3.72	C	9.80	45
23-000	V410 A24	18.7	1.22	5	1.58	...	...	...	...
24-000	V410 A24	18.7	1.22	5	1.58	...	...	...	...
23-029	V410 A25	0.64	0.46	5	3.08	...	...	...	...
24-027	V410 A25	0.64	0.46	5	3.08	...	...	...	...
23-032	V410 Tau ABC	2.74/5.22	1.51/ 0.12	58	2.31	1.94	48	71.00	48
24-028	V410 Tau ABC	2.74/5.22	1.51/ 0.12	58	2.31	1.94	48	71.00	48
23-033	DD Tau AB	4.47/4.47	0.30/ 0.30	58	1.03	...	...	...	...
24-029	DD Tau AB	4.47/4.47	0.30/ 0.30	58	1.03	...	...	...	...
23-035	CZ Tau AB	2.10	0.32	5	1.49	...	...	...	...
24-030	CZ Tau AB	2.10	0.32	5	1.49	...	...	...	...
23-036	IRAS 04154+2823	5.30	0.33	5	0.99	...	...	...	...
24-031	IRAS 04154+2823	5.30	0.33	5	0.99	...	...	...	...
23-037	V410 X2	0.44	0.56	5	3.90	...	...	...	...
24-032	V410 X2	0.44	0.56	5	3.90	...	...	...	...
23-045	V410 X4	0.06	0.27	5	3.56	...	...	...	...
24-038	V410 X4	0.06	0.27	5	3.56	...	...	...	...
23-047	V892 Tau	2.97	2.89	5	2.66	...	...	...	...
24-040	V892 Tau	2.97	2.89	5	2.66	...	...	...	...
23-048	LR 1	17.5	1.00	5	1.15	...	...	...	...
24-000	LR 1	17.5	1.00	5	1.15	...	...	...	...
23-050	V410 X7	1.90	0.50	5	1.69	...	...	...	...
24-042	V410 X7	1.90	0.50	5	1.69	...	...	...	...
23-000	V410 A20	20.7	1.01	5	1.06	...	...	...	...
24-000	V410 A20	20.7	1.01	5	1.06	...	...	...	...
23-056	Hubble 4	0.68	0.74	5	3.33	<13.17	C	12.80	48
24-047	Hubble 4	0.68	0.74	5	3.33	<13.17	C	12.80	48
23-000	KPNO-Tau 2	...	0.05	46	0.34	...	...	...	...
24-000	KPNO-Tau 2	...	0.05	46	0.34	...	...	...	...
23-000	CoKu Tau 1	29.1	0.70	59	0.81	<2.67	C	15.30	59
24-000	CoKu Tau 1	29.1	0.70	59	0.81	<2.67	C	15.30	59
23-061	V410 X6	0.71	0.18	5	1.60	...	...	...	...

Table 10. continued.

XEST	Name	Age <sup>a,b</sup> (Myr)	Mass <sup>a,c</sup> ( $M_{\odot}$ )	Refs.	Radius <sup>d</sup> ( $R_{\odot}$ )	$P$ (d)	Refs.	$v \sin i$ ( $\text{km s}^{-1}$ )	Refs.
24-054	V410 X6	0.71	0.18	5	1.60	...	...	...	...
23-063	V410 X5	4.21	0.14	5	1.03	...	...	...	...
24-055	V410 X5	4.21	0.14	5	1.03	...	...	...	...
23-067	FQ Tau AB	2.82/1.89	0.31/ 0.29	20	1.40	...	...	...	...
24-058	FQ Tau AB	2.82/1.89	0.31/ 0.29	20	1.40	...	...	...	...
28-100	BP Tau	1.91	0.75	27	1.97	7.60	48	7.80	48
23-074	V819 Tau AB	2.02	0.76	5	1.93	5.60	48	<15.00	48
24-061	V819 Tau AB	2.02	0.76	5	1.93	5.60	48	<15.00	48
16-000	IRAS 04166+2706	...	...	...	...	...	...	...	...
16-000	IRAS 04169+2702	...	...	...	...	...	...	...	...
11-000	CFHT-Tau 19	...	...	...	0.93	...	...	...	...
11-000	IRAS 04181+2655	0.90	0.68	10	2.80	<4.29	C	33.00	10
11-000	IRAS 04181+2654AB	...	...	...	...	...	...	...	...
11-023	2M J04213459	4.56	0.13	33	0.91	...	...	...	...
01-028	IRAS 04187+1927	...	...	...	...	...	...	...	...
11-037	CFHT-Tau 10	...	...	...	0.52	...	...	...	...
11-000	2M J04215450+2652	...	...	...	0.26	...	...	...	...
21-038	RY Tau	2.11	2.37	27	3.57	5.60	48	52.20	48
21-039	HD 283572	7.92	1.70	27	2.56	1.55	48	95.00	48
01-045	T Tau N(+Sab)	2.67	2.41	27	3.62	2.80	48	20.10	48
11-054	Haro 6-5 B	...	...	...	0.37	<0.91	C	20.70	59
11-057	FS Tau AC	17.2/3.06	0.61/ 0.28	20	0.93	...	...	...	...
21-044	LkCa 21	1.16	0.35	27	2.18	8.80	48	60.00	48
01-054	RX J0422.1+1934	...	...	...	...	...	...	...	...
01-062	2M J04221332+1934	...	...	...	0.59	...	...	...	...
11-079	CFHT-Tau 21	...	...	...	1.53	...	...	...	...
02-013	FV Tau AB	4.75/7.65	1.19/ 0.88	58	1.71	...	...	...	...
02-000	FV Tau/c AB	3.14/5.40	0.31/ 0.16	20, 58	1.22	...	...	...	...
02-016	KPNO-Tau 13	2.66	0.19	34	1.32	...	...	...	...
02-000	DG Tau B	0.30	0.72	10	4.90	...	...	...	...
02-022	DG Tau A	1.32	0.91	5	2.46	6.30	48	21.70	48
02-000	KPNO-Tau 4	...	0.01	46	0.35	...	...	...	...
02-000	IRAS 04248+2612AB	16.5	0.34	5	2.14	<6.61	C	16.40	59
15-020	JH 507	1.33	0.27	5	2.02	...	...	...	...
13-004	GV Tau AB	0.89	0.68	59	2.82	<5.63	C	25.30	59
13-000	IRAS 04264+2433	7.95	0.41	59	0.96	<1.52	C	32.00	10
15-040	DH Tau AB	1.53	0.47	5	1.82	7.00	48	10.00	48
15-042	DI Tau AB	1.07	0.56	5	2.24	7.70	48	10.50	48
15-044	KPNO-Tau 5	...	0.04	46	0.65	...	...	...	...
14-006	IQ Tau A	1.06	0.51	5	2.20	6.25	48	11.50	48
13-000	CFHT-Tau 20	...	...	...	1.31	...	...	...	...
14-000	KPNO-Tau 6	...	0.03	46	0.28	...	...	...	...
13-035	FX Tau AB	0.90	0.47	27	2.44	<12.34	C	10.00	48
14-057	DK Tau AB	1.32	0.74	5	2.31	8.40	48	11.40	48
14-000	KPNO-Tau 7	...	0.03	46	0.28	...	...	...	...
22-013	MHO 9	2.33	0.24	5	1.50	<7.66	C	9.90	48
22-021	MHO 4	...	0.10	58	0.88	<6.29	C	7.10	48
22-040	L1551 IRS5	4.80	1.58	10	2.34	<3.82	C	31.00	10
22-042	LkHa 358	0.12	0.21	5	2.74	<6.80	C	20.40	45
22-000	HH 30	...	0.52	59	0.19	...	...	<12.00	59
22-043	HL Tau	2.45	1.20	59	2.13	<7.21	C	15.00	48
22-047	XZ Tau AB	4.58/1.79	0.37/ 0.29	20	1.18	2.60	48	...	...
22-056	L1551 NE	...	...	...	...	...	...	...	...
03-005	HK Tau AB	1.76	0.51	5	1.75	<8.86	C	10.00	48
22-070	V710 Tau BA	1.69/1.69	0.51/ 0.40	5	1.78	<5.67	C	15.90	48
19-009	JH 665	0.60	0.19	5	1.82	...	...	...	...
22-089	L1551 51	5.42	0.80	5	1.39	2.43	48	27.00	48
22-097	V827 Tau	1.59	0.75	5	2.13	3.75	48	18.50	48
03-016	Haro 6-13	0.57	0.52	59	3.36	<7.30	C	23.30	59

Table 10. continued.

XEST	Name	Age <sup>a,b</sup> (Myr)	Mass <sup>a,c</sup> ( $M_{\odot}$ )	Refs.	Radius <sup>d</sup> ( $R_{\odot}$ )	$P$ (d)	Refs.	$v \sin i$ ( $\text{km s}^{-1}$ )	Refs.
22-100	V826 Tau	1.94	0.75	5	1.96	3.70	48	4.20	48
22-101	MHO 5	...	0.13	58	1.24	<7.84	C	8.00	48
03-017	CFHT-Tau 7	...	...	...	0.89	...	...	...	...
03-019	V928 Tau AB	0.73	0.50	5	2.77	<7.45	C	18.80	48
03-022	FY Tau	3.98	1.13	5	1.77	...	...	...	...
03-023	FZ Tau	1.08	0.56	5	2.23	...	...	...	...
17-002	IRAS 04295+2251	1.12	0.56	27	1.49	<1.48	C	51.00	10
19-049	UZ Tau E+W(AB)	2.26/1.43	0.47/ 0.39	5	1.54	<4.90	C	15.90	48
17-009	JH 112	4.06	0.95	27	1.62	...	...	...	...
03-031	CFHT-Tau 5	...	...	...	1.18	...	...	...	...
04-003	CFHT-Tau 5	...	...	...	1.18	...	...	...	...
03-035	MHO 8	0.47	0.15	5	1.54	<4.67	C	16.70	45
04-009	MHO 8	0.47	0.15	5	1.54	<4.67	C	16.70	45
04-010	GH Tau AB	2.04/2.03	0.42/ 0.38	58	1.56	<3.57	C	22.10	48
04-012	V807 Tau SNab	1.51/1.74	0.69/ 0.31	58	2.16	...	...	...	...
18-004	KPNO-Tau 14	1.09	0.13	34	1.24	...	...	...	...
17-000	CFHT-Tau 12	...	...	...	0.70	...	...	...	...
04-016	V830 Tau	2.48	0.76	5	1.79	2.75	48	29.10	48
18-000	IRAS S04301+261	...	...	...	0.36	...	...	...	...
17-000	IRAS 04302+2247	...	0.34	59	...	...	...	...	...
17-027	IRAS 04303+2240	0.50	0.51	59	3.62	<5.23	C	35.00	59
04-034	GI Tau	1.79	0.75	5	2.03	7.20	48	11.20	48
04-035	GK Tau AB	1.22	0.74	5	1.97	4.60	48	18.70	48
18-019	IS Tau AB	4.15/2.52	0.72/ 0.24	58	1.48	...	...	...	...
17-058	CI Tau	2.15	0.76	27	1.89	<9.20	C	10.40	48
18-030	IT Tau AB	4.75	1.67	5	2.16	...	...	...	...
17-066	JH 108	3.40	0.48	27	1.32	...	...	...	...
17-068	CFHT-BD Tau 1	...	...	...	0.53	...	...	...	...
25-026	AA Tau	2.40	0.76	5	1.81	8.22	48	11.40	48
09-010	HO Tau AB	9.13	0.52	5	0.96	...	...	...	...
08-019	FF Tau AB	2.95	0.77	5	1.68	...	...	...	...
12-040	DN Tau	1.05	0.56	5	2.25	6.30	48	8.10	48
12-000	IRAS 04325+2402AB	...	...	...	...	...	...	...	...
12-059	CoKu Tau 3 AB	0.92	0.46	5	2.41	...	...	...	...
09-022	KPNO-Tau 8	...	...	...	0.52	...	...	...	...
08-037	HQ Tau AB	...	...	...	...	...	...	...	...
09-026	HQ Tau AB	...	...	...	...	...	...	...	...
08-043	KPNO-Tau 15	3.86	0.32	34	1.05	...	...	...	...
09-031	KPNO-Tau 15	3.86	0.32	34	1.05	...	...	...	...
08-000	KPNO-Tau 9	...	...	...	0.19	...	...	...	...
09-000	KPNO-Tau 9	...	...	...	0.19	...	...	...	...
08-048	HP Tau AB	6.90	1.39	5	1.77	5.90	48	15.40	48
08-051a	HP Tau/G3 AB	2.83	0.77	5	1.71	...	...	...	...
08-051	HP Tau/G2	10.5	1.58	5	2.34	1.20	48	100.00	48
08-058	Haro 6-28 AB	10.0/18.5	0.35/ 0.21	20	0.79	<3.98	C	10.10	59
08-000	CFHT-BD Tau 2	...	...	...	0.45	...	...	...	...
08-080	CFHT-BD Tau 3	...	0.04	46	0.38	...	...	...	...
05-005	CFHT-Tau 6	...	...	...	0.65	...	...	...	...
05-000	IRAS 04361+2547	...	...	...	...	...	...	...	...
05-013	GN Tau AB	1.03	0.36	33	2.33	<11.80	C	10.00	48
05-017	IRAS 04365+2535	...	...	...	...	...	...	...	...
05-024	IRAS 04369+2539	...	...	...	7.24	...	...	<15.00	59
07-011	JH 223	4.27	0.37	33	1.12	...	...	...	...
07-022	Haro 6-32	3.33	0.09	33	1.18	...	...	...	...
07-000	ITG 33 A	12.9	0.26	33	0.65	...	...	...	...
07-000	CFHT-Tau 8	6.93	0.10	33	0.55	...	...	...	...
07-000	IRAS 04381+2540	...	...	...	...	...	...	...	...
07-041	IRAS 04385+2550AB	12.2	0.59	33	0.96	<2.13	C	22.80	59
10-017	CoKuLk332/G2 AB	5.60/4.50	0.52/ 0.34	20	0.99	<2.25	C	22.30	48

Table 10. continued.

XEST	Name	Age <sup>a,b</sup> (Myr)	Mass <sup>a,c</sup> ( $M_{\odot}$ )	Refs.	Radius <sup>d</sup> ( $R_{\odot}$ )	$P$ (d)	Refs.	$v \sin i$ ( $\text{km s}^{-1}$ )	Refs.
10-018	CoKuLk332/G1 AB	1.02/2.76	0.46/ 0.66	58	2.24	...	...	...	...
10-020	V955 Tau AB	24.1/6.70	0.90/ 0.47	58	0.98	<4.90	C	10.10	48
10-034	CIDA 7	...	...	...	...	...	...	...	...
10-045	DP Tau	7.24	0.52	33	1.05	<2.76	C	19.20	59
10-060	GO Tau	3.78	0.58	33	1.37	<3.96	C	17.50	48
26-012	2M J04552333+30	...	...	...	0.47	...	...	...	...
26-034	2M J04554046+30	9.39	0.11	33	0.51	...	...	...	...
26-043	AB Aur	4.00	2.70	9	2.31	<1.46	C	80.00	6
26-050	2MJ04554757/801	3.82	0.07	33	1.05	...	...	...	...
26-067	SU Aur	6.02	1.91	33	3.06	1.70	48	65.00	48
26-072	HBC 427	3.43	1.13	33	1.85	9.30	48	10.40	2
Additional sources from Chandra									
C1-0	KPNO-Tau 10	5.7	0.14	34	0.78	...	...	...	...
C1-1	IRAS 04158+2805	1.55	0.35	27	0.98	<2.15	C	23.10	59
C2-1	Haro 6-5 B	...	...	...	0.37	<0.91	C	20.70	59
C2-2	FS Tau AC	17.2/3.06	0.61/ 0.28	20	0.93	...	...	...	...
C3-1	FV Tau/c AB	3.14/5.40	0.31/ 0.16	20, 58	1.22	...	...	...	...
C3-2	DG Tau B	0.30	0.72	10	4.90	...	...	...	...
C4-1	GV Tau AB	0.89	0.68	59	2.82	<5.63	C	25.30	59
C5-2	HN Tau AB	41/14	0.78/ 0.07	58	0.81	<0.78	C	52.80	48
C5-1	L1551 55	8.2	0.81	5	1.23	6.20	48	<10.00	48
C5-4	HD 28867	3.71/3.76	2.85/ 2.82	65	2.60	<2.00	65	65.00	65
C5-3	DM Tau	3.26	0.47	5	1.33	<6.74	C	10.00	48
C6-1	CFHT-BD Tau 4	...	0.06	46	1.02	...	...	...	...
C6-0	L1527 IRS	...	...	...	...	...	...	...	...
C6-0	CFHT-Tau 17	...	...	...	0.94	...	...	...	...
C6-2	IRAS 04370+2559	...	...	...	...	...	...	...	...

<sup>a</sup> For binaries, first number refers to primary, second to secondary component (calculated from  $L_*$  and  $T_{\text{eff}}$ ).

<sup>b</sup> Ages derived after Siess et al. (2000) using the same principal parameters as for masses, quoted in Table 9).

<sup>c</sup> Masses derived after Siess et al. (2000) using principal parameters quoted in Table 9.

Exceptions quoted directly from literature: HBC 352 (XEST-27-115; ref. 2), 2M J04141188+28 (XEST-20-000; ref. 46), V410 A13 (XEST23-018 = XEST-24-000; ref. 58), KPNO-Tau 2 (XEST-23-000 = XEST-24-000; ref. 46), KPNO-Tau 4 (XEST-02-000; ref. 46), KPNO-Tau 5 (XEST-15-044; ref. 46), KPNO-Tau 6 (XEST-14-000; ref. 46), KPNO-Tau 7 (XEST-14-000; ref. 46), MHO 4 (XEST-22-021; ref. 58), HH 30 (XEST-22-000; ref. 59), IRAS 04302+2247 (XEST-17-000; ref. 59), CFHT-BD Tau 3 (XEST-08-080; ref. 46), CFHT-BD Tau 4 (C6-1; ref. 46).

<sup>d</sup> For multiples, radius is given only for primary if luminosity of primary is explicitly known.

Notes on individual objects:

- Mass/age calculations for references different from those of  $T_{\text{eff}}$  and  $L_*$  for CW Tau (XEST-20-046; ref. 27 instead of 5), and IRAS 04158+2805 (C1-1; ref. 27 instead of 59).
- V773 Tau = XEST-20-042: Multiple entries refer to A and C, respectively.
- V410 Tau = XEST-23-032 = XEST-24-028: Multiple entries refer to A and C, respectively.
- HD 28867 = C5-4: radius and  $P$  of G-type companion after Walter et al. (2003).



**Table 11.** Fundamental parameters of targets in XEST (5); Accretion and evolution.

XEST	Name	$\dot{M}$ (min/max) <sup>a</sup> ( $M_{\odot}\text{yr}^{-1}$ )	Refs.	$EW(\text{H}\alpha)$ <sup>b</sup> ( $\text{\AA}$ )	TTS type	Refs. <sup>c</sup>	IR <sup>d</sup> class	Refs. <sup>d</sup>	Type
27-115	HBC 352	...	...	0	W	29	III	1, 27	3
27-000	HBC 353	...	...	0	W	29	III	27	3
06-005	HBC 358 AB	<8.97	20	4–10	W/W	37, 29	III	1, 27	3
06-007	HBC 359	...	...	2–9	W	56, 37	III	27	3
06-059	L1489 IRS	-7.15	59	41–56	C*	59, 29	I	1, 59, 27	1
20-001	LkCa 1	<-9.72	58	3–4	W	29, 45	III	1, 27	3
20-005	Anon 1	<-8.94	58	1–3	W	37, 29	III	1, 27	3
20-000	IRAS 04108+2803 A	...	...	37	C	29	II	27	2
20-022	IRAS 04108+2803 B	...	...	...	...	...	I	59, 27	1
20-000	2M J04141188+28	-10.00	46	250	C	46	...	...	4
20-042	V773 Tau ABC	<-10.00	58	2–4	W	21, 24	II	1, 27	3
20-043	FM Tau	-8.87/ -8.45	58, 44	51–101	C	21, 29	II	1, 27	2
20-046	CW Tau	-7.99	58	135–140	C	7, 29	II	1, 59, 27	2
20-047	CIDA 1	-8.50	57	112–149	C	41, 29	II	27	2
20-056	MHO 2/1	<-8.48	59	58–88	C/C	59, 4	I;II	59; 22	2
20-058	MHO 3	...	...	16–21	C	4	II	22	2
20-069	FO Tau AB	-7.90/ -7.58	58, 20	116–137	C	29, 20	II	1, 27	2
20-073	CIDA 2	...	...	5–7	W	3, 29	III	1, 27	3
23-002	CY Tau	-8.86/ -8.12	58	55–70	C	58, 7	II	1, 27	2
24-002	CY Tau	-8.86/ -8.12	58, 44	55–70	C	58, 7	II	1, 27	2
23-004	LkCa 5	<-10.00	58	4	W	29	III	1, 27	3
24-004	LkCa 5	<-10.00	58	4	W	29	III	1, 27	3
23-008	CIDA 3	...	...	6	W	3	II	1, 27	3
24-008	CIDA 3	...	...	6	W	3	II	1, 27	3
23-015	V410 X3	<-9.30	57	14–30	W?	41, 57	...	...	3
24-015	V410 X3	<-9.30	57	14–30	W?	41, 57	...	...	3
23-018	V410 A13	-11.30	45	27–41	C?	45, 58	II	22	2
24-000	V410 A13	-11.30	45	27–41	C?	45, 58	II	22	2
23-000	V410 A24	...	...	...	...	...	...	...	9
24-000	V410 A24	...	...	...	...	...	...	...	9
23-029	V410 A25	...	...	...	...	...	...	...	9
24-027	V410 A25	...	...	...	...	...	...	...	9
23-032	V410 Tau ABC	<-8.80	58	2–3	W	29, 7	III	1, 27	3
24-028	V410 Tau ABC	<-8.80	58	2–3	W	29, 7	III	1, 27	3
23-033	DD Tau AB	-9.10/ -7.21	58, 20	90–206	C	37, 20	II	1, 27	2
24-029	DD Tau AB	-9.10/ -7.21	58, 20	90–206	C	37, 20	II	1, 27	2
23-035	CZ Tau AB	...	...	4–7	W	7, 29	II	1, 27	3
24-030	CZ Tau AB	...	...	4–7	W	7, 29	II	1, 27	3
23-036	IRAS 04154+2823	...	...	>17	C	59	FS;II	1; 59, 27	2
24-031	IRAS 04154+2823	...	...	>17	C	59	FS;II	1; 59, 27	2
23-037	V410 X2	...	...	...	...	...	...	...	9
24-032	V410 X2	...	...	...	...	...	...	...	9
23-045	V410 X4	...	...	...	...	...	...	...	9
24-038	V410 X4	...	...	...	...	...	...	...	9
23-047	V892 Tau	...	...	7–13	Ae	29, 7	II	1, 27	5
24-040	V892 Tau	...	...	7–13	Ae	29, 7	II	1, 27	5
23-048	LR 1	...	...	...	...	...	...	...	9
24-000	LR 1	...	...	...	...	...	...	...	9
23-050	V410 X7	...	...	3	W	58	III	22	3
24-042	V410 X7	...	...	3	W	58	III	22	3
23-000	V410 A20	...	...	...	...	...	...	...	9
24-000	V410 A20	...	...	...	...	...	...	...	9
23-056	Hubble 4	<-8.16	58	3–4	W	7, 29	III	1, 27	3
24-047	Hubble 4	<-8.16	58	3–4	W	7, 29	III	1, 27	3
23-000	KPNO-Tau 2	<-12.00	46	4–12	W	46, 5	...	...	4
24-000	KPNO-Tau 2	<-12.00	46	4–12	W	46, 5	...	...	4
23-000	CoKu Tau 1	-7.36	59	70–111	C	59, 29	II	59, 1, 27	2
24-000	CoKu Tau 1	-7.36	59	70–111	C	59, 29	II	59, 1, 27	2
23-061	V410 X6	...	...	13	W	38	...	...	3

Table 11. continued.

XEST	Name	$\dot{M}$ (min/max) <sup>a</sup> ( $M_{\odot}\text{yr}^{-1}$ )	Refs.	$EW(\text{H}\alpha)^b$ ( $\text{\AA}$ )	TTS type	Refs. <sup>c</sup>	IR <sup>d</sup> class	Refs. <sup>d</sup>	Type
24-054	V410 X6	...	...	13	W	38	...	...	3
23-063	V410 X5	...	...	11–19	W?	58, 45	II	22	3
24-055	V410 X5	...	...	11–19	W?	58, 45	II	22	3
23-067	FQ Tau AB	-7.98/-6.45	20, 44	81–114	C/C	29, 7	II	1, 27	2
24-058	FQ Tau AB	-7.98/-6.45	20, 44	81–114	C/C	29, 7	II	1, 27	2
28-100	BP Tau	-7.88/-7.54	58, 44	40–92	C	55, 29	II	1, 27	2
23-074	V819 Tau AB	<-8.86	58	2–3	W	60, 29	III	1, 27	3
24-061	V819 Tau AB	<-8.86	58	2–3	W	60, 29	III	1, 27	3
16-000	IRAS 04166+2706	...	...	...	...	...	I	59, 1, 27	1
16-000	IRAS 04169+2702	...	...	...	...	...	I	59, 1, 27	1
11-000	CFHT-Tau 19	...	...	442	C	18	...	...	2
11-000	IRAS 04181+2655	...	...	...	...	...	I	59, 27	1
11-000	IRAS 04181+2654AB	...	...	...	...	...	I	59, 27	1
11-023	2M J04213459	...	...	...	W	33	...	...	3
01-028	IRAS 04187+1927	...	...	...	...	...	II	27	2
11-037	CFHT-Tau 10	...	...	17	W	18	...	...	3
11-000	2M J04215450+2652	...	...	...	...	...	...	...	4
21-038	RY Tau	-7.19/-7.04	63	13	C	29	II	1, 27	2
21-039	HD 283572	...	...	0	W	29	III	1, 27	3
01-045	T Tau N(+Sab)	-7.50/-7.24	58, 63	41	C	29	II	59, 1, 27	2
11-054	Haro 6-5 B	-6.76	59	91	C*	59	(II)	(59)	1
11-057	FS Tau AC	-9.50/-8.09	58, 44	57–81	C/C	7, 58	FS;II	1; 27	2
21-044	LkCa 21	...	...	6	W	37	III	1, 27	3
01-054	RX J0422.1+1934	...	...	19	W	36	...	...	3
01-062	2M J04221332+1934	...	...	...	...	...	...	...	4
11-079	CFHT-Tau 21	...	...	45	C	18	...	...	2
02-013	FV Tau AB	-7.70/-6.23	58, 44	6–23	C/C	58, 7	II	23, 1, 27	2
02-000	FV Tau/c AB	<-8.70/<-8.60	20, 58	17–29	C/C	58, 21	II	23, 1, 27	2
02-016	KPNO-Tau 13	...	...	8–10	W	41, 34	...	...	3
02-000	DG Tau B	...	...	>276	C*	59	I/II;II	23; 59	1
02-022	DG Tau A	-7.34/-6.13	58, 59	63–125	C*	59, 29	I/II;II	23;59, 27	2
02-000	KPNO-Tau 4	<-12.00	46	38–158	C	46, 18	III	23	4
02-000	IRAS 04248+2612AB	-8.97	59	163	C*	59	I	59, 1, 27	1
15-020	JH 507	...	...	...	...	...	III	23, 1, 27	3
13-004	GV Tau AB	-6.71	59	86	C*	59	I	59, 1, 27	1
13-000	IRAS 04264+2433	-7.11	59	69–96	C*	29, 59	I	59, 27	1
15-040	DH Tau AB	-8.95/-8.30	58, 44	39–72	C	29, 58	II	23, 1, 27	2
15-042	DI Tau AB	...	...	1–2	W	14, 29	III;II	23; 27	3
15-044	KPNO-Tau 5	<-12.00	46	4–30	W	46, 5	III	23	4
14-006	IQ Tau A	<-8.32/-7.55	58, 44	8–17	C	29, 21	II	27	2
13-000	CFHT-Tau 20	...	...	114	C	18	...	...	2
14-000	KPNO-Tau 6	-11.40	46	41–350	C	46, 5	II	23	4
13-035	FX Tau AB	-8.65	44	10–15	C/W	7, 29	II	1, 27	2
14-057	DK Tau AB	-7.42	44	31–50	C/C	26, 29	II	1, 27	2
14-000	KPNO-Tau 7	-11.40	46	31–300	C	46, 5	II	23	4
22-013	MHO 9	<-9.70	57	3–6	W	41, 4	...	...	3
22-021	MHO 4	<-9.50	57	28–43	W?	57, 45	...	...	4
22-040	L1551 IRS5	...	...	83–412	C*	29, 59	I	59, 1, 27	1
22-042	LkHa 358	-8.50	45	47–87	C	7, 29	II	27	2
22-000	HH 30	-6.45	59	185–199	C	29, 59	II	59	2
22-043	HL Tau	-8.83/-6.80	58, 59	43–55	C*	59, 7	I;II	1,59; 27	1
22-047	XZ Tau AB	-8.90/-7.00	58, 20	62-274	C/C	58, 7	II	59, 27	2
22-056	L1551 NE	...	...	...	...	...	I	59, 1, 27	1
03-005	HK Tau AB	-7.65	59	29–54	C/C	7, 29	FS;I/II	1; 27, 59	2
22-070	V710 Tau BA	...	...	34–89	C/W	37, 26	II+II	1, 27	2
19-009	JH 665	...	...	...	...	...	III	27	3
22-089	L1551 51	<-9.50	58	1–2	W	37, 29	III	1, 27	3
22-097	V827 Tau	<-8.53	58	2–4	W	7, 29	III	1, 27	3
03-016	Haro 6-13	-7.54	59	34–88	C	59, 7	FS;I/II	1; 27, 59	2

Table 11. continued.

XEST	Name	$\dot{M}$ (min/max) <sup>a</sup> ( $M_{\odot}\text{yr}^{-1}$ )	Refs.	$EW(H\alpha)^b$ ( $\text{\AA}$ )	TTS type	Refs. <sup>c</sup>	IR <sup>d</sup> class	Refs. <sup>d</sup>	Type
22-100	V826 Tau	...	...	2–4	W	7, 29	III	1, 27	3
22-101	MHO 5	-10.80/<-9.70	45, 57	36–60	C?	57, 45	...	...	2
03-017	CFHT-Tau 7	...	...	9	W	18	...	...	3
03-019	V928 Tau AB	...	...	1–2	W	37, 29	III	1, 27	3
03-022	FY Tau	-7.48	58	48–73	C	58, 29	II	1, 27	2
03-023	FZ Tau	-7.70	58	181–204	C	29, 7	II	1, 27	2
17-002	IRAS 04295+2251	...	...	>11–66	C*	59, 29	I;FS	59, 27; 1	1
19-049	UZ Tau E+W(AB)	-8.70/-8.01	58, 20	65–82	C/C/C	29, 7	II+II	23	2
17-009	JH 112	...	...	16	C	29	II	1, 27	2
03-031	CFHT-Tau 5	...	...	30	W	18	...	...	4
04-003	CFHT-Tau 5	...	...	30	W	18	...	...	4
03-035	MHO 8	...	...	14–21	W	45, 4	II	22	3
04-009	MHO 8	...	...	14–21	W	45, 4	II	22	3
04-010	GH Tau AB	-8.90/-7.92	58, 44	10–31	C/C	20, 58	II	1, 27	2
04-012	V807 Tau SNab	<-8.68/-8.40	20, 58	5–16	C/W	21, 37	III;II	1; 27	2
18-004	KPNO-Tau 14	<-12.00	46	14–40	W?	46, 34	...	...	3
17-000	CFHT-Tau 12	...	...	80	C	18	...	...	2
04-016	V830 Tau	<-8.97	58	2–3	W	29, 60	III	1, 27	3
18-000	IRAS S04301+261	...	...	...	...	...	II	23, 27	2
17-000	IRAS 04302+2247	...	...	67	C*	59	I	59, 1, 27	1
17-027	IRAS 04303+2240	-6.63/-6.05	59	67–137	C	59	II	59, 27	2
04-034	GI Tau	-8.08	58, 44	15–21	C	29, 21	II	27	2
04-035	GK Tau AB	-8.19	58	15–35	C/C	21, 11	II	59, 1, 27	2
18-019	IS Tau AB	-8.10/-7.91	58, 20	10–26	C/C	20, 58	II	1, 23, 27	2
17-058	CI Tau	-7.59/-7.19	58, 44	77–102	C	29, 7	II	1, 27	2
18-030	IT Tau AB	...	...	10–22	C/C	29, 11	II	1, 23, 27	2
17-066	JH 108	...	...	3	W	29	III	1, 27	3
17-068	CFHT-BD Tau 1	...	...	7–19	W	41, 39	...	...	4
25-026	AA Tau	-8.48/-8.19	44, 58	37–46	C	7, 29	II	1, 27	2
09-010	HO Tau AB	-8.87	58	102–115	C	29, 7	II	1, 27	2
08-019	FF Tau AB	...	...	1–3	W	24, 29	III	1, 27	3
12-040	DN Tau	-8.73/-7.79	58, 59	12–87	C	7, 14	II	59, 1, 27	2
12-000	IRAS 04325+2402AB	...	...	...	...	...	I	59, 1, 27	1
12-059	CoKu Tau 3 AB	...	...	5	W	7	II	1, 27	3
09-022	KPNO-Tau 8	...	...	15–18	W	41, 5	...	...	3
08-037	HQ Tau AB	...	...	...	...	...	III	1	3
09-026	HQ Tau AB	...	...	...	...	...	III	1	3
08-043	KPNO-Tau 15	...	...	6	W	34	...	...	3
09-031	KPNO-Tau 15	...	...	6	W	34	...	...	3
08-000	KPNO-Tau 9	...	...	1	W	41	...	...	4
09-000	KPNO-Tau 9	...	...	1	W	41	...	...	4
08-048	HP Tau AB	...	...	20–35	C	29, 7	FS;II	1; 27	2
08-051a	HP Tau/G3 AB	...	...	1–2	W	21, 29	III	27	3
08-051	HP Tau/G2	...	...	0–5	W	21, 7	III	27	3
08-058	Haro 6-28 AB	-8.70/-8.10	45, 20	48–92	C/C	45, 7	FS;I	1; 59, 27	2
08-000	CFHT-BD Tau 2	...	...	7–13	W	41, 39	...	...	4
08-080	CFHT-BD Tau 3	<-12.00	46	11–55	W?	46, 39	...	...	4
05-005	CFHT-Tau 6	-11.30	46	64–102	C	18, 46	...	...	4
05-000	IRAS 04361+2547	...	...	...	...	...	I	59, 1, 27	1
05-013	GN Tau AB	-7.90	57	59–62	C	57, 3	II	1, 23, 27	2
05-017	IRAS 04365+2535	...	...	...	...	...	I	1, 23, 59, 27	1
05-024	IRAS 04369+2539	-6.20	59	22	C	59	II	1, 59, 27	2
07-011	JH 223	...	...	4	W	29	II	1, 23, 27	3
07-022	Haro 6-32	...	...	...	W	33	...	...	3
07-000	ITG 33 A	...	...	53	C	35	...	...	2
07-000	CFHT-Tau 8	...	...	52	C	18, 33	...	...	2
07-000	IRAS 04381+2540	...	...	...	...	...	I	23, 1, 59, 27	1
07-041	IRAS 04385+2550AB	-8.11	59	15–20	C	59, 29	II;I	23, 27; 59	2
10-017	CoKuLk332/G2 AB	<-8.58	20	2–3	W/W	20, 7	III	1, 23, 27	3

Table 11. continued.

XEST	Name	$\dot{M}$ (min/max) <sup>a</sup> ( $M_{\odot}\text{yr}^{-1}$ )	Refs.	$EW(H\alpha)^b$ ( $\text{\AA}$ )	TTS type	Refs. <sup>c</sup>	IR <sup>d</sup> class	Refs. <sup>d</sup>	Type
10-018	CoKuLk332/G1 AB	<8.20	58	0–5	W/W	58, 29	III;II	23, 1; 27	3
10-020	V955 Tau AB	<-8.97/ -8.50	20, 58	11–45	C/W	20, 58	II	23, 1, 27	2
10-034	CIDA 7	...	...	79	C	3	II	23, 1	2
10-045	DP Tau	-8.50/ -6.92	58, 59	74–102	C	59, 29	II	23, 1, 59, 27	2
10-060	GO Tau	-8.42	58	78–81	C	29, 7	II	23, 1, 27	2
26-012	2M J04552333+30	...	...	...	W	33	...	...	4
26-034	2M J04554046+30	...	...	...	W	33	...	...	3
26-043	AB Aur	...	...	22–44	Ae	15, 29	II	1, 27	5
26-050	2MJ04554757/801	...	...	25	C	46	...	...	2
26-067	SU Aur	-8.30/ -8.20	63	2–6	C	29, 14	II	1, 27	2
26-072	HBC 427	...	...	1	W	29	III	23, 1, 27	3
Additional sources from Chandra									
C1-0	KPNO-Tau 10	...	...	36	C	34	...	...	2
C1-1	IRAS 04158+2805	<-9.50	59	...	...	...	I;II	59; 27	2
C2-1	Haro 6-5 B	-6.76	59	91	C*	59	II	59	1
C2-2	FS Tau AC	-9.50/ -8.28	58, 20	57–81	C/C	7, 58	FS;II	1; 27	2
C3-1	FV Tau/c AB	<-8.70/ <-8.60	20, 58	17–21	C/C	58, 20	II	23, 1, 27	2
C3-2	DG Tau B	...	...	>276	C*	59	I/II;II	23; 59	1
C4-1	GV Tau AB	-6.71	59	86	C*	59	I	59, 1, 27	1
C5-2	HN Tau AB	...	...	138–163	C	21, 58	II	1, 27	2
C5-1	L1551 55	<-9.70	58	1	W	58	III	1, 27	3
C5-4	HD 28867	...	...	...	...	...	...	...	9
C5-3	DM Tau	-8.67	58	...	...	...	II	1, 27	2
C6-1	CFHT-BD Tau 4	-11.30	46	129–340	C	46, 39	...	...	4
C6-0	L1527 IRS	...	...	...	...	...	0	59, 23	0
C6-0	CFHT-Tau 17	...	...	7	W	18	...	...	3
C6-2	IRAS 04370+2559	...	...	...	...	...	II	27	2

<sup>a</sup> Range of  $\dot{M}$  reported in literature given. For multiple systems, numbers refer to primary or integrated system.<sup>b</sup> Range of  $EW$  reported in literature given. For multiple systems, numbers refer to primary or integrated system.<sup>c</sup> For  $EW$  range, first reference for minimum, second for maximum reported.<sup>d</sup> Infrared classification; double entries: '/' for transition objects, '+' for components, ';' for different types, ',' for different references. FS = flat-spectrum source.

## Appendix A: The XEST source catalog

The complete catalog of all X-ray sources detected in the XEST fields (not including the complementary fields from *Chandra*) is available in electronic form. The data reduction and source identification procedures are described in Sect. 4. The catalog also provides errors of the source counts and the count rates (not given in Table 4 for the “Scts” and “Rate” columns). The numbering scheme is as defined in Sect. 5, all TMC members being included. The catalog is sorted in right ascension and contains a total of 2347 identified X-ray sources. If two XEST fields overlap, the same sources may have been identified twice, with different XEST IDs assigned. The first ten entries of the catalog are given in the Table A.1 below for illustration. Note that if  $ML < 5$  in any band the number of counts and the count rate are 95% upper limits in that band. The HR is the upper (or lower) limit as appropriate in these situations with the low and high values set to  $-1$  and to the HR upper limit, respectively (or to the HR lower limit and to  $+1$ , respectively).

A description of the columns follows:

- 1 XEST ID (field-srcnr).
- 2 RA (J2000, X-ray, corrected for boresight shift).
- 3 Dec (J2000, X-ray, corrected for boresight shift).
- 4 RA as above, in decimal degrees.
- 5 Dec as above, in decimal degrees.
- 6  $1\sigma$  positional error (including rms of boresight shift).
- 7 Maximum likelihood of detection in the mosaicked EPIC full band (0.5–7.3 keV) image.
- 8 Number of detected counts in the mosaicked EPIC full band image. If  $ML_F < 5$ , then 95% upper limit is given.
- 9 Total PN-equivalent effective exposure time in s at source position in in the mosaicked EPIC full band image.

- 10 PN-equivalent source count-rate in the mosaicked EPIC full band image. If  $ML_F < 5$ , then 95% upper limit is given.
- 11  $1\sigma$  statistical error in PN-equivalent source count-rate in the mosaicked EPIC full band image (note: does not include uncertainty in spectrally-dependent PN/MOS sensitivity ratio).
- 12 Maximum likelihood of detection in the mosaicked EPIC soft band (0.5–2 keV) image.
- 13 Number of detected counts in the mosaicked EPIC soft band image. If  $ML_S < 5$ , then 95% upper limit is given.
- 14 Total PN-equivalent effective exposure time in s at source position in in the mosaicked EPIC soft band image.
- 15 PN-equivalent source count-rate in the mosaicked EPIC soft band image. If  $ML_S < 5$ , then 95% upper limit is given.
- 16  $1\sigma$  statistical error in PN-equivalent source count-rate in the mosaicked EPIC soft band image (note: does not include uncertainty in spectrally-dependent PN/MOS sensitivity ratio).
- 17 Maximum likelihood of detection in the mosaicked EPIC hard band (2–7.3 keV) image.
- 18 Number of detected counts in the mosaicked EPIC hard band image. If  $ML_H < 5$ , then 95% upper limit is given.
- 19 Total PN-equivalent effective exposure time in s at source position in in the mosaicked EPIC hard band image.
- 20 PN-equivalent source count-rate in the mosaicked EPIC hard band image. If  $ML_H < 5$ , then 95% upper limit is given.
- 21  $1\sigma$  statistical error in PN-equivalent source count-rate in the mosaicked EPIC hard band image (note: does not include uncertainty in spectrally-dependent PN/MOS sensitivity ratio).
- 22 Hardness ratio  $(Rate_H - Rate_S)/Rate_F$ . If object is not detected in soft (hard) band, then HR is upper (lower) limit. If object is detected only in full band, then  $HR = 0$ .
- 23  $1\sigma$  lower limit to hardness ratio.
- 24  $1\sigma$  upper limit to hardness ratio.
- 25 Comment concerning by-eye verification of source.

**Table A.1.** XEST Catalog (first ten entries).


XEST	RA <sub>X</sub>	Dec <sub>X</sub>	RA	Dec	Poserr	ML <sub>F</sub>	N <sub>F</sub>	T <sub>exp,F</sub>	Rate <sub>F</sub>	Err <sub>F</sub>
1	2	3	4	5	6	7	8	9	10	11
27-001	03 53 03.92	+31 53 02.8	58.266348	+31.884122	1.17	337.9	250	10386	0.0241	0.0017
27-002	03 53 05.45	+31 54 19.8	58.272721	+31.905495	1.47	20.0	41	11432	0.0037	0.0008
27-003	03 53 12.71	+31 55 12.7	58.302955	+31.920193	1.28	51.7	102	27319	0.0038	0.0005
27-004	03 53 15.59	+31 53 01.3	58.314976	+31.883708	1.20	138.4	180	29916	0.0060	0.0006
27-005	03 53 19.30	+31 47 39.1	58.330416	+31.794186	2.01	11.1	42	28822	0.0015	0.0004
27-006	03 53 19.73	+31 59 21.4	58.332216	+31.989273	1.95	9.2	38	25935	0.0015	0.0004
27-007	03 53 22.62	+31 56 24.9	58.344257	+31.940258	1.66	11.1	38	31200	0.0012	0.0003
27-008	03 53 24.35	+31 43 20.5	58.351468	+31.722347	1.83	8.0	24	10810	0.0022	0.0007
27-009	03 53 25.01	+31 42 13.4	58.354220	+31.703724	1.60	8.8	24	9042	0.0027	0.0008
27-010	03 53 25.25	+32 03 26.5	58.355189	+32.057353	1.21	80.4	129	10663	0.0121	0.0014


ML <sub>S</sub>	N <sub>S</sub>	T <sub>exp,S</sub>	Rate <sub>S</sub>	Err <sub>S</sub>	ML <sub>H</sub>	N <sub>H</sub>	T <sub>exp,H</sub>	Rate <sub>H</sub>	Err <sub>H</sub>	HR	HR <sub>low</sub>	HR <sub>high</sub>	Verif.
12	13	14	15	16	17	18	19	20	21	22	23	24	25
75.5	69	10471	0.0066	0.0009	271.6	181	10440	0.0174	0.0015	+0.448	+0.370	+0.526	
17.8	29	11517	0.0026	0.0006	3.0	21	11503	0.0019	0.0000	-0.160	-1.000	-0.160	
49.9	73	28589	0.0026	0.0004	9.1	32	27099	0.0012	0.0003	-0.372	-0.519	-0.224	
106.7	115	31188	0.0037	0.0004	34.4	62	29754	0.0021	0.0004	-0.265	-0.361	-0.169	
4.1	41	30079	0.0014	0.0000	6.4	24	28636	0.0009	0.0003	-0.226	-0.226	+1.000	
8.4	27	27223	0.0010	0.0003	1.7	34	25697	0.0013	0.0000	+0.142	-1.000	+0.142	
3.5	34	32427	0.0011	0.0000	5.8	22	31072	0.0007	0.0003	-0.178	-0.178	+1.000	
0.2	11	10896	0.0011	0.0000	10.5	21	10894	0.0020	0.0005	+0.297	+0.297	+1.000	
11.7	19	9131	0.0021	0.0006	0.3	19	9101	0.0022	0.0000	+0.006	-1.000	+0.006	
112.0	132	11298	0.0117	0.0012	0.1	11	10463	0.0011	0.0000	-0.825	-1.000	-0.825	

## Appendix B: The XEST atlas

Figures B.1–B.10 show all 28 XEST EPIC exposures. Each field of view is presented in two versions. The left panel shows a co-added EPIC image, logarithmically compressed in intensity and slightly smoothed. The colors code for hardness, where hardness increases from red to yellow to green to blue. Saturated (bright) stellar images are black. The right panel shows an unsmoothed, co-added EPIC image with an RA(J2000.0) – Dec(J2000.0) coordinate grid. Also plotted are the locations of known TMC members (red circles), labeled with their XEST source IDs (srcnr). The latter are located directly above the source circles except in crowded areas where labels may have been shifted.




**Fig. B.1.** Co-added EPIC images of field XEST-01, XEST-02, and XEST-03 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.




**Fig. B.2.** Co-added EPIC images of field XEST-04, XEST-05, and XEST-06 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.







**Fig. B.3.** Co-added EPIC images of field XEST-07, XEST-08, and XEST-09 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.




**Fig. B.4.** Co-added EPIC images of field XEST-10, XEST-11, and XEST-12 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.




**Fig. B.5.** Co-added EPIC images of field XEST-13, XEST-14, and XEST-15 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.




**Fig. B.6.** Co-added EPIC images of field XEST-16, XEST-17, and XEST-18 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.




**Fig. B.7.** Co-added EPIC images of field XEST-19, XEST-20, and XEST-21 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.



**Fig. B.8.** Co-added EPIC images of field XEST-22, XEST-23, and XEST-24 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.



**Fig. B.9.** Co-added EPIC images of field XEST-25, XEST-26, and XEST-27 (from top to bottom). *Left:* Smoothed images, color coded for hardness; *right:* coordinate grid and TMC identifications included.



**Fig. B.10.** Co-added EPIC images of field XEST-28. *Left:* Smoothed image, color coded for hardness; *right:* coordinate grid and TMC identifications included.