# Thermal-bath effects in quantum quenches within quantum critical regimes

Francesco Tarantelli

Dipartimento di Fisica dell'Università di Pisa and INFN, Largo Pontecorvo 3, I-56127 Pisa, Italy

Ettore Vicari

Dipartimento di Fisica dell'Università di Pisa, Largo Pontecorvo 3, I-56127 Pisa, Italy

(Dated: May 15, 2023)

We address the out-of-equilibrium dynamics arising from quantum-quench (QQ) protocols (instantaneous changes of the Hamiltonian parameters) in many-body systems within their quantum critical regime and in contact with (homogeneously coupled) thermal baths. We consider two classes of QQ protocols. In one of them the thermal bath is used to prepare the initial finite-temperature Gibbs state; then, after quenching, the thermal bath is removed and the dynamics of the system is unitary. We also address a more complex QQ protocol where the thermal bath is not removed after quenching, thus the quantum evolution is also driven by the interaction with the bath, which may be described by appropriate master equations for the density matrix of the system, where a further relevant time scale, or inverse decay rate, characterizes the system-bath coupling. Under these QQ protocols, the critical system develops out-of-equilibrium scaling behaviors, which extend those for isolated critical systems, by introducing further scaling variables proportional to the temperature of the thermal bath and the decay rate of the system-bath interactions. These out-of-equilibrium scaling behaviors are checked by analyzing QQ protocols within fermionic Kitaev wires, or equivalently quantum Ising chains, supplemented with a particular modelization of thermal bath that guarantees the asymptotic thermalization within the Lindblad master equation for the dynamics of open systems.

#### I. INTRODUCTION

Thanks to the recent experimental progress in the realization and control of the dynamics of quantum manybody systems, see e.g. Refs. [1, 2], the out-of-equilibrium quantum dynamics of many-body systems has become an important theoretical issue. In particular, out-ofequilibrium phenomena have been addressed within the critical regimes of many-body systems at continuous quantum transitions (CQTs) [3–5], where collective behaviors give rise to zero-temperature singularities in the equilibrium low-energy properties of the system, and the universal critical behaviors are determined by a limited number of relevant features, such as the global symmetry, the symmetry-breaking pattern, dimensionality, etc.. Within critical regimes and in the appropriate thermodynamic or finite-size scaling (FSS) limits, one can achieve a complete characterization of the complex dynamics of many-body systems by controlling a limited number of renormalization-group (RG) perturbations. The universal scaling behaviors at CQTs extend beyond the equilibrium conditions [5]. Indeed dynamic protocols entailing out-of-equilibrium evolutions develop scaling behaviors as well, in the appropriate limits, related to the universality class of the CQT. For example, out-of-equilibrium scaling behaviors emerge when analyzing the quantum evolutions arising from a quantum quench (QQ), see e.g. Refs. [5–11], or from slow changes of the Hamiltonian parameters across the transition point, such as the protocols associated with the so-called quantum Kibble-Zurek problem, see e.g. Refs. [5, 12–23].

These out-of-equilibrium issues have been mostly addressed within isolated many-body systems, unitarily driven by their Hamiltonian and the Schrödinger equation. In this paper we extend such studies to investigate how the interaction with a thermal bath, coupled homogeneously to the system, affects the out-ofequilibrium dynamics of many-body systems within the critical regime of a zero-temperature quantum transition, such as that arising from a QQ or a slow crossing of the quantum critical regime.

The role of the temperature within the equilibrium critical behavior at a CQT is generally associated with one of the relevant RG perturbations at the stable fixed point of the RG flow controlling the quantum criticality [3–5, 24]. Therefore, the quantum scaling behavior can be only observed in the zero-temperature limit. More precisely, the quantum scaling limit requires that the zero-temperature critical point is approached keeping the ratio  $T/\Delta$  fixed, where  $\Delta$  is the gap at the quantum critical point, which is generally power-law suppressed. For example, in the FSS limit the gap is suppressed as  $\Delta \sim L^{-z}$  at the critical point, where L is the size of the system and z > 0 is the universal dynamic exponent associated with universality class of the CQT. Within the equilibrium critical regime the temperature enters the asymptotic FSS laws through a further dependence of the scaling functions on the scaling variable  $\Xi \equiv TL^z \sim T/\Delta$ .

The role of the temperature becomes less definite when we consider out-of-equilibrium behaviors, because the temperature of the system is an equilibrium concept. However, one may consider the effects of thermal baths in contact with the system during its out-of-equilibrium dynamics. The main feature of a thermal bath is that it eventually drives the system toward thermalization at its temperature T, in the large-time limit of the evolution of the system in contact with the thermal bath. The thermalization process must somehow introduce a further time scale  $\tau$  in the problem, characterizing the approach of the system to the thermal state when it is put in contact with the thermal bath. Such time scale is expected to play an inportant role in the out-of-equilibrium dynamics of the system in contact with the thermal bath. In this paper we investigate these issues within the simplest dynamic protocols giving rise to out-of-equilibrium behaviors, i.e. those entailing instantaneous QQs of the Hamiltonian parameters starting from equilibrium thermal conditions.

A quench protocol is generally performed by suddenly varying a parameter within a family of Hamiltonians, such as

$$\hat{H}(w) = \hat{H}_c + w\hat{H}_p,\tag{1}$$

where  $\hat{H}_c$  and  $\hat{H}_p$  are independent of the parameter w, and  $[\hat{H}_c, \hat{H}_p] \neq 0$ . In a standard QQ protocol for closed systems, one usually starts from the ground state  $|\Phi_0, w_i\rangle$ of the Hamiltonian  $\hat{H}(w_i)$  associated with an initial value  $w_i$  of the parameter w, with corresponding density matrix  $\rho_i = |\Phi_0, w_i\rangle \langle \Phi_0, w_i|$ . At a given time, t = 0 say, the Hamiltonian parameter is suddenly changed from  $w_i$  to  $w \neq w_i$ , and the subsequent quantum evolution is supposed to be unitarily driven by the Hamiltonian  $\hat{H}(w)$ , that is  $|\Psi(t)\rangle = e^{-i\hat{H}(w)t} |\Phi_0, w_i\rangle$  (hereafter we set  $\hbar = 1$ ). Several interesting issues have been investigated within QQ dynamic protocols. They include the long-time relaxation and the consequent spreading of quantum correlations and entanglement, the statistics of the work, localization effects due to the mutual interplay of interactions and disorder, dynamical phase transitions, the dynamic scaling close to quantum transitions, effects of dissipation or of measurements due to interactions with an environment (see, e.g., Refs. [5, 9, 25–78]).

To focus on the out-of-equilibrium dynamics close to a quantum transition, we assume that the Hamiltonian  $H_c$  in Eq. (1) is critical, thus  $w = w_c = 0$  represents a quantum critical point. We recall that the critical behavior around the CQT point  $w_c = 0$  is characterized by a diverging length scale  $\xi \sim |w|^{-\nu}$  of the quantum critical modes, and the power-law suppression  $\Delta \sim \xi^{-z}$  of the gap. The out-of-equilibrium dynamics at CQTs develops scaling behaviors controlled by the universality class of the quantum transition, for example when the Hamiltonian parameters are slowly varied across the critical regime [5, 21, 23], and in the case of *soft* QQ protocols when both the initial and final values of the quenching parameters are such to maintain the system within the critical regime [5, 9, 59]. In particular, soft QOs require that the energy scale of the QQ [i.e. the difference of the energy  $\langle \Psi(t) | \hat{H}(w) | \Psi(t) \rangle$  of the evolving state  $| \Psi(t) \rangle$  for t > 0 and the ground state of  $\hat{H}(w)$  is sufficiently small. i.e. comparable with the energy gap  $\Delta \sim L^{-z}$  of the spectrum at the transition point in finite-size systems.

To study the effects of a thermal bath in the out-of-

equilibrium behavior arising from a QQ within the critical regime, we consider two protocols where the thermal baths are involved in different ways:

(i) Within the first protocol the thermal bath is used to prepare the system in a finite-temperature Gibbs state, described by the thermal density matrix (hereafter we set the Boltzmann constant  $k_B = 1$ )

$$\rho_t(w_i, T) = \sum_n e^{-E_n(w_i)/T} |\Phi_n, w_i\rangle \langle \Phi_n, w_i|, \quad (2)$$

where  $|\Phi_n, w_i\rangle$  are the eigenstates of  $\hat{H}(w_i)$ . Then the quantum evolution after the quench of the Hamiltonian parameters at t = 0 is unitary and driven by the Hamiltonian  $\hat{H}(w)$  only, i.e., the thermal bath is removed during the quantum evolution for t > 0. Therefore, the evolution of the density matrix is driven by the equation

$$\partial_t \rho(t) = -i[\hat{H}(w), \rho(t)], \qquad \rho(t=0) = \rho_t(w_i, T).$$
 (3)

(ii) In the second protocol the starting point is the same, i.e. the Gibbs state (2), but the thermal bath is not removed after quenching. Therefore, the out-of-equilibrium quantum evolution for t > 0 is not unitary anymore, but it is also driven by the interaction with the thermal bath. Under some conditions, discussed in Refs. [5, 79–84], the nonunitary evolution arising from the thermal baths can be described by a Lindbald master equation governing the time evolution of the density matrix of the system, which can be written as

$$\partial_t \rho = \mathcal{L}[\rho] \equiv -i \left[ \hat{H}(w), \rho \right] + \gamma \mathbb{D}_T[\rho], \tag{4}$$

where  $\mathcal{L}$  is a Liouvillian superoperator, and  $\mathbb{D}_T$  is a dissipative driving whose strength is controlled by the homogeneous coupling  $\gamma$ , playing the role of the decay rate (inverse time scale) associated with the interactions between the system and the bath. The operator  $\mathbb{D}_T$  is assumed to be such that the Lindbald master equation (4) drives the system toward an equilibrium Gibbs state at temperature T in the large-time limit.

We argue that, for both types of protocols and sufficiently small temperatures of the thermal baths, the out-of-equilibrium time evolution within the critical regime develop a nontrivial out-of-equilibrium FSS (OFSS) limit, with peculiar scaling behaviors, similar to those arising for closed systems. The effects of the thermal baths can be taken into account by appropriate extensions of the out-of-equilibrium zero-temperature scaling laws describing soft quantum QQs within the critical regime of isolated systems, already put forward by earlier works [5, 9]. As a theoretical laboratory to check our extended OFSS laws, we consider the quantum Ising chain [4], or the equivalent fermionic Kitaev wire [85], supplemented with a particular modelization of the thermal bath that guarantees the asymptotic thermalization within the Lindblad formulation of the dynamics of open systems with quadratic Hamiltonians [84, 86], such as the fermionic Kitaev wire.

Our analyses are developed within FSS frameworks, which generally simplify the study of the universal features of critical behaviors, with respect to studies in the thermodynamic limit. In the FSS limit the general requirement of a large length scale  $\xi$  of the critical correlations is not subject to further conditions on the system size L. It only requires that  $\xi \sim L$ , while critical behaviors in the thermodynamic limit requires  $\xi \ll L$ . Therefore much larger systems are necessary to probe analogous length scales  $\xi$  in the thermodynamic limit. Equilibrium and out-of-equilibrium FSS behaviors are often observed for systems of moderately large size. see e.g. Refs. [5, 9, 57, 87, 88]. Thus FSS behaviors should be more easily accessed by numerical computations and experiments where the quantum dynamics can be monitored for a limited number of particles or spins, such as experiments with quantum simulators in laboratories, e.g., by means of trapped ions [89, 90], ultracold atoms [91, 92], or superconducting qubits [93, 94].

The paper is organized as follows. In Sec. II we present the fermionic Kitaev wire, equivalent to the quantum Ising chain, and the model of thermal bath that we use as theoretical laboratory for our study; we also outline the QQ protocols that we consider and define the observables to monitor the quantum evolution after quenching. In Sec. III we outline the out-of-equilibrium scaling scenarios that are expected to be developed under the dynamic QQ protocols considered, and support them by numerical computations for the fermionic Kitaev wires in contact with the thermalizing bath. Finally, in Sec. IV we summarize, draw our conclusions, and add some remarks on the extension of this study to the dynamic Kibble-Zurek protocols slowly crossing quantum critical regimes. The appendix reports some details on the numerical computations for the QQ protocols within fermionic Kitaev wires in contact with a thermal bath.

## II. KITAEV FERMIONIC WIRES AND THERMAL BATHS

## A. The fermionic Kitaev chain

We consider fermionic Kitaev wires of L sites with open boundary conditions, whose quantum unitary dynamics is driven by the Hamiltonian [85]

$$\hat{H}_{\rm K} = -J \sum_{x=1}^{L-1} \left( \hat{c}_x^{\dagger} \hat{c}_{x+1} + \hat{c}_x^{\dagger} \hat{c}_{x+1}^{\dagger} + \text{h.c.} \right) - \mu \sum_{x=1}^{L} \hat{n}_x, \quad (5)$$

where  $\hat{c}_x$  is the fermionic annihilation operator associated with the site x of the chain,  $\hat{n}_x \equiv \hat{c}_x^{\dagger} \hat{c}_x$  is the particle density operator. In the following we assume J as the energy scale, thus we set J = 1.

The Hamiltonian (5) can be mapped into a quantum Ising chain, by means of the Jordan-Wigner transformation, see, e.g., Ref. 4. The corresponding spin model is the quantum Ising chain with open boundary conditions, i.e.

$$\hat{H}_{\rm Is} = -\sum_{x=1}^{L-1} \hat{\sigma}_x^{(1)} \hat{\sigma}_{x+1}^{(1)} - g \sum_{x=1}^{L} \hat{\sigma}_x^{(3)}, \tag{6}$$

 $\hat{\sigma}_x^{(k)}$  being the Pauli matrices and  $g = -\mu/2$ . In the following we prefer to stick with the Kitaev quantum wire, because the thermal baths and observables that we consider are best defined within the fermionic model. However, the general scaling scenarios that will emerge apply to both models.

The Kitaev model undergoes a CQT at  $\mu = \mu_c = -2$ (corresponding to  $g = g_c = 1$  in the quantum Ising chain), between a disordered quantum phase for  $\mu < \mu_c$ (corresponding to g > 1) and an ordered quantum phase for  $|\mu| < |\mu_c|$  (corresponding to |g| < 1). Thus, we define

$$w = \mu - \mu_c = \mu + 2, \tag{7}$$

so that one can easily see the correspondence between the Kitaev Hamiltonian (5) and the generic one reported in Eq. (1), i.e.  $\hat{H}_c$  corresponds to the Hamiltonian (5) for  $\mu = \mu_c$ , and  $\hat{H}_p = -\sum_{x=1}^L \hat{n}_x$ . The continuous transition at  $w = w_c$  belongs to the two-dimensional Ising universality class [4, 5], characterized by the length-scale critical exponent  $\nu = 1$ , related to the RG dimension  $y_w = 1/\nu = 1$  of the Hamiltonian parameter w. This implies that, approaching the critical point, the length scale  $\xi$  of the critical quantum fluctuations diverges as  $\xi \sim |w|^{-\nu}$ . The dynamic exponent z = 1 associated with the unitary quantum dynamics can be obtained from the power law  $\Delta \sim \xi^{-z}$  of the vanishing gap with increasing  $\xi$ . Moreover, the RG dimension of the fermionic operators  $\hat{c}_j$  and  $\hat{c}_j^{\dagger}$  at the CQT is  $y_c = 1/2$ , and that of the particle density operator  $\hat{n}_x$  is  $y_n = 1$  [4, 5].

#### B. Modelization of the thermal bath

In our study we consider a modelization of interaction with a thermal bath within the Lindblad master equation (4), whose asymptotic large-time behavior leads to a Gibbs density matrix at a given finite temperature T. In particular, we consider the proposal developed in Ref. [84] which applies to quantum models described by quadratic Hamiltonians, such as that of the fermionic Kitaev wires. This provides a relatively simple modelization of a thermal bath leading to thermalization in the largetime limit of the corresponding Lindblad master equation for the density matrix of the system.

The Kitaev Hamiltonian (5) with open boundary conditions can be diagonalized in the Nambu field space by a Bogoliubov transformation, see e.g. Refs. [84, 95, 96], so that we can rewrite it as

$$\hat{H}_{\rm K} = \sum_{k=1}^{L} \omega_k \, \hat{b}_k^{\dagger} \, \hat{b}_k, \qquad (8)$$

where  $\omega_k$  are values of the spectrum of the Bogoliubov eigenoperators  $\hat{b}_k$  (we are neglecting an irrelevant constant term). Note that both  $\omega_k$  and  $\hat{b}_k$  depend on the Hamiltonian parameter  $\mu$ . The relation between the fermionic operators  $\hat{c}_x$  and the Bogoliubov eigenoperators  $\hat{b}_k$  can be generally written as [84, 95, 96]

$$\hat{c}_x = \sum_{k=1}^{L} A_{xk} \,\hat{b}_k + B_{xk} \,\hat{b}_k^{\dagger}, \qquad (9)$$

where A and B are appropriate  $L \times L$  matrices depending on  $\mu$ . Following Refs. [84, 86], we write the dissipator  $\mathbb{D}_T[\rho]$  in the Lindblad master equation (4) in terms of the Bogoliubov eigenoperators as

$$\mathbb{D}_{T}[\rho] = \sum_{k} [1 - f(\omega_{k}, T)] \left( 2 \,\hat{b}_{k} \,\rho \,\hat{b}_{k}^{\dagger} - \{ \hat{b}_{k}^{\dagger} \hat{b}_{k}, \rho \} \right) \\ + \sum_{k} f(\omega_{k}, T) \left( 2 \,\hat{b}_{k}^{\dagger} \,\rho \,\hat{b}_{k} - \{ \hat{b}_{k} \hat{b}_{k}^{\dagger}, \rho \} \right), \quad (10)$$

where

$$f(\omega_k, T) = \left(1 + e^{\omega_k/T}\right)^{-1}.$$
 (11)

When using this homogeneous dissipator term, the Lindblad master equation (4) ensures the asymptotic largetime thermalization [84]. Therefore,

$$\lim_{t \to \infty} \rho(t) = \rho_t(w, T), \tag{12}$$

$$\rho_t(w,T) = \sum_n e^{-E_n(w)/T} |\Phi_n, w\rangle \langle \Phi_n, w|, \qquad (13)$$

where  $\rho_t(w,T)$  is the density matrix representing the thermal state,  $E_n(w)$  and  $|\Phi_n, w\rangle$  are the eigenvalues and eigenstates of  $\hat{H}(w)$ . The asymptotic approach to the thermal distribution is controlled by the decay-rate parameter  $\gamma$  [84]. Indeed the Liouvillian gap  $\Delta_{\mathcal{L}}$  that controls the exponential approach to the asymptotic stationary state of the Lindblad equation is proportional to the decay rate  $\gamma$ , i.e.

$$\Delta_{\mathcal{L}} \sim \gamma. \tag{14}$$

The above modelization of thermal baths provides a useful theoretical laboratory to investigate issues related to the out-of-equilibrium dynamics in the presence of thermal baths. Its derivation has been thoroughly discussed in Ref. [84]. We also mention that it has been employed in Refs. [86, 97]. Some details of the computations using the Lindblad master equation (4) with the dissipator (10) are reported in the appendix.

## C. Quantum-quench protocols

As already anticipated in Sec. I, we consider two protocols, differing for the absence or presence of the contact with the thermal bath during the quantum evolution after quenching, giving respectively rise to unitary or dissipative dynamics after quenching. We call them *unitary* and *dissipative* QQ protocols, respectively.

- Unitary QQ protocol: In this simplest QQ protocol the role of the thermal bath is limited to that of preparing the initial Gibbs state  $\rho_t(w_i, T)$  at t = 0, reported in Eq. (2). This can be obtained by keeping the thermal bath in contact with the system for a sufficiently long time  $t_{\rm th}$ , i.e  $t_{\rm th} \gg \gamma^{-1}$ . Then at t = 0 the Hamiltonian parameter is instantaneously quenched from  $w_i < 0$  to  $w \ge 0$  and the thermal bath is removed, so that the subsequent time evolution is that of a closed fermionic wire, i.e. it is unitary and only driven by the Hamiltonian of the system, cf. Eq. (3).
- Dissipative QQ protocol: The quantum evolution starts from the same initial Gibbs state  $\rho_t(w_i, T)$ , but the thermal bath is maintained in contact with the system after the QQ from  $w_i < 0$  to  $w \ge 0$ , at t = 0. Therefore, the quantum evolution for t > 0is driven by the Lindblad master equation (4) with the dissipator term (10). Note that this dynamic protocol entails a further time scale  $\tau = \gamma^{-1}$ , characterizing the asymptotic exponential approach to the large-time stationary Gibbs state associated with the Hamiltonian  $\hat{H}(w)$  and temperature T.

#### D. Observables monitoring the time evolution

To characterize the dynamic properties of the quantum evolution after the QQ at t = 0, we consider the subtracted particle-density average

$$n_s(t,L) = \frac{1}{L} \operatorname{Tr} \left[ \rho(t) \sum_{x=1}^{L} \hat{n}_x \right] - n_c(L), \qquad (15)$$

where  $n_c(L)$  is the ground-state energy density of the Kitaev wire of size L at the critical point  $w_c = 0$  (in the infinite-size limit  $n_c = 1/2 - 1/\pi$  [95]). Note that the particle density operator  $\hat{n}_x$  and the transverse spin component  $\hat{\sigma}_x^{(3)}$  of the quantum Ising chain (6) are trivially related, indeed  $\hat{\sigma}_x^{(3)} = 2\hat{n}_x$ . In the definition of  $n_s$ , the subtraction of  $n_c(L)$  simplifies the scaling behavior of  $n_s(t, L)$  within the critical regime, cancelling the leading analytical behavior [5, 24]. To monitor the spatial correlations, we also consider

$$P(x, y, t) = \operatorname{Tr}[\rho(t) \left(\hat{c}_x^{\dagger} \hat{c}_y^{\dagger} + \hat{c}_y \hat{c}_x\right)], \qquad (16)$$

$$C(x, y, t) = \text{Tr}[\rho(t) \left(\hat{c}_{x}^{\dagger} \hat{c}_{y} + \hat{c}_{y}^{\dagger} \hat{c}_{x}\right)].$$
(17)

Some details on the computation of the above quantities during the time evolution of the QQ protocols are reported in the appendix.

### III. OUT-OF-EQUILIBRIUM SCALING

We now discuss the out-of-equilibrium behaviors arising from the QQ protocols outlined in Sec. II C. We show that they develop OFSS behaviors where the effects of the thermal baths are taken into account by appropriate extensions of the out-of-equilibrium zerotemperature scaling laws describing soft QQs in closed systems within their critical regime, already put foward by earlier works [5, 9].



FIG. 1: The quantum evolution of the subtracted particle density  $n_s(t)$ , cf. Eq. (15), for the dissipative QQ protocol entailing a dissipative dynamics after the QQ at t = 0 of the Hamiltonian parameter w, describing the persistent interaction with the thermal bath, cf. Eqs. (4) and (10). These curves refer to a system of size L = 60, temperature T = 2 of the thermal bath, quenching from  $w_i = -0.01$  to w = 0, and various values of the decay rate  $\gamma$  (the case  $\gamma = 0$  corresponds to the evolution of the close system). We plot the difference  $n_s(t, L, T) - n_{s,eq}(L, T)$  which is expected to vanish in the large-time limit. In this figure and in the following ones, the unity that we use are such that  $\hbar = 1$ ,  $k_B = 1$ , and J = 1.

The OFSS behaviors that we put forward for QQ protocols considered are verified by numerical computations for the fermionic Kitaev wire up to relatively large sizes. See the appendix for details on such calculations.

As a preliminary example of out-of-equilibriun QQ behaviors that we want to address, in Fig. 1 we show some results for the quantum evolution of the subtracted particle density (15) along the dissipative protocol outlined in Sec. II C, after quenching a fermionic Kitaev wire of size L = 60, from  $w_i = -0.01$  to w = 0, in the presence of a thermal bath at a temperature T = 2, and various values of the decay rate  $\gamma$ . The quantum evolution turns out to have a significant dependence on the decay-rate parameter  $\gamma$  that characterized the interactions between the system and the thermal bath. Indeed, the curves of the subtracted particle density appear to approach its equilibrium value  $n_{s,eq}(w = 0, T = 2) \approx 0.0004601...$  (while at t = 0 we have  $n_{s,eq}(w = w_i, T = 2) \approx 0.126598...$ ), faster and faster with increasing  $\gamma$ , actually exponentially as  $\exp(-t/\tau)$  with  $\tau \sim \gamma^{-1}$ , conferming the role

of decay rate of the parameter  $\gamma$  within the Lindblad master equation, cf. Eq. (14). Analogous results are obtained for other observables, such as fermionic correlation functions defined in Sec. II D. In the following we put forward an out-of-equilibrium scaling theory for these out-of-equilibrium phenomena within the quantum critical regime.

## A. Zero-temperature scaling in quantum quenches

We now provide a brief summary of the out-ofequilibrium scaling theory for close systems, describing QQ protocols within the critical regime [5, 9]. The initial state is the ground state associated with an initial value  $w_i < 0$ , and, after the instantaneous quench at t = 0 from  $w_i$  to w, the quantum evolution is driven by the Schrödinger equation.

Out-of-equilibrium scaling laws can be obtained by extending those valid at equilibrium, allowing for a time dependence essentially controlled by the time scaling variable  $\Theta \sim t \Delta$ , which is obtained by assuming that the relevant time scale of the critical modes is proportional to the inverse energy difference  $\Delta$  of the lowest states. We refer to Ref. [5] for a through presentation of the scaling arguments leading to the asymptotic OFSS behaviors.

Let us consider the out-of-equilibrium evolution (after quenching) of generic observables, such as the expectation value O at time t of a local operator  $\hat{O}(\boldsymbol{x})$ and its fixed-time correlations  $G_O = \langle \hat{O}(\boldsymbol{x}) \hat{O}(\boldsymbol{y}) \rangle$ . The general working hypothesis underlying out-of-equilibrium FSS frameworks is that the expectation value of  $\hat{O}(\boldsymbol{x})$ and its correlation functions obey asymptotic homogeneous scaling laws [5], such as

$$O(t, \boldsymbol{x}, L, w_i, w) \approx b^{-y_o} \mathcal{O}(t/b^z, \boldsymbol{x}/b, L/b, b^{y_w} w_i, b^{y_w} w),$$
(18)

where b is an arbitrary (large) length scale,  $y_o$  is the RG dimension of the local operator  $\hat{O}_x$  and the RG exponents  $y_w$  and z are determined by the universality class of the CQT (they are the RG dimensions of the Hamiltonian parameter w and the temperature T, respectively). Thus both the initial and final values of w, i.e.  $w_i$  and w, take the same RG exponent  $y_w$ , being coupled to the RG perturbation  $\hat{H}_p$  within the Hamiltonian. Note that we do not assume translation invariance, which is generally broken by the presence of boundaries, such as those arising from open boundary conditions.

OFSS can be straightforwardly derived by fixing b = Lin the above homogenous scaling law. Then, we expect the OFSS of the expectation value O of a generic local operator  $\hat{O}_{\boldsymbol{x}}$ , of its spatial average  $\hat{O}_a = L^{-d} \sum_{\boldsymbol{x}} \hat{O}_{\boldsymbol{x}}$ , and its two-point correlation function  $G_O$ , develop the asymptotic OFSS behavior [5, 9]

$$O(t, \boldsymbol{x}, L, w_i, w) \approx L^{-y_o} \mathcal{O}(\Theta, \boldsymbol{X}, \Phi_i, \Phi),$$

$$O_a(t, L, w_i, w) \approx L^{-y_o} \mathcal{O}_a(\Theta, \Phi_i, \Phi),$$

$$G_O(t, \boldsymbol{x}_1, \boldsymbol{x}_2, L, w_i, w) \approx L^{-2y_o} \mathcal{G}_O(\Theta, \boldsymbol{X}_1, \boldsymbol{X}_2, \Phi_i, \Phi),$$
(19)

where the scaling variables appearing in the scaling functions  $\mathcal{O}$ ,  $\mathcal{O}_a$ , and  $\mathcal{G}_O$  are defined as

$$\Theta \equiv \frac{t}{L^z}, \quad \mathbf{X}_i \equiv \frac{\mathbf{x}_i}{L}, \quad \Phi_i \equiv L^{y_w} w_i, \quad \Phi \equiv L^{y_w} w. \quad (20)$$

The OFSS limit is obtained in the large-L and large-tlimit keeping the above scaling variables fixed. These conditions ensure that the system remains within the universal critical regime during the quantum evolution. Note that in the scaling law (20) the dynamic features are essentially encoded in the time dependence of the scaling variable  $\Theta \sim t \Delta$ . The other features, in particular when  $w_i = w$ , are analogous to those arising from equilibrium FSS at CQTs [5, 24], where the argument  $\Phi = L^{y_w} w$  of the scaling functions is controlled by the RG dimension  $y_w$  of the relevant parameters w at the RG fixed point associated with the CQT.

The above OFSS equations can be straightforwardly applied to the observables defined in Sec. IID, after a quench from  $w_i$  to w at t = 0, keeping into account that the RG dimension of the subtracted particle density is  $y_n = 1$ , and that of the fermionic operator  $\hat{c}_x$ is  $y_c = 1/2$ . Note that the dominant analytical contributions to the particle density [5, 24] coming from the analytical background are canceled in the difference  $n_s$ defined in Eq. (15), whose leading asymptotic behavior arises from the quantum critical modes, therefore it is analogous to that of  $O_a$  in Eq. (19), with  $y_o = y_n$ . Analogously one can apply the OFSS in Eq. (19) to observables and correlation functions constructed with the spin operators of the quantum spin chain (6). The OFSS functions are expected to be universal with respect to the microscopic details of the model, apart from nonuniversal multiplicative rescaling and normalizations of its arguments. Within isolated fermionc Kitaev wires and quantum Ising chains, the OFSS arising from soft QQs has been verified by numerical computations for various boundary conditions, and also along their quantum first-order transition line [5, 9].

The OFSS limit is expected to be approached with power-law suppressed corrections. There are various sources of scaling corrections when approaching the OFSS. Of course, they include those that are already present at equilibrium. In particular, the irrelevant RG perturbations are sources of scaling corrections for the asymptotic behavior of the free-energy density [5, 99]. In the case of one-dimensional quantum systems undergoing CQTs belonging to the two-dimensional Ising universality class, the leading scaling corrections from irrelevant RG perturbations are suppressed as  $L^{-\omega}$  with  $\omega = 2$  [24, 98]. However, other contributions may become more relevant [5, 24, 99], such as those arising from the presence of analytical backgrounds, from the presence of boundaries (which generally gives rise to O(1/L)) corrections), and, in the case of correlation functions, from RG mixings of the source fields [this for example happens in the case of the correlation functions of the fermionic field  $\hat{c}_x$ , for which corrections are O(1/L)].

These scaling corrections have been confirmed by numerical results [5, 24]. Therefore, we expect that the asymptotic OFSS of fermionic Kitaev wires and quantum Ising chains with open boundary conditions is generally approached with O(1/L) corrections.

## B. OFSS along the unitary QQ protocol

For the simplest unitary protocol reported in Sec. II C, where the quantum evolution is that of the isolated fermionic wire, the request that the dynamics remains within the critical regime implies that the temperature of the initial Gibbs state must be appropriately suppressed in the large-L OFSS limit, to obtain a nontrivial out-ofequilibrium critical limit. This is analogous to what happens within the equilibrium FSS, where one introduces the scaling variable [3–5]

$$\Xi \equiv L^z T, \tag{21}$$

to allow for a nonzero temperature in the FSS of the observables. Therefore, like equilibrium FSS, we conjecture that the temperature of the initial Gibbs state enters the OFSS associated with the unitary QQ protocol by adding a further dependence on  $\Xi$  in the scaling functions (19). In other words, a nontrivial asymptotic OFSS limit is expected to be realized in the large-*L* and large-*t* limits keeping also  $\Xi$  fixed, beside the scaling variables already defined in Eq. (20). Therefore, we expect that the OFSS of standard QQ protocols starting from ground states, cf. Eq. (19), changes into

$$O(t, \boldsymbol{x}, L, w_i, w, T) \approx L^{-y_o} \mathcal{O}(\Theta, \boldsymbol{X}, \Phi_i, \Phi, \Xi),$$
 (22)

and analogously for its spatial average  $O_a$  and the correlation function  $G_O$ .

The numerical analysis for the fermionic Kitaev wire under the unitary protocol fully support to this OFSS, obtained by extending the QQ FSS behaviors of closed systems starting from an initial ground state. This is clearly demonstrated by the curves reported in Fig. 2, associated with the quantum evolutions of the subtracted particle density  $n_s(t)$  and the fermionic correlation P(x, y, t) (the other fermionic correlation C(x, y, t)develops an analogous OFSS).

## C. OFSS along the dissipative QQ protocol

We now discuss the dynamics arising from the dissipative protocol outlined in Sec. II C, when the quantum evolution after quenching is described by the Lindblad master equation (4) with the thermal-like dissipator (10), to modelize the interaction with a thermal bath characterized by a temperature T (which does not change after quenching) and decay rate  $\gamma$ .

We expect that the temperature T of the thermal bath must be rescaled as in the case of the unitary QQ protocol, i.e. we must consider again the associated scaling



FIG. 2: OFSS behavior of the subtracted particle density (bottom) and the fermionic correlation function P(x = L/3, y = 2L/3, t), cf. Eq. (16), arising from the unitary QQ protocol, for various lattice sizes L, at fixed  $\Xi = L^z T = 1$ ,  $\Phi_i = L^{y_w} w_i = -1$  and  $\Phi = L^{y_w} w = 0$ , versus the time scaling variable  $\Theta = t/L^z$ . These computations nicely support the OFSS behaviors reported in Eq. (22). The inset of the bottom figure shows that the approach to the OFSS limit is consistent with O(1/L) corrections. Analogous results are obtained for other values of the scaling variables.

variable  $\Xi$  already defined in Eq. (21). However, since the QQ moves the system out-of-equilibrium, also the decay rate  $\gamma$ , and corresponding time scale  $\tau = \gamma^{-1}$ , associated with the interactions with the thermal bath is expected to play a relevant role to establish a corresponding non-trivial OFSS limit. This was already noted in Ref. [97] in the analysis of dynamic protocols entailing the variation of the temperature at the critical point.

When keeping  $\tau$  constant in the FSS limit where the scaling variable  $\Theta = t/L^z$  is kept fixed, in the large-L limit we have eventually that

$$t = \Theta L^z \gg \tau, \tag{23}$$

which is the condition ensuring thermalization for any finite value  $\Theta > 0$ . Therefore, when keeping  $\tau$  fixed, the quantum evolution is not expected to develop a non-trivial OFSS limit. Indeed, in the large-L limit, the



FIG. 3: Equilibrium FSS of the subtracted particle density  $n_{s,eq}$  at the critical point w = 0, versus the rescaled temperature  $\Xi = L^{z}T$ . With increasing L, the data show the expected convergence to the equilibrium FSS reported in Eq. (24) with  $y_{n} = 1$ .



FIG. 4: Quantum evolution of the subtracted particle density arising from the dissipative QQ protocol, when rescaling all quantities involved in the quench protocol, except for the decay rate  $\gamma$ . With increasing L, the curves appear to approach the equilibrium FSS value at finite temperature (where the temperature dependence enters through the scaling variable  $\Xi = L^{z}T$ ) faster and faster, reflecting a nonuniform convergence for any  $\Theta > 0$ . The dashed line shows the equilibrium value of  $n_{s}$  for  $\Phi = 0$  and  $\Xi = 1$ , which is asymptotically approached by the various curves.

system turns out to suddenly approach an equilibrium Gibbs state (associated with the Hamiltonian parameter w and temperature T) with respect to the rescaled time  $\Theta$ , without any further relevant evolution of the system for any  $\Theta > 0$ . Therefore, if the temperature is rescaled by keeping  $\Xi = L^{z}T$  fixed, we must recover the equilibrium FSS behavior in the presence of a thermal bath at temperature T, such as that associated with the subtracted particle density [5, 24]

$$n_{s,\text{eq}}(w,L,T) \approx L^{-y_n} \mathcal{N}(\Phi,\Xi),$$
 (24)

where  $\Phi = L^{y_w} w$ , and the temperature dependence enters through the associated scaling variable  $\Xi = L^z T$ . In Fig. 3 we show some equilibrium data at the critical point  $w = \Phi = 0$ , versus  $\Xi$ , showing the approach to the asymptotic large-*L* equilibrium FSS (24). The realization of the equilibrium FSS within the QQ protocol at fixed  $\gamma$  is demonstrated by the plots reported in Fig. 4, which show the somewhat trivial convergence toward the equilibrium FSS for any finite  $\Theta > 0$ .

The above results suggest that also the the decay rate  $\gamma$  of the system-bath interactions must be rescaled to observe a nontrivial OFSS limit as a function of the time scaling variable  $\Theta$ , to create the conditions for a balanced competition between the critical Hamiltonian driving and the interactions with the thermal bath. As already put forward in the case of other homogeneous dissipative terms in the Lindblad equation [5, 55, 100–102], for example associated with particle-decay or particle-pumping dissipative mechanisms, a nontrivial OFSS limit is obtained by rescaling the decay rate of the dissipative term, so that the scaling variable

$$\Gamma \equiv L^z \gamma \sim \gamma / \Delta \tag{25}$$

is kept fixed in the OFSS limit, where  $\Delta$  is the energy difference of the lowest eigenstates of  $\hat{H}(w)$  at the critical point  $w = w_c = 0$ . Then an OFSS behavior emerges from the nontrivial competition between the critical unitary dynamics and the dissipative driving arising from the thermal bath.

In conclusion, on the basis of the above scaling arguments, the OFSS arising from the dissipative QQ protocols in the presence of a thermal bath is expected to be given by

$$O_a(t, L, w_i, w, T, \gamma) \approx L^{-y_o} \mathcal{O}_a(\Theta, \Phi_i, \Phi, \Xi, \Gamma),$$
 (26)

and

$$G_O(t, \boldsymbol{x}_1, \boldsymbol{x}_2, L, w_i, w, T, \gamma) \approx (27)$$
$$L^{-2y_o} \mathcal{G}_O(\Theta, \boldsymbol{X}_1, \boldsymbol{X}_2, \Phi_i, \Phi, \Xi, \Gamma).$$

In the large- $\Gamma$  limit the above OFFS behaviors at fixed  $\Xi$  is expected to approach the corresponding equilibrium FSS, faster and faster in terms of  $\Theta$ , matching the behavior at finite  $\gamma$ . Moreover, we also expect that the equilibrium FSS is also approached in the large- $\Theta$  limit at fixed  $\Gamma$  and  $\Xi$ , independently of  $\Gamma$ , but faster and faster with increasing  $\Gamma$ .

Again, the numerical results for the particle density  $n_s(t)$  and correlation functions P and C fully support the above OFSS equations, i.e. Eq. (26) for  $n_s(t)$  with  $y_o = y_n = 1$ , and Eq. (27) for P and C with  $y_o = y_c = 1/2$ . Some results are reported in Fig. 5. We also stress that analogous results are expected for other observables, for example the correlation functions of the spin operator of the equivalent formulation provided by the quantum Ising chains.



FIG. 5: Quantum evolutions along the dissipative protocol, fully supporting the OFSS reported in Eqs. (26) and (27). We report curves for  $L n_s$  (bottom), L P(x = L/3, y = 2L/3, t)(middle), and C(x = L/3, y = 2L/3, t) (top), for various values of L, at fixed  $\Phi_i = -1$ ,  $\Phi = 0$ ,  $\Xi = 1$ , and two values of  $\Gamma = L^z \gamma$ , i.e.  $\Gamma = 1$ , 10 (except for the top figure where we only report data for  $\Gamma = 10$  to ensure a good readability). The inset of the top figure shows that the OFSS is approached with O(1/L) corrections. Analogous results are obtained for other values of the scaling variables.

### IV. CONCLUSIONS

We have reported a study of the effects of thermal baths to the out-of-equilibrium dynamics of many-body systems within their quantum critical regime close to a zero-temperature CQT. In particular, we analyze the outof-equilibrium quantum evolution arising from QQs of the Hamiltonian parameters within two different protocols involving a thermal bath coupled homogeneously to the system. Within the first protocol, named unitary QQ protocol, the thermal bath is used to prepare the system at t = 0 in a finite-temperature Gibbs state, then the dynamics after quenching of the Hamiltonian parameters is assumed unitary, i.e., the thermal bath is removed during the quantum evolution for t > 0. The second protocol, named dissipative QQ protocol, starts from the same initial condition, but the thermal bath is not removed after quenching, and the quantum evolution for t > 0 is assumed to be described by the Lindblad master equation (4). The dissipative term of the Lindblad equation is supposed to simulate a thermal bath, such that the manybody system is driven to a large-time finite-temperature Gibbs state. This dissipative protocol is characterized by a further time scale  $\tau = \gamma^{-1}$ , related to the decay rate of the interactions between the system and the bath.

Within OFSS frameworks, we argue that, when the thermal baths are associated with a sufficiently small temperature, their effects can be taken into account by appropriate extensions of the zero-temperature out-ofequilibrium scaling laws describing soft QQs of isolated systems within the critical regime. For the unitary QQ protocol, where the thermal bath only determines the initial Gibbs state and the evolution is unitary, a nontrivial OFFS limit is simply obtained by rescaling the temperature as  $T \sim L^{-z}$ , similarly to equilibrium FSS. Along the dissipative QQ protocol, where the thermal bath is not removed after quenching, the dynamics is more complicated, and the decay rate  $\gamma$  plays a relevant role. Indeed, in addition to the rescaling of the temperature Tassociated with thermal bath, one also needs to rescale  $\gamma$  as  $\gamma \sim L^{-z}$  to obtain a nontrivial OFSS. Otherwise, when keeping  $\gamma$  fixed, the dynamics converges toward the equilibrium FSS at finite temperature, which happens suddenly after quenching with respect to the time scale  $t_c \sim L^z$  of the critical regime. Therefore the scaling behavior when keeping  $\gamma$  fixed becomes somehow trivial, reproducing the equilibrium FSS for any rescaled time  $\Theta = L^{-z}t > 0$  in the large-L limit.

Our scaling arguments are supported by numerical results with the paradigmatic fermionic Kitaev model, or equivalently quantum Ising chain, at its CQT separating quantum disordered and ordered phases. We consider a particular modelization of the thermal bath that guarantees the asymptotic thermalization within the Lindblad formulation of the dynamics of open systems. However, we note that the scaling arguments used to arrive at the OFSS laws for critical QQs are general, and therefore we expect that the emerging out-of-equilibrium scenarios also apply to many-body systems at generic CQTs in contact with homogenous thermal baths, in any dimension.

We finally remark that the out-of-equilibrium scaling arguments we put forward, leading to the OFSS of QQs in the presence of a thermal bath, can be extended to other protocols giving rise to out-of-equilibrium dynamics. Another interesting class of dynamic protocols entails slow variations of the Hamiltonian parameters across the critical regime of a quantum transition, such as those associated with the quantum Kibble-Zurek (KZ) problem (see e.g. Refs. [5, 12–23]). In standard KZ protocols starting from the ground state for an initial parameter  $w_i < 0$ , the out-of-equilibrium quantum evolution arises from the linear time dependence of one Hamiltonian parameter,  $w(t) = t/t_s$  in Eq. (1), where  $t_s$  is the time scale of the KZ protocol. Since w(t) crosses the critical point at t = 0, the system passes through the quantum critical regime, moving it away from equilibrium even in the large- $t_s$  limit, and developing a peculiar out-of-equilibrium scaling behaviors. In particular, the interplay between the size L of the system and the time scale  $t_s$  of the protocol develops OFSS behaviors [5, 23] when  $t_s \to \infty$  and  $L \to \infty$ , keeping the scaling variables  $\Omega_t \equiv t/t_s^{\kappa} = t/t_s^{z/(y_w+z)}$  and  $\Upsilon \equiv t_s/L^{y_w+z}$  (thus  $\Omega_t = t/t_s^{1/2}$  and  $\Upsilon = t_s/L^2$  for the fermionic Kitaev wire or quantum Ising chain) fixed.

KZ-like protocols can be also extended to systems interacting with a thermal bath, such as that outlined in Sec. II B, starting from a Gibbs state for an initial  $w_i < 0$ and the temperature T of the thermal bath. Then we may consider a time evolution driven by the Lindblad master equation (4), with a time-dependent Hamiltonian H[w(t)] and the dissipator term (10), where also the Bogoliubov operators are assumed to be time dependent to adapt themselves to the time dependence of w. Analogously to the OFSS of QQs in contact with thermal baths, to define a nontrivial OFSS limit in KZ protocols, we expect that both the temperature T and the decay rate  $\gamma$  associated with the bath must be rescaled, as  $T \sim L^{-z}$  and  $\gamma \sim L^{-z}$ . If only the temperature of the thermal bath is rescaled as  $T \sim L^{-z}$ , while  $\gamma > 0$  is kept fixed, the time interval associated with a variation of  $\Omega_t$  in the KZ scaling limit, i.e.  $\Delta_{\Omega} t \sim t_s^{\kappa} \Delta \Omega_t$ , becomes eventually much larger than the time scale  $\tau \sim \gamma^{-1}$  of the interaction with the thermal bath. Since  $\tau/\Delta_{\Omega}t \to 0$  in the KZ limit, the system effectively thermalizes at each rescaled time  $\Omega_t$ . Therefore, in the KZ limit the quantum evolution is expected to pass through equilibrium finite-temperature states, thus effectively resulting into adiabatic evolutions reproducing the equilibrium finitetemperature FSS as a function of  $L^{y_w}w(t)$ . Therefore, like dissipative QQ protocols, the observation of a nontrivial OFSS in KZ protocols requires the simultaneous rescaling of the time scale  $\tau$  associated with the interaction with the thermal bath. The necessary rescaling of the decay rate  $\gamma$  of the dissipative term in the Lindblad master equation has been also put forward for KZ protocols in the presence of other dissipative mechanisms, such as those related to particle decay or pumping [100].

#### Acknowledgments

We thank Giulia Piccitto and Davide Rossini for interesting and useful discussions.

#### Appendix A: Details on the computations

In this section we provide some details of the computations for the fermionic Kitaev wire in the presence of a thermal bath.

### 1. Asymptotic thermal states

The dynamics of the system in contact with the thermal bath described by the Lindblad master equation (4) with the dissipator term (10) leads to thermal states, such as those described by the density matrix reported in Eq. (13). To compute the correlation functions of the fermionic operators  $\hat{c}_x$  in thermal states of the Hamiltonian  $\hat{H}(w)$ , one can use the relation with the Bogoliubov eigenoperators  $\hat{b}_k$ , cf. Eq. (9), and the thermal correlations of the Bogoliubov operators  $\hat{b}_k$ , i.e.

$$\langle b_k^{\dagger} b_q \rangle \equiv \text{Tr}[\rho_t(w,T) b_k^{\dagger} b_q] = \frac{\delta_{kq}}{1 + e^{\omega_k/T}},$$
 (A1)

corresponding to the standard Fermi-Dirac distribution function. Note also that the other correlations  $\langle b_k b_q \rangle$  and  $\langle b_k^{\dagger} b_q^{\dagger} \rangle$  vanish. Then the correlation functions of the original fermionic field  $\hat{c}_x$  can be straightforwadly obtained from Eq. (9).

## 2. Computations for the unitary protocol

In the unitary QQ protocol, one starts from a Gibbs state associated with the Hamiltonian parameter  $w_i$  and the temperature T, then at t = 0 one instantaneously changes  $w_i \to w$  and removes the contact with the thermal bath. Therefore the quantum evolution is unitary, described by the Schrödinger equation (3). One may easily obtain closed equations for the evolution of the correlation functions C and P defined in Eqs. (16) and (17).

We introduce the correlations

$$\mathscr{C}_{x,y} = \operatorname{Tr}\Big[\rho(t)\hat{c}_x^{\dagger}\hat{c}_y\Big], \quad \mathscr{P}_{x,y} = \operatorname{Tr}\Big[\rho(t)\hat{c}_x^{\dagger}\hat{c}_y^{\dagger}\Big], \quad (A2)$$

whose quantum evolution can be written as

$$\frac{d\mathscr{C}_{x,y}}{dt} = i \left[ \mathscr{C}_{x,y+1} - \mathscr{C}_{x-1,y} + \mathscr{C}_{x,y-1} - \mathscr{C}_{x+1,y} \right] - \\
-i \left( \mathscr{P}_{y,x-1}^{\dagger} - \mathscr{P}_{y,x+1}^{\dagger} \right) + i \left( \mathscr{P}_{x,y-1} - \mathscr{P}_{x,y+1} \right), \quad (A3)$$

$$\frac{d\mathscr{P}_{x,y}}{dt} = -i \left[ \mathscr{P}_{x,y+1} + \mathscr{P}_{x+1,y} + \mathscr{P}_{x,y-1} + \mathscr{P}_{x-1,y} \right] - \\
- 2i \mu \mathscr{P}_{x,y} - i \left( \delta_{x-1,y} - \delta_{x+1,y} \right) - \\$$

$$-i\left(\mathscr{C}_{x,y-1} - \mathscr{C}_{y,x-1} - \mathscr{C}_{x,y+1} + \mathscr{C}_{y,x+1}\right).$$
(A4)

The initial conditions are easily obtained by the relations with the thermal correlations of the Bogoliubov operators associated with the initial Gibbs state. Then the fermionic correlation function are obtained by

$$C(x, y, t) = 2 \operatorname{Re}\mathscr{C}_{x,y}(t), \ P(x, y, t) = 2 \operatorname{Re}\mathscr{P}_{x,y}(t).$$
 (A5)

The above differential equations are solved using the fourorder Runge-Kutta method. The particle density is obtained from the data of  $\mathscr{C}_{x,x} = \operatorname{Tr} \left[ \rho(t) \hat{c}_x^{\dagger} \hat{c}_x \right]$ .

## 3. Computations for the dissipative protocol

For the dissipative QQ protocol, where the thermal bath is kept in contact with the system, the evolution is driven by the Lindblad master equation (4), which can be equivalently written in terms of the time dependence of Heisenberg operators  $\hat{O}_{\rm H}(t)$ , i.e. [84, 86]:

$$\partial_t \hat{O}_{\mathrm{H}}(t) = i \left[ \hat{H}(w), \hat{O}_{\mathrm{H}}(t) \right] + \gamma \widehat{\mathbb{D}}_T[\hat{O}_{\mathrm{H}}(t)], \quad (A6)$$

where

$$\widehat{\mathbb{D}}_{T}[\hat{O}_{H}(t)] = \sum_{k} f(\omega_{k}) \left[ 2\hat{b}_{k}^{\dagger}\hat{O}_{H}(t)\hat{b}_{k} - \left\{ \hat{O}_{H}(t), \hat{b}_{k}\hat{b}_{k}^{\dagger} \right\} \right] + \sum_{k} (1 - f(\omega_{k})) \left[ 2\hat{b}_{k}\hat{O}_{H}(t)\hat{b}_{k}^{\dagger} - \left\{ \hat{O}_{H}(t), \hat{b}_{k}^{\dagger}\hat{b}_{k} \right\} \right], \quad (A7)$$

where  $\hat{b}_k$  are the Bogoliubov operators associated with the Hamiltonian  $\hat{H}(w)$ .

The initial state at t = 0 is the Gibbs state for the Hamiltonian parameter  $w_i$ . This state corresponds to the steady state solution of the Eq. (A6) with  $\hat{H}(w_i)$ . Then, the change of the Hamiltonian parameter to  $w \neq w_i$  leads to a change of the Bogoliubov operators diagonalizing the Hamiltonian. We call  $\{b'_k\}$  the operators which diagonalizes  $\hat{H}(w)$ ,

$$\hat{H}(w) = \sum_{k=1}^{L} \omega'_k \, \hat{b}'^{\dagger}_k \, \hat{b}'_k, \qquad (A8)$$

where  $\{\omega'_k\}$  is the Bogoliubov spectrum associated with  $\hat{H}(w)$ . To evaluate the correlations of the Bogoliubov

operatore  $\{b'_k\}$ , one can solve the Eq. (A6) for couples of operators  $\{b'_k\}$ , obtaining [84]

$$\begin{split} \langle b_k^{\dagger} b_k^{\prime} \rangle &= (1 - e^{-2\gamma t}) f(\omega_k^{\prime}) + e^{-2\gamma t} \langle b_k^{\dagger} b_k^{\prime} \rangle_0 \,, \\ \langle b_k^{\prime} b_q^{\prime} \rangle &= e^{i(\omega_k^{\prime} - \omega_q^{\prime})t - 2\gamma t} \langle b_k^{\prime} b_q^{\prime} \rangle_0 \,, \\ \langle b_k^{\prime} b_q^{\prime} \rangle &= e^{i(\omega_k^{\prime} + \omega_q^{\prime})t - 2\gamma t} \langle b_k^{\prime} b_q^{\prime} \rangle_0 \,, \\ \langle b_k^{\prime} b_q^{\prime} \rangle &= e^{-i(\omega_k^{\prime} + \omega_q^{\prime})t - 2\gamma t} \langle b_k^{\prime} b_q^{\prime} \rangle_0 \,. \end{split}$$
(A9)

The initial values  $\langle b_k^{\prime\dagger} b_q^{\prime} \rangle_0$  of the correlations is computed on the initial Gibbs state associated with  $w_i$ , and it can be obtained using the relations between  $\{b_k\}$ to  $\{b_k'\}$ . This relation can be formally derived as follows [84]. Introducing the fermionic Nambu field  $\mathbb{C}^{\dagger} =$  $(\hat{c}_1^{\dagger}, ..., \hat{c}_L^{\dagger}, \hat{c}_1, ..., \hat{c}_L)$ , their relations with the Bogoliubov operators  $\mathbb{B}(w)^{\dagger} = (\hat{b}_1^{\dagger}, ..., \hat{b}_L^{\dagger}, \hat{b}_1, ..., \hat{b}_L)$  corresponding to the Hamiltonian  $\hat{H}_{\mathrm{K}}(w)$  are obtained by a unitary transformation,  $\mathbb{C} = \mathbb{T}(w)\mathbb{B}(w)$ . See e.g. Ref. [84] for more

- I. Bloch, Quantum coherence and entanglement with ultracold atoms in optical lattices, Nature 453, 1016 (2008).
- [2] I. M. Georgescu, S. Ashhab, and F. Nori, Quantum simulation, Rev. Mod. Phys. 86, 153 (2014).
- [3] S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, Continuous quantum phase transitions, Rev. Mod. Phys. 69, 315 (1997).
- [4] S. Sachdev, *Quantum Phase Transitions*, (Cambridge University, Cambridge, England, 1999).
- [5] D. Rossini and E. Vicari, Coherent and dissipative dynamics at quantum phase transitions, Phys. Rep. 936, 1 (2021).
- [6] S. Bhattacharyya, S. Dasgupta, and A. Das, Signature of a continuous quantum phase transition in nonequilibrium energy absorption: Footprints of criticality on higher excited states, Sci. Rep. 5, 16490 (2015).
- [7] S. Roy, R. Moessner, and A. Das, Locating topological phase transitions using nonequilibrium signatures in local bulk observables, Phys. Rev. B 95, 041105(R) (2017).
- [8] M. Heyl, F. Pollmann, and B. Dora, Detecting Equilibrium and Dynamical Quantum Phase Transitions in Ising Chains via Out-of-Time-Ordered Correlators, Phys. Rev. Lett. **121**, 016801 (2018).
- [9] A. Pelissetto, D. Rossini, and E. Vicari, Dynamic finitesize scaling after a quench at quantum transitions, Phys. Rev. E 97, 052148 (2018).
- [10] P. Titum, J. T. Iosue, J. R. Garrison, A. V. Gorshkov, and Z.-X. Gong, Probing Ground-State Phase Transitions through Quench Dynamics, Phys. Rev. Lett. 123, 115701 (2019).
- [11] A. Pelissetto and E. Vicari, Scaling behaviors at quantum and classical first-order transitions, arXiv:2302.08238, to appear in 50 years of the renormalization group, dedicated to the memory of Michael E. Fisher, edited by A. Aharony, O. Entin-Wohlman, D. Huse, and L. Radzihovsky, World Scientific.

details. Therefore one can formally derive the relation between the Bogoliubov operators  $\hat{b}'_k$  and  $\hat{b}_k$ , corresponding to the Hamiltonian parameters  $w_i$  and w respectively, from the general relation

$$\mathbb{B}(w_2) = \mathbb{T}(w_2)^{\dagger} \mathbb{T}(w_1) \mathbb{B}(w_1).$$
 (A10)

Finally, to compute the time-dependent observables defined in Sec. II D, one can use the relations between the fermionic correlation functions associated with  $\hat{c}_x$  and those of the Bogoliubov operators  $\hat{b}_k$ , such as

$$C(x,y) = \sum_{k,q=1}^{L} \left[ A_{xk}^* A_{yq} \langle b_k^{\dagger} b_q \rangle + B_{xk}^* B_{yq} \langle b_k b_q^{\dagger} \rangle + A_{xk}^* B_{yq} \langle b_k^{\dagger} b_q^{\dagger} \rangle + B_{xk}^* A_{yq} \langle b_k b_q \rangle \right]$$
(A11)

where A and B are the matrices entering Eq. (9).

- [12] T. W. B. Kibble, Topology of Cosmic Strings and Domains, J. Phys. A 9, 1387 (1976).
- [13] T. W. B. Kibble, Some implications of a cosmological phase transition, Phys. Rep. 67, 183 (1980).
- [14] W. H. Zurek, Cosmological Experiments in Superfluid Helium?, Nature **317**, 505 (1985).
- [15] W. H. Zurek, Cosmological experiments in condensed matter systems, Phys. Rep. 276, 177 (1996).
- [16] W. H. Zurek, U. Dorner, and P. Zoller, Dynamics of a quantum phase transition, Phys. Rev. Lett. 95, 105701 (2005).
- [17] A. Polkovnikov and V. Gritsev, Breakdown of the adiabatic limit in low-dimensional gapless systems, Nature Phys. 4, 477 (2008).
- [18] J. Dziarmaga, Dynamics of a quantum phase transition and relaxation to a steady state, Adv. Phys. 59, 1063 (2010).
- [19] A. Dutta, G. Aeppli, B. K. Chakrabarti, U. Divakaran, T. F. Rosenbaum, and D. Sen, *Quantum phase tran*sitions in transverse field spin models: From statistical physics to quantum information, Cambridge University Press (2015).
- [20] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Colloquium: Nonequilibrium dynamics of closed interacting quantum systems, Rev. Mod. Phys. 83, 863 (2011).
- [21] A. Chandran, A. Erez, S. S. Gubser, and S. L. Sondhi, Kibble-Zurek problem: Universality and the scaling limit, Phys. Rev. B 86, 064304 (2012).
- [22] M. M. Rams, J. Dziarmaga, and W. H. Zurek, Symmetry Breaking Bias and the Dynamics of a Quantum Phase Transition, Phys. Rev. Lett. **123**, 130603 (2019).
- [23] F. De Franco and E. Vicari, Out-of-equilibrium finitesize scaling in generalized Kibble-Zurek protocols crossing quantum phase transitions in the presence of symmetry-breaking pertubations, Phys. Rev. B 107, 115175 (2023).
- [24] M. Campostrini, A. Pelissetto, and E. Vicari, Finite-

size scaling at quantum transitions, Phys. Rev. B 89, 094516 (2014).

- [25] T. Niemeijer, Some exact calculations on a chain of spins 1/2, Physica **36**, 377 (1967); Some exact calculations on a chain of spins 1/2 II, Physica **39**, 313 (1968).
- [26] E. Barouch, B. McCoy, and M. Dresden, Statistical mechanics of the XY model. I, Phys. Rev. A 2, 1075 (1970).
- [27] K. Sengupta, S. Powell and S. Sachdev, Quench dynamics across quantum critical points, Phys. Rev. A 69, 053616 (2004).
- [28] G. De Chiara, S. Montangero, P. Calabrese, and R. Fazio, Entanglement entropy dynamics of Heisenberg chains, J. Stat. Mech. P03001 (2006).
- [29] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, Spontaneous symmetry breaking in a quenched ferromagnetic spinor Bose–Einstein condensate, Nature 443, 312 (2006).
- [30] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, Relaxation in a Completely Integrable Many-Body Quantum System: An Ab Initio Study of the Dynamics of the Highly Excited States of 1D Lattice Hard-Core Bosons, Phys. Rev. Lett. 98, 050405 (2007).
- [31] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, Nature 452, 854 (2008).
- [32] M. Žnidarič, T. Prosen, and P. Prelovšek, Many-body localization in the Heisenberg XXZ magnet in a random field, Phys. Rev. B 77, 064426 (2008).
- [33] T. Prosen and M. Žnidarič, Matrix product simulations of non-equilibrium steady states of quantum spin chains, J. Stat. Mech. (2009) P02035.
- [34] F. Iglói and H. Rieger, Quantum Relaxation after a Quench in Systems with Boundaries, Phys. Rev. Lett. 106, 035701 (2011).
- [35] H. Rieger and F. Iglói, Semiclassical theory for quantum quenches in finite transverse Ising chains, Phys. Rev. B 84, 165117 (2011).
- [36] A. Gambassi and A. Silva, Large Deviations and Universality in Quantum Quenches, Phys. Rev. Lett. 109 250602 (2012).
- [37] P. Calabrese, F. H. Essler, and M. Fagotti, Quantum quench in the transverse field Ising chain: I. Time evolution of order parameter correlators, J. Stat. Mech. (2012) P07016.
- [38] P. Calabrese, F. H. Essler, and M. Fagotti, Quantum quenches in the transverse field Ising chain: II. Stationary state properties, J. Stat. Mech. (2012) P07022.
- [39] B. Blass, H. Rieger, and F. Iglói, Quantum relaxation and finite size effects in the XY chain in a transverse field after global quenches, Eur. Phys. Lett. 99, 30004 (2012).
- [40] J.-S. Caux and F. H. L. Essler, Time evolution of local observables after quenching to an integrable model, Phys. Rev. Lett. **110**, 257203 (2013).
- [41] M. Heyl, A. Polkovnikov, and S. Kehrein, Dynamical quantum phase transitions in the transverse-field Ising mode, Phys. Rev. Lett. **110**, 135704 (2013).
- [42] M. Fagotti, M. Collura, F. H. L. Essler, and P. Calabrese, Relaxation after quantum quenches in the spin-1/2 Heisenberg XXZ chain, Phys. Rev. B 89, 125101 (2014).
- [43] W. Fu, L.-Y. Hung, and S. Sachdev, Quantum quenches and competing orders, Phys. Rev. B 90, 024506 (2014).
- [44] J. Cardy, Thermalization and Revivals after a Quantum

Quench in Conformal Field Theory, Phys. Rev. Lett. **112**, 220401 (2014).

- [45] R. Nandkishore and D. A. Huse, Many body localization and thermalization in quantum statistical mechanics, Annu. Rev. Condens. Matter Phys. 6, 15 (2015).
- [46] A. Chiocchetta, M. Tavora, A. Gambassi, and A. Mitra, Short-time universal scaling and light-cone dynamics after a quench in an isolated quantum system in *d* spatial dimensions, Phys. Rev. B 94, 134311 (2016).
- [47] P. Calabrese and J. Cardy, Quantum quenches in 1 + 1 dimensional conformal field theories, J. Stat. Mech. (2016) 064003.
- [48] D. Bernard and B. Doyon, Conformal field theory out of equilibrium: a review, J. Stat. Mech. (2016) 064005.
- [49] E. Ilievski, M. Medenjak, T. Prosen, and L. Zadnik, Quasilocal charges in integrable lattice systems, J. Stat. Mech. (2016) 064008.
- [50] T. Langen, T. Gasenzer, and J. Schmiedmayer, Prethermalization and universal dynamics in near-integrable quantum systems, J. Stat. Mech. (2016) 064009.
- [51] R. Vasseur and J. E. Moore, Nonequilibrium quantum dynamics and transport: from integrability to manybody localization, J. Stat. Mech. (2016) 064010.
- [52] A. Nahum, J. Ruhman, S. Vijay, and J. Haah, Quantum Entanglement Growth under Random Unitary Dynamics, Phys. Rev. X 7, 031016 (2017).
- [53] M. Heyl, Dynamical quantum phase transitions: a review, Rep. Prog. Phys. 81, 054001 (2018).
- [54] D. Nigro, D. Rossini, and E. Vicari, Dynamic scaling of work fluctuations after quenches near quantum transitions, J. Stat. Mech. (2019) 023104.
- [55] D. Nigro, D. Rossini, and E. Vicari, Competing coherent and dissipative dynamics close to quantum criticality, Phys. Rev. A **100**, 052108 (2019); D. Rossini and E. Vicari, Scaling behavior of stationary states arising from dissipation at continuous quantum transitions, Phys. Rev. B **100**, 174303 (2019).
- [56] J. Surace, L. Tagliacozzo, and E. Tonni, Operator content of entanglement spectra after global quenches in the transverse field Ising chain, Phys. Rev. B 101, 241107(R) (2020).
- [57] D. Rossini and E. Vicari, Dynamics after quenches in one-dimensional quantum Ising-like systems, Phys. Rev. B 102, 054444 (2020).
- [58] O. A. Castro-Alvaredo, M. Lencsés, I. M. Szécsényi, and J. Viti, Entanglement oscillations near a quantum critical point, Phys. Rev. Lett. 124, 230601 (2020).
- [59] A. Pelissetto, D. Rossini, and E. Vicari, Scaling properties of the dynamics at first-order quantum transitions when boundary conditions favor one of the two phases, Phys. Rev. E **102**, 012143 (2020).
- [60] P. Ruggiero, P. Calabrese, L. Foini, and T. Giamarchi, Quenches in initially coupled Tomonaga-Luttinger liquids: a conformal field theory approach, SciPost Phys. 11, 055 (2021).
- [61] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, Quantum phase transition from a superfluid to a Mott insulator in a gas of ultracold atoms, Nature 415, 39 (2002).
- [62] T. Kinoshita, T. Wenger, and D. S. Weiss, A quantum Newton's cradle, Nature 440, 900 (2006).
- [63] S. Hofferberth, I. Lesanovsky, B. Fischer, T. Schumm, and J. Schmiedmayer, Non-equilibrium coherence dynamics in one-dimensional Bose gases, Nature 449, 324

(2007).

- [64] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, Probing the relaxation towards equilibrium in an isolated strongly correlated one-dimensional Bose gas, Nat. Phys. 8, 325 (2012).
- [65] M. Cheneau, P. Barmettler, D. Poletti, M. Endres, P. Scbauß, T. Fukuhara, C. Gross, I. Bloch, C. Kollath, and S. Kuhr, Light-cone-like spreading of correlations in a quantum many-body system, Nature 481, 484 (2012).
- [66] M. Gring, M. Kuhnert, T. Langen, T. Kitagawa, B. Rauer, M. Schreitl, I. Mazets, D. Adu Smith, E. Demler, and J. Schmiedmayer, Relaxation and Prethermalization in an Isolated Quantum System, Science **337**, 1318 (2012).
- [67] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider and I. Bloch, Observation of many-body localization of interacting fermions in a quasirandom optical lattice, Science 349, 842 (2015).
- [68] S. Braun, M. Friesdorf, S. S. Hodgman, M. Schreiber, J. P. Ronzheimer, A. Riera, M. del Rey, I. Bloch, J. Eisert, and U. Schneider, Emergence of coherence and the dynamics of quantum phase transitions, Proc. Natl. Acad. Sci. USA **112**, 3641 (2015).
- [69] Y.S. Patil, S. Chakram, and M. Vengalattore, Measurement-induced localization of an ultracold lattice gas, Phys. Rev. Lett. 115, 140402 (2015).
- [70] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated manybody system, Science **353**, 794 (2016).
- [71] J. Smith, A. Lee, P. Richerme, B. Neyenhuis, P. W. Hess, P. Hauke, M. Heyl, D. A. Huse, and C. Monroe, Many-body localization in a quantum simulator with programmable random disorder, Nat. Phys. 12, 907 (2016).
- [72] P. Bordia, H. Lüschen, U. Schneider, M. Knap, and I. Bloch, Periodically driving a many-body localized quantum system, Nat. Phys. 13, 460 (2017).
- [73] J. Zhang, G. Pagano, P. W. Hess, A. Kyprianidis, P. Becker, H. Kaplan, A. V. Gorshkov, Z.-X. Gong, and C. Monroe, Observation of a many- body dynamical phase transition with a 53-qubit quantum simulator, Nature 551, 601 (2017).
- [74] T. Tomita, S. Nakajima, I. Danshita, Y. Takasu, and Y. Takahashi, Observation of the Mott insulator to superfluid crossover of a driven-dissipative Bose-Hubbard system, Sci. Adv. 3, e1701513 (2017).
- [75] U. Mishra, H. Cheraghi, S. Mahdavifar, R. Jafari, and A. Akbari, Dynamical quantum correlations after sudden quenches, Phys. Rev. A 98, 052338 (2018).
- [76] R. Jafari, H. Johannesson, A. Langari, and M. A. Martin-Delgado, Quench dynamics and zero-energy modes: The case of the Creutz model, Phys. Rev. B 99, 054302 (2019).
- [77] T. Kohlert, S. Scherg, X. Li, H. P. Lüschen, S. Das Sarma, I. Bloch, and M. Aidelsburger, Observation of many-body localization in a one-dimensional system with a single-particle mobility edge, Phys. Rev. Lett. 122, 170403 (2019).
- [78] C. Maier, T. Brydges, P. Jurcevic, N. Trautmann, C. Hempel, B. P. Lanyon, P. Hauke, R. Blatt, and C. F. Roos, Environment-assisted quantum transport in a 10-

qubit network, Phys. Rev. Lett. 122, 050501 (2019).

- [79] G. Lindblad, On the generators of quantum dynamical semigroups, Commun. Math. Phys. 48, 119 (1976).
- [80] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of N-level systems, J. Math. Phys. 17, 821 (1976).
- [81] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, New York, 2002).
- [82] A. Rivas and S. F. Huelga, Open Quantum System: An Introduction (SpringerBriefs in Physics, Springer, 2012).
- [83] L. M. Sieberer, M. Buchhold, and S. Diehl, Keldysh field theory for driven open quantum systems, Rep. Prog. Phys. 79, 096001 (2016).
- [84] A. D'Abbruzzo and D. Rossini, Self-consistent microscopic derivation of Markovian master equations for open quadratic quantum systems, Phys. Rev. A 103, 052209 (2021).
- [85] A. Yu. Kitaev, Unpaired Majorana fermions in quantum wires, Phys. Usp. 44, 131 (2001).
- [86] G. Piccitto, M. Campisi, and D. Rossini The Ising critical quantum Otto engine, New Journal of Physics 24, 103023 (2022).
- [87] F. Tarantelli and E. Vicari, Out-of-equilibrium dynamics arising from slow round-trip variations of Hamiltonian parameters across quantum and classical critical points, Phys. Rev. B 105, 235124 (2022).
- [88] M. Schmitt, M. M. Rams, J. Dziarmaga, M. Heyl, and W. Zurek, Quantum phase transition dynamics in the two-dimensional transverse-field Ising model, Sci. Adv. 8, eabl6850 (2022).
- [89] R. Islam, E. E. Edwards, K. Kim, S. Korenblit, C. Noh, H. Carmichael, G.-D. Lin, L.-M. Duan, C.-C. Joseph Wang, J. K. Freericks, and C. Monroe, Onset of a quantum phase transition with a trapped ion quantum simulator, Nat. Commun. 2, 377 (2011).
- [90] S. Debnath, N. M. Linke, C. Figgatt, K. A. Landsman, K. Wright, and C. Monroe, Demonstration of a small programmable quantum computer with atomic qubits, Nature 536, 63 (2016).
- [91] J. Simon, W. S. Bakr, R. Ma, M. E. Tai, P. M. Preiss, and M. Greiner, Quantum simulation of antiferromagnetic spin chains in an optical lattice, Nature 472, 307 (2011).
- [92] H. Labuhn, D. Barredo, S. Ravets, S. de Leseleuc, T. Macri, T. Lahaye, and A. Browaeys, Tunable twodimensional arrays of single Rydberg atoms for realizing quantum Ising models, Nature 534, 667 (2016).
- [93] Y. Salathé, M. Mondal, M. Oppliger, J. Heinsoo, P. Kurpiers, A. Potocnik, A. Mezzacapo, U. Las Heras, L. Lamata, E. Solano, S. Filipp, and A. Wallraff, Digital quantum simulation of spin models with circuit quantum electrodynamics, Phys. Rev. X 5, 021027 (2015).
- [94] A. Cervera-Lierta, Exact Ising model simulation on a quantum computer, Quantum 2, 114 (2018).
- [95] P. Pfeuty, The one-dimensional Ising model with a transverse field, Ann. Phys. 57, 79 (1970).
- [96] J.-P. Blaizot and G. Ripka, Quantum Theory of Finite Systems. (MIT Press, 1986).
- [97] A. Bácsi and B. Dóra, Kibble-Zurek scaling due to environment temperature quench in the transverse field Ising model, Scientific Rep. 13, 1 (2023).
- [98] M. Caselle, M. Hasenbusch, A. Pelissetto, and E. Vicari, Irrelevant operators in the two-dimensional Ising model,

J. Phys. A **35**, 4861 (2002).

- [99] A. Pelissetto and E. Vicari, Critical phenomena and renormalization group theory, Phys. Rep. 368, 549 (2002).
- [100] D. Rossini and E. Vicari, Dynamic Kibble-Zurek scaling framework for open dissipative many-body systems crossing quantum transitions, Phys. Rev. Research 2, 023611 (2020).
- [101] F. Tarantelli and E. Vicari, Quantum critical systems with dissipative boundaries, Phys. Rev. B 104, 075140 (2021).
- [102] A. Franchi and F. Tarantelli, Liouvillian gap and out-ofequilibrium dynamics of a sunburst Kitaev ring: from local to uniform dissipation, arXiv:2303.04207.