Thermal Leptophilic Light Vector Dark Matter with Spinor Mediator and Muon (g-2) Anomaly

Seyed Yaser Ayazi^{*1} and Ahmad Mohamadnejad^{†2}

¹Physics Department, Semnan University, P.O. Box. 35131-19111, Semnan, Iran ²Department of Physics, Lorestan University, khorramabad, Iran

February 21, 2023

Abstract

Inspired by the recently new measurement of $(g-2)_{\mu}$ at FermiLab and reported upper bound for electron-dark matter (DM) recoil by the XENON1T collaboration, we revisited phenomenology of a light MeV scale vector dark matter in a leptophilic extension of standard model while a new spinor field plays the role of mediator. A viable parameter space is considered to discuss the possibility of light dark matter relic density as well as anomalous magnetic moment of the muon. We study DM-electron direct detection and cosmological bounds on the parameters space of the model. It is shown that although new bound of $(g-2)_{\mu}$ anomaly greatly confines the parametric space of the model, the thermal light dark matter can exist for $M_{\rm DM} \sim 10^{-1} - 10^1$ GeV.

1 Introduction

As it is well-known, the existence of DM on many scales in the universe has been widely established, but its nature remains an unsolved problem. Traditional DM candidates, in many beyond standard model (BSM) scenarios are weakly interacting massive particle (WIMP) with mass close to the electroweak scale and weak couplings with Standard Model (SM) particles. However the lack of any signal in direct, indirect detection experiments and search at the LHC, has opened the door to wide spreading DM models possibility covering small value ranges in DM masses and interaction coupling strengths.

Furthermore, physics of BSM may leave affect in low-energy precise measurements via possible deviations between the SM and experimental evidences. Some of these deviations include the LHCb [1] and the muon anomalous magnetic moment (AMM) $a_{\mu} = (g_{\mu} - 2)/2$ which was recently reported by FermiLab experiment (3.3 σ) [2] challenging lepton universality. Combining this result for AMM with the earlier measurement at the Brookhaven National Laboratory, this now amounts to a 4.2 σ deviation from the SM prediction [3]. This result leaves an open window to new particles which can contribute to muon g - 2 through direct coupling to the muon. Many models related to DM candidates and $(g - 2)_{\mu}$ anomaly have been proposed, including axion-like particle (ALP) explanation [4–7], SUSY-DM explanation [8–11], two Higgs doublet

^{*}syaser.ayazi@semnan.ac.ir

[†]mohamadnejad.a@lu.ac.ir

models (2HDM) plus singlet explanation [12], and others [13–27]. Also it should be noted that if we compare to the latest QCD lattice calculations this decrease to about 1.4 σ [28]. In order to clarify the origin of this difference, a new experiment aiming to measure the contribution of hadronic corrections is being prepared at CERN [29]. One should also note that the drastic change in Hadronic Vacuum Polarization (HVP) leads to tensions elsewhere [30–32].

The primary purpose of direct detection experiments is the search for thermal relic particles (WIMPs) with GeV-scale mass. However, recently, there has been significant interest in models of DM in which DM particle has a mass $m \leq \mathcal{O}(\text{GeV})$. This is appealing because such models evade nearly all constraints on DM imposed by nucleon-DM direct detection experiments, since direct detection experiments are insensitive to the small recoil energies characteristic of $m \leq \mathcal{O}(\text{GeV})$. An alternative scenario in direct search is the interaction of DM particles exclusively with the SM leptons and possibly having loop suppressed interactions with nucleons. However in this case, DM still obtain its relic abundance from Freez-out mechanism.

In this light, direct detection strategies are proposed for DM particles with KeV to MeV mass. In this range, DM scattering with electrons can cause single-electron ionization signals, which are detectable with current technology. Ultraviolet photons, individual ions, and heat are interesting alternative signals. For example, the light DM regime (1 MeV to 10 GeV) where DM interacts with atomic electrons via recoiling is now significantly constrained by recent data from the XENON1T [33], DarkSide50 [34] and SENSEI [35] experiments.

The goal of this paper is to broaden the scope of BSM physics. We propose light vector DM candidates with mass regime (1 MeV to 10 GeV). The interaction of DM with SM is leptophilic and DM particles only interact with SM leptons via spinor mediators. Scattering of vector DM from electron could be consistent with XENON1T [33] experiment bounds for DM-electron direct detection. The leptophilic models are particularly interesting when we look at the anomalous magnetic moment of the leptons. In this way, we attempt to find out parameter space of the model to explain the newest muon magnetic moment anomaly.

The outline of this paper is as follows: In Section 2, we introduce framework for a light vector DM. In Section 3, we study anomhous magnetic of muon in context of the model. We consider relic density and DM-electron recoil bounds on parameters of the model in section 4 and 5. The Section 6 contains the influence of various experimental constraints on the model parameters. Finally, we present combined result in last section.

2 The Model

Here, we present a vector DM model, which extends the SM with massive spinor mediators which couples to vector dark matter (VDM) particles and SM leptons. The only renormalizable interaction is taken as follows:

$$\mathcal{L} \supset \sum_{\ell=e,\mu,\tau} X_{\mu} \overline{\psi}_{l} \gamma^{\mu} [g_{s} + g_{p} \gamma^{5}] \ell + \text{H.C.}, \qquad (2.1)$$

where X_{μ} is the VDM candidate, and $\psi_l s$ are (Dirac) spinor mediators. Generally, the interaction terms $X_{\mu}\overline{\psi}_l\gamma^{\mu}[g_{s\psi}+g_{p\psi}\gamma^5]\psi_l$ and $X_{\mu}\overline{l}\gamma^{\mu}[g_{sl}+g_{pl}\gamma^5]l$ can also exist. However, these interactions

will lead to X_{μ} being unstable. To avoid such interactions we impose a Z_2 discrete symmetry under which both X_{μ} and ψ_l are odd but all SM fields are even. In our analysis, we have assumed universal couplings between the mediators and the SM leptons. For simplicity, we have eluded mixing between ψ_l s and lepton generations. If different lepton flavors couple with a same ψ and X_{μ} , then at one-loop level we should face the charged lepton flavor violation (CLFV) problem (for a review on the muon anomalous magnetic moment and the quest for lepton flavor violation see [36]). In such a scenario, at one-loop level, we should have muon to electron transition process $\mu \to e + \gamma$ which has been stringently constrained. In (2.1), to avoid CLFV, we have considered different ψ_l s for each lepton flavor. However, to keep the model minimal, we suppose all spinor mediators have the same mass. In order to avoid DM decay, we suppose $M_{\psi} > M_X$. By considering this constraint X_{μ} will be stable and can serve as DM. Since X_{μ} is neutral with no electric charge, for spin 1/2 mediators, the spinor ψ_l s have positive electric charges (and therefore couple to photon) that is equal but opposite to the charged leptons. Therefore, X_{μ} and neutrinos cannot couple together via charged spin 1/2 mediators. VDM stability commands that no tree level mixing between the Z-boson and X_{μ} is allowed. Therefore in our model, we have 4 independent parameters M_X , M_{ψ} , g_s and g_p . Finally, we emphasize that our approach in this paper, is quite phenomenological. We have not considered gauge invariance of Lagrangians under some gauge group, and the model is not UV-complete. For example, we supposed that ψ_{ls} are $SU(2)_{Weak}$ singlet while have $U(1)_{EM}$ charge; in constructing a gauge invariant model such assumption should be considered more carefully.

3 Anomalous Magnetic Moment (AMM) of the Muon

As it is well-known, there are some discrepancies between SM predictions and measurements in low energy experiments. One long-standing discrepancy is the anomalous magnetic dipole moment of muon:

$$a_{\mu} = (g_{\mu} - 2)/2. \tag{3.1}$$

Comparisons with experimental measurements a_{exp}^{l} results in studies of the magnetic moments of leptons being a powerful indirect search of new physics. The SM prediction of the anomalous magnetic moment is determined from the sum of all sectors of the SM which include the QED contributions, the electro-weak contributions, the hadronic vacuum polarization contributions and hadronic light-by-light scattering contributions. The uncertainty of a_{SM} is completely dominated by the hadronic contributions due to the non-perturbative nature of the low energy strong interaction. It is estimated from experimental measurements of the ratio of hadronic to muonic cross sections in electron-positron collisions [37–58]. The combined results of muon anomalous magnetic moment in two experiments at FermiLab (FNAL E989) [2], and Brookhaven National Laboratory (LabBNL E821) [3] yield overall deviation from the SM central value:

$$\Delta a_{\mu} = a_{\mu}^{exp} - a_{\mu}^{\rm SM} = (25.1 \pm 5.9) \times 10^{-10}, \qquad (3.2)$$

which corresponding to a significance of 4.2 σ .

In light of the results, we study constraints on the parameter space of the model. Since in our model, VDM particles interact with the SM leptons via a massive mediator, a significant effect on the anomalous magnetic moment of the leptons is expected.

In the present work, as we mentioned before, we consider universality for all interactions of leptons. Therefore, we only consider the magnetic moment of the muon and ignore weaker constraints on the AMMs of tau and electron. The model contribution to muon magnetic moment arises from the one-loop mediated by VDMs. The analytical expression for Δa^{ψ}_{μ} is [59]- [60]:

$$\Delta a^{\psi}_{\mu} = \frac{1}{8\pi^2} \frac{m^2_{\mu}}{M^2_X} \int \frac{g_s^2 P_s(x) + g_p^2 P_p(x)}{(1-x)(1-\lambda^2 x) + \epsilon^2 \lambda^2 x} dx$$
(3.3)

where

$$P_s(x) = 2x(1-x)(x-2(1-\epsilon)^2) + \lambda^2(1-\epsilon)^2 x^2(1+\epsilon-x)$$
$$P_p(x) = 2x^2(1+x+2\epsilon) + \lambda^2(1-\epsilon)^2 x(1-x)(x-\epsilon)$$

and $\epsilon = M_{\psi}/m_{\mu}$, $\lambda = m_{\mu}/M_X$ (m_{μ} is the muon mass, M_{ψ} and M_X are spinor mediator and DM mass). Therefore the contribution of a generic spinor mediator to the muon anomalous magnetic moment is given by:

$$\Delta a_{\mu}^{\psi} = \left(\frac{m_{\mu}}{2\pi M_X}\right)^2 \left\{ \left[\frac{M_{\psi}}{m_{\mu}} - \frac{2}{3}\right] g_s^2 + \left[-\frac{M_{\psi}}{m_{\mu}} - \frac{2}{3}\right] g_p^2 \right\},\tag{3.4}$$

where g_s and g_p are couplings of the SM leptons with new fields in accordance with interaction terms of Eq. (2.1). Figs. 1-a and b show allowed region in $M_X - M_{\psi}$ plane for random values of couplings which is consistent with BNL E821 and FNAL E989 experiments. As it is seen in figures, for masses of spinor and DM between a few hundred MeV to 100 GeV, the contribution of the model can explain a_{μ} anomaly. Note that in each plot, we turn on both couplings (g_s and g_p).

Before the next section, we comment on the electron g-2 result as a comparison. There is a 2.4 σ discrepancy between the theoretical prediction [61] and the existing experimental measurement [62,63] of the electron anomalous magnetic moment,

$$\Delta a_e = a_e^{exp} - a_e^{\rm SM} = (-87 \pm 36) \times 10^{-14}.$$
(3.5)

Given the negative sign of the above relation, to explain both electron and muon g-2 in this model, according to (3.4), one should consider different couplings for electron and muon generations. Otherwise, $\Delta a_{\mu}^{\psi} > 0$ leads to $\Delta a_{e}^{\psi} > 0$. However, considering the significance of 4.2 σ of muon anomalous magnetic moment and ignoring 2.4 σ electron g-2 anomaly, we continue to assume universal couplings between the mediators and the SM leptons.

4 Relic Density of DM

Cosmological observations point toward the existence of DM, whose cosmological relic abundance is approximately a fourth of the energy budget of the Universe. According to the recent measurements from the PLANCK [67] collaboration DM relic density is:

$$\Omega_{DM}h^2 = 0.120 \pm 0.001, \tag{4.1}$$



Figure 1: Scatter-points consistent with AMM of the Muon. We scanned over all free parameters of the model. Note that, the parameter space could be limited by other experiments, such as BaBar, Belle II and LEP II. However, maybe the case of a small mass difference of $M_{\psi}-M_X =$ $\delta > 0$ with $\delta/M_{\psi} \ll 1$ is still allowed by experiments (with large missing energy plus soft electrons/muons), like the case of supersymmetry. In addition, if $2M_{\psi}$ is lighter than the tau lepton m_{τ} , the decay spectrum and width of tau lepton will be changed. Thus, the parameter space left are smaller than that in this figure. To see how these experiments limit the parameter space see, e.g., [64–66]



Figure 2: Feynman diagrams for DM annihilation into (a) leptons and (b) spinor mediators. For every t channel there is associated u channel not depicted here. DM coannihilation is depicted in (c).

where $h \sim 0.7$ is the Hubble expansion rate at present times in units of 100 (km/s)/Mpc.

We suppose the mechanism that generates DM cosmological relic density is thermal freezeout described by a Boltzmann equation of the form

$$\frac{dn_{\chi}}{dt} + 3Hn_{\chi} = -\langle \sigma_{eff} | v_{rel} | \rangle (n_{\chi}^2 - n_{\chi,eq}^2), \qquad (4.2)$$

where n_{χ} is the number density of vector DM, H is the Hubble parameter, and $\langle \sigma_{eff} | v_{rel} | \rangle$ is the thermally averaged of effective cross section including both DM annihilation and coannihilation channels [68] (see figure 2).

The constraint from DM relic density on the vector DM have been determined by implementing the model in the numerical package MicrOMEGAs [69]. The allowed parameter space corresponding to this constraint is depicted in figure 3.

Note that the annihilation channel in Fig. 2 (b) is phase-space suppressed and the main



Figure 3: Dark matter relic density consistent with Planck data. We scanned over all free parameters of the model

channel is Fig. 2 (a). The s-wave annihilation $X X \to e^+ e^-$ is restricted by the CMB observation (see, e.g. [70]) for the mass of dark matter about < 10 GeV, and the AMS-02 observation (see, e.g. [71,72]) for the mass of dark matter from a few GeV to 100 GeV. There is nearly no parameter space left with joint constraints. Thus, the relic density of X equal to 0.12 seems to be in tension with CMB and AMS-02 observations. To resolve this issue, we consider coannihilation effect with $M_{\psi} - M_X = \delta > 0$ and $\frac{\delta}{M_{\psi}} \ll 1$. The particles X and ψ_l can be in thermal equilibrium for a while in the early universe via processes like Fig. 2 (b), and the abundance of X will be reduced by processes such as Fig. 2 (c). However, in our combined analysis, Sec. 7, we see that the relic density of X should be smaller than 0.12. Therefore, we consider $\Omega_X h^2 < 0.12$ as DM relic density constraint. This means X cannot be the only DM component.

5 Direct Detection and Electron recoil

Let us now turn our attention to the direct detection possibilities of DM in the framework. As it is mentioned, we consider the hypothesis that the VDM particle X_{μ} only couples directly to leptons in particular the electrons but not to quarks. For this reason, DM-nucleon interaction is absent at tree level. However, DM-electron elastic scattering of spin-independent type is feasible at tree level. In the following, we ignore the loop suppressed DM-nucleon interaction. In comparison to underground experiments focused on neutrino detection, that are typically sensitive to energy depositions above a few hundred keV, Xenon-based dark matter detectors provide the leading sensitivity for electronic energy depositions of 100 keV and below. In our model the relevant interaction is described by the Lagrangian 2.1. Electron recoil can occur corresponding to Feynman diagrams in Fig. 4. Therefore, in non-relativistic limit the elastic scattering cross section of the DM-electron has following form [73]:

$$\sigma_{DM-e} \approx \frac{g_s^4 \mu_{eX}^2}{2\pi M_X^2 M_\psi^2},\tag{5.1}$$

where μ_{eX} is the DM-electron reduced mass. The cross section that includes g_p is zero due to the odd number of γ^5 in the trace. So far, in direct detection experiments there is found no evidence of DM-electron elastic scattering. Nevertheless several experiments have set upper bounds on this cross section: XENON1T [33], DarkSide50 [34] and SENSEI [35]. We consider upper bound from the XENON1T experiment to search for DM interacting with electrons [33].



Figure 4: DM-electron Feynman diagram

6 Additional constraints on the leptophilic VDM interaction

Thus far in this paper, we have included discussion of the relic density, direct detection and $(g-2)_{\mu}$ constraints. In this section, we look at the various experimental constraints on the muon-VDM interaction in the model. Two kinds of these experimental constraints are as follows:

• Big Bang Nucleosynthesis (BBN) constrains the DM physics by three generic ways, the number of the excited relativistic degrees of freedom, energy injection due to annihilation or decay of heavy particles, and particle catalysis of nuclear reactions. For the first case, traditionally, this procedure is interpreted in terms of a bound on the effective number of neutrino. Note that in our model, X_{μ} is neutral and cannot couple neutrino via charged spin 1/2 mediators. In a more general way, this bound can be translated into a limit on dark radiation. In [74], it have been shown that BBN constraint as a limit on dark radiation which is massless or nearly-massless degrees of freedom that have the expected scaling as the Universe expands $\rho_{dr} \approx a^{-4}$. They assume that the additional relativistic component does not exchange energy with any SM species. This viewpoint can then be easily applied to the thermally decoupled extra degrees of freedom. For this reason, in our model in which new particles are not massless, the parameter space is not sensitive to the number of the excited relativistic degrees of freedom. Nevertheless, constraints on $\langle \sigma v \rangle$ from electromagnetic energy injection are proportional to temperature and dark matter mass. In [75], it was shown that BBN procedure rules out DM annihilation to e^+e^- for the DM mass region 30 MeV $\lesssim m_{\chi} \lesssim 77$ MeV.



Figure 5: Scatter-points consistent with Anomalous Magnetic Moment (AMM) of the Muon and DM relic density (RD).

• The observation of cosmic charge leptons have reported in wide range of the data (10 GeV-10 TeV); Fermi-LAT has reported a measurement of the CR electron-positron spectrum from 7 GeV to 2 TeV [76]. The PAMELA satellite experiment [77] released an abundance of the positron in the CR energy range of 15 - 100 GeV. Also the results of a CR electron-positron spectrum, between 10 GeV and 3TeV, have been reported based upon observations with CALET [78]. The DAMPE collaboration [79,80] has been presented a measurements of the electron-positron spectrum in the energy range 25 GeV to 4.6 TeV. These observations are both considered as signs of standard astrophysical sources or DM annihilation. In our model, DM has mass smaller than 10 GeV then the parameter space is not sensitive to these observations.

7 Combined Results

In this section, we present a combined analysis of all observational constraints described in the previous sections. The constraints discussed are summarized by Fig. 5. As it is seen, for narrow region in parameter space, all constraints are satisfied. Remarkable point is for light DM (\mathcal{O} (MeV)), implying we can have thermal DM which explain anomalous magnetic moment of the muon.

To obtain the parameter space consistent with all constraints, i.e., muon anomalous magnetic moment, DM relic density, and DM-electron direct detection constraints, first we solve Eq. (3.4) with respect to g_s :

$$g_s = \sqrt{\frac{12\pi^2 M_X^2}{m_\mu (3M_\psi - 2m_\mu)}} \Delta a_\mu^\psi + \frac{3M_\psi + 2m_\mu}{3M_\psi - 2m_\mu} g_p^2 \tag{7.1}$$

Since Δa^{ψ}_{μ} is positive, $M_{\psi} > \frac{2}{3}m_{\mu}$, and therefore $g_s > g_p$. Now, to obtain the parameter space



Figure 6: Scatter-points consistent with AMM of the Muon and DM relic density (RD). Some points will be excluded by XENON1T upper bound on DM-electron cross section. Points that are located in a narrow red area are excluded by BBN.

consistent with DM relic density ($\Omega_X h^2 < 0.12$), we scan over random points for M_{ψ} , g_p with the extra condition $M_{\psi} - m_e - \Delta < M_X < M_{\psi} - m_e$ (where M_X is chosen randomly and we put $\Delta = 10^{-6}$) and for g_s we use Eq. (7.1) where Δa^{ψ}_{μ} is chosen randomly in a domain given by Eq. (3.2). In this way, we obtain the parameter space which gives the correct value of both DM relic density and muon anomalous magnetic momont. Finally we keep the points that are also consistent with direct detection constraint. The result of our combined analysis is depicted in Fig. 6.

8 Conclusions

SM has achieved great success for its high accuracy to describe electroweak and strong interaction. However, there remains problems such as DM which SM can not explain. In addition, recent results about muon anomalous magnetic moment brings new challenges to the SM. The 4.2 σ discrepancy between experiment and SM prediction seems to indicate new physics behind $(g_{\mu} - 2)$ anomaly and gives possible hints to the BSM physics. In light of these results, we consider a vector DM model with a leptophilic spinor mediator coupled to the SM charged leptons. From phenomenological point of view, we probed the parameter space of the model for light DM \mathcal{O} (MeV) and sub-GeV spinor mediator. To conclude, the anomalies associated to the muon physics can be successfully addressed in light VDM regime. If the anomalies are confirmed by forthcoming experimental analyses, our results show that the model can be surveyed by DM-electron direct detection.

References

- LHCB collaboration, Test of lepton universality in beauty-quark decays, Nature Phys. 18 (2022) 277 [arXiv:2103.11769] [INSPIRE].
- MUON G-2 collaboration, Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm, Phys. Rev. Lett. 126 (2021) 141801 [arXiv:2104.03281] [INSPIRE].
- [3] MUON G-2 collaboration, Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL, Phys. Rev. D 73 (2006) 072003 [arXiv:hep-ex/0602035]
 [INSPIRE].
- [4] P. deNiverville, H.-S. Lee and M.-S. Seo, Implications of the dark axion portal for the muon g-2, B factories, fixed target neutrino experiments, and beam dumps, Phys. Rev. D 98 (2018) 115011 [arXiv:1806.00757] [INSPIRE].
- [5] C.-W. Chiang, M. Takeuchi, P.-Y. Tseng and T. T. Yanagida, Muon g 2 and rare top decays in up-type specific variant axion models, Phys. Rev. D 98 (2018) 095020
 [arXiv:1807.00593] [INSPIRE].
- [6] W.-Y. Keung, D. Marfatia and P.-Y. Tseng, Axion-Like Particles, Two-Higgs-Doublet Models, Leptoquarks, and the Electron and Muon g - 2, LHEP 2021 (2021) 209
 [arXiv:2104.03341] [INSPIRE].
- [7] B. Bhattacharya, A. Datta, D. Marfatia, S. Nandi and J. Waite, Axion-like particles resolve the $B \rightarrow \pi K$ and g 2 anomalies, Phys. Rev. D 104 (2021) 051701 [arXiv:2104.03947] [INSPIRE].
- [8] Q. Shafi and C. S. Ün, Sparticle Spectroscopy at LHC-Run3 and LSP Dark Matter in light of Muon g-2, arXiv:2107.04563 [INSPIRE].
- [9] A. K. Forster and S. F. King, Muon g-2, dark matter and the Higgs mass in no-scale supergravity, Nucl. Phys. B 976 (2022) 115700 [arXiv:2109.10802] [INSPIRE].
- [10] M. Frank, Y. Hiçyılmaz, S. Mondal, O. Özdal and C. S. Un, Electron and muon magnetic moments and implications for dark matter and model characterisation in non-universal U(1)' supersymmetric models, JHEP 10 (2021) 063 [arXiv:2107.04116] [INSPIRE].
- [11] J. S. Kim, D. E. Lopez-Fogliani, A. D. Perez and R. R. de Austri, The new (g-2)μ and right-handed sneutrino dark matter, Nucl. Phys. B 974 (2022) 115637
 [arXiv:2107.02285] [INSPIRE].
- [12] G. Arcadi, A. S. de Jesus, T. B. de Melo, F. S. Queiroz and Y. S. Villamizar, A 2HDM for the g-2 and dark matter, Nucl. Phys. B 982 (2022) 115882 [arXiv:2104.04456]
 [INSPIRE].
- [13] G. Cacciapaglia, C. Cot and F. Sannino, Naturalness of lepton non-universality and muon g-2, Phys. Lett. B 825 (2022) 136864 [arXiv:2104.08818] [INSPIRE].

- [14] L.-B. Jia, A dark leptophilic scalar with the updated muon g 2 anomaly, Eur. Phys. J. C 82 (2022) 321 [arXiv:2105.13805] [INSPIRE].
- [15] S. Singirala, S. Sahoo and R. Mohanta, Light dark matter, rare B decays with missing energy in Lμ-Lτ model with a scalar leptoquark, Phys. Rev. D 105 (2022) 015033
 [arXiv:2106.03735] [INSPIRE].
- [16] X. Qi, A. Yang, W. Liu and H. Sun, Scalar dark matter and muon g-2 in a model *, Chin. Phys. C 46 (2022) 083102 [arXiv:2106.14134] [INSPIRE].
- [17] NA64 collaboration, Probing the explanation of the muon (g-2) anomaly and thermal light dark matter with the semi-visible dark photon channel, Eur. Phys. J. C 81 (2021) 959
 [arXiv:2107.02021] [INSPIRE].
- [18] B. D. Sáez and K. Ghorbani, Singlet scalars as dark matter and the muon (g-2) anomaly, Phys. Lett. B 823 (2021) 136750 [arXiv:2107.08945] [INSPIRE].
- [19] D. Borah, M. Dutta, S. Mahapatra and N. Sahu, Lepton anomalous magnetic moment with singlet-doublet fermion dark matter in a scotogenic U(1)Lμ-Lτ model, Phys. Rev. D 105 (2022) 015029 [arXiv:2109.02699] [INSPIRE].
- [20] T. Hapitas, D. Tuckler and Y. Zhang, General kinetic mixing in gauged U(1)Lμ-Lτ model for muon g-2 and dark matter, Phys. Rev. D 105 (2022) 016014 [arXiv:2108.12440]
 [INSPIRE].
- [21] T. A. Chowdhury and S. Saad, Non-Abelian vector dark matter and lepton g-2, JCAP 10 (2021) 014 [arXiv:2107.11863] [INSPIRE].
- [22] K. Ghorbani, Light vector dark matter with scalar mediator and muon g-2 anomaly, Phys. Rev. D 104 (2021) 115008 [arXiv:2104.13810] [INSPIRE].
- [23] D. Borah, M. Dutta, S. Mahapatra and N. Sahu, Muon (g-2) and XENON1T excess with boosted dark matter in Lμ Lτ model, Phys. Lett. B 820 (2021) 136577
 [arXiv:2104.05656] [INSPIRE].
- [24] C.-T. Lu, R. Ramos and Y.-L. S. Tsai, Shedding light on dark matter with recent muon (g 2) and Higgs exotic decay measurements, JHEP 08 (2021) 073 [arXiv:2104.04503]
 [INSPIRE].
- [25] Y. Bai and J. Berger, Muon g 2 in Lepton Portal Dark Matter, arXiv:2104.03301
 [INSPIRE].
- [26] P. Das, M. K. Das and N. Khan, The FIMP-WIMP dark matter in the extended singlet scalar model, Nucl. Phys. B 975 (2022) 115677 [arXiv:2104.03271] [INSPIRE].
- [27] K. Kowalska and E. M. Sessolo, Minimal models for g-2 and dark matter confront asymptotic safety, Phys. Rev. D 103 (2021) 115032 [arXiv:2012.15200] [INSPIRE].

- [28] S. Borsanyi et al., Leading hadronic contribution to the muon magnetic moment from lattice QCD, Nature **593** (2021) 51 [arXiv:2002.12347] [INSPIRE].
- [29] MUONE collaboration, Status of the MUonE experiment, PoS ICHEP2020 (2021) 223
 [arXiv:2012.07016] [INSPIRE].
- [30] A. Crivellin, M. Hoferichter, C. A. Manzari and M. Montull, Hadronic Vacuum Polarization: (g - 2)_μ versus Global Electroweak Fits, Phys. Rev. Lett. **125** (2020) 091801
 [arXiv:2003.04886] [INSPIRE].
- [31] A. Keshavarzi, W. J. Marciano, M. Passera and A. Sirlin, Muon g 2 and $\Delta \alpha$ connection, Phys. Rev. D 102 (2020) 033002 [arXiv:2006.12666] [INSPIRE].
- [32] G. Colangelo, M. Hoferichter and P. Stoffer, Constraints on the two-pion contribution to hadronic vacuum polarization, Phys. Lett. B 814 (2021) 136073 [arXiv:2010.07943]
 [INSPIRE].
- [33] XENON collaboration, Excess electronic recoil events in XENON1T, Phys. Rev. D 102 (2020) 072004 [arXiv:2006.09721] [INSPIRE].
- [34] DARKSIDE collaboration, Constraints on Sub-GeV Dark-Matter-Electron Scattering from the DarkSide-50 Experiment, Phys. Rev. Lett. 121 (2018) 111303 [arXiv:1802.06998]
 [INSPIRE].
- [35] SENSEI collaboration, SENSEI: Direct-Detection Results on sub-GeV Dark Matter from a New Skipper-CCD, Phys. Rev. Lett. 125 (2020) 171802 [arXiv:2004.11378] [INSPIRE].
- [36] M. Lindner, M. Platscher and F. S. Queiroz, A Call for New Physics : The Muon Anomalous Magnetic Moment and Lepton Flavor Violation, Phys. Rept. 731 (2018) 1 [arXiv:1610.06587] [INSPIRE].
- [37] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, Reevaluation of the Hadronic Contributions to the Muon g-2 and to alpha(MZ), Eur. Phys. J. C 71 (2011) 1515
 [arXiv:1010.4180] [INSPIRE].
- [38] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, Complete Tenth-Order QED Contribution to the Muon g-2, Phys. Rev. Lett. 109 (2012) 111808 [arXiv:1205.5370]
 [INSPIRE].
- [39] T. Aoyama, T. Kinoshita and M. Nio, Theory of the Anomalous Magnetic Moment of the Electron, Atoms 7 (2019) 28 [INSPIRE].
- [40] A. Czarnecki, W. J. Marciano and A. Vainshtein, Refinements in electroweak contributions to the muon anomalous magnetic moment, Phys. Rev. D 67 (2003) 073006
 [arXiv:hep-ph/0212229] [INSPIRE].

- [41] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, The electroweak contributions to (g-2)_μ after the Higgs boson mass measurement, Phys. Rev. D 88 (2013) 053005
 [arXiv:1306.5546] [INSPIRE].
- [42] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon g - 2 and α(m²_Z) using newest hadronic cross-section data, Eur. Phys. J. C 77 (2017) 827 [arXiv:1706.09436] [INSPIRE].
- [43] A. Keshavarzi, D. Nomura and T. Teubner, Muon g 2 and $\alpha(M_Z^2)$: a new data-based analysis, Phys. Rev. D 97 (2018) 114025 [arXiv:1802.02995] [INSPIRE].
- [44] G. Colangelo, M. Hoferichter and P. Stoffer, Two-pion contribution to hadronic vacuum polarization, JHEP 02 (2019) 006 [arXiv:1810.00007] [INSPIRE].
- [45] M. Hoferichter, B.-L. Hoid and B. Kubis, Three-pion contribution to hadronic vacuum polarization, JHEP 08 (2019) 137 [arXiv:1907.01556] [INSPIRE].
- [46] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to α(m²_z), Eur. Phys. J. C 80 (2020) 241 [arXiv:1908.00921] [INSPIRE].
- [47] A. Keshavarzi, D. Nomura and T. Teubner, g 2 of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium, Phys. Rev. D 101 (2020) 014029 [arXiv:1911.00367] [INSPIRE].
- [48] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order, Phys. Lett. B 734 (2014) 144 [arXiv:1403.6400] [INSPIRE].
- [49] K. Melnikov and A. Vainshtein, Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment revisited, Phys. Rev. D 70 (2004) 113006
 [arXiv:hep-ph/0312226] [INSPIRE].
- [50] P. Masjuan and P. Sanchez-Puertas, Pseudoscalar-pole contribution to the $(g_{\mu} 2)$: a rational approach, Phys. Rev. D **95** (2017) 054026 [arXiv:1701.05829] [INSPIRE].
- [51] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, Dispersion relation for hadronic light-by-light scattering: two-pion contributions, JHEP 04 (2017) 161 [arXiv:1702.07347]
 [INSPIRE].
- [52] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S. P. Schneider, Dispersion relation for hadronic light-by-light scattering: pion pole, JHEP 10 (2018) 141 [arXiv:1808.04823]
 [INSPIRE].
- [53] A. Gérardin, H. B. Meyer and A. Nyffeler, Lattice calculation of the pion transition form factor with $N_f = 2 + 1$ Wilson quarks, Phys. Rev. D 100 (2019) 034520 [arXiv:1903.09471] [INSPIRE].

- [54] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, Short-distance constraints for the HLbL contribution to the muon anomalous magnetic moment, Phys. Lett. B 798 (2019) 134994 [arXiv:1908.03331] [INSPIRE].
- [55] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, Longitudinal short-distance constraints for the hadronic light-by-light contribution to $(g-2)_{\mu}$ with large-N_c Regge models, JHEP **03** (2020) 101 [arXiv:1910.13432] [INSPIRE].
- [56] T. Blum, N. Christ, M. Hayakawa, T. Izubuchi, L. Jin, C. Jung et al., Hadronic Light-by-Light Scattering Contribution to the Muon Anomalous Magnetic Moment from Lattice QCD, Phys. Rev. Lett. 124 (2020) 132002 [arXiv:1911.08123] [INSPIRE].
- [57] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, *Remarks on higher-order hadronic corrections to the muon g-2*, *Phys. Lett. B* **735** (2014) 90 [arXiv:1403.7512] [INSPIRE].
- [58] T. Aoyama et al., The anomalous magnetic moment of the muon in the Standard Model, Phys. Rept. 887 (2020) 1 [arXiv:2006.04822] [INSPIRE].
- [59] J. P. Leveille, The Second Order Weak Correction to (G-2) of the Muon in Arbitrary Gauge Models, Nucl. Phys. B 137 (1978) 63 [INSPIRE].
- [60] F. S. Queiroz and W. Shepherd, New Physics Contributions to the Muon Anomalous Magnetic Moment: A Numerical Code, Phys. Rev. D 89 (2014) 095024
 [arXiv:1403.2309] [INSPIRE].
- [61] T. Aoyama, T. Kinoshita and M. Nio, Revised and Improved Value of the QED Tenth-Order Electron Anomalous Magnetic Moment, Phys. Rev. D 97 (2018) 036001 [arXiv:1712.06060] [INSPIRE].
- [62] D. Hanneke, S. Fogwell and G. Gabrielse, New Measurement of the Electron Magnetic Moment and the Fine Structure Constant, Phys. Rev. Lett. 100 (2008) 120801
 [arXiv:0801.1134] [INSPIRE].
- [63] D. Hanneke, S. F. Hoogerheide and G. Gabrielse, Cavity Control of a Single-Electron Quantum Cyclotron: Measuring the Electron Magnetic Moment, Phys. Rev. A 83 (2011) 052122 [arXiv:1009.4831] [INSPIRE].
- [64] BABAR collaboration, Search for a muonic dark force at BABAR, Phys. Rev. D 94 (2016) 011102 [arXiv:1606.03501] [INSPIRE].
- [65] BABAR collaboration, Search for a Dark Leptophilic Scalar in e⁺e⁻ Collisions, Phys. Rev. Lett. 125 (2020) 181801 [arXiv:2005.01885] [INSPIRE].
- [66] BELLE-II collaboration, Search for an Invisibly Decaying Z' Boson at Belle II in e⁺e⁻ → μ⁺μ⁻(e[±]μ[∓]) Plus Missing Energy Final States, Phys. Rev. Lett. **124** (2020) 141801 [arXiv:1912.11276] [INSPIRE].

- [67] PLANCK collaboration, Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641 (2020) A6 [arXiv:1807.06209] [INSPIRE].
- [68] D. Hooper, Particle Dark Matter, in Theoretical Advanced Study Institute in Elementary Particle Physics: The Dawn of the LHC Era, pp. 709-764, 2010, arXiv:0901.4090
 [INSPIRE].
- [69] D. Barducci, G. Belanger, J. Bernon, F. Boudjema, J. Da Silva, S. Kraml et al., Collider limits on new physics within micrOMEGAs_4.3, Comput. Phys. Commun. 222 (2018) 327 [arXiv:1606.03834] [INSPIRE].
- [70] T. R. Slatyer, Indirect dark matter signatures in the cosmic dark ages. I. Generalizing the bound on s-wave dark matter annihilation from Planck results, Phys. Rev. D 93 (2016) 023527 [arXiv:1506.03811] [INSPIRE].
- [71] L. Bergstrom, T. Bringmann, I. Cholis, D. Hooper and C. Weniger, New Limits on Dark Matter Annihilation from AMS Cosmic Ray Positron Data, Phys. Rev. Lett. 111 (2013) 171101 [arXiv:1306.3983] [INSPIRE].
- [72] J. Liu, N. Weiner and W. Xue, Signals of a Light Dark Force in the Galactic Center, JHEP 08 (2015) 050 [arXiv:1412.1485] [INSPIRE].
- [73] R. Essig, J. Mardon and T. Volansky, Direct Detection of Sub-GeV Dark Matter, Phys. Rev. D 85 (2012) 076007 [arXiv:1108.5383] [INSPIRE].
- [74] M. Pospelov and J. Pradler, Big Bang Nucleosynthesis as a Probe of New Physics, Ann. Rev. Nucl. Part. Sci. 60 (2010) 539 [arXiv:1011.1054] [INSPIRE].
- [75] B. Henning and H. Murayama, Constraints on Light Dark Matter from Big Bang Nucleosynthesis, arXiv:1205.6479 [INSPIRE].
- [76] FERMI-LAT collaboration, Cosmic-ray electron-positron spectrum from 7 GeV to 2 TeV with the Fermi Large Area Telescope, Phys. Rev. D 95 (2017) 082007 [arXiv:1704.07195]
 [INSPIRE].
- [77] PAMELA collaboration, An anomalous positron abundance in cosmic rays with energies 1.5-100 GeV, Nature 458 (2009) 607 [arXiv:0810.4995] [INSPIRE].
- [78] CALET collaboration, Energy Spectrum of Cosmic-Ray Electron and Positron from 10 GeV to 3 TeV Observed with the Calorimetric Electron Telescope on the International Space Station, Phys. Rev. Lett. 119 (2017) 181101 [arXiv:1712.01711] [INSPIRE].
- [79] DAMPE collaboration, The DArk Matter Particle Explorer mission, Astropart. Phys. 95 (2017) 6 [arXiv:1706.08453] [INSPIRE].
- [80] DAMPE collaboration, Direct detection of a break in the teraelectronvolt cosmic-ray spectrum of electrons and positrons, Nature 552 (2017) 63 [arXiv:1711.10981] [INSPIRE].