

 Open access • Journal Article • DOI:10.1051/JP4:1995443

## Thermochemistry and Reaction Mechanisms of Nitromethane Ignition

— [Source link](#) 

C.F. Melius

**Institutions:** Sandia National Laboratories

**Published on:** 01 May 1995 - Journal De Physique Iv (EDP Sciences)

**Topics:** Ignition system, Elementary reaction, Nitromethane and Thermochemistry

Related papers:

- [Global kinetics for n-heptane ignition at high pressures](#)
- [Reduced kinetic mechanism for high-temperature propane ignition](#)
- [Numerical study of the effects of non-equilibrium plasma on the ignition delay of a methane–air mixture using detailed ion chemical kinetics](#)
- [Experimental and kinetic modeling study on 2,4,4-trimethyl-1-pentene ignition behind reflected shock waves](#)
- [Reduced Chemical Reaction Mechanism of Shock-Initiated Ignition of Methane and Ethane Mixtures with Oxygen](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/thermochemistry-and-reaction-mechanisms-of-nitromethane-2zx5s4h60a>



**HAL**  
open science

# Thermochemistry and Reaction Mechanisms of Nitromethane Ignition

C. Melius

► **To cite this version:**

C. Melius. Thermochemistry and Reaction Mechanisms of Nitromethane Ignition. Journal de Physique IV Proceedings, EDP Sciences, 1995, 05 (C4), pp.C4-535-C4-548. 10.1051/jp4:1995443 . jpa-00253749

**HAL Id: jpa-00253749**

**<https://hal.archives-ouvertes.fr/jpa-00253749>**

Submitted on 1 Jan 1995

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Thermochemistry and Reaction Mechanisms of Nitromethane Ignition

C.F. Melius

*Combustion Research Facility, Sandia National Laboratories, Livermore, CA 94551-0969, U.S.A.*

**Abstract:** The thermochemistry and reaction mechanisms of nitromethane initiation are modeled using detailed chemical kinetics. Initial conditions correspond to gaseous nitromethane at atmospheric and liquid-like densities and initial temperatures between 1100 and 2000 K. Global reactions as well as elementary reactions are identified for each of the two stages of ignition. The chemical steps to convert the nitro group to N<sub>2</sub> involve a complex set of elementary reactions. The time-dependence of the ignition steps (ignition delay times) as a function of temperature and pressure is used to determine effective activation energies and pressure dependencies to ignition. The ignition delay times range from several nanoseconds to tens of microseconds. At atmospheric conditions, the delay times for both ignition stages are in excellent agreement with observed experimental data. At the high densities, the ignition times at these elevated temperatures appear to be dominated by the same reaction mechanism that occurs for atmospheric gaseous nitromethane initiation. This is to be contrasted with lower temperature, condensed-phase ignition studies where it appears that solvent-assisted reactions dominate.

### 1. INTRODUCTION

The ability to identify the chemical reactions involved in the initiation of energetic materials is challenging due to the extremely short time scales involved as well as the high temperatures and pressures that are generated. Furthermore, the chemical intermediates of ignition are highly reactive. The chemistry of nitro compounds, such as nitramines and nitroaromatics, is further complicated by the fact that they may undergo two-stage ignition. We therefore have investigated the initiation of nitromethane as a prototype of an energetic material containing the nitro moiety. The ignition of nitromethane has been studied both experimentally [1-9] and computationally [1,10-15]. In particular, the ignition delay time of gas-phase ignition of nitromethane has been studied by Guirguis *et al.* [1] in which they were able to observe two stages of ignition. Several groups have modeled the ignition process using detailed chemical kinetics [1,9], including previous work by us [13,14]. The modeling was able to treat the first-stage ignition reasonably well, but the second-stage ignition delay times were off by several orders of magnitude. Recent work by Lin and coworkers [16-18] have improved our understanding of the elementary reactions involving the NO, HNO, NO<sub>2</sub>, and HCN species. This has enabled us to generate an improved detailed reaction mechanism for nitromethane. We use this improved mechanism to study the initiation of gaseous nitromethane at atmospheric conditions as well as at densities corresponding to that of

---

\*Work supported by the U. S. Department of Energy.

liquid nitromethane. We present the thermodynamics and reaction pathways of the ignition steps. We then use the ignition delay times to determine effective activation energies and pressure dependencies.

## 2. METHOD

The study of ignition in a uniformly heated system, such as in a shock tube, allows us to study the combustion chemistry of energetic materials in a environment that can provide the initial decomposition and initiation reaction mechanisms. The energetic material can be treated as a homogeneous gas mixture in a closed system, in which we follow the time-dependent evolution of the chemical species. For instance, we can treat an adiabatic system with constant pressure or volume, or we can follow the chemistry under shock detonation conditions. We use the SENKIN program [19], which includes sensitivity analysis, for the integration of the energy and mass conservation equations.

In this section, we study the chemistry of gaseous nitromethane initiation. The ignition of nitromethane is treated as being under constant volume. The governing equations are [14,19]

$$\frac{dY_k}{dt} = v \dot{\omega}_k W_k \quad (k = 1, \dots, K). \quad (1)$$

and

$$c_v \frac{dT}{dt} = -v \sum_{k=1}^K e_k \dot{\omega}_k W_k, \quad (2)$$

In the above equations,  $T$  is the temperature,  $Y_k$  the mass fraction of the  $k^{\text{th}}$  species,  $W_k$  is the molecular weight of the  $k^{\text{th}}$  species,  $t$  is time,  $v = V/m$  is the specific volume,  $e_k$  is the internal energy of the  $k^{\text{th}}$  species,  $c_v$  is the constant volume heat capacity of the mixture, and  $\dot{\omega}_k$  is the molar rate of production of the  $k^{\text{th}}$  species. The chemical reaction mechanism included 48 species and 312 elementary reactions. The thermochemical dynamic properties and chemical production rates are evaluated using the Chemkin package [19]. The 48 species and their heats of formation are presented in Table I.

**Table I.** The chemical species used in the detailed chemical kinetic modeling of nitromethane. Also given are the heats of formation at 298 K ( $\Delta H_{f298}^0$ ), where energies are given in kcal-mol<sup>-1</sup>.

1	CH <sub>3</sub> NO <sub>2</sub>	-16.84	2	CH <sub>3</sub> ONO	-15.25
3	NO	21.58	4	NO <sub>2</sub>	7.91
5	HNO	23.80	6	HONO	-18.34
7	N <sub>2</sub> O	19.61	8	H	52.10
9	H <sub>2</sub>	0.0	10	O	59.56
11	O <sub>2</sub>	0.0	12	OH	9.32
13	H <sub>2</sub> O	-57.8	14	HO <sub>2</sub>	2.50
15	N	112.96	16	N <sub>2</sub>	0.0
16	NH	85.2	18	NH <sub>2</sub>	45.50
19	NH <sub>3</sub>	-11.97	20	NNH	58.57
21	HNNO	55.20	22	HNOH	21.06
23	H <sub>2</sub> NO	15.82	24	N <sub>2</sub> H <sub>2</sub>	50.90
25	CO	-26.42	26	CO <sub>2</sub>	-94.06
27	HCO	10.40	28	CH <sub>2</sub> O	-27.70
29	CH <sub>3</sub> O	3.90	30	CH <sub>2</sub> OH	-4.10

**Table I.** (Cont.)

31	CH <sub>3</sub> OH	-48.06	32	CH <sub>3</sub>	34.82
33	CH <sub>4</sub>	-17.90	34	CN	104.01
35	HCN	31.89	36	H <sub>2</sub> CN	59.11
37	CH <sub>3</sub> NH	44.94	38	HCNO	38.43
39	HOCN	-3.53	40	HNCO	-28.22
41	NCO	31.51	42	NCN	107.60
43	CH <sub>3</sub> NO	18.95	44	H <sub>2</sub> CNO <sub>2</sub>	36.47
45	H <sub>2</sub> CNO	41.43	46	CH <sub>3</sub> NHO	13.32
47	H <sub>2</sub> CNOH	5.15	48	CH <sub>2</sub> NO <sub>2</sub> H	-2.86

### 3. RESULTS

In Figs. 1 and 2, we present the temperature and species profiles as a function of time for the initiation of 100 per cent gaseous CH<sub>3</sub>NO<sub>2</sub> at 7.25 atmospheres and 1131 K. These conditions are representative of the experimental conditions of Guirguis *et al.* [1]. The initiation undergoes two ignition stages, at 13.3 μsec and 25.8 μsec, respectively. The final temperature is 3598 K.

The chemistry can be subdivided by time into several stages: (I) a preliminary stage of slow decomposition followed by (II) a first-stage ignition that undergoes a rapid temperature rise followed by (III) an intermediate stage in which the temperature remains on a plateau followed by (IV) a second-stage ignition where the temperature again rises rapidly to form the final products. The reaction mechanism flow diagrams for the first-stage ignition (II), the intermediate stage (III), and the second-stage ignition (IV) are shown in Figs. 3-5. These reaction pathways show the dominating elementary reactions occurring in each of these stages, with thick arrows being the major pathways while thin arrows are the minor pathways. We have separately shown the reaction chemistry for the carbon atom and the nitrogen atom of nitromethane. The carbon atom must be oxidized, to form CO and, to a lesser extent, CO<sub>2</sub> (note that there is not enough oxygen in nitromethane to oxidize the carbon to CO<sub>2</sub> and the hydrogen to H<sub>2</sub>O). Meanwhile, the nitrogen, which is in an oxidized state as part of the nitro group moiety, must be reduced to N<sub>2</sub>. The reaction flow diagram for the preliminary decomposition stage (I) is similar to that of the first-stage ignition (II) and thus is not shown. The major difference is that the preliminary stage (I) occurs at the low end of the temperature range, giving rise to slower rates and resulting in a greater variety of minor intermediate species being formed.

In Figs. 6-8, we show the enthalpies and free energies of possible species corresponding to the composition C<sub>1</sub>H<sub>3</sub>N<sub>1</sub>O<sub>2</sub>. This represents the composition of nitromethane and includes tautomers of nitromethane as well as various product species. Besides the ΔH at 300K (see Table I for individual species heats of formation), we present the ΔG's at 300K and 2000K.

Fig. 1 showed the time profile for nitromethane initiation at 7.25 atmospheres and 1131 K, indicating two rapid temperature rises at 13.6 μsec and 26.1 μsec, designated τ<sub>1</sub> and τ<sub>2</sub> respectively. In Table II, we present these ignition delay times, τ<sub>1</sub> and τ<sub>2</sub>, for various initial temperatures and pressures. These results are used to determine activation energies of ignition, defined by the relation  $\ln(\tau_a/\tau_b) \propto (\Delta E/R)(1/T_a - 1/T_b)$ . The resulting ΔE's for the first-stage and second-stage ignitions are presented in Table III. In a similar manner, we can define pressure dependencies to the ignition delay time by the relation  $\ln(\tau_a/\tau_b) \propto n \ln(P_a/P_b)$ , which corresponds to τ being proportional to P<sup>n</sup>. The resulting pressure exponents are given in Table IV.




Fig. 1. Temperature vs. time profile for ignition of nitromethane. Initial conditions are 1131 K and 7.25 atm.





Fig. 2. Species vs. time profile for ignition of nitromethane. Initial conditions are 1131 K and 7.25 atm.

## Nitromethane First Stage Ignition Chemistry

### Carbon Mechanism



### Nitrogen Mechanism





Fig. 3. Reaction mechanism flow diagrams for the first stage-ignition of nitromethane. Thick arrows indicate major pathways. Dominant collision partners for bimolecular reactions are indicated next to arrows. Unimolecular decomposition reactions are indicated by the third-body notation +M. The chemistry of carbon- and nitrogen-containing species are presented separately.

## Nitromethane Intermediate Stage Chemistry

### Carbon Mechanism



### Nitrogen Mechanism




Fig. 4. Reaction mechanism flow diagrams for the intermediate stage (between first- and second-stage) ignition of nitromethane. See Fig. 3 for further caption descriptions.








Fig. 7. Free energies,  $\Delta G$ 's, at 300 K for nitromethane and its tautomers and other stoichiometric products. Free energies are in kcal-mol<sup>-1</sup>.




Fig. 8. Free energies,  $\Delta G$ 's, at 2000 K for nitromethane and its tautomers and other stoichiometric products. Free energies are in kcal-mol<sup>-1</sup>.

**Table II.** Ignition delay times for gaseous nitromethane for different initial temperatures and pressures.

T (K)	P (Atm)	$\tau_1$	$\tau_2$	$\tau_2 - \tau_1$
1131	5.25	14.9 $\mu$ s	35.5 $\mu$ s	20.6 $\mu$ s
1131	7.25	13.3 $\mu$ s	25.8 $\mu$ s	12.5 $\mu$ s
1331	5.25	1.6 $\mu$ s	25.3 $\mu$ s	23.7 $\mu$ s
1331	7.25	1.2 $\mu$ s	15.5 $\mu$ s	14.3 $\mu$ s
1331	8.53	1.2 $\mu$ s	12.3 $\mu$ s	11.1 $\mu$ s
1131	1574.4	4.352 $\mu$ s	4.403 $\mu$ s	.051 $\mu$ s
1131	1852.8	4.288 $\mu$ s	4.298 $\mu$ s	.010 $\mu$ s
1331	1574.4	0.208 $\mu$ s	0.219 $\mu$ s	.010 $\mu$ s
1331	1852.8	0.202 $\mu$ s	0.208 $\mu$ s	.006 $\mu$ s
1900	2644.8	2.52 ns	4.76 ns	2.24 ns
2000	2784.	1.59 ns	3.59 ns	2.00 ns

**Table III.** Activation energies of ignition (defined by  $\ln(\tau_a / \tau_b) \propto \Delta E / R (1/T_a - 1/T_b)$ ). Ignition delay times are taken from Table II. Energies in kcal-mol<sup>-1</sup>.

Constant Pressure	T(K)	P (Atm)	$\Delta E_1$	$\Delta E_2$
	1230	5.25	33.2	7.6
	1230	7.25	35.6	5.1
	1230	1574.4	45.4	44.9
	1230	1852.8	45.7	45.3
Constant Volume	T(K)	P (Atm)	$\Delta E_1$	$\Delta E_2$
	1230	5.25	35.7	11.0
	1230	1574.4	45.9	45.6
	1950	2644.8	34.8	21.3

**Table IV.** Pressure dependence of ignition delay times for gaseous nitromethane. Pressure dependence is defined by  $\tau \propto P^n$  or  $\ln(\tau_a / \tau_b) \propto n \ln(P_a / P_b)$ . Ignition delay times are taken from Table II.

Low Pressure	T (K)	$n_1$	$n_2$
	1131	-0.36	-0.99
	1331	-0.85	-1.52
Low to High	T (K)	$n_1$	$n_2$
	1131	-0.22	-0.37
	1331	-0.35	-0.82
High Pressure	T (K)	$n_1$	$n_2$
	1131	-0.09	-0.15
	1331	-0.19	-0.32

## 4. DISCUSSION

### 4.1 Chemical Reactions

Several points regarding the nitromethane thermochemistry can be observed from Figs. 6-8. First, from the  $\Delta H$ 's presented in Fig. 6, it is seen that indeed nitromethane is an unstable tautomer of  $C_1H_3N_1O_2$ , lying  $\sim 90$  kcal-mol<sup>-1</sup> above the most stable form, C(O)OHNH<sub>2</sub>. In fact, C(O)OHNH<sub>2</sub> is more stable than the major products of detonation, CO, H<sub>2</sub>O, N<sub>2</sub>, and H<sub>2</sub> (see Fig. 2).

Furthermore, as can be seen in Fig. 6, the lowest bond-breaking moiety for CH<sub>3</sub>NO<sub>2</sub> is the breaking of the C-N bond to form CH<sub>3</sub> + NO<sub>2</sub>, which is endothermic by nearly 59 kcal-mol<sup>-1</sup>. Thus, from an enthalpy view point, CH<sub>3</sub>NO<sub>2</sub> would appear to be a reasonable energetic material.

However, while one tends to think of the thermochemistry of the reactions in terms of enthalpies (e.g., heats of reaction), it is the free energy of the system that drives the reaction. Dividing the C<sub>1</sub>H<sub>3</sub>N<sub>1</sub>O<sub>2</sub> molecular species into two or more species lowers the free energy, particularly at higher temperatures. As one can see from Fig. 8, at 2000K, the  $\Delta G$  of CH<sub>3</sub> + NO<sub>2</sub> is more stable than that of CH<sub>3</sub>NO<sub>2</sub> itself. And the breakup of CH<sub>3</sub>NO<sub>2</sub> into its small product molecules, CO, H<sub>2</sub>O, N<sub>2</sub>, and H<sub>2</sub>, forming three molecules from one reactant molecule is extremely exothermic. Thus, it is necessary to have more than the thermostability to bond breaking to make an insensitive energetic material. As we shall see, it is necessary to have a reaction pathway which prevents direct conversion of the energetic material to products.

The overall reaction chemistry converting nitromethane to products is




with minor contributions of the following reactions




Contributions of R2-R4 relative to reaction R1 are given in parentheses. The heats of reaction  $\Delta H_{R_{xn}}$  are given in kcal-mol<sup>-1</sup>.


As stated above, the chemistry can be subdivided by time into several stages: (I) a preliminary stage of slow decomposition followed by (II) a first-stage ignition that undergoes a rapid temperature rise followed by (III) an intermediate stage in which the temperature remains on a plateau followed by (IV) a second-stage ignition where the temperature again rises rapidly forming the final products. This temporal division of chemical reactions, as indicated in Fig. 2, is consistent with the temporal division of the temperature into a preliminary, first-stage ignition, intermediate region, and second-stage ignition, as indicated in Fig. 1. The reaction chemistry during the preliminary stage (I) is




the reaction during the first-stage ignition (II) is



the reaction during the intermediate-stage ignition (III) between the first- and second-stage ignitions is




while the reaction during the second-stage ignition (IV) is



In order to relate the heats of reaction,  $\Delta H_{R_{xn}}$ , of the individual reactions to each other, we have normalized the  $\Delta H_{R_{xn}}$  for each of the reactions to the  $\text{CH}_3\text{NO}_2$  molecule. Thus, the  $\Delta H_{R_{xn}}$ 's are normalized per  $\text{CH}_3\text{NO}_2$  (R5 and R6) or NO (R7 and R8) molecule.

The global reactions for each stage in the nitromethane initiation can be further subdivided into subglobal reactions which begin to indicate the reaction mechanism of each stage. While this reaction subdivision is not unique, we have attempted to define those subglobal reactions which represent simple combinations of elementary reactions, as indicated in Figs. 3-5. In addition to the subglobal reaction, we also indicate the relative contribution of each reaction to the net reactions presented above. We also give the absolute heat of reaction,  $\Delta H_{R_{xn}}$ , for each of the subglobal reactions.

The preliminary stage (I) global reaction R5 is composed of the subglobal reactions



The first-stage global reaction R6 is composed of the subglobal reactions





The intermediate-stage global reaction R7 is composed of the subglobal reactions



The second-stage global reaction R8 is composed of the subglobal reactions



The net result of the preliminary (I) and first-stage ignition (II) reactions steps is the conversion of the  $\text{NO}_2$  moiety to  $\text{NO}$ . During the preliminary stage (I), the decomposition of  $\text{CH}_3\text{NO}_2$  is only slightly exothermic. In fact, the dominant decomposition reaction R5a is endothermic. Subsequent reactions involving the  $\text{HNO}$  species provide the exothermic heat release. The first-stage ignition (II) chemistry is slightly more exothermic. However, the major exothermic heat release of initiation arises during the second-stage ignition chemistry.

The net result of the intermediate (III) and second-stage ignition (IV) reactions is the conversion of  $\text{NO}$  to  $\text{N}_2$ . The  $\text{NO}$  to  $\text{N}_2$  conversion does not occur readily and requires the presence of hydrogen and  $\text{HCN}$ . The formation of  $\text{HCN}$  occurs primarily during the intermediate stage (III). This stage (III), which removes the  $\text{CH}_4$ , is also exothermic but slow. The rapid reaction rates of the second-stage ignition are a result of the high temperatures.

While the net reaction is conversion of  $\text{CH}_3\text{NO}_2$  to  $\text{CO}$ ,  $\text{H}_2\text{O}$ ,  $\text{N}_2$ , and  $\text{H}_2$ , the overall chemical process of ignition is complicated. Many steps are required to form the final exothermic products. The chemistry can be divided into two major efforts, (1) conversion of the  $\text{NO}_2$  moiety to  $\text{NO}$  and (2) conversion of the  $\text{NO}$  moiety to  $\text{N}_2$ . This process is generic to all nitro-containing energetic materials, including nitrate esters and nitramines [14]. The result is two separated ignition stages, denoted the first or primary stage and the secondary stage.

The primary-stage ignition is conversion of the  $\text{NO}_2$  to  $\text{NO}$ . This can be written stoichiometrically as  $\text{CH}_3\text{NO}_2 \rightarrow \text{CO} + \text{NO} + \frac{3}{2} \text{H}_2$ . Note that this reaction is itself endothermic (see reaction R6a). On the other hand, three and a half molecules are formed for each nitromethane molecule, so the entropy of this reaction is favorable, leading to a favorable free energy of reaction as the temperature increases. In order to gain heat release, some of the  $\text{NO}$  must be reduced. In the preliminary stage (I),  $\text{N}_2\text{O}$  is formed via the  $\text{HNO}$  intermediate. During the first-stage ignition (II), the temperature rise is sufficient to allow thermal decomposition of the  $\text{N}_2\text{O}$  to form  $\text{N}_2$ . The relative importance of  $\text{CH}_4$  and  $\text{CH}_3\text{OH}$  has changed from our previous investigation [14], with  $\text{CH}_4$  now playing the more significant role. The relative importance of  $\text{CH}_4$  to  $\text{CH}_3\text{OH}$  is extremely sensitive to the pressure dependence used for various elementary rate constants. In both cases, these species serve to create a temporary heat source (see reaction R6b). The formation of  $\text{CH}_4$  occurs earlier in the reaction mechanism (arising from  $\text{CH}_3$ ) than does formation of  $\text{CH}_3\text{OH}$  (arising from  $\text{CH}_3\text{O}$  as indicated in ref. 14).

One can see from Fig. 3 that even the oxidation of the carbon of nitromethane to  $\text{CO}$  is a multistep process. The major pathway has the  $\text{CH}_3\text{NO}_2$  breaking the C-N bond to form  $\text{CH}_3 + \text{NO}_2$  which recombines to form  $\text{CH}_3\text{ONO}$  which breaks the O-N bond to form  $\text{CH}_3\text{O} + \text{NO}$ . The  $\text{NO}$  then abstracts the H from  $\text{CH}_3\text{O}$  to form  $\text{CH}_2\text{O} + \text{HNO}$ . Subsequent abstraction of hydrogen atoms

from  $\text{CH}_2\text{O}$  lead to  $\text{CO}$ . As one can see from Figs. 6-8, this series of reactions is itself endothermic. Thus, the initial set of reactions do not produce significant heat release. This complex set of reactions plays an important role in keeping the nitro compounds from spontaneously decomposing. The heat release from any exothermic reactions is converted into net endothermic reactions. These series of elementary reactions are necessary to move the oxygen atom from the nitrogen atom to the carbon atom, which cannot occur until the C-N bond is first broken.

The secondary stage ignition is the reduction of  $\text{NO}$  to  $\text{N}_2$ . This occurs by first reducing one of the nitrogen atoms to a reduced state in  $\text{HCN}$ . This reduced nitrogen is then oxidized with another oxidized nitrogen of  $\text{NO}$  to form the formally neutral  $\text{N}_2$ . The formation of the reduced state of nitrogen,  $\text{HCN}$ , occurs during the intermediate stage (III). This step is necessary because the  $\text{NO}$  molecule is very stable, even at the intermediate temperatures.

As noted above, the large free energy release of the reaction is enhanced by the net increase in the entropy of the reaction. The majority of this entropy increase occurs during the preliminary and first-stage ignition, where the decomposition of each nitromethane molecule leads to between two and three new molecules. Thus, while the exothermicity of the preliminary and first stages of the reaction are small compared to that of the intermediate and second stages, the increase in entropy plays an important role in driving the initiation process, allowing the initial reactions to proceed.

## 4.2 Ignition Delay Times

As discussed in the above section on the reaction mechanism, the nitromethane molecule cannot "spontaneously" decompose to form exothermic products but rather must undergo a series of intermediate steps, many of which are endothermic. These reactions lead to finite ignition delay times, both for the primary and secondary stages. We can analyze these ignition delay times,  $\tau_1$  and  $\tau_2$ , in terms of energies of activation,  $\Delta E_1$  and  $\Delta E_2$ , and pressure dependencies  $n_1$  and  $n_2$ , as defined above and in Tables II-IV. The  $\Delta E$ 's and  $n_i$ 's are defined as derivatives of the ignition delay times  $\tau_i$ 's with respect to temperature and pressure, respectively, and thus, have meaning only in the near region of the temperature and pressure specified. We have calculated the ignition delay times for atmospheric pressures as well as a high pressure range corresponding to a nitromethane density of that of liquid nitromethane. The pressure values presented in Tables II-IV are an idealized gas effective pressure with the real gas pressure being much higher. For the constant volume calculations, it is better to view the calculations as constant density calculations.

The atmospheric conditions match the experimental data of Guirguis *et al.* [1]. The ignition delay times are in quite good agreement with the experimental data. The 7.25 atm, 1131 K conditions, for which the experimental was able to separate the two ignition stages, yielded a  $\tau_1$  and a  $\tau_2$  of 13.3 and 25.8  $\mu\text{sec}$ , which are in remarkable agreement with the experimental data of  $\sim 12$  and  $\sim 25$   $\mu\text{sec}$  respectively. These results give us confidence that the elementary reactions for modeling nitromethane are accurate. This represents an improvement over the reaction mechanism used previously [14], which had a reasonable first-stage ignition but a much delayed second stage ignition. The ignition delay times range from microseconds at the low pressure to hundreds of nanoseconds in the high density regime. At the high temperatures and densities, the  $\tau_1$ 's approach the nanosecond time scale.

The effective activation energies of ignition are presented in Table III. The resulting  $\Delta E$ 's indicate that the first-stage ignition has an activation energy  $\Delta E_1$  of  $\sim 33$ - $36$   $\text{kcal}\cdot\text{mol}^{-1}$  at atmospheric pressures, consistent with the experimental data [1]. At high densities, the  $\Delta E_1$  is larger,  $\sim 46$   $\text{kcal}\cdot\text{mol}^{-1}$  though at elevated temperatures (1950 K), the  $\Delta E_1$  is  $\sim 35$   $\text{kcal}\cdot\text{mol}^{-1}$ . These large activation energies are dominated by the initial C-N bond breaking process, which has a bond dissociation energy of  $59$   $\text{kcal}\cdot\text{mol}^{-1}$ . The second-stage ignition delay times,  $\tau_2$ 's, show a wide variation in activation energies. This occurs because the  $\tau_2$ 's decrease significantly faster than the  $\tau_1$ 's with increasing temperature and pressure. Ultimately, the  $\tau_2$ 's approach the  $\tau_1$ 's and are

dominated by the  $\tau_1$  times at high densities. However, the difference between the two stages,  $\tau_2 - \tau_1$ , is still several nanoseconds (see Table II).

The effective pressure dependencies of ignition (at constant temperature) are presented in Table IV. The resulting  $n_i$ 's exhibit varied behavior. At low temperature and pressure, the first-stage  $n_1$  pressure dependence is between zeroth order ( $n_1 = 0$ ) and first order ( $n_2 = 1$ ), corresponding to effective unimolecular and bimolecular behavior. The secondary stage  $n_2$  pressure dependence at the low temperature and pressure is first order and greater. At high pressures, both the  $n_1$  and  $n_2$  pressure dependencies approach zeroth order, indicating the dominance of the first-stage ignition delay time and the high effective activation energy representative of C-N bond breaking. It is difficult to relate these pressure dependencies to shock induced experiments where temperature changes occur simultaneously with pressure changes.

The comparison of our high density results with the experimental ignition delay times at kilobar pressures and low temperatures indicate some significant differences. As an example, we use the experimental ignition delay times measured by Lee *et al.* [3] for 10 and 50 kbar and 400-600 K. Their  $\tau$ 's range from 10 - 100 seconds and have an effective activation energy of  $\sim 16$  kcal-mol.<sup>-1</sup> While this activation energy is significantly lower than what we calculate for the higher temperature (1100-1300K) conditions ( $\sim 45$  kcal-mol.<sup>-1</sup>), our pre-exponential rate constants are much larger. Defining the rate constant  $k$  as the inverse of the ignition delay time, the effective high temperature rate constant is  $k_g = 1.4 \cdot 10^{14} \exp(-45.4/RT)$  while the effective low temperature rate constant (at 10 kbar) is  $2.7 \cdot 10^4 \exp(-15.7/RT)$ . That is, our ignition delay times are between 200 nsec - 4  $\mu$ sec while their extrapolated  $\tau$ 's are still longer than 100  $\mu$ sec. On the other hand, our  $\tau$ 's in the 400-600K range would be longer than theirs. This suggests a switch in mechanism in which at the lower temperatures, the reaction mechanism is "condensed phase"-like, involving proton transfer and hydrolysis reactions, while at high temperatures above 1000 K, the reaction mechanism is "gas phase"-like, involving radical species. We have investigated the inclusion of bimolecular as well as solvent-assisted proton transfer and hydrolysis reactions in our mechanism to mimic the condensed phase reaction process. The results are encouraging. The small pre-exponential rate constants for such reactions (possessing severe entropy constraints) make such reactions important in the 400-600 K temperature range while they have essentially no effect in the 1100-1900 K range. Considerable work remains in order to elucidate the actual elementary reactions for the low temperature, condensed-phase regime.

## 5. CONCLUSIONS

We have calculated the initiation process for nitromethane using detailed chemical kinetics. The starting conditions were gaseous nitromethane at atmospheric pressures as well as elevated pressures corresponding to liquid density and temperature above 1000 K. The results indicate that the chemical reaction mechanism requires multiple steps, in which many of the initial steps are endothermic. This complex chemical mechanism prevents the energetic nitro-compound from releasing its stored chemical energy immediately as a shock wave passes by. Ignition delay times indicate relatively high activation energies and low pressure dependencies, indicative of initial bond-breaking of the C-N bond. The agreement between our ignition delay times and the experimental data of Guirguis *et al.* [1] is encouraging for the modeling of other nitro-containing energetic material. These systems will have similar NO, HNO, N<sub>2</sub>O, and HCN chemistry, particularly for the second-stage ignition. These results indicate that a different mechanism is occurring at elevated temperatures (> 1000 K) compared to lower temperatures.

## REFERENCES

- [1] R. Guirguis, D. Hsu, D. Bogan, and E. Oran, *Comb. and Flames* **61**, 51 (1985).
- [2] D. S. Y. Hsu and M. C. Lin, *J. Ener. Mat.*, **3**, 95 (1985).
- [3] E. L. Lee, R. H. Sanborn, and H. D. Stromberg, *Proc. 5<sup>th</sup> Symp. (Int) Detonation*, 331 (1970).
- [4] G. J. Piermarini, S. Block, and P. J. Miller, *J. Phys. Chem.* **93**, 457 (1989).

- [5] C. P. Constantinou, T. Mukundan, and M. M. Chaudhri, *Phil. Trans. R. Soc. Lond A* 339, 403 (1992).
- [6] J. Connor, Chemistry and Physics of the Molecular Processes in Energetic Materials (S. N. Bulusu, ed.), ASI 309, p 545 (1990).
- [7] J. C. Mialocq, *J. de Phys.* C4, 163 (1987).
- [8] S. Zeman, *Thermochemica Acta*, in press.
- [9] A. Perche, J. C. Tricot, and M. Lucquin, *J. Chem. Res.* S, 116 (1979), *J. Chem. Res.* M, 1555 (1979).
- [10] R. D. Bardo, 8<sup>th</sup> Symp. (Int) Detonation, 855 (1989).
- [11] M. D. Cook and P. J. Haskins, 12<sup>th</sup> Int Pyro Symp., 43 (1987).
- [12] S. Odier, *J. de Phys.* C4, 225 (1987).
- [13] C. F. Melius, *Phil. Trans. R. Soc. Lond A* 339, 377 (1992).
- [14] C. F. Melius, Chemistry and Physics of the Molecular Processes in Energetic Materials (S. N. Bulusu, ed.), ASI 309, p51 (1990).
- [15] C. F. Melius, Chemistry and Physics of the Molecular Processes in Energetic Materials (S. N. Bulusu, ed.), ASI 309, p21 (1990).
- [16] C.-Y. Lin, H.-T. Wang, M. C. Lin, and C. F. Melius, *Int'l. J. Chem. Kinetics* 22, 455 (1990).
- [17] M. C. Lin, Y. He, and C. F. Melius, *Int'l. J. Chem. Kinetics* 24, 489 (1992).
- [18] M. C. Lin, Y. He, and C. F. Melius, *Int'l. J. Chem. Kinetics* 24, 1103 (1992).
- [19] A. E. Lutz, R. J. Kee, and J. A. Miller, Sandia Report SAND87-8248 (1987).