
Thin layer chromatography-application in qualitative analysis on presence of coumarins and flavonoids in plant material

Elvira E. Kovač-Bešović, Kemal Durić

Department of Pharmacognosy, Faculty of Pharmacy

University of Sarajevo, Čekaluša 90, Sarajevo, Bosnia and Herzegovina

Abstract

Drugs, natural medicinal plant, animals and mineral materials, have a large and various application in official pharmacy and medicine. Carriers of multilateral pharmacological effects that those drugs shown, are chemically define as active components that are present in them. Methods of qualitative and quantitative analysis are used for the chemical investigation of components that drugs contain. Method of thin layer chromatography has been shown as very reliable.

According to the chemical investigation of single drugs, it is possible to define a group of compound or single compound comparing them with standards. Relating to the usage of method of thin layer chromatography, it has been carried out investigation on presence of coumarins and flavonoids in domestic plant material that have wide everyday usage. Coumarins and flavonoids from the point of view of chemical belonging are phenol derivatives with important pharmacological effects.

Applying method of thin layer chromatography, it is detected presence of coumarins and flavonoids substances in plant material that has been tested. *Anethi graveolens fructus et folium* (fruit and leaf of dill), *Anethum graveolens L.*, *Apiaceae*, *Avenae sativae fructus* (fruit of oats), *Avena sativa L.*, *Poaceae* and *Asperulae odoratae herba* (sweet woodruff), *Asperula odorata L.*, *Rubiaceae*. Chromatograms are developed in systems cyclohexane-ethylacetat (13:7) and toluene-ether (1:1) saturated with 10% acetic acid, and visualisation by observing on UV lamp (254 and 366 nm), spraying with reagents KOH (10% ethanol solution) and diphenylboryloxyethylamine (1% methanol solution).

Key words: TLC, coumarins, flavonoids, dill, oats, sweet woodruff

Introduction

Chromatography belongs to the group of analytical separation methods that are applied for identification, purification and determination of contents of substances or preparations. The separation of different components using chromatography is based on dynamic distribution

of dissolve substance between two fazes that are unmixed and one of them must be mobile in relation to one other. The immobile faze slow down movement of the melted components, enabling single substances to be separated under certain conditions. Using chromatographic methods makes possible to separate substances, which have small difference from the point of view of the chemical structure, because they have similar physical and chemical properties, that chromatography put in advantage respect to other methods (1, 2).

Analysis of natural raw material, drugs and isolated substances, according to principles of pharmacognosy is carried out mostly by using methods of thin layer chromatography and column chromatography (3, 4). They were applied for qualitative and quantitative analysis of drugs, natural products and preparation with natural components.

A series of advantages, respect to the other chromatographic methods, gives the priority in usage of thin layer chromatography in pharmacognostic and phytochemical analysis of drugs. It distinguished, in addition to relatively fast application, well separation that can be archived using small quantities of samples, and with thin layer of adsorbents, and the results obtained are reproducible under standardised conditions.

In that sense it was carry out analysis of chemical composition of plant material using method of thin layer chromatography and that was shown in this work.

It was carry out the analysis of following drugs *Anethi graveolens fructus et folium* (fruit and leaf of dill), *Anethum graveolens L.*, *Apiaceae*, *Avenae sativae fructus* (fruit of oats), *Avena sativa L.*, *Poaceae* and *Asperulae odoratae herba* (sweet woodruff), *Asperula odorata L.*, *Rubiaceae*.

Dill is herbaceous annual plant that was grown as spice plant in the garden, but also in plantation. It is of use its mature fruit, leaf, but also the upper overgrown part of the plant in bloom as herbs. The fruits contain ethereous oil with carvone, limonene, than fatty oil, coumarin derivatives, and proteins. Leaf or herbs contain essential oil, vitamin C, pro-vitamin A, flavonoids. It is in use as carminative, stomachic, diuretic, bland sedative, lactogogue. Use of fruits extract as antispasmodic in light

forms of chronic coronary insufficiency, preventively against asthma and contraction of abdominal organs, is very important. As specie is in a large use (5, 6).


Oath is cultivated cereal that grows everywhere, even in a poor field. It is in use-husked fruit obtained from cultivated plant. It contain silicon acid, flavonoids, avenacin, avenacoside, coumarins, vitamin C and B-complex, amide, proteins, lecithin, fatty oils, mineral material, especially iron, iodine, wax and sugars. It is in use as food, dietetic and medicine. It use as sedative and diuretic in homeopathic medicine. It is very important as stimulant of immunity (6, 7).

Sweet woodruff is herbaceous plant wide spread in forest area especially in beech forests. It is useful over ground part of the plant during flowering period. Drug contains coumarins, iridoid glycosides, tannins, bitter substances and essential oil. Preparations with sweet woodruff are used as anti-inflammatory, lymphokinetic, antispasmodic and as diuretic, bland sedative and aromatic. The drug is not explored and used sufficiently (8, 9).

Investigation of the above mentioned drugs were carried out with thin layer chromatography method according to the principles of this method. Adsorbent was silica gel 60 F254, on which were applied methanolic extracts of drugs and respectively coumarins and flavonoids standards. Chromatograms were developed in two systems that literatures cite for analysis of coumarins and flavonoids phenolic derivatives.


In this way we would like to ascertain possibly optimisation in separation of these components that limit use of drugs that contain them (10, 11). Separated coumarins and flavonoids spots show blue-white and yellow fluorescence under UV-lamp on 254 and 366 nm that became more intense after spraying with reagents ethanolic potassium hydroxide and diphenylboryoxyethylamin. This is very evident after spraying with second reagent under UV-366 nm. The results we obtained were shown on photos and tables that contain Rf-values of separated spots.

Standard substances fraxin, fraxetin, aesculin, aesculetin were supplemented with coumarin substances obtained synthetically.




Their aim of usage is comparison between coumarin nucleus behaviours, such is contained in natural material and which is result of metabolic process of natural living

and plant environment, and those, which are, produced synthetically (12).



S-Synthetic samples



Standards of flavonoids compounds, quercetin and rutin were applied as natural raw material.

Material and methods

In this work the analysis have been carried out on presence of coumarins and flavonoids in selected plant material by using method of thin layer chromatography. This method was used for qualitative chemical analysis. Plant species we utilized for analysis are: dill, *Anetum graveolens L.*, *Apiaceae*, oats, *Avena sativa L.*, *Poaceae* and sweet woodruff *Asperula odorata L.*, *Rubiaceae*. All of three plant species give drugs that are: dill, *Anethi graveolens fructus*, fruit (Figure 1), *Anethi graveolens folium*, leaf (Figure 2), oats, *Avenae sativae fructus*, husked fruits (Figure 3) and sweet woodruff, *Asperulae herba*, over ground part (Figure 4). Fruit and leaf of dill had been collected in surroundings of Sarajevo and dried as drug. Leaf was dried binding over ground parts, herb, keeping it on dray and ventilated place. The fruit was collected in mature state, and was dried in thin layer on drayed and ventilated place. Along with draying of leaf of dill, it is necessary to conserve its green colour, while with its fruit the contact with humidity should be avoided. Mature fruits of oats were collected on ownership in surrounding of Sarajevo. Sweet woodruff was acquired as already prepared drug.

Method of thin layer chromatography was carried out under following conditions.

Extracts of plant material, of investigated drugs, were prepared in the following way: Powdered drug (1g) is extracted with methanol (10ml) for 30 minutes under reflux on the water bath; filtrate is evaporated to about 1 ml, and 10 μ l is used for TLC investigation. In this way,

Figure 1. *Anethi graveolens fructus*, dill's fruits, *Anethum graveolens L.*, Apiaceae



Figure 2. *Anethi graveolens folium*, (herba), (dill), *Anethum graveolens L.*, Apiaceae



extracts are prepared and bring on chromatogram by following sequence:

1. *Anethi graveolens fructus*,
2. *Anethi graveolens folium*,
3. *Avenae sativae fructus*,
4. *Asperulae herba*.

Methanolic solution (0.1%) of coumarins and flavonoids substances is used for standard, more precisely coumarin derivatives fraxin, fraxetin, aesculin, aesculetin and three coumarins as synthetic substances. Fraxin and aesculin are hetero-side compounds, fraxetin and aesculetin are their corresponding aglicons, which are liberated by hydrolysis of hetero-sides, during which glycolic part is separated. Synthetic coumarins are utilized as standards because of their coumarin structure which can give information in correlation with natural coumarin substances from their analytics as well affects point of view.

Figure 3. *Avenae sativae fructus*, oats's fruits, *Avena sativa L.*, Poaceae



Figure 4. *Asperulae odoratae herba*, sweet woodruff, *Asperula odorata L.*, Rubiaceae



For standards substances of flavonoid structure are used methanolic solution (0.1%) of quercetin ("Kemika", Zagreb) and rutin ("Merck", Darmstadt).

Pre-coated TLC plates with adsorbent silica gel 60 F254, layer thickness 0.25, "Riedel-de-Haën" Seelze, Germany, dimension 20x20 cm are used.

Extracts were brought on thin layer with capillary, dotted with Nanomat IV "Camag", Muttensz, Switzerland.

Chromatograms are developed in systems of solvents, which are specific for coumarins and flavonoids substances, cites in literature like that one (3).

Presence of those derivatives is investigated by developing in systems

- a) Cyclohexane-ethylacetate (13:7); developing values d_2 is 14.5 cm (travelling of mobile phase was 60 min).

- b) Toluene-ether (1:1), saturated with 10% acetic acid; developing values d_2 is 14.5 cm (travelling of mobile phase was 55 min).

Detection of developed spots is carrying out according to the principles of visualisation of chromatograms. First, chromatograms are observed under UV-lamp on 2 standard wavelengths, 254 nm and 366 nm. Fluorescence of separated spots in comparing to fluorescence of standards could be considered reliable information in identification of analysed coumarin and flavonoid derivatives. This substance gives intensive fluorescent spots of blue-white and yellow colour as standards and those corresponding spots got from drug's extracts.

Chromatograms are sprayed with reagents for identification, attaining reinforcement of intensity of fluorescence, thanks to them. Reagents are following:

- KOH-ethanolic (10%), reagents for identification of presence of coumarins substances
- Diphenylboryloxyethylamin (Merck, Darmstadt)-methanolic (1%), reagents for identification of flavonoids substances.

The chromatograms we obtained are presented in original photos made after developing and observing under UV-lamp before and after spaying with abovementioned reagents (Figure 5, 5a, 5b, 6, 6a, 6b, 7, 7a, 7b, 8, 8a, 8b).

Results and discussion

In this work chromatographic analysis have been carried out on presence of coumarins and flavonoids substances in selected plant material. Method of thin layer chromatography, which is quoted in pharmacopoeia's regulations for analysis of drugs but also other substances, it is utilised for qualitative chemical analysis of drugs (13, 14). The plant material we analysed, leaf and fruit of dill, fruit of oats and sweet woodruff, are drugs in everyday usage and that is why chemical study represents an important information. Taking food into man's organism is not the only importance of this studies but also the possibility of preparation of different pharmaceutical medicaments intended to the specific pharmacologic indication. From viewpoint of pharmacological affect, coumarins and flavonoids represent very important groups of pharmacologically active chemical compounds present in plant material, but also the substances that are obtained semi-synthetically or synthetically trying to meliorate pharmacological efficiency.

By chromatographic analysis of drugs, *Anethi graveolens fructus*, *Anethi graveolens folium*, *Avenae sativae fructus* and *Asperulae herba*, with corresponding standards, the chromatograms we obtained gives next results:

Figure 5. System, cyclohexane: ethylacetat (13:7); Visualization UV-254 nm




Figure 5a. System, cyclohexane: ethylacetat (13:7); Visualization UV-366 nm





Figure 5b. System, cyclohexane: ethylacetat (13:7); Visualization UV-366 nm, sprayed with KOH/EtOH (10%)



The presence of coumarin substances in analysed plant material under determinate conditions, after visualisation with UV-lamp and spraying chromatogram with ethanolic solution of potassium. Standard of hetero-side fraxin and its aglicon fraxetin, then hetero-side aesculin and its aglicon aesculetin belongs to the group of coumarin sub

Figure 6. System, toluen:ether (1:1), saturated with acetic acid (10%); Visualization UV-254 nm




Figure 6a. System, toluen:ether (1:1), saturated with acetic acid (10%); Visualization UV-366 nm




Figure 6b. System, toluen:ether (1:1), saturated with acetic acid (10%); Visualization UV-366 nm, sprayed with KOH/EtOH (10%)




Figure 7. System, cyclohexane:ethylacetat (13:7); Visualization UV-254 nm




Figure 7a. System, cyclohexane:ethylacetat (13:7); Visualization UV-366 nm





Figure 7b. System, cyclohexane:ethylacetat (13:7); Visualization UV-366 nm, sprayed with diphenylboryoxyethylamin



stances with basic coumarin nucleus. The same structure had the synthetic coumarin substances, which we used in experiment.

All samples we examined, on start line show the spots which correspond to standards of fraxin (A) and aesculin

(C), blue fluorescence (UV-254 nm), intensively blue-white (UV-366 nm), that reinforce after chemical treatment with ethanolic KOH, with $R_f=0,01$, values which are in correlation with literature data about this coumarin aglicons (3). Standard of fraxetin (B) has $R_f=0,05$, and the spot is blue-brown (UV-254), more exactly blue-dark

(UV-366 nm). Standard of aesculetin (D) has Rf-0.08, in zone of fluorescence 254 nm show blue spots, or blue-white fluorescents on 366 nm. From synthetic coumarin compounds, S3 show blue-brown fluorescence (UV-254 nm), Rf-0.05, as aesculetin. In system toluene-ether, saturated with acetic acid, synthesised compound remain on start line together with standards of fraxin and aesculin. As we mentioned before all comparative coumarin standards has the basic coumarin structure. For the separation of coumarin substances, the system we applied on had shown to be good, apposite to system toluene-ether, saturated with acetic acid, a little bit better then system cyclohexane-ethylacetat (Figure 5, 5a, 5b, 6, 6a, 6b, Table 1).

Methanolic extracts *Anethi graveolens fructus*, and *Anethi graveolens folium* in system cyclohexane-ethylacetat gives the spots of brown fluorescence (UV-254 nm) on start line, as standard quercetin, Rf-0.01. Dark-brown zone of fluorescence on Rf-0.15 show extract of *Asperulae herba* together with rutin standard (UV-254 nm). From extract of *Avenae sativae fructus* remain on start line spot of light-brown fluorescence (UV-254 nm), Rf-0.09, which is identical with behaviour of rutin standard, Rf-0.11.

In zone UV-366 nm, fluorescence of spots is blue (Rf-0,01), and after spraying with reagents diphenylboryoxyethylamin, the spots show intensive yellow fluorescence (Rf-0.01), and Rf-0.34 that is given from methanolic extract of *Asperulae herba*.

In system toluene-ether, saturated with acetic acid, from extract of *Avenae sativae fructus* the separated component in zone UV-366 nm show yellow fluorescence (Rf-0.16) the same as rutin (Rf-0.15), with close Rf values (Figure7, 7a,7b, 8, 8a, 8b, Table 2).

The results we obtained from chromatographic analysis of examined drugs, show that they contain coumarin and flavonoid substances, as part of there own metabolism, which is in correspondence with literature quotation From the type of coumarins and flavonoids point of view, especially their percentage contents, investigations continue, by using other necessary methods of qualitative and especially quantitative analysis.

The results obtained, also, show the justify utilize of this drugs as medicament natural raw material, and especially present contribution on better and bigger employment of sweet woodruff in pharmacy and medicine.

Figure 8. System, toluen:ether (1:1), saturated with acetic acid (10%); Visualization UV-254 nm




Figure 8a. System, toluen:ether (1:1), saturated with acetic acid (10%); Visualization UV-366 nm




Figure 8b. System, toluen:ether (1:1), saturated with acetic acid (10%); Visualization UV-254 nm, sprayed with diphenylboryoxyethylamin




Table 1. Rf-values of separated components (analyses of coumarins)

	Sample	Rf-values	
		System I*	System II**
1.	<i>Anethi graveolens fructus</i>	0.01	0.01
		0.15	0.14
		0.50	0.45
		0.84	0.80
2.	<i>Anethi graveolens folium</i>	0.01	0.01
		0.13	0.13
		0.30	0.44
		0.49	
3.	<i>Aveanae sativae fructus</i>	0.01	0.01
4.	<i>Asperulae odoratae herba</i>	0.01	0.01
		0.16	0.19
		0.32	0.27
		0.98	0.96
A	Fraxin	0.01	0.01
B	Fraxetin	0.05	0.07
C	Esculin	0.01	0.01
D	Esculetin	0.08	0.08
S₁	Synthetic coumarin	0.01	0.01
S₂	Synthetic coumarin	0.01	0.01
S₃	Synthetic coumarin	0.05	0.05

Table 2. Rf-values of separated components (analyses of flavonoids)

	Sample	Rf values	
		System I*	System II**
1.	<i>Anethi graveolens fructus</i>	0.01	0.01
		0.19	0.2
		0.32	0.24
		0.37	0.38
2.	<i>Anethi graveolens folium</i>	0.01	0.01
3.	<i>Aveanae sativae fructus</i>	0.09	0.16
4.	<i>Asperulae odoratae herba</i>	0.34	0.01
			0.30
E	Quercetin	0.01	0.01
F	Rutin	0.11	0.15

* System I - cyclohexane: ethylacetat (13:7)

** System II - toluene: ether (1:1), saturated with 10% acetic acid

References

1. Nyiredy, Sz. Planar Chromatography, A Retrospective View for the Third Millenium, Springer Scientific Publisher, Budakalász, Hungary, 2001
2. Nikolin, B. Analitika lijekova, Sarajevo Publishing, Sarajevo, 1998
3. Wagner, H., Bladt, S. Plant Drug Analysis, Springer-Verlag Berlin-Hedelberg-New York, 1996
4. Kovač-Bešović, E. Metode u farmakognoziji, Sarajevo Publishing, Sarajevo, 2001
5. Hassanzadeh, M.K., Rahimizadeh, M. and Vaheddie, L., Constituents of the essential oils of aerial parts of dill (*Anethum graveolens* L.) herb., *Oriental. J. Chemistry*, 2000; 14:2, 217-220
6. Hagers Handbuch der Pharmazeutischen Praxis, Springer-Verlag Berlin-Heidelberg-New York, 1973
7. British Herbal Pharmacopoeia, British Herbal Medicine Association, England, 1995
8. Sticher, O. Iridoidglucoside aus *Asperula odorata* L., *Galeopsis tetrahit* L. und *Thevetia peruviana* (PERS.) K. SCHUM., Habilitationsschrift ETH Zürich, 1970
9. Sticher, O. Isolierung von Monotropein aus *Asperula odorata* L. (Rubiaceae), *Pharm. Acta Helv.* 1971; 46, 121
10. Galand, N., Pothier, J., Mason, V. and Viel, C. Separation of flavonoids, coumarins and antocyanins in plant extracts by Overpressure Layer Chromatography, *Pharmayie*, 1999; 54:6, 468-471
11. Budzianowski, J. Coumarins, caffeoilytartaric acids and their artifactual methyl esters from *Taraxacum officinale* leaves, *Planta medica* 1997; 63:3, 288.
12. Kovač-Bešović, E., Završnik, D. And Durić, K. Phytochemical investigation of coumarines, 3rd Int. Symposium on Separation in BioSciences, Abstracts, Moscow, P85, 193, 2003
13. European Pharmacopeia, 3rd editon; Council of Europe, Strazburg, 1997
14. The United States Pharmacopeia XXII; The United states Pharmacopeial convention, Inc., Rockville, 1990