

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

AD-A024 604

THREE-DIMENSIONAL STRUCTURE OF THE EXTENDED SOLAR
MAGNETIC FIELD AND THE SUNSPOT CYCLE VARIATION IN
COSMIC RAY INTENSITY

STANFORD UNIVERSITY

PREPARED FOR
OFFICE OF NAVAL RESEARCH

MARCH 1976

THE EFFECTS OF THE
EXTENDED SOLAR MAGNETIC FIELD
AND THE SOLAR CYCLE VARIATION
ON COSMIC RAY INTENSITY


by
Lod Swegund
and
John M. Wilcox

March 1976

Reproduction in whole or in part
is authorized for any purpose of
the United States Government.

SLRP Report No. 252

Office of Naval Research
Contract N00014-76-C-0001
National Aeronautics and
Space Administration Grant
NGL-05-030-001, National
Science Foundation Grant
ATM76-1907 and Grant
DEB75-1901, and The Max
K. Pritzker Foundation



INSTITUTE FOR PLASMA RESEARCH
STANFORD UNIVERSITY, STANFORD, CALIFORNIA

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER SUIPR Report No. 562	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) Three-Dimensional Structure of the Extended Solar Magnetic Field and the Sunspot Cycle Variation in Cosmic Ray Intensity.		5. TYPE OF REPORT & PERIOD COVERED Scientific; Technical	
7. AUTHOR(s) Leif Svalgaard and John M. Wilcox		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Institute for Plasma Research Stanford University Stanford, California 94305		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0207	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Electronic Program Office Arlington, Virginia 22217		12. REPORT DATE March 1976	13. NO. OF PAGES 16
14. MONITORING AGENCY NAME & ADDRESS (if diff. from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this report) This document has been approved for public release and sale; its distribution is unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)			
18. SUPPLEMENTARY NOTES TECH; OTHER			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Cosmic rays Solar magnetic field Interplanetary magnetic field			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A principal cause for the eleven-year sunspot cycle variation in the primary cosmic ray intensity observed at earth may be a variation in the solid angle of the heliosphere occupied by the extended solar polar magnetic fields. We assume that galactic cosmic rays have relatively easy access to the inner solar system through the regular extended solar polar fields, and relatively difficult access through the irregular extended solar sector structure fields.			

THREE-DIMENSIONAL STRUCTURE OF THE EXTENDED SOLAR MAGNETIC FIELD
AND THE SUNSPOT CYCLE VARIATION IN COSMIC RAY INTENSITY

by

Leif Svølgaard
and
John M. Wilcox

Office of Naval Research
Contract N00014-76-C-0207

National Aeronautics and Space Administration
Grant NGR 05-020-559

National Science Foundation
Grant ATM74-19007
Grant DES75-15664

and


The Max C. Fleischmann Foundation

SUIPR Report No. 662

March 1976

Institute for Plasma Research
Stanford University
Stanford, California

Submitted to: Nature

ADDITIONAL TO	
NTIS	Work Order <input checked="" type="checkbox"/>
DOC	Buy Order <input type="checkbox"/>
UNPROCESSED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	APRIL 1976
	

Three-Dimensional Structure of the Extended Solar Magnetic Field
and the Sunspot Cycle Variation in Cosmic Ray Intensity

by

Leif Svalgaard

John M. Wilcox

Institute for Plasma Research

Stanford University

Stanford, California

The interplanetary magnetic field within several astronomical units of the sun appears to have one polarity in most of the hemisphere north of the solar equatorial plane and the opposite polarity in most of the hemisphere south of the equatorial plane.¹⁻⁷ The two hemispheres are separated by a curved current sheet that typically crosses the solar equatorial plane in either two or four places, thus dividing the equatorial region into either two or four sectors. Near sunspot minimum, at one astronomical unit the extent in latitude of the curved current sheet is typically $\pm 15^\circ$, so that the sector boundary (the current sheet separating the two hemispheres of opposed field polarity) is almost parallel to the solar equatorial plane. In the photosphere, on the other hand, the sector boundary makes an angle of approximately 90° with the equatorial plane.⁸ At 1.5 solar radii, in 1972 and 1973, the angle between the sector boundary and the equatorial plane was approximately 45° ,⁹ and at 3 to 10 solar radii the angle between boundary and plane was approximately 25° .¹⁰ A schematic of this structure for the case of four sectors is shown in Figure 1.

In the photosphere, near sunspot minimum, the sector magnetic fields cover a range in latitude of typically $\pm 40^\circ$,⁸ while at one astronomical unit the comparable range in latitude has been compressed to perhaps $\pm 15^\circ$. How is this compression in latitude accomplished? A typical magnitude of the sector magnetic fields in the photosphere is 0.5 gauss (P.H. Scherrer and T.L. Duvall, personal communication). This is a measure of the large-scale field that will dominate in the region a few solar radii above the photosphere,

where the smaller-scale but much stronger fields associated with active regions do not reach. In the polar regions of the sun the large-scale uni-directed photospheric field has a typical magnitude of perhaps 5 gauss (R. Howard, personal communication). Thus, at one or two solar radii above the photosphere, and above the region of influence of active regions, the magnetic pressure associated with the solar polar regions is two orders of magnitude larger than the magnetic pressure associated with the equatorial sector structure. This will compress the equatorial sector field structure into a narrow range of latitudes. This effect can be clearly seen in the sketch in Figure 2 made from an eclipse photograph taken at solar minimum in 1954.¹¹

During an interval near sunspot maximum, when the solar polar field polarities are reversing, the magnitude of the polar fields may be considerably less, and the resulting compression of the equatorial field structure also much less. The sector structure fields may then occupy a much larger fraction of the heliosphere. Since the sector structure fields reverse polarity typically four times or two times per solar rotation, a galactic cosmic ray headed toward the sun may encounter considerably more magnetic scattering from the complex sector structure field than from the uni-directional field that fills most of each solar hemisphere near sunspot minimum. This geometrical effect may be the principal cause of the eleven-year modulation of cosmic ray intensity observed at earth, since the solar wind velocity¹² and the magnitude of the interplanetary field¹³ observed near earth have not changed very much during the present sunspot cycle.

The fraction of the heliosphere occupied by sector structure fields as a function of time through an average sunspot cycle can be estimated in the following way. Because the current sheet shown in Figure 1 is warped with respect to the solar equatorial plane, during the half-year when the earth is north of the equatorial plane an interplanetary sector with the same polarity as the northern solar polar region is observed to be wider than it would be if observed when the earth is in the equatorial plane of

the sun. For example, assume that there are two sectors per solar rotation and that the extent in heliographic latitude of the sector structure near the earth is $\pm 20^\circ$. When the earth is at a latitude near 7°N (i.e. near September 7), the sector whose polarity is the same as the northern polar region will be observed to last 16.5 days, as compared with the 13.5 days it would last if observed when the earth is near the solar equatorial plane. For comparison, if the extent in heliographic latitude of the sector structure near the earth is $\pm 45^\circ$, then the sector with the same polarity as the northern polar region will have a length of 14.6 days when observed near the earth at 7° north. We see that if we measure the magnitude of this Rosenberg-Coleman effect^{14,15} through a sunspot cycle we can estimate the extent in heliographic latitude of the sector structure.

We have used a harmonic analysis to compute the average amplitude of the Rosenberg-Coleman effect as a function of years from the time of sunspot minimum for the four sunspot cycles whose minima were near 1934, 1944, 1954 and 1965. Interplanetary field polarities inferred from polar geomagnetic variations¹⁶ were used, and the resulting amplitude of the Rosenberg-Coleman effect was multiplied by 1.43 to correct for the approximately 85% accuracy^{17,18} of the inferred interplanetary field polarities.

Figure 3 shows the resulting value for the extent in heliographic latitude of the sector structure through an average sunspot cycle. Three-year running means were used to reduce the scatter. Year 0 is the average of the sunspot minimum years 1934, 1944, 1954 and 1965. The effect of uncertainties in the computation of the magnitude of the Rosenberg-Coleman effect will usually be to decrease the amplitude of the effect and, therefore, the latitude values shown in Figure 3 should be considered upper limits. We may expect considerable variation from the average effect shown in Figure 3, and in particular for intervals of several months near sunspot minimum the current sheet may occupy only a few degrees of latitude. As a first approximation to the sunspot cycle variation of cosmic ray intensity observed at earth resulting from the varying geometry described by Figure 3, we may consider that galactic cosmic rays have relatively difficult access to the inner solar system in the portion of the heliosphere occupied by the changing fields of the sector structure, and relatively easy access through the portions of the heliosphere

occupied by the extended solar polar fields. In Figure 4 we show the solid angle of the heliosphere occupied by the extended solar polar fields through an average sunspot cycle, where the latitude angles shown in Figure are used to compute the solid angles shown in Figure 4.

Also shown in Figure 4 are the monthly averages of the absolute intensity of primary cosmic rays of rigidity greater than 0.5 GV observed near Murmansk and at Mirny from 1958 to 1973.¹⁹ The total flux of such galactic cosmic rays in the interstellar medium in the vicinity of the earth is estimated to be $4000 \text{ m}^{-2} \text{ sec}^{-1} \text{ ster}^{-1}$.¹⁹ Therefore, in Figure 4 we have set the total solid angle 4π of the heliosphere as equivalent to a flux of $4000 \text{ m}^{-2} \text{ sec}^{-1} \text{ ster}^{-1}$, and the zero of the solid angle scale corresponds to zero flux. This corresponds to the assumption that galactic cosmic rays have easy access to the inner solar system through the solid angle of the heliosphere occupied by the extended solar polar fields, and difficult access through the solid angle occupied by the sector structure fields.

In Figure 4 we see that the average sunspot cycle variation of the solid angle of the extended solar polar fields is rather similar to the observed variation of the flux of primary cosmic rays of rigidity greater than 0.5 GV during 1961 to 1969. We should not expect a detailed agreement between the computed variation of solid angle averaged over four sunspot cycles and the observed cosmic ray flux around a single sunspot minimum. The similarity of the two curves in Figure 4 suggests that there may be some validity to the considerations advanced in this paper. The detailed computation of the diffusion lengths of cosmic rays related to these considerations is beyond the scope of this paper.

Acknowledgements

We thank the NASA-Skylab Workshop on Coronal Holes for helpful discussions. This work was supported in part by the Office of Naval Research under Contract N00014-76-C-0207, by the National Aeronautics and Space Administration under Grant NGR 05-020-559, by the Atmospheric Sciences

Section of the National Science Foundation under Grants ATM74-19007 and
DES75-15664, and by the Max C. Fleischmann Foundation.

References

- 1 Schatten K., Solar Wind (proceedings Solar Wind Conference, March 21-26, 1971, Asimolar Conference Grounds, Pacific Grove, California) 65-92; ed C.R. Sonnett, P.J. Coleman and J.M. Wilcox; Scientific and Technical Office, National Aeronautics and Space Administration (1972).
- 2 Schultz, M., Astrophys. and Space Sci., 24, 371-383 (1973).
- 3 Sawyer, C., Geophys. Res. Let., 1, 295-297 (1974).
- 4 Alfvén, H., Proceedings, Nobel-Symposium on the Physics of Hot Plasma in the Magnetosphere, Kiruna, Sweden, 1975.
- 5 Levy, E.H., Proceedings, 14th International Cosmic Ray Conference, 15-29 August 1975, Vol. 4 1215-1220 (1975). Max-Planck-Institut für Extraterrestrische Physik, München, Germany.
- 6 Svalgaard, L., Wilcox, J.M., Scherrer, P.H. and Howard, R., Solar Phys., 45, 83-91 (1975).
- 7 Saito, T., Sci. Rep. Tohoku Univ., Ser. 5, Geophysics 23, 37-54 (1975).
- 8 Wilcox, J.M., Space Sci. Rev., 8, 258-328 (1968).
- 9 Hansen, S. Sawyer, C. and Hansen, R.T., Geophys. Res. Let., 1, 13-15 (1974).
- 10 Howard, R.A. and Koomen, M.J., Solar Phys., 37, 469-475 (1974).
- 11 Vsekhsvjatsky, S.K., The Solar Corona (Proceedings of International Astronomical Union Symposium No. 16; Cloudcroft, New Mexico, 28-30 August 1961), 271-280; ed. J.W. Evans; Academic Press, New York (1963).
- 12 Diodato, L., Moreno, G., Signorini, C. and Ogilvie, K.W., J. Geophys. Res., 79, 5095-5108 (1974).

- 13 Barouch, E. and King, J.H., Proceedings, 14th International Cosmic Ray Conference, 15-29 August 1975, Vol. 3, 1036-1041 (1975).
Max-Planck-Institut fur Extraterrestrische Physik, Munchen, Germany.
- 14 Rosenberg, R.L. and Coleman, P.J., Jr., J. Geophys. Res., 74, 5611-5622 (1969).
- 15 Wilcox, J.M. and Scherrer, P.H., J. Geophys. Res., 77, 5385-5388 (1972).
- 16 Svalgaard, L., J. Geophys. Res., 77, 4027-4034 (1972).
- 17 Russell, C.T. and Rosenberg, R.L., Solar Phys., 37, 251-256 (1974).
- 18 Wilcox, J.M., L. Svalgaard and P.C. Hedgecock, J. Geophys. Res., 80, 3685-3688 (1975).
- 19 Vernov, S.O., Charakhchyan, A.N., Stozhkov, Ju.I. and Charakhchyan, T.N., Proceedings, 14th International Cosmic Ray Conference, 15-29 August 1975, Vol. 3, 1015-1024 (1975). Max-Planck-Institut fur Extraterrestrische Physik, Munchen, Germany.

Figure Captions

Figure 1 Schematic showing the warped current sheet in the inner solar system (inside 6 AU). This current sheet divides the interplanetary magnetic field in the heliosphere into two regions with oppositely directed field lines. In one region the field polarity is away from the sun (at present this region is north of the solar equator), in the other region the field polarity is toward the sun. The situation is shown for a four-sector structure, i.e. as the current sheet is rotated past a stationary observer in the course of a solar rotation, the observer will see four changes of magnetic polarity, suggesting that the interplanetary magnetic field is divided into four sectors of alternating polarity. Where the current sheet lies above the solar equatorial plane it is shown by full lines, while dashed lines indicate that the current sheet is below the equatorial plane. The extent in latitude of the current sheet was assumed to be $\pm 15^\circ$. The sun at the center is not shown to scale.

Figure 2 The structure of a sunspot minimum solar corona drawn from eclipse photographs¹¹ (30 June 1954) obtained in Kozeletsk.

Figure 3 Computed variation of the average extent in heliographic latitude of the extended solar sector magnetic fields. Year 0 is the average of the sunspot minimum years 1934, 1944, 1954 and 1965. The value 20° on the ordinate means that the extended sector fields are in the interval 20°N to 20°S , etc.

Figure 4 Galactic cosmic ray intensities with rigidity 0.5 GV observed near Murmask and at Mirny from 1958 to 1973.¹² Also shown is the computed average solid angle of the heliosphere occupied by the extended solar polar magnetic fields.




Figure 1




Figure 2




Figure 3




Figure 4