

Three new indole alkaloids from *Rauvolfia yunnanensis*

Yuan GAO,^{a,b,c} Fei WANG,^{a,b} Dong-Sheng ZHOU,^b Yan LI,^a and Ji-Kai LIU^{a,*}

^aState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China

^bBioBioPha Co., Ltd., Kunming 650201, China

^cGraduate University of Chinese Academy of Sciences, Beijing 100049, China

Received 11 October 2011; Accepted 23 November 2011

© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract: One rare tetracyclic macroline-type indole alkaloid, named rauvoyunine A (**1**), and two new picaline-type alkaloids rauvoyunines B and C (**2** and **3**) were isolated from the aerial parts of *Rauvolfia yunnanensis*. Their structures were elucidated on the basis of extensive spectroscopic analysis. Compounds **2** and **3** were evaluated for their *in vitro* cytotoxicity against five human tumor cell lines.

Keywords: *Rauvolfia yunnanensis*, indole alkaloid, rauvoyunine

Introduction

The plants of genus *Rauvolfia*, widely distributed in America, Africa, Asia, and Oceania,¹ are well-known rich sources of unique heterocyclic alkaloids with monoterpene indole skeletons. These alkaloids have attracted great interests from biological and therapeutic aspects, due to their anticancer,² antimalarial,³ antihypertensive,⁴ and sedative⁵ properties.

Rauvolfia yunnanensis Tsiang, belonging to the Apocynaceae, is indigenous to southwestern China, and its roots are utilized locally for the remedy of scab¹. Recently, several structurally interesting indole alkaloids were isolated from this plant.^{6–8} As one part of our research program exploring bioactive monoterpene indole alkaloids from Chinese species of *Rauvolfia*, phytochemical analysis from the methanol extract of aerial parts of *R. yunnanensis* led to the isolation of three new alkaloids, rauvoyunines A–C (**1**–**3**). This paper reports the isolation and structure elucidation of new compounds and cytotoxic evaluation of selected compounds.


Results and Discussion

Compound **1**, obtained as amorphous powder, has a molecular formula of C₂₁H₂₈N₂O₃ based on HRESIMS (pos.), showing a quasi-molecular ion peak at *m/z* 357.2185 (calc. for C₂₁H₂₉N₂O₃, 357.2178). The UV spectrum showed absorptions at λ_{max} 228, 282, and 303 nm typical of a substituted indole chromophore.⁹ The IR spectrum showed absorptions at 3432 and 1628 cm⁻¹, attributed to hydroxy and olefin groups,

respectively. The ¹H NMR spectrum (Table 1) showed signals for an aromatic AMX spin system at δ_H 6.73 (dd, *J* = 8.7 and 2.3 Hz), 6.85 (d, *J* = 2.3 Hz), and 7.17 (d, *J* = 8.7 Hz), typical of an indole moiety substituted by a hydroxy group at 10 or 11 position, signals characteristic of an ethylidene group at δ_H 1.44 (d, *J* = 6.8 Hz) and 5.58 (q, *J* = 6.8 Hz), singlets for N-methyl groups at δ_H 2.55 and 3.59, and resonances of two hydroxymethyl groups, one at δ_H 3.75 (dd, *J* = 10.5 and 3.7 Hz) and 3.91 (dd, *J* = 10.5 and 5.5 Hz), and another at δ_H 4.02 (2H, s). The ¹³C NMR (DEPT) spectrum (Table 1) exhibited six sp² quaternary carbons (δ_C 151.8, 140.3, 134.0, 132.3, 127.8, and 106.2), four sp² methines (δ_C 124.8, 112.2, 110.6, and 103.4), four sp³ methines (δ_C 58.7, 55.5, 47.7, and 30.2), four sp³ methylenes (δ_C 66.0, 64.1, 30.5, and 23.7), and three methyl groups (δ_C 41.5, 29.3, and 13.1). The above-mentioned data, together with the presence of nine degrees of unsaturation suggested that **1** was an indole alkaloid with tetracyclic macroline skeleton.^{10,11} This conclusion is supported by the HMBC (Figure 1) three-bond correlations of H-5 to C-3, C-15, and C-17, H-16 to C-6 and C-14, and H-15 to C-3, C-17, and C-19. In addition, the *m*-coupling doublet at δ_H 6.85 (*J* = 2.3 Hz) showed an HMBC correlation with C-7 and a ROESY correlation with H-6β, indicating the hydroxyl at C-10. HMBC cross-peaks of H-18 to C-19 and C-20, and H-19 to C-15 and C-20 located the ethylidene side chain at C-20, and cross-peaks of H-17 to C-5, C-15, and C-16, and H-21 to C-15, C-19, and C-20 revealed that the two hydroxymethyl group was adjacent to C-16 and C-20, respectively. The position of each methyl group was confirmed by HMBC correlations from N(1)-Me to C-2 and C-13, and N(4)-Me to C-3 and C-5.

The relative stereochemistry of **1** was elucidated by ROESY experiments (Figure 1). The ROESY spectrum exhibited correlations of H-16↔H-6β and H-3↔N(4)-Me↔H-5, fixing the two bridgehead hydrogen atoms as α orientation, revealing

*To whom correspondence should be addressed. E-mail: jkliu@mail.kib.ac.cn



the stereochemistry of H-16 as β -oriented. The ROESY correlation between H-14 α and the hydroxymethyl proton at C-20 indicated that C-20 was α -oriented and H-15 was β -oriented. The ROESY correlations of Me-18 \leftrightarrow H-15 and H-19 \leftrightarrow H-21 confirmed that the ethylidene side chain had an *E*-configuration. Consequently, the structure of compound **1** was determined as shown, named rauvoyunine A.




Figure 1. Key HMBC and ROESY correlations of **1**.

Compound **2** was obtained as amorphous powder. Its molecular formula was determined to be $C_{23}H_{26}N_2O_6$ on the basis of HRESIMS (pos.), showing a quasi-molecular ion peak at m/z 427.1868 (calc. for $C_{23}H_{27}N_2O_6$, 427.1869). The UV spectrum showed maximum absorption at 292 nm, and the IR spectrum implied the presence of carbonyl (1738 cm^{-1}) functionality. The ^1H NMR spectra (Table 1) showed the presence of three aromatic protons, an ethylidene side chain, a methyl ether, and an acetyl. The ^{13}C NMR spectra (Table 1) revealed 23 carbon signals due to six sp^2 quaternary carbons, four sp^2 methines, three sp^3 quaternary carbons, three sp^3 methines, four sp^3 methylenes, and three methyl groups. These spectroscopic features suggested that **2** had a picaline-type skeleton.^{13,14} The three observed aromatic resonances at δ_{H} 7.11 (d, $J = 8.8$ Hz), 6.20 (dd, $J = 8.8, 2.2$ Hz), and 6.19 (d, $J = 2.2$ Hz) combined with the fact that a hydroxylated carbon signal was detected at δ_{C} 158.6 (s) suggested that an OH group located at C-10 or C-11. Finding HMBC correlation from the *o*-coupling doublet at δ_{H} 7.11 ($J = 8.8$ Hz) to C-7 at δ_{C} 53.2 (s) and ROESY correlation between the doublet and H-6 α further confirmed placement of hydroxyl at C-11. Two geminally coupled proton signals at δ_{H} 3.85 and 4.52 (each d, $J = 11.1$ Hz) showed correlations with an *O*-acetyl carbonyl carbon at δ_{C} 171.9, indicating that an acetoxy group was located at C-17. An methoxy singlet at δ_{H} 3.66 showed correlation to C-22 at

δ_{C} 173.7, allowing assignment of methyl ether at C-22 position. The ROESY cross-peak between H-17 α and H-14 β indicated that C-17 was β -oriented, which was further supported by the presence of an NMR signal of the acetyl methyl group at δ_{H} 1.62 (the noticeable upfield shift being due to the shielding by the aromatic ring current).¹⁵ ROESY cross-peaks of H-14 α \leftrightarrow H-21 α \leftrightarrow H-3 and H-21 β \leftrightarrow H-5 revealed that the configuration of H-3 and H-5 were α and β , respectively. The ethylidene moiety was assigned to be *E*-type since the correlations of H-18 \leftrightarrow H-15 and H-19 \leftrightarrow H-21 were observed. Therefore, the structure of **2** was unambiguously elucidated as 11-hydroxypicaline, named rauvoyunine B.

Compound **3**, obtained as amorphous powder, possessed a molecular formula of $C_{32}H_{36}N_2O_9$, as evidenced by HRESIMS (pos.) at m/z 593.2496 (calc. for $C_{32}H_{37}N_2O_9$, 593.2499), in combination with ^1H and ^{13}C NMR (DEPT) spectra (Table 1), and appropriate for 16 degrees of unsaturation. The UV spectrum showed maximum absorption at 269 nm. Its IR spectrum revealed absorption bands of carbonyl (1722 cm^{-1}). The ^1H NMR data showed signals of three aromatic protons at δ_{H} 6.04 (br. d, $J = 8.3$ Hz), 6.26 (br. s), and 7.29 (d, $J = 8.3$ Hz), an ethylidene group at δ_{H} 1.62 (d, $J = 6.7$ Hz) and 5.48 (q, $J = 6.7$ Hz), two methoxy groups at δ_{H} 3.45 (s) and 3.64 (s), and a 3',4',5'-trimethoxybenzoyloxy unit, based on the presence of two aromatic protons at δ_{H} 6.94 (2H, s) and three overlapped methoxy groups at δ_{H} 3.87 (9H, s). The ^{13}C NMR (DEPT) data exhibited ten sp^2 quaternary carbons, six sp^2 methines, three sp^3 quaternary carbons, three sp^3 methines, four sp^3 methylenes, and six methyl groups. These spectroscopic features were closely related to those of **2**, but it was evident that **3** had a trimethoxybenzoyloxy unit instead of the acetoxy moiety at C-17 and a methoxyl group instead of the hydroxy group at C-11. This was also supported by HMBC correlations from H-17 to C-7' and correlation from methoxy protons at δ_{H} 3.45 (s) to C-11 at δ_{C} 160.1. The ROESY cross-peaks of Me-18 \leftrightarrow H-15 and H-19 \leftrightarrow H-21 confirmed that the ethylidene possessed an *E*-configuration. The cross-peak of 17 α \leftrightarrow H-14 β indicated that the stereochemistry of C-16 was identical to that of **2**. Since the $J_{3\ 14}$, $J_{14\ 15}$, and $J_{5\ 6}$ values were essentially unchanged compared to **2**, it is safe to deduce that the ring junction stereochemistry remain intact. Thus, the structure of **3** was established to be 11-methoxyburnamine-17-*O*-3',4',5'-trimethoxybenzoate and named as rauvoyunine C.

Compounds **2** and **3** were evaluated for their cytotoxicities against five human cancer cell lines (HL-60, SMMC-7721, A-549, MCF-7, and SW-480) using the MTT method as

Table 1. ^1H NMR and ^{13}C NMR spectroscopic data for rauvoyunines A–C (1–3) (δ in ppm, J in Hz).

No.	1 ^a		2 ^a		3 ^b	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}
2		132.3 (s)		109.0 (s)		107.5 (s)
3	4.39 (dd, 4.2, 2.4)	55.5 (d)	3.48 (br. d, 3.1)	52.7 (d)	3.98 (br. s)	51.9 (d)
5	3.72 (d, 7.2)	58.7 (d)	4.72 (d, 2.8)	88.1 (d)	5.15 (br. s)	87.4 (d)
6 α	3.35 (dd, 16.6, 7.2)		2.29 (dd, 14.2, 2.8)		2.52 (br. d, 14.7)	
6 β	2.63 (d, 16.6)	23.7 (t)	3.17 (d, 14.2)	45.0 (t)	3.17 (d, 14.7)	42.3 (t)
7		106.2 (s)		53.2 (s)		51.9 (s)
8		127.8 (s)		126.1 (s)		124.6 (s)
9	6.85 (d, 2.3)	103.4 (d)	7.11 (d, 8.8)	128.6 (d)	7.29 (d, 8.3)	127.4 (d)
10		151.8 (s)	6.20 (dd, 8.8, 2.2)	108.0 (d)	6.04 (br. d, 8.3)	106.3 (d)
11	6.73 (dd, 8.7, 2.3)	112.2 (d)		158.6 (s)		160.1 (s)
12	7.17 (d, 8.7)	110.6 (d)	6.19 (d, 2.2)	99.5 (d)	6.26 (br. s)	98.2 (d)
13		134.0 (s)		151.7 (s)		149.1 (s)
14 α	2.58 (ddd, 13.7, 13.3, 4.2)		2.00 (ddd, 15.1, 4.0, 3.1)			
14 β	1.73 (ddd, 13.7, 4.0, 2.4)	30.5 (t)	1.95 (br. d, 15.1)	22.5 (t)	2.11 (2H, s)	21.0 (t)
15	2.98 (ddd, 13.3, 5.6, 4.0)	30.2 (d)	3.36 (br. s)	36.3 (d)	3.46 (br. s)	34.5 (d)
16	1.86 (ddd, 5.6, 5.5, 3.7)	47.7 (d)		57.4 (s)		56.6 (s)
17a	3.91 (dd, 10.5, 5.5)	64.1 (t)	4.52 (d, 11.1)	68.2 (t)	4.61 (d, 11.2)	66.7 (t)
17b	3.75 (dd, 10.5, 3.7)		3.85 (d, 11.1)		4.17 (d, 11.2)	
18	1.44 (d, 6.8)	13.1 (q)	1.59 (d, 6.5)	13.7 (q)	1.62 (d, 6.7)	13.4 (q)
19	5.58 (q, 6.8)	124.8 (d)	5.43 (q, 6.5)	122.4 (d)	5.48 (q, 6.7)	123.6 (d)
20		140.3 (s)		137.7 (s)		132.0 (s)
21 α			3.72 (br. d, 17.7)		3.90 (br. d, 17.5)	
21 β	4.02 (2H, s)	66.0 (t)	3.20 (d, 17.7)	46.9 (t)	3.39 (d, 17.5)	45.9 (t)
22				173.7 (s)		171.9 (s)
NMe ₍₁₎	3.59 (s)	29.3 (q)				
NMe ₍₄₎	2.55 (s)	41.5 (q)				
COOMe			3.66 (s)	52.1 (q)	3.64 (s)	51.7 (q)
OMe ₍₁₁₎					3.45 (s)	55.0 (q)
CH ₂ COO			1.62 (s)	20.2 (q)		
CH ₂ COO				171.9 (s)		
OMe _(3', 5')					3.87 (s)	56.1 (q)
OMe _(4')					3.87 (s)	60.8 (q)
1'						124.3 (s)
2', 6'						106.7 (d)
3', 5'					6.94 (s)	152.5 (s)
4'						142.0 (s)
7'						164.6 (s)

^aRecorded in CD₃OD. ^bRecorded in CDCl₃.

reported previously,¹⁵ however, all tested compounds were inactive, and they showed IC₅₀ values > 40 μM .

Experimental Section

General Experimental Procedures. Optical rotations were measured on a Jasco P-1020 automatic digital polarimeter. IR spectra were obtained using a Bruker Tensor 27 FT-IR spectrometer with KBr pellets. NMR spectra were acquired with a Bruker DRX-500 instrument at room temperature. ESIMS (including HRESIMS) were measured on API QSTAR Pulsar i mass spectrometers. Silica gel (200–300 mesh, Qingdao Marine Chemical Inc., China) and Sephadex LH-20 (Amersham Biosciences, Sweden) were used for column chromatography. Fractions were monitored by TLC (Qingdao Marine Chemical Inc., China) in combination with reversed-phase HPLC (Agilent 1200, Extend-C18 column, 5 μm , 4.6 \times 150 mm).

Plant Material. The aerial parts of *R. yunnanensis* were collected in Xishuangbanna of Yunnan Province, China, in May 2008 and were identified by Prof. Yu Chen of Kunming Institute of Botany, Chinese Academy of Sciences. The voucher specimen was deposited at BioBioPha Co., Ltd.

Extraction and Isolation. The air-dried and powdered

aerial parts of *R. yunnanensis* (6.5 kg) were extracted three times with methanol (3 \times 50 L, each 3 days) at room temperature and filtered. The filtrate was evaporated under reduced pressure to get a residue (~480 g), which was fractionized by silica gel (200–300 mesh) CC, eluted with a gradient solvent system of petroleum ether-acetone and then MeOH to yield five fractions A–E. Fraction D, eluted by 100% acetone, was separated on silica gel (CHCl₃-MeOH, 100:1 \rightarrow 30:1) to give two subfractions D1 and D2, Fr. D2 was further isolated and purified by Sephadex LH-20, silica gel (CHCl₃-MeOH-ammonia, 100:1:0.1) to afford **3** (28 mg). Fraction E, eluted by 100% methanol, was separated on silica gel (CHCl₃-MeOH, 30:1 \rightarrow 10:1) to give four subfractions E1–E4, Fr. E4 was further isolated and purified by Sephadex LH-20, and then silica gel (CHCl₃-MeOH-ammonia, 30:1:0.1) to afford **1** (40 mg) and **2** (11 mg). The retention times (t_{R}) of **1–3** from analysis-type HPLC (50% \rightarrow 100% MeOH in H₂O over 6.0 min followed by 100% MeOH to 10 min, 1.0 ml/min, 20°C) were 5.8, 4.7, and 6.6 min, respectively.

Rauvoyunine A (1): yellowish, amorphous powder; $[\alpha]_{\text{D}}^{25}$ + 5.5 (c 0.10, MeOH); UV (MeOH) λ_{max} : 216 (sh), 228, 282, 303, 316 (sh) nm; IR (KBr) ν_{max} 3432, 2923, 1628, 1467, 1380, 1158, 1134 cm^{-1} ; ^1H and ^{13}C NMR data see Table 1; ESIMS (pos.): m/z 357 [M + H]⁺; HRESIMS (pos.): m/z 357.2185 (calc. for C₂₁H₂₉N₂O₃, 357.2178).

Rauvoyunine B (2): yellowish, amorphous powder; $[\alpha]_D^{25}$ –92.0 (*c* 0.19, MeOH); UV (MeOH) λ_{\max} : 231 (sh), 292 nm; IR (KBr) ν_{\max} 3440, 2951, 1738, 1622, 1454, 1385, 1236, 1146, 1114, 1048 cm^{-1} ; ^1H and ^{13}C NMR data see Table 1; ESIMS (pos.): *m/z* 427 $[\text{M} + \text{H}]^+$; HRESIMS (pos.): *m/z* 427.1868 (calc. for $\text{C}_{23}\text{H}_{27}\text{N}_2\text{O}_6$, 427.1869).

Rauvoyunine C (3): yellowish, amorphous powder; $[\alpha]_D^{25}$ –116.8 (*c* 0.18, MeOH); UV (MeOH) λ_{\max} : 269, 293 (sh) nm; IR (KBr) ν_{\max} 3433, 2945, 1722, 1625, 1590, 1503, 1459, 1384, 1416, 1335, 1226, 1127 cm^{-1} ; ^1H and ^{13}C NMR data see Table 1; ESIMS (pos.): *m/z* 593 $[\text{M} + \text{H}]^+$; HRESIMS (pos.): *m/z* 593.2496 (calc. for $\text{C}_{32}\text{H}_{37}\text{N}_2\text{O}_9$, 593.2499).

Cytotoxicity Assay. The cytotoxicity assay was performed according to the MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] method,¹⁶ by use of the following five human cancer cell lines: human myeloid leukemia HL-60, hepatocellular carcinoma SMMC-7721, lung cancer A-549, breast cancer MCF-7, and colon cancer SW-480. The IC50 values were calculated by Reed and Muench's method.¹⁷

Electronic Supplementary Material

Supplementary material is available in the online version of this article at <http://dx.doi.org/10.1007/s13659-011-0023-7> and is accessible for authorized users.

Acknowledgments

This work was financially supported by National Basic Research Program of China (973 Program) 2009CB522300, the “West Light” program of Chinese Academy of Sciences, and Natural Product Library Program of BioBioPha.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Editorial Committee of Flora Reipublicae Popularis Sinicae. *Flora Reipublicae Popularis Sinicae*; Academic Press: Beijing, 1977; Vol. 63, pp 46–58.
- [2] Bemis, D. L.; Capodice, J. L.; Gorroochurn, P.; Katz, A. E.; Buttyan, R. *Int. J. Oncol.* **2006**, *29*, 1065–1073.
- [3] Wright, C. W.; Phillipson, J. D.; Awe, S. O.; Kirby, G. C.; Warhurst, D. C.; Quetin-Leclercq, J.; Angenot, L. *Phytother. Res.* **1996**, *10*, 361–363.
- [4] Hiwada, K. *Ketsuatsu* **2006**, *13*, 325–333.
- [5] Neuss, N. *Indole and Biogenetically Related Alkaloids*; Academic Press: New York, **1980**; Chapter 17.
- [6] Hu, X. J.; He, H. P.; Zhou, H.; Di, Y. T.; Yang, X. W.; Hao, X. J.; Kong, L. Y. *Helv. Chim. Acta* **2006**, *89*, 1344–1350.
- [7] Cheng, B. H.; Chen, J. C.; Qiu, M. H. *Chin. Chem. Lett.* **2009**, *20*, 1221–1223.
- [8] Wang, H. B.; Liu, X. K. *Chem. J. Chin. Univ.* **2010**, *31*, 2005–2009.
- [9] Sheludko, Y.; Gerasimenko, I.; Kolshorn, H.; Stöckigt, J. *J. Nat. Prod.* **2002**, *65*, 1006–1010.
- [10] Ghedira, K.; Zeches-Hanrot, M.; Richard, B.; Massiot, G.; Le Men-Olivier, L.; Sevenet, T.; Goh, S. H. *Phytochemistry* **1988**, *27*, 3955–3962.
- [11] Kam, T. S.; Choo, Y. M. *J. Nat. Prod.* **2004**, *67*, 547–552.
- [12] Kam, T. S.; Choo, Y. M. *Tetrahedron* **2000**, *56*, 6143–6150.
- [13] Petitfrere-Auvray, N.; Vercauteren, J.; Massiot, G.; Lukacs, G.; Sevenet, T.; Le Men-Olivier, L.; Richard, B.; Jacquier, M. *J. Phytochemistry* **1981**, *20*, 1987–1990.
- [14] Hirasawa, Y.; Arai, H.; Zaima, K.; Oktarina, R.; Rahman, A.; Ekasari, W.; Widyawaruyanti, A.; Indrayanto, G.; Zaini, N. C.; Morita, H. *J. Nat. Prod.* **2009**, *72*, 304–307.
- [15] Pousset, J. L.; Poisson, J.; Olivier, L.; Le Men, J.; Janot, M. M. *C. R. Acad. Sci.* **1965**, *261*, 5538.
- [16] Mosmann, T. *J. Immunol. Methods* **1983**, *65*, 55–63.
- [17] Reed, L. J.; Muench, H. *Am. J. Hyg.* **1938**, *27*, 493–497.