

APR 21 1965

**MASTER**

NYO-3399-31  
1206-SU-31

A THREE-TRIPLET MODEL WITH DOUBLE SU(3) SYMMETRY

~~NOT FOR PUBLIC RELEASE - OFFICIAL  
DISTRIBUTION MAY BE MADE BY OFFICIAL  
REQUESTS MAY BE MADE BY REPORT  
CONTAINS NOTHING OF PATENT INTEREST  
PROCEDURES ON FILE IN RECEIVING  
SECTION~~

M. Y. Han

Department of Physics  
Syracuse University  
Syracuse, New York

and

Y. Nambu

The Enrico Fermi Institute for Nuclear Studies  
and the Department of Physics  
The University of Chicago, Chicago, Illinois

There is no objection from the patent  
point of view to the publication or  
dissemination of this document:  
NY Patent Group  
By *H. S. Potter*  
4/19 1965



Department of Physics  
**SYRACUSE UNIVERSITY**  
Syracuse, New York 13210

~~1965~~

## **DISCLAIMER**

**This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

A THREE-TRIPLET MODEL WITH DOUBLE SU(3) SYMMETRY \*

M. Y. Han

Department of Physics  
Syracuse University  
Syracuse, New York

and

Y. Nambu

The Enrico Fermi Institute for Nuclear Studies  
and the Department of Physics  
The University of Chicago, Chicago, Illinois

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information disclosed in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

This document is  
**PUBLICLY RELEASABLE**

Hugh Kinser Hugh Kinser

Authorizing Official

Date 10/19/68

\* Work supported in part by the U. S. Atomic Energy Commission under the Contract No. AT (30-1) -3399 and No. AT (11-1) -264.

## ABSTRACT

With a view to avoiding some of the kinematical and dynamical difficulties involved in the single triplet quark model, a model for the low lying baryons and mesons based on three triplets with integral charges is proposed, somewhat similar to the two-triplet model introduced earlier by one of us (Y. N.). It is shown that in a  $U(3)$  scheme of triplets with integral charges, one is naturally led to three triplets located symmetrically about the origin of  $I_3$ - $Y$  diagram under the constraint that Nishijima-Gell-Mann relation remains intact. A double  $SU(3)$  symmetry scheme is proposed in which the large mass splittings between different representations are ascribed to one of the  $SU(3)$ , while the other  $SU(3)$  is the usual one for the mass splittings within a representation of the first  $SU(3)$ .

## I. INTRODUCTION

Although the SU(6) symmetry strongly indicates that the baryon is essentially a three-body system built from some basic triplet field or fields, the quark model<sup>1</sup> is not entirely satisfactory from a realistic point of view, because (a) the electric charges are not integral, (b) three quarks in s states do not form the symmetric SU(6) representation assigned to the baryons, and (c) a simple dynamical mechanism is lacking for realizing only zero-triality states as the low lying levels.

These difficulties may be avoided if we introduce more than one basic triplet. Recently one of us (Y. N.) has attempted a two-triplet model<sup>2</sup> where the members of the triplets  $t_1$  and  $t_2$  had the charge assignment (1,0,0) and (0,-1,-1), as had been proposed earlier by Bacry et. al.<sup>3</sup> The baryon would be represented by the combination  $t_1 t_1 t_2$ , whereas the mesons would correspond to some combination  $\sim at_1 \bar{t}'_1 + bt_2 \bar{t}'_2$ . The triplets are assumed to have masses large compared to the baryon mass, which would mean that baryons and mesons have very large binding energies. A dynamical mechanism for this is provided by a neutral field coupled strongly to the "charm number"<sup>4</sup> C, which is 1 for  $t_1$  and -2 for  $t_2$ , and therefore C = 0 for baryons and mesons. In analogy with electrostatic energy, we can argue that the potential energy due to the charm field would be the lowest when the system is "neutral", namely C = 0. Thus all other unwanted configurations with C  $\neq$  0, which include among others triplet, sextet, etc. representations, would have high masses, and hence would not be easily observed.

There have been proposed two different ways in which to introduce basic triplet or triplets with integral charges. One approach essentially involves a modification of the Nishijima-Gell-Mann relation by way of introducing an additional quantum number, the triality quantum number,<sup>5</sup> and this has led to considerations

of higher symmetry schemes based on rank three Lie groups.<sup>6</sup> On the other hand, Okubo et. al.<sup>7</sup> have recently shown that the minimal group required for this purpose is actually the group  $U(3)$ .<sup>8</sup> It is shown that a triplet scheme may be defined in  $U(3)$  such that the triplet always possesses integral values of charge and hypercharge and satisfies the Nishijima-Gell-Mann relation without a modification. The  $U(3)$  triplet considered by Okubo et. al. is of Sakata-type; i.e. it consists of an iso-doublet and an iso-singlet. Actually the  $U(3)$  scheme is much more appealing than those of the rank three Lie groups on two accounts: firstly, the Nishijima-Gell-Mann relation is satisfied universally by triplets as by octets and decuplets and secondly, as far as the hitherto realized representations are concerned,  $U(3)$  is equivalent to  $SU(3)$ .<sup>9</sup>

In what follows, we show that the  $U(3)$  scheme, when fully utilized as described below, naturally and uniquely leads to a set of three basic triplets with integral charges, namely an I-triplet (iso-doublet and iso-singlet), a U-triplet (U-spin doublet and U-spin singlet) and a V-triplet (V-spin doublet and V-spin singlet).<sup>10</sup> These triplets arise from three different ways of defining charge  $Q$ , hypercharge  $Y$ , and a displaced isospin  $I_3$  in the  $U(3)$  group as opposed to the  $SU(3)$ , in such a way that the charge and hypercharge have integral values, while keeping the Nishijima-Gell-Mann relation intact and they differ from each other in their quantum number assignments as well as in their transformation properties under the Weyl reflections.<sup>11</sup> This is described in Section II. In Section III, a double  $SU(3)$  symmetry scheme is proposed based on the three-triplet model in which the large mass splittings between different representations are ascribed to one of the  $SU(3)$  and the other  $SU(3)$  is, as usual, responsible for the mass splittings within a representation. The low lying baryon and meson states may be taken as singlets with respect to one of the  $SU(3)$ . The extended symmetry group with respect to the  $SU(6)$  symmetry is briefly discussed.

II- THREE TRIPLETS

We shall denote the infinitesimal generators of U(3) by  $A_{\nu}^{\mu}$  which satisfies the following commutation relations:

$$\left[ A_{\beta}^{\alpha} , A_{\nu}^{\mu} \right] = \delta_{\beta}^{\mu} A_{\nu}^{\alpha} - \delta_{\nu}^{\alpha} A_{\beta}^{\mu} \quad (1)$$

where all indices take on the values 1, 2 and 3. The corresponding infinitesimal generators  $B_{\nu}^{\mu}$  of SU(3) are then given by

$$B_{\nu}^{\mu} = A_{\nu}^{\mu} - 1/3 \delta_{\nu}^{\mu} A_{\lambda}^{\lambda} \quad (2)$$

which satisfy the following equations:

$$\left[ B_{\beta}^{\alpha} , B_{\nu}^{\mu} \right] = \delta_{\beta}^{\mu} B_{\nu}^{\alpha} - \delta_{\nu}^{\alpha} B_{\beta}^{\mu} \quad (3)$$

and

$$B_{\lambda}^{\lambda} = 0 \quad (4)$$

Furthermore, the unitary restriction gives

$$(A_{\nu}^{\mu})^{\dagger} = A_{\mu}^{\nu} , \quad (B_{\nu}^{\mu})^{\dagger} = B_{\mu}^{\nu} . \quad (5)$$

Let us now briefly summarize the relevant results of Okubo et. al. In the SU(3) scheme, the charge Q, the hypercharge Y and the third component of isospin  $I_3$  are identified as follows:<sup>12</sup>



$$Q = -B_1^1 \quad (6a)$$

$$Y = B_3^3 = -B_1^1 - B_2^2 \quad (\text{by the relation (4)}) \quad (6b)$$

$$I_3 = 1/2 (B_2^2 - B_1^1) \quad (6c)$$

In the U(3) scheme, the corresponding quantities  $\tilde{Q}$ ,  $\tilde{Y}$  and  $\tilde{I}_3$  are defined as follows:

$$\tilde{Q} = -A_1^1 = Q - 1/3 \tau \quad (7a)$$

$$\tilde{Y} = -A_1^1 - A_2^2 = Y - 2/3 \tau \quad (7b)$$

$$\tilde{I}_3 = 1/2 (A_2^2 - A_1^1) = I_3 \quad (7c)$$

where

$$\tau = A_1^1 + A_2^2 + A_3^3 \quad (8)$$

With these definitions, the Nishijima-Gell-Mann relation is seen to be equally satisfied by the U(3) and SU(3) theories, i.e.

$$Q = I_3 + 1/2 Y \quad (9)$$

and

$$\tilde{Q} = \tilde{I}_3 + 1/2 \tilde{Y}, \quad (10)$$

respectively. Since the generators  $A_1^1$ ,  $A_2^2$  and  $A_3^3$  possess integral eigenvalues in any representation<sup>13</sup>, the identifications of  $\tilde{Q}$  and  $\tilde{Y}$  to be the charge and the

hypercharge respectively in U(3) theory shall always lead to integral values for the charge and the hypercharge. In particular, in the three-dimensional representation, the U(3) triplet has the eigenvalues

$$\tilde{Q} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{I}_3 = \begin{pmatrix} \frac{\sqrt{3}}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{Y} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (11)$$

This triplet corresponds to the Sakata triplet which we call an I-triplet for short.

We can now generalize the above constructions of the U(3) triplet in the following way. Comparing (6b) and (7b) we see that a particular choice has been made for  $\tilde{Y}$ . Had we defined  $\tilde{Y}$  to be  $A_3^3$ , it would still have integral eigenvalues but the relation (10) would have been violated. This is because  $B_\lambda^3 = 0$  in SU(3) but  $A_\lambda^3 \neq 0$  in general in U(3) and thus some care is needed in defining corresponding quantities in U(3). Making use of (4), the definition in (6) can be written more generally as:

$$Q = -B_1^1 = B_2^2 + B_3^3 \quad (12a)$$

$$Y = B_3^3 = -B_1^1 - B_2^2 \quad (12b)$$

$$I_3 = 1/2 (B_2^2 - B_1^1) = 1/2 (2B_2^2 + B_3^3) = -1/2 (2B_1^1 + B_3^3) \quad (12c)$$

As in (7), replacing  $B_\mu^\nu$ 's in (12) by corresponding  $A_\mu^\nu$ 's, we list all possible candidates for the corresponding quantities in U(3) which are now however not equivalent to each other (they are equivalent, of course, when reduced to SU(3)), i.e.

$$\tilde{Q}: -A_1^1, \quad A_2^2 + A_3^3 \quad (13a)$$

$$\tilde{Y}: A_3^3, \quad -A_1^1 - A_2^2 \quad (13b)$$

$$\tilde{I}_3: 1/2 (A_2^2 - A_1^1), \quad 1/2 (2A_2^2 + A_3^3), \quad -1/2 (2A_1^1 + A_3^3). \quad (13c)$$

To start with, the alternative choices in (13) provide twelve inequivalent ways in which to choose a set of three quantities  $\tilde{Q}$ ,  $\tilde{Y}$  and  $\tilde{I}_3$  for the U(3) scheme. In every choice  $\tilde{Q}$  and  $\tilde{Y}$  will have integral eigenvalues, but as can be easily checked the Nishijima-Gell-Mann relation will not be valid for all of them. In fact, there are only three cases for which it is valid and we are thus naturally led to three inequivalent triplets in the U(3) scheme; they are defined by the following three choices:

$$t_I: \tilde{Q} = -A_1^1, \quad \tilde{Y} = -A_1^1 - A_2^2, \quad \tilde{I}_3 = 1/2 (A_2^2 - A_1^1) \quad (14a)$$

$$t_U: \tilde{Q} = A_2^2 + A_3^3, \quad \tilde{Y} = A_3^3, \quad \tilde{I}_3 = 1/2 (2A_2^2 + A_3^3) \quad (14b)$$

$$t_V: \tilde{Q} = -A_1^1, \quad \tilde{Y} = A_3^3, \quad \tilde{I}_3 = -1/2 (2A_1^1 + A_3^3). \quad (14c)$$

Now the first one,  $t_I$ , for which

$$\tilde{Y} = -A_1^1 - A_2^2 \quad (15)$$

$$\tilde{I}_3 = 1/2 (A_2^2 - A_1^1) = 1/2 (B_2^2 - B_1^1) = I_3 \quad (16)$$

corresponds to the I-triplet mentioned above.

The structure of the remaining triplets  $t_U$  and  $t_V$  can be brought to much more transparent and symmetric forms in terms of the U-spin and V-spin subalgebras.<sup>10</sup> As in the case of relations (9) and (10) for SU(3) and U(3), we define the U- and V-spin of U(3) in exactly the same forms as in SU(3) except that all quantities are tilded quantities. From the SU(3) definitions<sup>12</sup>, we then have

$$\tilde{Y}_U = -\tilde{Q} = -A_2^2 - A_3^3 \quad (17)$$

$$\tilde{U}_3 = \tilde{Y} - 1/2\tilde{Q} = 1/2 (A_3^3 - A_2^2) = 1/2 (B_3^3 - B_2^2) = U_3 \quad (18)$$

for (14b), and

$$\tilde{Y}_V = \tilde{Q} - \tilde{Y} = -A_3^3 - A_1^1 \quad (19)$$

$$\tilde{V}_3 = -1/2 (\tilde{Y} + \tilde{Q}) = 1/2 (A_1^1 - A_3^3) = 1/2 (B_1^1 - B_3^3) = V_3 \quad (20)$$

for (14c). They correspond, therefore, to a U-triplet and a V-triplet respectively and hence the notations  $t_I$ ,  $t_U$  and  $t_V$ . With respect to the SU(3) triplet (quark), these U(3) triplets have their respective "hypercharges" (i.e.  $Y$ ,  $Y_U$  and  $Y_V$ ) shifted by the amount of  $2/3$  and as such they have quite different transformation properties under the Weyl reflections  $W_1$ ,  $W_2$  and  $W_3$ <sup>11</sup> which are reflections about the axis  $I_3 = 0$ ,  $U_3 = 0$  and  $V_3 = 0$  respectively. Whereas the SU(3) triplet is invariant under all three Weyl reflections, the U(3) triplets are not. They transform according to:

$$W_1: t_I \longrightarrow t_I, \quad t_U \longleftrightarrow t_V \quad (21a)$$

$$W_2: t_U \longrightarrow t_U, \quad t_I \longleftrightarrow t_V \quad (21b)$$

$$W_3: t_V \rightarrow t_V, \quad t_I \leftrightarrow t_U \quad . \quad (21c)$$

Figure 1a and Table 1a list the quantum numbers  $\tilde{I}_3$  and  $\tilde{Y}$  for the single triplet (quark) model; a possible assignment implied by the two-triplet model<sup>2</sup> is shown in Figure 1b and Table 1b; the corresponding quantum numbers for the three-triplet model are given in Figure 1c and Table 1c.

### III. DOUBLE SU(3) SYMMETRY

Let us call the three triplets  $t_1 (=t_I)$ ,  $t_2 (=t_U)$  and  $t_3 (=t_V)$ . Each triplet may be characterized in general by the average values,  $\bar{I}_3$  and  $\bar{Y}$ , of  $\tilde{I}_3$  and  $\tilde{Y}$  for its three members. This specifies the location of the center of the triplet in the  $\tilde{I}_3$ - $\tilde{Y}$  diagram. Since  $\bar{A}_1^1 = \bar{A}_2^2 = \bar{A}_3^3 = \bar{\tau}/3 = \tau/3$ , Eq. (14) gives for the three definitions of  $\tilde{I}_3$  and  $\tilde{Y}$

$$\bar{I}_3 = 0, (1/2)\tau, (-1/2)\tau \quad (22)$$

$$\bar{Y} = (-2/3)\tau, (1/3)\tau, (1/3)\tau$$

respectively, where  $\tau = -1$  for all the triplets. We may define new quantities  $I_3$ ,  $Y$  and  $Q = I_3 + Y/2$  by the relations

$$\begin{aligned} \tilde{I}_3 &= \bar{I}_3 + I_3 \\ \tilde{Y} &= \bar{Y} + Y \end{aligned} \quad (23)$$

$$\tilde{Q} = \bar{I}_3 + (1/2)\bar{Y} + I_3 + (1/2)Y = \bar{Q} + Q$$

It is clear that  $I_3$  and  $Y$  play the role of SU(3) generators within each triplet.

The charm number  $C$  defined in the two-triplet model<sup>2</sup> is then

$$(1/3)C = \bar{Q} = \bar{I}_3 + (1/2)\bar{Y} \quad (24)$$

Now it is interesting to note that according to Eq. (22) and Figure 1c, the centers of the three triplets form an anti-triplet, equivalent to an anti-quark, symmetrically located around the origin. Let us suppose that the nine members of the three triplets  $t_{1\alpha}, t_{2\alpha}, t_{3\alpha}, \alpha=1, 2, 3$  be combined into a single multiplet  $T = \{t_{i\alpha}\}, i = 1, 2, 3$ . We can then imagine two distinct sets of  $SU(3)$  operations on  $T$ . One is the  $SU(3)$  acting on the index  $\alpha$  for each triplet, while the other  $SU(3)$  acts on the index  $i$ , which mixes corresponding members of different triplets.  $T$  is then a representation  $(3, 3^*)$  of this group  $G \equiv SU(3)' \times SU(3)''$ .<sup>14</sup> The quantum numbers of  $SU(3)'$  and  $SU(3)''$  are identified as  $I_3' = I_3, Y' = Y, I_3'' = \bar{I}_3$  and  $Y'' = \bar{Y}$ . in Eq. (22), so that

$$\begin{aligned} \tilde{I}_3 &= I_3' + I_3'', \quad \tilde{Y} = Y' + Y'', \\ \tilde{Q} &= I_3' + I_3'' + (1/2)Y' + (1/2)Y'', \\ (1/3)C &= I_3' + (1/2)Y'' . \end{aligned} \quad (25)$$

A general representation of  $G$  may be characterized by four numbers  $p', q', p'', q''$  so that  $D(p', q', p'', q'') \sim D(p', q') \times D(p'', q'')$  where  $D(p, q)$  is a representation of  $SU(3)$ . However, in our scheme where the nonet  $T$  is the fundamental field, we do not get all the possible representations of  $G$ . This can be illustrated by means of the triality numbers<sup>5</sup>  $t' = p' - q' \pmod{3}, t'' = p'' - q'' \pmod{3}$ . The nonet  $T$  has  $t' = 1, t'' = -1$ . All representations constructed out of  $T$  and  $T^*$  then satisfy  $t' = -t''$ .

Let us next consider the meson and baryon states  $\sim TT^*$  and  $\sim TTT$ . The  $SU(3)' \times SU(3)''$  contents of these 81- and 729-plets are

$$\begin{aligned}
 (3, 3^*) \times (3^*, 3) &= (8, 1) + (1, 1) + (1, 8) + (8, 8), \\
 (3, 3^*) \times (3, 3^*) \times (3, 3^*) & \\
 &= (1, 1) + 2(8, 1) + 2(1, 8) + (1, 10^*) + (10, 1) + 2(8, 10^*) + 2(10, 8) + 4(8, 8) \\
 &\quad + (10, 10^*).
 \end{aligned}
 \tag{26}$$

It is an attractive possibility to postulate at this point that the energy levels are classified according to  $SU(3)''$ . The masses will then depend on the Casimir operators of  $SU(3)''$ . For Example, a simple linear form will be

$$m = m_0 + m_2 C_2'' + m_3 C_3'', \tag{27}$$

where  $C_2''$ ,  $C_3''$  are the eigenvalues of quadratic and cubic Casimir operators of  $SU(3)''$ . In particular we may assume that the main mass splitting comes from  $C_2''$ . Since this increases with the dimensionality of representation, the lowest mass levels will be  $SU(3)''$  singlets. This selects the low lying meson and baryon states to be  $(8, 1)$ ,  $(1, 1)$  and  $(8, 1)$ ,  $(1, 1)$ ,  $(10, 1)$  respectively. In general, all low lying states will have triality zero,  $t' = t'' = 0$ .

As for the baryon number assignment to the triplets, the simplest possibility would be to assign an equal baryon number, i.e.  $B = 1/3$ , to them. In this case the triplets themselves would be essentially stable, and their nine members would behave like an octet plus a singlet of "heavy baryons" as may be seen from Fig. 1c. Another simple possibility may be  $B = 1/3 + Y''$ , namely  $B = (1, 0, 0)$  for  $(t_1, t_2, t_3)$ . We expect a mass splitting depending on  $B$  or  $Y''$ , which may be the origin of the Okubo-Gell-Mann mass formula.

The advantage of the three-triplet model is that the  $SU(6)$  symmetry can be easily realized with s-state triplets. The extended symmetry group becomes now  $SU(6)' \times SU(3)''$ . Since an  $SU(3)''$  singlet is antisymmetric, the overall Pauli principle requires the baryon states to be the symmetric  $SU(6)$  56-plet. Other

SU(6) representations such as the 70, will be obtained by bringing in either the orbital angular momentum or the "  $\rho$  -spin" of the Dirac spinor triplets.

As in the two-triplet model mentioned in the Introduction, the mass formula of the type (27) may be derived dynamically. Instead of the charm number field, we introduce now eight gauge vector fields which behave as (1,8), namely as an octet in SU(3)", but as singlets in SU(3)'. Since their coupling to the individual triplets is proportional to  $\lambda_i''$  (the generators of SU(3)"), the interaction energy arising from the exchange of these vector fields will yield the first and second terms of Eq. (27). If these mesons obey again a similar type of mass formula, they will be expected to be massive compared to the ordinary mesons. However, it is not clear whether the resulting short range character of the interaction can be readily reconciled with the postulated largeness of the interaction energy.

We may characterize the hierarchy of interactions and their symmetries implied by the above model as follows. First, the superstrong interactions responsible for forming baryons and mesons have the symmetry SU(3)", and causes large mass splittings between different representations. The scale of mass involved would be comparable or large compared to the baryon mass, namely  $\approx 1$  Bev. The lowest states, i.e. SU(3)" singlet states, would split according to SU(3)', which would be the SU(3) group observed among the known baryons and mesons, with their strong interactions. The scale of mass splitting would then be  $\leq 1$  Bev.

When we go to the massive SU(3)" non-singlet states, there may very well be coupling between the two SU(3) groups similar to the L·S coupling. The levels should be classified in terms of the three sets of Casimir operators formed out of  $\lambda_i'$ ,  $\lambda_i''$  and  $\lambda_i = \lambda_i' + \lambda_i''$  respectively. The splitting due to the coupling would naturally be intermediate between the above two splittings, namely  $\sim 1$  Bev. Because of this coupling, the separate conservation of the two SU(3) spins,



$I_3'$  and  $Y'$  on the one hand, and  $I_3''$  and  $Y''$  on the other, would be destroyed, and only the sums  $I_3 = I_3' + I_3''$  and  $Y = Y' + Y''$  would be conserved under strong interactions. This in turn would mean that all the massive states are in general highly unstable, and decay strongly to the low lying states. (In the two-triplet model, we considered only weak decays of  $C \neq 0$  states. But strong decays are also a possibility as is contemplated here.)

We have discussed here a possible model of baryons and mesons based on three triplets. How can we distinguish this and other different models mentioned already? Certainly different models predict considerably different structure of massive states. These states are characterized by the triality for the quark model, by the charm number for the two-triplet model and by the  $SU(3)''$  representation for the present three-triplet model. If we restrict ourselves to the low lying states only, however, it seems difficult to distinguish them without making more detailed dynamical assumptions.

#### ACKNOWLEDGMENTS

One of us (M. Y. H.) wishes to thank Professor E. C. G. Sudarshan and Professor A. J. Macfalane for their encouragements and useful discussions and Professor L. O'Raifeiraigh and Mr. J. Kuriyan for helpful comments.

REFERENCES AND FOOTNOTES

1. M. Gell-Mann, Phys. Letters 8, 214 (1964); G. Zweig, CERN preprint.
2. Y. Nambu, Proc. of the Second Coral Gables Conference on Symmetry Principles at High Energy, University of Miami, January, 1965.
3. H. Bacry, J. Nuyts and L. van Hove, Phys. Letters 9, 279 (1964).
4. This name was originally used in connection with the SU(4) symmetry. B. J. Björken and S. L. Glashow, Phys. Letters 11, 255 (1964); A. Salam, Dubna Conference Report, 1964.
5. G. E. Baird and L. C. Biedenharn, Proc. of the First Coral Gables Conference on Symmetry Principles at High Energy, W. H. Freeman and Co., San Francisco (1964); C. R. Hagen and A. J. Macfarlane, Phys. Rev. 135, B432 (1964) and J. Math. Phys. 5, 1335 (1964).
6. For example, see I. S. Gerstein and M. L. Whippmann, Phys. Rev. 137, B1522 (1965). Earlier references are given in this paper.
7. S. Okubo, C. Ryan and R. E. Marshak, Nuovo Cimento 34, 759 (1964).
8. The use of U(3) in this connection has also been remarked by I. S. Gerstein and K. T. Mahanthappa, Phys. Rev. Letters 12, 570, 656(E) (1964).
9. S. Okubo, Phys. Letters 4, 14 (1963).
10. C. A. Levinson, H. J. Lipkin and S. Meshkov, Nuovo Cimento 23, 236 (1961); Phys. Letters 1, 44 (1962) and Phys. Rev. Letters 10, 361 (1963).
11. A. J. Macfarlane, E.C.C. G. Sudarshan and C. Dullemond, Nuovo Cimento 30, 845 (1963).
12. We use the sign convention of S. P. Rosen, J. Math. Phys. 5, 289 (1964).
13. For a derivation of this result, See Eq. (7) of Ref. 7.
14. Such a nonet provides a natural basis for the symmetry of SU(9). However, we will not consider it here.

FIGURE CAPTIONS

Figure 1a. The Single Triplet (quark) Model

Figure 1b. The Two-Triplet Model

Figure 1c. The Three-Triplet Model

(a)		quark		
	$\bar{I}_3$	1/2	-1/2	0
	$\bar{Y}$	1/3	1/3	-2/3
	$\bar{Q}$	2/3	-1/3	-1/3

(b)		$t_1$			$t_2$		
	$\bar{I}_3$	1/2	-1/2	0	1/2	-1/2	0
	$\bar{Y}$	1	1	0	-1	-1	-2
	$\bar{Q}$	1	0	0	0	-1	-1

(c)		$t_1(t_I)$			$t_2(t_U)$			$t_3(t_V)$		
	$\bar{I}_3$	1/2	-1/2	0	0	-1	-1/2	1	0	1/2
	$\bar{Y}$	1	1	0	0	0	-1	0	0	-1
	$\bar{Q}$	1	0	0	0	-1	-1	1	0	0

TABLE 1

QUANTUM NUMBER ASSIGNMENTS FOR

- (a) the quark model
- (b) the two-triplet model
- (c) the three-triplet model




FIG. 1a




FIG. 1b




FIG. 1c