

Thyroid Dysfunction in Central Referral Hospital, Sikkim

Sonam Choden Bhutia ^{1*}, Karma Loday Bhutia ¹, T.A. Singh ²

¹PhD, Assistant Professor, Department of Biochemistry, Sikkim Manipal Institute of Medical Sciences.

¹ MS, Assistant Professor, Department of Ophthalmology, Sikkim Manipal Institute of Medical Sciences.

² MD, FIIMSA, Professor, Department of Biochemistry, Sikkim Manipal Institute of Medical Sciences.

Research Article

Article Info:

Received on: 15/04/2016

Published on: 23/05/2016



QR Code for mobile



ABSTRACT :

The thyroid dysfunction is one of the most common endocrine disorders. Sikkim lies in the severely iodine deficient zone. This was a hospital based cross-sectional study. The serum samples were used to check the levels of thyroid stimulating hormone (TSH), triiodothyronine (T₃) and thyroxine (T₄). Out of 674 patients with suspected thyroid dysfunction, 83% were Euthyroid followed by 10 % of patients having subclinical hypothyroidism. The incidence subclinical hyperthyroidism (1%) were lowest. Females were found to be maximum with thyroid disorders. The data were represented as percentage and mean \pm SD. Thyroid hormones were compared among the different thyroid disorder by One way analysis of variance (ANOVA).

High incidence of thyroid dysfunction in females with subclinical hypothyroidism indicates that it still exists as a public health problem in Sikkim regardless of the implementation of iodized salt program since the last decade.

Keywords: Subclinical hypothyroidism, Subclinical hyperthyroidism, Tri-iodothyronine, Thyroid stimulating hormone, Thyroxine.

INTRODUCTION:

The thyroid disorders are the most common endocrine diseases in the world. Globally, India has the largest number of children born vulnerable to iodine-deficiency with 42 million people suffering from thyroid disorders [1, 2]. According to the Iodine status conducted by World Health Organization (WHO) in 2004, classified India as having an optimal iodine nutrition [3]. Sikkim, the small hilly state in the eastern Himalayas lies in the severely iodine deficient zone [4]. In 1989, the state government of Sikkim had implemented Iodine Deficiency Disorder control program in state. Various studies have shown that Iodine Deficiency Disorders is a significant public health problem for the state. Almost all of them have shown high prevalence of goitre and cretinism. According to a survey conducted in 1994, endemic cretinism was observed to be 77.9% in the state [5,6].

Factors like age, sex, ethnicity and geographical locations play an essential role on the prevalence of thyroid disorders and most commonly on iodine uptake [7]. TSH measurement is considered to be highly sensitive and specific for diagnosing hypothyroidism and hyperthyroidism. The relationship between serum TSH and serum thyroxine are logarithmic so an increase in thyroxine level produces a many fold change in TSH [8]. The signs and symptoms of hyper and hypothyroidism are well known and on the other hand subclinical thyroid condition may mimic other

diseases. So it is important to develop laboratory strategies to differentiate the various conditions for correct diagnosis and treatment [9].

The study aims to investigate the incidence of thyroid dysfunction in the local population visiting the outpatient department of Central Referral Hospital (CRH), Gangtok, Sikkim.

MATERIALS AND METHODS:

Study design

This was a hospital-based cross-sectional study conducted in the department of Biochemistry, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim. The duration of the study was the month of October 2013 to May 2014.

Sources of data

The suspected subjects with thyroid disorder were selected from the outpatient department, Central Referral Hospital, Gangtok, Sikkim for the study.

Sample collection:

Fasting venous blood samples (2ml) were collected from 643 subjects. The blood sample were collected in a plain vial, centrifuged at 3000 rpm for 15 minutes and the separated serum was stored at -20° C for hormone assay.

Assay

The serum samples were assayed to check the levels of thyroid stimulating hormone (TSH), triiodothyronine (T₃) and thyroxine (T₄) by using standard kits (RFCL, India).

*Corresponding author:

Dr. Sonam Choden Bhutia,

Assistant Professor,

Department of Biochemistry,

Sikkim Manipal Institute of Medical Sciences,

5th mile, Tadong Gangtok, Sikkim-737102

Email id : sonamcho07@hotmail.com

Mobile no: 09735974048

Conflict of interest: Authors reported none

submit your manuscript | www.jbiopharm.com



The procedures were followed as per the kit inserts. T3 and T4 were estimated by competitive enzyme linked immunoassay (ELISA) and TSH by Sandwich ELISA.

The subjects were categorized as euthyroid, subclinical hypothyroidism (Normal T3, T4 and increased TSH), overt hypothyroidism (decreased T3, T4 and increased TSH), subclinical

hyperthyroidism (Normal T3, T4 and decreased TSH), [10] and overt hyperthyroidism (increased T3, T4 and decreased TSH). The subjects were divided in four age groups to determine the occurrence of various thyroid disorders in different age groups.

Statistical Analysis :

Statistical analysis was done using the Statistical Package for the Social Sciences version 16.0 (SPSS Inc, Chicago, IL, USA). The datas were represented as percentage and mean ± SD. Comparison of thyroid hormone levels among the different thyroid disorder was analyzed by One way analysis of variance (ANOVA). P value <0.05 was considered to be significant.

RESULTS:

Table1. represents the normal values of thyroid hormone (T3, T4 and TSH) as per the kit insert.

Table 1. Reference range of different parameters of thyroid profile.

| Parameters | Reference range |
|------------|---|
| Serum T3 | 0.52-1.85ng/ml |
| Serum T4 | Male: 4.4-10.8 ug/dl Female: 4.8-11.6ug/dl |
| Serum TSH | 0.39-6.16 uIU/ml |

Normal values of the parameters as per the kit.

Over all distribution of patients with thyroid disorder is presented in Table 2 and figure 1. In the present study, out of total 674 patients with suspected thyroid disorder, 83% (n=560) subjects were found to be Euthyroid having normal levels of T₃, T₄ and TSH (females = 450, males = 129). 3.5% (n=195) belonged to overt hypothyroidism with elevated TSH levels and low T₃ and T₄ levels (female=19, males = 5). The patients with overt hyperthyroidism were 2.5% (n=17) (females =13, males=4). 10% (n=67) patients were suffering from subclinical hypothyroidism (females = 42, males=25). The remaining 1% (n=6) belonged to subclinical hyperthyroidism (females=, males=3). The females were found to be highest with thyroid disorder with 75.2% (n=507) and 24.6% were males (n=166).

Table 2. Overall distribution of thyroid disorder

| Thyroid disorder (total no:674) | Number of patients | | Total | Percentage (%) |
|------------------------------------|--------------------|------------|-------|----------------|
| | Female | Male | | |
| Euthyroid | 430 | 129 | 560 | 83.0% |
| Overt hypothyroidism | 19 | 5 | 24 | 3.5% |
| Overt hyperthyroidism | 13 | 4 | 17 | 2.5% |
| Subclinical hypothyroidism | 42 | 25 | 67 | 10.0% |
| Subclinical hyperthyroidism | 3 | 3 | 6 | 1.0% |
| Total* | 507 | 166 | | |




Figure 1. : Distribution of different thyroid disorder

Table 3 shows comparison of thyroid hormone levels among different thyroid dysfunction. By applying ANOVA test, there was a significant differences (p>0.001) among various groups of thyroid disorder. T₃ levels were found to be highest in hyperthyroidism (2.8±3.9) and lowest in hypothyroidism (0.8 ±0.512). Highest level of T₄ were seen in hyperthyroidism (9.4±5.2) with lowest in hypothyroidism (5.02±2.6). High level of TSH was observed in hypothyroidism (12.8±13.1) and lowest in hyperthyroidism (3.2±3.5).

Table 3. Comparison of thyroid hormone levels among various thyroid dysfunction.

| Thyroid hormones | Euthyroid (Mean±SD) | Hypo- thyroidism (Mean± SD) | Hyper- thyroidism (Mean± SD) | Subclinical Hypothyroidism (Mean± SD) | Subclinical Hyperthyroidism (Mean± SD) | p value |
|------------------|------------------------|--------------------------------------|---------------------------------------|---|--|---------|
| T3(ng/ml) | 1.1±0.554 | 0.8±0.512 | 2.8±3.9 | 1.05±0.4 | 1.05±0.17 | 0.001* |
| T4(ug/ml) | 6.8±2.4 | 5.02±2.6 | 9.4±5.2 | 5.7±2.0 | 6.7±2.0 | 0.001* |
| TSH(uIU/ml) | 3.7±3.6 | 12.8±13.1 | 3.2±3.5 | 9.4±8.0 | 9.4±8.0 | 0.001* |

p value <0.05 is considered significant*.

The distribution of thyroid disorder according to age is represented in table 4. According to the age group, maximum number of patients (n=195) were in the age group of 21-30 yrs with 29%, followed by 25.7% of patients within the age of group 31-40yrs (n=173). 14.7% of patients belonged to the age group of 41-50 yrs (n=99). The age group between 51- 60 yrs were 11.9% (n=80). 7.3% were found between the 61-70 yrs of age (n=49) and the remain-

ing 3% were above 71 years of age.

Table IV. Frequency distribution of thyroid disorder according to age

| Age | Female (Frequency) | Male | Total Number (N=674) | Total Percentage (%) |
|-----------|-----------------------|------|-------------------------|-------------------------|
| 11-20yrs | 41 | 16 | 57 | 8.4% |
| 21-30yrs | 167 | 28 | 195 | 28.9% |
| 31-40yrs | 131 | 42 | 173 | 25.7% |
| 41-50yrs | 80 | 19 | 99 | 14.7% |
| 51-60 yrs | 49 | 31 | 80 | 11.9% |
| 61-70yrs | 29 | 20 | 49 | 7.3% |
| >71yrs | 10 | 10 | 20 | 3.0% |

DISCUSSION:

In the present study, out of total 674 patients suspected with thyroid disorder, highest number (80%) of subjects were found to be Euthyroid with normal levels of T_3 , T_4 & TSH. A hospital based study conducted by Rebecca et al [11] also reported that about 80% of patients were found to be Euthyroid. Among the thyroid disorders, subclinical hypothyroidism (10%) was found to be most prevalent followed by overt hypothyroidism (3.5%) and overt hyperthyroidism (2.5%) and the results were found to be similar in one of the studies (12,13). Hyperthyroidism (overt hyperthyroidism=2.5% & subclinical hyperthyroidism =1%) had the lowest incidence among the thyroid dysfunction and is comparable to other studies [13,14]. The data regarding the prevalence of thyroid disorders in India is comparatively scanty, though many studies are carried out to determine the effectiveness of iodination program [13].

The numbers of females were highest (n=507) indicating a high prevalence of thyroid dysfunction in women than men. This finding is supported by few studies [15, 16]. Majority of female suffering from thyroid dysfunction belonged to subclinical hypothyroidism (n=42) and overt hypothyroidism (n=19). This may be due to the geographical locations and pattern of iodine deficiency in these regions. A population based study carried out in Cochin, India showed a higher prevalence of subclinical hypothyroidism in women (11.4%) as compared to men [17].

The thyroid disorder cases were divided in seven age groups to determine the occurrence of various thyroid disorders in different age groups (Table 4). It was observed that maximum number of patients were seen between the age group of 21-30 years (29%) followed by 31-40 yrs of age group (25.7%). Lowest was observed above 70 years of age (3%). One of the study conducted in Jammu and Kashmir, showed a similar result of having a maximum number of thyroid patients between the age group of 21-30 yrs [12].

CONCLUSION :

Our study reported a high incidence of hypothyroidism in women as compared to men with maximum number in the age group of 21-30yrs. Though the state has implemented the iodized salt programme and has made remarkable and measurable progress within the last decades controlling the Iodine deficiency disorder but it still exists as a public health problem in Sikkim.

ACKNOWLEDGEMENT:

Authors wish to express our gratitude to the North Eastern Region of NER-BMPC, DBT, India DBT India for the grant under the Institutional level Biotechnology Hub Project.

REFERENCES:

- Nimmy NJ, Aneesh PM, Narmadha MP, Udipi RH, Binu KM. A Survey on Prevalence of Thyroid Disorders Induced by Demography and Food Habits in South Indian population. *Indian Journal of Pharmacy Practice*. 2012, 5:49-52
- Iodine Network. Global scorecard 2010. Available from: <http://www.iodinenetwork.net/documents/scorecard-2010>.
- Tiwari BK, Ray I, Malhotra RL. Policy Guidelines on National Iodine Deficiency Disorders Control Programme-Nutrition and IDD Cell. Directorate of Health Services, Ministry of Health and Family Welfare. New Delhi: Government of India; 2006, 1-22.
- Correcting Iodine Deficiency: A Lesson from Sikkim, *IDD Newsletter*; Volume 16; Number 4; November, 2000. 49-52.
- R Sankar, T Pulger, Bimal Rai, S Gomathi, TR Gyatso, CS Pandav: Epidemiology of Endemic Goitre in Sikkim. *JAPI*. 1997 : 45;12 : 936 — 940 pp.
- Sankar R, Pulger T, Rai B, Gomathi S, Gyatso TR, Pandav CS. Epidemiology of endemic cretinism in Sikkim, India. 1998 ; 65(2):303-9.
- Kochupillai N. Clinical endocrinology in India. *Current Science*. 2000; 79: 1061-1067.
- Vahab Fatourech. Subclinical Hypothyroidism: An Update for Primary Care Physicians. *Mayo Clin Proc*. 2009 ;84(1): 65-71.
- Heuck CC, Kallner A, Kanagasabapathy AS, Riesen W. WHO Document (WHO/DIL/00.4) on Diagnosis and monitoring of diseases of the thyroid , 2000.
- Marwah RK, Tandon N, Desai A, Kanwar R, Grewal K, Aggarwal R, et al. Reference range of thyroid hormones in normal Indian school-age children. *Clin Endocrinol* . 2008;68:369-74.
- Rebecca A, V Srinavasa Murugan, P ukazhvanthen , SK Sen. Thyroid disorders in women of Puducherry. *Journal of Clinical Biochemistry*. 2009; 24(1):52-59.
- Rama J, Shivashankara AR, Vidya SP, Sameena. A hospital based study of prevalence of thyroid dysfunction in Srinagar. *Int J Med Sci Public Health*. 2015;4(2):151-154.
- Arindam B, Norman S, Nanda H, Dhananhay SC. A hospital based prevalence study on Thyroid Disorders in Malwa region of Central India. *Int.J.Curr.Microbiol.App.Sci* .2015; 4(6):604-611.
- Abraham R, Murugan VS, Pukazhvanthen P, Sen SK. Thyroid disorders in women of Puducherry. *Indian J Clin Biochem* ;2009 ;24: 52-59.
- RV Mahato, B Jha, KP Singh, BK Yadav, SK Shah ,M Lamsal. Status of Thyroid disorders in Central Nepal : A Tertiary care Hospital based study. *Int J App Sci Biotechnol* . 2015; 3(1): 199-122.
- Friedman MN. Screening for thyroid disease. *Ann. Med*. 1999;130: 161-162. DOI: <http://dx.doi.org/10.7326/0003-4819-130-2-199901190-00016>
- Menon UV, Sundaram KR, Unnikrishnan AG, Jayakumar RV, Nair V, Kumar H. High prevalence of undetected thyroid disorders in an iodine sufficient adult south Indian population. *J Indian Med Assoc*. 2009;107:72-7.

Cite this article as:

Sonam Choden Bhutia, Karma Loday Bhutia, T.A. Singh. Thyroid Dysfunction in Central Referral Hospital, Sikkim. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 6(56), 2016, 17-19.