

UC Davis

UC Davis Previously Published Works

Title

Tik-borne rickettsial pathogens in ticks and small mammals in Korea

Permalink

<https://escholarship.org/uc/item/7p60x6rn>

Journal

Applied and Environmental Microbiology, 72(9)

ISSN

0099-2240

Authors

Kim, Chul-Min
Yi, Ying-Hua
Yu, Do-Hyeon
et al.

Publication Date

2006-09-01

Peer reviewed

Tick-Borne Rickettsial Pathogens in Ticks and Small Mammals in Korea

Chul-Min Kim,¹ Ying-Hua Yi,¹ Do-Hyeon Yu,¹ Mi-Jin Lee,¹ Mae-Rim Cho,¹ Atul R. Desai,¹
Smriti Shringi,¹ Terry A. Klein,² Heung-Chul Kim,³ Jin-Won Song,⁴ Luck-Ju Baek,⁴
Sung-Tae Chong,³ Monica L. O'Guinn,⁵ John S. Lee,⁵ In-Yong Lee,⁶
Jin-Ho Park,¹ Janet Foley,⁷ and Joon-Seok Chae^{1*}

Bio-Safety Research Institute and College of Veterinary Medicine, Chonbuk National University, Jeonju 561-756, Korea¹;
Force Health Protection, 18th Medical Command, Unit 15821, Box 754, APO AP 96205-5281²; 5th Medical Detachment,
168th Medical Battalion (AS), 18th Medical Command, Unit 15247, APO AP 96205-5247³; Department of Microbiology,
College of Medicine, Korea University, 126-1, 5 Ka, Anam-Dong, Sungbuk-Gu, Seoul 136-705, Korea⁴; Department of
Virology, 1425 Porter Street, U.S. Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick,
Maryland 21702-5011⁵; Department of Parasitology, College of Medicine, Yonsei University, Seoul 120-749,
Korea⁶; and Center for Vector-Borne Diseases, School of Veterinary Medicine, University of
California, Davis, California 95616⁷

Received 21 February 2006/Accepted 16 June 2006

In order to investigate the prevalence of tick-borne infectious agents among ticks, ticks comprising five species from two genera (*Hemaphysalis* spp. and *Ixodes* spp.) were screened using molecular techniques. Ticks (3,135) were collected from small wild-caught mammals or by dragging/flagging in the Republic of Korea (ROK) and were pooled into a total of 1,638 samples (1 to 27 ticks per pool). From the 1,638 tick samples, species-specific fragments of *Anaplasma phagocytophilum* (1 sample), *Anaplasma platys* (52 samples), *Ehrlichia chaffeensis* (29 samples), *Ehrlichia ewingii* (2 samples), *Ehrlichia canis* (18 samples), and *Rickettsia rickettsii* (28 samples) were amplified by PCR assay. Twenty-one pooled and individual tick samples had mixed infections of two (15 samples) or three (6 samples) pathogens. In addition, 424 spleen samples from small captured mammals (389 rodents, 33 insectivores, and 2 weasels) were screened for selected zoonotic pathogens. Species-specific DNA fragments of *A. phagocytophilum* (110 samples), *A. platys* (68 samples), *E. chaffeensis* (8 samples), *E. ewingii* (26 samples), *E. canis* (51 samples), and *Rickettsia* sp. (22 samples) were amplified by PCR assay. One hundred thirty small mammals had single infections, while 4, 14, and 21 striped field mice (*Apodemus agrarius*) had mixed infections of four, three, and two pathogens, respectively. Phylogenetic analysis based on nucleotide sequence comparison also revealed that Korean strains of *E. chaffeensis* clustered closely with those from China and the United States, while the *Rickettsia* (rOmpA) sequences clustered within a clade together with a Chinese strain. These results suggest that these agents should be considered in differential diagnosis while examining cases of acute febrile illnesses in humans as well as animals in the ROK.

Ticks are notorious vectors of various pathogenic protozoa, rickettsiae, bacteria, and viruses that cause serious and life-threatening illnesses in humans and animals worldwide (2, 11, 15, 37, 40, 47). Screening of ticks for such pathogens by using molecular epidemiological tools may disclose the prevalence of tick-borne pathogens in particular geographic environments. Some of these agents, such as *Rickettsia prowazekii* (typhus fever), *Yersinia pestis* (plague), *Francisella tularensis* (tularemia), *Coxiella burnetii* (Q fever), West Nile virus, Rift Valley fever virus, and hantavirus, are now recognized as important emerging vector-borne infections as well as agents of bioterrorism worldwide. Recently, ehrlichial and rickettsial infections have been reported to exist in a broad band across Europe (15), Asia (9, 27, 29), Africa (3), and the Americas (13, 17). Other tick-borne organisms, including some *Borrelia* and *Bartonella* spp., have also been shown to cause disease in animals and humans (23, 44, 45).

In the United States and Korea, rodents (e.g., the white-footed mouse [*Peromyscus leucopus*]) and white-tailed deer (*Odocoileus virginianus*) are reservoirs of *Ehrlichia* and *Anaplasma* spp. (9, 12, 35, 51, 53). In Europe, several rodent species are implicated as natural reservoirs for *Ehrlichia* and *Anaplasma* spp. (38). Additionally, *Ehrlichia* spp. have been isolated from wild mice in Japan (25).

The examination of ticks for tick-borne pathogens by molecular tools such as PCR is commonly used for collecting and assessing data regarding the prevalence of these agents. Although no cases of human anaplasmosis (formerly called human granulocytic ehrlichiosis [HGE]) or human monocytic ehrlichiosis have been reported, seroepidemiological findings suggest the presence of human monocytic ehrlichiosis and HGE agents in the Republic of Korea (ROK) (18, 40). In 2000, the first suspected case of *Ehrlichia chaffeensis* was reported for an active-duty American soldier stationed in the ROK (43). Subsequently, Heo et al. (18) identified antibodies against *E. chaffeensis* and *Anaplasma phagocytophilum* among serum samples from patients with febrile illnesses of otherwise unknown etiology in the ROK by an indirect fluorescent antibody test and Western blotting. Recently, *Rickettsia japonica* was identified in *Haemaphysalis*

* Corresponding author. Present address: Center for Vector-Borne Diseases, Department of Veterinary Medicine and Epidemiology, School of Veterinary Medicine, University of California—Davis, One Shields Ave., Davis, CA 95616. Phone: (530) 754-9740. Fax: (530) 754-4837. E-mail: jschae@ucdavis.edu.

TABLE 1. Oligonucleotide primers used for detection of tick-borne pathogens

Target genus or species	Oligonucleotide primer	Primer sequence (5'-3')	Annealing temp (°C)	Target gene (expected amplicon size [bp])	Reference
<i>Ehrlichia</i> and <i>Anaplasma</i>	ESP-F	AGTCCACGCTGTAAACGATGAG	55	16S rRNA (116)	27
	ESP-R	TTCCITTTGAGTTTTAGTCTTGCGAC	55		
	ESP-P (probe)	FAM-ACGCGTTAAGCACTCCGCTGG-TAMRA	55		
<i>A. phagocytophilum</i>	EE-1	TCCTGGCTCAGAACGAACTGGCGGC	50	16S rRNA (926)	6
	EE-2	AGTCACTGACCCAACCTTAAATGGCTG	50		
	EE-3	GTCGAACGGATTATTCTTTATAGCTTGC	50		
	EE-4	CCCTTCGGTTAAGAAGGATCTAATCTCC	50		
	ECC (outer primer)	AGAACGAACGCTGGCGGCAAGC	65		
<i>A. platys</i> , <i>E. chaffeensis</i> , <i>E. ewingii</i> , and <i>E. canis</i>	ECB (outer primer)	CGTATTACCGCGGCTGCTGGCA	65	16S rRNA (450)	37
	EPLAT5 (inner primer)	TTTGTCTGCTGCTATGAT	55		
<i>A. platys</i>	EPLAT3 (inner primer)	CTTCTGTGGGTACCGTC	55	16S rRNA (359)	36
	HE1 (inner primer)	CAATTGCTTATAACCTTTTGGTTATAAAT	55		
<i>E. chaffeensis</i>	HE3 (inner primer)	TATAGGTACCGTCATTATCTTCCCTAT	55	16S rRNA (390)	37
	EE52 (inner primer)	CGAACAATTCTTAAATAGTCTCTGAC	55		
<i>E. ewingii</i>	HE3 (inner primer)	TATAGGTACCGTCATTATCTTCCCTAT	55	16S rRNA (350)	37
	HE3 (inner primer)	TATAGGTACCGTCATTATCTTCCCTAT	55		
<i>E. canis</i>	ECAN5 (inner primer)	CAATTATTTATAGCCTCTGGCTATAGGA	55	16S rRNA (365)	37
	HE3 (inner primer)	TATAGGTACCGTCATTATCTTCCCTAT	55		
<i>E. muris</i>	CAN M61F	TTATCTGTTTATGTTATATAAGC	50	<i>gluA</i> (288)	21
	MUR SPR1	TAAATCTACTATGTTATGTCC	50		
<i>B. burgdorferi</i>	BBOSPF	AAAGAATACATTAAGTGCGATATT	54	<i>ospC</i> (597)	52
	BBOSPR	GGGCTTGTAAGCTCTTAACTG	54		
<i>R. rickettsii</i>	Rr190k. 71p	TGGCGAATATTTCTCCAAA	48	<i>ompA</i> (532)	42
	Rr190k. 602n	AGTGAGCATTTCGCTCCCCCT	48		
<i>R. japonica</i>	Rj10	ATTCTAAAAACCATATACTG	57	17K antigen (357)	16
	Rj5	CGCCATTCTACGTTACTACC	57		

longicornis ticks by PCR (29). Antibodies against *R. japonica* were also detected by an indirect fluorescent antibody test in human patients with acute febrile illness in the ROK (24). We previously reported molecular evidence of the presence of *E. chaffeensis* and *A. phagocytophilum* by using genus-specific TaqMan PCR and a species-specific PCR with ticks collected from animals and grass vegetation in the ROK (27) and gave preliminary reports of other tick-borne pathogens, including *Anaplasma platys*, *Ehrlichia ewingii*, *Ehrlichia canis*, and *Ehrlichia muris*. In 2005, Lee et al. (30) identified *E. chaffeensis* in *H. longicornis* ticks from the ROK by PCR. Also, Kim et al. (26) reported the detection of *Bartonella* species in ticks, mites, and small mammals in Korea. However, the prevalence of tick-borne pathogens, including *Rickettsia rickettsii* and *R. japonica*, has yet to be determined by molecular methods.

Recently, advanced molecular techniques such as TaqMan PCR became widely used as rapid and effective tools for the detection and identification of tick-borne pathogens in arthropods, including ticks (19, 27, 31, 41). This study was therefore undertaken to investigate the prevalence of tick-borne pathogens in small mammals and various tick species from the ROK. Conventional and TaqMan PCR assays were applied for rapid screening of ticks for detection of selected pathogens, followed by specific identification of the pathogens using species-specific and genus-specific primers. Using these approaches, we report herein the infection rates for *A. phagocytophilum*, *A. platys*, *E. chaffeensis*, *E. ewingii*, and *Rickettsia* spp. in ticks as well as wild-collected small mammals.

The purpose of the present study is to provide a disease risk assessment for various tick-borne pathogens, including *Ehrlichia*, *Anaplasma*, *Borrelia*, and *Rickettsia* spp., for humans and animals based on their potential exposure to ticks and tick-borne disease pathogens in field environments.

MATERIALS AND METHODS

Sample collection. Ticks were collected by dragging and flagging grass vegetation with a 1 m-by-1 m cotton cloth and removing attached ticks from various species of live wild rodents and insectivores that were trapped near the demilitarized zone (DMZ) and other U.S. military installations in the ROK. During 2001 through 2003, a total of 3,135 ticks were collected from wild rodents (297 ticks) and grass vegetation (2,838 ticks) at 19 sites near or at U.S. military installations and training sites in the ROK. Based on microscopic examination, ticks were identified to the species level and classified morphologically into various developmental stages (54). Subsequently, different species of ticks were pooled by species and stage of development (larvae and nymphs) into 1,638 sample pools (1 to 27 ticks per sample pool; pools included ticks from wild rodents [40 sample pools] and those from grass vegetation [1,638 sample pools]) and stored at -70°C in 1.5-ml microcentrifuge tubes until assayed. Ninety-four mites were collected from wild-caught rodents and insectivores and pooled into 21 samples (four or five mites/sample pool).

During 2001 through 2004, a total of 3,564 small mammals belonging to 11 species and 9 genera were collected throughout the ROK at U.S. military installations and training sites as part of the 18th Medical Command Hantavirus Surveillance Program. Of those mammals, 389 rodents, 33 insectivores, and 2 weasels were selected for assessment of tick-borne pathogens. Live-caught rodents, insectivores, and weasels were transported to the central laboratory (Korea University), where they were killed in accordance with approved animal use protocols. After blood samples were taken, the abdominal cavity was opened aseptically, and spleen and other tissue samples were collected. A subsample of spleen samples were sent to Chonbuk National University packed in dry ice, where they were stored individually at -70°C until assayed. Samples were chosen to balance analysis to include all collection areas and mammalian species.

DNA preparation. For extraction of PCR-amplifiable DNA, the ticks were pooled into a total of 1,638 sample pools (1 to 27 ticks per pool), including 313 pools of larval stages (1 to 27 ticks per pools), 1,176 pools of nymphal stages (1 to 3 ticks/pool), and 151 adult ticks (assayed individually). Purified DNAs were used for the detection of tick-borne pathogens by standardized TaqMan PCR techniques. Individuals (adults) or pools (larvae/nymphs) of ticks and mites (four or five mites/pool) were mechanically homogenized in 1-ml cryovials by using sterile scissors. DNA extraction was performed with a DNeasy tissue kit (QIAGEN, Germany) according to the instructions provided by the manufacturer.

PCR for *Ehrlichia* and *Anaplasma* spp. The extracted DNAs were subjected to an initial screening by TaqMan PCR as previously described, using a set of




FIG. 1. Phylogenetic tree showing the position of *Ehrlichia chaffeensis* 0214 from Korea. The tree was made using PAUP* 4.0b software after alignment of 16S rRNA gene fragments obtained (390 bp) from GenBank and sequenced during this study by the ClustalX program. The scale represents 0.01 substitution per base per indicated horizontal distance. The numbers present at nodes of the tree represent the numbers of bootstrap replicates of 100 that display the indicated sequence groupings.

primers with a probe that amplified the 116-bp fragment of the 16S rRNA genes of bacteria within the family *Anaplasmataceae*, including the genera *Ehrlichia* and *Anaplasma* (27) (Table 1). The fluorescence data were analyzed using PE 7700 sequence detection system software (version 1.7; ABI).

TaqMan PCR-positive DNA samples were used for specific identification of *A. phagocytophilum*, *A. platys*, *E. chaffeensis*, *E. ewingii*, *E. canis*, and *E. muris* by nested and single-tube conventional PCRs, as previously described, using species-specific primers (Table 1). The 16S rRNA gene fragment of *A. phagocytophilum* was amplified by a nested PCR assay according to the procedure of Barlough et al. (6). PCR for *E. muris* was performed using primers CAN M61f and MUR SPR1, targeting the *gltA* gene of *E. muris* (21). The primer sets for nested PCRs for *A. platys*, *E. chaffeensis*, *E. ewingii*, and *E. canis* DNAs were derived from the 16S rRNA gene sequences, with the same pair of outer primers and different sets of inner primers (36, 37). The oligonucleotide sequences for each pair of genus- and species-specific primers are shown in Table 1. PCR amplification of genomic DNAs of *R. rickettsii*, *R. japonica*, and *Borrelia burg-*

dorferi was performed with species-specific primers (Table 1) as previously described (16, 42, 52).

Nested and single PCRs were performed as previously described in a total volume of 25 μ l. Each PCR mixture consisted of 2 pmol of each primer, a 200 μ M concentration of each deoxynucleoside triphosphate, PCR buffer I (Super Bio, Korea), 0.75 U of SuperTaq DNA polymerase (Super Bio, Korea), and 50 to 100 ng of sample DNA for the first PCR or 1 μ l of the first PCR product for the second PCR. PCR products were electrophoresed in 1% agarose gels, stained with ethidium bromide, and photographed using a still-video documentation system (Gel Doc 2000; Bio-Rad). Unless specified otherwise, PCR products were purified with a GFX PCR DNA purification kit for cloning and sequencing (Amersham Biosciences, United Kingdom). Prevention of cross-contamination and false-negative and -positive results was managed by using plugged tips, performing PCR in a separate room from that used for DNA extraction, and including a negative (water) control in each run. A positive control was run for each master mix batch.




FIG. 2. Phylogenetic tree showing the position of *Rickettsia* sp. strain 71-8 from Korea. The tree was made using PAUP* 4.0b software after alignment of *ompA* gene fragments obtained (532 bp) from GenBank and sequenced during this study by the ClustalX program. The scale represents 0.01 substitution per base per indicated horizontal distance. The numbers present at nodes of the tree represent the numbers of bootstrap replicates of 100 that display the indicated sequence groupings.

Cloning, nucleotide sequencing, and phylogenetic analysis. PCR products were purified using the Wizard Plus DNA purification system (Promega), ligated into the pGEM-T Easy vector (Promega), and transformed into TOP10F competent cells. The recombinant clones were verified by colony PCR for the respective clones. Two clones of each isolate were arbitrarily chosen for sequencing of the forward and reverse strands. Plasmid DNA for sequencing was prepared using the SV Minipreps DNA purification system (Promega) according to the manufacturer's instructions.

Amplified and purified DNAs were prepared for direct sequencing using a GFX PCR DNA purification kit (Amersham Biosciences, United Kingdom) and were sequenced by dideoxy termination with an automatic sequencer (ABI Prism 3700 DNA analyzer). Sequence data were collected using ABI Prism data collection software (version 2.1) and analyzed by ABI Prism sequence analysis software (version 2.1.1) and Chromas software (version 1.51; Technelysium Pty., Ltd., Mt. Gravatt Plaza, Queensland, Australia). Sequence homology searches were made via the National Center for Biotechnology Information (National Institutes of Health) BLAST network service. The sequences were aligned initially using ClustalX 1.60 (49). 16S rRNA and rickettsia *ompA* gene sequences were used for phy-

logenetic analyses. Aligned sequences were examined with a similarity matrix. Relationships between individuals were assessed by the neighbor-joining method with nucleotide distances (*P* distance) for 100 replications in the bootstrap test. Phylogenetic analyses based on the obtained sequences were conducted using the maximum likelihood method (PAUP* 4.0b for Macintosh) (48).

Nucleotide sequence accession numbers. The 16S rRNA gene sequences for the following organisms (with GenBank accession numbers) were used: *Ehrlichia* sp. strain ERM58 from *Rhipicephalus muhame* in Mali (AF311967), *Ehrlichia* (*Cowdria*) *ruminantium* Omatjenne from South Africa (U03776), *E. ruminantium* from Senegal (X62432), *Ehrlichia* sp. strain HF565 from Japan (AB024928), *Ehrlichia* sp. from *Ixodes ovatus* in Japan (AB028319), *E. muris* AS145 from *Eothenomys kageus* in the United States (U15527), *Ehrlichia* sp. strain TS37 from Japan (AB074459), *E. canis* from Madrid, Spain (AY394465), *Ehrlichia* sp. strain Germishuys from South Africa (U54805), *Ehrlichia ovina* from Turkey and The Netherlands (AF318946), *Ehrlichia chaffeensis* from *H. longicornis* in Korea (AY350424), *E. chaffeensis* from China (AF147752), *E. chaffeensis* from Arkansas (M73222), *E. chaffeensis* 0214 from Korea (DQ402484), *Ehrlichia* sp. strain Tibet from *Boophilum micropus* in China (AF414399), *Ehrlichia* sp. strain EBm52 from

TABLE 2. Results of TaqMan and species-specific tick-borne pathogen PCR for selected tick species and stages of development collected at or near U.S. military installations and training sites in the ROK, 2001 to 2003

Species	Stage	n	No. (%) of PCR-positive samples										
			<i>Ehrlichia/Anaplasma</i> ^h	<i>A. phagocytophilum</i>	<i>A. platys</i>	<i>E. chaffeensis</i>	<i>E. ewingii</i>	<i>E. canis</i>	<i>E. muris</i>	<i>B. burgdorferi</i> ⁱ	<i>Rickettsia</i> sp. ⁱ	<i>R. japonica</i> ⁱ	
<i>H. longicornis</i>	Larva pools ^a	274	229	0	20	15	0	10	0	0	0	0	0
	Nymph ^b	1,099	510	0	23	10	1	6	0	0	27	0	0
	Male	53	28	0	4	1	0	0	0	0	1	0	0
	Female	77	33	0	4	2	1	1	0	0	0	0	0
	Subtotal	1,503	800	0	51	28	2	17	0	0	28	0	0
<i>H. flava</i>	Larva pools ^c	11	1	0	0	0	0	0	0	0	0	0	0
	Nymph	53	21	0	0	0	0	0	0	0	0	0	0
	Male	7	2	0	0	0	0	0	0	0	0	0	0
	Female	3	0	0	0	0	0	0	0	0	0	0	0
	Subtotal	74	24	0	0	0	0	0	0	0	0	0	0
<i>I. turdus</i>	Nymph	8	4	0	0	0	0	1	0	0	0	0	0
	Female	1	0	0	0	0	0	0	0	0	0	0	0
	Subtotal	9	4	0	0	0	0	1	0	0	0	0	0
<i>I. nipponensis</i>	Nymph	13	3	0	0	0	0	0	0	0	0	0	0
	Male	4	1	0	0	0	0	0	0	0	0	0	0
	Female	3	1	0	0	0	0	0	0	0	0	0	0
	Subtotal	20	5	0	0	0	0	0	0	0	0	0	0
<i>I. persulcatus</i>	Male	2	2	1	1	0	0	0	0	0	0	0	0
	Female	1	1	0	0	1	0	0	0	0	0	0	0
	Subtotal	3	3	1	1	1	0	0	0	0	0	0	0
<i>Ixodes</i> spp.	Larva pools ^d	26	19	0	0	0	0	0	0	0	0	0	0
	Nymph	3	2	0	0	0	0	0	0	0	0	0	0
	Subtotal	29	21	0	0	0	0	0	0	0	0	0	0
Total	Larva pools ^e	313	245	0	20	15	0	10	0	0	0	0	0
	Nymph ^f	1,176	536	0	23	10	1	7	0	0	27	0	0
	Male	66	66	1	5	1	0	0	0	0	1	0	0
	Female	85	85	0	4	3	0	1	0	0	0	0	0
	Total ^g	1,638	857 (52.3)	1 (0.1)	52 (3.2)	29 (1.8)	2 (0.1)	18 (1.1)	0	0	28 (1.7)	0	0

^a Two to seven ticks per pool (1,445 ticks).

^b One to three ticks per pool (1,127 ticks).

^c Two to five ticks per pool (52 ticks).

^d One to 27 ticks per pool (284 ticks).

^e One to 27 ticks per pool (1,780 ticks).

^f One to three ticks per pool (1,204 ticks).

^g One to 27 ticks per pool (3,135 ticks).

^h TaqMan PCR-positive samples were tested for *A. phagocytophilum*, *A. platys*, *E. chaffeensis*, *E. ewingii*, *E. canis*, and *E. muris*, and the numbers and percentages were calculated based on the numbers of TaqMan PCR-positive samples for the respective species.

ⁱ The numbers and percentages of positive samples for *B. burgdorferi*, *R. rickettsii*, and *R. japonica* were calculated based on the total numbers of ticks tested.

B. micropus in Thailand (AF497581), *E. ewingii* HH591-2 from the United States (AY093440), *Ehrlichia* sp. strain EHT224 from *Hyalomma truncatum* in Niger (AF311968), *Ehrlichia* sp. strain EHF669 from *Hemaphysalis* sp. in Japan (AY309969), *A. platys* from a dog in Spain (AY530806), *Anaplasma bovis* AB-KGHL from *H. longicornis* in Korea (AF470698), *Anaplasma centrale* vaccine strain from The Netherlands (AF318944), *Anaplasma ovis* from sheep (AF318945), *Anaplasma marginale* Lushi from cattle in China (AJ633048), *Ehrlichia* sp. from Sweden (AJ242785), *A. ovis* from Germany (AY262124), *A. phagocytophilum* USG3 from the United States (AY055469), "Candidate *Neorickettsia mikurensis*" (AB074460), *Ehrlichia*-like sp. Schotti variant (AF104680), *Wolbachia* endosymbiont (AM180551), *Neorickettsia risticii* (M21290), and *Rickettsia rickettsii* (DQ150688) (Fig. 1).

The *ompA* gene sequences for the following rickettsiae (with GenBank accession numbers) were used: *Rickettsia* sp. strain 71-8 from Korea (DQ402485), *Rickettsia* sp. strain FUJ98 from China (AF169629), *Rickettsia heilongjiangii* HLJ-054 (AF179362), *Rickettsia hulunii* (AF179364), *R. japonica* (D28766), *Rickettsia mongolotimonae* (U43796), *Rickettsia sibirica* (U43807), *Rickettsia africana* (U43790), *Rickettsia parkeri* (U43802), *Rickettsia slovacica* (U43808), *Rickettsia honei* (AF018075), Israeli tick typhus rickettsia (AY197564), *Rickettsia conorii* (U43794), *R. rickettsii* (U55822), *Rickettsia rhipicephali* (U43803), *Rickettsia massiliae* (U43799), *Rickettsia aeschlimannii* (U43800), *Rickettsia amblyommii* (AY062007), *Rickettsia australis* (AF149108), and *Rickettsia montanensis* (U43801) (Fig. 2).

RESULTS

A total of 3,135 ticks, including five species from two genera (2,701 *H. longicornis*, 115 *H. flava*, 9 *Ixodes turdus*, 3 *Ixodes persulcatus*, 20 *Ixodes nipponensis*, and 287 *Ixodes* sp. ticks), were collected from wild rodents and insectivores (297 ticks) and from grass/vegetation (2,838 ticks) from 2001 through 2003 (Table 2). *H. longicornis* ticks were the most commonly collected species, and irrespective of species, most of the ticks were collected in the nymphal stage of development (Table 2). All ticks collected from wild rodents and insectivores during this period were *Ixodes* spp. All of the mesostigmatid mite samples included in this study were collected from rodents and insectivores and assayed for the same pathogens examined in ticks.

A total of 424 small mammals (six rodent species belonging to five genera [373 *Apodemus agrarius*, 3 *Apodemus peninsulae*, 1 *Cricetulus triton*, 9 *Eothenomys regulus*, 1 *Mus musculus*, and 2 *Rattus rattus* animals], one insectivore species [33 *Crosidura*

TABLE 3. Tick-borne pathogens identified by DNA analysis in small mammals collected at U.S. military installations and training sites in the ROK, 2001 to 2004

Species	n	No. (%) of PCR-positive samples									
		<i>Ehrlichia/Anaplasma</i>	<i>A. phagocytophilum</i>	<i>A. platys</i>	<i>E. chaffeensis</i>	<i>E. ewingii</i>	<i>E. canis</i>	<i>E. muris</i>	<i>B. burgdorferi</i>	<i>Rickettsia</i> sp.	<i>R. japonica</i>
<i>Apodemus agrarius</i>	373	270	88	68	8	26	50	0	0	22	0
<i>Crosidura lasiura</i>	33	26	21	0	0	0	1	0	0	0	0
<i>Eothenomys regulus</i>	9	0	0	0	0	0	0	0	0	0	0
<i>Apodemus peninsulae</i>	3	0	0	0	0	0	0	0	0	0	0
<i>Rattus rattus</i>	2	0	0	0	0	0	0	0	0	0	0
<i>Mustela sibirica</i>	2	1	0	0	0	0	0	0	0	0	0
<i>Cricetulus triton nestor</i>	1	1	0	0	0	0	0	0	0	0	0
<i>Mus musculus</i>	1	1	0	0	0	0	0	0	0	0	0
Total	424	299 (70.5)	110 (25.9)	68 (16)	8 (1.9)	26 (6.1)	51 (12.0)	0	0	22 (5.2)	0

lasiura animals], and one mustelid species [2 *Mustela sibirica* animals]) were collected from 2001 through 2004 at U.S. military installations and training sites near the DMZ and at other U.S. military installations south of Seoul, ROK. *A. agrarius* was the most commonly collected mammal and accounted for 88% (373/424) of the total samples from small mammals (Table 3).

TaqMan PCR signals for *Ehrlichia* and *Anaplasma* spp. were detected from all tick species sampled and from five of eight species of small wild mammals (*A. agrarius*, *C. lasiura*, *M. sibirica*, *C. triton nestor*, and *M. musculus*). The prevalence of *Ehrlichia* and *Anaplasma* parasites in ticks and all small mammals was 52.3% (857/1,638 tick pools), with minimum infection rates of 27.3% (857/3,135 total ticks) and 70.5% (299/424 mammals), respectively (Tables 2 and 3).

Species-specific PCR assays were conducted with DNA samples from 857 tick pools and 277 small mammals that were positive by TaqMan PCR. Except for *E. muris*, five of six tick-borne pathogens evaluated in this study were detected in ticks (*A. phagocytophilum* [1], *A. platys* [52], *E. chaffeensis* [29], *E. ewingii* [2], and *E. canis* [18]) and small mammals (*A. phagocytophilum* [110], *A. platys* [68], *E. chaffeensis* [8], *E. ewingii* [26], and *E. canis* [51]) by nested PCR (Tables 2 and 3). The most frequently isolated species in ticks was *A. platys* (52 [3.2%]), followed by *E. chaffeensis* (29 [1.8%]), *E. canis* (18

[1.1%]), *E. ewingii* (2 [0.1%]), and *A. phagocytophilum* (1 [$<0.1\%$]). In the case of small mammals, the most frequently detected species ($n = 424$) was *A. phagocytophilum* (110 [25.9%]), followed by *A. platys* (68 [16%]), *E. canis* (51 [12.0%]), *E. ewingii* (26 [6.1%]), and *E. chaffeensis* (8 [1.9%]). In an additional study, all of the DNA samples from 1,638 tick pools and 424 small wild mammals were examined for three tick-borne pathogens, *B. burgdorferi*, *R. rickettsii*, and *R. japonica*, directly by single PCRs with specific primers. Twenty-eight *H. longicornis* (1.7%) ticks and 22 *A. agrarius* (5.2%) animals were positive for *R. rickettsii*. However, DNA bands specific for *B. burgdorferi* and *R. japonica* were not observed for ticks or small mammals examined during this study. None of the 297 *Ixodes* spp. (40 pools) collected from small mammals were found to be infected with any of the nine tick-borne pathogens examined in this study. Among 21 mite samples collected from wild rodents, 4 (19%) were PCR positive for *Ehrlichia* and *Anaplasma* spp. by TaqMan PCR, while a specific DNA band was not observed for six *Ehrlichia* and *Anaplasma* spp., *B. burgdorferi*, *R. rickettsii*, or *R. japonica*.

A total of 21 pools/individual ticks represented multiple infections (Table 4). Pools of larvae and nymphs with multiple infections may represent individual infections from ticks within each of the samples. However, two and one adult ticks, assayed

TABLE 4. Mixed infections of tick-borne pathogens among individual ticks and pools of ticks collected at or near U.S. military installations and training sites near the DMZ, ROK, 2001 to 2003^a

Species	Stage	No. of infected samples					Total
		<i>A. phagocytophilum/ A. platys</i>	<i>E. chaffeensis/ E. canis</i>	<i>E. chaffeensis/ A. platys</i>	<i>E. chaffeensis/ E. canis/A. platys</i>	<i>E. chaffeensis/ E. canis/E. ewingii</i>	
<i>H. longicornis</i>	Larva pools	0	10	2	0	0	12
	Nymph	0	0	1	4	1	6
	Male	0	0	1	0	0	1
	Female	0	0	0	0	1	1
	Subtotal	0	10	4	4	2	20
<i>I. persulcatus</i>	Female	1	0	0	0	0	1
	Subtotal	1	0	0	0	0	1
Total	Larva pools	0	10	2	0	0	12
	Nymph	0	0	1	4	1	6
	Male	0	0	1	0	0	1
	Female	1	0	0	0	1	2
	Total	1	10	4	4	2	21

^a Adult male and female ticks were assayed individually. Pools may represent single infections among ticks from the same sample.

TABLE 5. Mixed infections of tick-borne pathogens in small mammals (*Apodemus agrarius*) collected at U.S. military installations and training sites, ROK, 2001 to 2004

Species	No. of mixed infections	Total no. of infections for group ^a
<i>A. phagocytophilum</i> / <i>A. platys</i> / <i>E. canis</i> / <i>E. ewingii</i>	3	
<i>A. platys</i> / <i>E. canis</i> / <i>E. ewingii</i> / <i>E. chaffeensis</i>	1	4
<i>A. platys</i> / <i>E. ewingii</i> / <i>E. canis</i>	7	
<i>A. platys</i> / <i>E. canis</i> / <i>E. chaffeensis</i>	2	
<i>A. phagocytophilum</i> / <i>E. ewingii</i> / <i>E. canis</i>	1	
<i>A. phagocytophilum</i> / <i>E. chaffeensis</i> / <i>E. ewingii</i>	4	14
<i>A. phagocytophilum</i> / <i>A. platys</i>	1	
<i>A. platys</i> / <i>E. ewingii</i>	1	
<i>A. platys</i> / <i>E. canis</i>	13	
<i>E. ewingii</i> / <i>E. canis</i>	2	
<i>A. phagocytophilum</i> / <i>E. ewingii</i>	1	
<i>A. phagocytophilum</i> / <i>R. rickettsii</i>	1	
<i>E. chaffeensis</i> / <i>R. rickettsii</i>	1	
<i>E. ewingii</i> / <i>R. rickettsii</i>	1	21
Total	39	39

^a Mixed infections are divided into groups of infections with four, three, or two organisms.

individually, were positive for two and three *Ehrlichia*/*Anaplasma* pathogens, respectively. One hundred thirty small mammals had single infections, while 4, 14, and 21 *A. agrarius* animals had mixed infections of four, three, or two pathogens, respectively (Table 5).

Among 19 tick collection sites, *Ehrlichia* and *Anaplasma* DNAs were detected from all sites, while selected pathogens tested by species-specific PCR in this study were recovered from only 13 sites. Of those, *A. platys* was detected at the most

sites (11), followed by *E. chaffeensis* (5 sites), *R. rickettsii* (3 sites), *E. canis* (2 sites), *A. phagocytophilum* (1 site), and *E. ewingii* (1 site).

During the 3-year period, monthly infection rates for *Ehrlichia* and *Anaplasma* spp. in ticks were higher in October (148/172 [86%]) than during earlier months (25% [2/8] in March, 42.8% [511/1,193] in June, 63.3% [19/30] in August, and 73.6% [170/231] in September), except for April (1/1) and July (3/3). Infection rates for small mammals were higher in February (46/49 [93.9%]) than in the other 5 months of sampling (60.3% [38/63] in March, 74.8% [92/123] in April, 83.9% [26/31] in May, 68.8% [44/64] in June, and 56.4% [53/94] in October). The infection rates among ticks positive for *Ehrlichia* or *Anaplasma* spp. were higher in September ($n = 231$) (28 *A. platys* infections [12.1%], 23 *E. chaffeensis* infections [10.0%], 2 *E. ewingii* infections [0.9%], and 16 *E. canis* infections [6.9%]), while positive rates for small mammals were higher in April ($n = 123$) (3 *A. phagocytophilum* infections [2.4%], 42 *A. platys* infections [34.1%], 2 *E. chaffeensis* infections [1.6%], 9 *E. ewingii* infections [7.3%], and 22 *E. canis* infections [1.9%]). *R. rickettsii* was observed in ticks collected during June (28) and in small mammals collected during March (19) and April (3).

E. chaffeensis sequences were compared with those of other isolates of *E. chaffeensis* available in GenBank. The *E. chaffeensis* 0214 sequence (present study) was homologous to those of other Korean (99.7%) (AY350424), U.S. (99.7%) (AF416764), and Chinese (99.7%) (AF147752) isolates. The homology level between the two nucleotide sequences determined in this study varied from 96.2% to 100%. Comparative analysis of nucleotide sequences of the Korean strains determined in this study and the 16S rRNA sequences of 14 known *Ehrlichia* spp. available in the GenBank database is shown in Table 6. Phylogenetic analysis showed that *E. chaffeensis* 0214

TABLE 6. Homology comparison of Korean *Ehrlichia chaffeensis* 16S rRNA gene fragment (390-bp) sequences with those of other strains

Sequence	% Identity or no. of nucleotide differences ^a														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1		99.7	99.7	99.7	98.2	98.2	97.9	97.4	97.4	97.2	97.2	97.2	96.7	96.2	96.2
2	1		100	100	98.5	98.5	98.2	97.7	97.7	97.4	97.4	97.4	96.9	96.4	96.4
3	1	0		100	98.5	98.5	98.2	97.7	97.7	97.4	97.4	97.4	96.9	96.4	96.4
4	1	0	0		98.5	98.5	98.2	97.7	97.7	97.4	97.4	97.4	96.9	96.4	96.4
5	7	6	6	6		100	97.7	99.2	99.2	99.0	99.0	99.0	96.2	96.4	96.4
6	7	6	6	6	0		97.7	99.2	99.2	99.0	98.5	96.7	96.2	96.4	96.4
7	8	7	7	7	9	9		97.4	97.4	96.9	98.2	97.2	96.7	96.7	97.2
8	9	8	8	8	3	3	10		100	98.2	97.7	96.9	96.4	96.1	96.6
9	9	8	8	8	3	3	10	0		98.2	97.7	96.9	96.4	96.1	96.6
10	11	8	8	8	4	3	12	7	7		97.4	96.2	95.9	95.6	95.6
11	11	9	9	9	4	5	7	9	9	10		95.9	95.4	95.9	96.4
12	11	9	9	9	4	12	11	12	12	16	16		99.5	95.9	95.1
13	13	9	9	9	15	15	13	14	14	18	18	2		95.4	94.6
14	15	12	12	12	13	14	12	15	15	17	16	15	18		95.4
15	15	14	14	14	14	14	11	13	13	17	14	19	22	18	

^a Percentages of identity between 16S rRNA gene fragment sequences are shown in the upper matrix. The lower matrix shows numbers of nucleotide differences. The following sequences were compared: 1, *E. chaffeensis* 0214 from *Haemaphysalis longicornis* in Korea; 2, AY350424 (*E. chaffeensis* from Korea); 3, AF416764 (*E. chaffeensis* from Arkansas); 4, AF147752 (*E. chaffeensis* from China); 5, AF414399 (*Ehrlichia* sp. from *Boophilus microplus* from Tibet (in China)); 6, AF497581 (*Ehrlichia* sp. strain EBm52 from *Boophilus microplus* in Thailand); 7, AY309969 (*Ehrlichia* sp. strain EHF669 from *Haemaphysalis* sp. in Japan); 8, AF311967 (*Ehrlichia* sp. strain ERm58 from Africa); 9, AF311967 (*E. chaffeensis* from France); 10, AF311968 (*Ehrlichia* sp. strain EHT224 from *Hyalomma truncatum* in Niger); 11, AB074459 (*Ehrlichia* sp. strain TS37 from Japan); 12, AB024928 (*Ehrlichia* sp. strain HF565); 13, AB028319 (*Ehrlichia* sp. strain Anan); 14, AF318946 (*E. ovina* [isolate from turkey ruminant in The Netherlands]); 15, U03776 (*C. ruminantium* from Omatjenne in South Africa).

TABLE 7. Homology comparison of *Rickettsia* sp. *ompA* gene fragment (532-bp) sequences

Sequence	% Identity or no. of nucleotide differences ^a														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1		99.4	94.7	94.5	94.3	93.9	93.8	93.8	93.8	93.8	93.8	93.8	93.2	92.6	92.4
2	3		94.5	94.3	93.8	93.4	93.6	93.2	93.6	93.2	93.2	93.2	92.6	92.0	91.8
3	26	28		99.8	94.5	94.7	97.3	93.9	93.9	94.1	94.3	93.8	94.1	93.0	92.2
4	27	29	1		94.3	94.5	97.1	93.8	93.8	93.9	93.9	93.6	93.9	92.8	92.0
5	28	32	28	28		98.6	93.6	97.9	97.5	98.0	98.2	98.0	96.9	93.4	93.2
6	28	34	27	27	6		93.8	97.7	98.2	98.6	98.8	98.2	96.7	93.4	93.2
7	28	33	14	15	33	32		93.4	93.0	93.2	93.4	92.8	93.2	92.2	91.4
8	32	35	31	32	11	12	34		96.5	97.5	97.7	97.1	96.3	93.2	93.0
9	32	33	31	31	13	9	35	18		97.5	98.0	96.9	95.5	92.6	92.4
10	32	35	29	31	10	7	35	11	12		99.0	97.3	96.5	93.2	93.0
11	32	35	28	30	9	6	33	12	9	5		97.5	96.7	93.4	93.2
12	32	35	32	33	10	9	37	15	15	13	13		96.5	92.8	92.6
13	32	38	29	31	16	17	35	19	23	18	16	18		92.4	92.2
14	40	41	35	37	34	34	38	35	37	35	34	37	39		98.8
15	40	42	39	41	35	35	43	36	38	36	35	38	41	6	

^a Percentages of identity between sequences of *ompA* gene fragments are shown in the upper matrix. The lower matrix shows the numbers of nucleotide differences. The following sequences were used: 1, *Rickettsia* sp. strain 71-8 from Korea; 2, AF169629 (*Rickettsia* sp. from China); 3, AF179364 (*Rickettsia hulinii* from France); 4, AF179362 (*Rickettsia heilongjiangii* from France); 5, U43808 (*Rickettsia slovacica* from France); 6, U43790 (*Rickettsia africana* from France); 7, D28766 (*Rickettsia japonica* from Japan); 8, AY197564 (Israeli tick typhus rickettsia from Italy); 9, U43802 (*Rickettsia parkeri* from France); 10, U43807 (*Rickettsia sibirica* from France); 11, U43796 (*Rickettsia mongolotimonae* from France); 12, AF018075 (*Rickettsia honei* from the United States); 13, U55822 (*Rickettsia rickettsii* from the United States); 14, U43803 (*Rickettsia rhipicephali* from France); 15, U43799 (*Rickettsia massiliae* from France); 16, AY062007 (*Rickettsia amblyommii* from the United States); 17, U43800 (*Rickettsia aeschlimannii* from France).

clustered together with Korean, Arkansan, and Chinese strains of *E. chaffeensis* (Fig. 1).

Sequence comparison and alignment of *ompA* gene nucleotide sequences from *Rickettsia* sp. strain 71-8 (present study) revealed >99% homology to previously reported sequences for Korean and Chinese strains of *Rickettsia* spp. (Table 7). The sequence similarity of *Rickettsia* sp. strain 71-8 with an *R. rickettsii* isolate from Japan was <94%. However, phylogenetic analysis showed that *Rickettsia* sp. strain 71-8 formed a single cluster with a *Rickettsia* sp. Chinese isolate, while an *R. rickettsii* U.S. isolate was placed in a different cluster. The close clustering of Chinese and Korean strains of *Rickettsia* spp. may indicate a close epidemiological link between these strains (Fig. 2).

DISCUSSION

An analysis of the prevalence of selected tick-borne pathogens in ticks and mites collected from wild-caught rodents/insectivores and by dragging and flagging vegetation at or near U.S. military installations in the ROK demonstrated a high rate of infection with *Ehrlichia* and *Anaplasma* sp. pathogens. Most *Ehrlichia* and *Anaplasma* spp. have been recovered from *Ixodes* sp. ticks in the Americas and Europe (1, 50), while in Asia, *Ehrlichia* spp. were identified from both *H.* and *Ixodes* sp. ticks (22, 27). *H. longicornis* is the most commonly collected species in tick drag-and-flag collections in the ROK, accounting for >98% of all ticks collected, especially in and around pastures for grazing cattle or where deer congregate. *H. flava* is more commonly collected in pine forests, accounting for >95% of all ticks collected in this environment. *Ixodes* spp. are infrequently collected in grassy vegetation and pine forests, generally making up <2% of the collected ticks (data not shown). However, in deciduous forests with an abundance of leaf litter, collection of *I. nipponensis* may exceed 5% of the collected ticks during tick dragging and flagging (H. C. Kim,

personal communication). Another sampling technique that used CO₂-baited traps proved to be unsuccessful for capturing ticks. In that investigation, all of the ticks (297) taken from wild rodents and insectivores belonged to the genus *Ixodes*. Recently, all ticks (2,760) taken from rodents and insectivores captured at the same sites during 2004 and 2005 were identified as *I. nipponensis*, except for one *H. flava* tick (Kim, unpublished data). Thus, it is likely that the ixodid ticks tested herein for zoonotic pathogens were *I. nipponensis*. Human cases of tick bite are more commonly reported for *Ixodes* spp. (20, 28). Thus, while small numbers of *Ixodes* spp. were collected through dragging and flagging vegetation, this method may represent a possible bias for attracting members of the genus *Hemaphysalis*, or the bites from ixodid ticks may be of greater severity and thus reported more frequently.

Simple and easy methods for the identification of tick-borne pathogens in ticks and wild animals are necessary for a rapid analysis of disease-causing agents in developing an accurate risk assessment for soldiers training in the field. TaqMan PCR has proven to be a relatively easy and rapid method for the detection and identification of microorganisms in field samples, such as ticks (32), and in a previous report, its sensitivity was identical to that of nested PCR (6). These techniques can also be applied for rapid diagnosis with other samples, e.g., blood from patients suffering from a febrile disease with an unknown etiology.

16S rRNA gene sequence analysis is a widely used method for characterization of pathogenic bacteria, especially for the genera *Ehrlichia* and *Anaplasma*. Primers ECC and ECB (12) were used to amplify a segment of the 16S rRNA gene, which includes the specific primer regions of four *Ehrlichia/Anaplasma*-related pathogens (*A. platys*, *E. chaffeensis*, *E. ewingii*, and *E. canis*) examined in this study (Table 1). For *A. phagocytophilum* and *E. muris*, we designed separate primers from ECC and ECB because there are no species-specific regions be-

tween the ECC and ECB sequences. While these primers effectively identified pathogens to the species level, approximately two-thirds of the *Anaplasma/Ehrlichia*-positive samples were unidentified. Some of these positive samples may represent other infectious agents previously identified in Korea, i.e., *E. bovis*, *A. marginale*, and *A. centrale*, or may represent unknown or previously unreported pathogens.

Ticks of the genera *Hemaphysalis* and *Ixodes* collected from grass were found to be infected with one or more of the six tick-borne pathogens tested (*A. phagocytophilum*, *A. platys*, *E. chaffeensis*, *E. ewingii*, *E. canis*, and *R. rickettsii*). Three pathogens identified in this study have been reported previously in the ROK (9, 27, 29, 30). To the best of our knowledge, this is the first report of *A. platys*, *E. ewingii*, and *E. canis* in the ROK (9). Screening of ticks as well as small mammalian spleens for the presence of *Rickettsia* sp. infections corroborated well with earlier findings in the ROK (Joon-Seok Chae, unpublished data). In this study, larval ticks were positive for various *Ehrlichia/Anaplasma* sp. infections. While transovarial transmission in *Amblyomma americanum* has not been reported, our results for transovarial transmission of *E. chaffeensis* parasites in female ticks is undetermined, as 15 pools of larval *H. longicornis* ticks were positive for *E. chaffeensis* (33). However, some larval ticks may have been collected either partially or fully engorged, and it is possible that the DNA detected might have been in the host's blood that was ingested by the ticks. While *H. longicornis* larvae were positive, there were no larval *Ixodes* spp. taken from rodents positive for *E. chaffeensis*, even though they had been attached and partially blood fed and 3% of the striped field mice were positive for this pathogen. More studies have to be conducted to determine the status of transovarial transmission of *E. chaffeensis* and other zoonotic tick-borne pathogens among various species of ticks found in Korea.

There were observed differences in the infection rates of ticks versus small-mammal tissues. For example, only 1 of 1,638 pools of ticks was positive for *A. phagocytophilum*, while 24% (88/373) of the *A. agrarius* and 64% (21/33) of *Crosodura lasiura* (shrew) animals were positive. However, this may be a result of the fact that most of the ticks sampled from tick dragging and flagging were from the genus *Hemaphysalis*, while nearly all ticks (>99.9%) taken from small mammals were *I. nipponensis*. Soft ticks, which were not sampled during this study, also may be vectors of various pathogens. More investigations are needed to resolve these differences.

While there is no evidence of *Rickettsia* sp. infections in animals and only a single case reported among Koreans, the results based on the present study, combined with previous serologic and molecular evidence (24, 29), suggest that cases of febrile illness by spotted fever group *Rickettsia* in the ROK could be missed during the diagnosis of such illnesses. This is supported by retrospective studies by Song et al. (46) and Baek et al. (5), who demonstrated that a large proportion of patients with suspected hemorrhagic fever with renal syndrome were serologically positive for spotted fever group agents while being negative for hemorrhagic fever with renal syndrome. Specific DNAs of *B. burgdorferi* and *R. japonica* were not amplified in this study, although there are previous reports of these infections in Korean patients and ticks (24, 39). Therefore, it appears that cases of febrile illness of unknown etiology in

humans as well as animals should always be considered possible tick-borne infections in differential diagnosis.

Since human granulocytic and monocytic ehrlichioses were first reported in 1994 and 1987, respectively, they have been found in many countries by molecular and serologic surveys (8, 10, 34). In particular, tick infestations of wild animals have often been investigated because these animals have a high risk of infections. Epidemiological, molecular, and serological studies provided evidence that wild rodents are the reservoir of *Ehrlichia* and *Anaplasma* spp. in the United States, Europe, and Asia. Natural reservoirs for *A. phagocytophilum* were demonstrated to be white-footed mice (*Peromyscus leucopus*), eastern chipmunks (*Tamias striatus*), southern red-backed voles (*Clethrionomys gapperi*), and insectivorous shrews (*Blarina brevicauda* and *Sorex cinereus*) in Minnesota (51). In Japan, while antibodies against *E. muris* were detected in *Apodemus speciosus* and *Apodemus argenteus* mice (25), they were observed neither in rodent tissues nor from ticks collected near the DMZ in Korea. Our results demonstrate that small mammals captured in the ROK were infected with *Ehrlichia/Anaplasma* and/or *Rickettsia* parasites. Infections with *Ehrlichia* and *Anaplasma* spp. have generally been considered host specific. However, our studies suggest that several *Ehrlichia* and *Anaplasma* spp. can be transmitted to a variety of hosts in nature. Therefore, additional efforts to define the spectrum of host susceptibility in domestic and wild animals are appropriate.

The spotted fever group rickettsiae in Korea were identified using a primer for *R. rickettsii*. However, phylogenetic analysis revealed that *Rickettsia* sp. strain 71-8 (present study) formed a single cluster with a Chinese *Rickettsia* sp., while a U.S. *R. rickettsii* strain was placed in a different cluster and most likely represents a distinct species. Close clustering with the *Rickettsia* sp. from China indicated that although Korean strains differed significantly from those of *R. rickettsii*, there is a potential epidemiological link between the Chinese and Korean strains. Similarly, *E. chaffeensis* 0214 clustered together with *E. chaffeensis* strains from Korea, Arkansas, and China, which also indicates the possibility of an epidemiological link between Korea and China. Further studies that provide for molecular characterization and identify the epidemiology, human infection rates, and potential human health hazards of these pathogens are needed.

Previous studies have implicated mites as potential vectors of *Ehrlichia* and *Anaplasma* pathogens in Spain (14). In Korea, mites have not been well studied to determine their potential impact on the zoonotic maintenance and transmission of *Ehrlichia*, *Anaplasma*, or *Rickettsia* pathogens. Of 21 mite samples assayed during this study, 4 (19%) were positive for *Ehrlichia* and *Anaplasma* spp. However, species-specific DNAs examined in this study were not amplified and may represent *Anaplasma* or *Ehrlichia* pathogens previously described from Korea but not evaluated here, unidentified emerging pathogens, or previously unreported pathogens. Further investigation is needed to determine the role of mites in maintenance and transmission of zoonotic vector-borne pathogens. However, the association between positive TaqMan PCR and negative PCR for nine tick-borne pathogens has not been determined.

Ixodid ticks play an important role as a reservoir for latent

infections with various tick-borne pathogens (4, 7). In 2003, three members of the family *Anaplasmataceae*, including *E. chaffeensis*, *A. phagocytophilum*, and *A. bovis*, were initially described in the ROK (27). Until now, there have been very few reports of epidemiological studies for tick-borne disease surveillance in the ROK, in contrast with numerous reports throughout the world. These studies have enabled us to provide further information on the epidemiology of tick-associated pathogens in the ROK, where little information on the subject exists. Further studies are required for a detailed understanding of these newly emerging tick-borne diseases in the ROK.

ACKNOWLEDGMENTS

Funding for portions of this work was provided by the U.S. Department of Defense Global Emerging Infections Surveillance and Response System, Silver Spring, MD, the Armed Forces Medical Intelligence Center, Ft. Detrick, MD, the international collaborative research fund of Chonbuk National University (2004), and the Brain Korea 21 Project in E007.

REFERENCES

- Adelson, M. E., R. V. S. Rao, R. C. Tilton, K. Cabets, E. Eskow, L. Fein, J. L. Occi, and E. Mordechai. 2004. Prevalence of *Borrelia burgdorferi*, *Bartonella* species, *Babesia microti*, and *Anaplasma phagocytophilum* in *Ixodes scapularis* ticks collected in northern New Jersey. *J. Clin. Microbiol.* **42**:2799–2801.
- Alekseev, A. N., H. V. Dubinina, I. van de Pol, and L. M. Schouls. 2001. Identification of *Ehrlichia* species and *Borrelia burgdorferi* in *Ixodes* ticks in the Baltic regions of Russia. *J. Clin. Microbiol.* **39**:2237–2242.
- Allsopp, M. T. E. P., and B. A. Allsopp. 2001. Novel *Ehrlichia* genotype detected in dogs in South Africa. *J. Clin. Microbiol.* **39**:4204–4207.
- Anderson, B. E., K. G. Sims, and J. G. Olson. 1993. *Amblyomma americanum*: a potential vector of human ehrlichiosis. *Am. J. Trop. Med. Hyg.* **49**:239–244.
- Baek, L. J., J. W. Song, and H. W. Lee. 1989. Serologic diagnosis of acute febrile hemorrhagic disease patients in Korea, 1988. *J. Korean Soc. Virol.* **19**:117–125.
- Barlough, J. E., J. E. Madigan, E. Derock, and L. Bigornia. 1996. Nested polymerase chain reaction for detection of *Ehrlichia equi* genomic DNA in horses and ticks (*Ixodes pacificus*). *Vet. Parasitol.* **63**:319–329.
- Cao, W. C., Y. M. Gao, and P. H. Zhang. 2000. Identification of *Ehrlichia chaffeensis* by nested PCR in ticks from southern China. *J. Clin. Microbiol.* **38**:2778–2780.
- Castro, M. B., W. L. Nicholson, L. C. Whitworth, J. C. Fox, and A. A. Kocan. 2001. Persistent infection in *Neotoma fuscipes* (Muridae: Sigmodontinae) with *Ehrlichia phagocytophilum* sensu lato. *Am. J. Trop. Med. Hyg.* **65**:261–267.
- Chae, J. S., C. M. Kim, E. H. Kim, E. J. Hur, T. A. Klein, T. K. Kang, H. C. Lee, and J. W. Song. 2003. Molecular epidemiological study for tick-borne disease (*Ehrlichia* and *Anaplasma* species) surveillance at selected U.S. military training sites/installations in Korea. *Ann. N. Y. Acad. Sci.* **990**:118–125.
- Chen, S. M., J. S. Dumler, J. S. Bakker, and D. H. Walker. 1994. Identification of a granulocytotropic *Ehrlichia* species as the etiologic agent of human disease. *J. Clin. Microbiol.* **32**:589–595.
- Chiba, N., M. Osada, K. Komoro, T. Mizutani, H. Kariwa, and I. Takashima. 1999. Protection against tick-borne encephalitis virus isolated in Japan by active and passive immunization. *Vaccine* **17**:1532–1539.
- Dawson, J. E., D. E. Stallknecht, E. W. Howerth, C. Warner, K. Biggie, W. R. Davidson, J. M. Lockhart, V. F. Nettles, J. G. Olson, and J. E. Childs. 1994. Susceptibility of white-tailed deer (*Odocoileus virginianus*) to infection with *Ehrlichia chaffeensis*, the etiologic agent of human ehrlichiosis. *J. Clin. Microbiol.* **32**:2725–2728.
- Dawson, J. E., K. L. Biggie, C. K. Warner, K. Cookson, S. Jenkins, J. F. Levine, and J. G. Olson. 1996. Polymerase chain reaction evidence of *Ehrlichia chaffeensis*, an etiologic agent of human ehrlichiosis, in dogs from southeast Virginia. *Am. J. Vet. Res.* **57**:1175–1179.
- Fernandez, S. P., S. R. Perez, and G. A. Encinas. 2001. Molecular detection of *Ehrlichia phagocytophilum* genogroup organisms in larvae of *Neotrombicula autumnalis* (Acari: Trombiculidae) captured in Spain. *J. Parasitol.* **87**:1482–1483.
- Fournier, P. E., and D. Raoult. 2004. Suicide PCR on skin biopsy specimens for diagnosis of rickettsioses. *J. Clin. Microbiol.* **42**:3428–3434.
- Furuya, Y., T. Katayama, Y. Yoshida, and I. Kaiho. 1995. Specific amplification of *Rickettsia japonica* DNA from clinical specimens by PCR. *J. Clin. Microbiol.* **33**:487–489.
- Galvao, M. A., J. A. Lamounier, E. Bonomo, M. S. Tropaia, E. G. Rezende, S. B. Calic, C. B. Chamone, M. C. Machado, M. E. Otoni, R. C. Leite, C. Caram, C. L. Mafra, and D. H. Walker. 2002. Emerging and reemerging rickettsiosis in an endemic area of Minas Gerais State, Brazil. *Cad. Saude Publica* **18**:1593–1597.
- Heo, E. J., J. H. Park, J. R. Koo, M. S. Park, M. Y. Park, J. S. Dumler, and J. S. Chae. 2002. Serologic and molecular detection of *Ehrlichia chaffeensis* and *Anaplasma phagocytophilum* (human granulocytic ehrlichiosis agent) in Korean patients. *J. Clin. Microbiol.* **40**:3082–3085.
- Hulinska, D., K. Langrova, M. Pejcoch, and I. Pavlasek. 2004. Detection of *Anaplasma phagocytophilum* in animals by real-time polymerase chain reaction. *APMIS* **112**:239–247.
- Im, K. I., I. Y. Lee, and W. J. Lee. 1998. A human case of tick bite by *Ixodes persulcatus*. *Korean J. Parasitol.* **36**:63–65.
- Inokuma, H., I. Brouqui, M. Drancourt, and D. Raoult. 2001. Citrate synthase gene sequence: a new tool for phylogenetic analysis and identification of *Ehrlichia*. *J. Clin. Microbiol.* **39**:3031–3039.
- Inokuma, H., T. Beppu, M. Okuda, Y. Shimada, and Y. Sakata. 2001. Detection of ehrlichial DNA in *Haemaphysalis* ticks recovered from dogs in Japan that is closely related to a novel *Ehrlichia* sp. found in cattle ticks from Tibet, Thailand, and Africa. *J. Clin. Microbiol.* **42**:1353–1355.
- Ishiguro, F., N. Takada, T. Masuzawa, and T. Fukui. 2000. Prevalence of Lyme disease *Borrelia* species in ticks from migratory birds on the Japanese mainland. *Appl. Environ. Microbiol.* **66**:982–986.
- Jang, W. J., J. H. Kim, Y. J. Choi, K. D. Jung, Y. G. Kim, S. H. Lee, M. S. Choi, I. S. Kim, D. H. Walker, and K. H. Park. 2004. First serologic evidence of human spotted fever group rickettsiosis in Korea. *J. Clin. Microbiol.* **42**:2310–2313.
- Kawahara, M., T. Ito, C. Suto, S. Shibata, Y. Rikihisa, K. Hata, and K. Hirai. 1999. Comparison of *Ehrlichia muris* strains isolated from wild mice and ticks and serologic survey of humans and animals with *E. muris* as antigen. *J. Clin. Microbiol.* **37**:1123–1129.
- Kim, C. M., J. Y. Kim, Y. H. Yi, M. J. Lee, M. R. Cho, D. H. Shah, T. A. Klein, H. C. Kim, J. W. Song, S. T. Chong, M. L. O'Guinn, J. S. Lee, I. Y. Lee, J. H. Park, and J. S. Chae. 2005. Detection of *Bartonella* species from ticks, mites and small mammals in Korea. *J. Vet. Sci.* **6**:327–334.
- Kim, C. M., M. S. Kim, M. S. Park, J. H. Park, and J. S. Chae. 2003. Identification of *Ehrlichia chaffeensis*, *Anaplasma phagocytophilum*, and *A. bovis* in *Haemaphysalis longicornis* and *Ixodes persulcatus* ticks from Korea. *Vector Borne Zoonotic Dis.* **3**:17–26.
- Ko, J. H., D. Y. Cho, B. S. Chung, and S. I. Kim. 2002. Two human cases of tick bite caused by *Ixodes nipponensis*. *Korean J. Parasitol.* **40**:199–203.
- Lee, J. H., H. S. Park, K. D. Jung, W. J. Jang, S. E. Koh, S. S. Kang, I. Y. Lee, W. J. Lee, B. J. Kim, Y. H. Kook, K. H. Park, and S. H. Lee. 2003. Identification of the spotted fever group rickettsiae detection from *Haemaphysalis longicornis* in Korea. *Microbiol. Immunol.* **47**:301–304.
- Lee, S. O., D. K. Na, C. M. Kim, Y. H. Li, Y. H. Cho, J. H. Park, J. H. Lee, S. K. Eo, T. A. Klein, and J. S. Chae. 2005. Identification and prevalence of *Ehrlichia chaffeensis* infection in *Haemaphysalis longicornis* ticks from Korea by PCR, sequencing and phylogenetic analysis based on 16S rRNA gene. *J. Vet. Sci.* **6**:151–155.
- Leutenegger, C. M., N. Pusteria, R. Wicki, and H. Lutz. 2002. New molecular biology detection methods for tick-borne infectious agents. *Schweiz Arch. Tierheilkd.* **144**:395–404.
- Lofits, A. D., R. F. Massung, and M. L. Levin. 2003. Quantitative real-time PCR assay for detection of *Ehrlichia chaffeensis*. *J. Clin. Microbiol.* **41**:3870–3872.
- Long, S. W., X. Zhang, J. Zhang, R. P. Ruble, P. Teel, and X. J. Yu. 2003. Evaluation of transovarial transmission and transmissibility of *Ehrlichia chaffeensis* (Rickettiales: Anaplasmataceae) in *Amblyoma americanus* (Acari: Ixodidae). *J. Med. Entomol.* **40**:1000–1004.
- Maeda, K., N. Markowitz, R. C. Hawley, M. Ristic, D. Cox, and J. E. McDade. 1987. Human infection with *Ehrlichia canis*, a leukocytic rickettsia. *N. Engl. J. Med.* **316**:853–856.
- Magnarelli, L. A., J. F. Anderson, K. C. Stafford, and J. S. Dumler. 1997. Antibodies to multiple tick-borne pathogens of babesiosis, ehrlichiosis, and Lyme borreliosis in white-footed mice. *J. Wildl. Dis.* **33**:466–473.
- Mathew, J. S., S. A. Ewing, G. L. Murphy, K. C. Kocan, R. E. Cortvet, and J. C. Fox. 1997. Characterization of a new isolate of *Ehrlichia plays* (order Rickettiales) using electron microscopy and polymerase chain reaction. *Vet. Parasitol.* **68**:1–10.
- Murphy, G. L., S. A. Ewing, L. C. Whitworth, J. C. Fox, and A. A. Kocan. 1998. A molecular and serologic survey of *Ehrlichia canis*, *E. chaffeensis*, and *E. ewingii* in dogs and ticks from Oklahoma. *Vet. Parasitol.* **79**:325–339.
- Ogden, N. H., K. Bown, and B. K. Horrocks. 1998. Granulocytic *Ehrlichia* infection in ixodid ticks and mammals in woodlands and uplands of the U.K. *Med. Vet. Entomol.* **12**:423–429.
- Oh, S. H., Y. H. Song, D. H. Yoo, S. Y. Kim, and H. Lee. 1993. Distribution of *Borrelia burgdorferi* specific antibody among patients with juvenile rheumatoid arthritis in Korea. *J. Korean Med. Sci.* **8**:405–412.
- Park, J. H., E. J. Heo, K. S. Choi, J. S. Dumler, and J. S. Chae. 2003. Detection of antibodies to *Anaplasma phagocytophilum* and *Ehrlichia chaffeensis* antigens in sera of Korean patients by Western immunoblotting

- and indirect immunofluorescence assays. *Clin. Diagn. Lab. Immunol.* **10**:1059–1064.
41. Pusterla, N., J. B. Huder, C. M. Leutenegger, U. Braun, E. John, J. E. Madigan, and H. Lutz. 1999. Quantitative real-time PCR for detection of members of the *Ehrlichia phagocytophila* genogroup in host animals and *Ixodes ricinus* ticks. *J. Clin. Microbiol.* **37**:1329–1331.
42. Regnery, R. L., C. L. Spruill, and B. D. Plikaytis. 1991. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J. Bacteriol.* **173**:1576–1589.
43. Sachar, D. S. 2000. *Ehrlichia chaffeensis* infection in an active duty soldier stationed in Korea. *MSMR* **6**:9–11.
44. Schouls, L. M., I. Van De Pol, S. G. Rijpkema, and C. S. Schot. 1999. Detection and identification of *Ehrlichia*, *Borrelia burgdorferi* sensu lato, and *Bartonella* species in Dutch *Ixodes ricinus* ticks. *J. Clin. Microbiol.* **37**:2215–2222.
45. Scola, B. L., Z. Liang, Z. Zeaiter, P. Houpijian, P. A. D. Grimont, and D. Raoult. 2002. Genotypic characteristics of two serotypes of *Bartonella henselae*. *J. Clin. Microbiol.* **40**:2002–2068.
46. Song, G., C. S. Hang, X. Z. Qui, D. S. Ni, H. X. Liao, G. Z. Gao, Y. L. Du, J. K. Xu, Y. S. Wu, J. N. Zhao, B. X. Kong, Z. S. Wang, Z. Q. Zhang, H. K. Shen, and N. Zhou. 1983. Etiologic studies of epidemic hemorrhagic fever (hemorrhagic fever with renal syndrome). *J. Infect. Dis.* **147**:654–659.
47. Sparagano, O. A. E., M. T. E. P. Allsopp, R. A. Mank, S. G. T. Rijpkema, J. V. Figueroa, and F. Jongejan. 1999. Molecular detection of pathogen DNA in ticks (Acari: Ixodidae): a review. *Exp. Appl. Acarol.* **23**:929–960.
48. Swofford, D. L. 2002. PAUP*. Phylogenetic analysis using parsimony (*and other methods), version 4. Sinauer Associates, Sunderland, Mass.
49. Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **24**:4876–4882.
50. von Loewenich, F. D., G. Stumpf, B. U. Baumgarten, M. Rollinghoff, J. S. Dumler, and C. Bogdan. 2003. A case of equine granulocytic ehrlichiosis provides molecular evidence for the presence of pathogenic *Anaplasma phagocytophilum* (HGE agent) in Germany. *Eur. J. Clin. Microbiol. Infect. Dis.* **22**:303–305.
51. Walls, J. J., B. Greig, D. F. Neitzel, and J. S. Dumler. 1997. Natural infection of small mammal species in Minnesota with the agent of human granulocytic ehrlichiosis. *J. Clin. Microbiol.* **35**:853–855.
52. Wang, I. N., D. E. Dykhulzen, W. Qiu, J. J. Dunn, E. M. Bosler, and B. J. Luft. 1999. Genetic diversity of *ospC* in a local population of *Borrelia burgdorferi* sensu stricto. *Genetics* **151**:15–30.
53. Yabsley, M. J., A. S. Varela, and C. M. Tate. 2002. *Ehrlichia ewingii* infection in white-tailed deer (*Odocoileus virginianus*). *Emerg. Infect. Dis.* **8**:668–671.
54. Yamaguti, N., V. J. Tipton, H. L. Keegan, and S. Toshioka. 1971. Ticks of Japan, Korea, and the Ryukyu Islands. Brigham Young Univ. Sci. Bull. Biol. Ser. **15**:1–225.