

# Tides in the Mandovi and Zuari estuaries, Goa, west coast of India

D SUNDAR and S R SHETYE

*Physical Oceanography Division, National Institute of Oceanography, Dona Paula, Goa 403 004, India.*

Mandovi and Zuari are two estuaries located in Goa, west coast of India. Variation of water level in the estuaries was monitored for a month at 13 locations using tide-poles during March–April 2003. Analysis of this data has provided for the first time, characteristics of how tidal constituents vary in the narrow and shallow estuaries, typical of those found along the west coast of India. At a distance of 45 km from the mouth the tidal range increased in both estuaries by approximately 20%. The tidal range at the upstream end of the two channels at the stations dropped sharply because of the increase in elevation of the channels.

## 1. Introduction

Located along the west coast of India are a number of estuarine channels that connect the Arabian Sea to the rivers that originate in the Western Ghats mountain range. The Western Ghats rise sharply from the coastline to an average height of about 1000 m. The runoff in the rivers is high during June–September when the Indian summer monsoon is active. At this time winds along the coast are approximately westerly and carry moisture picked up during their passage over the north Indian Ocean, including the Arabian Sea. As a result, the windward side of the Ghats receives high precipitation due to the topographic effect. The rainfall along the central west coast of India is about 250 cm during June–September.

The Mandovi and the Zuari are two estuaries that are located along the west coast, in Goa (figure 1). Each of these two estuaries is about 5 m deep. Their cross-sectional area decreases from mouth to head, and tides occur in the two estuaries up to a distance of about 50 km (Shetye *et al* 1995). The increase in elevation of the estuarine channels prevents tides from propagating beyond this distance. The runoff in the two estuaries is highly seasonal, just as is the precipitation. This can be seen from figure 2 that gives runoff measured by a gauge

located at Ganjem on the Mandovi river (figure 1). The gauge is located just beyond the distance up to which variation in water level due to tides is observed in the channel. The runoff recorded by the gauge is typical of the runoff observed in the rivers located along the west coast. The runoff peaks during the most active phase of the Indian summer monsoon, decreases rapidly after withdrawal of the monsoon and remains negligible from about January till the onset of the next monsoon which generally occurs during late May or early June.

The flow in the estuarine channels is primarily tidal after withdrawal of the monsoon, and continues to be so until onset of the next monsoon. In 1993, tide pole measurements were carried out at 15 locations in the two estuaries for three days. The inferences derived from analysis of these data were that the amplitude of the tide remains virtually unchanged in the channels, and that phase propagates from mouth to head with an average speed of about 6 m/s (Shetye *et al* 1995). However, the time-series of the data collected then, being only 3 days long, did not permit determination of characteristics of tidal constituents.

During March–April 2003, tide pole measurements similar to those during 1993 were carried out at 13 locations by recording the water level

**Keywords.** Harmonic analysis; tidal constituents; tide-pole; amplification; channel geometry.




Figure 1. A map of the Mandovi and Zuari estuaries. The locations where water level observations were carried out are marked with a number in a circle. The name of the location is given at the bottom of the map. **T** indicates the location up to which variation in water level has been reported to occur. The depth contours are in m.

once every fifteen minutes. In the present paper we report analysis of these data. The new time-series, being at least 30 days long, allows us to carry out tidal analysis of 34 constituents. The next section describes the data collected and the method of analysis. The results are presented in section 3. Section 4 provides a brief summary of the study.

## 2. Data collection and analysis

The 13 locations where tide pole measurements were carried out during March–April 2003 are shown in figure 1. The zero of each tide-pole used in the observations was referenced to a ‘local chart-datum’ for future reference. The tide pole readers worked in three shifts during a day. Each reader

was requested to visually average the water level for about a minute and then record the average water level. It is, however, not clear what the error associated with such a measurement is. Below we make an estimate of the error involved in the estimation of amplitude and phase lag of each constituent.

The distance of each of the 13 stations from the mouth, and start-and-end time of observation are given in table 1. The 15-minute data were analyzed using TASK-2000 (Tidal Analysis Software Kit 2000) of the Permanent Service for Mean Sea Level, Proudman Oceanography Laboratory, UK (Bell *et al* 1998) to determine tidal amplitude and phase lag of 26 major constituents and 8 related constituents. The analysis served as a tool for quality control too. Data points with large residuals were examined to determine if the residuals




Figure 2. Monthly-mean runoff ( $\text{m}^3/\text{s}$ ) at Ganjem in the Mandovi river (for location see figure 1). The top (bottom) of an error bar represents the maximum (minimum) discharge during a month. Climatological data (1980–97). Courtesy: Central Water Commission, New Delhi.

were due to errors in recording. If inferred to be so, the suspect data were removed. All computations are referenced to the Indian Standard Time which leads the Greenwich Mean Time by 330 minutes.

The main channels of the Mandovi and the Zuari are connected by the Cumbarjua Canal. Its cross-sectional area is, however, much too small to have a major impact on the characteristics of tidal

propagation in the two main channels. Of the 13 stations where the observations were carried out, 6 were located in the Mandovi main channel, 5 in the Zuari main channel, and 2 in the Cumbarjua Canal (figure 1). The quality controlled data at these 13 locations are shown in figure 3. Tidal analysis was carried out at only 11 stations. The water level variation at Ganjem and Sanguem was found to exhibit tidal effect only a part of the time. This is so because of the higher topographic elevation of these stations. We have discussed this point in section 3.2. The data from these stations were not subjected to tidal analysis.

Noting that the Mandovi–Zuari is the only estuarine system along the west coast of India where variation of amplitude and phase lag along the length of the estuarine channel is now known, we provide in tables 2 and 3 all the inferred tidal characteristics at the 11 locations. Of the 34 constituents whose amplitudes and phases were determined, 5 constituents –  $M_2$ ,  $S_2$ ,  $N_2$ ,  $K_1$ , and  $O_1$  – had amplitudes larger than 10 cm at the mouth of the two estuaries. In the next section we discuss the behaviour of these constituents.

In the rest of this section we make an estimate of the error associated with our determination of amplitude and phase lag of the 34 constituents listed in tables 2 and 3. The procedure followed for this purpose is as follows: The amplitudes and phase lags given in tables 2 and 3 were computed from time-series with 15 minute separation between consecutive values. The amplitudes and phase lags were recomputed using a time series in which the time interval between two consecutive values was two hours. As the original data are

Table 1. Distance of each of the 13 stations where the observations were carried out and the start-and-end time of the observations.

Station	Distance from mouth (km)	Start time	End time
<b>Mandovi</b>			
Verem	4.81	0000 hrs/10th March	2345 hrs/13th April
Britona	8.86	0000 hrs/10th March	2345 hrs/13th April
Akkada	20.31	0000 hrs/10th March	2345 hrs/13th April
Volvoi	32.91	0000 hrs/10th March	2345 hrs/13th April
Usgao	41.07	0000 hrs/10th March	2345 hrs/13th April
Ganjem	49.83	0000 hrs/10th March	2345 hrs/13th April
<b>Zuari</b>			
Dona Paula	0.00	0000 hrs/11th March	2345 hrs/13th April
Cortalim	11.70	0000 hrs/10th March	2345 hrs/13th April
Borim	21.02	0000 hrs/10th March	2345 hrs/13th April
Sanvordem	45.22	0000 hrs/10th March	2345 hrs/13th April
Sanguem	54.94	0000 hrs/11th March	2345 hrs/13th April
<b>Cumbarjua canal from Dona Paula through Zuari</b>			
Madkai	14.74	0000 hrs/10th March	2345 hrs/13th April
Banastarim	24.61	0000 hrs/10th March	2345 hrs/13th April




Figure 3. Observed sea level (cm) during the period of observation. The horizontal axis gives date and month.

separated by 15 minutes, we get 8 independent 2-hour time series. From these we get 8 values for amplitude and phase lag of each constituent at a station. We then computed standard deviation of both amplitude and phase lag. We take the

standard deviation to be indicative of the error associated with our determination of amplitudes and phase lag. Tables 4 and 5 give the values obtained through this exercise at one station, Akkada in the Mandovi (location 3 in figure 1).

Table 2. Amplitude (cm) of the tidal constituents at the 11 stations.

Constit.	Mandovi					Zuari					Cumbarjua	
	Verem	Britona	Akkada	Volvoi	Usgao	Dona Paula	Cortalim	Borim	Sanvordem	Madkai	Banastarim	
MM	5.7960	6.3502	6.9325	6.5066	8.5160	7.2330	7.1483	6.4123	4.1737	7.2916	6.7107	
MSF	4.3692	4.5059	4.8061	5.0299	4.8080	3.6859	2.3511	5.1460	4.2798	6.5520	5.0396	
Q1	3.8667	3.9718	4.2544	4.0092	3.9122	5.2581	6.5315	3.7028	5.4631	3.7558	4.0379	
O1	14.9116	14.7849	14.5834	14.3197	15.0861	16.9912	14.7654	13.9013	16.2402	14.1031	14.3811	
M1	0.8006	0.7554	0.8978	0.7999	1.1474	1.7379	1.8909	0.9846	1.9506	1.0011	0.8044	
PI1	0.6378	0.6418	0.6498	0.6761	0.6971	0.6169	0.6746	0.7376	0.7114	0.6627	0.6557	
P1	11.1120	11.1813	11.3199	11.7783	12.1439	10.7474	11.7531	12.8499	12.3925	11.5456	11.4235	
K1	33.5709	33.7804	34.1991	35.5839	36.6885	32.4694	35.5078	38.8214	37.4396	34.8810	34.5121	
PS11	0.2686	0.2702	0.2736	0.2847	0.2935	0.2598	0.2841	0.3106	0.2995	0.2790	0.2761	
PH11	0.4700	0.4729	0.4788	0.4982	0.5136	0.4546	0.4971	0.5435	0.5242	0.4883	0.4832	
J1	1.9320	1.4386	0.9917	1.2233	1.9873	1.4200	3.3896	1.4688	3.6067	0.5705	1.4530	
OO1	1.4072	1.3750	1.4471	1.7244	0.8233	2.0291	1.2337	2.4624	1.3564	2.4405	1.6128	
2N2	1.6407	1.6557	1.7157	1.9134	2.1119	1.6527	1.6418	2.1090	1.8569	1.9180	1.7820	
MU2	0.0780	1.1833	3.6507	4.6092	5.0406	0.9390	2.9748	3.2009	6.5673	2.2910	3.0707	
N2	12.3358	12.4486	12.8998	14.3861	15.8789	12.4265	12.3444	15.8573	13.9618	14.4210	13.3986	
NU2	2.3931	2.4150	2.5026	2.7909	3.0805	2.4107	2.3948	3.0763	2.7086	2.7977	2.5993	
M2	53.2327	52.4094	55.3923	61.1507	63.6825	54.0399	59.9542	63.8844	69.7756	57.6406	55.7980	
L2	3.0189	3.1784	3.7990	5.4268	6.4047	3.6251	2.3742	4.7417	10.4019	2.3000	4.0349	
T2	1.0749	1.0304	1.0576	1.1614	1.2212	1.2231	1.1700	1.2800	1.3560	1.1572	1.0909	
S2	18.2183	17.4640	17.9257	19.6847	20.6977	20.7309	19.8300	21.6953	22.9826	19.6141	18.4902	
K2	4.9554	4.7502	4.8758	5.3542	5.6298	5.6388	5.3938	5.9011	6.2513	5.3350	5.0293	
2SM2	0.3988	0.7077	1.3826	1.7766	1.3981	1.8490	1.9399	1.8844	4.0068	1.6173	1.3009	
MO3	0.1199	0.8459	1.9702	2.3468	2.6092	0.8148	3.1712	1.5370	1.6668	1.4653	1.9088	
M3	0.7559	0.5632	0.6335	0.7383	1.7180	1.4647	2.5311	1.4922	2.4183	0.6037	0.9923	
MK3	0.7339	0.3804	1.0762	2.2508	2.3297	1.4783	0.8182	1.2955	3.6295	1.8395	0.9406	
MN4	0.5157	0.7535	0.1326	0.8368	1.7633	0.5231	1.2053	1.4805	2.4239	0.2897	0.8368	
M4	0.6492	1.5864	1.2806	1.7017	2.3047	1.2956	2.7667	3.6605	4.9818	0.5086	2.0113	
SN4	0.2170	0.0397	0.4198	0.8743	1.7067	1.9948	1.6167	0.7135	2.5750	0.9806	0.4596	
MS4	0.5289	0.8120	0.6674	1.8081	2.5361	1.7895	0.5935	2.1739	4.5593	0.5092	1.0826	
2MN6	0.2133	0.3278	0.8060	1.5734	1.8862	0.5193	0.3044	0.6693	1.1902	1.2015	1.1943	
M6	0.4696	0.6528	0.9869	1.9981	2.2832	0.7767	0.4238	0.8482	1.8638	0.3634	1.5668	
MSN6	0.1348	0.3415	0.7338	0.6696	0.8154	0.8014	1.1674	0.7638	0.9139	0.4402	0.6953	
2MS6	0.1640	0.9609	1.2553	2.4757	2.8575	0.9450	1.4654	0.6079	1.9691	0.3685	1.6505	
2SM6	0.2228	0.2950	0.5249	1.0133	1.2801	0.5935	0.5194	0.6142	0.9933	0.4353	0.8669	

Table 3. Same as table 2 but for phase lag in degrees (referenced to IST). Legend of \* marks is in table 5.

Constit.	Mandovi				Zuari				Cumbarjua		
	Verem	Britona	Akkada	Volvoi	Usgao	Dona Paula	Cortalim	Borim	Sanvordem	Madkai	Banastarim
MM	93.168	84.506	85.333	80.715	70.841	97.584	64.113	83.451	54.975	115.161	91.583
MSF	342.181	349.031	359.897	5.155	354.315	308.930	354.020	2.417	15.762	343.093	3.236
Q1	59.817	66.967	71.599	77.692	83.746	59.489	55.732	78.458	61.577	81.155	74.609
O1	51.211	54.789	60.879	62.744	67.152	55.655	62.675	52.154	62.934	42.686	56.437
M1	57.990	71.966	62.661	87.009	92.405	24.973	91.983	66.167	140.184	33.878	81.993
*PII (K1)	61.568	67.092	75.001	77.568	83.265	67.726	63.174	70.087	78.365	73.526	72.002
*P1 (K1)	61.568	67.092	75.001	77.568	83.265	67.726	63.174	70.087	78.365	73.526	72.002
K1	61.568	67.092	75.001	77.568	83.265	67.726	63.174	70.087	78.365	73.526	72.002
*PSII(K1)	61.568	67.092	75.001	77.568	83.265	67.726	63.174	70.087	78.365	73.526	72.002
*PHII(K1)	61.568	67.092	75.001	77.568	83.265	67.726	63.174	70.087	78.365	73.526	72.002
J1	56.678	84.065	104.115	106.993	154.074	30.960	97.302	90.027	89.249	118.081	82.309
OO1	113.743	122.276	160.307	165.684	202.507	65.870	155.675	141.197	192.215	144.060	148.469
*2N2 (N2)	299.530	312.355	329.152	336.316	345.170	288.132	292.307	318.000	339.378	298.568	322.417
MU2	63.150	36.318	45.676	49.488	64.667	138.421	40.387	29.098	32.253	75.812	43.136
N2	299.530	312.355	329.152	336.316	345.170	288.132	292.307	318.000	339.378	298.568	322.417
*NU2 (N2)	299.530	312.355	329.152	336.316	345.170	288.132	292.307	318.000	339.378	298.568	322.417
M2	318.239	329.646	345.560	353.312	0.005	313.743	319.644	335.014	350.163	325.452	340.493
L2	318.400	308.134	310.170	308.044	298.184	335.463	251.625	288.256	276.407	321.333	306.208
*T2 (S2)	358.172	11.117	32.536	41.343	51.038	359.737	3.110	22.095	48.896	15.440	26.465
S2	358.172	11.117	32.536	41.343	51.038	359.737	3.110	22.095	48.896	15.440	26.465
*K2 (S2)	358.172	11.117	32.536	41.343	51.038	359.737	3.110	22.095	48.896	15.440	26.465
2SM2	162.165	184.527	210.337	218.747	266.609	69.198	166.952	179.735	225.894	153.167	204.297
MO3	110.857	51.665	28.116	46.791	68.197	23.564	13.571	64.874	41.802	30.289	50.146
M3	351.502	356.598	350.873	353.028	351.267	200.715	208.194	356.851	134.828	312.832	349.503
MK3	245.539	279.957	326.938	329.147	12.755	346.305	47.453	250.399	305.871	228.363	295.895
MN4	25.171	72.298	284.806	199.220	237.212	11.338	77.730	135.762	144.731	47.637	99.314
M4	54.141	91.561	158.405	201.016	259.898	40.858	117.115	140.600	157.348	149.673	116.918
SN4	278.840	63.432	358.221	1.241	18.997	274.202	267.853	311.854	267.327	277.105	306.023
MS4	166.005	132.658	253.495	292.914	344.131	3.921	136.841	198.032	244.340	193.924	210.151
2MN6	254.105	292.718	335.286	23.630	45.042	191.197	263.367	294.596	18.548	247.722	291.547
M6	301.019	292.527	5.050	44.078	67.567	242.441	284.022	327.921	44.350	270.603	318.245
MSN6	353.656	325.455	23.451	57.875	96.931	251.599	258.679	352.705	91.652	334.240	355.833
2MS6	25.959	6.022	50.616	86.870	102.193	263.795	315.571	8.959	80.296	43.345	0.637

Table 4. Estimation of error (cm) associated with the determination of amplitude (cm) at Akkada. From the original data at 15 min intervals, 8 data sets were derived by sub-sampling at 2-hour intervals. The column *x000* contains the data from values at 0000 hr, 0200 hr, 0400 hr and so on. The column *x015* contains the data from values at 0015 hr, 0215, 0415 and so on. Similar is the case for *x030*, *x045*, *x100*, *x115*, *x130* and *x145*. The mean and standard deviation were calculated using these eight columns. The values in the column *x* are from ALL the original values from data at 15 min intervals.

Constit.	x000	x015	x030	x045	x100	x115	x130	x145	Mean	Std. dev.	x
MM	6.88	7.03	6.99	6.89	6.89	6.94	6.90	6.94	6.93	0.052	6.93
MSF	4.79	4.94	4.82	4.75	4.83	4.85	4.75	4.73	4.81	0.065	4.81
Q1	4.20	4.21	4.29	4.35	4.33	4.18	4.24	4.25	4.26	0.058	4.25
O1	14.59	14.63	14.57	14.61	14.47	14.47	14.60	14.72	14.58	0.080	14.58
M1	0.87	0.91	0.89	0.84	0.93	0.93	0.91	0.91	0.90	0.028	0.90
PI1	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.001	0.65
P1	11.34	11.34	11.31	11.29	11.36	11.33	11.29	11.30	11.32	0.023	11.32
K1	34.25	34.26	34.16	34.12	34.31	34.24	34.12	34.12	34.20	0.070	34.20
PSI1	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.001	0.27
PHI1	0.48	0.48	0.48	0.48	0.48	0.48	0.48	0.48	0.48	0.001	0.48
J1	0.94	0.93	0.85	0.91	1.09	1.11	1.07	1.06	1.00	0.092	0.99
OO1	1.31	1.37	1.37	1.53	1.68	1.54	1.42	1.34	1.45	0.119	1.45
2N2	1.73	1.71	1.70	1.71	1.70	1.72	1.73	1.73	1.72	0.011	1.72
MU2	3.74	3.74	3.71	3.75	3.65	3.46	3.57	3.59	3.65	0.097	3.65
N2	12.98	12.83	12.79	12.86	12.81	12.91	13.03	12.99	12.90	0.086	12.90
NU2	2.52	2.49	2.48	2.49	2.48	2.50	2.53	2.52	2.50	0.017	2.50
M2	55.43	55.32	55.25	55.25	55.26	55.41	55.68	55.53	55.39	0.144	55.39
L2	3.73	3.87	3.96	3.79	3.89	3.78	3.72	3.67	3.80	0.090	3.80
T2	1.05	1.06	1.06	1.07	1.06	1.05	1.05	1.05	1.06	0.005	1.06
S2	17.86	17.96	17.96	18.10	17.98	17.85	17.86	17.84	17.93	0.084	17.93
K2	4.86	4.89	4.89	4.92	4.89	4.85	4.86	4.85	4.88	0.023	4.88
2SM2	1.55	1.52	1.31	0.99	1.26	1.49	1.55	1.50	1.39	0.183	1.38
MO3	1.81	2.04	2.04	2.00	2.02	2.02	1.96	1.87	1.97	0.079	1.97
M3	0.74	0.62	0.59	0.67	0.64	0.51	0.64	0.73	0.64	0.069	0.63
MK3	1.08	1.07	1.13	0.95	1.09	1.22	1.17	0.96	1.08	0.088	1.08
MN4	0.31	0.26	0.29	0.38	0.31	0.30	0.36	0.14	0.30	0.067	0.13
M4	1.37	1.29	1.26	1.19	1.17	1.26	1.30	1.42	1.28	0.079	1.28
SN4	0.25	0.27	0.39	0.56	0.63	0.64	0.41	0.29	0.43	0.151	0.42
MS4	0.86	0.62	0.69	0.56	0.53	0.58	0.74	0.80	0.67	0.111	0.67
2MN6	0.82	0.91	0.87	0.78	0.73	0.68	0.77	0.91	0.81	0.078	0.81
M6	1.10	1.01	1.00	0.89	0.90	0.94	1.07	1.08	1.00	0.075	0.99
MSN6	0.71	0.48	0.59	0.76	0.79	0.93	0.94	0.84	0.76	0.151	0.73
2MS6	1.45	1.41	1.40	1.21	1.11	1.13	1.21	1.14	1.26	0.130	1.26
2SM6	0.58	0.46	0.33	0.37	0.56	0.65	0.76	0.62	0.54	0.137	0.52

After examining the corresponding tables at all the stations, we concluded that for all constituents whose amplitude is greater than 3 cm, the upper limit of the error bar of the values listed in tables 4 and 5 is  $\pm 0.5$  cm for amplitude and  $\pm 5^\circ$  for phase lag.

### 3. Results

#### 3.1 Distribution of amplitude and phase lag

Variation of amplitude and phase lag with distance from mouth to head in an estuary with a

converging, narrow and shallow channel can be looked at as resulting from two waves: an incident shallow-water wave triggered by the tide at the mouth and a reflected shallow-water wave that gets generated as the incident wave propagates through the channel. In some channels the contribution of the reflected wave can be neglected (Friedrichs and Aubrey 1994). In general, however, both the waves need to be taken into account. When they are, the variation of amplitude and phase difference along the channel is a function of friction, geometry of the channel (rate of convergence of the channel from mouth

Table 5. Same as table 4 but for the phase lag in degrees (referenced to IST) at Akkada. The ‘star’ marked entries are related constituents which have not been computed independently, but computed in relation to the independent major constituents with larger amplitudes, given in brackets next to them. Hence is the same phase lag.

Constit.	x000	x015	x030	x045	x100	x115	x130	x145	mean	Std. dev.	x
MM	85.9	86.4	85.6	86.2	85.1	84.6	83.6	85.4	85.3	0.84	85.3
MSF	359.5	359.4	359.5	359.9	358.4	359.9	1.8	0.9	359.9	0.95	359.9
Q1	72.7	72.4	71.0	69.8	70.7	71.3	72.5	72.5	71.6	1.00	71.6
O1	61.0	60.9	60.9	60.8	61.0	60.6	61.0	60.8	60.9	0.12	60.9
M1	61.0	58.8	59.9	64.5	62.1	66.9	65.7	62.3	62.6	2.66	62.7
*PI1 (K1)	74.8	74.7	74.7	75.1	75.3	75.3	75.1	75.0	75.0	0.24	75.0
*P1 (K1)	74.8	74.7	74.7	75.1	75.3	75.3	75.1	75.0	75.0	0.24	75.0
K1	74.8	74.7	74.7	75.1	75.3	75.3	75.1	75.0	75.0	0.24	75.0
*PSI1 (K1)	74.8	74.7	74.7	75.1	75.3	75.3	75.1	75.0	75.0	0.24	75.0
*PHI1 (K1)	74.8	74.7	74.7	75.1	75.3	75.3	75.1	75.0	75.0	0.24	75.0
J1	100.6	97.3	103.5	112.7	107.3	106.1	104.1	100.9	104.1	4.44	104.1
OO1	161.9	157.7	157.4	161.5	160.0	159.9	162.3	161.8	160.3	1.78	160.3
*2N2 (N2)	329.9	329.0	329.3	328.9	328.6	328.3	329.1	330.2	329.2	0.60	329.2
MU2	44.8	44.7	45.8	48.1	46.0	45.1	44.6	46.4	45.7	1.11	45.7
N2	329.9	329.0	329.3	328.9	328.6	328.3	329.1	330.2	329.2	0.60	329.2
*NU2 (N2)	329.9	329.0	329.3	328.9	328.6	328.3	329.1	330.2	329.2	0.60	329.2
M2	345.8	345.6	345.6	345.8	345.5	345.2	345.3	345.7	345.6	0.18	345.6
L2	308.7	311.0	311.8	309.3	311.8	311.9	308.2	308.4	310.2	1.53	310.2
*T2 (S2)	32.5	32.1	32.1	32.6	32.8	32.8	32.7	32.7	32.5	0.26	32.5
S2	32.5	32.1	32.1	32.6	32.8	32.8	32.7	32.7	32.5	0.26	32.5
*K2 (S2)	32.5	32.1	32.1	32.6	32.8	32.8	32.7	32.7	32.5	0.26	32.5
2SM2	219.4	218.9	214.7	208.0	203.8	199.2	202.4	215.0	210.2	7.36	210.3
MO3	32.2	26.4	26.2	25.7	25.2	27.3	31.0	32.1	28.2	2.78	28.1
M3	0.1	355.5	345.5	338.2	335.2	347.3	0.6	1.7	350.5	9.79	350.9
MK3	330.8	321.7	320.0	323.3	325.9	325.7	332.5	336.7	327.1	5.37	326.9
MN4	206.3	230.0	276.5	278.0	307.4	343.2	39.3	143.2	228.0	91.99	284.8
M4	163.0	157.4	157.9	157.1	155.2	154.0	159.3	162.3	158.3	2.96	158.4
SN4	336.1	350.6	0.2	349.1	354.6	8.3	21.8	353.9	356.8	12.80	358.2
MS4	251.6	252.7	254.8	272.9	253.9	246.6	245.1	254.9	254.1	7.90	253.5
2MN6	343.4	344.9	345.4	336.2	326.6	329.3	322.9	329.9	334.8	8.32	335.3
M6	7.0	14.2	7.1	5.0	4.1	359.1	359.6	2.9	4.9	4.51	5.1
MSN6	40.9	19.8	13.3	4.2	7.1	20.0	33.2	38.4	22.1	13.12	23.5
2MS6	47.6	50.2	51.8	58.6	55.9	46.6	51.7	46.1	51.1	4.16	50.6
2SM6	120.5	99.6	97.2	59.7	80.5	64.4	87.3	100.3	88.7	18.83	89.6

to head) and tendency of the two waves to set up standing modes (Shetye 1999; Nayak and Shetye 2003).

Figure 4 shows how amplitude and phase lag of the most important tidal constituents (amplitude greater than 10 cm at the mouth) grow from mouth to head before topographic effect comes into play. Before the topographic influence is felt, amplitude of  $M_2$  increased by about 20% in the Mandovi from mouth to head, and by about 30% in the Zuari. The increase in  $K_1$  was by about 10% in the Mandovi and 15% in the Zuari. Phase lag increased from mouth to head for both diurnal and semi-diurnal constituents. However, this variation was

not linear with respect to distance from the mouth. A linear variation would have been expected if the reflected wave was absent. Further studies are needed to determine how the three processes – friction, geometric effect and co-oscillation of incident and reflected wave – set up the variation seen in figure 4.

### 3.2 Decay of tidal amplitude due to increase in channel elevation

Each of the 7 panels in figure 5 gives a section of the channel topography and water level along a line drawn through the middle of the Mandovi






Figure 4. Variation in amplitude (cm) and phase lag in degree (referenced to IST) of M<sub>2</sub>, S<sub>2</sub>, N<sub>2</sub>, K<sub>1</sub> and O<sub>1</sub> in the main channels of the Mandovi and Zuari estuaries.

channel. Both the topography and the water level are based on observations: the topography is the same as that described in Shetye *et al* (1995), and the water surface depicted in the figure is based on data collected during the 2003 observations. Two consecutive panels in the figure show how the water level changed during a period of two hours during spring tide. The panels together cover a typical 12-hour cycle during springs. The figure helps to visualize how the water level along the channel changes during a tidal cycle. And

hence, understand how the increase in channel elevation causes the tidal amplitude to drop at the upstream end in an estuarine channel. As seen from figure 3, the magnitude of water level variation drops sharply from Usgao to Ganjem in the Mandovi and from Sanvordem to Sanguem in the Zuari. From figure 5 it is seen that the tide reaches Ganjem in the Mandovi only when the tide is high enough to overcome the effect of increased channel elevation. The impact of the tide is felt at Ganjem in panels A, E, F, and G of the figure. In the rest




Figure 5. Each of the 7 panels in the figure gives an instantaneous section of the channel topography and water level along a line drawn through the middle of the Mandovi channel. Two consecutive panels are separated by two hours during spring tide. The water level at the upstream end is controlled by dynamics of the river flow.

of the panels no influence of the tide is seen at Ganjem and the water level is determined merely by whatever little runoff that exists in the river channel.

#### 4. Summary

The Mandovi and the Zuari are typical of the estuaries found along the west coast of India. The

estuaries are fed at the upstream end by rivers that originate on the slopes of the Western Ghats. In this paper we have described how the characteristics of tidal constituents change from mouth to head in these shallow ( $\sim 5$  m deep), narrow (usually less than few hundred meters wide, except near the mouth) and converging channels. To our knowledge this is the first time that such a description is available for the west coast estuaries. The exercise was carried out using tide-pole readers numbering about fifty. The present socio-economic conditions in this part of the world still make it more economical to employ tide-pole readers rather than self-reading tide-gauges. The estimated errors in the computed values of tidal constituents are small enough to conclude that the spatial pattern of variation in tidal constituents is dependable. Hence the method of measurement still remains attractive to gain insights into dynamics of the tide propagation in this estuary.

The results of this study show that both the diurnal and semi-diurnal tidal constituents amplify in the estuarine channels, the amplification being larger in the case of the semi-diurnal constituents. It is expected that the observed amplitude results from an incident and a reflected wave, both of which get altered by channel geometry and friction. Further research is needed to determine the contribution of each wave to the observed amplitude and phase lag.

## Acknowledgements

We are grateful to David Pugh and Philip Woodworth for reviewing the manuscript. Their comments have been very useful. We thank Shri Arun Y Mahale and Shri S Akerkar, DTP for their help in drawing figures 1 and 5. This is NIO contribution 4018.

## References

- Bell C, Vassie J M and Woodworth P L 1998 POL/PSMSL Tidal Analysis Software Kit 2000 (TASK-2000), Permanent Service for Mean Sea Level, UK, p. 21.
- Friedrichs C T and Aubrey D G 1994 Tidal propagation in strongly convergent channels; *J. Geophys. Res.* **99** C2, 3321–3336.
- Nayak R K and Shetye S R 2003 Tides in the Gulf of Khambhat, west coast of India; *Estuarine, Coastal and Shelf Science* **57** 249–254.
- Shetye S R 1999 Tides in the Gulf of Kutch, India; *Continental Shelf Research* **19** 1771–1782.
- Shetye S R, Gouveia A D, Singbal S Y, Naik C G, Sundar D, Michael G S and Nampoothiri G 1995 Propagation of tides in the Mandovi–Zuari estuarine network. *Proc. Indian Acad. Sci. (Earth Planet. Sci.)* **104**(4) 667–682.
- Unnikrishnan K, Shetye S R and Gouveia A D 1997 Tidal propagation in the Mandovi–Zuari estuarine network, west coast of India: Impact of fresh water influx; *Estuarine, Coastal and Shelf Science* **45** 737–744.