

REVIEW

Open Access

TIGAR, TIGAR, burning bright

Pearl Lee, Karen H Vousden* and Eric C Cheung*

Abstract

Cancers cells shift their metabolism towards glycolysis in order to help them support the biosynthetic demands necessary to sustain cell proliferation and growth, adapt to stress and avoid excessive reactive oxygen species (ROS) accumulation. While the p53 tumor suppressor protein is known to inhibit cell growth by inducing apoptosis, senescence and cell cycle arrest, recent studies have found that p53 is also able to influence cell metabolism. TIGAR is a p53 target that functions as a fructose-2,6-bisphosphatase, thereby lowering glycolytic flux and promoting antioxidant functions. By protecting cells from oxidative stress, TIGAR may mediate some of the tumor suppressor activity of p53 but could also contribute to tumorigenesis. Here we discuss the activities of TIGAR described so far, and the potential consequences of TIGAR expression on normal and tumor cells.

Keywords: Cancer metabolism, p53, TIGAR, PFK-2/FBPase-2

Review

Introduction

Cellular metabolism is a highly regulated process through which cell growth and survival are maintained. Nutrient availability promotes the production of biosynthetic compounds for cell growth and proliferation whereas starvation halts cell proliferation in order to conserve energy and assist in cell survival [1]. Many cancer cells shift their metabolism towards glycolysis - even under aerobic conditions - in order to provide a rapid production of energy and allow for the diversion of metabolic intermediates into anabolic pathways, as well as helping in the adaptation to challenging microenvironments. This increase in aerobic glycolysis is also known as the Warburg effect [2].

Alterations in metabolic pathways have been shown to play a role in tumorigenesis, with mutations as well as changes in the expression of metabolic enzymes contributing to metabolic transformation [3]. Activated oncogenes or the loss or inhibition of tumor suppressor proteins also directly influence cancer cell metabolism and cancer cell growth [4]. Oncogenic KRas, for example, enhances the flux of glycolytic intermediates to support anabolism [5] as well as influencing the transcriptional regulation of metabolic enzymes involved in glutamine metabolism [6]. Moreover, Ras-transformed cells can stimulate macropinocytosis in order to take up extracellular proteins and

provide amino acids for central carbon metabolism [7]. The transcription factor *MYC* directly activates genes involved in glucose metabolism [8,9] as well as those involved in glutamine metabolism such as glutaminase and glutamine transporters [10,11]. Hypoxia-inducible factor-1 (HIF-1), the major transcription factor involved in regulating the adaptation of cells to hypoxic conditions, also regulates the expression of many glycolytic genes [12] and can be activated in cancers even under normoxic (or pseudohypoxic) conditions in response to oncogenic signaling pathways or mutations in tumor suppressor proteins [13,14]. Cancers frequently show increased PI3K-Akt growth signaling and enhanced mammalian target of rapamycin (mTOR) activity. mTOR plays a central role in cellular metabolism by regulating growth-related processes such as protein synthesis, transcription and nutrient uptake, as well as autophagy, in response to changes in cellular nutrient and energy homeostasis. Many oncogenic events converge on the regulation of mTOR, including the loss of tumor suppressors such as PTEN [15], TSC1/TSC2 and LKB1 [16].

Oncogene activation, deregulated proliferation and altered metabolic activity in cancer cells can all generate increased levels of reactive oxygen species (ROS) [3,17]. While low levels of ROS can help to promote cell proliferation, oncogenic transformation promotes the production of excessive ROS, which would become toxic if not counteracted. Therefore, many cancer cells show an increased expression of antioxidant proteins such as Nrf2 [18,19],

* Correspondence: k.vousden@beatson.gla.ac.uk; e.cheung@beatson.gla.ac.uk
Cancer Research-UK Beatson Institute, Switchback Road, Glasgow G61 1BD, UK

which contribute to the survival and success of the tumor. Indeed, this dependence on antioxidants may make cancer cells more vulnerable to the inhibition of these detoxifying systems than normal cells, which do not carry such a high burden of oxidative stress [20-22].

p53 and cancer metabolism

The p53 tumor suppressor protein functions as a transcription factor and can initiate various cellular responses, including cell cycle arrest, senescence and apoptosis [23]. However, recent studies have suggested that none of these activities are essential to protect from cancer development [24], raising the possibility that other p53 functions are important for limiting tumorigenesis. Interest has now turned to the more recently described activities of p53 in regulating metabolism and allowing cells to adapt to and survive modest or transient periods of metabolic stress [25]. These survival activities of p53 have been linked with the promotion of catabolic pathways such as fatty acid oxidation and autophagy, which may provide alternative energy sources during starvation [26]. In addition, numerous activities of p53 that assist in limiting ROS and oxidative stress through the induction of target genes such as the tumor protein p53-induced nuclear protein 1 (TP53INP1) [27], glutaminase 2 (GLS2) [28,29], manganese superoxide dismutase (MnSOD) [30] and the sestrin family of proteins [31] also contribute to cell survival. It is not clear at present how, or even whether, these activities of p53 help prevent tumor development, although an ability to limit the accumulation of potentially oncogenic damage may be an important factor.

In contrast to its survival activity, the ability of p53 to induce senescence or cell death has been associated with an ability to induce oxidative stress. Several p53-inducible pro-oxidant genes have been described, and p53 can also limit the production of nicotinamide adenine dinucleotide phosphate (NADPH), which provides the major reducing power in cells in the form of reduced glutathione, by directly inhibiting the activity of glucose-6-phosphate dehydrogenase (G6PDH) [32] and repressing the expression of malic enzymes [33]. The anti- and pro-oxidant functions of p53 seem to mirror the ability to promote either survival or death - a complexity of the p53 response that is not fully understood. Current models suggest that these opposing functions of p53 reflect different roles in response to low or transient stress (where p53 protects cells and helps them survive and repair) and high or persistent stress (where p53 drives the elimination of the damaged cell) [34]. p53 is also activated by oxidative stress, resulting in protection from or exacerbation of damage through ROS, depending on the response. More recently, oxidative stress has been shown to drive the accumulation of p53 in the mitochondrial matrix, triggering the opening of the

mitochondrial permeability transition pore (PTP) through interaction with the PTP regulator cyclophilin D, leading to mitochondrial rupture and necrosis [35].

p53 therefore plays a complex but important role in the regulation of several metabolic pathways. Much like other cellular stress signals, metabolic stress can also activate p53. The activation of AMPK during low energy levels can lead to the induction of p53 activity [36] and PI3K-Akt growth signaling can inhibit p53 by activating MDM2 to promote the degradation of p53 [37]. While mTOR signaling inhibits p53 by promoting its dephosphorylation [38], a loss of the negative regulators of mTOR - and therefore, constitutive mTOR activity - can also promote p53 activity by enhancing translation [39]. Malate dehydrogenase has also been found to bind and activate p53 to mediate cell cycle arrest and apoptosis in response to glucose deprivation [40].

One important role of p53 that is beginning to emerge is its ability to help regulate the balance between glycolysis and oxidative phosphorylation. ATP and ADP can directly alter p53 activity, with ADP promoting and ATP inhibiting the ability of p53 to bind DNA [41]. p53 counteracts the elevation of glycolytic flux observed in cancer cells through inhibiting the expression of glucose transporters, GLUT1 and GLUT4 [42], as well as decreasing the levels of phosphoglycerate mutase 1 (PGAM1), the enzyme responsible for the conversion of 3-phosphoglycerate to 2-phosphoglycerate during glycolysis [43]. p53 can also promote oxidative phosphorylation through the activation of genes such as synthesis of cytochrome c oxidase 2 (SCO2) to increase mitochondrial respiration [44], as well as promote glutamine utilization through the activation of GLS2 [28,29]. Taken together, it would seem that p53 balances metabolic flux to allow for efficient energy production while blocking anabolic pathways necessary for cell growth. Indeed, loss of p53 has been suggested to be one of the mechanisms that contribute to the acquisition of the Warburg phenotype.

p53 also plays a role in preserving mitochondrial health with several activities likely to contribute to the maintenance of mitochondrial integrity. These include the induction of genes such as the ribonucleotide reductase subunit p53R2 [45-47], whose activity is required for the stability of mitochondrial DNA and the ability of p53 to contribute to the removal of damaged mitochondria [48,49]. While these results suggest that p53 helps to maintain mitochondrial quality, other studies have also demonstrated a role for p53 in the inhibition of mitophagy, an effect that would lead to increased mitochondrial dysfunction [50-52].

There have been many reviews of the role of p53 in regulating metabolic pathways, reflecting the complex interplay between p53-mediated responses that promote cell survival and those that induce cell death [53]. Here we

will focus on one aspect of the p53 response: the induction of TIGAR. Of note, it has recently become clear that the expression and activity of TIGAR can be uncoupled from the p53 response and the contribution of TIGAR to cancer development may depend on the manner by which it is regulated.

TIGAR: a fructose-2,6-bisphosphatase

TIGAR (*TP53*-induced glycolysis and apoptosis regulator) was discovered through microarray analysis of gene expression following the activation of p53 [54,55]. The human *TIGAR* gene is located on chromosome 12p13-3 and contains six coding exons and two p53 binding sites, one upstream of the first exon (BS1) and one within the first intron (BS2). Of the two sites, BS2 is much more efficient in binding p53. In the mouse genome, *Tigar* shows a similar genomic organisation but only possesses one p53 binding site, located upstream of the first exon. TIGAR is highly conserved through vertebrate species and shares similarities with the glycolytic enzyme phosphofructokinase-2/fructose-2,6-bisphosphatase (PFK-2/FBPase-2) [55].

PFK-2/FBPase-2 is a bifunctional protein containing a kinase domain within the NH₂-terminus and a bisphosphatase domain at the COOH-terminus. These two enzymatic activities are regulated through the formation of a dimer stabilized by interactions at the kinase domain [56]. Four different genes encode the PFK-2/FBPase-2 family of enzymes, *PFKFB1* to *PFKFB4*. While their catalytic domains are highly conserved, there are notable differences between different isoforms, including tissue specificity and preferential catalytic activity [57]. Moreover, cells have been shown to co-express different PFK-2/FBPase-2 isoforms, suggesting they each have distinct functions [58]. Both the expression and activity of PFK-2/FBPase-2 can be regulated by hormones and metabolites [59,60].

Notably, TIGAR only shares similarities with the bisphosphatase domain of PFK-2/FBPase-2 [55], with clear structural similarities despite limited amino acid conservation [55,61]. Thus, TIGAR, like FBPase-2, acts to degrade intracellular fructose-2,6-bisphosphate (F-2,6-P₂), which is a powerful allosteric activator of phosphofructokinase-1 (PFK-1). PFK-1 catalyses the conversion of fructose-6-phosphate (F-6-P) to fructose-1,6-bisphosphate (F-1,6-P₂) and in doing so, drives glycolysis. In addition, F-2,6-P₂ also acts as an inhibitor of fructose-1,6-bisphosphatase (FBP1) [62], which opposes the activity of PFK-1 by converting fructose-1,6-bisphosphate to fructose-6-phosphate.

By lowering F-2,6-P₂ levels, TIGAR decreases the activity of PFK-1 and reduces glycolytic flux downstream of this point. Several studies have shown that depletion of TIGAR results in increased levels of F-2,6-P₂ and increased flux through glycolysis [55,63,64], consistent with a model

in which the expression of TIGAR results in a dampening, rather than a complete inhibition, of the pathway. A number of consequences of TIGAR activity can therefore be predicted, including a diversion of the glycolytic metabolites to alternative metabolic fates, such as the hexosamine pathway to support glycosylation and the oxidative or non-oxidative branches of the pentose phosphate pathway (PPP) (Figure 1). The PPP plays a key role in generating ribose-5-phosphate to be used as an intermediate in nucleotide synthesis. Furthermore, the oxidative arm of the PPP allows for the production of NADPH, which supports antioxidant function and is required for anabolic pathways such as fatty acid synthesis (Figure 1).


The antioxidant activities of TIGAR

A dampening of glycolytic flux, either through the regulation of F-2,6-P₂ levels [65], glycosylation of PFK-1 [66], or ROS-induced inhibition of the M2 isoform of pyruvate kinase (PKM2) [67], has been shown to lead to an elevation of NADPH and antioxidant activity, which reflects an increase in the PPP. By analogy, the FBPase-2 activity of TIGAR should result in a similar response and a number of studies have shown that the downregulation of TIGAR is associated with decreased levels of NADPH [68-70], lower levels of reduced glutathione [55,69,71] and, consequently, an increase in ROS [72].

However, the antioxidant effect of TIGAR appears to reflect more than just its FBPase-2 activity. During hypoxia, a fraction of TIGAR was found to relocate to the mitochondria and associate with hexokinase 2 (HK2), resulting in enhanced HK2 activity, lower mitochondrial membrane potential and decreased ROS [73]. This activity displays some similarity to PFK-2/FBPase-2, where the FBPase-2 domain is able to bind and activate glucokinase (also known as hexokinase 4) [74,75]. Low oxygen availability can influence many cellular responses associated with tumor development, including angiogenesis and metastasis. In particular, hypoxia can regulate the metabolic activity of cells and induce glycolysis through the activation of HIF-1, which controls the expression of many metabolic enzymes, including *PFKFB3* and *PFKFB4* [76-78]. Notably, mutant TIGAR protein lacking FBPase-2 activity retains the ability to bind and enhance HK2 activity and the full antioxidant function of TIGAR under low-oxygen conditions depends on both HK2 binding and catalytic activity [73,79].

TIGAR under stress

The consequences of TIGAR expression on glycolysis and ROS regulation can depend, in part, on cell type and context. For example, cytokine-dependent lymphoid cells showed a decreased growth in response to TIGAR expression, possibly in response to decreased glycolysis [55],




and TIGAR was found to contribute to cell death in cardiac myocytes, an outcome that is also linked to a decrease in glycolysis [63]. However, in most cells where TIGAR functions to limit ROS, the effect of TIGAR expression was closely associated with protection against ROS-induced cell death [55,69-71,80]. More confusing is the association of TIGAR with senescence, where loss of TIGAR can induce senescence in glioblastoma cells [64] but can also inhibit this process in adult T-cell leukaemia cells [81].

A clearer understanding of the physiological role of TIGAR can be provided by the analysis of the role of TIGAR *in vivo* (Figure 2). Unlike many metabolic enzymes, which are essential for normal development [82,83], TIGAR deficient mice showed no profound developmental defect [79]. However, these mice have revealed

a role for TIGAR in the response to various forms of stress, such as cancer and heart failure.

Cardiac myocytes are known to undergo cell death following ischaemia-reperfusion injury, where greater tissue damage occurs due to the return of oxygenated blood following an ischaemic period, resulting in inflammation and oxidative stress. Both p53 and TIGAR protein expression are induced after myocardial infarction surgery, and both have been linked to an increase in apoptosis due to a decrease in glycolysis, resulting in decreased levels of phosphocreatine (a high energy phosphate important in tissues, such as muscles, with high fluctuating energy demands) [63]. In addition, the role of p53 and TIGAR following cardiac damage was also suggested to be due to their ability to inhibit autophagy, particularly in the form of mitophagy. p53- or TIGAR-deficient animals



were able to increase mitophagy after cardiac injury to reduce the number of damaged mitochondria and, hence, showed increased recovery in these tissues. In this case, the increase of ROS, due to the lack of TIGAR, functions as a signal to increase Bnip3 expression, resulting in an increase in mitophagy [51]. While a role for p53 in mediating adverse pathologies through the induction of cell death has been suggested in several diseases such as diabetes and ischaemia, protection due to a lack of TIGAR in this response is unanticipated [51,63].

More consistent with the antioxidant functions of TIGAR as protective for cell survival, as described *in vitro*, is the role of TIGAR in promoting recovery from stress-induced damage during intestinal tissue regeneration. Following ablation of the intestinal epithelium through whole body irradiation or genotoxic stress, mice deficient for TIGAR showed reduced regenerative capacity in their intestinal crypts [79]. Similarly, in a model of ulcerative colitis in the colon [84], mice that were deficient for TIGAR showed poorer recovery. As seen in cultured cells, a loss of TIGAR expression was accompanied by an increase in ROS. A lack of TIGAR compromised the ability of cells to undergo proliferation in order to regenerate the intestinal epithelium after ablation [79]. Further investigation using an *in vitro* intestinal crypt culture model [85] showed that organoids lacking TIGAR are less able to form crypt structures in a three-dimensional tissue culture model. These defects in *TIGAR*^{-/-} cells could be rescued following the addition of nucleosides or the antioxidant *N*-acetyl L-cysteine (NAC), suggesting that TIGAR acts to provide antioxidants and precursors for nucleic acid synthesis for intestinal growth [79].

TIGAR in cancer

The remodelling of metabolic pathways to help the control of redox homeostasis and provide intermediates needed for cell growth is of particular importance in tumor development. The identification of TIGAR as a p53 target gene indicates some role in tumor suppression, and the antioxidant functions of TIGAR would be consistent with a role in the protective p53 response to transient or repairable stress. Indeed, while TIGAR is induced during the early stages of a p53 response, a fall in TIGAR protein levels was shown to accompany the switch to apoptosis in cells under persistent p53-activating stress [55]. These results suggest that TIGAR levels must be tightly regulated during a p53 response, and there is now growing evidence that the deregulated expression of TIGAR may contribute to cancer development.

Studies on the PFK-2/FBPase-2 family have already revealed a role for these enzymes in tumor development. All *PFKFB* mRNAs have been reported to be overexpressed in human lung cancers [86] and PFKFB3, which has predominantly kinase activity, has been suggested to

promote tumorigenesis by enhancing PFK-1 activity and glycolytic flux [87]. Moreover, a recent study found a role for PFKFB3 in the proliferation of stalk endothelial cells and vessel sprouting by influencing the formation of filopodia/lamellipodia as well as cell migration. The loss of PFKFB3 in endothelial cells resulted in vascular defects *in vivo*, illustrating the importance of glycolysis in regulating vessel branching [88]. On the other hand, PFKFB4 plays an essential role in the survival of glioma stem-like cells and loss of PFKFB4 induced apoptosis in these cells [89]. Similarly in prostate cancer cells, loss of *PFKFB4* is detrimental to cell viability and resulted in a decrease in F-2,6-P₂ [65]. PFKFB4 shows predominantly bisphosphatase activity, leading to the suggestion that these cancer cells rely on PFKFB4 to dampen glycolytic flux, promote the PPP and manage ROS accumulation - very similar to the proposed action of TIGAR. However, this response to PFKFB4 expression may be more complicated than simply functioning to inhibit the PFK-1 step in glycolysis. While the inhibition of PFK-1 activity through glycosylation has been shown to promote the PPP and growth of cancer cells [66], loss of FBP1, whose activity directly opposes that of PFK-1 by converting F-1,6-P₂ to F-6-P, has also been observed in human liver, colon, gastric and breast cancers [90,91]. Interestingly, in this context, FBP1 expression is associated not only with decreased glycolysis and enhanced flux through the TCA cycle, but also with decreased PPP flux, and thereby an increase in ROS [92]. At first glance, these results seem contradictory to the model proposed for TIGAR and PFKFB4 expression, both of which also dampen glycolysis but appear to promote the PPP. While it is difficult to compare different models and tissue types in this way, these results may reflect the functions of TIGAR that are additional to the regulation of the PFK-1/FBP1 step of glycolysis. Most clearly, the ability of TIGAR to bind to and activate HK2 [73] could profoundly influence the availability of glucose intermediates for use in pathways such as the PPP. Consistently, HK2 was found to be important in maintaining tumor proliferation in a mouse model of *KRas*-driven lung cancer by promoting the PPP [93].

Given the activities of TIGAR in lowering ROS and promoting anabolic pathways, and the contribution of these pathways to cancer development, it is perhaps not surprising that overexpression of TIGAR has been described in a number of tumor types. Increase in TIGAR protein expression was observed in primary colon cancer and associated metastases [79], as well as in invasive breast cancer when compared to normal tissue [94]. Glioblastoma have been found to show a high expression of TIGAR compared to normal brain tissue [71,95], and knockdown of TIGAR resulted in radiosensitisation in glioma cells through an accumulation

of ROS, leading to DNA damage and cellular senescence [64]. Inhibition of transketolase-like 1, an enzyme involved in the PPP, was able to reverse the beneficial effects of TIGAR in these cells, further supporting the importance of the PPP in this response [71].

The role of TIGAR in balancing redox state in cancer cells has also been implicated in multiple myeloma cells, where inhibition of the oncoprotein MUC1-C resulted in a downregulation of TIGAR protein, lower levels of NADPH and in turn, increased ROS and cell death [69]. Moreover, in nasopharyngeal cancer cells, inhibition of c-Met, a tyrosine kinase whose overexpression has been associated with poor patient survival and metastasis [96], resulted in lower TIGAR expression, decreased NADPH and increased cell death [68].

In an intestinal adenoma model where APC is deleted in LGR5⁺ intestinal stem cells [97], mice deficient in TIGAR showed a reduction in total tumor burden and average tumor size in the small intestine compared to wild-type mice. TIGAR is also highly expressed in these adenomas when compared to the surrounding normal tissue, supporting the importance of TIGAR in proliferating tissue. A similar contribution of TIGAR to tumor progression was also observed in the colon, and importantly, the decrease in tumor burden observed in TIGAR-deficient mice correlated with a greater survival in these mice. *In vitro*, the defective growth of TIGAR-null tumor crypts could be rescued with antioxidants and nucleosides. The PPP has been shown to be of particular importance in redox homeostasis under hypoxic conditions, and TIGAR-deficient crypts were found to be more sensitive to hypoxia than wild-type crypts [79].

While the ability of TIGAR to promote cancer development might appear counterintuitive to its function in the p53 tumor suppressor pathway, it is important to note that in tumor cells overexpressing TIGAR, expression of TIGAR is uncoupled from p53 expression. Indeed, closer analysis in tumor cell lines showed that the basal expression of TIGAR is not dependent on the maintenance of wild-type p53 [79]. The ability of a p53-target protein to become oncogenic when no longer properly controlled has also been described for other mediators of the p53 survival response, such as carnitine palmitoyltransferase 1C [98]. Understanding how these genes are regulated will be critical in determining their role in cancer development.

The regulation of TIGAR

Initial studies identified TIGAR as a p53-responsive gene [51,54,55,63,71,81]. However, p53-independent expression of TIGAR has also been seen in several human cancer cell lines [79], and its expression in human breast cancer was inversely correlated to the expression of p53 [94]. Little is known about p53-independent regulation of TIGAR, which could be transcriptional, translational, through

the control of protein stability or through other post-translational modifications of the protein. Other members of the p53 family, p63 and p73, can activate promoters of several p53 target genes such as p21 and Bax [99,100], and could, therefore, also be capable of regulating TIGAR expression. While the TAp73 isoform has recently been found to be able to increase PPP activity through direct activation of G6PDH to support tumor cell proliferation [101], it is possible that the regulation of TIGAR also contributes to this response. In addition, mutant forms of p53 often display an oncogenic gain of function that can also involve modulation of tumor cell metabolism. While mutant p53s generally lose the ability to activate wild-type p53 target genes, they retain the ability to control transcription, such as the activation of genes involved in the mevalonate pathway in breast cancer cells [102]. As TIGAR expression is preserved in tumor cells that carry mutations in p53 [79], it is possible that some mutant p53s retain the ability to influence the expression of TIGAR and so help to promote tumorigenesis. Another transcription factor, SP1, has been found to regulate the basal level of TIGAR expression in liver cancer cell lines [103]. While *PFKFB3* expression can be induced by HIF-1 [76-78], TIGAR expression levels are not controlled by hypoxia. However, as discussed above, the activity of TIGAR is clearly modulated under conditions of low oxygen. Moreover, the loss of FBP1 observed in a number of human cancers and breast cancer cell lines was found to be due to promoter DNA methylation, demonstrating that epigenetic regulation also plays an important role in governing metabolism in cancer [90-92]. There is still much to be learnt about how TIGAR expression and activity are controlled under normal as well as stressed conditions.

Conclusions

As we gain further insight into the roles of TIGAR under normal and disease conditions, we can begin to make predictions about the benefit of modulating TIGAR for therapeutic intervention. *In vivo* studies have shown that the expression of TIGAR appears to be beneficial in certain circumstances, as seen in allowing for the recovery of intestinal epithelium following damage-induced ablation, but can also be detrimental, for example in promoting cardiac damage following ischaemic stress.

The situation seems somewhat clearer in cancer development, where overexpression of TIGAR is found in several tumor types and the deletion of TIGAR corresponds to a delay in cancer development. Indeed, using conditionally expressed TIGAR alleles, TIGAR loss was beneficial subsequent to tumor establishment, providing some indication that TIGAR may be a useful therapeutic target [71,79,94]. While these effects of TIGAR loss are consistent with the observation that inhibition of other antioxidants

can lead to excessive ROS and cell death in several cancer types [20-22], further investigation into TIGAR's activity, regulation, localisation and possible post-translational modifications are required to fully understand the role of TIGAR in the control of normal and disease pathologies.

Abbreviations

AMPK: AMP-activated protein kinase; APC: Adenomatous polyposis coli; Bnip3: BCL2/adenovirus E1B 19 kDa interacting protein 3; F-6-P: Fructose-6-phosphate; F-1,6-P₂: Fructose-1,6-bisphosphate; F-2,6-P₂: Fructose-2,6-bisphosphate; FBP1: Fructose-1,6-bisphosphatase; FBPase2: Fructose-2,6-bisphosphatase; G6PDH: Glucose-6-phosphate dehydrogenase; GLS2: Glutaminase 2; HIF-1: Hypoxia-inducible factor-1; HK2: Hexokinase 2; GLUT1: Glucose transporter 1; GLUT4: Glucose transporter 4; LGR5: Leucine-rich repeat containing G-protein coupled receptor 5; LKB1: Liver kinase B1; MDM2: Mouse double minute 2 homolog; MnSOD: Manganese superoxide dismutase; mTOR: Mammalian target of rapamycin; MUC1-C: Mucin 1 C-terminal subunit; NAC: N-acetyl L-cysteine; NADPH: Nicotinamide adenine dinucleotide phosphate; Nrf2: Nuclear factor (erythroid-derived 2)-like 2; PFK-1: Phosphofructokinase-1; PFK-2: Phosphofructokinase-2; PGAM1: Phosphoglycerate mutase 1; PI3K: Phosphatidylinositol 3-kinase; PKM2: Pyruvate kinase M2 isoform; PPP: Pentose phosphate pathway; PTEN: Phosphatase and tensin homolog; PTP: Permeability transition pore; ROS: Reactive oxygen species; SCO2: Synthesis of cytochrome c oxidase; SP1: Specificity protein 1; TCA: Tricarboxylic acid; TIGAR: TP53-induced glycolysis and apoptosis regulator; TP53INP1: Tumor protein p53-inducible nuclear protein 1; TSC1/TSC2: Tuberous sclerosis 1/tuberous sclerosis 2.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PL, KHV and ECC wrote the article. All authors read and approved the final manuscript.

Acknowledgements

We are grateful for funding from Cancer Research UK, ERC Grant 322842-METABOp53 and MRC.

Received: 20 September 2013 Accepted: 26 November 2013

Published: 3 January 2014

References

- Dang CV: Links between metabolism and cancer. *Genes Dev* 2012, **26**:877-890.
- Vander Heiden MG, Cantley LC, Thompson CB: Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 2009, **324**:1029-1033.
- Cairns RA, Harris IS, Mak TW: Regulation of cancer cell metabolism. *Nat Rev Cancer* 2011, **11**:85-95.
- Jones RG, Thompson CB: Tumor suppressors and cell metabolism: a recipe for cancer growth. *Genes Dev* 2009, **23**:537-548.
- Ying H, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sanankone E, Locasale JW, Son J, Zhang H, Colloff JL, Yan H, Wang W, Chen S, Viale A, Zheng H, Paik JH, Lim C, Guimaraes AR, Martin ES, Chang J, Hezel AF, Perry SR, Hu J, Gan B, Xiao Y, Asara JM, Weissleder R, Wang YA, Chin L, Cantley LC, et al: Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* 2012, **149**:656-670.
- Son J, Lyssiotis CA, Ying H, Wang X, Hua S, Ligorio M, Perera RM, Ferrone CR, Mullarky E, Shyh-Chang N, Kang Y, Fleming JB, Bardeesy N, Asara JM, Haigis MC, DePinho RA, Cantley LC, Kimmelman AC: Glutamine supports pancreatic cancer growth through a KRAS-regulated metabolic pathway. *Nature* 2013, **496**:101-105.
- Commisso C, Davidson SM, Soydaner-Azeloglu RG, Parker SJ, Kamphorst JJ, Hackett S, Grabocka E, Nofal M, Drebin JA, Thompson CB, Rabinowitz JD, Metallo CM: Vander Heiden MG, Bar-Sagi D: Macropinositosis of protein is an amino acid supply route in Ras-transformed cells. *Nature* 2013, **497**:633-637.
- Kim JW, Gao P, Liu YC, Semenza GL, Dang CV: Hypoxia-inducible factor 1 and dysregulated c-Myc cooperatively induce vascular endothelial growth factor and metabolic switches hexokinase 2 and pyruvate dehydrogenase kinase 1. *Mol Cell Biol* 2007, **27**:7381-7393.
- Osthus RC, Shim H, Kim S, Li Q, Reddy R, Mukherjee M, Xu Y, Wonsey D, Lee LA, Dang CV: Dereglulation of glucose transporter 1 and glycolytic gene expression by c-Myc. *J Biol Chem* 2000, **275**:21797-21800.
- Gao P, Tchernyshyov I, Chang TC, Lee YS, Kita K, Ochi T, Zeller KI, De Marzo AM, Van Eyk JE, Mendell JT, Dang CV: c-Myc suppression of miR-23a/b enhances mitochondrial glutaminase expression and glutamine metabolism. *Nature* 2009, **458**:762-765.
- Wise DR, DeBerardinis RJ, Mancuso A, Sayed N, Zhang XY, Pfeiffer HK, Nissim I, Daikhin E, Yudkoff M, McMahon SB, Thompson CB: Myc regulates a transcriptional program that stimulates mitochondrial glutaminolysis and leads to glutamine addiction. *Proc Natl Acad Sci U S A* 2008, **105**:18782-18787.
- Majmundar AJ, Wong WHJ, Simon MC: Hypoxia-inducible factors and the response to hypoxic stress. *Mol Cell* 2010, **40**:294-309.
- Maxwell PH, Pugh CW, Ratcliffe PJ: Activation of the HIF pathway in cancer. *Curr Opin Genet Dev* 2001, **11**:293-299.
- Semenza GL: HIF-1: upstream and downstream of cancer metabolism. *Curr Opin Genet Dev* 2010, **20**:51-56.
- Li J, Yen C, Liaw D, Podyspanina K, Bose S, Wang SI, Puc J, Milaresis C, Rodgers L, McCombie R, Bigner SH, Giovanella BC, Iltmann M, Tycko B, Hibshoosh H, Wigler MH, Parsons R: PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 1997, **275**:1943-1947.
- Guertin DA, Sabatini DM: Defining the role of mTOR in cancer. *Cancer Cell* 2007, **12**:9-22.
- Schumacker PT: Reactive oxygen species in cancer cells: Live by the sword, die by the sword. *Cancer Cell* 2006, **10**:175-176.
- DeNicola GM, Karreth FA, Humpton TJ, Gopinathan A, Wei C, Frese K, Mangal D, Yu KH, Yeo CJ, Calhoun ES, Scrimieri F, Winter JM, Hruban RH, Iacobuzio-Donahue C, Kern SE, Blair IA, Tuveson DA: Oncogene-induced Nrf2 transcription promotes ROS detoxification and tumorigenesis. *Nature* 2011, **475**:106-112.
- Mitsuishi Y, Taguchi K, Kawatani Y, Shibata T, Nukiwa T, Aburatani H, Yamamoto M, Motohashi H: Nrf2 redirects glucose and glutamine into anabolic pathways in metabolic reprogramming. *Cancer Cell* 2012, **22**:66-79.
- Raj L, Ide T, Gurkar AU, Foley M, Schenone M, Li X, Tolliday NJ, Golub TR, Carr SA, Shamji AF, Stern AM, Mandinova A, Schreiber SL, Lee SW: Selective killing of cancer cells by a small molecule targeting the stress response to ROS. *Nature* 2011, **475**:231-234.
- Ren DM, Villeneuve NF, Jiang T, Wu TD, Lau A, Toppin HA, Zhang DD: Brusatol enhances the efficacy of chemotherapy by inhibiting the Nrf2-mediated defense mechanism. *Proc Natl Acad Sci U S A* 2011, **108**:1433-1438.
- Trachootham D, Zhou Y, Zhang H, Demizu Y, Chen Z, Pelicano H, Chiao PJ, Achanta G, Arlinghaus RB, Liu J, Huang P: Selective killing of oncogenically transformed cells through a ROS-mediated mechanism by beta-phenylethyl isothiocyanate. *Cancer Cell* 2006, **10**:241-252.
- Vousden KH, Lu X: Live or let die: the cell's response to p53. *Nat Rev Cancer* 2002, **2**:594-604.
- Bieganski KT, Attardi LD: Deconstructing p53 transcriptional networks in tumor suppression. *Trends Cell Biol* 2012, **22**:97-106.
- Vogelstein B, Lane D, Levine AJ: Surfing the p53 network. *Nature* 2000, **408**:307-310.
- Vousden KH, Ryan KM: p53 and metabolism. *Nat Rev Cancer* 2009, **9**:691-700.
- Cano CE, Gommeaux J, Pietri S, Culcasi M, Garcia S, Seux M, Barelier S, Vasseur S, Spoto RP, Pébusque MJ, Dusetti NJ, Iovanna JL, Carrier A: Tumor protein 53-induced nuclear protein 1 is a major mediator of p53 antioxidant function. *Cancer Res* 2009, **69**:219-226.
- Hu W, Zhang C, Wu R, Sun Y, Levine A, Feng Z: Glutaminase 2, a novel p53 target gene regulating energy metabolism and antioxidant function. *Proc Natl Acad Sci U S A* 2010, **107**:7455-7460.
- Suzuki S, Tanaka T, Poyurovsky MV, Nagano H, Mayama T, Ohkubo S, Lokshin M, Hosokawa H, Nakayama T, Sugano S, Sato E, Nagao T, Yokote K, Tatsuno I, Prives C: Phosphate-activated glutaminase (GLS2), a p53-inducible regulator of glutamine metabolism and reactive oxygen species. *Proc Natl Acad Sci U S A* 2010, **107**:7461-7466.
- Lussain SP, Amstad P, He P, Robles A, Lupold S, Kaneko I, Ichimiya M, Sengupta S, Mechanic L, Okamura S, Hofseth LJ, Moake M, Nagashima M, Forrester KS, Harris CC: p53-induced up-regulation of MnSOD and GPx but not catalase increases oxidative stress and apoptosis. *Cancer Res* 2004, **64**:2350-2356.

31. Budanov AV, Karin M: **p53 target genes sestrin1 and sestrin2 connect genotoxic stress and mTOR signaling.** *Cell* 2008, **134**:451–460.
32. Jiang P, Du W, Wang X, Mancuso A, Gao X, Wu M, Yang X: **p53 regulates biosynthesis through direct inactivation of glucose-6-phosphate dehydrogenase.** *Nat Cell Biol* 2011, **13**:310–316.
33. Jiang P, Du W, Mancuso A, Wellen KE, Yang X: **Reciprocal regulation of p53 and malic enzymes modulates metabolism and senescence.** *Nature* 2013, **493**:689–693.
34. Gottlieb E, Vousden KH: **p53 regulation of metabolic pathways.** *Csh Perspect Biol* 2010, **2**:4.
35. Vaseva AV, Marchenko ND, Ji K, Tsirka SE, Holzmann S, Moll UM: **p53 opens the mitochondrial permeability transition pore to trigger necrosis.** *Cell* 2012, **149**:1536–1548.
36. Jones RG, Plas DR, Kubek S, Buzzai M, Mu J, Xu Y, Birnbaum MJ, Thompson CB: **AMP-activated protein kinase induces a p53-dependent metabolic checkpoint.** *Mol Cell* 2005, **18**:283–293.
37. Ogawara Y, Kishishita S, Obata T, Isazawa Y, Suzuki T, Tanaka K, Masuyama N, Gotoh Y: **Akt enhances Mdm2-mediated ubiquitination and degradation of p53.** *J Biol Chem* 2002, **277**:21843–21850.
38. Kong M, Fox CJ, Mu J, Solt L, Xu A, Cinali RM, Birnbaum MJ, Lindsten T, Thompson CB: **The PP2A-associated protein alpha4 is an essential inhibitor of apoptosis.** *Science* 2004, **306**:695–698.
39. Lee CH, Inoki K, Karbowniczek M, Petroulakis E, Sonenberg N, Henske EP, Guan KL: **Constitutive mTOR activation in TSC mutants sensitizes cells to energy starvation and genomic damage via p53.** *Embo J* 2007, **26**:4812–4823.
40. Lee SM, Kim JH, Cho EJ, Youn HD: **A nucleocytoplasmic malate dehydrogenase regulates p53 transcriptional activity in response to metabolic stress.** *Cell Death Differ* 2009, **16**:738–748.
41. Okorokov AL, Milner J: **An ATP/ADP-dependent molecular switch regulates the stability of p53-DNA complexes.** *Mol Cell Biol* 1999, **19**:7501–7510.
42. Schwartzenberg-Bar-Yoseph F, Armoni M, Karnieli E: **The tumor suppressor p53 down-regulates glucose transporters GLUT1 and GLUT4 gene expression.** *Cancer Res* 2004, **64**:2627–2633.
43. Kondoh H, Leonart ME, Gil J, Wang J, Degan P, Peters G, Martinez D, Carnero A, Beach D: **Glycolytic enzymes can modulate cellular life span.** *Cancer Res* 2005, **65**:177–185.
44. Matoba S, Kang JG, Patino WD, Wragg A, Boehm M, Gavrillova O, Hurley PJ, Bunz F, Hwang PM: **p53 regulates mitochondrial respiration.** *Science* 2006, **312**:1650–1653.
45. Tanaka H, Arakawa H, Yamaguchi T, Shiraishi K, Fukuda S, Matsui K, Takei Y, Nakamura Y: **A ribonucleotide reductase gene involved in a p53-dependent cell-cycle checkpoint for DNA damage.** *Nature* 2000, **404**:42–49.
46. Nakano K, Balint E, Ashcroft M, Vousden KH: **A ribonucleotide reductase gene is a transcriptional target of p53 and p73.** *Oncogene* 2000, **19**:4283–4289.
47. Bourdon A, Minai L, Serre V, Jais JP, Sarzi E, Aubert S, Chrétien D, de Lonlay P, Paquis-Flucklinger V, Arakawa H, Nakamura Y, Munnich A, Röttig A: **Mutation of RRM2B, encoding p53-controlled ribonucleotide reductase (p53R2), causes severe mitochondrial DNA depletion.** *Nat Genet* 2007, **39**:776–780.
48. Kitamura N, Nakamura Y, Miyamoto Y, Miyamoto T, Kabu K, Yoshida M, Futamura M, Ichinose S, Arakawa H: **Mieap, a p53-Inducible Protein. Controls Mitochondrial Quality by Repairing or Eliminating Unhealthy Mitochondria.** *PLoS One* 2011, **6**:1.
49. Miyamoto Y, Kitamura N, Nakamura Y, Futamura M, Miyamoto T, Yoshida M, Ono M, Ichinose S, Arakawa H: **Possible existence of lysosome-like organelle within mitochondria and its role in mitochondrial quality control.** *PLoS One* 2011, **6**:1.
50. Hoshino A, Mita Y, Okawa Y, Ariyoshi M, Iwai-Kanai E, Ueyama T, Ikeda K, Ogata T, Matoba S: **Cytosolic p53 inhibits Parkin-mediated mitophagy and promotes mitochondrial dysfunction in the mouse heart.** *Nat Commun* 2013, **4**:2308.
51. Hoshino A, Matoba S, Iwai-Kanai E, Nakamura H, Kimata M, Nakaoka M, Katamura M, Okawa Y, Ariyoshi M, Mita Y, Ikeda K, Ueyama T, Okigaki M, Matsubara H: **p53-TIGAR axis attenuates mitophagy to exacerbate cardiac damage after ischemia.** *J Mol Cell Cardiol* 2012, **52**:175–184.
52. Tasdemir E, Maiuri MC, Galluzzi L, Vitale I, Djavaheri-Mergny M, D'Amelio M, Criollo A, Morselli E, Zhu C, Harper F, Nannmark U, Samara C, Pinton P, Vicencio JM, Carnuccio R, Moll UM, Madeo F, Paterlini-Brechot P, Rizzuto R, Szabadkai G, Pierron G, Blomgren K, Tavernarakis N, Codogno P, Cecconi F, Kroemer G: **Regulation of autophagy by cytoplasmic p53.** *Nat Cell Biol* 2008, **10**:676–687.
53. Puzio-Kuter AM: **The role of p53 in metabolic regulation.** *Genes Cancer* 2011, **2**:385–391.
54. Jen KY, Cheung VG: **Identification of novel p53 target genes in ionizing radiation response.** *Cancer Res* 2005, **65**:7666–7673.
55. Bensaad K, Tsuruta A, Selak MA, Vidal MN, Nakano K, Bartrons R, Gottlieb E, Vousden KH: **TIGAR, a p53-inducible regulator of glycolysis and apoptosis.** *Cell* 2006, **126**:107–120.
56. Rider MH, Bertrand L, Vertommen D, Michels PA, Rousseau GG, Hue L: **6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase: head-to-head with a bifunctional enzyme that controls glycolysis.** *Biochem J* 2004, **381**:561–579.
57. Okar DA, Manzano A, Navarro-Sabate A, Riera L, Bartrons R, Lange AJ: **PFK-2/FBPase-2: maker and breaker of the essential biofactor fructose-2,6-bisphosphate.** *Trends Biochem Sci* 2001, **26**:30–35.
58. Minchenko O, Opentanova I, Caro J: **Hypoxic regulation of the 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene family (PFKFB-1-4) expression in vivo.** *FEBS Lett* 2003, **554**:264–270.
59. Bartrons R, Hue L, Van Schaftingen E, Hers HG: **Hormonal control of fructose 2,6-bisphosphate concentration in isolated rat hepatocytes.** *Biochem J* 1983, **214**:829–837.
60. Frenzel J, Schellenberger W, Eschrich K, Hofmann E: **Control of the fructose 6-phosphate/fructose 2,6-bisphosphate cycle by sn-glycerol 3-phosphate.** *Biomed Biochim Acta* 1988, **47**:461–470.
61. Li H, Jogl G: **Structural and biochemical studies of TIGAR (TP53-induced glycolysis and apoptosis regulator).** *J Biol Chem* 2009, **284**:1748–1754.
62. Okar DA, Lange AJ: **Fructose-2,6-bisphosphate and control of carbohydrate metabolism in eukaryotes.** *Biofactors* 1999, **10**:1–14.
63. Kimata M, Matoba S, Iwai-Kanai E, Nakamura H, Hoshino A, Nakaoka M, Katamura M, Okawa Y, Mita Y, Okigaki M, Ikeda K, Tsumumi T, Matsubara H: **p53 and TIGAR regulate cardiac myocyte energy homeostasis under hypoxic stress.** *Am J Physiol Heart Circ Physiol* 2010, **299**:H1908–H1916.
64. Pena-Rico MA, Calvo-Vidal MN, Villalonga-Planells R, Martinez-Soler F, Gimenez-Bonafe P, Navarro-Sabate A, Tortosa A, Bartrons R, Manzano A: **TP53 induced glycolysis and apoptosis regulator (TIGAR) knockdown results in radiosensitization of glioma cells.** *Radiother Oncol* 2011, **101**:132–139.
65. Ros S, Santos CR, Moco S, Baenke F, Kelly G, Howell M, Zamboni N, Schulze A: **Functional metabolic screen identifies 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 4 as an important regulator of prostate cancer cell survival.** *Cancer Discov* 2012, **2**:328–343.
66. Yi W, Clark PM, Mason DE, Keenan MC, Hill C, Goddard WA 3rd, Peters EC, Driggers EM, Hsieh-Wilson LC: **Phosphofructokinase 1 glycosylation regulates cell growth and metabolism.** *Science* 2012, **337**:975–980.
67. Anastasiou D, Pouligiannis G, Asara JM, Boxer MB, Jiang JK, Shen M, Bellinger G, Sasaki AT, Locasale JW, Auld DS, Thomas CJ, Vander Heiden MG, Cantley LC: **Inhibition of pyruvate kinase M2 by reactive oxygen species contributes to cellular antioxidant responses.** *Science* 2011, **334**:1278–1283.
68. Lui VW, Wong EY, Ho K, Ng PK, Lau CP, Tsui SK, Tsang CM, Tsao SW, Cheng SH, Ng MH, Ng YK, Lam EK, Hong B, Lo KW, Mok TS, Chan AT, Mills GB: **Inhibition of c-Met downregulates TIGAR expression and reduces NADPH production leading to cell death.** *Oncogene* 2011, **30**:1127–1134.
69. Yin L, Kosugi M, Kufe D: **Inhibition of the MUC1-C oncoprotein induces multiple myeloma cell death by down-regulating TIGAR expression and depleting NADPH.** *Blood* 2012, **119**:810–816.
70. Lui VW, Lau CP, Cheung CS, Ho K, Ng MH, Cheng SH, Hong B, Tsao SW, Tsang CM, Lei KI, Yamasaki Y, Mita A, Chan AT: **An RNA-directed nucleoside anti-metabolite, 1-(3-C-ethynyl-beta-d-ribo-pentofuranosyl)cytosine (ECyd), elicits antitumor effect via TP53-induced glycolysis and apoptosis regulator (TIGAR) downregulation.** *Biochem Pharmacol* 2010, **79**:1772–1780.
71. Wanka C, Steinbach JP, Rieger J: **TP53-induced glycolysis and apoptosis regulator (TIGAR) protects glioma cells from starvation-induced cell death by up-regulating respiration and improving cellular redox homeostasis.** *J Biol Chem* 2012, **287**:33436–33446.
72. Bensaad K, Cheung EC, Vousden KH: **Modulation of intracellular ROS levels by TIGAR controls autophagy.** *Embo J* 2009, **28**:3015–3026.
73. Cheung EC, Ludwig RL, Vousden KH: **Mitochondrial localization of TIGAR under hypoxia stimulates HK2 and lowers ROS and cell death.** *Proc Natl Acad Sci U S A* 2012, **109**:20491–20496.
74. Baltrusch S, Lenzen S, Okar DA, Lange AJ, Tiedge M: **Characterization of glucokinase-binding protein epitopes by a phage-displayed peptide library. Identification of 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase as a novel interaction partner.** *J Biol Chem* 2001, **276**:43915–43923.
75. Massa L, Baltrusch S, Okar DA, Lange AJ, Lenzen S, Tiedge M: **Interaction of 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase (PFK-2/FBPase-2)**

- with glucokinase activates glucose phosphorylation and glucose metabolism in insulin-producing cells. *Diabetes* 2004, **53**:1020–1029.
76. Obach M, Navarro-Sabate A, Caro J, Kong XG, Duran J, Gomez M, Perales JC, Ventura F, Rosa JL, Bartrons R: **6-phosphofructo-2-kinase (pfkfb3) gene promoter contains hypoxia-inducible factor-1 binding sites necessary for transactivation in response to hypoxia.** *J Biol Chem* 2004, **279**:53562–53570.
77. Minchenko O, Opentanova I, Minchenko D, Ogura T, Esumi H: **Hypoxia induces transcription of 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase-4 gene via hypoxia-inducible factor-1 alpha activation.** *FEBS Lett* 2004, **576**:14–20.
78. Bobarykina AY, Minchenko DO, Opentanova IL, Moenner M, Caro J, Esumi H, Minchenko OH: **Hypoxic regulation of PFKFB-3 and PFKFB-4 gene expression in gastric and pancreatic cancer cell lines and expression of PFKFB genes in gastric cancers.** *Acta Biochim Pol* 2006, **53**:789–799.
79. Cheung EC, Athineos D, Lee P, Ridgway RA, Lambie W, Nixon C, Strathdee D, Blyth K, Sansom OJ, Vousden KH: **TIGAR is required for efficient intestinal regeneration and tumorigenesis.** *Dev Cell* 2013, **25**:463–477.
80. Ye L, Zhao X, Lu J, Qian G, Zheng JC, Ge S: **Knockdown of TIGAR by RNA interference induces apoptosis and autophagy in HepG2 hepatocellular carcinoma cells.** *Biochem Biophys Res Commun* 2013, **437**:300–306.
81. Hasegawa H, Yamada Y, Iha H, Tsukasaki K, Nagai K, Atogami S, Sugahara K, Tsuruda K, Ishizaki A, Kamihira S: **Activation of p53 by Nutlin-3a, an antagonist of MDM2, induces apoptosis and cellular senescence in adult T-cell leukemia cells.** *Leukemia* 2009, **23**:2090–2101.
82. Piruat JJ, Pintado CO, Ortega-Saenz P, Roche M, Lopez-Barneo J: **The mitochondrial SDHD gene is required for early embryogenesis, and its partial deficiency results in persistent carotid body glomus cell activation with full responsiveness to hypoxia.** *Mol Cell Biol* 2004, **24**:10933–10940.
83. Yang H, Brosel S, Acin-Perez R, Slavkovich V, Nishino I, Khan R, Goldberg JJ, Graziano J, Manfredi G, Schon EA: **Analysis of mouse models of cytochrome c oxidase deficiency owing to mutations in Sco2.** *Hum Mol Genet* 2010, **19**:170–180.
84. Cooper HS, Murthy SN, Shah RS, Sedergran DJ: **Clinicopathologic study of dextran sulfate sodium experimental murine colitis.** *Lab Invest* 1993, **69**:238–249.
85. Sato T, Vries RG, Snippert HJ, van de Wetering M, Barker N, Stange DE, van Es JH, Abo A, Kujala P, Peters PJ, Clevers H: **Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche.** *Nature* 2009, **459**:262–265.
86. Minchenko OH, Ogura T, Opentanova IL, Minchenko DO, Ochiai A, Caro J, Komisarenko SV, Esumi H: **6-Phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene family overexpression in human lung tumor.** *Ukr Biokhim Zh* 2005, **77**:46–50.
87. Atsumi T, Chesney J, Metz C, Leng L, Donnelly S, Makita Z, Mitchell R, Bucala R: **High expression of inducible 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase (iPFK-2; PFKFB3) in human cancers.** *Cancer Res* 2002, **62**:5881–5887.
88. De Bock K, Georgiadou M, Schoors S, Kuchnio A, Wong BW, Cantelmo AR, Quaegebeur A, Ghesquière B, Cauwenberghs S, Eelen G, Phng LK, Betz I, Tembuyser B, Brepoels K, Welts J, Geudens I, Segura I, Cruys B, Bifari F, Decimo I, Blanco R, Wyns S, Vangindertael J, Rocha S, Collins RT, Munck S, Daelemans D, Imamura H, Devlieger R, Rider M, et al: **Role of PFKFB3-driven glycolysis in vessel sprouting.** *Cell* 2013, **154**:651–663.
89. Goidts V, Bageritz J, Puccio L, Nakata S, Zapata M, Barbus S, Toedt G, Campos B, Korshunov A, Momma S, Van Schaftingen E, Reifemberger G, Herold-Mende C, Lichter P, Radlwimmer B: **RNAi screening in glioma stem-like cells identifies PFKFB4 as a key molecule important for cancer cell survival.** *Oncogene* 2012, **31**:3235–3243.
90. Chen M, Zhang J, Li N, Qian Z, Zhu M, Li Q, Zheng J, Wang X, Shi G: **Promoter hypermethylation mediated downregulation of FBP1 in human hepatocellular carcinoma and colon cancer.** *PLoS One* 2011, **6**:e25564.
91. Bigl M, Jandrig B, Horn LC, Eschrich K: **Aberant methylation of human L- and M-fructose 1,6-bisphosphatase genes in cancer.** *Biochem Biophys Res Commun* 2008, **377**:720–724.
92. Dong C, Yuan T, Wu Y, Wang Y, Fan TW, Miriyala S, Lin Y, Yao J, Shi J, Kang T, Lorkiewicz P, St Clair D, Hung MC, Evers BM, Zhou BP: **Loss of FBP1 by snail-mediated repression provides metabolic advantages in basal-like breast cancer.** *Cancer Cell* 2013, **23**:316–331.
93. Patra KC, Wang Q, Bhaskar PT, Miller L, Wang Z, Wheaton W, Chandel N, Laakso M, Muller WJ, Allen EL, Jha AK, Smolen GA, Clasquin MF, Robey RB, Hay N: **Hexokinase 2 is required for tumor initiation and maintenance and its systemic deletion is therapeutic in mouse models of cancer.** *Cancer Cell* 2013, **24**:213–228.
94. Won KY, Lim SJ, Kim GY, Kim YW, Han SA, Song JY, Lee DK: **Regulatory role of p53 in cancer metabolism via SCO2 and TIGAR in human breast cancer.** *Hum Pathol* 2012, **43**:221–228.
95. Sinha S, Ghildiyal R, Mehta VS, Sen E: **ATM-NFkappaB axis-driven TIGAR regulates sensitivity of glioma cells to radiomimetics in the presence of TNFalpha.** *Cell Death Dis* 2013, **4**:e615.
96. Comoglio PM, Giordano S, Trusolino L: **Drug development of MET inhibitors: targeting oncogene addiction and expedience.** *Nat Rev Drug Discov* 2008, **7**:504–516.
97. Barker N, Ridgway RA, van Es JH, van de Wetering M, Begthel H, van den Born M, Danenberg E, Clarke AR, Sansom OJ, Clevers H: **Crypt stem cells as the cells-of-origin of intestinal cancer.** *Nature* 2009, **457**:608–611.
98. Zaugg K, Yao Y, Reilly PT, Kannan K, Kiarash R, Mason J, Huang P, Sawyer SK, Fuerth B, Faubert B, Kalliomäki T, Elia A, Luo X, Nadeem V, Bungard D, Yalavarthi S, Growney JD, Wakeham A, Moolani Y, Silvester J, Ten AY, Bakker W, Tsuchihara K, Berger SL, Hill RP, Jones RG, Tsao M, Robinson MO, Thompson CB, Pan G, et al: **Carnitine palmitoyltransferase 1C promotes cell survival and tumor growth under conditions of metabolic stress.** *Genes Dev* 2011, **25**:1041–1051.
99. Yang A, Kaghad M, Wang Y, Gillett E, Fleming MD, Dotsch V, Andrews NC, Caput D, McKeon F: **p63, a p53 homolog at 3q27–29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities.** *Mol Cell* 1998, **2**:305–316.
100. Lee CW, La Thangue NB: **Promoter specificity and stability control of the p53-related protein p73.** *Oncogene* 1999, **18**:4171–4181.
101. Du W, Jiang P, Mancuso A, Stonestrom A, Brewer MD, Minn AJ, Mak TW, Wu M, Yang X: **Tap73 enhances the pentose phosphate pathway and supports cell proliferation.** *Nat Cell Biol* 2013, **15**:991–1000.
102. Freed-Pastor WA, Mizuno H, Zhao X, Langerød A, Moon SH, Rodriguez-Barrueco R, Barsotti A, Chicas A, Li W, Polotskaia A, Bissell MJ, Osborne TF, Tian B, Lowe SW, Silva JM, Børresen-Dale AL, Levine AJ, Bargonetti J, Prives C: **Mutant p53 disrupts mammary tissue architecture via the mevalonate pathway.** *Cell* 2012, **148**:244–258.
103. Zou S, Gu Z, Ni P, Liu X, Wang J, Fan Q: **SP1 plays a pivotal role for basal activity of TIGAR promoter in liver cancer cell lines.** *Mol Cell Biochem* 2012, **359**:17–23.

doi:10.1186/2049-3002-2-1

Cite this article as: Lee et al.: TIGAR, TIGAR, burning bright. *Cancer & Metabolism* 2014 **2**:1.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

