

## Time-periodic Poiseuille flow in a pipe for some classes of fluids

Alessandra Borrelli · Maria Cristina Patria

Published online: 6 November 2007  
© Università degli Studi di Ferrara 2007

**Erratum to: Ann Univ Ferrara (2007) 53(1):13–28**  
**DOI 10.1007/s11565-007-0002-4**

The original version of this article unfortunately contained a mistake.

In page 19, lines 18–19, the sentence “Then the problem (2.1), (3.1), (3.2)” should be changed to “Then the problem (2.1) with  $\Gamma(t) = \Gamma(t)\mathbf{h}$  and  $\Phi(t) = \int_{\Sigma} \mathbf{h} \cdot \mathbf{V}(\mathbf{x}, t) d\Sigma$ ”

In page 20, line 7, equation “ $\mathbf{V} \equiv \mathbf{V}(\mathbf{x}) = \frac{\Phi}{a_{01}} \varphi_0(\mathbf{x})$ ” should be changed to  
“ $\mathbf{V} \equiv \mathbf{V}(\mathbf{x}) = \frac{\Phi}{\mathbf{h} \cdot \mathbf{a}_0} \varphi_0(\mathbf{x})$ ”.

---

The online version of the original article can be found under doi:10.1007/s11565-007-0002-4.

---

A. Borrelli (✉) · M. C. Patria  
Dipartimento di Matematica, Università di Ferrara, via N. Machiavelli 35,  
44100 Ferrara, Italy  
e-mail: brs@unife.it

M. C. Patria  
e-mail: pat@unife.it