

## **Erratum to: Time to flowering of temperate pulses in vivo and generation turnover in vivo–in vitro of narrow-leaf lupin accelerated by low red to far-red ratio and high intensity in the far-red region**

**Janine S. Croser<sup>1</sup> · Maria Pazos-Navarro<sup>1</sup> · Richard G. Bennett<sup>1</sup> · Sabrina Tschirren<sup>1</sup> · Kylie Edwards<sup>1</sup> · William Erskine<sup>1</sup> · Robert Creasy<sup>1</sup> · Federico M. Ribalta<sup>1</sup>**

Published online: 7 November 2016  
© Springer Science+Business Media Dordrecht 2016

### **Erratum to: Plant Cell Tiss Organ Cult DOI [10.1007/s11240-016-1092-4](https://doi.org/10.1007/s11240-016-1092-4)**

In the original publication, the name of the model/manufacture of the blue LED light used in Environments 1 and 2 is not correct. Environment 1 contained blue LED light from 18w-T8 LED tubes (model 108D18-V12; S-Tech Lighting, Australia) and Environment 2 contained a combination of blue LED light used in E1 plus far red-enriched LED light from AP67 Valoya L series (Helsinki, Finland).

---

The online version of the original article can be found under  
doi:[10.1007/s11240-016-1092-4](https://doi.org/10.1007/s11240-016-1092-4).

---

✉ Federico M. Ribalta  
[federico.ribalta@uwa.edu.au](mailto:federico.ribalta@uwa.edu.au)

<sup>1</sup> Centre for Plant Genetics and Breeding, The University of Western Australia, 35 Stirling Hwy, Crawley, WA 6009, Australia