

Available online on 15.03.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

To perform phytochemical screening and study the antioxidant potential of isolated compound from *Hemidesmus indicus*

Nutan*¹, Manas Kr Das², Gaurav Saxena³, Nitin Kumar¹¹ HIMT College of Pharmacy, Greater Noida (U P) India² Deptt of Pharmacy, IEC Group of Institutions, Greater Noida (U P) India³ GNIT College of Pharmacy, Greater Noida (U P) India

ABSTRACT

"Anantmul" is an important and widely used medicinal plant. The study aimed to determine the physicochemical composition, bioactive compounds and antioxidant activity of *Hemidesmus indicus* (Asclepiadaceae). The roots of the plant were collected and sequentially extracted using petroleum ether, ethyl acetate, and methanol. The preliminary phytochemical screening of extracts was carried out and found to be a good source of bioactive compounds. Biological activities of flavonoid and phenolic compounds have been discovered in several latest studies. Further phytochemical isolation was carried out, and Lupeol was isolated. Phenols are shown to be multifunctional antioxidants which will perform as singlet oxygen quenchers. Lupeol was evaluated for in-vitro antioxidant activity. It showed a correlation with antioxidant activity by DPPH ($IC_{50} = 0.52$, $P < 0.05$) and H_2O_2 ($IC_{50} = 0.43$, $P < 0.05$). The results show promising perspectives for the exploitation and use of anantmul rhizome as a constituent of anti-aging as well as anticancer diet.

Keywords: Anantmul, Lupeol, Antioxidant, DPPH**Article Info:** Received 28 Jan 2019; Review Completed 03 March 2019; Accepted 06 March 2019; Available online 15 March 2019**Cite this article as:**Nutan, Das MK, Saxena G, Kumar N, To perform phytochemical screening and study the antioxidant potential of isolated compound from *Hemidesmus indicus*, Journal of Drug Delivery and Therapeutics. 2019; 9(2):188-191
<http://dx.doi.org/10.22270/jddt.v9i2.2401>***Address for Correspondence:**

Nutan, HIMT College of Pharmacy, Greater Noida (U P) India

INTRODUCTION

Plants drugs are traditionally utilized in customary therapeutic frameworks; ethnomedicine, folk remedy, and herbalism give a rational and obvious source of contenders for focused identification of lead substances with unique structures, combinations, and mechanisms of action.

They moreover have the additionally preferred standpoint that, as a medicament, their safety and efficacy profiles are appropriately settled through old use or long time span human experience¹.

Hemidesmus indicus universally accepted as Indian sarsaparilla (figure 1, 2 and 3), belonging to family Asclepiadaceae. Its vernacular name "Anantmul" may be an Indic word which implies 'endless root'².

Figure 1 *Hemidesmus indicus* plant



Figure 2 *Hemidesmus indicus* flower



Figure 3 *Hemidesmus indicus* root

H. indicus may be a slender laticiferous, twig, typically prostrate or semi-erect woody plant, occurring in greater part of Indian Subcontinent. Anantmul is often distinguished by its slender, twisted, rigid, cylindrical and aromatic root and rhizome. Its bark is rust-colored and corky.

Furthermore, it has wrinkled with annulated cracks. Its stems are varied, slender, terete, thickened at the nodes. Leaves are opposite, variable, elliptic-oblong to linear-lanceolate, usually variegated with white above and pubescent beneath. Flowers are light-green outside and deep purple within huddled in subsessile axillary cymes. Follicles are slender, four inches long, cylindrical, typically arciform and divaricate. Its seeds are numerous, black planate. Phytoconstituents of *H. indicus* ranges from hydrocarbons, glycosides, oligoglycosides, and terpenoids to steroids³.

In folk medicine, the root of *Hemidesmus indicus* R. Br. is reported as aphrodisiac, antipyretic, anti-diarrheal, alleviates leprosy, leucoderma, skin diseases, and useful in piles. Further, it is also used as diuretic; in the joints- pain, syphilis, and leucoderma. The leaves are good for vomiting, cold, wounds, leucoderma- the stem has a bitter bad taste; diaphoretic, diuretic, laxative; lessens inflammation; good for diseases of the brain, the liver, the kidney; useful in syphilis, gleet and urinary discharges, uterine complaints, leucoderma, paralysis, cough, asthma; gargle good for toothache.²

H. indicus roots have been reported for many pharmacological actions, most notably antimicrobial activity⁴, antioxidant⁵, wound healing activity⁶,

antihyperglycemic, antidyslipidemic⁷, anti-arthritic activity⁸, Cytotoxic activity⁹ to cite a few.

This study aimed to perform the phytochemical screening and characterization of the isolated compound from *H. indicus* root, followed by the evaluation of its antioxidant activity.

MATERIALS AND METHODS

Collection and Authentication of Plant Material

Sample collection: Roots and rhizome powder of *Hemidesmus indicus* R.Br. (locally called Anantmul) were obtained and authenticated from NISCAIR-PUSA (Ref. no. NISCAIR/ RHMD/ consult/2013/2224/05).

Preparation of Anantmul Extracts

Dried powder of *H indicus* (100 gm) was exhaustively extracted with 500 ml petroleum ether and then with methanol in Soxhlet apparatus for 24 hours and dark brown residue (3.7 gm) was obtained after evaporation of the solvent. The dried extract (HIME) was stored in amber colored airtight container at 2.0°C temperature¹⁰.

Preliminary Phytochemical Study

For the identification of various phytochemical constituents, the different extracts were subjected to qualitative tests as per the standard procedure¹¹.

Isolation and purification of the compound.

A small quantity of HIME was dissolved in chloroform, and the solution was spotted on TLC plates. The plates were developed using several solvent systems; notably, Hexane / Chloroform (9:1) and Chloroform / Ethylacetate (5:1) gave better separation of the components and were used in the TLC monitoring of the Column Chromatography. 10g of the chloroform fraction (CF) was subjected to column chromatography on a silica gel (60 - 120 mesh) with gradient elution using Hexane and Chloroform^{12,13}

Spectroscopic Characterization

Different spectroscopic methods were used to elucidate the structure of HM₁, including IR, ¹H NMR, and ¹³C NMR techniques. The IR spectrum was recorded on a Jasco FTIR V 460 plus spectrometer using Diffuse Reflectance Attachment, ¹H NMR spectra were recorded on Varian Mercury YH 300 (300 MHz FT NMR), and ¹³C NMR spectra were recorded on a JEOL GSX 400 NB, 400 MHz FT NMR spectrometer in deuterated chloroform with TMS as an internal standard.

Antioxidant Activity Assessment

In-Vitro Antioxidant Activity conducted on *H indicus* extracts was DPPH (2, 2-diphenyl-picryl-hydrazine) test as per Silva¹⁴ and H₂O₂ assay as per Yang¹⁵ using ascorbic acid as standard. All the studies were carried out in triplicate.

RESULTS

Preliminary Phytochemical Analysis

The study results showed a spectrum of secondary metabolites (Table 1). It was also determined that extracts of *H indicus* contained a high concentration of secondary metabolites like Terpenoids, Saponins Flavonoids, Glycosides, Phytosterols, Tannins, all of which were reported to have antioxidant as well as cytotoxic properties.¹⁶

Table 1: Phytochemical Analysis of *Hemidesmus indicus*

S. No	Compounds	Analysis
1	Alkaloids	++
2	Carbohydrate	-
3	Fats & Oils	-
4	Flavonoids	++
5	Glycosides	++
6	Protein & amino acid	+
7	Phenols	+
8	Phytosterol	++
9	Resins	+
10	Saponin	+++
11	Tannins	++
12	Terpenoids	+++

Isolation and purification of the compound

A total of seventy collections, based on their TLC profiles, were made and pooled into half - dozen major fractions. Fraction four suggested a greater proportion of the interest compound and was further purified by preparatory TLC using the hexane/ethyl acetate solvent system (9:1). A uniform spot with two different Hexane / Ethylacetate solvent systems (9:1) and (5:1) was obtained on TLC. This compound, coded (HM₁), appeared as white needles and was subjected to spectral analysis.


Chemical Characterization result by various spectroscopy:

Different spectroscopic methods were used to elucidate the structure of HM₁. The Spectral data is presented in Table 2.

Table 2: Various Spectroscopy techniques result data

S. N.	Spectroscopy technique	Data
1	FTIR (CDCl ₃)	3139.59cm ⁻¹ (br, OH), 2945.31 cm ⁻¹ , 2872.60 cm ⁻¹ (C-H Str. In CH ₃ and CH ₂), 1637.98 cm ⁻¹ (C=C Str.) 1453.15 cm ⁻¹ (C-H deformation in CH ₂ /CH ₃), 1400.31 cm ⁻¹ (C-H deformation in gem dimethyl), 1043.20cm ⁻¹ (C-O Str. Of secondary alcohol), 880.06cm ⁻¹ (exocyclin CH ₂)
2	¹ HNMR (CDCl ₃)	δ 4.665(s,1H, H-29), δ 4.583(s,1H,H-27), δ 3.202(d,1H, H-3), δ 2.370(m,1H,H-19), δ 1.938(m,1H, H-21), δ 1.704(s,3H,H-30), δ 1.679(t,1H,H-13), δ 1.626(s,3H,H-2A), δ 1.542(s,3H,H-2B), δ 1.519(d,1H,H-11), δ 1.418(d,1H,H-14), δ 1.389(q,1H,H-6), δ 1.335(s,1H,H-21), δ 1.287(s,1H,H-9), δ 1.253(s,1H,H-9), δ 1.055(s,1H,H-23), δ 1.018(d,1H,H-15), δ 0.989(s,3H,H-23), δ 0.966(s,3H,H-27), δ 0.905(t,1H,H-18), δ 0.879(s,3H,H-25), δ 0.787(s,3H,H-28), δ 0.691(d,1H,H-5)
3	¹³ C NMR (CDCl ₃)	δ 151.178, δ 109.481, δ 79.135, δ 55.477, δ 48.493, δ 43.014, δ 41.031, δ 37.354, δ 34.463, δ 30.019, δ 27.626, δ 25.334, δ 21.108, δ 19.490, δ 14.760

These assignments are in good agreement for the structure of lupeol as per Jain¹⁷.

**Figure 4: Lupeol : pentacyclic tri-terpenoid****Antioxidant Activity Assessment**

Mostly because of the complex nature of phytochemicals, the antioxidant effects of plant products must be measured by incorporating two or more different in vitro assays to acquire satisfactory data. Each of these tests is based on one feature of the antioxidant activity, such as the ability to scavenge free radicals, or the metal ion chelation. The results are presented in Table 3 and 4. Overall, our study indicates that the high antioxidant properties of lupeol obtained from *H. indicus* root extract and may inhibit cellular lipid peroxidation and ameliorate other oxidative damage caused by free radicals¹⁸.

Table 3: Antioxidant activity of Lupeol by H₂O₂ Free Radical Scavenging Activity

S.No.	Dose µg/ml.	% Inhibitor	IC ₅₀
1	Lupeol 10	35.76±0.963	30 µg/ml
2	Lupeol 25	45.62±1.043	30µg/ml
3	Lupeol 50	74.76±1.873	30µg/ml

Table 4: Antioxidant activity of Lupeol by DPPH Radical Scavenging Activity.

S.No.	Dose µg/ml.	% Inhibitor	IC ₅₀
1	Lupeol 10	33.12±0.934	30 µg/ml
2	Lupeol 25	41.53±1.768	30 µg/ml
3	Lupeol 50	78.76±1.532	30 µg/ml

CONCLUSION

Phytochemical screening of the HIME showed the presence of triterpenoids, tannins, glycosides, flavonoids, polyphenols etc. Our findings strongly suggest that the Anantmul roots are promising sources of natural antioxidants, as indicated by their high contents of polyphenols, flavonoids, tannins, etc. All these classes of compounds have good antioxidant potential and their effects on human nutrition and health are significant. Also the considerable DPPH free radical-scavenging activities and H₂O₂ values of Lupeol further support these finding. Further studies may be carried out to evaluate its in-vivo anti cancer potential.

Conflict of Interest: None

REFERENCES

- 1 Predes FS, Ruiz AL, Carvalho JE, Foglio MA, Dolder H. Antioxidant and in vitro antiproliferative activity of *Arctium lappa* root extracts. BMC Complem Altern Med. 2011; 11(25).
- 2 Gupta NS. The Ayurvedic System of Indian Medicine, New Delhi, Bharatiya Kala Prakashan. 2006; 1:96-97
- 3 Sethi A, Srivastav SS, Srivastav S. Pregnane glycoside from *Hemidesmus indicus*. Indian J Heterocycl Chem 2006; 16:191-192.
- 4 Pandey KK, Dwivedi M, Urinary Tract Infection and its Management by Renalka. The Antiseptic. 2001; 98:295- 296.

- 5 Kumar S, Pooja M, Harika K, Haswitha E, Nagabhushanamma G, Vidyavathi N. In vitro Antioxidant Activities, Total Phenolics and Flavonoid Contents of whole plant of *Hemidesmus indicus*, Asian Journal of Pharmaceutical and Clinical Research. 2013; 6(2):249-251.
- 6 Kurupati VK, Nishteswar, K Phytochemical and clinical evaluation of Sariba (*Hemidesmus indicus*) on wound healing. International Research Journal of Pharmacy. 2012; 3(3):277-281.
- 7 Subramanian S, Abarna A, Thamizhiniyan V. Antihyperglycemic, Antioxidant and Antidyslipidemic properties *Hemidesmus indicus* R. Br. root extract studied in Alloxan -Induced Experimental Diabetes in Rats; IJPSR. 2012; 3(1):227-234
- 8 Mehta A, Sethiya NK, Mehta C, Shah GB. Anti-arthritis activity of roots of *Hemidesmus indicus* R.Br. (Anantmul) in rats, Asian Pacific Journal of Tropical Medicine. 2012, 130-135
- 9 Fimognari C, Lenzi M, Ferruzzi L, Turrini E, Scartezzini P et al., Mitochondrial Pathway Mediates the Antileukemic Effects of *Hemidesmus indicus*, a Promising Botanical Drug. Plos one. 2011; 6(6):e21544.
- 10 Nutan, Kumar, N , Saxena, Cytotoxic effect of *Hemidesmus indicus* R. Br. on HCT 116 human colon cell lines. 2019; 8(1):86-89.
- 11 Harborne JB. Phytochemical methods, London. Chapman and Hall, Ltd, 1973, 49-188.
- 12 Stahl E., Thin Layer Chromatography: A Laboratory Handbook, Second Edition, Springer, 1969, 206-258.
- 13 Chatterjee, I, Chakravarty , AK, Gomes, A. Daboia russellii and Naja kaouthia venom neutralization by lupeol acetate isolated from the root extract of Indian sarsaparilla *Hemidesmus indicus* R.Br. Journal of Ethnopharmacology 2006; 106:38-43.
- 14 Silva BM, Andrade PB, Valentão P, Ferreres F, Seabra RM, Ferreira MA. Quince, (*Cydonia oblonga* Miller) fruit (Pulp, Peel and Seed) and jam: Antioxidant activity. J Agr. Food Chem. 2004, 52:4705-4712.
- 15 Yang GJ, Yuan J. In vitro antioxidant properties of rutin, Lebensm. Wiss Technol. 2008; 41:1060-1066.
- 16 Dai J, Mumper RJ. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. Molecules. 2010; 15:7313-52..
- 17 Jain, P. S. and Bari, S. B., 'Isolation of Lupeol, Stigmasterol and Campesterol from Petroleum Ether Extract of Woody Stem-Bark of *Wightia tinctoria*', Asian Journal of Plant Sciences, 2010; 9(3):163-167
- 18 Paul , S., Hossen, M. S. , Tanvir, E. et al., "Antioxidant properties of *Citrus macroptera* fruit and its in vivo effects on the liver, kidney and pancreas in wistar rats," International Journal of Pharmacology, 2015; 11(8):899-909.

