

TOI-1338

Kostov, Veselin B.; Orosz, Jerome A.; Feinstein, Adina D.; Welsh, William F.; Cukier, Wolf; Haghhighipour, Nader; Quarles, Billy; Martin, David V.; Montet, Benjamin T.; Torres, Guillermo; Triaud, Amaury H. M. J.; Barclay, Thomas; Boyd, Patricia; Briceno, Cesar; Cameron, Andrew Collier; Correia, Alexandre C. M.; Gilbert, Emily A.; Gill, Samuel; Gillon, Michael; Haqq-Misra, Jacob

DOI:

[10.3847/1538-3881/ab8a48](https://doi.org/10.3847/1538-3881/ab8a48)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kostov, VB, Orosz, JA, Feinstein, AD, Welsh, WF, Cukier, W, Haghhighipour, N, Quarles, B, Martin, DV, Montet, BT, Torres, G, Triaud, AHMJ, Barclay, T, Boyd, P, Briceno, C, Cameron, AC, Correia, ACM, Gilbert, EA, Gill, S, Gillon, M, Haqq-Misra, J, Hellier, C, Dressing, C, Fabrycky, DC, Furesz, G, Jenkins, J, Kane, SR, Kopparapu, R, Hodzic, VK, Latham, DW, Law, N, Levine, AM, Li, G, Lintott, C, Lissauer, JJ, Mann, AW, Mazeh, T, Mardling, R, Maxted, PFL, Eisner, N, Pepe, F, Pepper, J, Pollacco, D, Quinn, SN, Quintana, EV, Rowe, JF, Ricker, G, Rose, ME, Seager, S, Santerne, A, Segransan, D, Short, DR, Smith, JC, Standing, MR, Tokovinin, A, Trifonov, T, Turner, O, Twicken, JD, Udry, S, Vanderspek, R, Winn, JN, Wolf, ET, Ziegler, C, Ansgore, P, Barnet, F, Bergeron, J, Hutten, M, Pappa, G & Straeten, TVD 2020, 'TOI-1338: TESS' first transiting circumbinary planet', *The Astronomical Journal*, vol. 159, no. 6, 253. <https://doi.org/10.3847/1538-3881/ab8a48>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

TOI-1338: TESS' First Transiting Circumbinary Planet

VESELIN B. KOSTOV,^{1,2} JEROME A. OROSZ,³ ADINA D. FEINSTEIN,^{4,*} WILLIAM F. WELSH,³ WOLF CUKIER,⁵
NADER HAGHIGHIPOUR,⁶ BILLY QUARLES,⁷ DAVID V. MARTIN,^{4,†} BENJAMIN T. MONTET,^{8,4} GUILLERMO TORRES,⁹
AMAURY H.M.J. TRIAUD,¹⁰ THOMAS BARCLAY,^{1,11} PATRICIA BOYD,¹ CESAR BRICENO,¹² ANDREW COLLIER CAMERON,¹³
ALEXANDRE C.M. CORREIA,¹⁴ EMILY A. GILBERT,^{4,15,16,17} SAMUEL GILL,¹⁸ MICHAËL GILLON,¹⁹ JACOB HAQQ-MISRA,²⁰
COEL HELLIER,²¹ COURTNEY DRESSING,²² DANIEL C. FABRYCKY,⁴ GABOR FURESZ,²³ JON JENKINS,²⁴ STEPHEN R. KANE,²⁵
RAVI KOPPARAPU,¹ VEDAD KUNOVAC HODŽIĆ,¹⁰ DAVID W. LATHAM,⁹ NICHOLAS LAW,²⁶ ALAN M. LEVINE,²³ GONGJIE LI,⁷
CHRIS LINTOTT,²⁷ JACK J. LISSAUER,²⁸ ANDREW W. MANN,²⁹ TSEVI MAZEH,³⁰ ROSEMARY MARDLING,³¹
PIERRE F.L. MAXTED,²¹ NORA EISNER,²⁷ FRANCESCO PEPE,³² JOSHUA PEPPER,^{33,23} DON POLLACCO,¹⁸
SAMUEL N. QUINN,³⁴ ELISA V. QUINTANA,¹ JASON F. ROWE,³⁵ GEORGE RICKER,²³ MARK E. ROSE,²⁸ S. SEAGER,^{23,36,37}
ALEXANDRE SANterne,³⁸ DAMIEN SÉGRANSAN,³² DONALD R. SHORT,³ JEFFREY C. SMITH,³⁹ MATTHEW R. STANDING,¹⁰
ANDREI TOKOVININ,¹² TRIFON TRIFONOV,⁴⁰ OLIVER TURNER,³² JOSEPH D. TWICKEN,^{2,24} STÉPHANE UDRY,³²
ROLAND VANDERSPEK,²³ JOSHUA N. WINN,⁴¹ ERIC T. WOLF,⁴² CARL ZIEGLER,⁴³ PETER ANSORGE,⁴⁴ FRANK BARNET,⁴⁴
JOEL BERGERON,⁴⁴ MARC HUTEN,⁴⁴ GIUSEPPE PAPPÀ,⁴⁴ AND TIMO VAN DER STRAETEN⁴⁴

¹NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771, USA

²SETI Institute, 189 Bernardo Ave, Suite 200, Mountain View, CA 94043, USA

³Department of Astronomy, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182, USA

⁴Department of Astronomy and Astrophysics, University of Chicago, 5640 S. Ellis Ave, Chicago, IL 60637, USA

⁵Scarsdale High School, 1057 Post Rd, Scarsdale, NY 10583, USA

⁶Institute for Astronomy, University of Hawaii-Manoa, Honolulu, HI 96822, USA

⁷Center for Relativistic Astrophysics, School of Physics, Georgia Institute of Technology, Atlanta, GA 30332, USA

⁸School of Physics, University of New South Wales, Sydney NSW 2052, Australia

⁹Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

¹⁰School of Physics & Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

¹¹University of Maryland, Baltimore County, 1000 Hilltop Cir, Baltimore, MD 21250, USA

¹²Cerro Tololo Inter-American Observatory, CTIO/AURA Inc., Casilla 603, La Serena, Chile

¹³Centre for Exoplanet Science, SUPA, School of Physics and Astronomy, University of St Andrews, St Andrews KY16 9SS, United Kingdom

¹⁴CFisUC, Department of Physics, University of Coimbra, 3004-516 Coimbra, Portugal

¹⁵The Adler Planetarium, 1300 South Lakeshore Drive, Chicago, IL 60605, USA

¹⁶NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

¹⁷GSFC Sellers Exoplanet Environments Collaboration

¹⁸Department of Physics, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, United Kingdom

¹⁹Astrobiology Research Unit, Université de Liège, 19C Allée du 6 Août, 4000 Liège, Belgium

²⁰Blue Marble Space Institute of Science, Seattle, Washington, USA

²¹Astrophysics Group, Keele University, Staffordshire, ST5 5BG, United Kingdom

²²Department of Astronomy, University of California at Berkeley Berkeley, CA 94720, USA

²³Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

²⁴NASA Ames Research Center, Moffett Field, CA, 94035, USA

²⁵Department of Earth and Planetary Sciences, University of California, Riverside, CA 92521, USA

²⁶Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3255, USA

²⁷Oxford University, United Kingdom

²⁸NASA Ames Research Center, Moffett Field, CA 94035, USA

²⁹Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

³⁰Department of Astronomy and Astrophysics, Tel Aviv University, 69978 Tel Aviv, Israel

³¹School of Physics & Astronomy, Monash University, Victoria, 3800, Australia

³²Observatoire Astronomique de l'Université de Genève, 51 chemin des maillettes, CH-1290 Sauverny, Switzerland

³³Department of Physics, Lehigh University, 16 Memorial Drive East, Bethlehem, PA 18015, USA

³⁴*Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA*

³⁵*Bishops University, 2600 College St, Sherbrooke, QC J1M 1Z7, Canada*

³⁶*Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

³⁷*Department of Aeronautics and Astronautics, MIT, 77 Massachusetts Avenue, Cambridge, MA 02139, USA*

³⁸*Aix Marseille Univ, CNRS, CNES, LAM, Marseille*

³⁹*SETI Institute/NASA Ames Research Center, 189 Bernardo Ave, Suite 200, Mountain View, CA, 94043, US*

⁴⁰*Max Planck Institut fuer Astronomie, Heidelberg, Germany*

⁴¹*Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA*

⁴²*Department of Atmospheric and Oceanic Sciences, Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, Colorado, USA, USA*

⁴³*Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, Ontario M5S 3H4, Canada*

⁴⁴*Planet Hunters TESS*

(Accepted AJ)

ABSTRACT

We report the detection of the first circumbinary planet found by TESS. The target, a known eclipsing binary, was observed in sectors 1 through 12 at 30-minute cadence and in sectors 4 through 12 at two-minute cadence. It consists of two stars with masses of $1.1 M_{\odot}$ and $0.3 M_{\odot}$ on a slightly eccentric (0.16), 14.6-day orbit, producing prominent primary eclipses and shallow secondary eclipses. The planet has a radius of $\sim 6.9 R_{\oplus}$ and was observed to make three transits across the primary star of roughly equal depths ($\sim 0.2\%$) but different durations—a common signature of transiting circumbinary planets. Its orbit is nearly circular ($e \approx 0.09$) with an orbital period of 95.2 days. The orbital planes of the binary and the planet are aligned to within $\sim 1^{\circ}$. To obtain a complete solution for the system, we combined the TESS photometry with existing ground-based radial-velocity observations in a numerical photometric-dynamical model. The system demonstrates the discovery potential of TESS for circumbinary planets, and provides further understanding of the formation and evolution of planets orbiting close binary stars.

Keywords: Exoplanet Astronomy (486), Eclipsing Binary Stars (444)

1. INTRODUCTION

One of the most exciting breakthroughs from the *Kepler* mission was the discovery of circumbinary planets (CBPs). Four years of continuous observations of several thousand eclipsing binary stars (EBs, Prša et al. 2011; Slawson et al. 2011; Kirk et al. 2016), led to the discovery of 13 transiting CBPs orbiting 11 *Kepler* EBs (Doyle et al. 2011; Welsh et al. 2012; Orosz et al. 2012a,b; Schwamb et al. 2013; Kostov et al. 2013, 2014; Welsh et al. 2015; Kostov et al. 2016; Orosz et al. 2019; Socia et al. 2020). These discoveries spanned a number of firsts—e.g. the first transiting CBP, the first CBP in the Habitable Zone (HZ), the first CBP in a quadruple star system, and the first transiting multiplanet CBP system. In addition to opening a new chapter in studies of extrasolar planets, Kepler’s CBPs have confirmed theoretical predictions that planet formation in circumbinary configurations is a robust process, and suggest that many such planetary systems must exist (e.g. Pierens & Nelson 2013; Kley & Haghighipour 2015). Indeed, recent studies argue that the occurrence rate of giant, Kepler-like CBPs is comparable to that of giant planets in single-star systems ($\sim 10\%$; Armstrong et al. 2014; Martin & Triaud 2014; Li et al. 2016; Martin et al. 2019).

As exciting as the CBPs discovered from Kepler are, however, the present sample is small, likely hindered by observational biases, and leaves a vast gap in our understanding of this new class of worlds. This is not unlike the state of exoplanet science 20 years ago, when only a handful of hot Jupiter exoplanets were known. Pressing questions remain regarding the formation and migration efficiency of CBPs, their orbital architectures and occurrence rates, the formation, evolution and population characteristics of their host binary stars. Addressing these questions requires more CBP discoveries—which require continuous observations of a large number of EBs for prolonged periods of time. NASA’s Transiting Exoplanet Survey Satellite (TESS, Ricker et al. 2015) will assist CBP discovery by observing

* NSF Graduate Research Fellow

† Fellow of the Swiss National Science Foundation

roughly half a million EBs continuously for timespans between one month and one year for the nominal mission (Sullivan et al. 2015). This motivated us to continue our search for transiting CBPs by examining light curves of EBs observed by TESS.

Here we report the discovery of the first CBP from TESS—TOI-1338, a Saturn-size planet orbiting the known eclipsing binary star EBLM J0608-59¹ approximately every 95 days. At the time of this writing, this is the longest-period confirmed planet discovered by TESS. Below we present the details of our discovery, and discuss some of the characteristics of this newly-found CBP that allow us to place its discovery in a broader context.

This paper is organized as follows. In Section 2 we briefly outline the discovery of the system, and describe the TESS data for the target star and the detection of the CBP transits. In Section 3 we present the complementary observations and data analysis; Section 4 outlines the photometric-dynamical analysis of the system. Section 5 presents a discussion of the results. We draw our conclusions in Section 6.

2. DISCOVERY

2.1. TESS Mission

The primary goal of the TESS mission is to identify transiting planets around nearby bright stars that are amenable to follow-up characterization. TESS will observe about 85% of the sky during its two-year primary mission (Ricker et al. 2015), using four cameras that provide a $24^\circ \times 96^\circ$ field-of-view (FOV); a sector is a \sim month-long observation of a single FOV. Most of the stars in the Full-Frame Images (FFIs) will be observed at 30-minute cadence, and 200,000 pre-selected stars (spread over the whole sky) will be observed at 2-min cadence. TESS observes 13 sectors per hemisphere, per year for at least ≈ 27 days; where the sectors overlap near the ecliptic poles, in the two Continuous Viewing Zones (CVZs), TESS observes for up to ≈ 350 days. The CVZs are especially valuable places to search for exoplanets as the longer baseline enables the detection of smaller and/or longer-period planets—like the CBP presented here—and also overlaps with the James Webb Space Telescope CVZ (Ricker et al. 2015). Half way through its primary mission, TESS has already discovered a number of confirmed planets and identified more than a thousand planet candidates (e.g. Huang et al. 2018; Vanderspek et al. 2019; Kostov et al. 2019, and references therein), vetted by the TESS Data Validation initiative (Twicken et al. 2018; Li et al. 2019) (Guerrero et al., in prep.)

2.2. Discovery of the Host Eclipsing Binary

TOI-1338 was identified as an eclipsing binary in 2009 as a part of the EBLM (Eclipsing Binary Low Mass; Triaud et al. 2013) project, a survey constructed using the false-positives of the WASP survey for transiting hot-Jupiters (Pollacco et al. 2006; Collier Cameron et al. 2007; Triaud 2011). As the observed eclipse depth is comparable to that of an inflated hot Jupiter, at the time it was not possible to distinguish between an eclipsing late-type M-dwarf and a transiting hot-Jupiter from the photometric signature alone. Follow-up observations with the CORALIE² high-resolution spectrograph obtained near quadrature revealed a semi-amplitude of 21.6 km s^{-1} , confirming that the target is a low-mass eclipsing binary (Triaud et al. 2017a). As part of the EBLM project to improve the M-dwarf mass-radius-temperature-luminosity relation, TESS observations at two-minute cadence were obtained for the target under a Cycle 1 Guest Observer program for Sectors 4-12 (G011278—PI O. Turner).

2.3. Detection of the Circumbinary planet

Based on the two-minute cadence observations from TESS, the target was flagged as an eclipsing binary on the Planet Hunters TESS platform (Eisner et al. 2020)³ by user Pappa on subject 31326051. This is a citizen science project where, in addition to primarily flagging transit-like features, volunteers may flag targets as various phenomena including eclipsing binaries, variable stars, etc. Planet Hunters has had successful contribution to the field of CBPs through the independent discovery of Kepler-64 (also known as Planet Hunters-1, Kostov et al. 2013; Schwamb et al. 2013).

As part of our search for CBPs around eclipsing binaries from TESS, we have been performing a visual inspection of targets tagged as potential eclipsing binaries (with a hashtag “eclipsingbinary”) on the Planet Hunters TESS Talk.

¹ The target also has the designations TIC 260128333, TYC 8533-00950-1, Gaia DR2 5494443978353833088. Its Right Ascension and Declination are 06:08:31.97 and -59:32:28.08, respectively. It has TESS magnitude $T = 11.45 \pm 0.02$ mag, and $V = 11.72 \pm 0.02$ mag.

² CORALIE is a fibre-fed échelle spectrograph, mounted on the Swiss Euler 1.2 metre telescope at La Silla, Chile. It has a resolving power of $R = 55,000$, and achieves long-term stability through a thermally-stabilised housing and nightly calibrations with respect to a Thorium-Argon reference spectrum (Lovis & Pepe 2007).

³ <https://www.zooniverse.org/projects/nora-dot-eisner/planet-hunters-tess>

We note that TOI-1338 was also listed on exo.MAST⁴ as two Threshold Crossing Events with a period of ~ 14.61 days.

During the examination of the light curves on Planet Hunters TESS Talk, one of us (WC) noticed that in one of the partial light curves of TOI-1338 there was both a prominent primary eclipse and an additional feature. The latter was of similar depth to the secondary eclipse, but not at the expected time according to exo.MAST. To confirm that the feature is real and not a false positive caused by instrumental artifacts, we next extracted the 2-min cadence light curve using the `Lightkurve` software package (Lightkurve Collaboration et al. 2018) and confirmed that there are indeed two genuine transit-like features in Sectors 6 and 10 that are not associated with the secondary eclipses, separated by ≈ 95 days, and with different durations (~ 0.3 and ~ 0.6 days respectively). Further analysis of the 30-min cadence data, extracted with `eleanor` (Feinstein et al. 2019), revealed the presence of a third transit in Sector 3, ≈ 93 days before the transit in Sector 6, with a duration of ~ 0.4 days, further strengthening the CBP interpretation. Overall, the three transits exhibit the trademark “smoking gun” signatures of transiting CBPs (Welsh & Orosz 2018) where (i) the transit durations vary depending on the orbital phase of the host EB such that transits across the primary star occurring near primary eclipses have shorter durations than transits occurring near secondary eclipses, and (ii) the transit times vary significantly from a linear ephemeris where specifically in this case the interval between the first and second transits (~ 93 days) is significantly different than the interval between the second and third transits (~ 95 days).

We note that it is highly unlikely for the three CBP transits to be a false positive scenario due, for example, by an unresolved eclipsing binary star. First, because the three transits have different durations, there would be only two plausible scenarios. One scenario, S1, involves two unresolved eclipsing binaries where one (hereafter EB1) would produce the first and third transits as primary and secondary eclipses, and the other (hereafter EB2) produces only the second transit as either a primary or a secondary eclipse. The other scenario, S2, requires an unresolved triple star system consisting of EB1 (producing the first and the third transits as a primary and secondary eclipse) and a long-period third star (hereafter EB3) producing the second transit as an eclipse across either the primary or the secondary star of EB1. For circular orbits, EB1 needs to have an orbital period of about 400 days (twice the time between the first and third transit) regardless of the scenario, EB2 for scenario S1 should have an orbital period greater than about 200 days (so that it would not produce a second eclipse during the TESS observations), and EB3 would need to have a dynamically-stable orbit around the ~ 400 -day EB1. Second, because the three CBP transits have a depth of $\sim 0.2\%$, such background EBs cannot be fainter than $T \sim 18$ mag (i.e. $\Delta T \sim 7$ mag difference compared to TOI-1338). We note that spectroscopy observations of the target do not show any signs of a second or third EB, albeit not as strong as 7 magnitudes difference. Below we explore this further using rough approximations to estimate the order of magnitude of the probability.

While the TESS Input Catalog indicates that there are 98 contaminating sources for TOI-1338, Gaia shows that there are only 6 sources within $\Delta T \sim 7$ mag⁵ inside the entire 13×13 TESS pixel array of the target, although none of them is inside the TESS aperture of the target. Assuming that these 6 sources are representative of the field of view, the density of sources within $\Delta T \sim 7$ mag of the target is then ~ 0.036 sources/pixel, i.e. $\sim 10^{-4}$ sources/sq. arcsec. The contrast sensitivity of Gaia DR2 for $\Delta G \sim 7$ mag (and thus $\Delta T \sim 7$ mag) is ~ 3 arcsec (Brandeker & Cataldi 2019). Thus the probability of having one $\Delta T \sim 7$ mag source unresolved by Gaia within 9 sq. arcsec of the target is $P_{\text{unresolved}} \sim 10^{-3}$.

Using the results of Raghavan et al. (2010), we estimate that the probability of EB1 having an orbital period of 400 days is $P_{\text{EB1,per}} \sim 20\%$; the probability of EB2 having an orbital period of 200 days is $P_{\text{EB2,per}} \sim 85\%$. Using the results of Tokovinin (2014), we estimate the probability of a triple star to be $P_{\text{triple}} \sim 10\%$. The probability that EB1, EB2, and EB3 are eclipsing is roughly $(R_1 + R_2)/a$. Assuming similar stars (because of equal depth eclipses) of solar-radius and mass (because larger stars are more rare and smaller stars would be too faint to produce the required contamination) for both scenarios, the corresponding probabilities are $P_{\text{EB1,ecl}} \sim (2 \times 0.0046)/1.34 \sim 0.006$ and $P_{\text{EB2,ecl}} \sim (2 \times 0.0046)/0.84 \sim 0.011$. For scenario S2, we used Rebound’s IAS15 integrator (Rein & Spiegel 2015) to test that the orbital period of EB3 would need to be greater than ~ 2000 days to be dynamically stable. Thus the probability that EB3 is eclipsing EB1 is $P_{\text{triple,ecl}} \sim (2 \times 0.0046)/4.48 \sim 0.002$.

⁴ <https://exo.mast.stsci.edu>

⁵ Gaia magnitude is similar to TESS magnitude, i.e. $G = T + 0.43$ (Stassun et al. 2019)

We also note that the orbital phases of EB1, EB2, and EB3 would need to be such that in ~ 325 days (duration of TESS observations) EB1 would produce one primary and one secondary eclipse, and EB2 and EB3 would produce a single eclipse—within a window of a few days for its duration to agree with the CBP model. This introduces additional constraints. Namely, for orbital periods of 400-days, 200-days, and 2000-days for EB1, EB2, and EB3 respective, and assuming said window is ~ 2 days, the corresponding probabilities are $P_{\text{EB1,phase}} \sim 80\%$ (i.e. $\sim 325/400$), $P_{\text{EB2,phase}} \sim 1\%$ (i.e. $\sim 2/200$), and $P_{\text{EB3,phase}} \sim 0.1\%$ (i.e. $\sim 2/2000$).

Putting all this together, the combined probability for scenarios S1 and S2 is as follows:

$$P_{\text{S1}} = P_{\text{unresolved,EB1}} \times P_{\text{unresolved,EB2}} \times P_{\text{EB1,per}} \times P_{\text{EB2,per}} \times P_{\text{EB1,ecl}} \times P_{\text{EB2,ecl}} \times P_{\text{EB1,phase}} \times P_{\text{EB2,phase}} = \quad (1)$$

$$= 10^{-3} \times 10^{-3} \times 0.2 \times 0.85 \times 0.006 \times 0.011 \times 0.8 \times 0.01 \sim 10^{-13} \quad (2)$$

$$P_{\text{S2}} = P_{\text{unresolved}} \times P_{\text{triple}} \times P_{\text{EB1,per}} \times P_{\text{EB1,ecl}} \times P_{\text{EB3,ecl}} \times P_{\text{EB1,phase}} \times P_{\text{EB3,phase}} = \quad (3)$$

$$= 10^{-3} \times 0.1 \times 0.2 \times 0.006 \times 0.002 \times 0.8 \times 0.001 \sim 2 \times 10^{-13} \quad (4)$$

To convert these numbers to a false positive probability (FPP), we compare them to the probability that TOI-1338 b is a circumbinary planet. Specifically, from the Kepler dataset the probability that any given star has a transiting CBP is $\sim 10/200,000 = 5 \times 10^{-5}$. Thus the CBP hypothesis is $\sim 5 \times 10^{-5}/10^{-13} = 5 \times 10^{-8}$ times more likely than the false positive hypothesis. The FPP is therefore about 10^{-8} .

These false positive scenarios can be further argued against based on the durations and depths of the three transits. Specifically, as the depths of the first and third transits are similar, then the two stars of EB1 should have comparable sizes. As discussed above, assuming Sun-like stars the semi-major axis of EB1 would be ~ 1.34 AU and the corresponding duration of the primary eclipse of EB1 (for circular orbit, $R_1 = R_2 = R_{\odot}$, and impact parameter $b = 0$) would be ~ 0.9 days, i.e. nearly twice the duration of the observed transits, thus further ruling out this specific scenario. While an eccentric orbit of EB1 might alleviate this tension to some degree, this would require special orbital elements – in addition to the requirement imposed by the special orbital phase as discussed above. Thus overall we consider false positive scenarios S1 and S2 to be highly unlikely.

2.4. TESS Lightcurve

TESS telemeters data in two modes: postage stamps, i.e. small regions, around roughly 20,000 stars at 2-minute cadence every sector⁶ as well as the Full-Frame Images, which contain about a million stars each (brighter than $T = 15$ mag), at 30-minute cadence. TOI-1338 was observed by TESS in 30-min cadence in Sectors 1-12, and in 2-min cadence in Sectors 4-12.

The TESS light curve of TOI-1338 is shown in Figure 1, where the 30-min cadence data extracted with `eleanor` using aperture photometry is shown for Sectors 1-12 and 2-min cadence SAPFLUX measurements from the standard processing provided by the TESS mission is shown for Sectors 4-12. The nominal units of the observation times are days from BJD 2,457,000. The `eleanor` software package performs background subtraction, aperture photometry, and detrending for a given source on the FFIs. It also provides the opportunity to use a custom aperture and if desired can use models of the point spread function (PSF) to extract the light curves.

Each full-frame image delivered by the TESS project is barycentric corrected. However, there is only a single correction applied to each CCD, each of which covers more than 100 square degrees of the sky. As a result, the barycentric-corrected times in the raw FFI data can be discrepant by up to a minute. The `eleanor` software corrects for this potential offset, removing the barycentric correction and applying a more accurate value given the actual position of the target in question. We verified the `eleanor` timestamps were accurate, comparing them to the midpoint of the 15 2-minute images that make up a single FFI. We find the two are consistent at the 2-second level, sufficiently precise for the photodynamical modeling we employ in Section 4.

⁶ For sectors 1-3 slightly less than 16,000 targets received 2-min cadence observations. The number of 2-min cadence targets was increased to 20,000 after predicted compressibility was demonstrated in flight and the SPOC demonstrated in a ground segment test that 20,000 targets could be handled in the pipeline (Jenkins et al. 2016).




Figure 1. Upper panel: Sectors 1-12 of 30-min cadence TESS *eleanor* PSF-extracted lightcurve (black). Some of the eclipses were missed due to data gaps. The CBP transits are highlighted in green; Middle panel: Sectors 4-12 of 2-min cadence SAPFLUX lightcurve (gray); Lower panels: 5-day sections of the lightcurve centered on the three CBP transits (highlighted in green). The middle and right panel also show primary and secondary eclipses near days 1483 and 1577, respectively.

In addition to the prominent stellar eclipses the light curve of TOI-1338 contains several small, transit-like events that required further scrutiny. Specifically, we noticed four events near days 1390, 1391, 1403, and 1404 (Sector 3), one event near day 1484 (Sector 6), and an event near day 1579 (Sector 10). Using the PSF fitting built into *eleanor* we showed that the events near days 1391, 1484, and 1579 are astrophysical in origin—namely, the three transits of the CBP—while the remaining events are artifacts caused by pointing “jitter” (mainly before reaction wheel “momentum dumps”).

The 30-min cadence light curve from Sector 3 was particularly difficult to extract because the target fell near an edge of the detector as shown in Figure 2 (the target was well away from the detector edges in the remaining Sectors). Additionally, the light curve extraction is especially prone to systematic errors as occasional spacecraft pointing jitter during Sector 3 moved some of the light from the target off of the detector, thereby producing transit-like events in the observed flux. As discussed below, the PSF of the target changes during some of the events listed above, which helps us rule out an astrophysical origin.

The PSF can be modeled in *eleanor* using either a two-dimensional Gaussian or a Moffat profile, and in this case both models perform equally well. Briefly, the analysis proceeded as follows. At each cadence, we fit parameters that describe the shape of the stellar PSF, assuming all stars in a 13×13 pixel region share the same PSF. We then optimized the flux of each star, the shared PSF parameters, and a single background level across this region. Based on this analysis, we found that during the events near days 1390, 1403 and 1404 in Sector 3, the shape of the PSF changed, affecting how much of the star falls in the optimal aperture and therefore how much flux is observed. It is likely that the pointing became “looser” during the times of these events, and the PSF-fitting package interpreted the resulting images as an increase in the size of the PSF. The final pointing algorithm was implemented on board the spacecraft after Sector 4, and TESS experienced sporadic pointing errors more frequently and of higher amplitude prior to that. Owing to an unfortunate set of coincidences, the spurious transit events in Sector 3 happen to have similar depths and durations as the real CBP transits and secondary eclipses. Given the PSF change, we then built a linear model from the out-of-eclipse data that predicts the flux of the star at every cadence from the PSF parameters alone. Figure 3 shows part of the Sector 3 light curve from *eleanor* using aperture photometry (top curve) and the model light curve predicted from changes in the PSF (bottom curve). We found that the PSF-model predicts transit-like features near days 1390, 1402, and 1405. However, this model does *not* predict transit-like events at the time of a secondary eclipse (days 1387 and 1402) or near day 1391 (CBP transit). This is further demonstrated in Figure 4, showing a section of the per-pixel light curve for the two events near days 1402 and 1405—the events are anti-correlated among some

of the core pixels such that they appear as transit-like in some pixels and anti-transit-like in others, indicating data artifacts. We thus confirmed the reality of the day 1391 transit-like event (and also the secondary eclipses near days 1387 and 1402) and conclude that the apparent transit-like events near days 1390, 1403 and 1404 are instrumental artifacts. A similar analysis confirmed the reality of the two CBP transits in Sectors 6 and 10. We found the long- and short-cadence light curves to have different eclipse depths in some sectors because of different levels of background subtraction. In many cases, the short cadence data overestimated the background, causing many “sky” pixels near the star to record negative flux values and the eclipses to appear artificially deep. We re-fit a background model for the short cadence data using the full-frame images, interpolating to the times of each short cadence exposure, leading to consistent eclipse depths between the two datasets. We also tested a variety of different apertures, finding the choice of aperture did not make a significant difference on the ultimate photometry, and in most sectors we used the pipeline default aperture.

We achieved the best photometric precision for TOI-1338 using the PSF-based photometry built into *eleanor*. Our final adopted light curve is a combination of 30-min cadence data extracted from the FFIs for Sectors 1, 2, and 3, and 2-min cadence data extracted from the target pixels in the “postage stamps” for this particular target for the remaining sectors. This final light curve was detrended and normalized in the manner described in [Orosz et al. \(2019\)](#). As part of this iterative detrending process, we measure the durations of all eclipse and transit events. Most of the out-of-eclipse portions of the light curve are trimmed, and we keep only the out-of-eclipse regions that are within 0.25 days or 1.25τ of each event (which ever is greater), where τ is the duration of the event. The last primary eclipse in Sector 12 had unusually large residuals (for unknown reasons), and consequently we excluded it from further analysis.

We measured the times of the eclipses and transits by fitting a simple model to trimmed and normalized light curves for each event using the Eclipsing Light Curve (ELC) code of [Orosz & Hauschildt \(2000\)](#). This model has nine free parameters: the period P , the conjunction time T_{conj} , the inclination i , the primary radius R_1 , the ratio of the radii R_1/R_2 , two limb darkening parameters for the quadratic limb darkening law $q_{1,1}$, $q_{2,1}$, and two eccentricity parameters $\sqrt{e}\cos\omega$ and $\sqrt{e}\sin\omega$. The goal was to find a smooth and symmetric curve that best fit each segment, so no attempt was made to optimize more than one segment at a time. For each segment, we found the best-fitting model using the Differential Evolution Monte Carlo Markov Chain (DE-MCMC) algorithm of [Ter Braak \(2006\)](#) with 80 chains that was run for 5000 generations. Using the best-fitting model, the individual uncertainties on each point were scaled in such a way to get $\chi^2_{\text{best}} = N$, where N is the number of points. The scale factors for the first five primary eclipses (observed in long cadence) were 0.347, 0.353, 0.626, 0.690, and 0.770, respectively. For primary eclipses observed in long cadence, the scale factors ranged from 1.042 to 1.376 with a median of 1.121. After the uncertainties on the data in each segment were scaled, the DE-MCMC code was run using 80 chains for 40,000 generations. Posterior samples were selected starting at generation 400, and sampling every 400th generation thereafter. The median of each posterior sample was adopted as the eclipse time, and the rms of the sample was taken to be the 1σ uncertainty. The measured primary and secondary cycle numbers and times are given in [Table 1](#). The cycle numbers for the secondary eclipses are given as fractional values of the form NN.45345 (the mean phase of the secondary eclipses is not 0.5 owing to the orbital eccentricity).

To make the final light curve that was modeled using the photodynamical model described in [Section 4](#) below, we simply combined the individual segments with scaled uncertainties from all of the eclipse and transit events. Since each segment has had its uncertainties scaled individually, no one event should be given undue weight owing to underestimated uncertainties for the photometric measurements. The scale factors for the segments in long cadence were smaller than one, which suggests the photometric errors were overestimated. On the other hand, the scale factors for the event observed in short cadence were all larger than one, which suggests the photometric errors were underestimated.

3. COMPLEMENTARY OBSERVATIONS

3.1. Radial Velocity Characterization

After its initial classification as an eclipsing binary, EBLM J0608-59 was observed with the CORALIE spectrograph to constrain the component masses. Between December 2009 and April 2012, 19 radial velocity observations were performed to map out the Keplerian orbit of the binary ([Triaud et al. 2017a](#)). Exposures were typically 600 s long, yielding a median precision of 34 m s^{-1} . Owing to the small mass ratio of the binary, and indeed of the entire EBLM sample by construction, the system appears as a single-line spectroscopic binary. We expect a V magnitude difference of 9.8 between the primary and the secondary, given that $M_1 \sim 1.1 M_\odot$, $M_2 \sim 0.3 M_\odot$, respectively ([Triaud et al.](#)




Figure 2. Nominal *eleanor* apertures for Sectors 3, 6, and 10. The target was on the edge of the detector in Sector 3, as seen in the left panel. Hence in this case a custom aperture is needed.




Figure 3. Aperture photometry for Sector 3 data (upper curve) with the flux model predicted from changes in the shape of the PSF at each cadence inferred by *eleanor* PSF modeling (lower curve). We expect that spurious events caused by changes in the PSF to appear in both curves, whereas astrophysical events, which occur independently of the instrumental PSF, will only appear in the upper curve. Thus, the apparent events near days 1390, 1403 and 1404 are caused by changes in the PSF shape which occur as parts of the star fall off and on the detector owing to pointing jitter. The purple- and green-shaded regions correspond to the locations of secondary eclipses and a transit of the planet, respectively. These events are apparent only in the upper curve, which is a strong indication of their validity. Finally, we note the deep feature near day 1395 is a primary eclipse.

2017a). As a consequence of the large flux ratio between the primary and secondary, the spectral lines of the secondary are not noticeable in the observations, thereby allowing high-accuracy measurements of the primary’s radial velocity. Likewise, the secondary star is too faint for TESS to be able to detect transits of the CBP across it.

3.1.1. The BEBOP radial velocity search for circumbinary planets

In late 2013 the BEBOP (Binaries Escorted By Orbiting Planets) program was created as a radial velocity survey for the detection of circumbinary planets. The BEBOP sample is exclusively constructed from eclipsing single-line spectroscopic binaries to avoid contamination effects that severely hinder planet detection in double-line spectroscopic binaries (Konacki et al. 2009). An initial target list of roughly 50 binaries was created from the larger EBLM sample. Selection criteria include the obtainable radial-velocity precision and the lack of stellar activity. EBLM J0608-59 was included in this initial BEBOP selection. Between November 2014 and August 2015 we acquired 17 additional CORALIE measurements with longer exposures of 1800 s to increase the precision of the radial velocity measurements (the median uncertainty was 25 ms^{-1}). This second set of observations coincided with a change to a new octagonal fibre. The new fibre provided greater long-term stability compared to the original circular one, but at the cost of $\sim 10\%$ of the incoming flux. Such a fibre change may also induce a small velocity offset (Triaud et al. 2017b). Consequently, when modelling the data we treat the CORALIE data sets as if they were acquired from two different instruments, with a free parameter for the offset.




Figure 4. A section of the TESS *eleanor* per-pixel light curve for Sector 3 near days 1402 and 1405. The panels with black light curves represent the core pixels used for light curve extraction. The two events highlighted in green appear as transit-like in some of the core pixels and anti-transit-like in others. This demonstrates that these two events are data artifacts (see Fig. 3).

The combined radial velocities from both the EBLM and BEBOP surveys are presented in (Martin et al. 2019). No circumbinary planet was detected in the radial velocity of this system within the sensitivity of the observations, which were adequately fit by a single Keplerian orbit ($\chi^2_{\nu} \sim 1$). This ruled out (at 95% confidence) the presence in the system of a CBP more massive than $\sim 0.7M_{\text{Jup}}$ and with orbital period of roughly six times shorter than that of the binary.

Starting in April 2018 the BEBOP survey was extended to the HARPS high-resolution spectrograph on the ESO 3.6-m telescope (Prog.ID 1101.C-0721, PI A. Triaud; Pepe et al. 2002), also at La Silla, Chile⁷. Compared to CORALIE, HARPS benefits from a larger telescope aperture, a higher resolving power of $R = 115,000$, and greater radial-velocity stability by being both thermally-stabilized *and* operated under vacuum. Seven HARPS spectra of the target have been acquired to date. They were reduced with the HARPS pipeline, which has been shown to achieve remarkable precision and accuracy (e.g. Mayor et al. 2009; López-Morales et al. 2014). The radial-velocities were computed by using a binary mask corresponding to a G2 spectral-type template (Baranne et al. 1996). We achieve a median radial-velocity precision of 5.9 m^{-1} in our measurements. A fit to the complete set of radial-velocities (CORALIE and HARPS) produces a fit of reduced $\chi^2_{\nu} \sim 1$ mostly because the number of HARPS measurements is close to the number of free parameters for the binary orbit. The model adjusts to the HARPS measurements first, because they have the greatest weights. CORALIE’s precision is not sufficient to detect the additional planetary signal. The HARPS measurements indicate that this system has very little activity, making it optimal for radial-velocity measurements. The BEBOP survey is ongoing and more spectra from both HARPS and ESPRESSO will be obtained for TOI-1338/EBLM J0608-59, and published in a subsequent paper.

3.2. Spectroscopic Characterization

The seven extracted HARPS spectra were co-added onto a common wavelength axis reaching a signal-to-noise ratio of approximately 73. The resulting spectrum was analysed with the spectral analysis package ISPEC (Blanco-Cuaresma et al. 2014). We used the synthesis method to fit individual spectral lines of the co-added spectra. The radiative transfer code SPECTRUM (Gray & Corbally 1994) was used to generate model spectra with MARCS model atmospheres (Gustafsson et al. 2008), version 5 of the GES (GAIA ESO survey) atomic line list provided within ISPEC and solar abundances from Asplund et al. (2009). Macroturbulence is estimated using equation (5.10) from Doyle (2015) and microturbulence was accounted for at the synthesis stage using equation (3.1) from the same source. The $\text{H}\alpha$, Na I D and Mg I b lines were used to infer the effective temperature T_{eff} and gravity $\log g$ while Fe I and Fe II lines were used to determine the metallicity $[\text{Fe}/\text{H}]$ and the projected rotational velocity $v \sin i_*$. Trial synthetic

⁷ BEBOP also surveys the northern skies, using SOPHIE, at the Observatoire de Haute-Provence, Prog.ID 19A.PNP.SANT, PI A. Santerne. The northern EBLM sample will also be included in the TESS 2 minute cadence under proposal G022253, PI D. Martin.




Figure 5. Left: radial velocities for TOI-1338, phase-folded on the binary orbital period of 14.61 days, taken with the CORALIE instrument (blue and orange) and HARPS (red). Right: residuals to the single-Keplerian radial velocity fit over time. The change in CORALIE residuals between blue and orange measurements coincide with a fibre change (see text for details).

model spectra were fit until an acceptable match to the data was found. Uncertainties were estimated by varying individual parameters until the model spectrum was no longer well-matched to the spectra of TOI-1338. For the primary star, we find an effective temperature of $T_{\text{eff},1} = 6050 \pm 80$ K, a metallicity of $[\text{Fe}/\text{H}]_1 = 0.01 \pm 0.05$, a gravity of $\log g_1 = 4.0 \pm 0.08$ dex, and a projected rotational velocity of $v \sin i_{*,1} = 3.6 \pm 0.6$ km s $^{-1}$. These measurements are summarized in Table 2.

As a check on the spectroscopic temperature of the primary, we first gathered brightness measurements from the literature in the Johnson, Tycho-2, 2MASS, and Sloan systems, constructed 10 non-independent color indices, and corrected each for reddening using the extinction law of Cardelli et al. (1989) with a value of $E(B - V) = 0.020 \pm 0.010$ derived from the extinction map of Schlafly & Finkbeiner (2011) and the *Gaia* (*Gaia* Collaboration et al. 2016, 2018) distance of 405 ± 3 pc. These colors are unaffected by the secondary star because it is so faint. We then used color-temperature calibrations from Casagrande et al. (2010) and Huang et al. (2015) to infer a mean photometric temperature of 5990 ± 110 K, in good agreement with the spectroscopic value.

We estimated the radius of the primary star in three different ways. One method used the procedure outlined by Stassun et al. (2019) for the preparation of the TIC-8 catalog, involving the *Gaia* parallax and G magnitude, an extinction correction, the spectroscopic temperature, and the G -band bolometric correction. Another method we used was based on a fit to the spectral energy distribution (SED) performed with EXOFASTv2 (Eastman et al. 2019) and the MIST bolometric correction tables⁸, using brightness measurements in the *Gaia* and WISE systems in addition to those mentioned earlier. Suitable priors were placed on the distance, extinction, temperature, metallicity, and $\log g_1$. A third method we used was also based on an SED fit but used instead the NextGen model atmospheres of Allard et al. (2012). The three procedures gave very similar results. We adopt the value $R_1 = 1.345 \pm 0.046 R_{\odot}$ in the following to use as a prior for the photometric-dynamical modeling described below.

3.2.1. Image Analysis

The relatively large sizes of the TESS pixels, approximately $21''$ on a side, leave the target susceptible to photometric contamination from nearby stars, including additional wide stellar companions. The nearest object (TIC 260128336) to TOI-1338 after proper motion correction is separated by $53.6''$. The SPOC Data Validation centroid offsets for the Sectors 1-12 multi-sector run indicated that the primary and secondary eclipses for TOI-1338 originate from the target itself (Jenkins et al. 2016); complementary analysis with the photocenter module of DAVE (Kostov et al. 2019) confirmed this. We also searched for nearby sources with SOAR speckle imaging (see Tokovinin et al. 2018, 2019, for details of the instrumentation) on 17 March 2019 UT, observing in a similar visible bandpass as TESS. We detected no nearby sources within $3''$ of TOI-1338. The detection sensitivity and speckle auto-correlation function of the SOAR observations are plotted in Figure 6.

4. PHOTOMETRIC-DYNAMICAL ANALYSIS OF THE SYSTEM

⁸ http://waps.cfa.harvard.edu/MIST/model_grids.html




Figure 6. The 5σ detection limits of the SOAR speckle imaging of TOI-1338; the inset is the speckle auto-correlation function. No nearby stars were detected within $3''$ of the target.

Due to the rich dynamical interactions between the two stars and a planet in a CBP system, the deviations of the planet’s orbit from a strictly periodic one are much more pronounced compared to a single-star system. The secular evolution in a CBP system can occur on a timescale as short as a decade instead of thousands of years (like in the Solar System). Thus measurable changes in quantities such as the inclination of the planet’s orbit can be observed with missions like *Kepler* and TESS. Given the relatively rapid secular evolution, the orbits in a CBP system are not simple Keplerians. A complete description of a CBP system relies on a large number of parameters—e.g. masses, radii, and orbital parameters for the two stars and the planet(s), radiative parameters for the two stars, etc. Stellar eclipses, both primary and secondary, allow for precise measurements of times of conjunction. In addition, transits of the CBP across the primary and/or the secondary star can provide precise position measurements of both the stars and the planet at times other than the times stellar conjunction. Thus, the dynamical complexity of the system—while computationally challenging—enables precise measurements of the system’s parameters. For example, the stellar masses and radii of the two stars in the Kepler-16 CBP system have been measured to sub-percent precision (Doyle et al. 2011).

4.1. ELC Modeling

To obtain a complete solution for the TOI-1338 system, we carried out a photometric-dynamical analysis with the ELC code (Orosz & Hauschildt 2000), utilizing the photometry from TESS and the precise radial velocity measurements from CORALIE and HARPS. For this task, the ELC code combines N-body simulations, modified to include tidal interaction and general relativistic effects, with a photometric model for the stellar eclipses and planetary transits, to reproduce a light curve of the system. These modifications to ELC to allow for modelling stellar triple and higher-order systems, and CBP systems have been described in Welsh et al. (2015) and Orosz et al. (2019). This code has been used extensively for the analysis and confirmation of *Kepler* CBPs; for the sake of completeness we outline it below.

Briefly, given instantaneous orbital parameters (e.g. the orbital period, the eccentricity, the inclination, etc.) at some reference epoch and the masses of each body, ELC solves the Newtonian equations of motion using a symplectic

integrator, in this case a 12th order Gaussian Runge Kutta integrator [Hairer et al. \(2002\)](#). When necessary, the Newtonian equations of motion can be modified to account for general relativistic precession and tidal effects (see [Mardling & Lin 2002](#)). The solution of the dynamical equations enables the positions of all the bodies to be specified at any given time. Then, given the positions of the bodies on the plane of the sky, their radii, and their radiative properties, the observed flux is computed for any number of overlapping bodies using the algorithm discussed in [Short et al. \(2018\)](#). Likewise, the solution to the dynamical equations gives the radial velocity of each body at any given time. It is also possible to include other observable quantities that do not depend on time (for example the surface gravity of the primary star) in the fitting process.

For TOI-1338, we initially had the following 25 free parameters. The binary orbit is specified by the time of conjunction $T_{\text{conj,bin}}$, the period P_{bin} , the eccentricity parameters $\sqrt{e_{\text{bin}} \cos \omega_{\text{bin}}}$ ⁹ and $\sqrt{e_{\text{bin}} \sin \omega_{\text{bin}}}$ (where e_{bin} is the eccentricity and ω_{bin} is the argument of periastron), and the inclination i_{bin} . We fix the initial nodal angle of the binary orbit to $\Omega_{\text{bin}} = 0$. The stellar masses are specified by the primary mass M_1 (in units of M_{\odot}) and the mass ratio $Q = M_2/M_1$, the stellar radii—by the primary radius R_1 and the radius ratio R_1/R_2 , and the stellar temperatures—by $T_{\text{eff},1}$ and the temperature ratio $T_{\text{eff},2}/T_{\text{eff},1}$. We assume a quadratic limb darkening law with the [Kipping \(2013\)](#) “triangular” resampling, for a total of four additional parameters. Finally, to account for tidal precession we use two additional free parameters—the apsidal constants $k_{2,1}$ and $k_{2,2}$ for the primary and secondary, respectively. For the planet’s orbit, we fit for the sum and differences of the period and time of barycentric conjunction, i.e. $\zeta_p = (P_{\text{pl}} + T_{\text{conj,pl}})$, $\zeta_m = (P_{\text{pl}} - T_{\text{conj,pl}})$, the eccentricity parameters $\sqrt{e_{\text{pl}} \cos \omega_{\text{pl}}}$, $\sqrt{e_{\text{pl}} \sin \omega_{\text{pl}}}$, the inclination i_{pl} , and the nodal angle Ω_{pl} . The planet’s mass is specified by M_3 , where the units are M_{\oplus} . Finally, the radius of the planet is specified by the ratio R_1/R_{pl} .

The observables for the TOI-1338 system are the trimmed and normalized light curve from TESS, and the radial velocity measurements from CORALIE and HARPS. The radial velocities span three distinct data sets where an offset velocity is found for each, namely the “early” CORALIE data before the fibre change (roughly days 200 to 1000 in units of BJD - 2,455,000), the “late” CORALIE data post fibre change (roughly days 1900 to 2250), and HARPS (roughly days 3200 to 3550). Another set of observables is the spectroscopically-measured gravity for the primary star ($\log g_1 = 4.0 \pm 0.08$ dex in cgs units), and the radius of the primary star ($R_1 = 1.345 \pm 0.046 R_{\odot}$) as determined from multi-color photometry and the distance. We use the usual χ^2 statistic for the likelihood, where the χ^2 contributions for the light curve, the velocity curve, the spectroscopic gravity, and the computed radius are combined. In general, model parameters are drawn uniformly within specified bounds. The use of the primary radius R_1 as both a fitting parameter and an observed parameter effectively gives that parameter a Gaussian prior. Finally, we also include estimates of the times of the planet transits with generous uncertainties (0.0050 days for all events) in the overall χ^2 . Using estimates of the transit times is very important early in the fitting process as models with transits that are very far away from the observed times would have the same χ^2 contribution from the light curve as models that are much closer to the observed times since the out-of-transit parts of the light curve are flat. Including observed transit times in the overall χ^2 gives a larger penalty to the models that miss the observed times by larger amounts compared to models that have smaller differences between the model transit times and the observed ones. As the models converge to the “true” solution, the χ^2 contribution from the eclipse times becomes very small, leaving the actual fit to each individual transit profile in the light curve to determine the χ^2 contribution to the overall total χ^2 .

The reference time for the osculating orbital parameters was chosen to be day 186 in units of BJD - 2,455,000. The ending time of the numerical integration was day 3750 in the same units. For these integrations we include the effects of both precession due to general relativity and precession due to tides. As some measure of the round-off errors that afflict nearly every numerical integrator, we measure the position and velocity of the system center of mass as a function of time. These two quantities start out at the origin [e.g. at the coordinates (0,0,0)] by default, and should remain at the origin for the ideal integrator. We find that the center of mass at day 3750 is offset by $\approx 3.40 \times 10^{-13}$ AU (≈ 5.09 cm), and that the velocity at this time is offset by $\approx 2.00 \times 10^{-16}$ AU per day ($\approx 3.46 \times 10^{-10}$ meters per second). Thus we conclude that numerical round-off error is not an issue for these integrations.

ELC has a number of optimizers available, and the two that proved to be the most useful for the analysis of TOI-1338 were the genetic algorithm of [Charbonneau \(1995\)](#) and the Differential Evolution Monte Carlo Markov Chain (DE-MCMC) algorithm of [Ter Braak \(2006\)](#). The spectroscopic parameters of the binary were well-known from analytic models of the radial velocities. The other parameters, especially the orbital parameters for the planet, were initially

⁹ Combinations and/or ratios of individual parameters are often more convenient to use.

not well constrained. Thus our fitting proceeded by iteration. The genetic code was run where an initial “population” of 100 models was evolved for a few thousand generations. Next, the top 80 models were evolved using the DE-MCMC code for a few thousand more generations. The top few models, randomly chosen to be between one and five, were put back into the genetic code along with random models and evolved. After a few iterations of this process, a reasonably good model was found.

After an initial good model was found, we performed several pilot runs of the DE-MCMC code. The initial population of models consisted of the best model and mutated copies of the best model where several randomly chosen parameters were offset from their optimal values by small amounts drawn from a normal distribution. The standard deviation of the normal distribution was chosen to be between 0.001 and 0.01 times the parameter value. We found that the chains quickly spread out, and achieved their final overall spreads after a burn-in period of typically 1000 generations. During these pilot runs, we checked to confirm that the prior ranges included support for the entire range with non-trivial likelihood, and that the likelihood falls to extremely small values by the time model parameters reach any of these boundaries (with possible exception of hard physical boundaries). We also found that the two apsidal parameters and the limb darkening coefficients for the secondary star were not constrained. Consequently, we fixed their values at representative values ($k_{2,1} = 0.01$, $k_{2,2} = 0.10$ for the apsidal parameters and $q_{1,2} = 0.06$, $q_{2,2} = 0.41$ for the secondary’s quadratic law limb darkening coefficients), reducing the number of free parameters to 21.

For the final step leading to the adopted parameters and their uncertainties, we used a brute-force “grid search” algorithm to find optimal models with the third body mass fixed at values from 0.3 to $117.3 M_{\oplus}$ in steps of $1 M_{\oplus}$ as “seed” models for the final runs of the DE-MCMC code. We then ran the DE-MCMC code 8 separate times, each with the same seed models, but with a different initial random number seed. All 8 runs used 120 chains, and each was run for at least 8800 generations (the longest run had 38,300 generations). The posterior samples were drawn starting at generation 3000, with subsequent draws that skipped every 2000 generations until the chains ended. The individual posterior samples were combined into single samples (with $N = 12120$) for each fitting parameter and for several derived parameters of interest. Table 3 provides the parameters for the best-fitting model, the mode of the posterior sample (found using 50 bins), the median of the posterior sample, and the $+1\sigma$ and -1σ uncertainties. Table 4 provides several derived parameters of astrophysical interest using a similar format as Table 3. In the discussion that follows we use the posterior medians as the adopted parameter values. The model fits to the stellar eclipses are shown in Figure 7, and the model fits to the planet transits are shown in Figure 8.

There is a slight difference between the depths of the long-cadence primary eclipses and the model, such that the former are deeper. To address this issue, we reran the fit using a negative contamination factor for the long cadence data in order to make the model deeper and match the data. The best-fit dilution factor is -0.0336 ± 0.0022 , indicating that there may be a variable flux offset between the long- and short- cadence data. To first order, this suggests that either the sky background in the long cadence data was over-subtracted by 3.4%, or the sky background in the short-cadence data was under-subtracted by 3.4%. This dilution term seems to be somewhat on the high side of what one might expect based on the number of counts in the actual data. It is also hard to determine to which data set it has to be applied (long cadence only, short cadence only, or a combination of both). New TESS observations in Cycle 3 will help address this issue. Regardless of the specific reason and approach, the key parameters (mass, radius, etc.) do not change significantly and the results are consistent between the models with and without a dilution factor. The radial velocities are fit quite well, and Figure 10 shows the residuals for the “early” CORALIE, the “late” CORALIE, and the HARPS measurements. Finally, in Table 5 we give the initial dynamical parameters and Cartesian coordinates for the best-fitting model to full machine precision.

4.2. Planet Mass

As noted in Section 2.3, there is undoubtedly a transiting circumbinary object in the TOI-1338 system. Our analysis here shows that the mass of this object is well within the planetary regime, i.e. $M_3 = 33 \pm 20 M_{\oplus}$. Here we give a brief discussion of what feature(s) in the data allow us to make this determination. The top of Figure 11 shows the posterior distribution of the planet mass M_3 in units of M_{\oplus} . The minimum and maximum values in the posterior sample are $0.01 M_{\oplus}$ and $117.05 M_{\oplus}$, respectively. The bottom of Figure 11 shows $\chi^2 - \chi_{\min}^2$ for all computed models, where $\chi_{\min}^2 = 14587.93$. We see that $\Delta\chi^2$ is about 4 when $M_3 = 0$, and about 18 when $M_3 = 117 M_{\oplus}$. As a reminder, the total χ^2 has contributions from the TESS light curve, the three radial velocity sets, the measured gravity, and the measured radius of the primary star. For the best-fitting model, these contributions are 14542.39 (TESS), 13.88 (early CORALIE), 17.11 (late CORALIE), 5.44 (HARPS), 8.07 ($\log g$), and 1.01 (R_1). For the best model with $M_3 = 120 M_{\oplus}$,

those values are 14545.53 (TESS), 17.67 (early CORALIE), 20.31 (late CORALIE), 14.85 (HARPS), 9.72 ($\log g$), and 0.02 (R_1). There is hardly any change in the χ^2 fit to the TESS light curve between the two models. The data set with the largest change in χ^2 is HARPS, where the χ^2 changed by about 9.4. When the planet mass is fixed at $150 M_\oplus$, the total χ^2 is 14626.61 (38.68 larger than the overall best model), and the individual contributions are 14553.20 (TESS), 17.96 (early CORALIE), 19.46 (late CORALIE), 25.32 (HARPS), 10.05 ($\log g$), and 0.04 (R_1). Although the χ^2 for the TESS light curve got slightly worse, it seems that the HARPS radial velocity measurements have the most sensitivity to the planet mass. The bottom of Figure 10 shows the residuals of the HARPS measurements for the best overall model, the best model with $M_3 = 120 M_\oplus$, and the best model with $M_3 = 150 M_\oplus$. The residuals of the first measurement near day 3220 and the last two measurements near days 3560 and 3570 show the most variation with the changing planet mass. Thus in the near term the most effective way to better constrain the mass of the planet would be to obtain more radial velocity measurements with a quality similar to or better than the HARPS measurements presented here.

A circumbinary planet can perturb the binary and give rise to eclipse time variations (ETVs). The interaction between the planet and the binary can cause the binary orbit to precess, which leads to changes in the phase difference between the primary and secondary eclipses. When one attempts to fit the primary and secondary eclipse times to a common ephemeris, the O-C (Observed minus Computed) values for the primary eclipses will have the opposite slope that the O-C values for the secondary eclipses have. Figure 12 shows the Common Period O-C diagram for the model primary and secondary eclipses over the whole time span of the radial velocity and photometric observations. For our overall best-fitting model, the argument of periastron ω changes by 0.0005715 degrees per cycle. The contribution of this precession from General Relativity (GR) is 0.0001132 degrees per cycle, and the contribution from tides is 0.0000055 degrees per cycle. This precession causes a divergence between the primary and secondary O-C curves of about 2 minutes over the roughly 10 year time span of the data. When the planet mass is fixed at $150 M_\oplus$, the best-fitting model for that mass has a change in ω of 0.002129 degrees per cycle. This results in a divergence between the primary and secondary O-C curves of about 9 minutes. As a practical matter, we only have measurements of eclipse times over the last 1.5 years or so, and the uncertainties are relatively large: 0.36 minutes for the primary eclipses and 5.40 minutes for the secondary eclipses. Unless the measurements of the eclipse times can be vastly improved, we would need many more years of eclipse time measurements before the time baseline is long enough to accumulate a measurable divergence in the Common Period O-C diagram.

4.3. Dynamical Evolution

The large tidal potential produced by the inner binary causes the orbital elements of the CBP to vary with time. Indeed, the best-fit osculating orbital elements (Table 4) represent only a snapshot at the reference epoch. These variations have consequences for both the stability and observability of CBPs (see Sect. 4.4). Dynamical studies of CBPs indicate that a critical regime exists, such that CBPs with periods less than P_{crit} are unstable, predominantly scattering onto an unbound orbit, or occasionally colliding with either star (Dvorak 1986; Holman & Wiegert 1999; Sutherland & Fabrycky 2016; Lam & Kipping 2018; Quarles et al. 2018). The process by which this instability occurs is resonant overlap (Mudryk & Wu 2006; Sutherland & Kratter 2019). The value of P_{crit} is primarily a function of the dynamical mass ratio of the host stars $\mu = M_2/(M_1 + M_2)$, binary period P_{bin} and eccentricity e_{bin} , but also has a dependence on the mutual inclination between the two orbits, ΔI .

To investigate the dynamical evolution of the TOI-1338 system, we integrated the orbit of the CBP for $\sim 10^5$ yr (corresponding to $\sim 7 \times 10^3$ orbits of the binary) using the best-fit photometric-dynamical solution in Table 4, and the IAS15 integrator in the REBOUND integration package (Rein & Liu 2012; Rein & Spiegel 2015). Figure 13 shows the evolution of the system for 40,000 yr, using as the initial condition the best-fit photometric-dynamical solution from Table 5. As seen from the figure, both the CBP (black) and binary (red) semi-major axes are practically constant with time, indicating that the system is stable. Over the course of these integrations, the eccentricity of CBP varies in a small range from 0.0695 to 0.1763 (Figure 13b).

To explore whether the planet's eccentricity will continue to increase, and if that will affect the stability of its orbit, we used a modified version of the `mercury6` integration package Chambers et al. (2002) and integrated the system for 10^7 yr. Our results showed that the extrema for the planetary eccentricity extends by only an additional, but insignificant amount of 0.0004, confirming that the orbit of the CBP is long-term stable. Figure 14 demonstrates a more global range of stability using the `mercury6` integrator for binaries, tracking the extrema of planetary eccentricity (Dvorak et al. 2004; Ramos et al. 2015). As shown here, the orbit of the CBP (green dot) lies between 6:1 and 7:1




Figure 7. The folded primary and secondary eclipses are shown with the best-fitting model (solid red line). Upper left: primary eclipse in with residuals. Upper right: Primary eclipse in and 30-min cadence with residuals. Lower left: secondary eclipse in with residuals. Lower right: secondary eclipse in 30-min cadence with residuals. There is a slight trend of the model being shallower than the long-cadence primary eclipses.




Figure 8. The transits of the CBP across the primary in Sector 3 (left), Sector 6 (middle), and Sector 10 (right) and the best-fitting model. The Sector 6 and 10 events were fit using 2-min cadence data, but for clarity we show the data binned to the 30-min cadence.

mean-motion resonances (MMRs) (downward ticks, top axis) with the binary. This is important for long-term orbital stability, and is well below the eccentricity-dependent stability limit (dashed line; Quarles et al. (2018)). This is an expected result and a consequence of the fact that CBPs form at large distances away from the binary and migrate to their current orbits (e.g. Pierens & Nelson 2013; Kley & Haghighipour 2015). Those that maintain stable orbits are trapped between two $N : 1$ MMR with the binary. This has indeed been observed in all *Kepler* CBPs as it is critical for long-term orbital stability. Figure 14 also shows that, although the orbit of the CBP is stable, small changes in its semimajor axis or eccentricity may result in a more chaotic orbit by situating the planet near a region of instability corresponding to $N : 1$ MMRs with the binary.

As an independent test to examine the stability of the planet, we used the results of our numerical integrations in the context of the scheme developed by Quarles et al. (2018), and identified a region around the binary where the orbit of the CBP will certainly be unstable. Our analysis shows that the outer boundary of this unstable region corresponds to $P_{crit} = 64.3$ days ($a_{crit} = 0.36$ AU). The observed planetary period is $\approx 50\%$ longer than this critical value¹⁰, once again confirming that the orbit of the CBP is stable. Additionally, we also used a frequency analysis (Laskar 1993) to obtain a quasi-periodic decomposition of the orbital perturbations of the CBP. We found these to be a combination of the five fundamental frequencies—i.e. the mean motions of the orbit of the binary, the CBP, the apsidal precession of the binary and CBP orbits, and the nodal precession—and fully consistent with the numerical simulations.

Our numerical simulations also show indications of both apsidal and nodal precessions in the orbit of the CBP. Figures 13c and 13d show the x -components of the planet’s eccentricity (apsidal) and inclination (nodal) vectors. As seen here, many secular precession cycles of the planet occur within the span of 40,000 yr. The figures show a mode

¹⁰ The planetary semi-major axis is $\approx 30\%$ larger than the critical semi-major axis.




Figure 9. Same as Fig. 7, but for a model with a dilution factor for the long cadence data as an additional parameter to address the trend seen in that figure. While this model suggests that there is a dilution of $\sim 3.4\%$ between the long- and short-cadence data, the best-fit parameters of the system do not change significantly.




Figure 10. The residuals for the radial velocities measured by CORALIE (top and middle, black points) and HARPS (bottom, black points) for the best-fitting photodynamical model. The red and blue points show the residuals of the HARPS measurements for models with the planet mass fixed at $120 M_{\oplus}$ and $150 M_{\oplus}$, respectively.

with a $\sim 14,286$ year period, and variations that occur on a much shorter timescale of decades. We use the Fast Fourier Transform (FFT) routine within `scipy` to produce the periodograms shown in Figures 13e and 13f, where the system was evolved for 100,000 yr. These periodograms show strong peaks at ~ 23 yr (8375 days) for the planetary apsidal precession period and 21.4 yr (7816 days) for the planetary nodal precession period. This nodal precession period differs slightly from the analytical result (8980 days) derived from formula given in Farago et al. (2009) since the stellar binary’s orbit is non-circular.

Similar to the CBP, the orbit of the binary also experiences nodal precession. This precession is predominantly due to the perturbation of the CBP, with the tidal precession and general relativistic effect being secondary. The planetary apsidal and nodal precessions occur with similar periods but in opposite directions, as expected from Lee & Peale (2006). The longer period mode ($\sim 14,286$ yr) for the planet is in phase with the binary secular precession period,




Figure 11. Top: The posterior distribution of the planet mass in units of M_{\oplus} . The median value is $33.0 M_{\oplus}$, and the largest value is $117.05 M_{\oplus}$. Bottom: $\chi^2 - \chi_{\min}^2$ vs. the planet mass in M_{\oplus} for all computed models.




Figure 12. A Common Period O-C diagram that shows the eclipse timing variations (ETVs) in minutes for the model primary eclipses (black lines) and the model secondary eclipse (red lines). The two inner lines are for the overall best-fitting model with $M_3 = 33 M_\oplus$, and the two outer lines are for the best model with the planet mass fixed at $M_3 = 150 M_\oplus$. The black point and error bar show the median uncertainty of 0.36 minutes for the measured times of the primary eclipses, and the red point and error bar show the median uncertainty of 5.40 minutes for the measured times of the secondary eclipses. TESS observed the system between days ≈ 3336 and ≈ 3634 .

where the binary causes an oscillation in the planetary argument of pericenter by $\pm 28^\circ$ per binary precession cycle (Andrade-Ines & Robutel 2018).

4.4. Transit and Detection Probabilities

The nodal precession of the planet’s orbit has important consequences for the long-term detectability of its transits as these can only occur when the projected path of the planet on the plane of the sky intersects with that of the stars. Nodal precession alters the planet’s path, even to the extent that transits disappear for long periods of time. This was predicted by Schneider (1994), and observationally confirmed by the transits of the CBP Kepler-413b (Kostov et al. 2014). In this context, Martin & Triaud (2014) discussed the “transitability” CBPs, and found that transits of CBPs could also occur in non-eclipsing binaries.

In Figure 15 we show how the impact parameter of the planet, and therefore its transitability, varies over time due to the orbital evolution of the planet. As seen from the figure, there are two windows of transitability per nodal precession cycle, each roughly ~ 1000 days wide as indicated by the points between the horizontal gray lines in the range $-1 \leq b \leq 1$. Figure 15b illustrates the motion of the planet on the sky plane for the 2000 days after the starting epoch, where the points (color-coded) are spaced by ~ 0.7 days and vary in size (larger, opaque is towards the




Figure 13. Evolution of the orbital elements of the CBP using the best-fit parameters from Table 5 for 40,000 years (BJD - 2,455,000). The top-left panel (a) shows small variations in the semimajor axis of the planet (black) and binary (red). The top-right panel shows the variations of orbital eccentricities, indicating that the system is stable. The evolution of the x -components of the eccentricity (c) and inclination vectors (d) illustrate apsidal and nodal precession, respectively. Panels (e) and (f) show Fourier periodograms using a 100,000 year simulation, where the peak values are the planetary apsidal and nodal precession periods for the respective vectors.

observer). The horizontal ellipses represent the orbits of the primary (black) and secondary (gray) stars with respect to the center of mass at the origin. The vertical ellipses indicate the cross section that each stellar disk takes up on the sky. When the larger, opaque points overlap with the vertical ellipse then transits are possible. As seen from Figure 15a, the planet transits 29.7% of the time. This is valid both for the best-fit solution from the posterior of the photometric-dynamical analysis (see Section 4), and for the solutions taken from the overall sample from the posterior, and propagated for 155,000 days (~ 20 nodal precession cycles), where the planet transits $29.6\% \pm 7.6\%$ of the time. This is typical of the *Kepler* CBPs, for which the mean primary transitability across the first ten discovered planets was $\sim 36\%$ (Martin 2017).

4.5. Future transits

As an aid to enable further observations of the TOI-1338b transits, we present in Table 6 the predictions of the times, impact parameters, and durations of future transit events. These three quantities were computed using 9000 models from the posterior sample. The quoted values are the sample medians, and the quoted uncertainties are the sample r.m.s. Transits will certainly occur on 2020 January 12, April 14, July 19, and October 19 since all 9000 models from the posterior had transits at these dates. Starting on 2021 April 26, not all models from the posterior produce transits at that time, so the transits become less likely. At first, the fraction of missed transits is rather small (a few percent), but then starting with the conjunction on 2023 May 20, the fraction of missed transits is large ($\sim 30\%$) and grows larger and larger thereafter. After the 2025 September 12 conjunction, the transits have $\lesssim 30\%$ chance of occurring, and if they do occur, their impact parameters will likely be close to -1 .




Figure 14. Stability map exploring the range of planetary eccentricity oscillation ($e_{\max} - e_{\min}$; color-coded) considering a wide range of initial values in planetary semi-major axis a_p (in AU) and eccentricity e_p . The white cells indicate unstable initial conditions on a 100,000 yr timescale, where the vertical dips denote the locations of $N : 1$ mean motion resonances (MMRs) with the inner binary. The current planetary parameters are indicated by a green dot, where the planet lies approximately midway between the 6:1 and 7:1 MMRs. The dashed curve shows the boundary of stability from Quarles et al. (2018), where the top axis marks the planetary period (in days) along with the location of the $N : 1$ MMRs.

The primary TESS mission ends in 2020 July, but fortunately the mission has been extended through at least the year 2022¹¹. Depending on the exact pointing schedule, there is a good chance that TESS can observe transits again on 2020 October 19, 2021 January 23, 2021 April 26, and possibly July 30.

For the sake of completeness, we also calculate the extent of the Habitable Zone (HZ Kasting et al. 1993) of the binary (Figure 16) using the Multiple Star Habitable Zone website developed by Müller & Haghighipour (2014). The inner boundary of orbital stability is shown in red and the orbit of the CBP is shown in blue. TOI-1338 b is substantially interior to the HZ, receiving more than 9 times the Sun-Earth insolation.

5. DISCUSSION

5.1. Comparison with stellar evolution models

We compared the best-fit stellar masses, sizes and temperatures of the primary and secondary stars of TOI-1338 against model isochrones from the MIST series (Choi et al. 2016) for the measured metallicity of the system. The fitted masses and sizes of both stars are in excellent agreement with a 4.4 Gyr isochrone model (see Figure 17). It is interesting to note that the M dwarf secondary of TOI-1338 does not seem to be significantly inflated compared to standard models, as may be the case for some CBP systems with similar secondary stars (e.g. Kepler-38 and Kepler-47, Orosz et al. 2012b, 2019). However, this is in line with the rest of objects detected during the EBLM Project, which show no systematic radius inflation for fully-convective low mass stars (von Boetticher et al. 2019). The effective temperature of the secondary star is also consistent with model predictions, within the errors.

¹¹ <https://tess.mit.edu/news/nasa-extends-the-tess-mission/>




Figure 15. Evolution of the impact parameter (a) due to nodal precession of the planetary orbit for 20,000 days (BJD - 2,455,000). Motion of the planet on the sky plane (b), where each point (color-coded) represents a small increment in the planetary orbit (~ 0.7 days). The larger, opaque points indicate when the planet lies between the binary and observer, while the smaller, translucent points are when the planet lies behind the binary orbit. The horizontal ellipses show the orbits of the primary (black) and secondary (gray) star about the center of mass, while the vertical ellipses illustrate the cross-section for transits to occur across each stellar disk.

5.2. Stellar rotation

Starspots create modulations in the light curves of eclipsing binaries (typically seen in the out-of-eclipse regions), which affect measurements of eclipse times and thus photodynamical models (see e.g. Kepler-453, Welsh et al. 2018). The systematic errors present in the TESS light curve of TOI-1338 (see Fig. 1) preclude measurement of the intrinsic stellar variability from the TESS data alone which, in turn, precludes determination of the stellar spin period based on starspot modulations.

However, if we assume that the stellar rotation axis is approximately perpendicular to the line of sight, then our measurement of $v \sin i_*$ combined with the stellar radius estimated above imply a rotation period for the primary star of $P_{\text{rot}} = 19 \pm 3$ days. Given that the binary orbit is eccentric, and that the timescale for synchronization (~ 2.5 Gyrs) is comparable to the estimated age (~ 4.4 Gyrs), we can expect the rotation period to be closer to the pseudosynchronous period (~ 12.7 days). If this star is magnetically active then we expect to see modulation of the light curve at frequencies $1/P_{\text{rot}}$ and/or $2/P_{\text{rot}}$, depending on the distribution of active regions on the stellar surface at the time of observation. To search for such modulations and periodic signals in the WASP light curve of TOI-1338, we used the sine-wave fitting method described in Maxted et al. (2011). The WASP light curve contains 26,492 observations obtained with the same CCD camera and 200-mm lens over 3 observing seasons. We calculated the




Figure 16. The extent of the system HZ, calculated based on the properties shown in Table 4. The conservative HZ is shown in dark green and the optimistic extension to the HZ is shown in light green. The known planet is substantially interior to the boundaries of the system HZ and close to the stability boundary.

periodogram over 32,768 uniformly spaced frequencies from 0 to 1.5 cycles/day. The false alarm probability (FAP) is calculated using a boot-strap Monte Carlo method also described in Maxted et al. (2011). The periodogram is shown in Fig. 18. From the boot-strap Monte Carlo simulations and the lack of any significant signal in this periodogram we can put an upper limit of approximately 1 mmag to the semi-amplitude of any signal due to rotational modulation. A similar analysis for the 3 seasons of WASP data separately is consistent with this conclusion.

To further this point, we also obtained historical data from the All-Sky Automated Survey for Supernovae (ASAS-SN) project (Shappee et al. 2014; Kochanek et al. 2017) in order to search for evidence of star spot modulation. Our analysis indicates that ~ 1600 days of ASAS-SN V-band data shows no significant photometric modulation either (Fig. 19). Additionally, the HARPS spectra have a dispersion of only a few m s^{-1} , compatible with a chromospherically quiet star. The bisectors of the cross-correlation function likewise show no variability (Queloz et al. 2001). These considerations further strengthen our assumption that the stellar activity and starspot-induced bias in the measured eclipse times is negligible, and that their effect on the photodynamical solution is minimal. If additional photometric observations should reveal the primary star to be heavily spotted, then the planet mass determination may need to be revised.

5.3. The Planet

With a mass of $33.0 \pm 20.0 M_{\oplus}$, a radius of $6.85 \pm 0.19 R_{\oplus}$, and a bulk density of $0.56 \pm 0.34 \text{ g cm}^{-3}$ (Table 4, where the quoted values are from the sample medians), the closest Solar System analogue of TOI-1338b is perhaps Saturn, where $M = 95.16 M_{\oplus}$, $R = 9.14 R_{\oplus}$, and $\rho = 0.69 \text{ g cm}^{-3}$. Among the known CBPs, TOI-1338b has bulk properties similar to those of Kepler-16b (Doyle et al. 2011) and Kepler-34b (Welsh et al. 2012).

The large radius of TOI-1338b is consistent with the predictions of planet-formation models in circumbinary disks, a trend that has also been observed among CBPs detected by the *Kepler* telescope. Combined with the small orbital inclination, this indicates that TOI-1338b formed at larger distances from its host binary and migrated to its current orbit through planet-disk interaction (e.g. Pierens & Nelson 2013; Kley & Haghighipour 2015).

5.4. TOI-1338 within the context of the Kepler CBP systems




Figure 17. Upper panel: Mass and radius of TOI-1338 compared against MIST isochrones. Dotted lines represent isochrones for ages between 2 and 6 Gyr in 1 Gyr increments, and the solid line is the best-fit isochrone corresponding to 4.4 Gyr. The inset shows an enlargement around the location of the secondary star, whose properties are seen to be consistent with the best-fit model. Lower panel: Same as upper panel but for mass and temperature.

The TOI-1338 system follows the trends established by *Kepler* CBPs. Namely, these are gas giant planets (radius larger than $3R_{\oplus}$) with low-eccentricity, nearly co-planar orbits¹² with periods longer than ~ 50 days, and orbit around binary stars with $P_{\text{bin}} \sim 7.5 - 40$ days (Welsh & Orosz 2018; Martin et al. 2019, and references therein)¹³. The near co-planarity of the planetary orbits is also consistent with the observational results of circumbinary disks around short period stellar binaries (Czekala et al. 2019). TOI-1338 is similar to the *Kepler*-38 CBP system (Orosz et al. 2012a) in terms of both the orbital periods and orbital period ratio (see Figure 20). The orbital precession timescale of TOI-1338b is comparable to that of the CBP *Kepler*-413b, where the time scale is ~ 22 years for the former compared to ~ 11 years for the latter Kostov et al. (2014).

Overall, TOI-1338b has a relatively long period for a transiting planet, particularly when compared with other TESS candidates. This is demonstrated in Figure 21, where it resides on the very tail of the TESS planet candidate period distribution. We note that the current lack of small CBPs is likely an observational bias, since unique challenges have inhibited their detection to date, largely as a result of the transit timing variations induced by the barycentric binary motion and the orbital dynamics (Armstrong et al. 2013, Armstrong et al. 2014, Martin 2019a, Windemuth et al. 2019). The circumbinary planet population is yet to be constrained below $4R_{\oplus}$ (Armstrong et al. 2014), and we expect that the large quantity and brightness of the TESS stars will enable the expansion of this parameter space.

¹² $e_{\text{CBP}} < 0.15$, mutual orbital inclination $\Delta i < 5^{\circ}$.

¹³ Whereas most of *Kepler*'s EBs have orbital periods shorter than ~ 3 days (Muñoz & Lai 2015; Martin et al. 2015; Hamers et al. 2016; Fleming et al. 2018).




Figure 18. Periodogram of the WASP lightcurve for TIC 260128333. The dashed line indicates a false-alarm probability level of 0.01.




Figure 19. Approximately 1600 days of ASAS-SN V-band photometry showing no significant rotationally-induced photometric modulation.

6. CONCLUSIONS

We presented the discovery of the first transiting circumbinary planet from TESS, TOI-1338. The target was observed by TESS in 12 sectors of 30-min cadence data (Sectors 1 through 12), and 9 sectors of 2-min cadence data (Sectors 4 through 12). In addition to stellar eclipses, three transit events were observed in Sectors 3, 6, and 10. These extra transit events show the hallmark characteristics of a circumbinary object where their duration depends on the binary phase and their times have significant deviations from a simple linear ephemeris. Blending is not an issue with TOI-1338 as the nearest source is $53''$ away, and speckle imaging observations from SOAR rule out nearby sources with a magnitude difference of $\Delta I \sim 4$ down to $0.5''$ from the target. Radial velocity measurements are available as




Figure 20. (a) Ratio of the planet and binary orbital periods for Kepler CBPs (blue squares) and TESS (purple downwards triangle). The diagonal black dashed lines indicate constant period ratios. (b) Corresponding ratio of the planet periastron and the binary apoastron, which together represent the shortest separation between the planet and the binary.




Figure 21. Radius and period of the TESS circumbinary planet (purple downwards triangle) and Kepler circumbinary planets (blue squares) compared with the TESS planet candidates (yellow upwards triangles), Kepler planet candidates (green circles) and the 5 innermost Solar System planets (red diamonds). This circumbinary discovery is the longest-period confirmed TESS planet to date, and exists in a parameter space similar to the Kepler circumbinary planets.

the host eclipsing binary has been monitored for more than three years by CORALIE and HARPS as part of the EBLM project. To solve for the parameters of the system, we combined the TESS data with the radial velocities into the photometric-dynamical model ELC. Our analysis confirms that the circumbinary object is indeed a planet, with a mass of $33.0 \pm 20.0 M_{\oplus}$, a radius of $6.85 \pm 0.19 R_{\oplus}$, and a bulk density of $0.56 \pm 0.34 \text{ g cm}^{-3}$. The planet’s orbit is within $\sim 1^{\circ}$ of being coplanar with the binary, has a period of 95.2 days and small eccentricity, and is safely beyond the boundary for stability. The host eclipsing binary (with $P = 14.6$ days and $e \approx 0.16$) consists of G+M stars with masses $1.1 M_{\odot}$ and $0.3 M_{\odot}$, and radii of $1.3 R_{\odot}$ and $0.3 R_{\odot}$ respectively. Based on the stellar parameters, we estimate an age of 4.4 Gyr for the system.

This manuscript includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA Science Mission directorate. We acknowledge the use of public TESS data from pipeline at the TESS Science Processing Operations Center. The manuscript includes data from CORALIE, an instrument mounted on the Euler 1.2m telescope, a project of the University of Geneva, funded by the Swiss National Science Foundation. Furthermore, our analysis includes spectra obtained with HARPS, an instrument mounted on the ESO-3.6m telescope at La Silla. Those data were obtained under Prog.ID 1101.C-0721 (PI Triaud). They are or will become available through the ESO public archive. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center for the production of the SPOC data products. This research was supported in part through research cyberinfrastructure resources and services provided by the Partnership for an Advanced Computing Environment (PACE) at the Georgia Institute of Technology. This research made use of Lightkurve, a Python package for Kepler and TESS data analysis (Lightkurve Collaboration, 2018). This research has made use of the Exoplanet Follow-up Observation Program website, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program. We are also grateful to the observer support staff at CTIO, at ESO/HARPS and Swiss Euler Telescope/CORALIE. This work has made use of data from the European Space Agency (ESA) mission Gaia (<https://www.cosmos.esa.int/gaia>), processed by the Gaia Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement. WFW and JAO thank John Hood, Jr. for his generous support of exoplanet research at SDSU. Support was also provided and acknowledged through NASA Habitable Worlds grant 80NSSC17K0741 and NASA XRP grant 80NSSC18K0519. This work is partly supported by NASA Habitable Worlds grant 80NSSC17K0741. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. (DGE-1746045). AHMJT has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement n° 803193/BEBOP), and from a Leverhulme Trust Research Project Grant n° RPG-2018-418. AC acknowledges support by CFisUC strategic project (UID/FIS/04564/2019). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. E.A.G. thanks the LSSTC Data Science Fellowship Program, which is funded by LSSTC, NSF Cybertraining Grant #1829740, the Brinson Foundation, and the Moore Foundation; her participation in the program has benefited this work. E.A.G. is thankful for support from GSFC Sellers Exoplanet Environments Collaboration (SEEC), which is funded by the NASA Planetary Science Divisions Internal Scientist Funding Model.

Facilities: TESS; SOAR 4.0-m; WASP; Swiss Euler Telescope (CORALIE); ESO 3.6-m (HARPS)

Software: astropy (Astropy Collaboration et al. 2013); scipy (Virtanen et al. 2019); eleanor (Feinstein et al. 2019); Lightkurve (Lightkurve Collaboration et al. 2018); ispec (Blanco-Cuaresma et al. 2014); REBOUND (Rein & Liu 2012; Rein & Spiegel 2015); Mercury6 (Chambers et al. 2002)

Table 1. Observed eclipse times for TOI-1338

Cycle	Observed Time ^a	Model Time ^a	O-C ^b	Cycle	Observed Time ^a	Model Time ^a	O-C ^b
Primary				Secondary			
0	...	3322.21319	...	0.45345	3328.83160 ± 0.00557	3328.83833	-9.71
1	3336.82137 ± 0.00027	3336.82171	-0.48	1.45345	3343.44848 ± 0.00970	3343.44687	2.32
2	3351.43036 ± 0.00026	3351.43028	0.12	2.45345	3358.05700 ± 0.00285	3358.05537	2.34
3	3366.03902 ± 0.00023	3366.03888	0.21	3.45345	3372.65843 ± 0.00410	3372.66390	-7.87
4	3380.64698 ± 0.00059	3380.64735	-0.52	4.45456	3387.27302 ± 0.00379	3387.27251	0.73
5	3395.25589 ± 0.00044	3395.25598	-0.12	5.45345	3401.86970 ± 0.01653	3401.88102	-16.29
6	...	3409.86452	...	6.45345	3416.49508 ± 0.00375	3416.48957	7.93
7	...	3424.47302	...	7.45345	3431.09816 ± 0.00340	3431.09811	0.06
8	3439.08165 ± 0.00020	3439.08157	0.13	8.45345	3445.70014 ± 0.00459	3445.70662	-9.34
9	3453.69010 ± 0.00019	3453.69017	-0.09	9.45345	3460.32094 ± 0.00364	3460.31513	8.36
10	...	3468.29871	...	10.45345	3474.92301 ± 0.00324	3474.92372	-1.02
11	3482.90704 ± 0.00022	3482.90721	-0.24	11.45345	3489.53315 ± 0.00300	3489.53225	1.30
12	3497.51571 ± 0.00021	3497.51582	-0.15	12.45345	...	3504.14078	...
13	3512.12410 ± 0.00022	3512.12432	-0.31	13.45345	3518.73613 ± 0.00491	3518.74934	-19.02
14	3526.73304 ± 0.00019	3526.73285	0.29	14.45345	...	3533.35786	...
15	3541.34140 ± 0.00022	3541.34142	-0.02	15.45345	3547.97016 ± 0.00444	3547.96636	5.48
16	...	3555.95001	...	16.45345	3562.57901 ± 0.00349	3562.57490	5.91
17	...	3570.55847	...	17.45345	3577.18768 ± 0.00349	3577.18350	6.02
18	3585.16712 ± 0.00032	3585.16712	0.02	18.45345	3591.78979 ± 0.00254	3591.79201	-3.19
19	...	3599.77564	...	19.45345	3606.40434 ± 0.00371	3606.40057	5.42
20	3614.38474 ± 0.00030	3614.38415	0.85	20.45345	3621.01281 ± 0.00399	3621.00911	5.32
21	3628.99361 ± 0.00038	3628.99271	1.31	21.45345	3635.61695 ± 0.00322	3635.61762	-0.97
22	3643.60114 ± 0.00036	3643.60131	-0.23	22.45345	3650.22268 ± 0.00518	3650.22613	-4.97

^aBJD - 2,455,000^bObserved time minus model time in minutes.

Table 2. Photometric and spectroscopic parameters of the system prior to the photo-dynamical solution

Parameter	Value	Uncertainty	Unit	Source
RV semi-amplitude, K_1	21.619	0.007	km s ⁻¹	Martin et al. (2019)
Gravity of Primary, $\log g_1$	4.0	0.08	cgs	Spectroscopy, this work
Metallicity of Primary, $[\text{Fe}/\text{H}]_1$	0.01	0.05	dex	Spectroscopy, this work
Projected Rotational Velocity of Primary, $v \sin i$	3.6	0.6	km s ⁻¹	Spectroscopy, this work
Reddening, $E(\text{B}-\text{V})$	0.02	0.01	mag	Gaia + Photometry, this work
Effective Temperature of Primary, $T_{\text{eff},1}$	6050	80	K	Spectroscopy, this work
Effective Temperature of Primary, $T_{\text{eff},1}$	5990	110	K	Gaia + Photometry, this work
Radius of Primary, R_1	1.345	0.046	R_\odot	Gaia+Photometry, this work
Age	4.4	0.2	Gyr	This work

Table 3. Fitted Parameters for TOI-1338

Parameter ^a	Best	Mode	Median	+1 σ	-1 σ
$T_{\text{conj,bin}}$	3336.8245	3336.8237	3336.8242	0.0025	0.0023
P_{bin} (d)	14.608561	14.608559	14.608559	0.000013	0.000012
$\sqrt{e_{\text{bin}}} \cos \omega_{\text{bin}}$	-0.18275	-0.18270	-0.18272	0.00040	0.00040
$\sqrt{e_{\text{bin}}} \sin \omega_{\text{bin}}$	0.35015	0.35008	0.35020	0.00036	0.00035
M_1 (M_\odot)	1.038	1.149	1.127	0.068	0.069
$Q \equiv M_2/M_1$	0.2865	0.2752	0.2774	0.0069	0.0062
i (deg)	89.658	89.649	89.696	0.178	0.114
$T_{\text{eff},1}$ (K)	5990.7	6072.4	6040.8	98.1	91.5
$T_{\text{eff},2}/T_{\text{eff},1}$	0.5537	0.5548	0.5516	0.0042	0.0047
R_1 (R_\odot)	1.299	1.338	1.331	0.024	0.026
R_1/R_2	4.307	4.310	4.308	0.013	0.014
$q_{1,1}$	0.255	0.245	0.255	0.048	0.043
$q_{2,1}$	0.308	0.299	0.310	0.058	0.050
ζ_p (d)	3436.57	3437.26	3437.32	0.83	0.78
ζ_m (d)	-3246.29	-3246.84	-3246.97	0.71	0.77
$\sqrt{e_{\text{pl}}} \cos \omega_{\text{pl}}$	-0.035	-0.051	-0.058	0.018	0.024
$\sqrt{e_{\text{pl}}} \sin \omega_{\text{pl}}$	-0.303	-0.285	-0.290	0.009	0.011
i_{pl} (deg)	89.22	89.29	89.37	0.35	0.00
Ω_{pl} (deg)	0.87	0.83	0.91	0.35	0.35
M_3 (M_\oplus)	30.2	33.2	33.0	20.3	19.6
R_1/R_3	21.12	21.20	21.17	0.47	0.44
γ_1^b (km s ⁻¹)	30.74769	30.74766	30.74778	0.00089	0.00084
γ_2^c (km s ⁻¹)	30.74621	30.74618	30.74641	0.00128	0.00125
γ_3^d (km s ⁻¹)	30.75866	30.75836	30.75857	0.00098	0.00098

^aOsculating parameters valid at BJD 2,455,186.0000.^bRelative velocity offset, “early” CORALIE data.^cRelative velocity offset, “late” CORALIE data.^dRelative velocity offset, HARPS data.

Table 4. Derived Parameters for TOI-1338

Parameter ^a	Best	Mode	Median	+1 σ	-1 σ
Bulk Properties					
M_1 (M_\odot)	1.038	1.149	1.127	0.068	0.069
R_1 (R_\odot)	1.299	1.338	1.331	0.024	0.026
M_2 (M_\odot)	0.2974	0.3168	0.3128	0.0113	0.0118
R_2 (R_\odot)	0.3015	0.3102	0.3089	0.0056	0.0060
M_3 (M_\oplus)	30.2	33.2	33.0	20.3	19.6
R_3 (R_\oplus)	6.71	6.83	6.85	0.19	0.19
ρ_3 (g cm^{-3})	0.55	0.57	0.56	0.34	0.33
Binary Orbit					
P_{bin} (d)	14.608561	14.608559	14.608559	0.000013	0.000012
$T_{\text{conj,bin}}$	3336.8245	3336.8237	3336.8242	0.0025	0.0023
K_{bin} (km s^{-1})	21.6248	21.6246	21.6247	0.0034	0.0032
e_{bin}	0.15601	0.15603	0.15603	0.00015	0.00015
ω_{bin} (deg)	117.561	117.568	117.554	0.072	0.074
a_{bin} (AU)	0.1288	0.1319	0.1321	0.0024	0.0025
true anomaly (deg)	111.217	111.246	111.226	0.071	0.069
mean anomaly (deg)	93.882	93.897	93.889	0.065	0.065
mean longitude (deg)	228.779	228.783	228.779	0.020	0.020
i_{bin} (deg)	89.658	89.649	89.696	0.178	0.114
Ω_{bin} (deg)	0.0	0.0	0.0	0.0	0.0
Planet Orbit					
P_{pl} (d)	95.141	95.175	95.174	0.031	0.035
$T_{\text{conj,pl}}$	3341.43	3342.11	3342.15	0.80	0.74
e_{pl}	0.0928	0.0861	0.0880	0.0043	0.0033
ω_{pl} (deg)	263.3	260.3	258.6	3.7	4.8
a_{pl} (AU)	0.4491	0.4639	0.4607	0.0084	0.0088
true anomaly (deg)	136.0	141.0	141.7	5.2	4.5
mean anomaly (deg)	128.3	134.1	135.2	6.1	5.4
mean longitude (deg)	400.2	401.6	401.2	0.9	1.1
i_{pl} (deg)	89.22	89.29	89.37	0.35	0.26
Ω_{pl} (deg)	0.87	0.83	0.91	0.35	0.35
I^b (deg)	0.98	0.92	0.99	0.31	0.28

^aOsculating parameters valid at BJD 2,455,186.0000.

^bMutual inclination between orbital planes.

REFERENCES

- Allard, F., Homeier, D., & Freytag, B. 2012, in IAU Symposium, Vol. 282, From Interacting Binaries to Exoplanets: Essential Modeling Tools, ed. M. T. Richards & I. Hubeny, 235–242
- Andrade-Ines, E., & Robutel, P. 2018, *Celestial Mechanics and Dynamical Astronomy*, 130, 6
- Armstrong, D. J., Osborn, H. P., Brown, D. J. A., et al. 2014, *MNRAS*, 444, 1873
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Baranne, A., Queloz, D., Mayor, M., et al. 1996, *A&AS*, 119, 373
- Blanco-Cuaresma, S., Soubiran, C., Heiter, U., & Jofré, P. 2014, *A&A*, 569, A111

Table 5. Initial Dynamical Parameters^a

parameter ^b	binary orbit	planet orbit	
Period (days)	1.46085607280931704E + 01	9.51407742682822573E + 01	
$e \cos \omega$	-7.21833507304852212E - 02	-1.07723606253091915E - 02	
$e \sin \omega$	1.38301268444568498E - 01	-9.22020142859296676E - 02	
i (rad)	1.56482063218343082E + 00	1.55715854420219579E + 00	
Ω (rad)	0.00000000000000000E + 00	1.52632127894949884E - 02	
T_{conj} (days) ^c	3.33682449014051008E + 03	3.34142761690665247E + 03	
a (AU)	1.28783121829547487E - 01	4.49132971740733966E - 01	
e	1.56005374830665650E - 01	9.28291720948987292E - 02	
ω (deg)	1.17561331987513597E + 02	2.63336097695842795E + 02	
true anomaly (deg)	1.11217395295404202E + 02	1.36038415640448932E + 02	
mean anomaly (deg)	9.38816596366525289E + 01	1.28272133285056185E + 02	
mean longitude (deg)	2.28778727282917799E + 02	4.00249031010939916E + 02	
i (deg)	8.96576179191039415E + 01	8.92186126155212662E + 01	
Ω (deg)	0.00000000000000000E + 00	8.74517674648163101E - 01	

parameter ^d	body 1	body 2	body 3
Mass (M_{\odot})	1.03784970719363567E + 00	2.97388770751337850E - 01	9.06017229632760055E - 05
x (AU)	1.95196590778876217E - 02	-6.82335050435495666E - 02	3.68709659800795730E - 01
y (AU)	1.32648896371446897E - 04	-4.65900424342245798E - 04	9.75628498103004414E - 03
z (AU)	2.22880532978689296E - 02	-7.78747286204819755E - 02	3.02645766911772141E - 01
v_x (AU day ⁻¹)	-7.66586168788931707E - 03	2.67578221143398472E - 02	-1.61533175726252566E - 02
v_y (AU day ⁻¹)	5.45640776560390306E - 05	-1.90441050119945319E - 04	6.31089229491667655E - 05
v_z (AU day ⁻¹)	9.13006908476301365E - 03	-3.18697187907663951E - 02	2.27034214206392367E - 02

^aReference time = 186.00000, integration step size = 0.05000 days

^bJacobian instantaneous (Keplerian) elements

^cTimes are relative to BJD 2,455,000.000

^dBarycentric Cartesian coordinates

Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, in IAU Symposium, Vol. 135, Interstellar Dust, ed. L. J. Allamandola & A. G. G. M. Tielens, 5–10

Casagrande, L., Ramírez, I., Meléndez, J., Bessell, M., & Asplund, M. 2010, *A&A*, 512, A54

Chambers, J. E., Quintana, E. V., Duncan, M. J., & Lissauer, J. J. 2002, *AJ*, 123, 2884

Charbonneau, P. 1995, *ApJS*, 101, 309

Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102

Collier Cameron, A., Wilson, D. M., West, R. G., et al. 2007, *MNRAS*, 380, 1230

Czekala, I., Chiang, E., Andrews, S. M., et al. 2019, *ApJ*, 883, 22

Doyle, A. P. 2015, PhD thesis, Keele University

Doyle, L. R., Carter, J. A., Fabrycky, D. C., et al. 2011, *Science*, 333, 1602

Dvorak, R. 1986, *A&A*, 167, 379

Dvorak, R., Pilat-Lohinger, E., Schwarz, R., & Freistetter, F. 2004, *A&A*, 426, L37

Eastman, J. D., Rodriguez, J. E., Agol, E., et al. 2019, arXiv e-prints, arXiv:1907.09480

Farago, F., Laskar, J., & Couetdic, J. 2009, *Celestial Mechanics and Dynamical Astronomy*, 104, 291

Feinstein, A. D., Montet, B. T., Foreman-Mackey, D., et al. 2019, *PASP*, 131, 094502

Fleming, D. P., Barnes, R., Graham, D. E., Luger, R., & Quinn, T. R. 2018, *ApJ*, 858, 86

Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1

Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1

Gray, R. O., & Corbally, C. J. 1994, *AJ*, 107, 742

Gustafsson, B., Edvardsson, B., Eriksson, K., et al. 2008, *A&A*, 486, 951

Table 6. Times, durations, and impact parameters of future planet transits

BJD - 2,455,000	Year	Month	Day	UTC	Impact parameter	Duration (hr)	Transit fraction
3861.1845 ± 0.0161	2020	Jan	12	16:25:40.7	0.143 ± 0.128	6.74 ± 0.16	100.0%
3954.1353 ± 0.0214	2020	Apr	14	15:14:50.2	0.206 ± 0.108	9.80 ± 0.23	100.0%
4049.6930 ± 0.0196	2020	Jul	19	04:37:53.2	-0.006 ± 0.189	7.91 ± 0.25	100.0%
4142.3991 ± 0.0224	2020	Oct	19	21:34:40.9	0.055 ± 0.166	7.46 ± 0.18	100.0%
4237.9220 ± 0.0236	2021	Jan	23	10:07:40.6	-0.131 ± 0.236	9.73 ± 0.63	99.9%
4330.9984 ± 0.0264	2021	Apr	26	11:57:42.7	-0.128 ± 0.232	6.31 ± 0.38	99.9%
4425.7194 ± 0.0319	2021	Jul	30	05:15:54.7	-0.224 ± 0.266	11.78 ± 1.20	99.6%
4519.6951 ± 0.0328	2021	Nov	1	04:40:55.7	-0.315 ± 0.285	5.94 ± 0.79	98.7%
4708.3278 ± 0.0389	2022	May	8	19:52:05.0	-0.468 ± 0.308	6.06 ± 1.27	92.9%
4801.0174 ± 0.0461	2022	Aug	9	12:24:59.3	-0.417 ± 0.304	8.44 ± 1.58	95.5%
4896.7585 ± 0.0461	2022	Nov	13	06:12:14.7	-0.566 ± 0.310	6.78 ± 1.72	82.4%
4989.3540 ± 0.0509	2023	Feb	13	20:29:49.0	-0.540 ± 0.313	6.04 ± 1.46	85.2%
5084.8165 ± 0.0693	2023	May	20	07:35:48.3	-0.624 ± 0.310	8.19 ± 2.34	72.6%
5177.9790 ± 0.0600	2023	Aug	21	11:29:44.2	-0.633 ± 0.317	4.93 ± 1.45	70.0%
5272.3566 ± 0.1296	2023	Nov	23	20:33:30.0	-0.650 ± 0.311	9.79 ± 2.97	64.5%
5366.6770 ± 0.0742	2024	Feb	26	04:14:52.4	-0.670 ± 0.318	4.79 ± 1.52	54.2%
5459.7304 ± 0.1788	2024	May	29	05:31:47.0	-0.665 ± 0.315	8.76 ± 2.76	55.7%
5555.2836 ± 0.0967	2024	Sep	1	18:48:23.7	-0.690 ± 0.320	5.26 ± 1.77	42.9%
5647.6474 ± 0.1677	2024	Dec	3	03:32:17.7	-0.687 ± 0.322	6.27 ± 2.08	44.5%
5743.6224 ± 0.1383	2025	Mar	9	02:56:12.7	-0.707 ± 0.320	6.35 ± 2.21	35.5%
5836.0445 ± 0.1543	2025	Jun	9	13:04:05.3	-0.700 ± 0.329	4.93 ± 1.67	34.2%
5931.4246 ± 0.2222	2025	Sep	12	22:11:25.1	-0.713 ± 0.320	8.39 ± 3.01	31.0%
6024.6162 ± 0.1615	2025	Dec	15	02:47:16.6	-0.694 ± 0.333	4.71 ± 1.64	27.7%
6118.5043 ± 0.3348	2026	Mar	19	00:06:09.6	-0.702 ± 0.325	9.86 ± 3.45	27.4%
6213.1325 ± 0.1991	2026	Jun	21	15:10:46.2	-0.687 ± 0.333	5.22 ± 1.81	23.9%
6305.7169 ± 0.3200	2026	Sep	22	05:12:19.8	-0.698 ± 0.333	7.40 ± 2.63	23.8%
6401.3824 ± 0.2847	2026	Dec	26	21:10:36.5	-0.685 ± 0.332	6.52 ± 2.34	22.4%
6493.7147 ± 0.2750	2027	Mar	29	05:09:06.7	-0.673 ± 0.340	5.53 ± 1.88	20.7%
6589.0498 ± 0.4693	2027	Jul	2	13:11:42.7	-0.678 ± 0.329	8.97 ± 3.27	21.5%
6682.0668 ± 0.2875	2027	Oct	3	13:36:11.6	-0.665 ± 0.342	5.11 ± 1.70	19.6%
6775.8467 ± 0.7051	2028	Jan	5	08:19:17.2	-0.675 ± 0.332	10.95 ± 3.69	20.4%
6870.3964 ± 0.3605	2028	Apr	8	21:30:48.3	-0.672 ± 0.345	5.65 ± 1.98	20.7%
6962.8655 ± 0.5975	2028	Jul	10	08:46:15.3	-0.666 ± 0.340	7.74 ± 2.61	19.6%
7058.3711 ± 0.5322	2028	Oct	13	20:54:25.6	-0.672 ± 0.341	7.42 ± 2.70	23.1%
7150.8050 ± 0.4580	2029	Jan	14	07:19:14.6	-0.661 ± 0.351	5.64 ± 1.88	21.1%
7245.5142 ± 0.8968	2029	Apr	19	00:20:24.3	-0.682 ± 0.341	10.53 ± 4.07	26.0%
7339.0599 ± 0.4556	2029	Jul	21	13:26:12.1	-0.664 ± 0.361	5.33 ± 1.87	25.6%
7431.8295 ± 1.0387	2029	Oct	22	07:54:30.9	-0.685 ± 0.351	10.20 ± 3.65	27.5%
7527.1701 ± 0.5751	2030	Jan	25	16:04:55.6	-0.696 ± 0.361	6.19 ± 2.34	36.4%

- Hairer, E., Lubich, C., & Wanner, G. 2002, Springer Series in Computational Mathematics, Vol. 31, Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations (Berlin: Springer)
- Hamers, A. S., Perets, H. B., & Portegies Zwart, S. F. 2016, MNRAS, 455, 3180
- Holman, M. J., & Wiegert, P. A. 1999, AJ, 117, 621
- Huang, C. X., Burt, J., Vanderburg, A., et al. 2018, ApJL, 868, L39
- Huang, Y., Liu, X. W., Yuan, H. B., et al. 2015, MNRAS, 454, 2863
- Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9913, Proc. SPIE, 99133E
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, Icarus, 101, 108
- Kipping, D. M. 2013, MNRAS, 435, 2152
- Kirk, B., Conroy, K., Prša, A., et al. 2016, AJ, 151, 68
- Kley, W., & Haghighipour, N. 2015, A&A, 581, A20
- Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, PASP, 129, 104502
- Konacki, M., Muterspaugh, M. W., Kulkarni, S. R., & Helminiak, K. G. 2009, ApJ, 704, 513
- Kostov, V. B., McCullough, P. R., Hinse, T. C., et al. 2013, ApJ, 770, 52
- Kostov, V. B., McCullough, P. R., Carter, J. A., et al. 2014, ApJ, 784, 14
- Kostov, V. B., Orosz, J. A., Welsh, W. F., et al. 2016, ApJ, 827, 86
- Kostov, V. B., Schlieder, J. E., Barclay, T., et al. 2019, AJ, 158, 32
- Lam, C., & Kipping, D. 2018, MNRAS, 476, 5692
- Laskar, J. 1993, Celestial Mechanics and Dynamical Astronomy, 56, 191
- Lee, M. H., & Peale, S. J. 2006, Icarus, 184, 573
- Li, G., Holman, M. J., & Tao, M. 2016, ApJ, 831, 96
- Li, J., Tenenbaum, P., Twicken, J. D., et al. 2019, PASP, 131, 024506
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, Lightkurve: Kepler and TESS time series analysis in Python, Astrophysics Source Code Library, , , ascl:1812.013
- López-Morales, M., Triaud, A. H. M. J., Rodler, F., et al. 2014, ApJL, 792, L31
- Lovis, C., & Pepe, F. 2007, A&A, 468, 1115
- Mardling, R. A., & Lin, D. N. C. 2002, ApJ, 573, 829
- Martin, D. V. 2017, MNRAS, 467, 1694
- Martin, D. V., Mazeh, T., & Fabrycky, D. C. 2015, MNRAS, 453, 3554
- Martin, D. V., & Triaud, A. H. M. J. 2014, A&A, 570, A91
- Martin, D. V., Triaud, A. H. M. J., Udry, S., et al. 2019, A&A, 624, A68
- Maxted, P. F. L., Anderson, D. R., Collier Cameron, A., et al. 2011, PASP, 123, 547
- Mayor, M., Udry, S., Lovis, C., et al. 2009, A&A, 493, 639
- Muñoz, D. J., & Lai, D. 2015, Proceedings of the National Academy of Science, 112, 9264
- Mudryk, L. R., & Wu, Y. 2006, ApJ, 639, 423
- Müller, T. W. A., & Haghighipour, N. 2014, ApJ, 782, 26
- Orosz, J. A., & Hauschildt, P. H. 2000, A&A, 364, 265
- Orosz, J. A., Welsh, W. F., Carter, J. A., et al. 2012a, ApJ, 758, 87
- . 2012b, Science, 337, 1511
- Orosz, J. A., Welsh, W. F., Haghighipour, N., et al. 2019, AJ, 157, 174
- Pepe, F., Mayor, M., Rupprecht, G., et al. 2002, The Messenger, 110, 9
- Pierens, A., & Nelson, R. P. 2013, A&A, 556, A134
- Pollacco, D. L., Skillen, I., Collier Cameron, A., et al. 2006, PASP, 118, 1407
- Prša, A., Batalha, N., Slawson, R. W., et al. 2011, AJ, 141, 83
- Quarles, B., Satyal, S., Kostov, V., Kaib, N., & Haghighipour, N. 2018, ApJ, 856, 150
- Queloz, D., Henry, G. W., Sivan, J. P., et al. 2001, A&A, 379, 279
- Ramos, X. S., Correa-Otto, J. A., & Beaugé, C. 2015, Celestial Mechanics and Dynamical Astronomy, 123, 453
- Rein, H., & Liu, S.-F. 2012, A&A, 537, A128
- Rein, H., & Spiegel, D. S. 2015, MNRAS, 446, 1424
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003
- Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103
- Schneider, J. 1994, Planet. Space Sci., 42, 539
- Schwamb, M. E., Orosz, J. A., Carter, J. A., et al. 2013, ApJ, 768, 127
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, ApJ, 788, 48
- Short, D. R., Orosz, J. A., Windmiller, G., & Welsh, W. F. 2018, AJ, 156, 297
- Slawson, R. W., Prša, A., Welsh, W. F., et al. 2011, AJ, 142, 160
- Socia, Q. J., Welsh, W. F., Orosz, J. A., et al. 2020, arXiv e-prints, arXiv:2001.02840
- Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, AJ, 158, 138
- Sutherland, A. P., & Fabrycky, D. C. 2016, ApJ, 818, 6
- Sutherland, A. P., & Kratter, K. M. 2019, MNRAS, 487, 3288

- Ter Braak, C. J. F. 2006, *Statistics and Computing*, 16, 239
- Tokovinin, A., Mason, B. D., Hartkopf, W. I., Mendez, R. A., & Horch, E. P. 2018, *AJ*, 155, 235
- Tokovinin, A., Mason, B. D., Mendez, R. A., Horch, E. P., & Briceño, C. 2019, *AJ*, 158, 48
- Triaud, A. H. M. J. 2011, PhD thesis, Observatoire Astronomique de l'Université de Genève
- Triaud, A. H. M. J., Hebb, L., Anderson, D. R., et al. 2013, *A&A*, 549, A18
- Triaud, A. H. M. J., Martin, D. V., Ségransan, D., et al. 2017a, *A&A*, 608, A129
- Triaud, A. H. M. J., Neveu-VanMalle, M., Lendl, M., et al. 2017b, *MNRAS*, 467, 1714
- Twicken, J. D., Catanzarite, J. H., Clarke, B. D., et al. 2018, *PASP*, 130, 064502
- Vanderspek, R., Huang, C. X., Vanderburg, A., et al. 2019, *ApJL*, 871, L24
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2019, arXiv e-prints, arXiv:1907.10121
- von Boetticher, A., Triaud, A. H. M. J., Queloz, D., et al. 2019, *A&A*, 625, A150
- Welsh, W. F., & Orosz, J. A. 2018, *Two Suns in the Sky: The Kepler Circumbinary Planets*, 34
- Welsh, W. F., Orosz, J. A., Carter, J. A., et al. 2012, *Nature*, 481, 475
- Welsh, W. F., Orosz, J. A., Short, D. R., et al. 2015, *ApJ*, 809, 26