TOLERANCE RELATIONS AND QUANTIZATION

FRANCESCO D'ANDREA, GIOVANNI LANDI, FEDELE LIZZI

ABSTRACT. It is well known that "bad" quotient spaces (typically: non-Hausdorff) can be studied by associating to them the groupoid C*-algebra of an equivalence relation, that in the "nice" cases is Morita equivalent to the C*-algebra of continuous functions vanishing at infinity on the quotient space. It was recently proposed in [8] that a similar procedure for relations that are reflexive and symmetric but fail to be transitive (i.e. *tolerance relations*) leads to an operator system. In this paper we observe that such an operator system carries a natural product that, although in general non-associative, arises in a number of relevant examples. We relate this product to truncations of (C*algebras of) topological spaces, in the spirit of [10], discuss some geometric aspects and a connection with positive operator valued measures.

CONTENTS

1. Introduction	1
2. Tolerance relations	2
3. From relations to algebras	4
3.1. Finite-dimensional case	5
3.2. Magmas acting on sets	7
3.3. Truncations	8
4. State spaces	11
4.1. Operator systems	11
4.2. Positivity in a tolerance algebra	13
4.3. Pure states of a tolerance algebra	15
5. Positive operator valued measures: a curious example	19
References	22

1. INTRODUCTION

A standard construction in Noncommutative Geometry is that of the C*-algebra associated to an equivalence relation R on a locally compact Hausdorff space X: one can construct the groupoid \mathcal{G} of the relation R and, under some technical conditions — for example if \mathcal{G} is étale —, one can construct the convolution algebra of \mathcal{G} and complete it to a C*-algebra (see e.g. [31]). A source of examples comes from group actions: if G is a discrete group with a continuous action on X, the action groupoid (corresponding to the relation "being on the same orbit") is étale; if X is compact, the full groupoid C*algebra is isomorphic to the crossed product $C(X) \rtimes G$; if, in addition, the action is free and proper — which means that X/G is a compact Hausdorff space, and even a smooth manifold if X is a smooth manifold and the action is smooth — then the above crossed product is strongly Morita equivalent to the commutative C*-algebra C(X/G) (this is a

Date: May 2022.

²⁰²⁰ Mathematics Subject Classification. Primary: 58B34; Secondary: 17A99; 81R60.

Key words and phrases. Tolerance relations, non-associative quantization, operator systems.

```
eps = 50*$MachineEpsilon;
a = 1;
b = a + eps;
c = b + eps;
{a == a, b == b, c == c, a == b, b == a, b == c, c == b, a == c, c == a}
(* Out: {True, True, True, True, True, True, False, False} *)
```

TABLE 1. Equality in Wolfram Mathematica.

special case of, e.g., "situation 2" in [27]). Applications of this construction cover foliations, orbifolds, tilings of the plane, dynamical systems arising in number theory such as the Bost-Connes system, just to name a few [5, 6]. Notice that associativity of the groupoid partial composition law, and then of the convolution product, is a consequence of transitivity of R.

It is natural to wonder how much of the above picture can be generalized to relations R that are reflexive and symmetric but not transitive, i.e. those that in the literature are called "tolerance relations". It was observed by A. Connes and W.D. van Suijlekom in [7] that from any tolerance relation one can get an *operator system*, which they propose as a starting point for a generalization of one of the main ingredients of Noncommutative Geometry, namely *spectral triples* [5, 19, 16]. The connection between tolerance relations and operator systems was then developed in [8]. It turns out that these operator systems possess, in fact, the additional structure of an algebra. However, in general, this algebra would not be associative.

The aim of this paper is to study the convolution algebra of a tolerance relation. We start, in §2, with a short introduction to tolerance relations and present a few motivating examples. In §3 we discuss the construction of convolution algebras of tolerance relations: we study finite-dimensional examples, infinite-dimensional ones associated to étale tolerance relations, and special actions of magmas on sets replacing, in this picture, the above-mentioned action groupoids. We will also relate this construction to truncations of (C*-algebras of) topological spaces, in the spirit of [10]. In §4 we focus on state spaces and discuss some geometric aspects. In particular, we will give an explicit description of the set of pure states of the finite-dimensional systems from §3. Finally, §5 is devoted to a curious example of a positive operator valued measure (POVM) that comes from a natural tolerance relation on the circle.

2. Tolerance relations

It was Poincaré who first observed, in a discussion about mathematical and physical continuum, that "equality" in the real world is not always transitive, due to potential measurement errors [25, Chap. 2]. A beautiful concrete example with Wolfram Mathematica was suggested in [21]. The code is reported in Table 1: here we define $b := a + \varepsilon$ and $c := b + \varepsilon$ with ε sufficiently small. Due to the finite storage capacity for numbers, the computer considers a = b and b = c, but recognizes that $a \neq c$. This is exactly the behaviour one would expect in a physical experiment when using an instrument with finite resolution.

A relation that is reflexive and symmetric, but not necessarily transitive, is called a *tolerance relation*. The name was coined in [37], while a mathematical theory was later developed in [33].




FIGURE 1. Causality is not transitive.

The one in Table 1 is an example of "proximity" relation: on a metric space (X, d) we can fix an $\varepsilon > 0$ and consider the relation $R \subset X \times X$ defined by

$$(x,y) \in R \iff d(x,y) < \varepsilon.$$
(1)

Such a relation in general fails to be transitive, except in special cases: for example if d is an ultrametric (which means $d(x, z) \leq \max\{d(x, y), d(y, z)\} \forall x, y, z \in X$), then (1) is an equivalence relation.

Another important example is the tolerance relation associated to a cover. Let X be a set and \mathcal{U} a collection of subsets covering X. The relation $R \subset X \times X$ given by

$$(x,y) \in R \iff \{ \exists S \in \mathcal{U} : x, y \in S \}$$

$$(2)$$

is clearly a tolerance relation. From a physical point of view, we can imagine that this models an experiment where two points contained in the same set of the cover are not distinguishable one from the other. For $X = \mathbb{R}^n$ with Euclidean distance, the example (1) corresponds to the cover of \mathbb{R}^n by open balls of diameter ε .

It is worth mentioning that the idea of using finite covers to approximate a topological space is not new: it can be traced back to Sorkin and his work on posets [32], and to [2, 20] in the framework of Noncommutative Geometry.

A notable example of tolerance relation in physics is *causality* in special relativity: the set of pairs (x, y) of points in Minkowski space that are causally connected, that means x - y is either timelike or lightlike, is reflexive and symmetric but not transitive. This is illustrated pictorially in Figure 1, where $(x, y), (y, z) \in R$ and $(x, z) \notin R$.

Finally, we mentioned that a source of examples of equivalence relations is given by group actions on sets. Given an action $\alpha : G \times X \to X$ of a group G on a set X, the property

$$\alpha_g(\alpha_h(x)) = \alpha_{gh}(x) \quad \forall \ g, h \in G, x \in X, \tag{3}$$

ensures that the relation $R := \{(x, \alpha_g x) : g \in G, x \in X\}$ on X is transitive. There are a number of examples, however, where the condition (3) is not satisfied.

Suppose, for example, that we have a morphism from a group G to the group

$$\operatorname{Out}(A) := \operatorname{Aut}(A) / \operatorname{Inn}(A)$$

of outer automorphisms of some unital C*-algebra A. In the Noncommutative Geometry approach to particle physics, when A is almost commutative, this corresponds to an action of G as diffeomorphisms of the underlying manifold (cf. [6, 35] and references therein, or the recent physically-oriented review [11]). In such a situation, we can lift the above map to a map $\alpha : G \to \operatorname{Aut}(A)$ by picking up a representative for each class in $\operatorname{Out}(A)$. Such a map α , however, will be only an action modulo inner automorphism. If A has only trivial inner automorphisms, given by the adjoint action of the unitary group U(A), then

$$\alpha_g \circ \alpha_h = \operatorname{Ad}_{f(q,h)} \circ \alpha_{gh} \tag{4}$$

for all $g, h \in G$ and for a suitable function $f : G \times G \to U(A)$. This reduces to (3) if f has image in the center of A, and was one of the motivating examples in [4] to develop a theory of non-associative crossed products.

Another class of examples where the condition (3) fails is given by group *quasi-actions*, that are a natural coarse generalization of isometric group actions: here (3) is replaced by the weaker condition

$$d(\alpha_g \alpha_h(x), \alpha_{gh}(x)) \le \varepsilon \quad \forall \ g, h \in G, x \in X,$$

where $\varepsilon > 0$ is a constant and d is a distance on X (see e.g. [22]).

In §3.2 we will discuss a general setting where one can pass from an action to a tolerance relation.

3. FROM RELATIONS TO ALGEBRAS

In this section we mimic the construction of the convolution algebra of an étale groupoid associated to an equivalence relation and explain how to pass from a tolerance relation to a complex *-algebra, possibly non-associative. We will then study two special cases: tolerance relations on finite sets, and coming from suitable actions of magmas on sets.

Non-associative algebras naturally arise in the framework of cochain quantization, which consists in constructing a new product on a module algebra A over an Hopf algebra Husing an invertible element $F \in H \otimes H$. See e.g. [1] for a description of octonions as a cochain quantization of the group algebra of the group $(\mathbb{Z}/2\mathbb{Z})^3$. A similar construction was used in [9] to describe the modules of "sections of line bundles" on the noncommutative torus as U(1)-spectral subspaces of a suitable non-associative formal deformation of the algebra of smooth functions on a Heisenberg manifold. In deformation quantization, nonassociative algebras arise if one tries to quantize a manifold M equipped with a 2-form ω which is non-degenerate but not closed, or more generally a manifold M equipped with a skew-symmetric bi-derivation $\{, \}: C^{\infty}(M) \times C^{\infty}(M) \to C^{\infty}(M)$ which does not satisfy Jacobi identity; a similar phenomenon is encoutered in string theory in the presence of a B-field background [34].

Let X be a topological space and $R \subset X \times X$ a tolerance relation. We will say that R is *étale* if there exists a topology on R such that the projection $pr_1 : R \ni (x, y) \mapsto x \in X$ onto the first component is a local homeomorphism.¹ We stress that the topology on R is not necessarily the relative topology coming from the product topology on $X \times X$. In fact, it is usually finer [13]. See also Example 2.

Lemma 1. If R is Hausdorff and étale, the space $C_c(R)$ of compactly-supported complexvalued continuous functions on R is a *-algebra with product and involution given by:

$$(f \star g)(x, z) := \sum_{y \in X: (x, y), (y, z) \in R} f(x, y)g(y, z)$$
(5)

$$f^*(x,z) := \overline{f(z,x)} \tag{6}$$

for all $(x, z) \in R$.

¹If R is an equivalence relation, this simply means that the groupoid of the equivalence relation is étale.

Proof. We wish to show that only finitely many elements in the sum (5) are non-zero, so that the product is well-defined. We can rewrite the right hand side of (5) as a sum of $f(\xi)g(\eta)$ over all $\xi, \eta \in R$ such that $\operatorname{pr}_1(\xi) = x$, $\operatorname{pr}_2(\eta) = z$ and $\operatorname{pr}_2(\xi) = \operatorname{pr}_1(\eta)$. It is then enough to show that the set

$$(\mathrm{pr}_1)^{-1}(x) \cap \mathrm{supp}(f) \tag{7}$$

is finite. But local homeomorphisms have discrete fibers. Thus (7) is the intersection of the closed discrete subset $(\text{pr}_1)^{-1}(x)$ of R and the closed compact subset supp(f) of R (since R is Hausdorff). Thus, (7) is a discrete compact space in the topology induced from R, hence finite.

A neutral element for the product is given by the characteristic function of the diagonal $\Delta := \{(x, x) : x \in X\}$, which is continuous iff Δ is clopen in R. However, since the diagonal map $X \to R$ is in general not continuous, even if X is compact such a characteristic function may be not compactly supported (see Example 2).

Example 2. Let X := [-1,1] and R be the equivalence relation generated by $x \sim -x$. Thus, R is a cross. With the topology induced by $X \times X$, the projection onto the first axis is not a local homeomorphism (the cross is not locally Euclidean). On the other hand we can write $R = (R \setminus \{O\}) \sqcup \{O\}$, put on $R \setminus \{O\}$ and $\{O\}$ the subspace topology from \mathbb{R}^2 , and on R the disjoint union topology. With such a topology, R is étale.

Observe that the diagonal map $X \to R$, $x \mapsto (x, x)$, is not continuous, since $\{O\}$ is closed in R but $\{0\}$ is not closed in X.

The diagonal Δ is clopen but not compact in R, since $\Delta \setminus \{O\}$ is closed in Δ and noncompact. Hence the characteristic function of the interval is continuous but not compactly supported, and the convolution algebra is not unital (even if X is compact).

Example 3. Let X be a discrete space and R any discrete tolerance relation on X. For $(i, j) \in R$, let E_{ij} be the function on R that is 1 at the site (i, j) and zero everywhere else. These functions form a basis of $C_c(R)$ (compactly means finitely supported in the present case). In such a basis, the product is explicitly given by:

$$E_{ij} \star E_{kl} = \begin{cases} \delta_{jk} E_{il} & if (i,l) \in R\\ 0 & otherwise \end{cases}$$
(8)

The algebra is unital if and only if X is a finite set (i.e. the diagonal Δ is compact), with unit given by

$$1 = \sum_{i \in X} E_{ii} \; .$$

When R is an equivalence relation, it is well known that the convolution product is associative, and one can complete the convolution algebra using an injective bounded *-representation. However, if (A, \star) is not associative, there is no injective homomorphism $(A, \star) \to \mathcal{B}(H)$ into bounded operators on a Hilbert space H. There is no good notion of "representation" that works for arbitrary non-associative algebras.

Observe that the definition of convolution product makes sense if we replace complexvalued functions by functions with values in any field \mathbb{K} . However, for the sake of simplicity, we will only study the case $\mathbb{K} = \mathbb{C}$.

3.1. Finite-dimensional case. It is useful to have a pictorial representation of tolerance relations. Observe that $R \subset X \times X$ is a tolerance relation if and only if $\Gamma := (X, R \setminus \Delta)$ is

an undirected graph with no loops. Of course, from any undirected graph with no loops we get a tolerance relation by adding a loop to each vertex. Thus for example the graph:

$$\begin{array}{c}
2 \\
1 \\
3
\end{array}$$
(9)

represents the relation on $X = \{1, 2, 3\}$ with $1 \sim 2$ and $2 \sim 3$, but $1 \not\sim 3$ (where, as costumary, we write $i \sim j$ to mean that $(i, j) \in R$).

A tolerance relation is transitive if, whenever there is an edge between x and y and between y and z, there is also an edge between x and z. One immediately realizes by induction that:

Remark 4. Let X be a finite set. Then, a tolerance relation R on X is an equivalence relation if and only if each connected component of its graph is a complete graph.

In the rest of this section, we will assume that $X := \{1, \ldots, n\}$ is finite and that the topology on both X and R is the discrete one. The convolution product is then given by (8) in the basis of delta functions. We will denote by A(R) the algebra $C_c(R)$ with product (8), and call it the *tolerance algebra* associated to R.

Observe that, if (i, j) and (k, l) are edges in different connected components of the graph of R, the product of the corresponding basis elements of A(R) is zero. The algebra A(R) is then a direct sum of algebras associated to connected components of the graph of R. If R is an equivalence relation with connected graph, which means $R = X \times X$ by Remark 4, if we identify E_{ij} with the $n \times n$ matrix with 1 in position (i, j) and zero everywhere else, we see that (8) is just the usual matrix product, the involution (6) is the usual Hermitian conjugation, and A(R) is *-isomorphic to $M_n(\mathbb{C})$. In general, if R is an equivalence relation, A(R) is isomorphic to a direct sum of matrix algebras (as it is well-known).

Example 5. Let us denote by \mathfrak{A}_3 the tolerance algebra of the relation (9). We think of its elements as matrices $a = (a_{ij}) \in M_3(\mathbb{C})$ with $a_{13} = a_{31} = 0$, and the product of two such elements is the matrix product composed with the natural projection $M_3(\mathbb{C}) \to \mathfrak{A}_3$ (which kills the (1,3) and (3,1) matrix elements).

The algebra \mathfrak{A}_3 is not power associative (which in particular means that it is neither commutative nor associative). Indeed, if

$$a := \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix},$$

then

$$(a \star a) \star a = \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 2 \\ 0 & 1 & 0 \end{bmatrix} \neq a \star (a \star a) = \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & 1 \\ 0 & 2 & 0 \end{bmatrix}.$$

Lemma 6. Let R be a tolerance relation on $X = \{1, ..., n\}$. Then, R is an equivalence relation if and only if A(R) has no subalgebra isomorphic to \mathfrak{A}_3 .

Proof. If R is an equivalence relation, then A(R) cannot have a subalgebra isomorphic to \mathfrak{A}_3 , since the former algebra is associative and the latter is not. If R is not an equivalence relation, then there exists three distinct elements $x, y, z \in X$ such that $x \sim y$ and $y \sim z$ but $x \not\sim z$; the vector subspace of A(R) spanned by the elements $E_{xx}, E_{yy}, E_{zz}, E_{xy}, E_{yz}, E_{yx}, E_{zy}$ is a subalgebra isomorphic to \mathfrak{A}_3 , with isomophism given by (with an abuse of notations): $x \to 1, y \to 2, z \to 3$.

Proposition 7. Let R be a tolerance relation on $\{1, ..., n\}$ and A(R) the tolerance algebra of R. The following properties are equivalent:

- (i) A(R) is associative;
- (ii) A(R) is power associative;
- (iii) R is an equivalence relation.

Proof. If R is not an equivalence relation, then A(R) contains a subalgebra isomorphic to \mathfrak{A}_3 (Lemma 6). Since \mathfrak{A}_3 is not power associative, A(R) cannot be power associative. Thus (ii) \Rightarrow (iii). The implications (iii) \Rightarrow (i) \Rightarrow (ii) are obvious.

It follows from the previous proposition that an algebra that is power associative but not associative cannot be obtained from the above construction. Thus, for example, algebras obtained from \mathbb{R} by a iterated Cayley–Dickson construction with dimension ≥ 8 (octonion, sedenions, etc.) are not tolerance algebras. The same is true for split octionions. Division algebras (such as quaternions) also cannot be obtained from the above construction: since $E_{ii} \star E_{jj} = 0$ for all $i, j \in X$ with $i \neq j$, the algebra A(R) is a division algebra if and only if X is a singleton, i.e. $A(R) = \mathbb{C}$.

3.2. Magmas acting on sets. Let G be a set with a binary operation $*: G \times G \to G$ (the "multiplication"), a unary operation $(-)^{-1}: G \to G$ (the "right inversion"), and an element $1 \in G$ (the "right inverse"), such that:

$$(g * h) * h^{-1} = g * 1 = g \qquad \forall g, h \in G.$$
 (10)

We call G a magma with a right inverse and unit.

Example 8. Consider the set $\mathbb{R} \setminus \{0\}$ with operation x * y := x/y and with standard inversion and unit. Observe that $(x * y) * y^{-1} = xy^{-1}y = x$ even if $y * y^{-1}$ is not 1.

A right action of G on a set X is a map $X \times G \to X$, $(x, g) \mapsto x \triangleleft g$, such that:

$$(x \triangleleft g) \triangleleft g^{-1} = x \triangleleft 1 = x \qquad \forall \ g \in G, x \in X.$$

$$(11)$$

Given such an action, the image R(G, X) of the canonical map:

$$X \times G \to X \times X, \qquad (x,g) \mapsto (x, x \triangleleft g), \tag{12}$$

is a tolerance relation. Indeed, $(x, x) = (x, x \triangleleft 1) \in R(G, X)$ for all $x \in X$, and if $(x, y) = (x, x \triangleleft g) \in R(G, X)$, then $(y, x) = (y, y \triangleleft g^{-1}) \in R(G, X)$ as well.

The action \triangleleft is called *free* if (12) is injective, *transitive* if (12) is surjective.

Remark 9. The action of G on itself by right multiplication is free if and only if G is left-cancellative, i.e. for all $x, g, h \in G$:

$$x * g = x * h \implies g = h.$$

If $g^{-1} * (g * h) = h$ for all $g, h \in G$, then the action is both free and transitive. In Example 8, the right action of G on itself is both free and transitive.

Example 10. If G is a Moufang loop [23], the inverse properties of Moufang loops read: $g^{-1} * (g * h) = h = (h * g) * g^{-1}$ for all $g, h \in G$. They guarantee that G satisfies (10) and that the right action of G on itself is both free and transitive.

If G and X are topological space and G acts freely on X, the bijection $X \times G \to R(G, X)$ given by (12) defines a topology on R that we will call *standard* (and in general is not the subspace topology from $X \times X$). The projection $X \times G \to X$ onto the first component is a local homeomorphism if and only if G is discrete.² Thus:

Remark 11. Let G and X be topological spaces with G acting freely on X. The tolerance relation R(G, X) is étale if and only if G is discrete.

A celebrated example where G is a group and its action is a group action (in the standard sense) is the following.

Example 12. Let $G := \mathbb{Z}$ with action on $X := \mathbb{S}^1 := \mathbb{R}/\mathbb{Z}$ by translations of a multiple of a fixed $\theta \in \mathbb{R} \setminus \mathbb{Q}$. The associated relation R(G, X) is then given by $x \sim y \iff x - y \in \theta \mathbb{Z}$. Since θ is irrational, the action is free and R(G, X) with the topology of $\mathbb{S}^1 \times \mathbb{Z}$ is étale. Let us identify R with $\mathbb{S}^1 \times \mathbb{Z}$. Every $f \in C_c(R)$ is a finite sum

$$f = \sum_{j} f_{j} V^{j} \tag{13}$$

where V^j is the delta function $V^j(k) = \delta_{j,k}$ and $f_j \in C(\mathbb{S}^1)$. The convolution product (5) of two elements fV^j and gV^k is

$$fV^{j} \star gV^{k}(x,n) = f(x)g(x+j\theta)\delta_{j+k,n} .$$
⁽¹⁴⁾

In particular $V^j \star V^k = V^{j+k}$, thus justifying the notations.

Let
$$U \in C(\mathbb{S}^1)$$
 be the function $U(x) = e^{2\pi i x}$. One easily checks that

$$V \star U = e^{2\pi \mathrm{i}\theta} U \star V.$$

We recognize a dense *-subalgebra of the C*-algebra of the noncommutative torus [26].

One can construct many examples where G is not a group. If $X \simeq S^7$ is the set of unit octonions (hence a Moufang loop) and $G \subset X$ a discrete subset containing 1 and closed under inversion, then right multiplication of X by G defines a free action. In particular, G can be a sub-loop, such as the loop of integer vectors in $S^7 \subset \mathbb{R}^8$, or the set of basis vectors and their additive inverses, which are both Moufang loops of finite order, so that the associated tolerance relation is étale.

In general, if K is any group with a free right action \triangleleft on a set X and $G \subset K$ is a subset containing 1 and closed under inversion, then the restriction of \triangleleft to G is a free action in our more general sense, and the associated relation may be non-trivial (not transitive) even if the multiplication in K is associative. For example, if $K = S_3$ is the group of permutations of $X = \{1, 2, 3\}$ and $G \subset S_3$ is the subset containing the identity and the transpositions $1 \leftrightarrow 2$ and $2 \leftrightarrow 3$, then the associated relation is the one with graph (9).

3.3. **Truncations.** It is a common attitude in Noncommutative Geometry to regard the Dirac operator of a spectral triple (A, H, D) as a kind of inverse of the line element of Riemannian geometry:

$$D \sim ds^{-1}$$
.

The precise meaning of this statement is explained for example in [5]. Given a spectral triple, for example the one canonically associated to a Riemannian spin manifold M, one can imagine to implement a cut-off on large energies (small distances) by using the spectral projection $P_{\Lambda} := \chi_{[-\Lambda,\Lambda]}(D)$, with $\Lambda > 0$. One can then replace the original spectral triple

²Observe that no continuity assumption on the operations of G is needed. When G is discrete, they will automatically be continuous but the action on X might still not be continuous.

by the truncated one $(P_{\Lambda}AP_{\Lambda}, P_{\Lambda}H, P_{\Lambda}DP_{\Lambda})$ and think of it as some kind of "coarse graining" of the manifold we started from. If D has compact resolvent, e.g. if the spectral triple is unital, the truncations will be finite-dimensional. In the example of a compact Riemannian spin manifold M, one obtains matrix geometries "approximating" M (see, e.g., [7, 10]).

The issue of convergence of matrix geometries in the case of coadjoint orbits was studied in several seminal papers by Rieffel, see e.g. the most recent one [29] and references therein. The metric convergence of truncations in the $\Lambda \to \infty$ limit were recently studied in [36]: a sufficient condition, there, for convergence is the existence of a " C^1 -approximate order isomorphism", see [36, Def. 2]. In the case of a compact Riemannian spin manifold M, when $A = C^{\infty}(M)$ and D is the Dirac operator of the spin structure acting on the Hilbert space H of L^2 -spinors, it was shown in [15] that one can choose a suitable set of "localized" states on each $P_{\Lambda}AP_{\Lambda}$ forming a sequence of metric spaces (equipped with the distance induced by the truncated Dirac operator) that converge to M in the Gromov-Hausdorff sense. A numerical study through computer simulations is in [14]. The question of convergence of the full state space was solved in [36] in the case of the circle, and is still open for a general (compact) Riemannian spin manifold.

A spectral projection as above defines a linear map $A \to P_{\Lambda}AP_{\Lambda}$ that allows one to put a non-associative product on its image, and is a special case of the following natural construction.

Let B be a C*-algebra and $T : B \to B$ a linear map (our "cut-off" operator). On A := Im(T) we can define a product \star by

$$a \star b := T(ab) \; \forall \; a, b \in A,\tag{15}$$

where the one on the right hand side is the product of B, and get a (possibly non associative) algebra (A, \star) . If $T(a^*) = T(a)^*$ for all $a \in A$, then (A, \star) is a *-algebra. If B is commutative, then (A, \star) is commutative (and then power associative). We will call (A, \star) a "truncation" of B.

Let us observe that for a truncation as above one has two notions of positivity: we could call an element $a \in A$ positive if there exists a finite set of elements $b_1, \ldots, b_k \in A$ such that $a = \sum_{i=1}^k b_i \star b_i^*$, or we could call it positive if it is a positive element of the C*-algebra B. As shown in the next examples, these two notions do not always coincide. If $T: B \to B$ is a positive map, then $b \star b^* = T(bb^*)$ is positive in B for all $b \in A$ and the former notion of positivity implies the latter.

A characteristic property of a truncation map defined by a "sandwich" with a spectral projection — our motivating example — is that it is *idempotent*. That is:

$$T(T(b)) = T(b) \qquad \forall \ b \in B.$$
(16)

From property (16) it follows that, in such a case, $B = A \oplus \ker T$.

Theorem II.6.10.11 of [3] provides a sufficient condition for (15) to be associative: if $T: B \to B$ is idempotent and a completely positive³ contraction, then \star is associative and makes A into a C*-algebra with its involution and with the norm inherited from B (even if it is not, in general, a C*-subalgebra of B).

Example 13. Let B be a C*-algebra and $P \in B$ a projection. Then, the map $T : B \to B$ defined by T(b) := PbP is a completely positive idempotent contraction. Observe, however,

³Here completely positive means that the map id $\otimes T : M_k(\mathbb{C}) \otimes B \to M_k(\mathbb{C}) \otimes B$ is positive for all $k \geq 1$.

that in such a case associativity of (15) can be proved by a direct computation. Indeed, for every $a_1 = T(b_1)$ and $a_2 = T(b_2)$:

$$a_1 \star a_2 = P(Pb_1P)(Pb_2P)P = (Pb_1P)(Pb_2P) = a_1a_2.$$

Thus, \star is the restriction of the product of B to the C*-subalgebra A := Im(T). (See also Example 3.3 of [10].)

Example 13 is a special case of the following notion. Let A be a C*-subalgebra of a C*-algebra B. A linear map $T: B \to B$ with Im(T) = A is called a *conditional expectation* from B to A if one of the two equivalent conditions is satisfied:

- (i) T is idempotent with norm 1;
- (ii) T is positive, idempotent and an A-bimodule map.

Both (i) and (ii) imply that T is a completely positive contraction (see [3, §II.6.10]). However, the setting is less general than that of [3, Theorem II.6.10.11] since here we assume that A is a C*-subalgebra of B, while the above mentioned theorem guarantees that \star is associative, but it is not necessarily the restriction to A of the product in B.

In the next examples, the map T fails to be either positive or idempotent.

Example 14. Let $n \ge 2$ be fixed, $X := \mathbb{S}^1$, let u be the unitary generator of B := C(X), given by $u(t) := e^{it}$, and define:

$$T(f) := \sum_{k=-n+1}^{n-1} u^k \widehat{f}(k) = \frac{1}{2\pi} \sum_{k=-n+1}^{n-1} u^k \int_0^{2\pi} u^{-k}(t) f(t) dt.$$
(17)

Observe that, in this example, the product (15) is power associative but not associative: $(u^* \star u^{n-1}) \star u = u^{n-1}$ while $u^* \star (u^{n-1} \star u) = 0$. Moreover, in spite of power associativity, in general $f^{\star k} \neq T(f^k)$. For $f := u^{-n+1} + u^{n-1} \in A$, for example, one has

$$T(f^3) = 3f \neq f^{\star 3} = 2f$$
 .

The map (17) satisfies (16), but not positivity. Take for example n = 2 and $f = u + u^* + 1$. Then $T(f^2) = 3 + 2(u + u^*)$, i.e. $T(f^2)(t) = 3 + 4\cos t$, proving that the function $T(f^2)$ is not everywhere greater or equal than zero.

Since the Fouries series of a continuous function on \mathbb{S}^1 may be divergent, a smart idea is to replace the partial sum of the Fourier series (17) by the corresponding *n*-th Cesàro sum. As shown in the next example, we loose the property (16) but gain positivity.

Example 15. Let $n \ge 2$ be fixed, $X := \mathbb{S}^1$, let u be the unitary generator of B := C(X), given by $u(t) := e^{it}$, and define:

$$T(f) := \frac{1}{n} \sum_{j=1}^{n} \sum_{k=-j+1}^{j-1} u^k \widehat{f}(k).$$
(18)

From

$$T(u^k) = \begin{cases} \frac{n-|k|}{n}u^k & \text{if } |k| < n\\ 0 & \text{if } |k| \ge n \end{cases}$$

one sees that the maps in (17) and (18) have the same range, and that the latter is not idempotent. From the well-known formula in terms of Fejér kernel:

$$T(f)(t) = \frac{1}{2\pi n} \int_{\mathbb{S}^1} f(t-\tau) \frac{\sin^2(n\tau/2)}{\sin^2(\tau/2)} d\tau$$

it is evident that if f is non-negative, then T(f) is a non-negative function as well.

The product defined by the map (18) is non-associative, as one can check, and in general $f^{\star k} \neq T(f^k)$. For example if $f := u^{-n+1} + u^{n-1}$, then $T(f^3) = \frac{3}{n}f \neq f^{\star 3} = \frac{2}{n}f$.

An interesting although simple observation is that every finite-dimensional tolerance algebra is a truncation of a matrix algebra.

Example 16. Let R be a tolerance relation on $X = \{1, ..., n\}$ and $T : M_n(\mathbb{C}) \to M_n(\mathbb{C})$ be the map:

$$T(b) = \sum_{(i,j)\in R} E_{ii}bE_{jj} \qquad \forall \ b \in B.$$
⁽¹⁹⁾

Then, the product (15) coincides with the convolution product (8) and $(A, \star) = A(R)$ is the tolerance algebra of §3.1.

Observe that such a T is idempotent but, in general, not positive. Let us start with n = 3 and R the relation (9). If:

$$a := \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \ then \ a \star a^* = T(aa^*) = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

has determinant -1, hence $T(aa^*)$ is not a positive semidefinite matrix and $T: M_3(\mathbb{C}) \to M_3(\mathbb{C})$ is not a positive map.

For a general tolerance relation R, arguing as in the proof of Lemma 6, one can show that T is positive if and only if R is an equivalence relation.

The vector space underlying the algebra A(R) is called in [18] the quantum relation on $M_n(\mathbb{C})$ associated to the relation R. See Sect. 1 and in particular Proposition 1.4 of [18].

4. State spaces

4.1. **Operator systems.** Let H be a complex Hilbert space. A concrete unital operator system is a complex vector subspace $A \subset \mathcal{B}(H)$ containing the identity operator 1 of $\mathcal{B}(H)$ and the adjoint of all its elements: $a^* \in A$ for all $a \in A$ (we will only consider unital operator systems). This is very close to the notion of concrete order unit space, which is a real vector subspace of the set of selfadjoint operators on H containing the identity. We refer to [7] for a recollection of properties of operator systems and their use in Noncommutative Geometry, and to [28] for order unit spaces and their use in the theory of compact quantum metric spaces.

If P_n is the spectral projection of the Dirac operator D of \mathbb{S}^1 on eigenspaces with eigenvalue $1 \leq \lambda \leq n$, the truncation $C(\mathbb{S}^1)^{(n)} := P_n C(\mathbb{S}^1) P_n$ is the operator system of $n \times n$ Toeplitz matrices $(n \times n \text{ matrices that are constant on descending diagonals})$ studied in [7, 36]. A dual operator system $C(\mathbb{S}^1)_{(n)}$ is given by the truncation of $C(\mathbb{S}^1)$ in Examples 14-15 (see [7] for the exact meaning of operator system duality, and [12] for the proof that in the example above there is a unital complete order isomorphism of operator systems).

The complex vector space underlying the algebra A(R) in §3.1 is obviously a finitedimensional operator system. Note, however, that not every finite-dimensional operator system is of this form. The vector space A(R) always contains all diagonal matrices, so for example, for $n \ge 2$, the operator system $C(\mathbb{S}^1)^{(n)}$ does not come from a tolerance relation.

If A is an operator system, there is a notion of *state* very much as for C*-algebras, as a positive bounded functional $\varphi : A \to \mathbb{C}$ satisfying $\varphi(1) = 1$. The set of all states, denoted by $\mathcal{S}(A)$, is a convex subset of the Banach dual of A, compact in the weak* topology; its extremal points are called *pure states*. The set of all pure states of A is denoted by $\mathcal{P}(A)$.

Even if we will not deal with convergence problems here, it is worth mentioning the result in [36] that the state spaces of both operator systems $C(\mathbb{S}^1)^{(n)}$ and $C(\mathbb{S}^1)_{(n)}$, with the obvious metric induced by the Dirac operator of \mathbb{S}^1 , converge to $\mathcal{S}(C(\mathbb{S}^1))$ in the Gromov-Hausdorff distance for $n \to \infty$. Surprisingly, it was shown in [17] that the pure state spaces $\mathcal{P}(C(\mathbb{S}^1)^{(n)})$, with metric induced by the Dirac operator of \mathbb{S}^1 , converge to $\mathcal{S}(C(\mathbb{S}^1))$ and not to $\mathcal{P}(C(\mathbb{S}^1)) \simeq \mathbb{S}^1$ as one would have guessed.

Suppose, now, that B is a C*-algebra and $A \subset B$ an operator system. By Hahn-Banach theorem every state of A can be extended to a state of B, and we get a surjective map

$$F: \mathcal{S}(B) \twoheadrightarrow \mathcal{S}(A) \tag{20}$$

given by the restriction of states of B to A. The first two points in the next lemma, that we prove just for the sake of completeness, appeared already in [7, Fact 2.9].

Lemma 17. Let F be the map (20) and $\varphi \in \mathcal{S}(A)$. Then:

- (i) The fiber $F^{-1}(\varphi)$ is a convex subset of $\mathcal{S}(B)$.
- (ii) If φ is pure, then extremal points of $F^{-1}(\varphi)$ are pure states of B.
- (iii) If $F^{-1}(\varphi) = \{\psi\}$ is a singleton and $\psi \in \mathcal{P}(B)$, then $\varphi \in \mathcal{P}(A)$.

Proof. (i) Let $f, g \in F^{-1}(\varphi)$. Then for any $\lambda \in [0, 1]$ the combination

$$\psi := \lambda f + (1 - \lambda)g \tag{21}$$

is still a state of B and $F(\psi) = \lambda \varphi + (1 - \lambda)\varphi = \varphi$, so that $\psi \in F^{-1}(\varphi)$.

(ii) Let now φ be a pure state and ψ an extremal point of $F^{-1}(\varphi)$. Suppose, by contradiction, that there exists $\lambda \in [0, 1[$ and $f, g \in \mathcal{S}(B)$ such that ψ can be decomposed like in (21). Then

$$\varphi = F(\psi) = \lambda F(f) + (1 - \lambda)F(g).$$

But φ is pure, so it must be $F(f) = F(g) = \varphi$, that means $f, g \in F^{-1}(\varphi)$, contradicting the hypothesis that ψ is an extremal point of $F^{-1}(\varphi)$.

(iii) Let $\varphi = \lambda \varphi_1 + (1 - \lambda) \varphi_2$ with $\lambda \in [0, 1]$ and $\varphi_1, \varphi_2, \in \mathcal{S}(A)$. Choose any $\psi_1 \in F^{-1}(\varphi_1)$ and $\psi_2 \in F^{-1}(\varphi_2)$ (recall that F is surjective). Observe that

$$\lambda\psi_1 + (1-\lambda)\psi_2 \in F^{-1}(\varphi),$$

which implies $\psi = \lambda \psi_1 + (1 - \lambda)\psi_2$. But ψ is pure, hence $\lambda = 0$ or $\lambda = 1$. This implies that φ is pure as well.

A natural question concerns injectivity of (20), or at least injectivity of its restriction to pure states. We will see a natural example of a convex-linear map whose restriction to pure states is injective in §5, when discussing informationally complete positive operator valued measures (IC POVM).

In general, injectivity of a convex-linear map on extremal points does not guarantee injectivity on the full domain, as shown next.

Example 18. Let Δ_3 be the standard simplex

$$\Delta_3 := \left\{ p = (p_1, \dots, p_4) \in \mathbb{R}^4 : p_i \ge 0 \ \forall \ i \ and \ \sum p_i = 1 \right\}$$

Let $F: \Delta_3 \to [0,1]^2$ be the map

$$F(p) := (p_1 + p_3, p_2 + p_3)$$
.

It gives a bijection between the vertices of the tetrahedron and the vertices of the square. Since every convex set is the closed convex hull of its extremal points, $F(\Delta_3) \supset [0,1]^2$. The opposite inclusion is obvious, thus $F(\Delta_3) = [0, 1]^2$. The map is clearly not a bijection, since for example $F(\frac{1}{2}, \frac{1}{2}, 0, 0) = F(0, 0, \frac{1}{2}, \frac{1}{2})$.

For the operator system of a tolerance relation on $\{1, \ldots, n\}$ it is easy to verify that the map (20) is not injective, not even on pure states, unless R is an equivalence relation. (This could be derived from Corollary 3.9 in [8] where L is the adjacency matrix of the graph of R, but we prefer to give an independent proof.)

Proposition 19. Let $B := M_n(\mathbb{C})$, R a tolerance relation on $\{1, \ldots, n\}$ and A = A(R) its tolerance algebra. The following are equivalent:

- (i) the map (20) is injective;
- (ii) the restriction of (20) to pure states is injective;
- (iii) $A = B = M_n(\mathbb{C}).$

Proof. (iii) \Rightarrow (i) \Rightarrow (ii) is obvious. We now prove that the negation of (iii) implies the negation of (ii). Suppose that R is not an equivalence relation, which means that there exists $x, y, z \in X$ such that $x \sim y, y \sim z$ and $x \not\sim z$. We can assume x = 1, y = 2, z = 3 (relabeling the elements of X is equivalent to reordering the rows and columns of a matrix). For $u \in \mathbb{C}$ with $|u| \leq 1$, consider the state with density matrix:

$$\rho_u = \frac{1}{2} \begin{bmatrix} 1 & 0 & u & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 \\ u^* & 0 & 1 & 0 & \cdots & 0 \\ \hline 0 & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 \end{bmatrix}$$

Since ρ_u is a rank 1 projection, the associated state φ_u , given by $\varphi_u(a) = \text{Tr}(\rho_u a)$ for all $a \in B$, is pure. Now we observe that, if $a \in A$:

$$\varphi_u(a) = \operatorname{Tr}(\rho_u T(a)),$$

where T is the truncation map in Example 16 and we used the fact that T is the identity on A, since $T \circ T = T$. By cyclicity of the trace and symmetry of R:

$$\varphi_u(a) = \sum_{(i,j)\in R} \operatorname{Tr}(\rho_u E_{ii} a E_{jj}) = \sum_{(i,j)\in R} \operatorname{Tr}(E_{ii} \rho_u E_{jj} a) = \operatorname{Tr}(T(\rho_u) a) .$$
(22)

But $T(\rho_u)$ is independent of u, since the truncation map kills the matrix elements in position (1,3) and (3,1). Thus, all states φ_u have the same image under the map (20).

4.2. **Positivity in a tolerance algebra.** As in the previous section, let R be a tolerance relation on the set $\{1, \ldots, n\}$, $A = A(R) \subset M_n(\mathbb{C})$ the vector space in Example 16, T the map (19), \star the product (8). There are two natural partial orders on A(R), which we will denote by \geq and \succeq , defined as follows. For $a \in A(R)$ we will write $a \geq 0$ if a is a positive semidefinite matrix, and $a \succeq 0$ if there exists $b \in M_n(\mathbb{C})$ such that $b \geq 0$ and a = T(b).

Observe that, since T is idempotent, this in particular means that, for all $a \in A$, $a \ge 0$ implies $a \succeq 0$. On the other hand, for every element $a \in A$ of the form

$$a = \sum_{i=1}^{k} b_i \star b_i^*, \tag{23}$$

with $b_1, \ldots, b_k \in A(R)$, one has $a = T(\sum_{i=1}^k b_i b_i^*) \succeq 0$ even if in general (23) may be not a positive semidefinite matrix (see Example 16).

Elements (23) are the natural candidates for positive elements in a *-algebra. The relation with the partial order \succeq is illustrated in the next proposition.

Recall that in an undirected graph $\Gamma = (V, E)$, a *dominant vertex* is a vertex that is adjacent to all other vertices of the graph.

Proposition 20. Assume that in each connected component of the graph of R there is a dominant vertex. Then, for all $a \in A(R)$, one has $a \succeq 0$ if and only if a is of the form (23) for some $b_1, \ldots, b_k \in A(R)$.

Proof. We have to prove the implication \Rightarrow , the other being always true. Clearly every $a \succeq 0$ is of the form $T(\sum_{i=1}^{k} b_i b_i^*)$ for some $b_1, \ldots, b_k \in M_n(\mathbb{C})$ (in fact, even with k = 1). The non-trivial part is to show that one can choose b_1, \ldots, b_k belonging to A(R).

Up to a permutation of rows and columns, which preserves both the partial order \succeq and the decompositions of the form (23), we can think of $A(R) \subset M_n(\mathbb{C})$ as subset of block diagonal matrices where each block corresponds to a connected component of the graph of R. It is then enough to prove the proposition for each block, i.e. under the assumption that the graph of R is connected.

By hypothesis, there exists j_0 such that $(i, j_0) \in R$ for all $i \in \{1, \ldots, n\}$. Let a = T(c) with $c \in M_n(\mathbb{C})$ positive semidefinite. Since T is linear and every positive semidefinite matrix is a linear combination with positive coefficients of rank 1 projections, it is enough to prove the statement when c is itself a rank 1 projection. Thus, $c = \sum_{i,j=1}^{n} v_i v_j^* E_{ij}$ for some unit vector $v = (v_1, \ldots, v_n) \in \mathbb{C}^n$. Let $b := \sum_{i=1}^{n} v_i E_{ij_0}$ and observe that $b \in A(R)$. Since $bb^* = c$, this concludes the proof.

We will see that the assumption in Prop. 20 is necessary (cf. Prop. 31).

As recalled in the previous section, we use the partial order \geq to define $\mathcal{S}(A)$. Explicitly, we define a partial order \geq on the dual vector space A^* by setting $\varphi \geq 0$ if and only if $\varphi(a) \geq 0$ for all $a \geq 0$. A state $\varphi \in \mathcal{S}(A)$ will then be a positive element of A^* with normalization $\varphi(1) = 1$. With such a definition, the restriction of a state of $M_n(\mathbb{C})$ to Ais a state, and (20) is well defined.

Even if we use \geq to define states, when trying to describe states in terms of matrices the other partial order \succeq pops up, as explained below. This stems from the next crucial observation.

Proposition 21. For $x \in M_n(\mathbb{C})$, let $\varphi_x : M_n(\mathbb{C}) \to \mathbb{C}$ be the linear map defined by

$$\varphi_x(a) := \operatorname{Tr}(x^*a) \qquad \forall \ a \in M_n(\mathbb{C}).$$

Then, the map $(A, \succeq) \to (A^*, \geq)$ sending $x \in A$ to $\varphi_x|_A$ is an isomorphism of ordered vector spaces.

Recall that an isomorphism of ordered complex vector spaces is an isomorphism of the underlying real vector spaces that is compatible with the partial orders (this remark is necessary since $x \mapsto \varphi_x$ is an antilinear map), cf. [30]. Notice that the isomorphism above transforms the partial order \succeq , defined using the non-associative product, into the partial order \geq , which only depends on the underlying operator system.

Proof. For $a, b \in M_n(\mathbb{C})$ let $\langle a, b \rangle := \operatorname{Tr}(a^*b)$ be the Hilbert-Schmidt inner product. Its restriction to A is still an inner product, and from its non-degeneracy it follows that the map $A \to A^*$, $x \mapsto \varphi_x = \langle x, . \rangle$, is injective. Since the vector spaces A and A^* have the

same dimension, such a map is surjective as well, hence an isomorphism of real vector spaces. It remains to show that, for all $x \in A$, one has

$$x \succeq 0 \iff \varphi_x \ge 0. \tag{24}$$

By Hahn-Banach theorem, every positive element in A^* is the restriction of a positive linear functional $M_n(\mathbb{C}) \to \mathbb{C}$. Thus $\varphi_x \ge 0$ if and only if there exists $b \in M_n(\mathbb{C})$ such that $b \ge 0$ and

$$\varphi_x(a) = \varphi_b(a) = \operatorname{Tr}(ba) \quad \forall a \in A$$

Repeating the proof of (22) one finds that

$$\varphi_b(a) = \operatorname{Tr}(T(b)a) \quad \forall a \in A,$$

thus $\langle x - T(b), a \rangle = 0$ for all $a \in A$, which implies x = T(b). Since $b \ge 0$, it follows that $x \succeq 0$. This proves " \Leftarrow ".

Conversely, if $x \succeq 0$, then x = T(b) for some positive $b \in M_n(\mathbb{C})$ and $\varphi_x = \varphi_{T(b)}|_A = \varphi_b|_A$ is the restriction to A of the positive linear functional $\varphi_b : M_n(\mathbb{C}) \to \mathbb{C}$, hence it is positive itself.

Definition 22. Let us denote by $\mathcal{D}(A)$ the set of all $\rho \in A$ such that $\rho \succeq 0$ and $\operatorname{Tr}(\rho) = 1$.

If $A = M_n(\mathbb{C})$ (thus, R is the trivial relation $i \sim j$ for all $i, j = \{1, \ldots, n\}$), the one above is the standard definition of *density matrix*.

In general, since the relation \geq is contained in the relation \succeq , every positive semidefinite matrix with trace 1 (every density matrix) belongs to $\mathcal{D}(A)$, but there may be matrices in $\mathcal{D}(A)$ that are not positive semidefinite. We may call an element $\rho \in A$ with trace 1 a *strict* density matrix if $\rho \geq 0$, and a *weak* density matrix if $\rho \succeq 0.^4$

Example 23. Let R be the relation (9) and

$$\rho = \frac{1}{3} \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \frac{1}{3} T \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

This is a weak density matrix for A(R), but not a strict one since $det(\rho) < 0$.

An immediate consequence of Prop. 21 is that states on A in the sense of operator systems are in bijection with weak density matrices in the sense of (non-associative) tolerance algebras. The same conclusion could be reached using Prop. 3.10 of [8].

Corollary 24. The map $\mathcal{D}(A) \to \mathcal{S}(A)$, $\rho \mapsto \varphi_{\rho}$, is a bijection (in fact, it is a homeomorphism if we put the norm topology on $\mathcal{D}(A)$ and the weak-* topology on $\mathcal{S}(A)$).

4.3. **Pure states of a tolerance algebra.** Let R be a tolerance relation on the set $\{1, \ldots, n\}$, A(R) and T as in the previous section. Here we want to give an explicit description of pure states of A(R).

Given a non-zero vector $v = (v_1, \ldots, v_n) \in \mathbb{C}^n$, then

$$R_v := \{ (i,j) \in R : v_i v_j \neq 0 \}$$
(25)

is a tolerance relation on the (non-empty) subset of $i \in \{1, ..., n\}$ such that $v_i \neq 0$. In terms of graphs, R_v is the subgraph of R obtained by removing all vertices i such that $v_i = 0$, and all edges that are incident on such vertices.

⁴Observe that, in the physics language, a strict density matrix is a density matrix, while a weak density matrix is not.




FIGURE 2. Some tolerance relations discussed in the text.

Definition 25. A non-zero vector $v \in \mathbb{C}^n$ is called R-tolerant if the graph of R_v is connected. A pure state of $M_n(\mathbb{C})$ is called R-tolerant if its density matrix is a projection in the direction of an R-tolerant vector.

Notice that in the previous definition we do not assume that the graph of R is connected.

Pictorially, some examples are illustrated in Figure 2, where vertices are labelled by the components of v. In the first example (on the left), since v_3 is adjacent to every other vertex, if $v_3 \neq 0$ the graph of R_v is connected; if $v_3 = 0$ and $v_4 \neq 0$, R_v is disconnected; if $v_3 = v_4 = 0$, then R_v is connected. Similarly in the third example (on the right), R_v is disconnected if and only if $v_2 = 0$ and both v_1 and v_3 are non-zero. In the second example (in the middle), we see the opposite phenomenon: if both v_1 and v_2 are non-zero, $R_v = R$ is disconnected; if $v_1 = 0$ or $v_2 = 0$, R_v is connected.

Inspired by these "experiments", let us record some more general examples.

Example 26. If $A(R) = M_n(\mathbb{C})$, then every non-zero vector $v \in \mathbb{C}^n$ is R-tolerant (R_v is connected for all $v \neq 0$, since a complete graph cannot be disconnected by deleting vertices).

If $A(R) = \mathbb{C}^n$, then R-tolerant vectors are those with only one non-zero component (corresponding to diagonal projection matrices).

If R is the relation (9), then $v \in \mathbb{C}^3$ is R-tolerant except when $v_2 = 0$ and $v_1, v_3 \neq 0$.

Under the isomorphism between a vector subspace of $M_n(\mathbb{C})$ and its dual, given by the Hilbert-Schmidt inner product, the map $F : \mathcal{S}(M_n(\mathbb{C})) \to \mathcal{S}(A(R))$ in (20) becomes the restriction to density matrices of the truncation map $T : M_n(\mathbb{C}) \to A(R)$.

If $v \in \mathbb{C}^n$ is a unit vector, let us denote by $P_v := (v_i v_j^*) \in M_n(\mathbb{C})$ the corresponding projection. Because of Lemma 17(ii), every pure state of A(R) is the restriction of a pure state of $M_n(\mathbb{C})$, hence it has weak density matrix $T(P_v)$ for some unit vector $v \in \mathbb{C}^n$. The remaining part of this section is devoted to the proof that:

Proposition 27. $T(P_v)$ is the weak density matrix of a pure state of A(R) if and only if $v \in \mathbb{C}^n$ is an R-tolerant unit vector. More precisely, F gives a bijection between the subset of pure states of $M_n(\mathbb{C})$ that are R-tolerant and pure states of A(R).

Before we prove the proposition, let us note that the definition of R-tolerant vector is very close to the one of state with ϵ -connected support in [8, Sect. 4.2]. However, the latter concerns proximity relations on a metric space, while our more general Definition 25 is for arbitrary tolerance relations on a finite set. There is, of course, a non-empty intersection between the two cases. On this intersection, Prop. 27 is analogous to Theorem 4.18 [8].

We need a few preliminary lemmas.

Lemma 28. Let $k \ge 3$, let $w \in \mathbb{C}^k$ be a vector with $w_i \ne 0$ for all $i \in \{1, \ldots, k\}$, and let $a = (a_{ij}) \in M_k(\mathbb{C})$ be a positive semidefinite matrix such that $a_{ij} = w_i w_j^*$ for all $|i-j| \le 1$. Then $a_{ij} = w_i w_j^*$ for all $i, j \in \{1, \ldots, k\}$.

Proof. We are claiming that a is uniquely determined by its main diagonal and the two adjacent descending diagonals. We will prove the statement by induction on k. Let then $a \in M_k(\mathbb{C})$ be a matrix with elements $a_{ij} = w_i w_j^*$ if $|i - j| \leq 1$ and call $\lambda_{ij} := a_{ij}/w_i w_j^*$ if |i - j| > 1. If k = 3, one has

$$a = \begin{pmatrix} |w_1|^2 & w_1w_2^* & \lambda_{13}w_1w_3^* \\ w_2w_1^* & |w_2|^2 & w_2w_3^* \\ \lambda_{13}^*w_3w_1^* & w_3w_2^* & |w_3|^2 \end{pmatrix}.$$

Since

$$\det(a) = (\lambda_{13} + \lambda_{13}^* - |\lambda_{13}|^2 - 1)|w_1w_2w_3|^2$$

and $w_1w_2w_3 \neq 0$, the condition $\det(a) \geq 0$ implies $\lambda_{13} + \lambda_{13}^* - |\lambda_{13}|^2 - 1 \geq 0$. The unique solution of such an inequality is $\lambda_{13} = 1$, which is what we wished to prove.

Now let $k \ge 4$ be arbitrary and assume by inductive hypothesis that the lemma is true for matrices of size k - 1. If $a = (a_{ij}) \in M_k(\mathbb{C})$ is positive semidefinite and $a_{ij} = w_i w_j^*$ for all $|i - j| \le 1$, the matrix $a' = (a_{ij}) \in M_{k-1}(\mathbb{C})$ obtained from a by removing the last row and column is positive semidefinite and satisfies $a'_{ij} = w_i w_j^*$ for all $|i - j| \le 1$. By inductive hypothesis $a'_{ij} = w_i w_j^*$ for all $i, j \in \{1, \ldots, k-1\}$. Thus, the principal minor of a given by elements in the intersection of the rows and columns $\{1, k - 1, k\}$ is:

$$\det \begin{pmatrix} |w_1|^2 & w_1 w_{k-1}^* & \lambda_{1k} w_1 w_k^* \\ w_{k-1} w_1^* & |w_{k-1}|^2 & w_{k-1} w_k^* \\ \lambda_{1k}^* w_k w_1^* & w_k w_{k-1}^* & |w_k|^2 \end{pmatrix} = (\lambda_{1k} + \lambda_{1k}^* - |\lambda_{1k}|^2 - 1)|w_1 w_{k-1} w_k|^2.$$

The determinant above is ≥ 0 if and only if $\lambda_{1k} = 1$, which completes the inductive step.

Lemma 29. If $\rho \in M_n(\mathbb{C})$ is a density matrix, $v \in \mathbb{C}^n$ an *R*-tolerant unit vector and $T(\rho) = T(P_v)$, then $\rho = P_v$.

Proof. Let $v \in \mathbb{C}^n$ be an *R*-tolerant unit vector and $\rho = (\rho_{ij}) \in M_n(\mathbb{C})$. The condition $T(\rho) = T(P_v)$ gives $\rho_{ij} = v_i v_j^*$ for all $(i, j) \in R$. Using the condition $\rho \ge 0$ we now prove that $\rho_{ij} = v_i v_j^*$ for all $(i, j) \notin R$ as well. Since we are working with Hermitian matrices, it is enough to consider i < j. Thus, in the rest of the proof $(i, j) \notin R$ and i < j, and we assume that $\rho \ge 0$.

If $v_i = 0$, since the submatrix:

$$\begin{pmatrix} \rho_{ii} & \rho_{ij} \\ \rho_{ji} & \rho_{jj} \end{pmatrix} = \begin{pmatrix} 0 & \rho_{ij} \\ \rho_{ij}^* & |v_j|^2 \end{pmatrix}$$

must have non-negative determinant, we deduce that $\rho_{ij} = 0$. But $v_i v_j^* = 0$ as well, so $\rho_{ij} = v_i v_j^*$. The case $v_j = 0$ is analogous.

If both v_i and v_j are non-zero, since v is R-tolerant, there exists a path in R_v from v_i to v_j . That is, there exists non-zero components v_{l_1}, \ldots, v_{l_k} such that $l_1 = i$, $l_k = j$ and $(l_r, l_{r+1}) \in R$ for all $1 \leq r < k$. Since $(i, j) \notin R$, it must be $k \geq 3$.

For $r, s \in \{1, \ldots, k\}$, call $a_{rs} := \rho_{l_r l_s}$ and $w_r := v_{l_r}$. Observe that $a = (a_{rs}) \in M_k(\mathbb{C})$ and $w = (w_r) \in \mathbb{C}^k$ satisfy the hypotheses of Lemma 28. We conclude that $a_{rs} = w_r w_s^*$ for all $r, s \in \{1, \ldots, k\}$. In particular, for r = 1 and s = k we get $\rho_{ij} = a_{1k} = w_1 w_k^* = v_i v_j^*$, that is exactly what we wished to prove. **Lemma 30.** Let $w = (w_1, \ldots, w_k) \in \mathbb{C}^k$ be a unit vector with $w_i \neq 0$ for all $i \in \{1, \ldots, k\}$, R' a tolerance relation on $\{1, \ldots, k\}$ and $T : M_k(\mathbb{C}) \to A(R')$ the corresponding truncation map. If $T'(P_w)$ is the weak density matrix of a pure state of A(R'), then the graph of R' must be connected.

Proof. By contradiction, assume that the graph of R' is not connected. After renaming the vertices and reordering rows and columns we can assume that, for some $m \in \{1, \ldots, k-1\}$, the first m vertices are disconnected from the last k - m, that is: if $i \leq m < j$ then $(i, j) \notin R'$. This means that $T(P_w)$ is a block diagonal matrix of the form:

$$T(P_w) = \left(\begin{array}{c|c} Q' \\ \hline \\ \hline \\ Q'' \end{array}\right)$$

where $Q' \in M_m(\mathbb{C}), Q'' \in M_{k-m}(\mathbb{C})$, and off-diagonal blocks are zero. Let

$$t := \sum_{i=1}^m |w_i|^2$$

and observe that, since w has unit norm, $1-t = \sum_{i=m+1}^{k} |w_i|^2$. Since w has all components different from zero, both t and 1-t are non-zero, that means 0 < t < 1. Consider the unit vectors:

$$w' := t^{-1/2}(w_1, \dots, w_m, 0, \dots, 0)$$
 and $w'' := (1-t)^{-1/2}(0, \dots, 0, w_{m+1}, \dots, w_k).$

Clearly

$$\left(\begin{array}{c|c} Q' \\ \hline \\ \hline \\ \hline \\ \end{array}\right) = t T(P_{w'}) \quad \text{and} \quad \left(\begin{array}{c|c} 0 \\ \hline \\ \hline \\ \hline \\ \end{array}\right) = (1-t) T(P_{w''}).$$

Thus $T(P_w) = t T(P_{w'}) + (1 - t) T(P_{w''})$, and the corresponding state of A(R') is not pure.

We are now ready to prove Proposition 27.

Proof of Prop. 27. Let $v \in \mathbb{C}^n$ be a unit vector and denote by φ the state of A(R) with weak density matrix $T(P_v)$.

If v is R-tolerant, it follows from Lemma 29 that $F^{-1}(\varphi) = \{\psi\}$ is a singleton, with ψ the state with density matrix P_v . Thus F is injective on R-tolerant pure states of $M_n(\mathbb{C})$. From Lemma 17(iii), it also follows that φ is a pure state of A(R). It remains to prove that if v is not R-tolerant, then φ is not a pure state of A(R).

Without loss of generality, we can assume that the non-zero components of v are the first k ($k \ge 1$). We can identify $A(R_v)$ (risp. $M_k(\mathbb{C})$) with the vector subspace of A(R) (risp. $M_n(\mathbb{C})$) of matrices with the last n - k rows and columns filled with zeros. We now apply Lemma 30 to $R' = R_v$ and $w = (v_1, \ldots, v_k)$. Since R_v is not connected, $T'(P_w)$ is not the weak density matrix of a pure state of A(R'). That means there are two distinct density matrices $\rho_1, \rho_2 \in M_k(\mathbb{C})$ and a $t \in [0, 1[$ such that $T'(P_w) = tT'(\rho_1) + (1-t)T'(\rho_2)$. But T(a) = T'(a) for all $a \in M_k(\mathbb{C})$, hence $T(P_v) = T'(P_w) = tT(\rho_1) + (1-t)T(\rho_2)$ is a non-trivial convex combination of two weak density matrices. We conclude that the state φ is not pure.

As a corollary of the above results, we now show that the assumption in Prop. 20 is necessary.

Proposition 31. Let R be a tolerance relation on $X := \{1, ..., n\}$ and assume that each $a \succeq 0$ in A(R) is of the form (23), for some $b_1, ..., b_k \in A(R)$. Then the graph of R has a dominant vertex in each connected component.

Proof. Similarly to the proof of Prop. 20, it is enough to prove the statement when the graph of R is connected. Since R is connected, w := (1, 1, ..., 1) is an R-tolerant vector. Called $P := n^{-1}(w_i w_j^*)$, we now prove that a := T(P) can be written in the form (23) only if there the graph of R has a dominant vertex (at least one).

Assume that $a = T(\sum_{i=1}^{k} b_i b_i^*)$ for some $b_1, \ldots, b_k \in A(R)$. From Lemma 29 it follows

$$\sum_{i=1}^{k} b_i b_i^* = P.$$
 (26)

From this we deduce that

$$||b_i^*v||^2 = \langle v, b_i b_i^*v \rangle \le \langle v, Pv \rangle = 0$$

— i.e. $b_i v = 0$ — for all i = 1, ..., k and for every v in the kernel of P. Because of (26), not all matrices $b_i b_i^*$'s can be zero: for some $i_0 \in \{1, ..., k\}$, one has $b_{i_0} b_{i_0}^* \neq 0$. Since $b_{i_0} b_{i_0}^*$ vanishes on the kernel of the rank 1-projection P, it must be $b_{i_0} b_{i_0}^* = \lambda P$ for some $\lambda > 0$. Called $b := \lambda^{-1/2} b_{i_0}$, we proved that

$$P = bb^*$$

for some $b \in A(R)$. Since ker $P = \text{ker } b^*$, the matrix b^* must be of rank 1, hence $b^* = n^{-1/2}(u_i w_i^*)$ for some unit vector $u \in \mathbb{C}^n$, and with w the vector defined above. That is:

$$\sqrt{n} b^* = \begin{pmatrix} u_1 & u_1 & \dots & u_1 \\ u_2 & u_2 & \dots & u_2 \\ \vdots & \vdots & & \vdots \\ u_n & u_n & \dots & u_n \end{pmatrix}.$$

At least one row of b^* , say j_0 , must be non-zero. Since $b^* \in A(R)$ and in the row j_0 all its elements are different from zero, this means that $(i, j_0) \in R$ for all i = 1, ..., n. Thus, j_0 is a dominant vertex in the graph of R.

5. Positive operator valued measures: a curious example

Given an observable, represented by a Hermitian matrix P, and a physical system in a state described by a density matrix ρ of the same size of P, a measurement of such observable returns an eigenvector of P. Order the eigenvalues and label them with an integer, say from 1 to k. The outcome of a measurement is then an integer $i \in \{1, \ldots, k\}$, telling us which eigenvalue comes out of our measurement apparatus. Performing the experiment many times gives the frequency of the outcome i, that is an estimate of the probability $\rho^{P}(i)$ that the state collapses in an eigenstate associated to the eigenvalue i. Such a probability is expressed by the formula:

$$\rho^P(i) := \operatorname{Tr}(\rho P_i),\tag{27}$$

where P_i is the spectral projection of the *i*-th eigenvalue of P. The spectral projections are an example of a positive operator valued measure (POVM), which in the simplest case are a collection of k positive semidefinite $n \times n$ matrices P_i , with $i = 1, \ldots, k \leq n$, such that $\sum_{i=1}^{k} P_i = 1$. The complex linear span of the P_i 's is a concrete operator system $A \subset M_n(\mathbb{C})$. From now on, we will denote by P a collection of positive semidefinite $n \times n$ matrices P_1, \ldots, P_k . POVMs arise in the study of physical systems of which we have partial knowledge ("open" quantum systems), and an excellent textbook reference is [24].

Given a POVM, $P := (P_i)_{i=1}^k$, the formula (27) defines a convex-linear map

$$\rho \mapsto \rho^P \tag{28}$$

from density matrices to probability distributions on k points, and a natural question is whether this map is injective: i.e. the knowledge of the probabilities $\rho^P(i)$ allows one to reconstruct the state ρ . If the map (28) is injective, the POVM P is called "informationally complete" (IC). Since, in such a case, the map gives a smooth embedding of a $n^2 - 1$ smooth manifold (the interior of the set of $n \times n$ density matrices) into the interior of a k - 1 simplex, this forces $k \ge n^2$.

Let us consider on $M_n(\mathbb{C})$ the inner product $\langle a, b \rangle := \operatorname{Tr}(a^*b)$, and extend (28) to a linear map $L: M_n(\mathbb{C}) \to \mathbb{R}^k$ in the obvious way:

$$L(a) := (\langle P_1, a \rangle, \dots, \langle P_k, a \rangle).$$
⁽²⁹⁾

Lemma 32. Let $P = (P_i)_{i=1}^k$ be a POVM. Then, the convex-linear map (28) is injective if and only if the linear map (29) is.

Proof. The implication " \Leftarrow " is clear. We now prove " \Rightarrow ". Every $a \in \ker L$ has $\operatorname{Tr}(a) = \sum_{i=1}^{k} \langle a, P_i \rangle = 0$. Assume that $\ker L \neq \{0\}$. Then we can find a non-zero Hermitian matrix a in the kernel, that can be written as $a = b_1 - b_2$ with b_1 and b_2 two distinct positive semidefinite matrices with $\operatorname{Tr}(b_1) = \operatorname{Tr}(b_2) =: \lambda$. The trace of a positive semidefinite matrix is zero if and only if the matrix is zero, but b_1 and b_2 cannot be both zero, thus $\lambda \neq 0$. Hence $\rho_1 := \lambda^{-1}b_1$ and $\rho_2 := \lambda^{-1}b_2$ are two distinct density matrices such that $\rho_1^P = \rho_2^P$, and (28) is not injective.

If A is the operator system associated to our POVM, since ker $L = A^{\perp}$, an immediate corollary of the previous lemma is the well-known fact that the POVM is IC if and only if $A = M_n(\mathbb{C})$.

We now change setting and pass to diagonal matrices, that is we consider a finite *classical* system. Notice that, in fact, Lemma 32 holds for any C*-subalgebra of $M_n(\mathbb{C})$ (direct sum of matrix algebra), and in particular for diagonal matrices, i.e. finite classical systems.

Let us consider the following simple classical example. Fix two integers $1 \le k \le n$. Suppose we have *n* points on a circle and a detector with *n* LEDs at those points (Figure 3). Only groups of *k* consecutive LEDs can turn on simultaneously, and when a group is on we know that a particle passed throught one of those *k* points, but not which one. The detector cannot distinguish between *k* consecutive points, meaning that we have a tolerance relation:




FIGURE 3

$$i \sim j \iff |i - j| < k$$

This is clearly not transitive if $k \notin \{1, n\}$.

The POVM describing this detector is $P = (\frac{1}{k}Q_i)_{i=1}^n$, with $Q_i \in M_n(\mathbb{C})$ the diagonal matrix with 1 in positions $i, i+1, \ldots, i+k-1 \pmod{n}$ and zero everywhere else. For example, for k = 3 and n = 5 one has:

$$Q_{1} = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 0 \end{bmatrix} \quad Q_{2} = \begin{bmatrix} 0 & & \\ & 1 & \\ & & 1 \end{bmatrix} \quad Q_{3} = \begin{bmatrix} 0 & & \\ & 0 & \\ & & 1 \end{bmatrix} \quad Q_{4} = \begin{bmatrix} 1 & & & \\ & 0 & & \\ & & 1 \end{bmatrix} \quad Q_{5} = \begin{bmatrix} 1 & & & \\ & 0 & & \\ & & 0 & & 1 \end{bmatrix},$$

where the zeros outside the main diagonal are omitted. If

$$\rho = \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$$

is a density matrix (a probability distribution on n points), then

$$\rho^P(i) = \frac{1}{k} \sum_{j=i}^{i+k-1} \lambda_j$$

where the index in the sum is defined mod n. Since the case k = 1 is trivial, from now on let us assume that $k \ge 2$.

Proposition 33. The POVM $P = (\frac{1}{k}Q_i)_{i=1}^n$ is IC if and only if k and n are coprime.

Proof. It follows from Lemma 32 that (28) is injective if and only if the map $L : \mathbb{C}^n \to \mathbb{C}^n$, sending $(\lambda_1, \ldots, \lambda_n)$ to the tuple with elements $\frac{1}{k} \sum_{j=i}^{i+k-1} \lambda_j$, $1 \le i \le n$, is injective. The representative matrix of L is the $n \times n$ matrix:

where here and in the following we denote by C the *clock* and by S the *shift*, given by

$C := \begin{bmatrix} q & 0 & 0 & \dots & 0 \\ 0 & q^2 & 0 & \dots & 0 \\ 0 & 0 & q^3 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & q^n \end{bmatrix}$	S :=	0 0	0 1 : :		$\begin{array}{c} 0 \\ \vdots \\ 1 \end{array}$,
--	------	-----	------------	--	---	---

and q is a primitive n-th root of 1. The shift can be transformed into the clock by a unitary transformation, so that M is conjugated to the matrix $C^0 + C^1 + \ldots + C^{k-1}$. Such a matrix is diagonal, with element in position (i, i) given by

$$\sum_{j=0}^{k-1} q^{ij} \tag{30}$$

for all i = 1, ..., n. The coefficient matrix is invertible if and only if (30) is different from zero for all i = 1, ..., n.

Observe that, if i = n, then $q^i = 1$ and (30) equals k. If i < n, since q is a primitive n-th root of 1 we have $q^i \neq 1$ and

$$\sum_{j=0}^{k-1} q^{ij} = \frac{1-q^{ik}}{1-q^i}$$

Thus, (30) is zero if and only if q^i is a k-th root of unity, that is ik is a multiple of n. Hence, L is injective if and only if there is no integer ik < nk that is a multiple of n, i.e. there is no integer less than nk that is a common multiple of n and k. But this exactly one of the characterizations of coprime numbers.

When k and n are not coprime, even if the POVM is not IC, the map (28) is injective on pure states. In fact, we can strengthen the claim: **Proposition 34.** If k < n, fibers of the map (28) which contain pure states are singletons.

Proof. Observe that if $\rho = E_{kk}$, then

$$\rho^P = \frac{1}{k} \underbrace{(\overbrace{1,\ldots,1}^{k \text{ times}}, \overbrace{0,\ldots,0}^{n-k \text{ times}})}_{(31)}.$$

The image of every other pure state is obtained from the formula above with a cyclic permutation, and they are all different if $n-k \ge 1$, i.e. if at least one component is zero (if the last 1 in ρ^P is in position *i*, then necessarily $\rho = E_{ii}$). Thus, the map (28) is injective on pure states.

Now we wish to prove that if ρ^P is a cyclic permutation of (31), then the state ρ is necessarily pure. It is enough to show that, if ρ is not pure, then $k\rho^P$ has (at least) one entry that is neither 0 nor 1. If ρ is not pure, it means that it has (at least) one entry $\lambda_{i_0} = \lambda$ which is neither 0 nor 1. Thus

$$k\rho^{P}(i_{0}+1) = \sum_{j=i_{0}+1}^{i_{0}+k} \lambda_{j} \le \sum_{j \ne i_{0}} \lambda_{j} = 1 - \lambda < 1$$

(since k < n, λ_{i_0} does not appear in the former sum). If $\rho^P(i_0 + 1) \neq 0$ the proof is concluded. If $\rho^P(i_0 + 1) = 0$, then $\lambda_j = 0$ for all $i_0 < j \le i_0 + k$. Therefore $k\rho^P(i_0) = \lambda_{i_0} = \lambda$ is neither 0 nor 1.

Acknowledgements

We thank Alain Connes and Walter van Suijlekom for useful discussions. FL and FDA acknowledge support form the INFN Iniziativa Specifica GeoSymQFT. FL acknowledges financial support from the State Agency for Research of the Spanish Ministry of Science and Innovation through the "Unit of Excellence Maria de Maeztu 2020–2023" award to the Institute of Cosmos Sciences (CEX2019-000918-M) and from PID2019-105614GB-C21 and 2017-SGR-929 grants. GL acknowledges partial support from INFN, Iniziativa Specifica GAST, from INdAM-GNSAGA and from the INDAM-CNRS IRL-LYSM.

References

- [1] H. Albuquerque and S. Majid, Quasialgebra Structure of the Octonions, J. Algebra 220 (1999) 188-224.
- [2] A.P. Balachandran, G. Bimonte, E. Ercolessi, G. Landi, F. Lizzi, G. Sparano and P. Teotonio-Sobrinho, Noncommutative lattices as finite approximations, J. Geom. Phys. 18 (1996) 163–194.
- [3] B. Blackadar, Operator algebras: Theory of C*-Algebras and von Neumann Algebras, Encyclopaedia of Mathematical Sciences vol. 122, Springer, 2006.
- [4] P. Bouwknegt, K. Hannabuss and V. Mathai, Nonassociative Tori and Applications to T-duality, Commun. Math. Phys. 264 (2006), 41–69.
- [5] A. Connes, Noncommutative Geometry, Academic Press, 1994.
- [6] A. Connes and M. Marcolli, Noncommutative Geometry, Quantum Fields and Motives, Colloquium Publications, vol. 55, AMS, 2008.
- [7] A. Connes and W.D. van Suijlekom, Spectral Truncations in Noncommutative Geometry and Operator Systems, Commun. Math. Phys. 383 (2020), 2021–2067.
- [8] A. Connes and W.D. van Suijlekom, *Tolerance relations and operator systems*, arXiv:2111.02903 [math.OA].
- [9] F. D'Andrea and D. Franco, Non-Associative Geometry of Quantum Tori, SIGMA 12 (2016) 015.
- [10] F. D'Andrea, F. Lizzi and P. Martinetti, Spectral geometry with a cut-off: topological and metric aspects, J. Geom. Phys. 82 (2014) 18–45.
- [11] A. Devastato, M. Kurkov and F. Lizzi, Spectral Noncommutative Geometry Standard Model and all that, Int. J. Mod. Phys. A34 (2019) 1930010.
- [12] D. Farenick, The operator system of Toeplitz matrices, Trans. AMS Series B, 8 (2021), 999–1023.

- [13] T. Giordano, I. Putnam and C. Skau, Affable equivalence relations and orbit structure of Cantor dynamical systems, Ergod. Th. & Dynam. Sys. 24 (2004) 441–475.
- [14] L. Glaser and A.B. Stern, Understanding truncated non-commutative geometries through computer simulations, J. Math. Phys. 61 (2020) 033507.
- [15] L. Glaser and A.B. Stern, Reconstructing manifolds from truncations of spectral triples, J. Geom. Phys. 159 (2021) 103921.
- [16] J.M. Gracia-Bondía, J.C. Varilly and H. Figueroa, *Elements of Noncommutative Geometry*, Birkhäuser, 2001.
- [17] E.-J. Hekkelman, Truncated Geometry on the Circle, arXiv:2111.13865 [math.QA].
- [18] G. Kuperberg and N. Weaver, A von Neumann algebra approach to quantum metrics/quantum relations, AMS, 2012.
- [19] G. Landi, An Introduction to Noncommutative Spaces and their Geometry, Springer-Verlag, 1997.
- [20] G. Landi and F. Lizzi, Projective Systems of Noncommutative Lattices as a Pregeometric Substratum, in "Quantum Groups, Noncommutative Geometry and Fundamental Physical Interactions", D. Kastler and M. Rosso Eds., Nova Science Publisher, Commack NY, 1999, pp 85–113, arXiv:math-ph/9810011.
- [21] M. McClure (https://math.stackexchange.com/users/21361/mark mcclure). Are there real-life relations which are symmetric and reflexive but not transitive? https://math.stackexchange.com/q/ 1566311 (version: 2015-12-08).
- [22] L. Mosher, M. Sageev and K. Whyte, Quasi-actions on trees I. Bounded valence, Ann. of Math. 158 (2003), 115–164.
- [23] R. Moufang, Zur Struktur von Alternativkörpern, Math. Ann. 110 (1935), 416–430.
- [24] A. Peres, Quantum Theory: Concepts and Methods, Springer, 2002.
- [25] J.H. Poincaré, La Science et l'Hypothèse, Ernest Flammarion, 1902.
- [26] M.A. Rieffel, C*-Algebras Associated with Irrational Rotations, Pacific J. Math. 93 (1981) 415-429.
- [27] M.A. Rieffel, Applications of strong Morita equivalence to transformation group C*-algebras, Proc. Symp. Pure Math. 38 (1982), 299–310.
- [28] M.A. Rieffel, Gromov-Hausdorff Distance for Quantum Metric Spaces, Mem. Amer. Math. Soc. 168 (2004) no. 796, 1–65.
- [29] M.A. Rieffel, Dirac operators for matrix algebras converging to coadjoint orbits, arXiv:2108.01136 [math.OA].
- [30] H.H. Schaefer and M.P. Wolff, *Topological Vector Spaces*, 2nd ed., Springer-Verlag 1999.
- [31] A. Sims, G. Szabó and D. Williams, Operator Algebras and Dynamics: Groupoids, Crossed Products, and Rokhlin Dimension, Birkhäuser, 2020.
- [32] R.D. Sorkin, Finitary substitute for continuous topology, Int. J.. Theo. Phys. 30 (1991) 923–947.
- [33] A.B. Sossinsky, Tolerance space theory and some applications, Acta Appl. Math. 5 (1986) 137–167.
- [34] R. Szabo, Higher Quantum Geometry and Non-Geometric String Theory, PoS(CORFU2017)151.
- [35] W.D. van Suijlekom, Noncommutative Geometry and Particle Physics, Springer, 2015.
- [36] W.D. van Suijlekom, Gromov-Hausdorff convergence of state spaces for spectral truncations, J. Geom. Phys. 162 (2021) 104075.
- [37] E.C. Zeeman, The topology of the brain and visual perception, in "Topology of 3-manifolds and related topics", pp. 240–256, Prentice-Hall, 1962.
 - (F. D'Andrea) UNIVERSITÀ DI NAPOLI FEDERICO II AND I.N.F.N. SEZIONE DI NAPOLI COMPLESSO MSA, VIA CINTIA, 80126 NAPOLI, ITALY

Email address: francesco.dandrea@unina.it

(G. Landi) UNIVERSITÀ DI TRIESTE, VIA A. VALERIO, 12/1, 34127 TRIESTE, ITALY INSTITUTE FOR GEOMETRY AND PHYSICS (IGAP) TRIESTE, ITALY I.N.F.N. SEZIONE DI TRIESTE, ITALY

Email address: landi@units.it

(F. Lizzi) UNIVERSITÀ DI NAPOLI FEDERICO II AND I.N.F.N. SEZIONE DI NAPOLI COMPLESSO MSA, VIA CINTIA, 80126 NAPOLI, ITALY INSTITUT DE CIÉNCIES DEL COSMOS, UNIVERSITAT DE BARCELONA

INSTITUT DE CIENCIES DEL COSMOS, UNIVERSITAT DE DARCELONA

Email address: fedele.lizzi@na.infn.it