



TITLE:

Toric varieties and smooth convex approximation of a polytope

AUTHOR(S):

BATYREV, VICTOR V.

CITATION:

BATYREV, VICTOR V.. Toric varieties and smooth convex approximation of a polytope. 数理解析研究所講究録 1992, 776: 20-20

ISSUE DATE:

1992-03

URL:

<http://hdl.handle.net/2433/82441>

RIGHT:

Toric varieties and smooth convex approximations of a polytope

VICTOR V. BATYREV

Let V be a projective toric variety, \mathcal{L} an ample T -linearized invertible sheaf on V with T -invariant metric q whose curvature form is positive. If s is a global section of \mathcal{L} which nonvanishes on T , then $f(x) = \log \|s(x)\|_q^{-1}$ can be approximated by a piecewise linear function as x tends to some point in $V \setminus T$. This observation gives an explicit formula for some convex approximation of an arbitrary convex polytope in a finite dimensional real space.

Let $P \subset \mathbf{R}^d$ be a convex d -dimensional polytope defined by inequalities

$$\langle p, \gamma_i \rangle \leq a_i, \quad 1 \leq i \leq n,$$

where γ_i are linear functions on \mathbf{R}^d . We assume that the zero $0 \in \mathbf{R}^d$ is in the interior of P , so that all $a_i \neq 0$. After a normalization we get

$$P = \{p \in \mathbf{R}^d \mid \langle p, \alpha_i \rangle \leq 1, \quad 1 \leq i \leq n\},$$

where $\alpha_i = \gamma_i/a_i$. Consider the following two functions on \mathbf{R}^d :

$$F(p) = \frac{1}{2} \log \left(\sum_{1 \leq i \leq n} e^{2\langle p, \alpha_i \rangle} \right),$$

$$L(p) = \max_{1 \leq i \leq n} (\langle p, \alpha_i \rangle).$$

Proposition 1. $F(p)$ satisfies the following conditions

- (i) $F(p)$ is a convex function;
- (ii) $F(p) > L(p)$ for all $p \in \mathbf{R}^d$.

For any positive real number t , define the following convex sets:

$$Q_t = \{p \in \mathbf{R}^d \mid F(tp) \leq t\},$$

$$P_t = \{p \in \mathbf{R}^d \mid L(tp) \leq t\}.$$

Clearly, for all t , one has $P_t = P$. It follows from the proposition 1 that Q_t is a convex body with a smooth boundary, and $Q_t \subset P$ for all t .

Proposition 2. $\lim_{t \rightarrow \infty} Q_t = P$.