

## TORSION AND CRITICAL METRICS ON CONTACT THREE-MANIFOLDS

BY DOMENICO PERRONE<sup>1)</sup>

**1. Introduction.** Let  $M$  be a compact orientable manifold of class  $C^\infty$ . It is well known that a Riemannian metric  $g$  on  $M$  is a critical point of the functional “integral of the scalar curvature  $\int_M r dv$ ” defined on the set of all Riemannian metrics of the same total volume on  $M$ , if and only if  $g$  is an Einstein metric.

Now let  $(M, \omega)$  be a compact contact three-manifold. Then there exists a unique vector field  $X_0$  on  $M$  such that  $\omega(X_0)=1$  and  $d\omega(X_0, \cdot)=0$ . Consider the following functional

$$\mathcal{F}(g)=\int_M r dv \quad g \in \mathcal{M}(\omega)$$

where  $\mathcal{M}(\omega)$  denotes the space of all associated Riemannian metrics to the contact form  $\omega$ . This functional was studied by Blair and Ledger [2] in general dimension. However the three-dimensional case has many special features to merit a separate study. Chern and Hamilton [7] introduced the torsion  $\tau=L_{X_0}g$ , namely the Lie derivative of  $g$  with respect to  $X_0$ , in their study of compact contact three-manifolds, and studied the Dirichlet energy

$$\mathcal{E}_c(g)=\int_M c^2 dv \quad g \in \mathcal{M}(\omega), \quad c^2=\frac{1}{2}|\tau|^2$$

over the set of “ $CR$ -structures” on  $M$  (see also Tanno [15]). Goldberg, the present author and Toth [10] studied the geometry of a compact contact Riemannian three-manifold  $(M, \omega, g)$  with  $g$  critical metric of  $\mathcal{E}_c$ .

The main purpose of this paper is to study compact Riemannian three-manifolds  $(M, \omega, g)$  with  $g$  critical metric of the functional  $\mathcal{F}$ .

In §3, we show that a point  $g$  of  $\mathcal{M}(\omega)$  is a critical point of  $\mathcal{F}$ , if and only if

$$(1.1) \quad \nabla_{X_0}\tau=0.$$

---

*1980 Mathematics Subject Classification.* Primary 53C15; Secondary 53C20.

*Key words and phrases.* Contact three-manifolds, torsion, critical metrics, curvature.

<sup>1)</sup> Supported by funds of the Ministero Pubblica Istruzione.

Received April 22, 1989; revised July 17, 1989

This condition is related to some interesting geometric properties, for example it is equivalent to the condition that the sectional curvatures of all planes, at a given point, perpendicular to  $B=\ker\omega$  are equal to  $(1-c^2/4)$ . The (1.1) was incorrectly obtained in [7] as the condition for the critical metrics of the functional  $\mathcal{E}_c$ . Note that  $X_0$  is a Killing vector field with respect to  $g$ , if and only if  $g$  is a critical point of  $\mathcal{E}_c$  and  $\mathcal{F}$ . Besides if  $g$  is  $\omega$ -Einstein or locally symmetric, then it is a critical point of  $\mathcal{F}$ .

In §4 we show that a metric  $g$  of  $\mathcal{M}(\omega)$  is  $\omega$ -Einstein if, and only if, the following hold: (1)  $g$  is a critical point of  $\mathcal{F}$ , and (2) the  $\phi$ -torsion  $\phi=-\tau\cdot\phi$  is perpendicular to the orbit of  $g$  under the group of diffeomorphisms of  $M$ . Moreover we give some properties of a tensor  $S_1$  which measures the deviation from the  $\omega$ -Einstein structure, for example  $S_1=-\frac{1}{2}\nabla_{X_0}\tau$  if and only if the above condition (2) holds.

In §5, we extend some results of [8] and [10]. Precisely we show that the metric of a compact contact Riemannian three-manifold  $(M, \omega, g, X_0)$  whose characteristic vector field  $X_0$  is of Killing, may be deformed to a contact metric of positive sectional curvature if either the Ricci curvature is greater than  $-2g$  or the  $\phi$ -sectional curvature is greater than  $-3$ . Hence if, in addition,  $M$  is simply connected, then by [11] it is diffeomorphic with the three-sphere.

**2. Contact manifolds.** A  $(2n+1)$ -dimensional manifold  $M$  is said to be a *contact manifold* if it carries a global 1-form  $\omega\neq 0$  with the property that  $\omega\wedge(d\omega)^n\neq 0$  everywhere. It has an underlying almost contact structure  $(X_0, \omega, \phi)$ , where  $\omega(X_0)=1, \phi X_0=0$  and  $\phi^2=-I+\omega\otimes X_0$ . A metric  $g$ , called an *associated metric*, can then be found such that  $\omega=g(X_0, \cdot), d\omega(X, Y)=g(\phi X, Y)$  and hence  $g(\phi X, Y)=-g(X, \phi Y)$ . These metrics are constructed by the polarization of  $d\omega$  evaluated on a local orthonormal basis of an arbitrary metric on the sub-bundle  $B$  of  $TM$  defined by  $\ker\omega$ . We refer to  $(\omega, g)$  or  $(\omega, g, X_0, \phi)$  as a *contact Riemannian structure*. All metrics  $g$  of  $\mathcal{M}(\omega)$ , namely associated to the contact form  $\omega$ , have the same volume element  $(1/2^n n!)\omega\wedge(d\omega)^n$ , and hence we will write  $dv$  instead of  $dv_g$ . Given a contact metric structure  $(\omega, g, X_0, \phi)$ , the torsion  $\tau=L_{X_0}g$  satisfies (cf. [9]):

$$(2.1) \quad \tau(X_0, \cdot)=0, \quad \tau(X, Y)=\tau(Y, X)$$

$$(2.2) \quad \tau(\phi X, Y)=\tau(X, \phi Y), \quad \tau(\phi X, \phi Y)=-\tau(X, Y).$$

Moreover (see for example formula (3.1))  $\tau(X, Y)=2g(\phi X, hY)$  where  $h=\frac{1}{2}L_{X_0}\phi$ .

So  $h$  is a symmetric operator which anticommutes with  $\phi$ . If  $X_0$  is a Killing vector field with respect to  $g$ , the contact metric structure is said to be *K-contact*. It is easy to see that a contact metric structure is *K-contact* if and only if  $\tau=0$  (or equivalently  $h=0$ ). The reader is referred to [3] for details and other properties of contact manifolds. In the sequel we denote by  $R, S, r$  and  $K$ , respectively, the curvature tensor, the Ricci tensor, the scalar curvature

and the sectional curvature of a given contact Riemannian manifold; moreover for tensor fields  $U$  and  $V$  of the same type, we put

$$\langle U, V \rangle = U^i \dots V_{i, \dots} \quad \text{and} \quad |U|^2 = \langle U, U \rangle.$$

**3. Torsion and critical metrics.** Let  $M(\omega, g, X_0, \phi)$  be a  $(2n+1)$ -dimensional contact Riemannian manifold and  $\nabla$  the Riemannian connection with respect to  $g$ . First we give the following.

**PROPOSITION 3.1.** *The tensor field  $\nabla_{X_0}\tau$  satisfies the following properties:*

(i)  $(\nabla_{X_0}\tau)(X, Y) = (\nabla_{X_0}\tau)(Y, X),$

(ii)  $(\nabla_{X_0}\tau)(X_0, \cdot) = 0,$

(iii)  $(\nabla_{X_0}\tau)(\phi X, \phi Y) = -(\nabla_{X_0}\tau)(X, Y),$

(iv) for  $E$  in  $B$ ,  $|E|=1$ , the sectional curvature  $K(X_0, E)$  is given by

$$K(X_0, E) = -\frac{1}{2}(\nabla_{X_0}\tau)(E, E) + 1 - |h(E)|^2,$$

(v)  $\nabla_{X_0}\tau = 0$  if, and only if,  $K(X_0, E') - K(X_0, E) = |h(E)|^2 - |h(E')|^2$  for every  $E, E'$  in  $B$ ,  $|E|=|E'|=1$ ,

(vi) if  $n=1$ , then

$$\begin{aligned} (\nabla_{X_0}\tau)(X, Y) &= S(\phi X, \phi Y) - S(X, Y) + \omega(X)S(X_0, Y) + \omega(Y)S(X_0, X) \\ &\quad - S(X_0, X_0)\omega(X)\omega(Y). \end{aligned}$$

*Proof.* (i)  $\nabla_{X_0}\tau$  is symmetric because  $\tau$  is symmetric.

(ii)  $(\nabla_{X_0}\tau)(X_0, \cdot) = X_0\tau(X_0, \cdot) - \tau(\nabla_{X_0}X_0, \cdot) - \tau(X_0, \nabla_{X_0}\cdot) = 0$  because  $\nabla_{X_0}X_0 = 0$  (cf. [3]) and  $\tau(X_0, \cdot) = 0$ .

(iii) follows from  $\nabla_{X_0}\phi = 0$  (cf. [3]), i.e.  $\nabla_{X_0}\phi(X) = \phi(\nabla_{X_0}X)$ , and (2.2).

(iv) Since  $L_{X_0}d\omega = 0$  (cf. [3]), we have

$$\begin{aligned} 0 &= X_0d\omega(X, Y) - d\omega([X_0, X], Y) - d\omega(X, [X_0, Y]) \\ &= X_0g(X, \phi Y) - g([X_0, X], \phi Y) - g(X, \phi[X_0, Y]) \\ &= \tau(X, \phi Y) + g(X, [X_0, \phi Y] - \phi[X_0, Y]) = \tau(X, \phi Y) + 2g(X, hY) \end{aligned}$$

and hence

$$(3.1) \quad \tau(X, Y) = 2g(\phi X, hY).$$

Since  $\nabla_{X_0}\phi = 0$ , from (3.1) it follows that

$$(3.2) \quad (\nabla_{X_0}\tau)(X, Y) = 2g(\phi X, (\nabla_{X_0}h)Y).$$

Moreover we have the following formula (cf. (3.3) of [4])

$$(3.3) \quad \nabla_{X_0}h = \phi - \phi h^2 - \phi R(\cdot, X_0)X_0.$$

From (3.3) and (3.2), since  $h$  is symmetric and anticommutes with  $\phi$ , we get

$$\begin{aligned} K(X_0, E) &= g(R(E, X_0)X_0, E) = g(-(\nabla_{X_0}h)E + \phi E - \phi h^2 E, \phi E) \\ &= -\frac{1}{2}(\nabla_{X_0}\tau)(E, E) + 1 - g(hE, hE). \end{aligned}$$

(v) If  $\nabla_{X_0}\tau=0$ , from (iv) we have

$$K(X_0, E') - K(X_0, E) = |h(E)|^2 - |h(E')|^2$$

for  $E, E'$  in  $B$ ,  $|E|=|E'|=1$ . Conversely if this formula holds, then (iv) implies  $(\nabla_{X_0}\tau)(E, E) = (\nabla_{X_0}\tau)(E', E')$ . Choosing  $E' = \phi E$ , by (iii), we obtain

$$(\nabla_{X_0}\tau)(E, E) = 0 \quad \text{for } E \text{ in } B, |E|=1.$$

So, by (ii),  $\nabla_{X_0}\tau=0$ .

(vi) For  $E$  in  $B$ ,  $|E|=1$ , since  $h\phi = -\phi h$ , we have  $|hE| = |h\phi E|$  and hence (iv) implies

$$(3.4) \quad K(X_0, E) - K(X_0, \phi E) = -(\nabla_{X_0}\tau)(E, E)$$

(see also Lemma 7.1 of [15]). Since  $\dim M=3$ , from (3.4) it follows that

$$S(\phi E, \phi E) - S(E, E) = (\nabla_{X_0}\tau)(E, E).$$

Consequently for  $E, E'$  in  $B$ ,

$$S(\phi(E+E'), \phi(E+E')) - S(E+E', E+E') = (\nabla_{X_0}\tau)(E+E', E+E')$$

implies

$$(3.5) \quad S(\phi E, \phi E') - S(E, E') = (\nabla_{X_0}\tau)(E, E').$$

Finally for  $X$  and  $Y$  in  $TM$ ,  $\phi X$  and  $\phi Y$  are in  $B$  and  $\phi^2 X = -X + \omega(X)X_0$ ,  $\phi^2 Y = -Y + \omega(Y)X_0$ , therefore by (iii) and (3.5) we get the property (vi).

**THEOREM 3.2.** *Let  $(M, \omega)$  be a compact contact three-manifold. Then a metric  $g$  in  $\mathcal{M}(\omega)$  is a critical point of the functional  $\mathcal{F}$  if and only if*

$$\nabla_{X_0}\tau=0.$$

*Proof.* Let  $g(t)$  be a smooth curve in  $\mathcal{M}(\omega)$  such that  $g(0)=g$ . We calculate  $d\mathcal{F}/dt$  at  $t=0$ , where

$$\mathcal{F}(t) = \mathcal{F}(g(t)) = \int_M r(t) dv.$$

We put

$$g(t) = g + tk + [t^2]$$

where  $[t^2]$  denotes a set of terms of higher order ( $\geq 2$ ) in  $t$ , and  $k$  is a second order symmetric tensor that satisfies (see [6] p. 304)

$$k(X_0, \cdot) = 0 \quad \text{and} \quad k(\phi X, \phi Y) = -k(X, Y).$$

Moreover the scalar curvature  $r(t)$  is given (see [15] § 13) by

$$r(t) = r + t\{\operatorname{div} - \langle k, S \rangle\} + [t^2]$$

where  $\operatorname{div}$  denotes a term which is a divergence. So, by Green's Theorem, we get

$$\frac{d\mathcal{F}}{dt}(0) = \left\{ \int_M \frac{dr(t)}{dt} dv \right\}(0) = - \int_M \langle k, S \rangle dv.$$

Since  $k(X_0, \cdot) = 0$ , we can write  $\langle k, S \rangle = \langle k, T \rangle$  where

$$T = S - S(X_0, \cdot) \otimes \omega - \omega \otimes S(X_0, \cdot) + S(X_0, X_0) \omega \otimes \omega$$

Moreover, since  $k(\phi E, \phi E) = -k(E, E)$  for  $E$  in  $B$ ,  $\langle k, T \rangle = \langle k, V \rangle$  where  $V = \frac{1}{2}T - \frac{1}{2}S(\phi \cdot, \phi \cdot)$ . On the other hand by Proposition 3.1 property (vi),

$$\frac{1}{2} \nabla_{X_0} \tau = \frac{1}{2} \{S(\phi \cdot, \phi \cdot) - T\} = -V.$$

Therefore

$$(3.6) \quad \frac{d\mathcal{F}}{dt}(0) = \frac{1}{2} \int_M \langle k, \nabla_{X_0} \tau \rangle dv.$$

So if  $\nabla_{X_0} \tau = 0$ , then  $g$  is a critical point of  $\mathcal{F}$ . Conversely assume that  $g$  is critical for  $\mathcal{F}$ . We put  $k = \nabla_{X_0} \tau$ , then by Proposition 3.1  $k$  is symmetric,  $k(X_0, \cdot) = 0$ , and  $k(\phi X, Y) = -k(X, \phi Y)$ . Consequently, by [6] p. 304 (see also [15]),  $g(t) = ge^{tk^*}$ ,  $-\varepsilon < t < \varepsilon$ , is a smooth curve in  $\mathcal{M}(\omega)$  such that  $g(0) = g$  where  $k^* = (k^i_j)$  and  $g(t)(X, Y) = g(X, e^{tk^*} Y)$ . Applying (3.6) to this deformation, we get  $\nabla_{X_0} \tau = 0$ .

*Remark 3.1.* (i) Blair and Ledger [2] proved, in general dimension, that a metric  $g$  in  $\mathcal{M}(\omega)$  is a critical point of  $\mathcal{F}$  if and only if the Ricci operator and  $\phi$  when restricted to the contact distribution, commute.

(ii)  $\nabla_{X_0} \tau = 0$  is the condition incorrectly obtained in [7] (cf. Theorem 5.4) for a metric  $\mathcal{E}_c$ -critical. Therefore, by our Theorem 3.2, the main result of [9] holds replacing the assumption  $g$   $\mathcal{E}_c$ -critical by  $g$   $\mathcal{F}$ -critical.

(iii) The condition  $\nabla_{X_0} \tau = 0$ , in general dimension, was studied by Tanno (see [15] § 7) because it is related to some interesting properties. For example he proved that the conditions:  $\nabla_{X_0} \tau = 0$ ,  $\nabla_{X_0} \nabla X_0 = 0$  and  $\nabla_{X_0} T^* = 0$ , are equivalent, where  $T^*$  is the torsion tensor of the generalized Tanaka connection. From Proposition 3.1 we obtain

$$\nabla_{X_0} \tau = 0 \text{ iff } K(X_0, E') - K(X_0, E) = |hE|^2 - |hE'|^2 \text{ for } E, E' \text{ in } B, |E| = |E'| = 1.$$

Hence, when  $\nabla_{X_0} \tau = 0$ , we have at a given point

$K(X_0, E') > K(X_0, E)$  (resp.  $=$ ) iff  $|hE| > |hE'|$  (resp.  $=$ ).

In particular if  $E'$  is a unit vector of the plane generated by  $\{E, \phi E\}$ , we have  $|hE'| = |hE|$ . So, for three-dimensional manifolds, the condition  $\nabla_{X_0}\tau = 0$  is equivalent to the condition that the sectional curvatures of all planes at a given point perpendicular to  $B$  be equal.

*Remark 3.2.* Let  $(M, \omega)$  be a compact contact three-manifold. Chern and Hamilton [7] studied also the following energy

$$\mathcal{E}_w(g) = \int_M W dv \quad g \in \mathcal{M}(\omega)$$

where  $W = \frac{1}{8} \left( r + \frac{c^2}{2} + 2 \right)$  is the Webster scalar curvature (see [7] p. 284). If  $g(t)$  is a smooth curve in  $\mathcal{M}(\omega)$  with  $g(0) = g$ , then  $8\mathcal{E}_w(g(t)) = \mathcal{F}(g(t)) + \frac{1}{2}\mathcal{E}_c(g(t)) + 2\text{vol}(M, g)$  and hence

$$(3.7) \quad 8(d\mathcal{E}_w/dt)(0) = (d\mathcal{F}/dt)(0) + \frac{1}{2}(d\mathcal{E}_c/dt)(0).$$

Tanno proved (see [15] § 5) that

$$(3.8) \quad (d\mathcal{E}_c/dt)(0) = - \int_M \langle k, \nabla_{X_0}\tau - 2\tau \cdot \phi \rangle dv$$

From (3.6), (3.7) and (3.8), we get

$$(3.9) \quad (d\mathcal{E}_w/dt)(0) = \frac{1}{8} \int_M \langle k, \tau \cdot \phi \rangle dv.$$

If  $\tau = 0$ , then  $g$  is a critical point of  $\mathcal{E}_w$ . Conversely assume that  $g$  is a critical point of  $\mathcal{E}_w$ , defining  $k = \tau \cdot \phi$ ,  $g(t) = ge^{tk}$  is a smooth curve in  $\mathcal{M}(\omega)$  (see [6] p. 304 or [15]) with  $g(0) = g$ . Applying (3.9) to this deformation we have  $\tau = 0$ . Therefore we obtain the following.

**THEOREM 3.3 (Chern-Hamilton).** *Let  $(M, \omega)$  be a compact contact three-manifold. Then a metric  $g$  in  $\mathcal{M}(\omega)$  is a critical point of  $\mathcal{E}_w$  if and only if the characteristic vector field  $X_0$  is of Killing with respect to  $g$ .*

This Theorem was obtained in [7] (see Theorem 5.2) where  $\mathcal{E}_w$  was studied as a functional on  $\mathcal{M}(\omega)$  regarded as the set of "CR-structures" on  $M$ .

*Examples of critical metrics.* Let  $(\omega, g)$  be a contact Riemannian structure on a compact three-manifold  $M$ .

(i) Note that:  $M$  is  $K$ -contact if and only if  $g$  is a critical metric for  $\mathcal{F}$  and  $\mathcal{E}_c$ .

(ii) If  $g$  is  $\omega$ -Einstein (in particular if  $g$  is of constant sectional curvature),

then  $g$  is critical for  $\mathcal{F}$  (see Theorem 4.3). If  $g$  is of constant sectional curvature  $K=0$ , then it is not  $K$ -contact and so this metric is critical for  $\mathcal{F}$  but not for  $\mathcal{E}_c$ .

(iii) If  $g$  is locally symmetric, then  $g$  is a critical metric for  $\mathcal{F}$ . In fact, since  $g$  is locally symmetric, by Lemma 1 of [5] we have  $\nabla_{X_0}h=0$ , and so by formula (3.2) we get  $\nabla_{X_0}\tau=0$ .

(iv) The natural contact Riemannian structure of the tangent sphere bundle of a compact Riemannian 2-manifold with constant curvature  $-1$  is critical for  $\mathcal{E}_c$  but not for  $\mathcal{F}$  (combine (i), Theorem of [4] and a result of Tashiro [3] p. 136).

(v) Let  $N$  be a compact orientable surface of constant negative curvature  $-1$ . Let  $(\theta^1, \theta^2)$  be an orthonormal coframe and  $\Omega_2^1$  the connection 1-form. Chern and Hamilton [7] defined on the unit tangent bundle  $T_1N$  a contact Riemannian structure  $(\omega, g')$  by

$$\omega = \frac{1}{2}\Omega_2^1 \quad \text{and} \quad g' = \frac{1}{4}\{\theta^1 \otimes \theta^1 + \theta^2 \otimes \theta^2 + 4\omega \otimes \omega\}.$$

It is not difficult to see that the Ricci curvature in the direction of  $X_0$  and the scalar curvature of  $(T_1N, \omega, g')$  are given by

$$S(X_0, X_0)=2 \quad \text{and} \quad r=\text{const.}=-10.$$

Recall that a contact Riemannian three-manifold is  $K$ -contact iff  $S(X_0, X_0)=2$  (see [3] p. 65). Also recall the result of Tanno [14] that a locally symmetric  $K$ -contact manifold is of constant curvature. So the metric  $g'$  is critical for  $\mathcal{F}$  and  $\mathcal{E}_c$  but is not locally symmetric.

**4. Contact  $\omega$ -Einstein spaces of dimension three.** Let  $M$  be a  $(2n+1)$ -dimensional manifold with contact metric structure  $(\omega, g, X_0, \phi)$ .  $M$  is said to be  $\omega$ -Einstein if the Ricci tensor  $S$  is of the form

$$(4.1) \quad S = ag + b\omega \otimes \omega$$

where  $a$  and  $b$  are functions on  $M$ . It is known that if  $M$  is a  $K$ -contact  $\omega$ -Einstein  $(2n+1)$ -manifold, with  $n>1$ , then the functions  $a$  and  $b$  are constant. Moreover every  $K$ -contact three-manifold is  $\omega$ -Einstein and the Ricci tensor is given by

$$S = \left(\frac{r}{2}-1\right)g + \left(-\frac{r}{2}+3\right)\omega \otimes \omega.$$

However, we know nothing about the geometry of contact  $\omega$ -Einstein three-manifolds. Note that the connected sum of two non-simply connected closed three-manifolds has never  $K$ -contact structure (see [13]), while every compact orientable three-manifold has a contact structure (see [12]). In this section we give a characterization of contact  $\omega$ -Einstein three-manifolds in terms of critical metrics of  $\mathcal{F}$ . Moreover we give some properties of a tensor  $S_1$  which measures

the deviation from the  $\omega$ -Einstein structure.

**PROPOSITION 4.1.** *Let  $M$  be a  $(2n+1)$ -dimensional manifold with contact metric structure  $(\omega, g, X_0, \phi)$ . If  $M$  is  $\omega$ -Einstein, then the Ricci tensor is given by*

$$(4.2) \quad S = \left\{ \frac{r}{2n} - \left( 1 - \frac{c^2}{4n} \right) \right\} g + \left\{ -\frac{r}{2n} + (2n+1) \left( 1 - \frac{c^2}{4n} \right) \right\} \omega \otimes \omega$$

If, in addition,  $n=1$ , then the curvature tensor is given by

$$(4.3) \quad R(X, Y)Z = \left\{ \frac{r}{2} - 2 \left( 1 - \frac{c^2}{4} \right) \right\} \cdot \{g(Y, Z)X - g(X, Z)Y\} + \left\{ 3 \left( 1 - \frac{c^2}{4} \right) - \frac{r}{2} \right\} \cdot \{g(Y, Z)\omega(X)X_0 + \omega(Y)\omega(Z)X - g(X, Z)\omega(Y)X_0 - \omega(X)\omega(Z)Y\}.$$

*Proof.* Let  $(X_0, E_i, \phi E_i)$  be an orthonormal  $\phi$ -basis. From (4.1),  $S(X_0, X_0) = a + b$ . Besides (see [3])

$$\begin{aligned} S(X_0, X_0) &= 2n - \text{trace} \left( \frac{1}{2} L_{X_0} \phi \right)^2 = 2n - \frac{1}{4} |L_{X_0} \phi|^2 \\ &= 2n - \frac{1}{4} |L_{X_0} g|^2 = 2n \left( 1 - \frac{c^2}{4n} \right). \end{aligned}$$

Consequently

$$(4.4) \quad a + b = 2n - c^2/2.$$

Moreover (4.1) implies

$$(4.5) \quad r = S(X_0, X_0) + 2 \sum_1^n S(E_i, E_i) = (2n+1)a + b.$$

From (4.4) and (4.5) we get

$$a = \frac{r}{2n} - 1 + \frac{c^2}{4n} \quad \text{and} \quad b = -\frac{r}{2n} + (2n+1) \left( 1 - \frac{c^2}{4n} \right).$$

Finally, when  $n=1$ , the curvature tensor is given by

$$(4.6) \quad R(X, Y)Z = S(Y, Z)X + g(Y, Z)Q(X) - S(X, Z)Y - g(X, Z)Q(Y) - \frac{r}{2} \{g(Y, Z)X - g(X, Z)Y\}$$

where  $Q = S^* = (S^\flat)$  is the Ricci curvature operator. So (4.3) follows from (4.2) and (4.6).

*Remark 4.1.* If  $M$  is an Einstein contact manifold, by (4.2) the scalar curvature  $r \leq 2n(2n+1)$  where the equality holds if and only if  $M$  is a  $K$ -contact Einstein manifold.



PROPOSITION 4.2. *Let  $M$  be a three-manifold with contact metric structure  $(\omega, g, X_0, \phi)$ . Let  $S_1$  be the tensor defined by*

$$(4.7) \quad S_1 = S - ag - b\omega \otimes \omega,$$

where

$$a = \left( \frac{r}{2} - 1 + \frac{1}{4}c^2 \right) \quad \text{and} \quad b = \left( -\frac{r}{2} + 3 - \frac{3}{4}c^2 \right).$$

Then

- (j)  $|S_1|^2 = 2|\sigma|^2 + \frac{1}{4}|\nabla_{X_0}\tau|^2$  where  $\sigma = S(X_0, \cdot)_{|B}$ ;
- (jj)  $\langle S_1, \tau \rangle = \langle S, \tau \rangle = -\frac{1}{2}\langle \nabla_{X_0}\tau, \tau \rangle$ ;
- (jjj) if  $\nabla_{X_0}\tau = 0$  or  $\nabla_{X_0}\tau = 2\tau \cdot \phi$  holds, then  $S$  and  $S_1$  are perpendicular to  $\tau$ ;
- (jv)  $\langle S_1, \nabla_{X_0}\tau \rangle = \langle S, \nabla_{X_0}\tau \rangle = -\frac{1}{2}|\nabla_{X_0}\tau|^2$ .

*Proof.* (j) By a direct computation we get

$$|S_1|^2 = |S|^2 + 3a^2 + b^2 - 2ar - 4b\left(1 - \frac{1}{4}c^2\right) + 2ab$$

and hence

$$(4.8) \quad |S_1|^2 = |S|^2 - \frac{1}{2}\left(r - 2 + \frac{1}{2}c^2\right)^2 - 4\left(1 - \frac{1}{4}c^2\right)^2.$$

If  $(E, \phi E, X_0)$  is an arbitrary  $\phi$ -basis, from (3.4) we get

$$(4.9) \quad S(\phi E, \phi E) - S(E, E) = (\nabla_{X_0}\tau)(E, E).$$

From (3.5), by putting  $E' = \phi E$ , we obtain

$$(4.10) \quad S(E, \phi E) = -\frac{1}{2}(\nabla_{X_0}\tau)(E, \phi E)$$

Moreover the scalar curvature is given by

$$r = S(E, E) + S(\phi E, \phi E) + S(X_0, X_0) = 2S(E, E) + (\nabla_{X_0}\tau)(E, E) + S(X_0, X_0),$$

from which

$$S(E, E) = \frac{r}{2} - \left(1 - \frac{1}{4}c^2\right) - \frac{1}{2}(\nabla_{X_0}\tau)(E, E)$$

$$S(\phi E, \phi E) = \frac{r}{2} - \left(1 - \frac{1}{4}c^2\right) + \frac{1}{2}(\nabla_{X_0}\tau)(E, E).$$

It follows that

$$\begin{aligned} |S|^2 &= S(X_0, X_0)^2 + S(E, E)^2 + S(\phi E, \phi E)^2 + 2S(E, \phi E)^2 + 2S(X_0, E)^2 + 2S(X_0, \phi E)^2 \\ &= 4\left(1 - \frac{1}{4}c^2\right)^2 + \frac{1}{2}\left(r - 2 + \frac{1}{2}c^2\right)^2 + \frac{1}{4}|\nabla_{X_0}\tau|^2 + 2|\sigma|^2. \end{aligned}$$

So, by (4.8) we obtain (j).

(jj) Let  $(X_0, E, \phi E)$  be an arbitrary  $\phi$ -basis. Using (2.1), (2.2), (4.7), (4.9) and (4.10) we get

$$\begin{aligned} \langle S_1, \tau \rangle &= \langle S, \tau \rangle - a\langle g, \tau \rangle - b\langle \omega \otimes \omega, \tau \rangle = \langle S, \tau \rangle \\ &= S(E, E)\tau(E, E) + 2S(E, \phi E)\tau(E, \phi E) + S(\phi E, \phi E)\tau(\phi E, \phi E) \\ &= -\tau(E, E)\{S(\phi E, \phi E) - S(E, E)\} + 2S(E, \phi E)\tau(E, \phi E) \\ &= -\tau(E, E)(\nabla_{X_0}\tau)(E, E) - \tau(E, \phi E)(\nabla_{X_0}\tau)(E, \phi E) = -\frac{1}{2}\langle \nabla_{X_0}\tau, \tau \rangle. \end{aligned}$$

(jjj) is a consequence of (jj) and (2.2).

(jv) is obtained like (jj) by using (i)-(iii) of Proposition 3.1.

Combining Theorem 3.2, Proposition 4.1 and (j) of Proposition 4.2 we obtain the following result.

**THEOREM 4.3.** *Let  $M$  be a compact three-manifold with contact metric structure  $(\omega, g)$ . Then  $g$  is  $\omega$ -Einstein if, and only if,  $g$  is a critical point of  $\mathcal{F}$  and  $\sigma=0$ .*

*Remark 4.2.* The condition  $\sigma=0$  means (see [9] p. 372) that in the space  $\mathcal{R}$  of all Riemannian metrics on  $M$ , the tangent vector  $\phi \in T_g(\mathcal{R})$ ,  $\phi(X, Y) = -\tau(X, \phi Y)$ , is perpendicular to the orbit of  $g$  under the group of diffeomorphisms of  $M$ .

**THEOREM 4.4.** *Let  $(M, \omega, g)$  be a contact Riemannian three-manifold. Then  $\sigma=0$  if and only if  $S_1 = -\frac{1}{2}\nabla_{X_0}\tau$ , that is, the Ricci tensor is given by*

$$(4.11) \quad S = -\frac{1}{2}\nabla_{X_0}\tau + \left(\frac{r}{2} - 1 + \frac{1}{4}c^2\right)g + \left(-\frac{r}{2} + 3 - \frac{3}{4}c^2\right)\omega \otimes \omega.$$

*Proof.* Let  $T$  be the tensor defined by

$$T = S_1 + \frac{1}{2}\nabla_{X_0}\tau.$$

Then

$$|T|^2 = |S_1|^2 + \frac{1}{4}|\nabla_{X_0}\tau|^2 + \langle \nabla_{X_0}\tau, S_1 \rangle,$$

and hence (j) and (jv) of Proposition 4.2 imply

$$|T|^2 = 2|\sigma|^2.$$

So Theorem 4.4 follows from (4.7).

**THEOREM 4.5.** *Let  $(M, \omega, g)$  be a compact contact Riemannian three-manifold. Then  $g$  is a critical metric for  $\mathcal{E}_c$  and  $\sigma=0$  if and only if  $S_1=\phi$ , that is, the Ricci tensor is given by*

$$(4.12) \quad S = \phi + \left(\frac{r}{2} - 1 + \frac{1}{4}c^2\right)g + \left(-\frac{r}{2} + 3 - \frac{3}{4}c^2\right)\omega \otimes \omega.$$

*Proof.* If  $g$  is a critical metric for  $\mathcal{E}_c$  and  $\sigma=0$ , from Theorem 4.4 above and Theorem 5.1 of [15] we get (4.12). Conversely assume (4.12), since  $\phi(X_0, \cdot) = 0$ , we have  $\sigma=0$ . So (4.11), (4.12) and Theorem 5.1 of [15] imply that  $g$  is critical for  $\mathcal{E}_c$ .

**5. Curvature of  $K$ -contact three-manifolds.** In [10] the following was proved.

**THEOREM 5.1.** *Let  $M$  be a compact three-manifold with  $K$ -contact metric structure  $(\omega, g)$ . If the scalar curvature  $r > -2$  or the Webster curvature  $W > 0$ , then  $M$  admits a  $K$ -contact metric structure  $(\tilde{\omega} = a\omega, \tilde{g} = ag + (a^2 - a)\omega \otimes \omega)$  of positive sectional curvature for some  $a, 0 < a \leq 1$ .*

This result is relative to the question posed by S.S. Chern (cf. appendix of [7]) of determining those compact three-manifolds admitting a contact metric structure  $(\omega, g)$  for which the torsion invariant  $|\tau|$  is identically zero (i. e. the contact metric structure is  $K$ -contact). In this section we extend Theorem 5.1; precisely we give the following.

**THEOREM 5.2.** *Let  $M$  be a compact three-manifold with  $K$ -contact metric structure  $(\omega, g)$ . If one of the following four conditions holds: (a)  $W > 0$ , (b)  $r > -2$ , (c)  $S + 2g > 0$ , (d) the  $\phi$ -sectional curvature  $H > -3$ , then  $M$  admits a  $K$ -contact metric structure  $(\tilde{\omega}, \tilde{g})$  of positive sectional curvature.*

This Theorem is a consequence of Theorem 5.1 and of the following Proposition.

**PROPOSITION 5.3.** *Let  $M$  be a contact  $\omega$ -Einstein three-manifold with contact metric structure  $(\omega, g, X_0, \phi)$ . Then, if  $c < 2$ , the following five conditions are equivalent:*

- (a)  $W > c^2/8$ , (b)  $r + 2 > c^2/2$ , (c)  $S + 2g > (c^2/2)g$ ,
- (d) the sectional curvature  $K > -3(1 - c^2/4)$ ,
- (e) the  $\phi$ -sectional curvature  $H > -3(1 - c^2/4)$ .

*Proof.* Since  $8W = r + 2 + c^2/2$ , (a) and (b) are equivalent. If  $X$  is vertical,

that is, if  $X=tX_0$ , then

$$\{S+2(1-c^2/4)g\}(X, X)=4t^2(1-c^2/4)>0.$$

If  $X$  is horizontal, that is, if  $\omega(X)=0$ , then by (4.2)

$$\{S+2(1-c^2/4)g\}(X, X)=(r/2+1-c^2/4)g(X, X)>0.$$

On the other hand  $S(X_0, \cdot)_{,B}=\sigma=0$ , so (b) and (c) are equivalent. For each point  $x \in M$ , we consider an arbitrary plane  $P$  of  $T_x(M)$  and an orthonormal basis  $(X, Y)$  of  $P$  with  $Y=P \cap B$ . Then, by (4.3), the sectional curvature  $K(P)$  at  $x$  is given by

$$\begin{aligned} K(P) &= g(R(X, Y)Y, X) = (r/2 - 2 + c^2/2) + (-r/2 + 3 - 3c^2/4)g(X_0, X)^2 \\ &= (r/2 - 2 + c^2/2)\sin^2(X, X_0) + (1 - c^2/4)\cos^2(X, X_0) \end{aligned}$$

and hence

$$(5.1) \quad K(P) = (r/2 - 2 + c^2/2)\cos^2\alpha + (1 - c^2/4)\sin^2\alpha$$

where  $\alpha$  is the angle between  $P$  and  $B$ . By (5.1) we get that (b) implies (d). The converse is trivial. Moreover the scalar curvature at  $x$  is given by

$$r = \text{trace } S = 2S(X_0, X_0) + 2g(R(E, \phi E)\phi E, E),$$

where  $E$  is an unit vector of  $B$ , that is

$$(5.2) \quad r = 4(1 - c^2/4) + 2H.$$

Therefore (b) and (e) are equivalent.

*Remark 5.1.* (i) The main result of [7] says that every contact structure on a compact orientable three-manifold has a contact metric whose Webster curvature  $W$  is either  $>0$  or  $W = \text{const.} \leq 0$ .

(ii) Hamilton [11] showed that a metric  $g$  of positive Ricci curvature on a compact three-manifold can be deformed to a metric of (positive) constant curvature. Hence in Theorem 5.2 if, in addition,  $M$  is simply-connected, then it is diffeomorphic with the three-sphere. This extends Corollary of [8] (cf. p. 654).

(iii) The formula (5.2) holds for every metric  $g \in \mathcal{M}(\omega)$ . So the conditions on the scalar curvature given in [10] can be replaced by conditions on the  $\phi$ -sectional curvature  $H$ .

(iv) Let  $(M, g)$  be a compact Riemannian manifold and  $\mathcal{S}^2$  the space of all symmetric tensor fields of type  $(0, 2)$ . Berger and Ebin (cf. [1] §6) introduced a zero-order differential operator  $\mathcal{K}: \mathcal{S}^2 \rightarrow \mathcal{S}^2$  which is related to the rough Laplacian and to the Lichnerowicz operator. They proved that the operator  $\mathcal{K}$  is positive definite on  $TZ = \{D \in \mathcal{S}^2 : \text{trace } D = 0\}$  if  $(M, g)$  is of strictly positive

sectional curvature. Now observe that if  $(M, \omega, g)$  is a compact three-manifold as in Theorem 5.2, then  $M$  admits a contact metric structure  $(\tilde{\omega}, \tilde{g})$  for which the corresponding operator  $\tilde{K}$  is positive definite on  $T_{\tilde{g}}(\mathcal{N}(\tilde{\omega}))=TZ$ , where  $\mathcal{N}(\tilde{\omega})$  is the set of all Riemannian metrics on  $M$  which have the same volume element of the metrics of  $\mathcal{M}(\tilde{\omega})$ .

## REFERENCES

- [ 1 ] M. BERGER AND D. EBIN, Some decomposition of the space of symmetric tensors on a Riemannian manifold, *J. Diff. Geom.* **3** (1969), 379-392.
- [ 2 ] D. E. BLAIR AND A. J. LEDGER, Critical associated metrics on contact manifolds II, *J. Austral. Math. Soc.* **41** (1986), 404-410.
- [ 3 ] D. E. BLAIR, Contact manifolds in Riemannian Geometry, *Lect. Notes in Math.*, 509, Springer-Verlag, Berlin and New York, 1976.
- [ 4 ] D. E. BLAIR, Critical associated metrics on contact manifolds III, preprint.
- [ 5 ] D. E. BLAIR, When is the tangent sphere bundle locally symmetric ?, preprint.
- [ 6 ] D. E. BLAIR, On the set of metrics associated to a symplectic or contact form, *Bull. Inst. Math. Acad. Sinica* **11** (1983), 297-308.
- [ 7 ] S. S. CHERN AND R. S. HAMILTON, On Riemannian metrics adapted to three-dimensional contact manifolds, *Lect. Notes in Math.*, 1111, Springer-Verlag, Berlin and New York, 1985, 279-305.
- [ 8 ] S. I. GOLDBERG, Nonnegatively curved contact manifolds, *Proc. A. M. S.* **96** (1986), 651-656.
- [ 9 ] S. I. GOLDBERG AND G. TOTH, Torsion and deformation of contact metric structures on 3-manifolds, *Tohoku Math. J.* **39** (1987), 365-372.
- [10] S. I. GOLDBERG, D. PERRONE AND G. TOTH, Contact three-manifolds with positive generalized Tanaka-Webster scalar curvature, *C. R. Math. Rep. Acad. Sci. Canada* vol. X, **6** (1988), 255-260.
- [11] R. S. HAMILTON, Three-manifolds with positive Ricci curvature, *J. Diff. Geom.* **17** (1982), 255-306.
- [12] J. MARTINET, Formes de contact sur les variétés de dimension 3, *Proc. Liverpool Singularities Sympos. II*, Springer Lect. Notes in Math. **209** (1971), 142-163.
- [13] H. SATO, Remarks concerning contact manifolds, *Tohoku Math. J.* **29** (1977), 577-584.
- [14] S. TANNO, Locally symmetric  $K$ -contact Riemannian manifolds, *Proc. Japan Acad.* **43** (1968), 581-583.
- [15] S. TANNO, Variational problems on contact Riemannian manifolds, *Trans. Amer. Math. Soc.* **314** (1989), 349-379.

DIPARTIMENTO DI MATEMATICA  
 FACOLTÀ DI SCIENZE  
 UNIVERSITÀ DEGLI STUDI DI LECCE  
 VIA ARNESANO  
 73100 LECCE, ITALY