

# Towards extracellular $\text{Ca}^{2+}$ sensing by MRI: synthesis and calcium-dependent $^1\text{H}$ and $^{17}\text{O}$ relaxation studies of two novel bismacrocylic $\text{Gd}^{3+}$ complexes

Kirti Dhingra · Petra Fousková · Goran Angelovski ·  
Martin E. Maier · Nikos K. Logothetis · Éva Tóth

Received: 20 July 2007 / Accepted: 29 August 2007 / Published online: 15 September 2007  
© SBIC 2007

**Abstract** Two new bismacrocylic  $\text{Gd}^{3+}$  chelates containing a specific  $\text{Ca}^{2+}$  binding site were synthesized as potential MRI contrast agents for the detection of  $\text{Ca}^{2+}$  concentration changes at the millimolar level in the extracellular space. In the ligands, the  $\text{Ca}^{2+}$ -sensitive BAPTA-bisamide central part is separated from the DO3A macrocycles either by an ethylene ( $\text{L}^1$ ) or by a propylene ( $\text{L}^2$ ) unit [ $\text{H}_4\text{BAPTA}$  is 1,2-bis(*o*-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid;  $\text{H}_3\text{DO}_3\text{A}$  is 1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid]. The sensitivity of the  $\text{Gd}^{3+}$  complexes towards  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  was studied by  $^1\text{H}$  relaxometric titrations. A maximum relaxivity increase of 15 and 10% was observed upon  $\text{Ca}^{2+}$  binding to  $\text{Gd}_2\text{L}^1$  and  $\text{Gd}_2\text{L}^2$ , respectively, with a distinct selectivity of

$\text{Gd}_2\text{L}^1$  towards  $\text{Ca}^{2+}$  compared with  $\text{Mg}^{2+}$ . For  $\text{Ca}^{2+}$  binding, association constants of  $\log K = 1.9$  ( $\text{Gd}_2\text{L}^1$ ) and  $\log K = 2.7$  ( $\text{Gd}_2\text{L}^2$ ) were determined by relaxometry. Luminescence lifetime measurements and UV–vis spectrophotometry on the corresponding  $\text{Eu}^{3+}$  analogues proved that the complexes exist in the form of monohydrated and nonhydrated species;  $\text{Ca}^{2+}$  binding in the central part of the ligand induces the formation of the monohydrated state. The increasing hydration number accounts for the relaxivity increase observed on  $\text{Ca}^{2+}$  addition. A  $^1\text{H}$  nuclear magnetic relaxation dispersion and  $^{17}\text{O}$  NMR study on  $\text{Gd}_2\text{L}^1$  in the absence and in the presence of  $\text{Ca}^{2+}$  was performed to assess the microscopic parameters influencing relaxivity. On  $\text{Ca}^{2+}$  binding, the water exchange is slightly accelerated, which is likely related to the increased steric demand of the central part leading to a destabilization of the Ln–water binding interaction.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00775-007-0296-9) contains supplementary material, which is available to authorized users.

K. Dhingra · G. Angelovski · N. K. Logothetis (✉)  
Max Planck Institute for Biological Cybernetics,  
Tübingen, Germany  
e-mail: nikos.logothetis@tuebingen.mpg.de

P. Fousková · É. Tóth (✉)  
Centre de Biophysique Moléculaire,  
CNRS, rue Charles Sadron,  
45071 Orléans Cedex 2, France  
e-mail: eva.jakabtoth@cnrs-orleans.fr

M. E. Maier  
Institut für Organische Chemie,  
Universität Tübingen,  
Tübingen, Germany

N. K. Logothetis  
Imaging Science and Biomedical Engineering,  
University of Manchester,  
Manchester, UK

**Keywords** Relaxivity · MRI contrast agents · Calcium sensitive · Lanthanide complexes · Luminescence

## Introduction


Magnetic resonance imaging (MRI) is one of the most versatile techniques in clinical and experimental in vivo imaging. It provides excellent three-dimensional spatial resolution down to the 10- $\mu\text{m}$  range, whole body coverage and the opportunity to noninvasively measure additional physiological parameters [1]. Its sensitivity is, however, 2–6 orders of magnitude lower than that of optical imaging. This can be overcome by using appropriate MRI contrast agents. Nowadays, more than 30% of

MRI examinations benefit from the use of paramagnetic contrast agents [2]. By accelerating the longitudinal or transverse relaxation of water protons, they increase the signal-to-noise ratio, which, in turn, positively influences the image contrast. The currently used clinical contrast agents are mainly  $\text{Gd}^{3+}$  chelates, making use of the high magnetic moment (seven unpaired electrons) and slow electronic relaxation of this paramagnetic lanthanide ion [3, 4]. These contrast agents are, however, mostly non-specific and provide only anatomical information. Nowadays, there is an active search for new classes of contrast agents capable of reflecting a change in biological activity in their local environment. A new field, molecular imaging, is being developed, with the aim of in vivo visualization of molecular events. Any molecular imaging procedure requires an imaging probe that is specific for a given molecular event. Examples of such “smart” potential contrast agents, sensing variables like pH [5–7], partial oxygen pressure [8, 9], ion and metabolite concentration [10–14], or enzyme activity [15, 16] have been described in the literature. They function on the basis of the modulation of one or more microscopic parameters of the  $\text{Gd}^{3+}$  chelate determining relaxivity, induced by a change in the local environment of the agent. Among the variables that influence the relaxivity of a contrast agent, the most important are the applied magnetic field ( $B_0$ ), the number of water molecules directly bound to the paramagnetic center ( $q$ ), their exchange rate ( $k_{ex}$ ), the water proton– $\text{Gd}^{3+}$  distance ( $r_{\text{GdH}}$ ), the electron spin relaxation times ( $T_{1e}$ ,  $T_{2e}$ ), the rotational correlation time ( $\tau_R$ ), and the presence and the number of water molecules in the so-called second coordination sphere ( $q^{2\text{nd}}$ ) [2, 4, 17, 18].

Calcium ( $\text{Ca}^{2+}$ ) is one of the most important metal ions for life as it controls muscular contraction, neural cell communication, hormonal secretion, etc. [19]. In the brain,  $\text{Ca}^{2+}$  acts as an important second messenger [20]. Synaptic transmission depends on the entry of calcium ions at the presynaptic terminal to cause fusion and release of vesicles. Significant changes in  $\text{Ca}^{2+}$  concentration take place during neuronal activity [21]. Tracking the dynamics of  $\text{Ca}^{2+}$  concentration changes could thus contribute to the understanding of basic aspects of neuronal regulation or to highlight the abnormalities in the diseased state [22]. Optical imaging based on fluorescent dyes possessing a calcium-dependent response has been widely used at the cellular or cell population level [23]. However, the intrinsic depth limitations associated with optical imaging restrict these applications to superficial regions. In contrast to optical imaging, MRI has no depth penetration limits and could be more adapted for tracking  $\text{Ca}^{2+}$  changes. Li et al. [11, 24] have proposed a potential MRI contrast agent, GdDOPTA, whose relaxivity was reported to be

sensitive to  $\text{Ca}^{2+}$  concentration in the 0.1–10- $\mu\text{M}$  range. The two macrocyclic subunits of the dimeric ligand DOPTA serve as lanthanide(III) ion chelators, while the  $\text{BAPTA}^{4-}$  subunit bridging the macrocycles is known for its high selectivity towards  $\text{Ca}^{2+}$  [ $\text{H}_4\text{BAPTA}$  is 1,2-bis(*o*-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid]. The mechanism proposed for the  $\text{Ca}^{2+}$ -dependent relaxivity change of GdDOPTA accounts for changes in the number of water molecules directly bound to the paramagnetic metal ion ( $q$ ) upon  $\text{Ca}^{2+}$  binding to the  $\text{BAPTA}^{4-}$  moiety.  $\text{BAPTA}^{4-}$  was proposed a long time ago as a selective fluorescent indicator for  $\text{Ca}^{2+}$  [25]. It presents several advantages for biological  $\text{Ca}^{2+}$  detection by optical methods: integrated aromatic chromophore moiety and pH insensitivity of  $\text{Ca}^{2+}$  complexation thanks to the low values of the protonation constants (6.36 and 5.47, both below physiological pH). More recently, another calcium-sensitive MRI probe was proposed by Atanasijevic et al. [20]. It is based on superparamagnetic iron oxide nanoparticles and calmodulin. In this approach, calcium-dependent protein–protein interactions lead to particle clustering and consequent changes in transverse relaxivity. Similarly to GdDOPTA, this agent had calcium-response in the micromolar  $\text{Ca}^{2+}$  concentration range. It was therefore proposed to detect elevated intracellular  $\text{Ca}^{2+}$  levels, provided it is combined to a cellular delivery system.

With the objective of detecting *extracellular*  $\text{Ca}^{2+}$  concentration changes by MRI methods, we synthesized two novel potential bismacrocyclic MRI contrast agents (Structure 1). Two  $\text{Gd}^{3+}$ -loaded 1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid (DO3A) derived units were connected by a BAPTA-derived linker capable of  $\text{Ca}^{2+}$  binding. However, in comparison with  $\text{BAPTA}^{4-}$ , the  $\text{Ca}^{2+}$  binding moiety in the central part of the ligand was designed to have a reduced affinity towards  $\text{Ca}^{2+}$  in order to shift to higher  $\text{Ca}^{2+}$  concentrations the range where the probe is expected to have a relaxivity response. The conditional stability constant of GdDOPTA–Ca at physiological pH was reported to be  $1.0 \times 10^6 \text{ M}^{-1}$  [11]. Similarly, the fluorescent  $\text{Ca}^{2+}$  indicators used in biological applications and based on  $\text{BAPTA}^{4-}$  are adapted to assess the cytosolic free  $\text{Ca}^{2+}$  concentration in the micromolar range. In contrast, the extracellular concentration of  $\text{Ca}^{2+}$  is in the millimolar range. The relaxivity response of a probe with affinity of  $K \sim 10^6 \text{ M}$  for  $\text{Ca}^{2+}$  would have already leveled off at the millimolar level. The  $\text{BAPTA}^{4-}$  ligand was therefore modified by replacing two carboxylates by the less strong donor amide functions. As for the macrocyclic  $\text{Gd}^{3+}$  binding site, it was designed to respond to the presence of  $\text{Ca}^{2+}$  by modulation of the hydration number with consequent change in the relaxivity.

**Structure 1**

The proton relaxivities of the two new complexes were studied by relaxometric titrations at different  $\text{Ca}^{2+}$  concentrations and the relaxivity modulation by the  $\text{Ca}^{2+}$  concentration changes was confirmed. We specifically aimed to identify the microscopic factors responsible for the  $\text{Ca}^{2+}$ -dependent relaxivity response. To this end, a detailed mechanistic study was carried out. We used UV–vis absorption and luminescence lifetime measurements on the corresponding  $\text{Eu}^{3+}$  complexes to assess the number of inner-sphere water molecules and their variation on  $\text{Ca}^{2+}$  addition. The  $\text{Gd}_2\text{L}^1$  complex was characterized by variable-temperature  $^{17}\text{O}$  NMR spectroscopy and variable-field relaxivity measurements (nuclear magnetic relaxation dispersion, NMRD). The experimental data were evaluated on the basis of the Solomon–Bloembergen–Morgan theory, extended with a second-sphere contribution.

## Experimental

### Materials and methods

All chemicals were purchased at the purest grade commercially available and were used without further purification. Cyclen was purchased from Strem, France. Bromoethylamine hydrogen bromide, bromopropylamine hydrobromide, *tert*-butyl bromoacetate, *N*-methylpyrrolidone (dry), pyridine,  $\text{H}_4\text{BAPTA}$ , acetic anhydride and *N,N*-dimethylformamide (extra dry) were purchased from Sigma–Aldrich, Germany. Toluene, acetonitrile, dichloromethane and methanol were purchased as analytical grade solvents from Acros Organics, Germany. Lanthanide(III) chlorides were prepared by dissolving metal(III) oxides (Aldrich) in a slight excess of HCl (Carlo Erba, 37%) followed by evaporation of solvents and dissolution in  $\text{H}_2\text{O}$ .

High-performance liquid chromatography (HPLC) was performed at room temperature using a Varian (Australia) PrepStar instrument equipped with a PrepStar 335 photodiode array detector at 254 nm. Reversed-phase analytical

HPLC was performed in a stainless steel ChromSep (length 250 cm, internal diameter 4.6 mm, outside diameter 3/8 in. and particle size 8  $\mu\text{m}$ ) C18 column (Varian). The ligands were purified using the following gradient:

Method A: 40% solvent A (methanol) and 60% solvent B (water) to 100% solvent B in 5 min running isocratically at 100% solvent B for 10 min and then to 60% solvent B in the next 2 min.

Method B: 95% solvent A (acetonitrile, 0.1% HCOOH) and 5% solvent B (water, 0.1% HCOOH) to 70% solvent B in 10 min and then 100% in the next 8 min running isocratically for 12 min after that and then to 5% in next 2 min.

The flow rate used for analytical HPLC was 1  $\text{mL min}^{-1}$  and for preparative 65  $\text{mL min}^{-1}$ . All solvents used were HPLC grade filtered through a 0.45- $\mu\text{m}$  nylon-66 Millipore filter prior to use.

$^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded using a Bruker 400 MHz spectrometer ( $^1\text{H}$ —internal reference  $\text{CDCl}_3$  at 7.27 ppm or  $\text{D}_2\text{O}$  at 4.75 ppm;  $^{13}\text{C}$ —internal reference  $\text{CDCl}_3$  at 77.0 ppm or tetramethylsilane at 0 ppm). All the experiments were performed at 298 K. Electrospray ionization (ESI) low-resolution mass spectrometry (LRMS) spectra were obtained using an SL 1100 system (Agilent, Germany) with ion-trap detection in positive and negative ion mode. IR spectra were recorded with a Nicolet Impact 400 D spectrometer using neat compounds as disks with KBr and only the major bands were noted. UV–vis spectra of  $^5\text{D}_0 \leftarrow ^7\text{F}_0$  transitions of  $\text{Eu}_2\text{L}^1$  were obtained with a PerkinElmer Lambda 19 spectrometer in the region 577–581 nm with data steps of 0.05 nm [26]. The concentrations of the samples were approximately 0.02 M and the temperature dependence was measured in the interval 288–323 K in the absence and presence of  $\text{Ca}^{2+}$ . To maintain a constant temperature, thermostatizable cells with a 10-cm optical length were used. The luminescence measurements were performed with a Varian Eclipse spectrofluorimeter, equipped with a 450-W xenon arc lamp, a microsecond flash lamp and a red-sensitive photomultiplier (300–850 nm). The luminescence spectra were obtained after excitation of the  $^5\text{L}_6 \leftarrow ^7\text{F}_0$  band (394 nm). An aqueous solution (0.02 M) of  $\text{Eu}_2\text{L}^1$  was measured at 298 K in the absence and presence of  $\text{Ca}^{2+}$ .

The  $^1\text{H}$  NMRD profiles were recorded at the Laboratory of Inorganic and Bioinorganic Chemistry, Ecole Polytechnique Fédérale de Lausanne, Switzerland, using a Stellar Spinmaster FFC fast-field-cycling relaxometer covering magnetic fields from  $2.35 \times 10^{-4}$  to 0.47 T (proton Larmor frequency range 0.01–20 MHz). The temperature was controlled by a VTC90 temperature control unit and fixed by a gas flow. At higher fields, the relaxivity was recorded using Bruker Minispecs mq30 (30 MHz), mq40

(40 MHz) and mq60 (60 MHz), on a Bruker 4.7 T (200 MHz) cryomagnet connected to a Bruker Avance 200 console and with a Bruker Avance 500 spectrometer (500 MHz). The temperature was measured by a substitution technique [27] or via a preliminary calibration using methanol and ethylene glycol standards [28]. The longitudinal ( $1/T_1$ ) and transverse ( $1/T_2$ )  $^{17}\text{O}$  NMR relaxation rates were measured in the temperature range 277–344 K. The data were recorded using a Bruker Avance 500 (11.75 T, 67.8 MHz) spectrometer. The temperature was calculated according to a previous calibration with ethylene glycol and methanol [28]. The samples were measured in 5-mm NMR tubes and were enriched with *tert*-butanol to allow for the bulk magnetic susceptibility correction [29]. The  $1/T_1$  data were obtained by the inversion recovery method, while the  $1/T_2$  data were measured by the Carr–Purcell–Meiboom–Gill spin-echo technique. Acidified water ( $\text{HClO}_4$ , pH 3.8) was used as an external reference. Analyses of the  $^{17}\text{O}$  NMR and  $^1\text{H}$  NMRD experimental data were performed with the Visualiseur/Optimiseur programs running on a MATLAB platform version 6.5 [30, 31].

## Synthesis

Compound **4a** was synthesized according to a previously reported procedure [32]. With use of an analogous method, compound **4b** was synthesized via **3b** from the substrate **1b** [33].

Compound **3b**.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz),  $\delta$  (ppm): 6.97–6.84 (m, 5H), 4.65 (s, 2H), 2.98–2.89 (m, 2H), 2.86–2.70 (m, 5H), 2.70–2.60 (m, 3H), 2.58–2.17 (m, 7H), 2.16–1.75 (m, 9H), 1.32 (br s, 2H), 1.06–1.03 (m, 27H).  $^{13}\text{C}$  NMR 172.2, 171.2, 155.2, 152.9, 135.6, 127.1, 126.4, 81.3, 81.0, 65.7, 64.7, 55.4, 50.3, 48.9, 38.5, 37.8, 28.2, 26.7, 26.5, 25.2. ESI–MS  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{37}\text{H}_{63}\text{N}_5\text{O}_8$ , 705.5; found, 706.5.

Compound **4b**.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz),  $\delta$  (ppm): 8.19 (br s, 2H), 3.42–2.99 (m, 8H), 2.85–2.20 (m, 16H), 1.61 (br s, 2H), 1.41 (s, 27H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 100 MHz),  $\delta$  (ppm): 172.0, 169.8, 81.8, 81.1, 57.2, 55.9, 50.1, 49.4, 48.9, 38.5, 27.2, 27.1, 22.9. ESI–MS  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{29}\text{H}_{57}\text{N}_5\text{O}_6$ , 571.4; found, 572.4.

## General method for the synthesis of compounds **6a** and **6b**

Compound **4a** or **4b** (2 mmol) was dissolved in 5 mL of dry *N*-methylpyrrolidinone with 50  $\mu\text{L}$  of dry  $\text{Et}_3\text{N}$  and heated at 333 K for 15 min. Anhydride **5** [34] was then added to this reaction mixture (295 mg, 0.67 mmol) in small lots under  $\text{N}_2$ . After complete addition of **5**, the

solution was kept under continuous stirring at 333 K overnight. It was then evaporated, dissolved in dichloromethane and extracted with water. The organic layer was collected and evaporated to get a yellow oil. The crude product was then purified by reversed-phase HPLC using method A to get a light yellow fluffy solid.

1,2-Bis{[2-{{[1-[1,4,7-tris(*tert*-butoxycarbonylmethyl)-1,4,7,10-tetraazacyclododecane-10-yl]eth-2-yl]amino}carbonyl]methyl}-(carboxymethyl)amino]phenoxy}ethane, **6a**. Yield: 573 mg (55%)  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz),  $\delta$  (ppm): 7.01–6.98 (m, 2H), 6.92–6.89 (m, 2H), 6.77–6.74 (m, 4H), 4.56 (d,  $J = 6.4$  Hz, 2H, *CHHO*), 4.34 (d,  $J = 15.5$  Hz, 2H, *CHHCONH*), 4.01 (d,  $J = 17.8$  Hz, 2H, *CHHCOOH*), 3.91 (d,  $J = 6.4$  Hz, 2H, *CHHO*), 3.48 (d,  $J = 17.8$  Hz, 2H, *CHHCOOH*), 3.36–3.21 (m, 22H), 3.08–2.80 (m, 12H), 2.78–2.69 (m, 6H), 2.76–2.56 (m, 18H), 1.47 (s, 36H), 1.43 (s, 18H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 100 MHz),  $\delta$  (ppm): 175.2 (*CONH*), 175.0 (*CONH*), 170.2, 170.1, 149.4, 139.2, 120.6, 119.5, 115.4, 112.1, 81.8, 81.5, 66.4 (*CH}\_2\text{O}*), 60.7 (*NCH}\_2\text{CONH}*), 58.5 (*CH}\_2\text{COOH}*), 56.5, 54.8, 53.5, 51.3, 50.3, 49.7, 47.8, 32.1 (*CH}\_2\text{CH}\_2\text{CONH}*), 29.6, 28.1. ESI–MS  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{78}\text{H}_{130}\text{N}_{12}\text{O}_{20}$ , 1,554.9; found, 1,555.9.

1,2-Bis{[3-{{[1-[1,4,7-tris(*tert*-butoxycarbonylmethyl)-1,4,7,10-tetraazacyclododecane-10-yl]prop-3-yl]amino}carbonyl]methyl}-(carboxymethyl)amino]phenoxy}ethane, **6b**. Yield: 424 mg (40%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz),  $\delta$  (ppm): 7.08–7.06 (m, 2H), 6.91–6.89 (m, 2H), 6.85–6.82 (m, 4H), 4.49 (br s, 2H), 4.31 (br s, 4H), 4.08–3.73 (m, 8H), 3.57–3.09 (m, 20H), 3.04–2.47 (m, 32H), 2.40 (br s, 4H), 2.01 (br s, 2H), 1.53 (s, 18H), 1.48 (s, 36H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 100 MHz),  $\delta$  (ppm): 175.7, 173.4, 170.9, 170.2, 150.2, 139.8, 120.3, 120.2, 115.7, 112.2, 82.2, 82.0, 66.6, 61.0, 58.7, 56.5, 55.3, 49.6, 49.1, 47.6, 35.5, 28.6, 28.4, 22.2. ESI–MS  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{80}\text{H}_{134}\text{N}_{12}\text{O}_{20}$ , 1,582.9; found, 1,584.0.

## General method for the synthesis of $L^1$ and $L^2$

Neat trifluoroacetic acid (70 mL) was added to the previously obtained compound **6a** or **6b** (0.32 mmol) and the reaction was kept at room temperature for 24 h. Trifluoroacetic acid was then evaporated, and the residue was dried under vacuum and purified by reversed-phase HPLC using method B.

1,2-Bis{[2-{{[1-[1,4,7-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecane-10-yl]eth-2-yl]amino}carbonyl]methyl}-(carboxymethyl)amino]phenoxy}ethane,  $L^1$ . Yield: 220 mg (60%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz),  $\delta$  (ppm): 7.06–6.99 (m, 4H), 6.94 (t,  $J = 7.0$  Hz, 2H), 6.87–6.85 (m, 2H), 4.27 (s, 4H), 3.83 (s, 4H), 3.77 (s, 4H), 3.58 (br s, 8H), 3.45 (s, 4H), 3.28–3.12 (m, 24H), 3.02–2.97 (m, 4H),

2.82–2.75 (m, 12H).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 100 MHz),  $\delta$  (ppm): 177.6, 174.7, 170.5, 169.9, 150.1, 138.9, 122.5, 121.4, 117.4, 113.0, 67.0, 57.9, 56.7, 56.1, 55.3, 51.0, 50.6, 50.5, 49.2, 47.6, 33.5. IR ( $\text{cm}^{-1}$ ): 3,426 (vs), 2,964 (m), 2,929 (m), 2,860 (m), 1,718 (s), 1,637 (vs), 1,384 (s), 1,355 (s), 1,328 (m), 1,240 (s), 1,203 (s), 762 (m), 694 (m). ESI-MS  $m/z$ :  $[\text{M}-\text{H}]^-$  calcd for  $\text{C}_{54}\text{H}_{82}\text{N}_{12}\text{O}_{20}$ , 1,218.57; found, 1,217.7.

1,2-Bis{[3-[[[1-[1,4,7-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecane-10-yl]prop-3-yl]amino]carbonyl]methyl}-(carboxymethyl)amino]phenoxy}ethane,  $\text{L}^2$ . Yield: 126 mg from 400 mg of **6b** (40%).  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 400 MHz),  $\delta$  (ppm): 6.98–6.91 (m, 4H), 6.88 (t,  $J = 7.4$ , 2H), 6.82–6.80 (m, 2H), 4.20 (s, 4H), 3.78 (s, 4H), 3.74 (s, 4H), 3.57 (s, 4H), 3.39–3.31 (m, 8H), 3.27–3.20 (m, 8H), 3.13–2.92 (m, 20H), 2.89–2.83 (m, 8H), 2.65 (br s, 4H), 1.61–1.53 (m, 4H).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 100 MHz),  $\delta$  (ppm): 177.7, 174.4, 149.9, 138.5, 122.3, 121.5, 117.4, 113.2, 67.1, 57.6, 56.2, 56.1, 55.4, 50.9, 50.5, 49.6, 49.2, 36.4, 22.6. ESI-MS  $m/z$ :  $[\text{M}-\text{H}]^-$  calcd for  $\text{C}_{56}\text{H}_{86}\text{N}_{12}\text{O}_{20}$ , 1,246.6; found, 1,245.6.

#### Preparation of solutions of lanthanide(III) complexes

The complexes used for  $^1\text{H}$  and  $^{17}\text{O}$  relaxometric and UV–vis measurements were prepared by mixing a slight excess (5%) of the ligand solution with the appropriate lanthanide(III) chloride solution. The pH was adjusted to 7 using KOH solution and the reaction mixture was stirred at 323 K overnight. The absence of free  $\text{Ln}^{3+}$  was checked by xylenol orange indicator in HCl/urotropine buffer (pH 5.8). For each  $\text{Gd}_2\text{L}$  sample, the  $\text{Gd}^{3+}$  concentration was determined by measuring the bulk magnetic susceptibility shifts [35].

$\text{Gd}_2\text{L}^1$ . ESI-LRMS  $m/z$ :  $[\text{M}-\text{H} + \text{Na}]^-$  calcd for  $\text{C}_{78}\text{H}_{124}\text{Gd}_2\text{N}_{12}\text{O}_{20}$ , 1,525.1; found, 1,547.6. IR: 3,448 (vs), 1,602 (vs), 1,560 (s), 1,400 (m), 1,323 (m), 1,242 (m), 1,203 (vs), 1,085 (m), 762 (m), 723(m).

$\text{Eu}_2\text{L}^1$ . ESI-LRMS  $m/z$ :  $[\text{M}-\text{H}]^-$  calcd for  $\text{C}_{54}\text{H}_{76}\text{Eu}_2\text{N}_{12}\text{O}_{20}$ , 1,156.4; found 1,155.5.

$\text{Gd}_2\text{L}^2$ . ESI-LRMS  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{56}\text{H}_{80}\text{Gd}_2\text{N}_{20}\text{O}_{12}$ , 1,553.1; found, 1,553.6.

#### Relaxometric $\text{Ca}^{2+}$ titrations of $\text{Gd}_2\text{L}^1$ and $\text{Gd}_2\text{L}^2$

The titrations were performed at 11.75 T, 298 K and pH 7.3 [maintained by *N*-(2-hydroxyethyl)piperazine-*N'*-ethanesulfonic acid buffer]. A solution of  $\text{CaCl}_2$  of known concentration was added stepwise to the complex solution and the longitudinal proton relaxation time  $T_1$  was measured after each  $\text{Ca}^{2+}$  addition. The initial  $\text{Gd}^{3+}$  concentrations

were 7.4 and 4.0 mM for  $\text{Gd}_2\text{L}^1$  and  $\text{Gd}_2\text{L}^2$ , respectively. The relaxivity  $r_1$  was calculated from Eq. 1, using the actual  $\text{Gd}^{3+}$  concentration at each point of the titration:

$$\frac{1}{T_{1,\text{obs}}} = \frac{1}{T_{1,\text{d}}} + r_1[\text{Gd}], \quad (1)$$

where  $T_{1,\text{obs}}$  is the observed longitudinal relaxation time,  $T_{1,\text{d}}$  is its diamagnetic contribution in the absence of the paramagnetic substance and  $[\text{Gd}]$  is the concentration of  $\text{Gd}^{3+}$ .

#### Relaxometric $\text{Mg}^{2+}$ titration of $\text{Gd}_2\text{L}^1$

The titration was done analogously to the  $\text{Ca}^{2+}$  titration ( $B = 11.75$  T, 298 K; initial  $\text{Gd}^{3+}$  concentration of 2.6 mM). When the  $\text{Mg}^{2+}$ -to-complex ratio was 23 (in the  $\text{Ca}^{2+}$  titration, the plateau was already reached at this  $\text{Ca}^{2+}$ -to- $\text{Gd}_2\text{L}^1$  ratio), a  $\text{Ca}^{2+}$  solution was added stepwise and the relaxivity was measured.

## Results and discussion


### Synthesis of ligands

The ligands were prepared from the appropriate anhydride and amine precursors in a facile two-step procedure as shown in Scheme 1. Compounds **1a**, **1b**, **2** and **5** were synthesized according to reported procedures [32–34]. Protected amines **3a** and **3b** were synthesized by alkylating **2** with **1a** or **1b** in acetonitrile. Hydrogenation of the resulting products in methanol using 10% Pd on carbon as a catalyst yielded **4a** and **4b**. Bismacrocycles **6a** and **6b** were obtained by coupling of amines **4a** or **4b** to an anhydride **5** in *N*-methylpyrrolidinone and triethylamine. After the purification, they were hydrolyzed with neat trifluoroacetic acid and purified by reversed-phase HPLC giving the final ligands  $\text{L}^1$  and  $\text{L}^2$ . It is interesting to note that an additional proof for the extremely high steric constraints in the bismacrocyclic systems studied was observed in 2D NMR (correlation spectroscopy, heteronuclear single quantum coherence) spectra of **6a** (Figs. S13, S14). The splitting of the signals occurs on all methylene groups not belonging to the macrocyclic ring; this disappears upon the removal of *tert*-butyl protective groups in  $\text{L}^1$ .

### Relaxometric $\text{Ca}^{2+}$ titrations of $\text{Gd}_2\text{L}^1$ and $\text{Gd}_2\text{L}^2$

The titration curves are represented as the variation of the relaxivity of the  $\text{Gd}^{3+}$  complex solution versus the  $\text{Ca}^{2+}$  concentration (Fig. 1). The maximum relaxivity increase

**Scheme 1** Synthesis of the ligands and their lanthanide(III) complexes. Reagents and conditions: *a* benzyl chloroformate,  $K_2CO_3$ , water/dioxane; *b* *tert*-butylbromoacetate,  $NaHCO_3$ , MeCN, 70%; *c* **1a** or **1b**,  $K_2CO_3$ , MeCN, 60%; *d*  $H_2$ , Pd/C, MeOH; *e*  $Ac_2O$ , pyridine, 75%; *f* *N*-methylpyrrolidinone,  $Et_3N$ ; *g* trifluoroacetic acid; *h*  $LnCl_3 \cdot 6H_2O$




upon  $Ca^{2+}$  addition was 15% for  $Gd_2L^1$  and 10% for  $Gd_2L^2$ . The saturation of both curves occurs at relatively high  $Ca^{2+}$  concentrations (approximately 20 equiv of  $Ca^{2+}$  for  $Gd_2L^1$  and approximately 5 equiv of  $Ca^{2+}$  for  $Gd_2L^2$ ). The titration curves were fitted to obtain the apparent association constants, which are  $\log K = 1.9 \pm 0.2$  and  $\log K = 2.7 \pm 0.2$  for  $Gd_2L^1$  and  $Gd_2L^2$ , respectively. Similarly to the case for previously investigated BAPTA-type complexes [11, 25], 1:1 binding stoichiometry has been assumed. These association constants are to be compared with the conditional stability constant,  $\log K_{cond} = 6.9$  of  $CaBAPTA^{2-}$  at pH 7.0, calculated by taking into account the protonation constants of  $BAPTA^{4-}$  [25]. The constants obtained for our systems are 3–4 orders of magnitude lower. This difference can be rationalized by the fact that even on  $Ca^{2+}$  binding, the amide groups of the ligand remain coordinated to the lanthanide as indicated by proton relaxivity and luminescence data on the  $Gd^{3+}$  and  $Eu^{3+}$  complexes, respectively (see below). Therefore, in comparison with  $CaBAPTA^{2-}$ , there are two carboxylate donors fewer coordinating to the  $Ca^{2+}$  ion, which is responsible for the considerably reduced stability. It is interesting to note that the stability of the  $Gd_2L^2$ -Ca complex is somewhat higher than that of  $Gd_2L^1$ -Ca. This is likely related to the higher flexibility of the  $Ca^{2+}$  binding

site in  $Gd_2L^2$ -Ca, where the propylene linker between the macrocycle and the  $Ca^{2+}$  binding site induces fewer steric constraints, hence ensuring a better “overwrapping” of the cation by the chelator. We have to note that for  $GdDOPTA$ -Ca, possessing an integral  $BAPTA^{4-}$  unit for  $Ca^{2+}$  binding, Li et al. [11] reported a much higher apparent association constant,  $K = 1.0 \times 10^6$  M.

#### Relaxometric $Mg^{2+}$ titration of $Gd_2L^1$

The relaxivity remains approximately constant upon addition of  $Mg^{2+}$ , then it increases upon  $Ca^{2+}$  addition even in the presence of a large amount of  $Mg^{2+}$  (Fig. 2). The relaxivity increase after  $Ca^{2+}$  addition (16%) was similar to that observed in the  $Ca^{2+}$  titration without the presence of  $Mg^{2+}$ . The insensitivity of  $Gd_2L^1$  towards  $Mg^{2+}$  is due to the lower stability of BAPTA-derived complexes with  $Mg^{2+}$  compared with  $Ca^{2+}$ . The titration curve shows that, though the affinity of our chelator for  $Ca^{2+}$  is diminished in comparison with  $BAPTA^{4-}$ , the selectivity versus  $Mg^{2+}$  is conserved. The discrimination of BAPTA-type ligands towards  $Ca^{2+}$  compared with  $Mg^{2+}$  is supposed to stem from the right size of the binding cavity for the larger-sized  $Ca^{2+}$ , which is already too big for  $Mg^{2+}$  and cannot




**Fig. 1** Relaxometric  $\text{Ca}^{2+}$  titration curves of  $\text{Gd}_2\text{L}^1$  (a) and  $\text{Gd}_2\text{L}^2$  (b) obtained at 298 K and 11.75 T. The lines correspond to the fit as explained in the text

constrict further to envelop snugly this smaller cation (the association constant of  $\text{MgBAPTA}^{2-}$  is  $\log K = 1.8$ ) [25]. Spectrophotometric measurements on the  $\text{Mg}^{2+}$ –BAPTA system proved that  $\text{Mg}^{2+}$  coordination affects only half of the ligand, in contrast to  $\text{Ca}^{2+}$  coordination, where the entire ligand is involved [25].

Luminescence and UV–vis absorption studies to assess the hydration state of  $\text{Eu}_2\text{L}^1$

In order to determine the number of water molecules coordinated to the lanthanide ion before and after  $\text{Ca}^{2+}$  binding to the complex, we performed luminescence lifetime measurements on  $\text{Eu}_2\text{L}^1$  in  $\text{H}_2\text{O}$  and  $\text{D}_2\text{O}$  solutions [36–38]. In the absence of  $\text{Ca}^{2+}$ , the luminescence lifetimes are  $\tau_{\text{H}_2\text{O}} = 0.328$  ms and  $\tau_{\text{D}_2\text{O}} = 0.408$  ms. The hydration number,  $q = 0.4$ , was calculated according to the revised equation of Beeby et al. [38] which involves a correction for the contribution of second-sphere water molecules (Eq. 2):




**Fig. 2** Relaxometric titration curve of  $\text{Gd}_2\text{L}^1$  with  $\text{Mg}^{2+}$  followed by addition of  $\text{Ca}^{2+}$  (298 K, 11.75 T)

$$q = A'(\Delta k_{\text{H}_2\text{O}} - k_{\text{D}_2\text{O}})_{\text{corr}}, \quad (2)$$

where  $A'$  is 1.2 ms and the correction factor for the contribution of the second sphere is  $-0.25 \text{ m s}^{-1}$ . The longitudinal  $^{17}\text{O}$  and  $^1\text{H}$  relaxation rates (vide supra) indicate that we cannot neglect the effect of second-sphere water molecules. The noninteger hydration number suggests the presence of both  $q = 1$  and  $q = 0$  species. Upon the addition of  $\text{Ca}^{2+}$  to the  $\text{Eu}^{3+}$  complex, the hydration number determined by luminescence increases to  $q \approx 0.7$  ( $\tau_{\text{H}_2\text{O}} = 0.422$  ms and  $\tau_{\text{D}_2\text{O}} = 0.661$  ms) (Fig. 3). Such an increase of  $q$  can be interpreted in terms of a shift towards the monohydrated species.


To investigate further the hydration state, variable-temperature UV–vis measurements were performed on  $\text{Eu}_2\text{L}^1$ . In general, the presence of a hydration equilibrium of europium(III) species leads to the appearance of two absorption bands for the  $^5\text{D}_0 \leftarrow ^7\text{F}_0$  transition with peak separations of more than 0.5 nm [39, 40]. High-resolution UV–vis absorption spectra of a  $\text{Eu}_2\text{L}^1$  aqueous solution (pH 7) were recorded in the 577–581-nm region at various temperatures and  $\text{Ca}^{2+}$  concentrations. The measurements revealed two temperature-invariant absorption bands, which could be deconvoluted into two symmetrical peaks (Fig. 4). The separation of those peaks is about 0.5 nm and lies in the range typical of different coordination environments of  $\text{Eu}^{3+}$  [41, 42]. We assume that in the absence of  $\text{Ca}^{2+}$ , there are two nine-coordinate species present; in both species the amines and carboxylates of the macrocycle as well as the amide oxygen are coordinated to the lanthanide, and either carboxylate from the central part or one water molecule completes the coordination sphere to a coordination number of nine, which is the usual coordination number for this type of  $\text{Ln}^{3+}$  complex (Fig. 5). The coordination of amide oxygens is supported by IR measurements performed on the ligand  $\text{L}^1$  and its  $\text{Gd}^{3+}$  complex: the comparison of the two IR spectra (Figs. S5, S6) shows that the amide




**Fig. 3** Dependence of the relaxivity  $r_1$  and of the hydration number  $q$  on the  $\text{Ca}^{2+}$ -to-complex ratio for  $\text{Ln}_2\text{L}^1$ . Squares correspond to the relaxivity values obtained from the relaxometric titration of  $\text{Gd}_2\text{L}^1$  (298 K, 11.75 T). Filled triangles represent the hydration number obtained from luminescence lifetime studies on  $\text{Eu}_2\text{L}^1$  and open triangles correspond to the molar fraction of the monohydrated species determined for  $\text{Eu}_2\text{L}^1$  by UV-vis spectroscopy (ratio of the integrals of the absorption bands corresponding to monohydrated and nonhydrated complex; see text)

absorption is affected by complexation to the lanthanide ion.

Among analogous structures, lanthanide complexes of monomeric DO3A derivatives bearing  $-(\text{CH}_2)_n\text{-NHCO-R}$  ( $n = 2, 3$ ) amide units have been reported in the literature [43]. With  $\text{Eu}^{3+}$  and  $\text{Gd}^{3+}$ , the  $n = 2$  ligand forms monohydrated complexes, while the  $n = 3$  ligand forms nonhydrated complexes. In contrast to this,  $\text{Eu}_2\text{L}^1$  contains a carboxylate in the central part as a potential donor which can partially replace the water molecule in the inner coordination sphere. This leads to the presence of the two



**Fig. 4** Experimental and fitted UV-vis spectra (323 K, pH 7) of  $\text{Eu}_2\text{L}^1$  in the absence and presence of  $\text{Ca}^{2+}$ . On addition of  $\text{Ca}^{2+}$ , the relative intensity of the band at lower wavelength increases, while that of the band at higher wavelength decreases



**Fig. 5** Proposed structures present in aqueous solution of  $\text{Ln}_2\text{L}^1$ . R stands for the remaining part of the bismacrocycle complex

structures as proposed in Fig. 5, and consequently to a reduced hydration number.

Upon  $\text{Ca}^{2+}$  addition to a  $\text{Eu}_2\text{L}^1$  solution, the relative intensity of the two UV-vis absorption bands changes: the band at higher wavelengths decreases, while the other band increases (Fig. 4) [44]. In parallel, the luminescence measurements indicate an increase of the hydration number. Therefore, we conclude that the band which decreases in intensity (at 579.9 nm) on  $\text{Ca}^{2+}$  addition can be attributed to the  $q = 0$  complex, while the band which increases in intensity (579.3 nm) can be attributed to the monohydrated complex. We assume that both in the presence and in the absence of  $\text{Ca}^{2+}$ , the amide oxygen remains coordinated to the lanthanide ion to preserve the overall coordination number of nine. If the amide oxygen participated also in the  $\text{Ca}^{2+}$  coordination, the concomitant increase in the hydration number and in relaxivity would be more prominent than what is experimentally observed. The  $\text{Ca}^{2+}$  binding in the central part of the complex demands the coordination of the central carboxylates, which, by leaving the coordination environment of  $\text{Ln}^{3+}$ , allow a water molecule to bind to the lanthanide ion. Therefore, the  $\text{Ca}^{2+}$  binding in the central part will favor the formation of the monohydrated complex (Fig. 5).

#### Relation between the $\text{Ca}^{2+}$ -dependent hydration number $q$ and relaxivity

Upon stepwise addition of  $\text{Ca}^{2+}$ , we observe a good correlation between the relaxivity increase of the  $\text{Gd}_2\text{L}^1$  complex and the increase in  $q$  determined from the luminescence lifetime measurements on the  $\text{Eu}^{3+}$  analogue (Fig. 3). In addition, we also calculated the ratio of the integrals of the two UV-vis absorption bands (attributed to  $q = 0$  and  $q = 1$ ) of  $\text{Eu}_2\text{L}^1$  at various  $\text{Ca}^{2+}$  concentrations. The  $q$  values obtained in this way overlap with those measured by luminescence. This supports our hypothesis that the complex  $\text{Ln}_2\text{L}^1$  exists in the form of two differently hydrated species, each with an overall coordination number of nine. As was proved by the UV-vis studies, their ratio is temperature-independent but it shifts towards the monohydrated species



with increasing  $\text{Ca}^{2+}$  concentration, resulting in a relaxivity increase.

The relaxivity of the  $\text{Gd}_2\text{L}^2$  complex is lower than that of the  $\text{L}^1$  analogue, which suggests that  $q$  is also lower. In the case of the monomeric DO3A derivatives bearing  $-(\text{CH}_2)_n\text{-NHCO-R}$  amide units, with the propyl-linked ( $n = 3$ ) amide  $q = 0$  was determined for the  $\text{Eu}^{3+}$  and  $\text{Gd}^{3+}$  complexes [43]. The extra steric demand associated with the enlargement of the chelate ring incorporating the amide carbonyl group was found to be sufficient to suppress water coordination. For  $\text{Gd}_2\text{L}^2$ , the relaxivities suggest  $q > 0$ . Moreover, the relaxivity increases on  $\text{Ca}^{2+}$  addition, though to a smaller extent than for  $\text{Gd}_2\text{L}^1$ . Therefore, we assume that two differently hydrated species exist also for  $\text{Ln}_2\text{L}^2$  complexes, with an increased proportion of the nonhydrated species compared with  $\text{Ln}_2\text{L}^1$ . The relaxivity change observed on  $\text{Ca}^{2+}$  addition indicates that the central carboxylate participates in the lanthanide coordination; on  $\text{Ca}^{2+}$  binding this carboxylate is removed from the lanthanide ion and replaced by a water molecule. The small relaxivity change shows that the participation of this carboxylate in the lanthanide coordination is limited with respect to  $\text{Gd}_2\text{L}^1$ .

For  $\text{GdDOPTA}$ , Li et al. [11, 24] reported a more important variation of  $q$  and a correspondingly greater relaxivity change on  $\text{Ca}^{2+}$  binding. This is likely related to the more flexible nature of the central  $\text{BAPTA}^{4-}$  part of the DOPTA ligand in contrast to the BAPTA-bisamide moiety in our case, where the monoamide functionalities are integrated in the central skeleton of the ligand and have much less flexibility to change coordination from  $\text{Gd}^{3+}$  to  $\text{Ca}^{2+}$ .


$^1\text{H}$  and  $^{17}\text{O}$  relaxation studies of  $\text{Gd}_2\text{L}^1$ : evaluation of the parameters influencing proton relaxivity

$^1\text{H}$  NMRD profiles were recorded for  $\text{Gd}_2\text{L}^1$  with and without  $\text{Ca}^{2+}$ . In the presence of  $\text{Ca}^{2+}$ , an increase in the relaxivity was observed at all frequencies, related to an increase in the hydration number. The relaxivity at 20 MHz and 298 K in the absence and presence of  $\text{Ca}^{2+}$  is 5.74 and  $6.13 \text{ mM}^{-1} \text{ s}^{-1}$ , respectively, slightly higher than the relaxivities of currently used MRI contrast agents [2, 4, 19, 45].

The transverse  $^{17}\text{O}$  relaxation rates indicate a relatively slow water exchange (Fig. 6), which is visible from the curve of  $\ln(1/T_{2r})$  versus inverse temperature: at low temperatures,  $1/T_{2r}$  decreases with decreasing temperature. The reduced chemical shifts ( $\Delta\omega_r$ ) in the absence of  $\text{Ca}^{2+}$  are smaller than would be expected for a  $q = 0.4$  complex. Such small chemical shifts have previously been observed in systems with a significant second-sphere contribution

[46]. Therefore, the chemical shifts were not included in the final fitting. For the  $\text{Ca}^{2+}$ -free system, the transverse and longitudinal  $^{17}\text{O}$  relaxation rates and the  $^1\text{H}$  NMRD data were analyzed simultaneously (Fig. 6) on the basis of the Solomon–Bloembergen–Morgan approach, extended by a second-sphere contribution [46, 47]. The presence of free carboxylates in the complex induces a second-sphere contribution that affects both  $^1\text{H}$  and  $^{17}\text{O}$  longitudinal relaxation. In fact, by fitting the  $^{17}\text{O}$   $1/T_1$  values without second-sphere contribution, we obtained inconceivably high rotational correlation times. The inner-sphere hydration number  $q$  was fixed to 0.4, the value found by luminescence measurements on  $\text{Eu}_2\text{L}^1$ . Since  $\text{Eu}^{3+}$  and  $\text{Gd}^{3+}$  have similar ionic radii, we expect a similar hydration mode for  $\text{Gd}_2\text{L}^1$ . To describe the second-sphere contribution, one water molecule per  $\text{Gd}^{3+}$  ( $q^{2\text{nd}} = 1$ ) was considered during the fitting. The distance between the  $\text{Gd}^{3+}$  ion and the second-sphere water proton and oxygen was fixed to  $r_{\text{Gd-H}}^{2\text{nd}} = 3.5 \text{ \AA}$  and  $r_{\text{Gd-O}}^{2\text{nd}} = 4.1 \text{ \AA}$ , respectively, the enthalpy of activation to  $\Delta H^{\#2\text{nd}} = 35 \text{ kJ mol}^{-1}$ , and the second-sphere water residence time to  $\tau_{\text{m}}^{2\text{nd}} = 50 \text{ ps}$  [46, 47]. Other parameters were also fixed during the fitting in order to add some constraints. They are as follows: the hyperfine coupling constant,  $A/\hbar = -3.8 \text{ MHz}$ ; the distance between  $\text{Gd}^{3+}$  and the oxygen and the proton of the first-sphere water molecule,  $r_{\text{GdO}} = 2.5 \text{ \AA}$  and  $r_{\text{GdH}} = 3.1 \text{ \AA}$ , respectively. The distance of the closest approach of outer-sphere water molecules to  $\text{Gd}^{3+}$ ,  $a$ , was fixed to  $3.6 \text{ \AA}$ . The quadrupolar coupling constant [ $\chi(1 + \eta^2/3)^{1/2}$ ] was set to  $7.58 \text{ MHz}$ , the value of pure water. The activation energy  $E_v$  had to be fixed to  $1 \text{ kJ mol}^{-1}$ , otherwise the fit converged to negative values. In the fitting procedure, we used a model which considers different rotational correlation times for  $\text{Gd-O}$  and  $\text{Gd-H}$  rotating vectors ( $\tau_{\text{RO}}^{298}$  and  $\tau_{\text{RH}}^{298}$ , respectively) [48]. For geometrical reasons, their ratio ( $\tau_{\text{RH}}^{298}/\tau_{\text{RO}}^{298}$ ) has to lie between 0.65 and 1. The most important parameters obtained from the best simultaneous least-squares fit to the experimental data are listed in Table 1. For the complete fitting results, see Table S9.


The water exchange of  $\text{Gd}_2\text{L}^1$  is slightly slower than that found for gadolinium 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetate ( $\text{GdDOTA}$ ) [49]. It is interesting to note that for the monohydrated  $\text{Gd}^{3+}$  complex of the DO3A ligand bearing an N-linked  $\text{CH}_2\text{CH}_2\text{NHCO}$ -pyridyl pendant arm, much faster water exchange has been reported ( $k_{\text{ex}}^{298} = 1.1 \times 10^8 \text{ s}^{-1}$ ) [43]. This fast water exchange has been explained by the steric destabilization of the  $\text{Ln}$ -water binding interaction by the presence of the bulky substituent. Such a destabilization effect is expected to be less important in  $\text{Gd}_2\text{L}^1$  since the amide carbonyl is linked to a flexible  $-\text{CH}_2\text{-N}-$  moiety in contrast to the direct attachment to a pyridyl group in the previous case. This limited steric constraint around the water binding site will then be



**Fig. 6** Variable-temperature  $^{17}\text{O}$  NMR [a;  $\ln(1/T_{1r})$  diamonds,  $\ln(1/T_{2r})$  squares] and  $^1\text{H}$  nuclear magnetic relaxation dispersion (b; 298 K squares, 310 K circles) data of  $\text{Gd}_2\text{L}^1$  in the absence of  $\text{Ca}^{2+}$ . The curves correspond to the simultaneous fit as explained in the text

translated by a more sluggish water exchange as observed for  $\text{Gd}_2\text{L}^1$ . The more than 1 order of magnitude difference in the water exchange rate between  $\text{Gd}_2\text{L}^1$  and the  $\text{Gd}^{3+}$  complex of the DO3A- $\text{CH}_2\text{CH}_2\text{NHCO}$ -pyridyl ligand is a good example of the importance of the steric compression around the water binding site. Steric compression is indeed the main factor determining the rate of exchange in a dissociative water exchange mechanism, characteristic of nine-coordinate complexes [50]. The dissociative activation mode for  $\text{Gd}_2\text{L}^1$  is indicated by the positive value of the activation entropy.

The higher  $\tau_r$  value of  $\text{Gd}_2\text{L}^1$  compared with that of GdDOTA is rationalized by the higher molecular weight and greater rigidity of the molecule. In spite of the slow rotational motion and the presence of a second hydration sphere in the bismacrocyclic system, the relaxivities are only slightly higher than those of GdDOTA, owing to the low hydration number of  $\text{Gd}_2\text{L}^1$  ( $q = 0.4$ ).



**Fig. 7** Variable-temperature, reduced longitudinal (squares) and transverse (circles)  $^{17}\text{O}$  relaxation rates for  $\text{Gd}_2\text{L}^1$ , in the presence of 1 M  $\text{Ca}^{2+}$ . The curves correspond to the fit as explained in the text

**Table 1** Kinetic and structural parameters obtained from the fit of  $^{17}\text{O}$  NMR and NMR dispersion (NMRD) data for the  $\text{Gd}_2\text{L}^1$  complex in the absence and in the presence of  $\text{Ca}^{2+}$ , compared with those of gadolinium 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (GdDOTA)

Parameters	$\text{Gd}_2\text{L}^1$ $^{17}\text{O}$ NMR + NMRD	$\text{Gd}_2\text{L}^1 + \text{Ca}^{2+}$ $^{17}\text{O}$ NMR	GdDOTA $^{17}\text{O}$ NMR + NMRD <sup>a</sup>
$k_{\text{ex}}^{298}$ ( $10^6 \text{ s}^{-1}$ )	$2.4 \pm 0.2$	$7.5 \pm 1.6$	4.1
$\Delta H^\ddagger$ ( $\text{kJ mol}^{-1}$ )	$43.6 \pm 3.3$	43.6	49.8
$\Delta S^\ddagger$ ( $\text{J mol}^{-1} \text{ K}^{-1}$ )	+23.5	+33.0	–
$\tau_{\text{RO}}^{298}$ (ps)	$350 \pm 50$	$1,150 \pm 250$	77
$E_r$ ( $\text{kJ mol}^{-1}$ )	$24 \pm 1$	$21 \pm 6$	16.1
$\tau_v^{298}$ (ps)	$20.6 \pm 2.7$	$0.13 \pm 0.02$	11
$\Delta^2$ ( $10^{20} \text{ s}^{-2}$ )	$0.46 \pm 0.10$	$0.50 \pm 0.05$	0.16
$q$	0.4	0.7	1
$q^{2 \text{ ns}}$	1	1	–
$r_1^{298}$ ( $\text{mM}^{-1} \text{ s}^{-1}$ ) 20 MHz	5.74	6.13	4.74

Parameters in *italics* were fixed during the fitting

<sup>a</sup> Reference [49]

The simultaneous fit of the  $^1\text{H}$  and  $^{17}\text{O}$  relaxation rates also supplies parameters that describe the electron spin relaxation of the  $\text{Gd}^{3+}$  complexes, such as  $\tau_v$ , the correlation time for the modulation of the zero field splitting, its activation energy,  $E_v$ , and the mean zero field splitting energy,  $\Delta^2$ . The values obtained for  $\text{Gd}_2\text{L}^1$  are in the usual range for similar complexes [49].

The  $^1\text{H}$  NMRD and  $^{17}\text{O}$  NMR data of the  $\text{Gd}_2\text{L}^1$  system containing  $\text{Ca}^{2+}$  could not be fitted simultaneously because the conditions for  $^{17}\text{O}$  NMR and  $^1\text{H}$  NMRD samples differed significantly in terms of the ionic strength and viscosity, and also in the  $\text{Ca}^{2+}$ -to- $\text{Gd}^{3+}$  ratio of the samples. Only the  $^{17}\text{O}$   $\ln(1/T_{1r})$  and  $\ln(1/T_{2r})$  data were analyzed,

and the relaxivities are given in the electronic supplementary material. The  $^{17}\text{O}$  relaxation rates were interpreted by using the same set of equations as for the system without  $\text{Ca}^{2+}$  (Fig. 7). Analogously, the same parameters were fixed in the fit (except for the hydration number,  $q = 0.7$ , obtained from luminescence studies). The best fitting was obtained with the parameters listed in Tables 1 and S10. On  $\text{Ca}^{2+}$  addition, the most flagrant change is observed for the rotational correlation time as obtained from the  $^{17}\text{O}$  longitudinal relaxation rates ( $\tau_{\text{R}} = 350$  compared with 1,150 ps without and with  $\text{Ca}^{2+}$ , respectively). We attribute this significant increase mainly to the high ionic strength and viscosity of the  $^{17}\text{O}$  NMR sample containing 1 M  $\text{CaCl}_2$ , though some rigidification of the molecule on  $\text{Ca}^{2+}$  is also expected. The water exchange rate triples on  $\text{Ca}^{2+}$  binding ( $k_{\text{ex}}^{298} = 2.4 \times 10^6$  compared with  $7.5 \times 10^6 \text{ s}^{-1}$ ). The decreasing negative charge of the  $\text{Ca}^{2+}$ -bound  $\text{Gd}_2\text{L}^1$ -Ca complex would be rather expected to diminish the exchange rate in a dissociatively activated process. Among other factors that can compensate this effect, the increased steric demand of the central part after  $\text{Ca}^{2+}$  binding can lead to a steric destabilization of the Ln-water binding interaction. Overall, the relaxivity increase upon  $\text{Ca}^{2+}$  addition is mainly related to an increase in the hydration number.

## Conclusion

With the objective of  $\text{Ca}^{2+}$  sensing by MRI in the extracellular space, we synthesized two novel bismacrocylic ligands containing a  $\text{Ca}^{2+}$ -sensitive BAPTA-bisamide moiety (Scheme 1). Their  $\text{Gd}^{3+}$  complexes exhibit relaxivities comparable to that of currently used monomeric MRI contrast agents, in accordance with their larger size but lower hydration numbers ( $q < 1$ ). Upon  $\text{Ca}^{2+}$  addition, the relaxivities of  $\text{Gd}_2\text{L}^1$  and  $\text{Gd}_2\text{L}^2$  increase by 15 and 10%, respectively. These changes are likely insufficient for in vivo MRI detection.  $\text{Gd}_2\text{L}^1$  is practically insensitive towards  $\text{Mg}^{2+}$ . The apparent association constants for the  $\text{Ca}^{2+}$  interaction were obtained from the relaxometric titration curves. They are several orders of magnitude lower for our complexes than those determined for BAPTA $^{4-}$  itself or other tetracarboxylate BAPTA $^{4-}$  analogues. The hydration number of  $\text{Eu}_2\text{L}^1$  was determined from luminescence lifetime measurements in the absence ( $q = 0.4$ ) and presence ( $q = 0.7$ ) of  $\text{Ca}^{2+}$ . UV-vis measurements confirmed the presence of two coordination environments that we attribute to a monohydrated and a nonhydrated state (Fig. 5). Upon the coordination of  $\text{Ca}^{2+}$  in the central part of the ligand, the molecule undergoes a conformational change and an acetate arm, originally coordinated to the  $\text{Ln}^{3+}$  ion, binds to  $\text{Ca}^{2+}$  and leaves space

for the coordination of a water molecule to the lanthanide, which shifts the hydration state towards the more hydrated species.

**Acknowledgements** We are grateful to Lothar Helm, Ecole Polytechnique Fédérale de Lausanne, Switzerland, for allowing us to use their relaxometric facilities. The authors would like to thank A.J. Meixner (University of Tübingen) for providing access to his fluorescence spectrometer and I. Mamedov (MPI for Biological Cybernetics, Tübingen) for his help with luminescence lifetime experiments. This work was carried out in the framework of the EC COST D38 Action and was supported by the Ministère de l'Éducation Nationale et de la Recherche (France), the Centre National pour la Recherche Scientifique, the Hertie foundation and the Max Planck Society.

## References

- Persigehl T, Heindel W, Bremer C (2005) *Abdom Imaging* 30:342–354
- Merbach AE, Tóth É (eds) (2001) *The chemistry of contrast agents in medical MRI*. Wiley, Chichester
- Aime S, Botta M, Terreno E (2005) *Adv Inorg Chem* 57:173–237
- Caravan P, Ellison JJ, McCurry TJ, Lauffer RB (1999) *Chem Rev* 99:2293–2352
- Lowe MP, Parker D, Reany O, Aime S, Botta M, Castellano G et al (2001) *J Am Chem Soc* 123:7601–7609
- Aime S, Barge A, Delli Castelli D, Fedeli F, Mortillaro A, Nielsen FU et al (2002) *Magn Reson Med* 47:639–648
- Zhou J, Lal B, Wilson DA, Larterra J, van Zijl PC (2003) *Magn Reson Med* 50:1120–1126
- Aime S, Botta M, Gianolio E, Terreno E (2000) *Angew Chem Int Ed Engl* 39:747–750
- Sun PZ, Schoening ZB, Jasanoff A (2003) *Magn Reson Med* 49:609–614
- Que EL, Chang CJ (2006) *J Am Chem Soc* 128:5942–5943
- Li W, Fraser SE, Meade TJ (1999) *J Am Chem Soc* 121:1413–1414
- Hanaoka K, Kikuchi K, Urano Y, Narazaki M, Yokawa T, Sakamoto S et al (2002) *Chem Biol* 9:1027–1032
- Aime S, Delli Castelli D, Fedeli F, Terreno E (2002) *J Am Chem Soc* 124:9364–9365
- Hanaoka K, Kikuchi K, Urano Y, Nagano T (2001) *J Chem Soc Perkin Trans 2* 1840–1843
- Louie AY, Huber MM, Ahrens ET, Rothbacher U, Moats R, Jacobs RE et al (2000) *Nat Biotechnol* 18:321–325
- Perez JM, Josephson L, O'Loughlin T, Hogemann D, Weissleder R (2002) *Nat Biotechnol* 20:816–820
- Woods M, Zhang S, Ebron VH, Sherry AD (2003) *Chem Eur J* 9:4634–4640
- Zech SG, Eldredge HB, Lowe MP, Caravan P (2007) *Inorg Chem* 46:3576–3584
- Krause W (ed) (2002) *Contrast agents I. Magnetic resonance imaging. Topics in current chemistry*, vol 221. Springer, Heidelberg
- Atanasijevic T, Shusteff M, Fam P, Jasanoff A (2006) *Proc Natl Acad Sci USA* 103:14707–14712
- Meldolesi J (1998) *Nature* 392:863–866
- Burgoyne RD (2007) *Nat Rev Neurosci* 8:182–193
- Tsien RY, Waggoner A (1995) In: Pawley JB (ed) *Handbook of biological confocal microscopy*, 2nd edn. Plenum, New York, pp 267–280
- Li W, Parigi G, Fragai M, Luchinat C, Meade TJ (2002) *Inorg Chem* 41:4018–4024

25. Tsien RY (1980) *Biochemistry* 19:2396–2404
26. Graeppi N, Powell DH, Laurenzcy G, Zékány L, Merbach AE (1995) *Inorg Chim Acta* 235:311–326
27. Amman C, Meyer P, Merbach AE (1982) *J Magn Reson* 46:319–321
28. Raiford DS, Fisk CL, Becker ED (1979) *Anal Chem* 51:2050–2051
29. Zitha-Bovens E, Vander Elst L, Muller RN, van Bekkum H, Peters JA (2001) *Eur J Inorg Chem* 12:3101–3105
30. Yerly F (1999) Visualiseur 2.3.4. Institut de Chimie Moléculaire et Biologique, EPFL-BCH, Lausanne
31. Yerly F (1999) Optimiseur 2.3.4. Institut de Chimie Moléculaire et Biologique, EPFL-BCH, Lausanne
32. Wong W, Li C (2005) *Int Patent Appl WO* 2005003105 A1 20050113
33. Wei W, Tomohiro T, Kodaka M, Okuno H (2000) *J Org Chem* 65:8979–8987
34. Adam SR, Kao JPY, Gryniewicz G, Minta A, Tsien RY (1988) *J Am Chem Soc* 110:3212
35. Corsi DM, Platas C, Van Bekkum H, Peters JA (2001) *Magn Res Chem* 39:723
36. Horrocks WD Jr, Sudnick DR (1979) *J Am Chem Soc* 101:334–340
37. Supkowski RM, Horrocks WD Jr (2002) *Inorg Chim Acta* 340:44–48
38. Beeby A, Clarkson IM, Dickins RS, Faulkner S, Parker D, Royle L, de Sousa AS, Williams JAG, Woods M (1999) *J Chem Soc Perkin Trans 2* 493–503
39. Albin M, Farber GK, Horrocks WD Jr (1984) *Inorg Chem* 23:1648–1651
40. Geier G, Jorgensen CK (1971) *Chem Phys Lett* 9:263–264
41. Tóth É, Ni Dhubghaill OM, Besson G, Helm L, Merbach AE (1999) *Magn Res Chem* 37:701–708
42. Yerly F, Dunand FA, Tóth É, Figueirinha A, Kovács Z, Sherry AD, Geraldés CFGC, Merbach AE (2000) *Eur J Inorg Chem* 1001–1006
43. Congreve A, Parker D, Gianolio E, Botta M (2004) *Dalton Trans* 1441–1445
44. Frey ST, Horrocks WD Jr (1995) *Inorg Chim Acta* 229:383–390
45. Caravan P (2006) *Chem Soc Rev* 35:512–523
46. Lebdušková P, Hermann P, Helm L, Tóth É, Kotek J, Binnemans K, Rudovský J, Lukeš I, Merbach AE (2007) *Dalton Trans* 493–501
47. Lebdušková P, Sour A, Helm L, Tóth É, Kotek J, Lukeš I, Merbach AE (2006) *Dalton Trans* 3399–3406
48. Dunand FA, Borel A, Merbach AE (2002) *J Am Chem Soc* 124:710–716
49. Powell DH, Dhubghaill OMN, Pubanz D, Helm L, Lebedev YS, Schlaepfer W, Merbach AE (1996) *J Am Chem Soc* 118:9333–9346
50. Laus S, Ruloff R, Tóth É, Merbach AE (2003) *Chem Eur J* 9:3555