

Toxic Effect of Herbicide 2,4-D on the Earthworm *Eutyphoeus waltoni* Michaelsen

Vandana Singh · Keshav Singh

Received: 13 August 2014 / Accepted: 20 October 2014 / Published online: 22 January 2015
© Springer International Publishing Switzerland 2015

Abstract The toxicity of herbicide 2,4-D (2'-4' dichloro phenoxy acetic acid) was studied on earthworms in different combinations of biological wastes and different types of soils as feed materials under laboratory conditions. *Eutyphoeus waltoni* were exposed to different concentrations of 2,4-D (200, 300, 400, 450 mg/kg) in feed material (i.e., buffalo dung, wheat straw and gram bran) and different concentrations of 2,4-D (150, 300, 350, 400 mg/kg) in different types of soil (i.e., loamy soil, clay soil and sandy soil). Observations were recorded from 24 h up to 240 h. It was observed that the toxic effect of herbicide 2,4-D on the earthworm *Eutyphoeus waltoni* was both time and dose dependent. Maximum toxicity was observed in the sandy soil, whereas minimum in the feed material of buffalo dung with gram bran. There was no mortality observed in control.

Keywords Herbicide · 2,4-D · *Eutyphoeus waltoni* · Soils · Buffalo dung · Agro-wastes · Toxicity

1 Introduction

Soil environments are contaminated by the indiscriminate use of pesticides and herbicides, which affect the soil flora and fauna population (Gobi and Gunasekaran 2010). Earthworms were used as model experimental organisms for toxicity as well as bioaccumulation assessment (Nuseti et al. 1999). 2,4 dichlorophenoxy acetic acid (2,4-D) is a low cost, easily available, and early and extensively effective used herbicide worldwide. Ville et al. (1997) reported that the 2,4-D has toxic effects on mammals, including neurological dysfunction, pulmonary oedema, hepatic and renal dysfunctions or symptoms of tetanus. Generally 2,4-D is used in agricultural fields for the purpose of controlling broad-leaf weeds (Munro et al. 1992). The present work studies the toxicity of 2,4-D on earthworm *Eutyphoeus waltoni* in soil, because this herbicide has poor biodegradability, several metabolic alterations and tissue necrosis in non-target organisms, including important members of the food chain organisms, such as fish (Gallagher and Di Giulio 1991; Chingombe et al. 2006).

The abundant use of herbicides does not affect only selected weeds; herbicides can also destroy non-target species in the agricultural fields and also affect the texture and physico-

V. Singh · K. Singh (✉)
Vermiculture Research Laboratory, Department of Zoology, D.D.U. Gorakhpur University,
Gorakhpur 273 009 UP, India
e-mail: keshav26singh@rediffmail.com

chemical properties of soils (Robidoux et al. 1999). The use of specific herbicides, fungicides, insecticides in the agricultural field can be highly toxic to earthworm population (Williamson 2000; Zhou et al. 2007). The exclusive major role of earthworms in paedogenesis is through mixing of the particles during digesting, depositing their casts throughout the soil column, and improving aeration and drainage of the agricultural soils (Kavitha et al. 2011). Earthworms are also important contributors to the recycling of carbon and nitrogen in the ecosystem, so, they are used as bioindicators (Callahan 1988; Goats and Edwards 1988).

Correia and Moreira (2010) reported that the effects on growth, survival and reproduction rates of earthworms were verified for different exposure times. Earthworms kept in glyphosate treated soil remained alive in all evaluations, but showed gradual and significant reduction in mean weight (50 %) at all test concentrations. The combinations of animal dung with different agro-wastes are the best suitable feed material for better growth and development of earthworm *Eisenia fetida*. The combination of buffalo dung with wheat straw and gram bran resulted in maximum biomass, weight and length (Nath et al. 2009; Chauhan and Singh 2012; Kumar and Singh 2013). Singh and Kumar (2014) reported that the earthworm *Eutyphoeus waltoni* is found abundantly in agricultural fields of different localities of eastern Uttar Pradesh. Correia and Moreira (2010) reported that 100 % mortality of epigeic earthworm was observed in soil treated with 500 and 1000 mg/kg of 2,4-D; at 14 days, 30–40 % mortality levels were observed for all treated concentrations. No cocoons or juveniles were found in soil treated with the herbicide. Glyphosate and 2,4-D demonstrated severe effects on the development and reproduction of *Eisenia foetida* in laboratory tests in the range of test concentrations (Correia and Moreira 2010).

The aim of the present study was to investigate the toxic effect of herbicide 2,4-D on the earthworm *Eutyphoeus waltoni* in different soils, and in combination of feed material of buffalo dung with agro-wastes and soil alone as feed materials in laboratory conditions.

2 Materials and Methods

Collection of the Earthworm: The cultured earthworm *Eutyphoeus waltoni* was used in the experiment.

Collection of animal dung and agro-wastes: The buffalo dung, wheat straw and gram bran were collected from different part of Gorakhpur district of U.P. India.

Herbicide: The commercially available herbicide 2,4-D (2'-4' dicholoro phynoxy acetic acid), purchased from EarthCare Pvt. Ltd India, was used in the experiment at different concentrations. The composition of 2,4-D is 2,4-D Acid Technical 59 % w/w (based on 97 % w/w ai.a), dimethyl amine 30 % w/w, Sequesting agent (lignin sulphonate) 1 % w/w, Divent acqua sufficient 10 % w/w.

Table 1 Concentrations used for toxicity determination against earthworm (*Eutyphoeus waltoni*)

Name	Combinations	Concentration (mg/kg)
2,4-D	BD+Ws	200, 300, 400, 450
	BD+Gb	200, 300, 400, 450
	BD+Ws+Gb	200, 300, 400, 450
	Loamy soil	150, 250, 350, 400
	Clay soil	150, 250, 350, 400
	Sandy soil	150, 250, 350, 400

Table 2 Toxicity of herbicide 2,4-D against earthworm *Eutyphoeus waltoni* in different feed materials

Period	Combinations	LC ₅₀	Lower limit	Upper limit	Slope value	t-ratio	g-value	Heterogeneity
24 h	BD+Ws	540.20	477.06	696.39	4.06	4.19	0.218	0.28
	BD+Gb	540.44	475.89	673.04	3.44	6.30	0.097	0.14
	BD+Ws+Gb	510.81	458.33	609.59	3.77	6.71	0.085	0.09
	Loamy soil	470.38	416.14	572.38	3.43	6.95	0.079	0.08
	Clay soil	490.92	427.67	617.82	3.18	6.61	0.088	0.10
	Sandy soil	445.56	392.93	543.98	2.99	6.79	0.083	0.09
48 h	BD+Ws	456.72	416.55	523.62	3.67	7.19	0.074	0.11
	BD+Gb	460.73	414.85	543.85	3.19	6.51	0.090	0.08
	BD+Ws+Gb	474.57	425.11	567.12	3.17	6.37	0.095	0.07
	Loamy soil	392.55	349.76	466.22	2.74	6.79	0.083	0.12
	Clay soil	399.48	356.15	474.41	2.82	6.87	0.081	0.08
	Sandy soil	380.35	338.24	452.59	2.59	6.58	0.088	0.18
72 h	BD+Ws	399.54	368.19	445.09	3.48	7.33	0.071	0.12
	BD+Gb	404.11	364.83	468.91	2.74	6.02	0.106	0.12
	BD+Ws+Gb	409.61	370.43	474.44	2.83	6.15	0.102	0.12
	Loamy soil	310.52	278.56	354.04	2.45	6.58	0.088	0.15
	Clay soil	300.84	271.86	337.55	2.64	7.05	0.077	0.12
	Sandy soil	313.15	280.08	359.27	2.38	6.41	0.093	0.37
96 h	BD+Ws	342.65	312.78	378.72	2.94	6.60	0.088	0.07
	BD+Gb	353.42	314.08	408.84	2.24	5.13	0.145	0.05
	BD+Ws+Gb	343.99	311.58	384.30	2.70	6.12	0.102	0.10
	Loamy soil	250.07	219.84	280.22	2.34	6.46	0.092	0.24
	Clay soil	244.28	216.61	271.06	2.56	7.03	0.078	0.26
	Sandy soil	251.15	222.52	279.81	2.47	6.77	0.084	0.42
120 h	BD+Ws	289.06	261.97	313.90	3.26	7.34	0.071	0.04
	BD+Gb	304.16	265.79	341.67	2.30	5.31	0.136	0.06
	BD+Ws+Gb	299.74	267.23	330.74	2.70	6.17	0.101	0.11
	Loamy soil	212.82	182.97	238.13	2.48	6.82	0.083	0.28
	Clay soil	201.05	171.64	225.25	2.56	7.01	0.078	0.38
	Sandy soil	210.34	179.75	235.89	2.45	6.73	0.085	0.21
240 h	BD+Ws	252.63	225.33	275.08	3.51	7.79	0.063	0.04
	BD+Gb	271.54	230.52	304.43	2.34	5.40	0.131	0.10
	BD+Ws+Gb	268.97	233.40	297.99	2.66	6.09	0.103	0.07
	Loamy soil	190.72	160.07	215.17	2.53	6.89	0.081	0.28
	Clay soil	182.97	156.61	204.38	2.92	7.76	0.064	0.78
	Sandy soil	176.60	160.01	191.04	4.58	10.57	0.034	0.47

Mortality was determined every 24 h. Each set of experiment was replicated six times. Product momentum correlation showed that there was significant negative coefficient ($p < 0.05$) observed between exposure time and different value of LC₅₀ of 2,4-D

Determination of LC₅₀: The toxicity experiment was performed following the method of Agarwal and Singh (1988). Ten adult earthworms were kept in vermibeds of one kg feed material. The vermibeds were exposed to different concentration of herbicides (Table 1). Six




Fig. 1 The toxicity of 24 h exposure of herbicide 2,4-D against earthworm *Eutyphoeus waltoni*

vermibeds were set up for each dose of herbicides. Vermibeds without any treatment were used as control. Mortality was recorded at different exposure periods. Lethal concentration (LC_{50}) value, its upper and lower confidence limits (UCL and LCL) and slope value were calculated according to the method of POLO computer programmers of Russel et al. (1977).

Statistical Analysis: All the studies were replicated at least six times. Product momentum correlation coefficient was determined between exposure time and different values of LC. Analysis of variance was used to analyze the significant difference between different combinations and exposure time. (Sokal and Rohlf 1973).

3 Results

Laboratory toxicity was evaluated for different concentrations of 2,4-D against earthworm *Eutyphoeus waltoni* in different combination of buffalo dung with agro-wastes




Fig. 2 The toxicity of 48 h exposure of herbicide 2,4-D against earthworm *Eutyphoeus waltoni*

72 exposure period

Eutyphoeus waltoni




Fig. 3 The toxicity of 72 h exposure of herbicide 2,4-D against earthworm *Eutyphoeus waltoni*

and soil as feed materials. The toxicity of 2,4-D was both time and dose dependent against earthworm *Eutyphoeus waltoni* in all the exposures. There was a negative significant product momentum coefficient ($P < 0.05$) observed between exposure time and different value of LC_{50} of 2,4-D (Tables 2 and 4; Figs. 1, 2, 3, 4, 5 and 6). The toxicity of 2,4-D against *Eutyphoeus waltoni* was higher in sandy soil in exposure periods of 24 h to 240 h, whereas, minimum in the feed material of combination of buffalo dung with gram bran (Tables 2 and 4; Fig. 1).

Toxicity of 24 h exposure of 2,4-D against the earthworm *Eutyphoeus waltoni* was higher in sandy soil than other combinations of 2,4-D at different exposure periods against *Eutyphoeus waltoni* as 445.56 mg/kg LC_{50} . The order of 24 h

96h exposure period

Eutyphoeus waltoni




Fig. 4 The toxicity of 96 h exposure of herbicide 2,4-D against earthworm *Eutyphoeus waltoni*




Fig. 5 The toxicity of 120 h exposure of herbicide 2,4-D against earthworm *Eutyphoeus waltoni*

exposure toxicity of 2,4-D was sandy soil > loamy soil > clay soil > BD+Ws+Gb > BD+Ws > BD+Gb. The slope value given in Tables 2 and 3 was steep and separate estimation of LC of different combinations was found to be within the 95 % confidence limits. The t-ratio was greater than 1.96 and heterogeneity less than 1. The g-value was less than 0.5 at all probability levels. A significant correlation coefficient (r) between exposure time and LC₅₀ value of 2,4-D was obtained (Tables 2 and 4). There was no mortality recorded after 240 h exposure period in all the treatments (Tables 2 and 4; Fig. 7). Analysis of variance (ANOVA) was used to analyse the significant difference between different combinations of feed materials and exposure period (Table 3).




Fig. 6 The toxicity of 240 h exposure of herbicide 2,4-D against earthworm *Eutyphoeus waltoni*

Table 3 Summary of computation of analysis of variance (ANOVA) of the data of Tables 2 and 4

Source of variation	Toxicity of different combination of wastes				
	DF	SS	σ^2	F	P
Between treatment	5	62909.1	12581.8	33.6	$P < 0.001$
Between Time	5	337386.3	67477.3	180.0	$P < 0.001$
Error	10	3748.7	374.9		
Total	20	404044.1			

4 Discussion

It is evident from the results that the studied toxicity of 2,4-D was time and dose dependent against earthworm *Eutyphoeus waltoni*. Thus, the earthworms have been used as model animals for studying the effects of agrochemicals on soil fauna (Cock et al. 1980; Gobi et al. 2004). Herbicides have been reported to have adverse effect on the survival of earthworms (van Gestrel and van Dis 1988; Robidoux et al. 1999). The herbicide acetochlor caused adverse effect on the sperm number and DNA of *Eisenia fetida* (Xiao et al. 2006). Several other studies have demonstrated the lethal activity of herbicides and pesticides on earthworms and histopathological effects (Gupta and Sundaraman 1988; Sorour and Larink 2001; Lydy and Linck 2003; Gobi et al. 2004; Rombke et al. 2007; Mosieh 2009). Correia and Moreira (2010) reported that 100 % mortality was observed a few hours after exposure of those organisms when exposed to 1000 mg/kg of 2,4-D. 2,4-D is toxic against earthworms according to Roberts and Dorough (1984). Brown (1978) also reported that some herbicides are directly toxic to earthworms.

Herbicides affect the feeding behaviour of earthworms, which was reflected in the weight loss and reproduction capacity (Venter et al. 1988; Bustos–Obregon and Goicochea 2002). Indiscriminate use of pesticides may affect non-target organisms in the soil and can cause serious damage to ecosystem (Reinecke and Reinecke 2007). Smith et al. (1992) reported that soil animals, especially earthworms, are one of the best bioindicators of pesticide contamination. The agrochemical concentration is higher in surface layers; earthworm activity is very much reduced in the soil surface layer (Keogh and Whitehead 1975; Cock et al. 1980).

The toxicity of 2,4-D against earthworm *Eutyphoeus waltoni* was higher in the sandy soil at 24 h up to 240 h exposure periods, because sandy soil have less organic content and other nutritional components than other combinations of agro-wastes and soil. Nath et al. (2009) reported that the feed material of buffalo dung with gram bran have rich organic nutrients. The combination of buffalo dung with gram have minimum toxicity of 2,4-D against *Eutyphoeus waltoni* because it is possible that this combination have rich amount of organic nutrients which tolerate the toxicity of 2,4-D. There was no mortality recorded after 240 h exposure period in all treatments, which may be due to the development of tolerance power against 2,4-D in earthworms.

It is evident from the results, that the herbicide 2,4-D has toxic effects against earthworm population, but the use of vermicompost of different combinations of buffalo dung with agro-wastes in soil protect the earthworms against toxic effects of herbicides. The direct use of different combinations of buffalo dung with agro-wastes

Table 4 Lethal concentrations of herbicide 2,4-D against earthworm *Eutyphoeus waltoni* in different feed materials

Period	Combinations/Soils	LC ₁₀	LC ₅₀	LC ₉₀
24 h	BD+Ws	261.30	540.28	1116.75
	BD+Gb	229.70	540.44	1271.51
	BD+Ws+Gb	233.75	510.81	1116.36
	Loamy soil	199.35	470.38	1109.90
	Clay soil	194.40	490.92	1239.70
	Sandy soil	166.33	445.56	1193.49
48 h	BD+Ws	204.71	456.72	1018.96
	BD+Gb	183.00	460.73	1159.96
	BD+Ws+Gb	187.58	474.57	1200.64
	Loamy soil	133.90	392.55	1150.81
	Clay soil	140.52	399.48	1135.67
	Sandy soil	121.77	380.35	1188.06
72 h	BD+Ws	171.51	399.54	930.74
	BD+Gb	137.88	404.11	1184.40
	BD+Ws+Gb	144.59	409.61	1160.38
	Loamy soil	93.39	310.52	1032.44
	Clay soil	98.61	300.84	917.82
	Sandy soil	91.03	313.15	1077.23
96 h	BD+Ws	125.91	342.65	932.45
	BD+Gb	94.85	353.42	1316.38
	BD+Ws+Gb	115.61	343.99	1023.49
	Loamy soil	71.03	250.06	880.38
	Clay soil	77.44	244.28	778.58
	Sandy soil	76.27	251.15	827.01
120 h	BD+Ws	117.12	289.06	713.44
	BD+Gb	84.32	304.16	1097.17
	BD+Ws+Gb	100.49	299.74	893.99
	Loamy soil	64.83	212.82	698.66
	Clay soil	63.74	201.05	634.18
	Sandy soil	63.18	218.34	700.22
240 h	BD+Ws	109.02	252.63	585.42
	BD+Gb	76.98	271.54	957.81
	BD+Ws+Gb	88.77	268.97	814.96
	Loamy soil	59.43	190.72	612.04
	Clay soil	66.74	182.97	501.65
	Sandy soil	92.77	176.60	336.18

In all cases t-ratio is greater than 1.96, heterogeneity factor is less than 1.0 and g-values were less than 0.5 at all probability levels

in the agricultural fields are suitable feed materials for earthworms which provide better nourishment to tolerate the toxic effect of the herbicides. Production and use of vermicompost from biological wastes promotes plant growth and productivity, as well as protects the earthworms, and helps the growth and development of its population.




Fig. 7 Effect of different combinations of feed material of exposure of herbicide 2,4-D on earthworm *Eutyphoeus waltoni* in different exposure period

References

- Agarwal RA, Singh DK (1988) Harmful gastropods and their control. *Acta Hydrochim Hydrobiol* 16:113–138
- Brown AWA (1978) *Ecology of pesticides*. John Wiley and Sons, New York
- Bustos–Obregon E, Goicochea RI (2002) Pesticides soil contamination mainly affects earthworm male reproductive parameters. *Asian J Androl* 4:195–199
- Callahan CA (1988) Earthworms as ecotoxicological assessment tools. In: Edwards CA, Neuhauser EF (eds) *Earthworms in waste and environmental assessment*. SPB Academic Publishing, The Hague, pp 295–301
- Chauhan HK, Singh K (2012) Effect of binary combinations of buffalo, cow and goat dung with different agro wastes on reproduction and development of earthworm *Eisenia foetida*. *World J Zool* 7(1):23–29
- Chingombe P, Saha B, Wakeman RJ (2006) Effect of surface modification of activated carbon on the sorption of 2,4-dichlorophenoxy acetic acid and benazolin from water. *J Colloid Interface Sci* 297:434–442
- Cock AG, Critchley BRV, Perfect JJ, Yeadon E (1980) Effect of cultivation and DDT on earthworm activity in a forest soil in the subhumid tropics. *J Appl Ecol* 17:21–29
- Correia FV, Moreira JC (2010) Effects of glyphosate and 2,4-D on earthworms (*Eisenia foetida*) in laboratory tests. *Bull Environ Contam Toxicol* 85:264–268
- Gallagher E, di Giulio R (1991) Effects of 2,4-D dichlorophenoxyacetic acid and picloran on biotransformation, peroxisomal and serum enzyme activities in channel catfish (*Ictalurus punctatus*). *Toxicol Lett* 57:65–72
- Goats GC, Edwards CA (1988) Prediction of field toxicity of chemicals to earthworms by laboratory methods. In: Edwards CA, Neuhauser EF (eds) *Earthworms in waste and environmental assessment*. Academic Publishing, The Hague, pp 283–294
- Gobi M, Gunasekaran P (2010). Effect of butachlor herbicide on earthworm *Eisenia fetida*—its histological perspicuity. *Appl Environ Soil Sci*, Article ID 850758, doi: 10.1155/2010/850758
- Gobi M, Suman J, Ganesan SV (2004) Sublethal toxicity of the herbicide butachlor on the earthworm *Perionyx sansibaricus* and its histological changes. *J Soil Sediment* 5(2):62–86
- Gupta SK, Sundaraman V (1988) Carbaryl induced changes in the earthworm *Pheretima posthumus*. *Indian J Exp Biol* 26:688–693
- Kavitha V, Shoba V, Ramalingam R (2011) Histopathological changes in the intestine of the earthworm *Lampito mauritii* (Kingberg) exposed to sublethal concentration of Monocrotophos. *Int J Recent Sci Res* 2(12):302–305
- Keogh RG, Whitehead PH (1975) Observation on some effect of pasture spraying with benomyl and carbendazim on earthworm activity and litter removal from pasture. *N Z J Exp Agric* 3:103–104

- Kumar Y, Singh K (2013) Distribution of earthworm in different block of Gorakhpur district in eastern Uttar Pradesh. *World Appl Sci J* 21(9):1379–1385
- Lydy MJ, Linck SL (2003) Assessing the impact of triazine herbicides on organophosphate insecticide toxicity to the earthworm *Eisenia foetida*. *Arch Environ Contam Toxicol* 45(3):343–349
- Mosieh YY (2009) Assessing the toxicity of the herbicide isoproturon on *Aporrectodea caliginosa* (Oligochaeta) and its fate in soil ecosystem. *Environ Toxicol* 24(3):396–403
- Munro IA, Carlo GL, Orr JC, Sund KG, Wilson RM, Kennepohl E, Lynch BS, Jablinske M, Lee NL (1992) A comprehensive, integrated review and evaluation of the scientific evidence relating to the safety of the herbicide 2,4-D. *J Am Coll Toxicol* 11:559–664
- Nath G, Singh K, Singh DK (2009) Effect of different combinations of animal dung, and agro/kitchen wastes on growth and development of earthworm *Eisenia foetida*. *Aust J Basic Appl Sci* 3(4):3672–3676
- Nusetti O, Parejo E, Esclapés MM, Rodríguez-Grau J, Marcato L (1999) Acute-sublethal copper effects on phagocytosis and lysozyme activity in the earthworm *Amyntas hawayanus*. *Bull Environ Contam Toxicol* 63:350–356
- Reinecke SA, Reinecke AJ (2007) The impact of organophosphate pesticides in orchards on earthworms in the Western Cape, South Africa. *Ecotoxicol Environ Saf* 66:244–251
- Roberts BL, Dorough HW (1984) Relative toxicity of chemicals to the earthworm. *Environ Toxicol Chem* 3:67–78
- Robidoux PY, Hawari J, Thiboutot S, Ampleman G, Sunahara GI (1999) Acute toxicity of 2,4,6-trinitrotoluene in earthworm (*Eisenia andrei*). *Ecotoxicol Environ Saf* 44(3):11–321
- Rombke J, Garcia MV, Scheffczyk L (2007) Effect of the fungicide Benomyl on earthworm in laboratory tests under tropical conditions. *Arch Environ Contam Toxicol* 53(4):590–598
- Russel RM, Robertson JL, Savin NE (1977) Polo: a new computer programme for profit analysis. *Bull Ent Soc Am* 23:209–213
- Singh K, Kumar Y (2014) Earthworm diversity and ecology. Gene-Tech Books, New Delhi. ISBN 978-81-89729-33-0
- Smith G, Becker H, Edwards PJ, Heimbach F (1992) *Ecotoxicology of earthworms* Intercept, Andover
- Sokal RR, Rohlf FJ (1973) *Introduction of Biostatistics*. W. H. Freeman & Co, San Francisco
- Sorour J, Larink G (2001) Toxic effects of benomyl on the ultrastructure during spermatogenesis of the earthworm *Eisenia foetida*. *Ecotoxicol Environ Saf* 50(3):180–186
- van Gestrel CAM, van Dis WA (1988) The influence of soil characteristics on the toxicity of four chemicals to the earthworm *Eisenia foetida andrei* (Oligochaeta). *Biol Fertil Soil* 6(3):262–265
- Venter JM, Reinecke AJ (1988) Sublethal ecotoxicological effect of Dieldrin on the earthworm *E. foetida* (Oligochaeta). In: Edwards CA, Neuhauser EF (ed) *Earthworms in Wastes and Environmental Management* pp 337–353
- Ville P, Roch P, Cooper EL, Narbonne JF (1997) Immuno-modulator effects of Carbaryl and 2,4 D in the earthworm *Eisenia fetida andrei*. *Arch Environ Contam Toxicol* 32:291–297
- Williamson C (2000) *Earthworms in Turf*, University of Wisconsin garden facts
- Xiao N, Jling B, Ge E, Liu L (2006) The fate of herbicide acetochlor and its toxicity to *Eisenia foetida* under laboratory conditions. *Chemosphere* 62(8):1366–1373
- Zhou SP, Duan CP, FU H, Chen YH, Wang XH, Yu ZE (2007) Toxicity assessment for chlorpyrifos contaminated soil with three different earthworm test methods. *J Environ Sci* 19(7):854–858