

***Toxoplasma gondii* Inhibits Apoptosis in Infected Cells by Caspase Inactivation and NF- κ B Activation**

Ji-Young Kim, Myoung-Hee Ahn, Hye-Sun Jun, Jai-Won Jung, Jae-Sook Ryu, and Duk-Young Min

Department of Parasitology, Institute of Biomedical Science, Hanyang University College of Medicine, Seoul, Korea.

Our experiments aimed to clarify the mechanism by which host cell apoptosis is inhibited by infection with the intracellular protozoan parasite, *Toxoplasma gondii* (*T. gondii*). Mouse spleen cells were cultured in 6-well plates with RPMI 1640/10% FBS at 37°C, in a 5% CO₂ atmosphere. Apoptosis of spleen cells was induced by actinomycin-D (AD) treatment for 1 h prior to infection with *T. gondii*. A variety of assays were used to assess the progression of apoptosis: DNA size analysis on agarose gel electrophoresis, flow cytometry with annexin V/PI staining, and analysis of expression levels of Bcl-2 family and NF- κ B mRNA and proteins by RT-PCR, Western blotting, and EMSA. Additionally, transmission electron microscopy (TEM) was performed to observe changes in cell morphology. Fragmentation of DNA was inhibited in spleen cells treated with AD and *T. gondii* 5 h and 18 h post infection, respectively, and flow cytometry studies showed a decreased apoptotic rates in AD and *T. gondii* treated spleen cells. We observed decreased expression of Bax mRNA and protein, while levels of Bcl-2 mRNA remained constant in spleen cells treated with AD and *T. gondii*. Caspase 3 and PARP were inactivated in cells treated with AD and *T. gondii*, and increased levels of cleaved caspase 8 were also observed. Analysis of EMSA and Western blot data suggests that activation of transcription factor NF- κ B may be involved in the blockade of apoptosis by *T. gondii*. TEM analysis showed nuclear fragmentation and chromatin condensation occurring in spleen cells treated with AD; however, such apoptosis-associated morphological changes were not observed in cells treated with both AD and *T. gondii* tachyzoites. Together, these data show that *T. gondii* infection inhibits AD induced apoptosis via caspase inactivation and NF- κ B activation in

mouse spleen cells.

Key Words: *Toxoplasma*, apoptosis, Bcl-2 family, caspase, NF- κ B, TEM

INTRODUCTION

Toxoplasma gondii (*T. gondii*) is an obligate intracellular protozoan parasite that causes congenital infections in newborns and opportunistic infections in immunocompromised patients, such as those with AIDS. *T. gondii* infects all nucleated cells, eventually causing the host cell to rupture following replication of the tachyzoite.

Cell survival or death, as induced by apoptosis, is a physiological response to cell damage caused by external and internal stimuli, and the exact fate of a cell depends on the balance of pro-apoptotic and anti-apoptotic proteins. Apoptosis limits the growth of infectious agents and plays a crucial role in the immune response. There are two pathways of apoptosis of cells: one triggered by death signals, and another involving the mitochondria. Activation of death receptors triggers the cleavage of caspase 8, whereas release of cytochrome c from mitochondria activates caspase 9; both pathways eventually converge to activate the execution caspase, caspase 3.¹ While the pathogens *Acanthamoeba* sp. and *E. histolytica* can induce host cell apoptosis,^{2,4} the anti- or pro-apoptotic activities associated with *T. gondii* remain controversial. Induction of apoptosis in Peyer's patch T cells and the ocular tissue of mice infected with *T. gondii* has been reported.^{5,6} The Fas/Fas L pathway may play a crucial role in the induction of IFN- γ mediated apoptosis in *N. caninum* and *T. gondii* infected cells,⁷ and infection with *T. gondii* (RH

Received February 3, 2006

Accepted June 7, 2006

This work was supported by the Anti-Communicable Diseases Programs of the National Institute of Health (NIH 348-6111-215) and Ministry of Health and Welfare, Korea.

Reprint address: requests to Dr. Myoung-Hee Ahn, Department of Parasitology, Institute of Biomedical Science, Hanyang University College of Medicine, 17 Haengdang-dong, Seongdong-gu, Seoul 133-791, Korea. Tel: 82-2-2220-0682, Fax: 82-2-2281-6519, E-mail: mhahn@hanyang.ac.kr

strain) results in an over production of IFN- γ and high levels of apoptosis.⁸ Conversely, some intracellular protozoa have the ability to inhibit host cell apoptotic responses. *T. gondii* activates a survival response in infected cells by increasing host resistance to multiple apoptotic stimuli.⁹ In the present study, we focus on the inhibition of apoptosis in mouse spleen cells infected with *T. gondii* using morphological, molecular, and biochemical approaches to characterize the role of NF- κ B in the anti-apoptotic response.

MATERIALS AND METHODS

Toxoplasma gondii

T. gondii tachyzoites (RH) were maintained in the peritoneal cavity of ICR mice (Dae-Han BioLink Co., Eumseong-gun, Chungcheongbuk-do, Korea) for three to four days. Tachyzoites were harvested in RPMI 1640 medium (Invitrogen, Carlsbad, CA, USA) from the peritoneal cavity and separated from peritoneal cells by repeated low and high speed centrifugation ($40 \times g$ for 5 min and $650 \times g$ for 10 min).

Treatment of mouse spleen cells with *T. gondii* infection and actinomycin-D

Spleen cells in RPMI 1640 media were harvested from BALB/c mice and pelleted by centrifugation ($650 \times g$ for 10 min). Cells were resuspended in lysis buffer (0.83% NH_4Cl -0.01M Tris-HCl) in order to remove red blood cells. Mouse spleen cells (5×10^6 cells/well) were cultured in 6-well plates with RPMI 1640 media containing 2 mM L-glutamine, 2 mg/mL NaHCO_3 , 10% fetal bovine serum (FBS, Cambrex, Walkersville, MD, USA), 100 U/mL penicillin, and 100 $\mu\text{g}/\text{mL}$ streptomycin at 37°C , 5% CO_2 atmosphere. Cells were infected with 5×10^7 *T. gondii* tachyzoites and/or treated with 5-10 $\mu\text{g}/\text{mL}$ actinomycin-D (AD, Sigma-Aldrich, St. Louis, MO, USA) for 5 h and 18 h, respectively.

DNA size analysis by agarose gel electrophoresis

Infected and non-infected mouse spleen cells

were harvested by centrifugation and resuspended in NET buffer containing 10% SDS, DNA-free Rnase, and proteinase K. The cells were then centrifuged at $12,000 \times g$ for 5 min at 4°C , and DNA was extracted from the cell pellet with phenol, phenol/isoamyl alcohol, and chloroform, followed by precipitation with 100% ethanol and 10 M NH_4OAc . The DNA was dissolved in TE buffer and quantified by A260 on a spectrophotometer (Perkin Elmer, Wellesley, MA, USA). DNA samples were analyzed on a 2% agarose gel stained with 1 $\mu\text{g}/\text{mL}$ ethidium bromide.

Flow cytometry

Cells were stained with annexin V using the Annexin V/PI kit (R&D systems, Minneapolis, MN, USA), and apoptotic cells identified and quantified by flow cytometry. Briefly, spleen cells (1×10^6) were washed in PBS and incubated with 10 X binding buffer, propidium iodide (PI), annexin V-FITC, and dH_2O (total 100 μL) for 15 min at room temperature. The apoptotic cells were analyzed on a fluorescence-activated cell sorter (FACS, Becton Dickinson-FACScalibur, Cockeysville, MD, USA).

Reverse transcriptase (RT)-PCR

Bcl-2 family messenger RNA was identified by RT-PCR using message specific primers. Mouse spleen cells were incubated with AD and AD/*T. gondii* for 30 min, 60 min or 120 min, and total cellular RNA extracted using TRIzol (Invitrogen), chloroform, and isopropyl alcohol. Reverse transcription was performed with an oligo dT primer, and cDNAs were amplified by PCR using the primers listed in Table 1. PCR products were resolved on a 2% agarose gel stained with 1 $\mu\text{g}/\text{mL}$ ethidium bromide.

Western blot analysis of Bcl-2 family proteins and caspases

The abundance of Bcl-2 family proteins (Bax, Bcl-2, Bcl-X_L), caspase 3, caspase 8, PARP and actin in mouse spleen cells was analyzed by Western blotting. Spleen cells (5×10^5) were resuspended in lysis buffer (20 mM Tris-HCl, PH

Table 1. Primer Sequences and RT-PCR Conditions

Gene of interest	Primer	PCR conditions
Bax (S)	5'-AAGCTGAGCGAGTGTCTCCGGCG-3'	95°C (10 min) - 95°C (30 s) - 64°C (30 s) -72°C (30 s)
(AS)	5'-GCCACAAAGATGGTCTGCC-3'	- 72°C (7 min)
Bcl-2 (S)	5'-CAGCTGCACCTGACGCCCTT-3'	95°C (10 min) - 95°C (1 min) - 52°C (1min) -72°C (1 min)
(AS)	5'-GCCTCCGTTATCCTGGATCC-3'	- 72°C (7 min)
Bcl-X _L (S)	5'-TGGAGTAAACTGGGGTTCGCATCG-3'	95°C (10 min) - 95°C (1 min) - 50°C (1 min) - 72°C (1 min)
(AS)	5'-AGCCACCGTCATTCCCCTCAGG-3'	- 72°C (7 min)
β-actin (S)	5'-CCAGAGCAAGAGAGGTATCC-3'	95°C (10 min) - 95°C (45 s) - 58°C (2 min 30 s)
(AS)	5'-CTGTGGTGGTGAAGCTGTAG-3'	- 72°C (10 min)

7.5, 60 mM beta-glycerophosphate, 10 mM EDTA, 10 mM MgCl₂, 10 mM NaF, 2 mM DTT, 1 mM Na₃VO₄, 1 mM APMSF, 1% Np-40, 5 µg/mL leupeptin). Following a 40 min incubation on ice, the lysates were combined with 2 X sample buffer (0.125 M Tris-HCl, pH 6.8, 4% SDS, 20% glycerol, 10% 2-mercaptoethanol, 0.2% bromphenol blue) and boiled for 5 min, samples were resolved on a SDS-polyacrylamide gel. After semi-dry transfer to a PVDF membrane (Amersham Pharmacia Biotech, Uppsala, Sweden), ponceau S staining was used to confirm loading of equal amounts of total protein loaded in each lane. The membranes were blocked (5% nonfat dried milk, 0.1% Tween 20, 100 mM NaCl, 10 mM Tris-HCl, PH 7.6) for 1 h and incubated with the appropriate primary antibody: monoclonal mouse anti-Bax antibody (BD PharMingen, San Diego, CA, USA, 2 µg/mL), polyclonal rabbit anti-Bak antibody (BD PharMingen, 2 µg/mL), monoclonal mouse anti-Bcl-2 antibody (Oncogene, San Diego, CA, USA, 2 µg/mL), polyclonal rabbit anti-caspase 3 antibody (Cell Signaling, Beverly, MA, USA, 1:100), monoclonal mouse anti-poly (ADP-ribose) polymerase (PARP) antibody (BD PharMingen, 1:200), polyclonal rabbit anti-caspase 8 antibody (Santa Cruz Biotechnology, Santa Cruz, CA, USA, 1:200), or polyclonal rabbit anti-actin antibody (Santa Cruz Biotechnology, 1:200). Bound primary antibodies were visualized with an ECL anti-mouse IgG peroxidase-linked species-specific F(ab')₂ fragment (Amersham Pharmacia Biotech, 1:10,000) or an ECL anti-rabbit IgG peroxidase-linked species-specific F(ab')₂ fragment (1:200,000) and the ECL PLUS chemiluminescence detection system em-

ployed according to the manufacture's instructions (Amersham Pharmacia Biotech).

Western blot for NF-κB and IκBα

Nuclear and cytoplasmic fractions were prepared from spleen cells.¹⁰ Monoclonal mouse IκBα antibody (Cell signaling, 1:100), rabbit phospho-NF-κB p65 antibody (Cell signaling, 1:50), NF-κB p65 antibody (Cell signaling, 1:100) and mouse anti-actin antibody (Chemicon international, Temecula, CA, USA, 1:1,000) were used for detection of NF-κB and IκBα by Western blot as described above.¹¹

Electrophoresis motility shift assay (EMSA)

Nuclear extracts from spleen cells were prepared as previously described.¹⁰ Protein concentrations of the nuclear extracts were determined by Bradford assay, and EMSAs were performed according to the manufacture's instructions (Promega, Madison, WI, USA). Briefly, 5 µg of nuclear extract was incubated with a (γ³²P)-labeled oligonucleotide probe corresponding to the consensus NF-κB binding site for 30 min at room temperature. After incubation, bound and free DNAs were resolved on a 5% polyacrylamide gel¹² run at 100 V for 3 h, dried, and subjected to autoradiography for 12 h to 48 h.¹³

Transmission electron microscopy (TEM)

The apoptotic changes of mouse spleen cells treated with AD (10 µg/mL) and/or infected with

T. gondii (3×10^6 , cell:tachyzoite = 1:10) for 6 h in 6 well plates were visualized by TEM performed using standard methods. Briefly, spleen cells were fixed in 3% glutaraldehyde (PH 7.8) for 3 h at room temperature and post-fixed in 1% osmium tetroxide for 1.5 h at 4°C. After serial dehydration with 50%, 70% and 100% ethanol, specimens were embedded in Epon resin. Ultrathin sections were loaded onto copper grids; stained with uranyl acetate and lead citrate, and visualized with a transmission electron microscope H-7600s (Hitachi, Tokyo, Japan).

RESULTS

DNA fragmentation in spleen cells was suppressed by *T. gondii*

Spleen cells were incubated with AD and/or *T. gondii* tachyzoites for 5 h and 18 h respectively at 37°C. DNA was then extracted from spleen cells, resolved on a 2% agarose gel, and stained with ethidium bromide. DNA laddering was observed in mouse spleen cells treated with AD for 5 h (data not shown) and 18 h (Fig. 1, lanes 1 & 2).




Fig. 1. Fragmentation patterns of DNA from mouse spleen cells treated with actinomycin-D and/or *T. gondii* for 18 h. Fragmentation of DNA was down regulated in cells treated with AD/*T. gondii* (lane 6). Spleen cells infected with *T. gondii* exhibited only weak DNA fragmentation (lane 5). Lane 1. spleen cells treated with actinomycin-D (5 μ g/mL); 2. spleen cells treated with actinomycin-D (10 μ g/mL); 3. normal spleen cells for 18h incubation; 4. fresh normal spleen cells; 5. *T. gondii* infected spleen cells (cell:parasite = 1:10); 6. spleen cells infected with *T. gondii* and treated with actinomycin-D (5 μ g/mL). In repeated experiments, similar results were shown.

Apoptosis was down regulated in mouse spleen cells treated with AD and infected with *T. gondii* (Fig. 1, lane 6). Spleen cells infected with *T. gondii* only showed weak DNA fragmentation (Fig. 1, lane 5). DNA laddering was observed in spleen cells for 18h incubation, but not in freshly isolated cells (Fig. 1, lanes 3 & 4). Similar results were found upon repeated experiments.

Inhibition of apoptosis in spleen cells infected with *T. gondii* was observed by flow cytometric analysis

Mouse spleen cells treated with AD and/or *T. gondii* infection were pooled, incubated with annexin-FITC V and PI, and analyzed by flow cytometry. Early apoptotic cells have increased annexin V, but not PI staining. Late apoptotic or necrotic cells stain with both annexin V and PI. Mouse spleen cells cultured with AD, AD/*T.*




Fig. 2. Flow cytometric analysis of apoptosis in mouse spleen cells treated with actinomycin-D and/or *T. gondii* for 18 h. (%) indicates level of apoptosis. (A) actinomycin-D (5 μ g/mL), 80.3%; (B) actinomycin-D (10 μ g/mL), 81.2%; (C) normal spleen cells for 18h incubation, 39.6%; (D) freshly isolated spleen cells, 23.8%; (E) *T. gondii* infected, 18.2%; (F) *T. gondii* and actinomycin-D treated spleen cells, 30.1%. Flow cytometry was repeated more than three times and similar results were observed.

gondii and freshly isolated cells showed apoptosis of 43.6%, 37.0%, and 15.8%, respectively after incubation for 5 h (data not shown). The quantity of apoptotic spleen cells were 80.3%, 30.1%, and 23.8%, respectively, after incubation for 18 h (Fig. 2, A, D, F). Similar results were found upon repeated experiments.

Levels of Bax mRNA and protein expression in spleen cells were reduced upon treatment with *T. gondii*

RT-PCR analysis of Bcl-2 family genes was performed on mouse spleen cells incubated with AD, or AD/*T. gondii* for 30 min, 60 min or 120 min at 37°C. Expression of pro-apoptotic Bax mRNA was down regulated in AD/*T. gondii* infected spleen cells (Fig. 3). Western blot analysis showed that the production of Bax protein was decreased in spleen cells treated with AD and infected with *T. gondii*, while Bak protein levels remained unchanged (Fig. 4). Expression levels of anti-apoptotic Bcl-2 and Bcl-X_L mRNAs remained constant in spleen cells treated with AD and infected with *T. gondii* (Fig. 3). Spleen cells infected with *T. gondii* showed indistinct Bax and Bcl-2 mRNA levels. Moreover, Bcl-2 protein did not appear in AD/*T. gondii* infected cells.

Caspase 3 and PARP protein activation was suppressed in *T. gondii* infected spleen cells

Activation of caspase 3 and PARP proteins was




Fig. 3. RT-PCR analysis of Bcl-2 family genes in mouse spleen cells. Expression of pro-apoptotic Bax mRNA was down regulated and levels of anti-apoptotic Bcl-2 and Bcl-X_L mRNA remained constant in spleen cells treated with AD/*T. gondii*. Spleen cells were co-cultured with either actinomycin-D and/or *T. gondii* for 30 min, 60 min, or 120 min followed by extraction of total cellular RNA. (M): marker; (-): negative; (AD): actinomycin-D (5 µg/mL); (AD + *T.g*): actinomycin-D + *T. gondii* infection.

suppressed in spleen cells treated with AD and infected with *T. gondii*. Down regulated, cleaved caspase 3 at 120 min post infection and reduced production of PARP in *T. gondii* infected cells were detected by Western blotting (Fig. 4). Although cleaved caspase 8 was observed in spleen cells treated with AD and infected with *T. gondii*, it was not expressed in spleen cells treated with AD only (Fig. 4).

Activation of NF-κB in spleen cells infected with *T. gondii*

We examined the role of NF-κB in the anti-apoptotic response of *T. gondii* infected spleen cells by EMSA and Western blotting. NF-κB activity was apparent at 120 min in spleen cells treated with AD and infected with *T. gondii*, which decreased by 180 min (Fig. 5A). *T. gondii* infected spleen cells showed more activated NF-κB than uninfected cells at 60 min on EMSA. A similar pattern of NF-κB activity in total cell and nuclear extracts was observed by Western blot (Fig. 5B). Decreased IκBα was associated with increased NF-κB activity.




Fig. 4. Western blot analysis of Bcl-2 family proteins, caspase 3, caspase 8 and PARP in mouse spleen cells in the presence or absence of *T. gondii* and actinomycin-D after incubation for 0 min, 60 min or 120 min. Production of Bax protein was decreased in spleen cells treated with AD/*T. gondii*, while levels of Bak protein was unchanged. Down regulation of cleaved caspase 3 at 120 min post infection and reduced production of PARP in *T. gondii* infected cells was detected by Western blotting. (AD): actinomycin-D (5 µg/mL); (AD + *T.g*): actinomycin-D + *T. gondii* infection.




Fig. 5. Increased NF- κ B and decreased I κ B α of mouse spleen cells by *T. gondii* infection was observed. (A) NF- κ B activity was apparent at 120 min in spleen cells treated with AD and infected with *T. gondii* on EMSA. (B) Decreased cytosolic I κ B α , pNF- κ B and increased nuclear NF- κ B proteins were detected in spleen cells treated with AD/*T. gondii* by Western blot. (+): TNF- α (20 ng/mL); (AD): actinomycin-D; (AD + *T. gondii*): actinomycin-D + *T. gondii*; (CP): cytosolic proteins; (NP): nuclear proteins.

TEM shows inhibition of apoptosis in *T. gondii* infected spleen cells

Condensation of nuclear chromatin and nuclear fragmentation was observed in spleen cells treated with AD (Fig. 6A & B); however, these two events were not observed in spleen cells treated with AD and infected with *T. gondii* tachyzoites (Fig. 6C & D). In spleen cells infected with *T. gondii* only, neither nuclear condensation nor fragmentation was observed (Fig. 6E).

DISCUSSION

Apoptosis, a mechanism of programmed cell death, is a result of both physical and pathological events. Transduction of pro-apoptotic stimuli via signaling pathways results in the activation of the caspase family of cysteine proteases. Bcl-2 family pro-apoptotic proteins promote cell death by inducing mitochondrial damage, cytochrome c release, and subsequent caspase activation, and




Fig. 6. Ultrastructural morphology of mouse spleen cells treated with actinomycin-D and *T. gondii* after cultivation in vitro for 6h. Condensation of nuclear chromatin and fragmentation of the nucleus were observed in spleen cells treated with AD (A & B); however, these events were not observed in cells treated with AD/*T. gondii* tachyzoites (C & D) or *T. gondii* only(E). A, B: Spleen cells treated AD (5 μ g/mL); C, D: Spleen cells treated with AD and *T. gondii*; E: Spleen cells treated with *T. gondii*; F: Normal spleen cells. A, B, E and F: 6000 \times , C: 7000 \times , D: 4000 \times , Bar = 2 μ m. The arrows and arrow heads indicate apoptotic spleen cells (A & B) and *T. gondii* tachyzoites, respectively (C, D & E).

the Bcl-2 family of anti-apoptotic proteins inhibit apoptosis by preventing the release of cytochrome c; however, the mechanism by which parasites utilize this method of apoptotic inhibition has not been elucidated. Down-regulation of host cell apoptosis may also provide an explanation for the persistence of intracellular pathogens. *T. gondii* has several strategies for inhibiting the initiation of the apoptotic cascade. In the present study, we found that apoptosis was not induced in mouse spleen cells infected with *T. gondii* when compared to uninfected controls (data not shown); however, apoptosis was induced in spleen cells by treatment with AD only. One biochemical analysis of apoptosis activity is the presence of oligonu-

cleosome-sized fragments of DNA, which produce ladders when analyzed with agarose gel electrophoresis. Our results indicated that the induction of DNA fragmentation in AD treated spleen cells was inhibited by *T. gondii* infection (Fig. 1). Apoptotic cells can be detected by the binding of fluorescent labeled annexin V to the plasma membrane. As such, early and late apoptotic cells were measured by flow cytometric analysis after double staining with FITC-conjugated annexin V and PI.¹⁴ While flow cytometric analysis could not differentiate late apoptosis and necrosis of cells, it demonstrated that concomitant infection with *T. gondii* abrogated AD induced apoptosis. Specifically, in the AD treated population, 80.3% of cells were apoptotic, as opposed to 30.1% of the cells treated with both AD and *T. gondii* after incubation for 18 h (Fig. 2). Proteins of the Bcl-2 family with pro-apoptosis or anti-apoptotic functions reside in the outer mitochondrial membrane. Bax, a pro-apoptotic protein, triggers the release of cytochrome c from mitochondria.¹ Conversely, Bcl-2 and Bcl-X_L are capable of blocking the release of cytochrome c, thereby preventing the activation of caspase 9. The induction of apoptosis by *T. vaginalis* is associated with a reduction of expression of anti-apoptotic Bcl-X_L protein, but not Bcl-2, and the activation of caspase 9 suggests that mitochondria play a role in *T. vaginalis*-induced RAW 264.7 programmed cell death.¹³ RT-PCR analysis demonstrated that *T. gondii* infection promotes the activation of anti-apoptotic Bcl-2 family members but not pro-apoptotic pathways.¹⁵ Our studies showed that the expression of Bax mRNA and protein was indistinct in AD/*T. gondii*-treated mouse spleen cells, compared to uninfected cells (Fig. 3). Likewise, expression of Bcl-2 and Bcl-X_L mRNA and Bak protein were unchanged in mouse spleen cells treated with AD and infected with *T. gondii* (Figs. 3 & 4). These results suggested that the anti-apoptotic Bcl-2 protein was not affected in *T. gondii* infection and that the other anti-apoptotic protein, Bcl-X_L may be involved. Generally *T. gondii* infection promotes the suppression of pro-apoptotic Bax, but not anti-apoptotic Bcl-2 proteins (Fig. 4).

Caspase 3 is a critical downstream member of the caspase family, and is an essential effector of apoptosis. The resistance of *T. gondii* infected cells

to apoptosis may be attributed to prevention of caspase 3 activity.^{17,16} In the present study, inactivation of caspase 3 at 120 min post infection and decreased PARP were present in cells treated AD and infected with *T. gondii* as determined by Western blotting (Fig. 4). Cleaved caspase 8, which was present in cells treated with AD and *T. gondii* infection, may have induced a weak apoptosis response. We propose that AD did not trigger apoptosis via the caspase 8 pathway and that infection by the parasite induced apoptosis slightly. Activation of the caspase cascade leads to cleavage of a variety of target proteins including PARP, nuclear lamins, and protein kinase C (PKC); both PARP protein levels and cleavage of nuclear target proteins are down regulated by *T. gondii*.¹⁶ Our results showed that *T. gondii* mediates inhibition of apoptosis by a blockade of Bax expression and caspase 3 activation. Inhibition of apoptosis by *T. gondii* is regulated at the transcriptional level of NF- κ B, as infection with *T. gondii* induces the activation of NF- κ B and the transcription of anti-apoptotic genes.¹⁷⁻²¹ Phosphorylation of I κ B activates and promotes NF- κ B translocation to the nucleus, where NF- κ B associates with the transcription of proinflammatory mediators. Further, expression of NF- κ B is correlated with anti-apoptotic protein expression after *T. gondii* infection. In this experiment, spleen cells treated with AD and infected with *T. gondii* showed increased NF- κ B DNA binding activity by EMSA (Fig. 5). Condensation of chromatin, disruption of nuclear architecture, vacuolation of the cytoplasm, membrane blebbing, detachment from substrate, and cell shrinkage are some of the morphological changes associated with apoptosis. We observed some of these morphologic changes in cells treated with AD and/or *T. gondii* using TEM. Condensed chromatin and fragmented nuclei of cells were apparent 6 h after treatment of the cells with AD; however, there were no apoptotic changes in cells treated with AD and *T. gondii* (Fig. 6C & D). In contrast to the inhibition of apoptosis in *T. gondii* infected cells, parasite infection induced apoptosis has been reported in neighboring non-invaded cells.^{5,6,22} However, we were unable to identify such changes in uninfected mouse spleen cells on TEM (Fig. 6D & E).

In conclusion, infection with *T. gondii* induces

an anti-apoptotic effect, which was associated with inactivation of caspase 3 and PARP. Host cell mitochondria around the parasitophorous vacuole may be involved in anti-apoptotic activity by *T. gondii*.¹⁸ To survive in host cells, parasitic protozoa modulate their host's apoptotic response. These findings suggest that mouse spleen cells infected with *T. gondii* resisted AD treatment, an apoptotic stimuli, by inactivating the caspase cascade and suppressing Bax protein expression through NF- κ B activation. These results provide direction for further investigation of the mechanism of inhibition of apoptosis by *T. gondii* in host cells.

REFERENCES

- Payne JM, Molestina RE, Sinai AP. Inhibition of caspase activation and requirement for NF- κ B function in the *Toxoplasma gondii*-mediated blockade of host apoptosis. *J Cell Sci* 2003;116:4345-58.
- Shin HJ, Cho MS, Kim HI, Lee M, Park S, Sohn S, et al. Apoptosis of primary-culture rat microglial cells induced by pathogenic *Acanthamoeba* spp. *Clin Diagn Lab Immunol* 2000;7:510-4.
- Sim S, Kim KA, Yong TS, Park SJ, Im KI, Shin MH. Ultrastructural observation of human neutrophils during apoptotic cell death triggered by *Entamoeba histolytica*. *Korean J Parasitol* 2004;42:205-8.
- Sim S, Yong TS, Park SJ, Im KI, Kong Y, Ryu JS, et al. NADPH oxidase-derived reactive oxygen species-mediated activation of ERK1/2 is required for apoptosis of human neutrophils induced by *Entamoeba histolytica*. *J Immunol* 2005;174:4279-88.
- Liesenfeld O, Kosek JC, Suzuki Y. Gamma interferon induces Fas-dependent apoptosis of Peyer's patch T cells in mice following peroral infection with *Toxoplasma gondii*. *Infect Immun* 1997;65:4682-9.
- Hu MS, Schwartzman JD, Yeaman GR, Collins J, Seguin R, Khan IA, et al. Fas-FasL interaction involved in pathogenesis of ocular toxoplasmosis in mice. *Infect Immun* 1999;67:928-35.
- Nishikawa Y, Makala L, Otsuka H, Mikami T, Nagasawa H. Mechanisms of apoptosis in murine fibroblasts by two intracellular protozoan parasites, *Toxoplasma gondii* and *Neospora caninum*. *Parasite Immunol* 2002;24:347-54.
- Gavrilescu LC, Denkers EY. IFN- γ overproduction and high level apoptosis are associated with high but not low virulence *Toxoplasma gondii* infection. *J Immunol* 2001;167:902-9.
- Nash PB, Purner MB, Leon RP, Clarke P, Duke RC, Curiel TJ. *Toxoplasma gondii*-infected cells are resistant to multiple inducers of apoptosis. *J Immunol* 1998;160:1824-30.
- Kim JS, Kim JM, Jung HC, Song IS. Caspase-3 activity and expression of Bcl-2 family in human neutrophils by *Helicobacter pylori* water-soluble proteins. *Helicobacter* 2001;6:207-15.
- Elewaut D, DiDonato JA, Kim JM, Truong F, Eckmann L, Kagnoff MF. NF- κ B is a central regulator of the intestinal epithelial cell innate immune response induced by infection with enteroinvasive bacteria. *J Immunol* 1999;163:1457-66.
- Kim JM, Oh YK, Kim YJ, Cho SJ, Ahn MH, Cho YJ. Nuclear factor-kappa B plays a major role in the regulation of chemokine expression of HeLa cells in response to *Toxoplasma gondii* infection. *Parasitol Res* 2001;87:758-63.
- Chang JH, Ryang YS, Morio T, Lee SK, Chang EJ. *Trichomonas vaginalis* inhibits proinflammatory cytokine production in macrophages by suppressing NF- κ B activation. *Mol Cells* 2004;18:177-85.
- Kim JM, Eckmann L, Savidge TC, Lowe DC, Witthoft T, Kagnoff MF. Apoptosis of human intestinal epithelial cells after bacterial invasion. *J Clin Invest* 1998;102:1815-23.
- Goebel S, Luder CG, Gross U. Invasion by *Toxoplasma gondii* protects human-derived HL-60 cells from actinomycin D-induced apoptosis. *Med Microbiol Immunol* 1999;187:221-6.
- Goebel S, Gross U, Luder CG. Inhibition of host cell apoptosis by *Toxoplasma gondii* is accompanied by reduced activation of the caspase cascade and alterations of poly (ADP-ribose) polymerase expression. *J Cell Sci* 2001;114:3495-505.
- Heussler VT, Kuenzi P, Rottenberg S. Inhibition of apoptosis by intracellular protozoan parasites. *Int J Parasitol* 2001;31:1166-76.
- Molestina RE, Payne TM, Coppens I, Sinai AP. Activation of NF-kappaB by *Toxoplasma gondii* correlates with increased expression of antiapoptotic genes and localization of phosphorylated IkkappaB to the parasitophorous vacuole membrane. *J Cell Sci* 2003;116:4359-71.
- Denkers EY, Butcher BA, Del Rio L, Kim L. Manipulation of mitogen-activated protein kinase/nuclear factor-kappa B-signaling cascades during intracellular *Toxoplasma gondii* infection. *Immunol Rev* 2004;201:191-205.
- Carmen JC, Hardi L, Sinai AP. *Toxoplasma gondii* inhibits ultraviolet light-induced apoptosis through multiple interactions with the mitochondrion-dependent programmed cell death pathway. *Cell Microbiol* 2006;8:301-15.
- Sinai AP, Payne TM, Carmen JC, Hardi L, Watson SJ, Molestina RE. Mechanisms underlying the manipulation of host apoptotic pathways by *Toxoplasma gondii*. *Int J Parasitol* 2004;34:381-91.
- James ER, Green DR. Manipulation of apoptosis in the host-parasite interaction. *Trends Parasitol* 2004;20:280-7.