

Transcriptional regulation of the Th17 immune response by IKK α

Li Li,¹ Qingguo Ruan,¹ Brendan Hilliard,¹ Jennifer DeVirgiliis,¹ Michael Karin,² and Youhai H. Chen¹

¹Department of Pathology and Laboratory Medicine, University of Pennsylvania School of Medicine, Philadelphia, PA 19104

²Department of Pharmacology, University of California, San Diego, San Diego, CA 92093

Th17 cells are a subset of T cells that play crucial roles in the pathogenesis of many inflammatory diseases. We report here the identification of IKK α (inhibitor of NF- κ B kinase- α) as a key transcriptional regulator of the Th17 lineage. T cells expressing a nonactivatable form of IKK α were significantly compromised in their ability to produce IL-17 and to initiate neural inflammation. IKK α is present in the nuclei of resting CD4⁺ T cells. Upon Th17 differentiation, IKK α selectively associated with the *Il17a* locus, and promoted its histone H3 phosphorylation and transcriptional activation in a NF- κ B-independent manner. These findings indicate that nuclear IKK α maintains the Th17 phenotype by activating the *Il17a* gene.

CORRESPONDENCE

Youhai H. Chen:
yhc@mail.med.upenn.edu

Abbreviations used: ChIP, chromatin immunoprecipitation; EAE, experimental autoimmune encephalomyelitis; hnRNA, heterogeneous nuclear RNA; IKK α , inhibitor of NF- κ B kinase- α ; MOG, myelin oligodendrocyte glycoprotein; ROR, retinoid-related orphan receptor; Stat3, signal transducer and activator of transcription 3.

For many years, CD4⁺ T helper (Th) cells have been classified into two major types, Th1 and Th2 cells (Romagnani, 1997). Th1 cells express IFN- γ and control cellular immunity, whereas Th2 cells produce IL-4, IL-5, and IL-13 and regulate humoral immunity. Recently, a new helper T cell subset, Th17 (also known as Th1), which produces IL-17A, IL-17E, IL-21, and IL-22, but not IFN- γ or IL-4, has been defined (Cua et al., 2003; Langrish et al., 2005; Veldhoen and Stockinger, 2006; Ivanov et al., 2006; Weaver et al., 2006; Bettelli et al., 2006; Sutton et al., 2006). Unlike Th1 and Th2 cells, Th17 cells are considered to be “proinflammatory” because they are involved primarily in mediating inflammatory diseases and immune defense against extracellular bacteria (Langrish et al., 2005; Bettelli et al., 2006; Ivanov et al., 2006; Veldhoen and Stockinger, 2006; Weaver et al., 2006; Sutton et al., 2006). Th17 cells can be generated in vitro by activating naive T cells in the presence of IL-6/IL-21 and TGF- β (Weaver et al., 2006; Bettelli et al., 2007). IL-6 acts in a signal transducer and activator of transcription 3 (Stat3)-dependant manner to induce IL-21, IL-23 receptor, retinoid-related orphan receptor (ROR) γ T, and ROR α expression (Yang et al., 2007; Dong, 2008; Yang et al., 2008). Upon binding to IL-23, which is normally produced by macrophages and dendritic cells, IL-23 receptor promotes the survival of Th17 cells and maintains its differentiated phenotype (Cua et al., 2003;

Langrish et al., 2005). Transcriptionally, ROR γ T and ROR α are considered to be master regulators of Th17 differentiation, as T-bet and GATA3 are to Th1 and Th2 cells, respectively (Ivanov et al., 2006; Dong, 2008; Yang et al., 2008). Moreover, similar to Th1 and Th2 cells in which the *Ifng* and *Il4* loci are selectively activated, respectively, differentiated Th17 cells exhibit unique epigenetic modifications of the *Il17a* locus (Akimzhanov et al., 2007). However, the nuclear factors that are responsible for *Il17a* locus activation are not well understood.

The inhibitor of nuclear factor- κ B kinase- α (IKK α) is a member of the IKK family, which regulates multiple biological processes through either NF- κ B-dependent or -independent mechanisms (Häcker and Karin, 2006). IKK α can phosphorylate NF- κ B2 (p100), leading to the generation of p52, which dimerizes with RelB, to activate target genes involved in lymphoid organ development (Senftleben et al., 2001). However, it has recently been recognized that IKK α can also regulate gene expression in an NF- κ B-independent manner. Unlike IKK β , IKK α contains a nuclear localization sequence. It was suggested that in the nucleus, IKK α phosphorylates histone H3 at serine (Ser) 10

© 2011 Li et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

position, a prerequisite event for subsequent histone acetylation and gene transcription (Anest et al., 2003; Yamamoto et al., 2003). However, H3 Ser10 phosphorylation may simply serve as an indicator of an active “open” chromatin structure, and its dependence on IKK α may indicate that IKK α is required for establishment of the active chromatin state. More recently, IKK α kinase activity was shown to be required in the nucleus for repression of certain genes (Sil et al., 2004; Luo et al., 2007). Additionally, IKK α can also regulate epidermal keratinocyte differentiation through a kinase-independent mechanism (Hu et al., 2001). The NF- κ B-independent functions of IKK α remain to be fully established.

To determine whether IKK α is required for T cell differentiation and T cell-mediated autoimmunity, we studied myelin oligodendrocyte glycoprotein (MOG)-induced experimental autoimmune encephalomyelitis (EAE) in *Ikk α ^{AA}* knock-in mice (Senfleben et al., 2001). The knock-in allele specifies expression of a variant IKK α protein, in which the activating phosphorylation sites, Ser176 and Ser180, are replaced by two alanines (AA), thereby abolishing the activation of its kinase activity (Bonizzi et al., 2004; Lawrence et al., 2005). We found that *Ikk α ^{AA}* mutant mice were refractory to EAE, and *Ikk α ^{AA}* CD4⁺ T cells were defective in their Th17 cell differentiation. We then discovered that IKK α controls Th17 lineage commitment by maintaining the activation state of the *Il17a* locus, in an NF- κ B-independent manner.

RESULTS

Autoimmune encephalomyelitis is markedly reduced in *Ikk α ^{AA}* knock-in mice

To determine the roles of IKK α in T cell-mediated inflammation, we immunized WT and *Ikk α ^{AA}* C57BL/6 mice with MOG peptide, and monitored daily for clinical signs of EAE (Fig. 1 a). Although all WT mice developed EAE, only 77.8% of *Ikk α ^{AA}* mice developed the disease. The disease severity was also significantly reduced in the *Ikk α ^{AA}* group (maximal clinical score, 1.2 \pm 0.8) as compared with the WT group (maximal clinical score, 3.3 \pm 0.4; *P* < 0.05). The day of disease onset was increased from 11.9 \pm 0.9 d in the WT group to 15 \pm 1.8 d in the *Ikk α ^{AA}* group (*P* < 0.01; Fig. 1 a). Consistent with these clinical findings, histological examination of spinal cord sections revealed significant differences in the degree of inflammation between the two groups. In the WT group, multiple inflammatory foci were observed, with extensive leukocyte infiltration in the white matter (Fig. 1 d, top). In contrast, leukocyte infiltration in *Ikk α ^{AA}* spinal cords was much less pronounced (Fig. 1 d, bottom). Thus, IKK α kinase activity contributes to the development of EAE.

Loss of IKK α kinase activity in CD4⁺ T cells is responsible for the abrogated EAE development in *Ikk α ^{AA}* mice

Ikk α ^{AA} mice have severe defects in secondary lymphoid organogenesis and develop only rudiments of certain lymph nodes (Senfleben et al., 2001). This defect is caused by the loss of IKK α activity in nonhematopoietic stromal cells, other than hematopoietic cells (Senfleben et al., 2001; Bonizzi et al., 2004). To separate the effect of IKK α on lymphoid organogenesis from its effect on EAE, we studied the disease in irradiated WT C57BL/6 mice that




Figure 1. IKK α expressed by T cells is required for the development of autoimmune encephalomyelitis. WT (*n* = 9) and *Ikk α ^{AA}* (*n* = 6) C57BL/6 mice (a), WT C57BL/6 mice that received WT (*n* = 8) or *Ikk α ^{AA}* (*n* = 8) bone marrow (b), and *Rag1*^{-/-} C57BL/6 mice that received WT (*n* = 3) or *Ikk α ^{AA}* (*n* = 3) CD4⁺ T cells (c) were immunized with MOG to induce EAE as described in the Materials and methods. Data presented are means \pm SD of EAE scores. The differences between two groups are statistically significant (*P* < 0.01) for all panels. (d) Mice were sacrificed at the end of the experiments, and their spinal cords and brains were sectioned and stained with luxol fast blue and cresyl violet. (d, top) The spinal cord of a WT mouse from panel a with a clinical score of 4; (bottom) the spinal cord of an *Ikk α ^{AA}* mouse with no signs of EAE. Data presented are representative of three independent experiments.

had received bone marrow from either WT or *Ikk α ^{AA}* mice (Fig. 1 b). In the chimeric mice, 80–90% of leukocytes were derived from donor bone marrow as determined by flow cytometry (unpublished data). Importantly, mice that received *Ikk α ^{AA}* bone marrow developed significantly less severe EAE than those reconstituted with WT cells (maximal disease score, 3.0 ± 0.5 vs. 4.3 ± 0.5 ; $P < 0.01$). Disease onset was also delayed from 11.0 ± 0 in the WT to 14.3 ± 0.9 d ($P < 0.01$) in the *Ikk α ^{AA}* group (Fig. 1 b). Therefore, loss of IKK α kinase activity in hematopoietic cells alone is sufficient to compromise the development of EAE.

Although MOG-induced EAE is a T cell-dependent disease, other hematopoietic cell types also contribute to the development of the disease. To directly test the T cell-specific function of IKK α in EAE, we studied disease development in *Rag1*^{-/-} mice that had received WT or *Ikk α ^{AA}* CD4⁺ T cells (Fig. 1 c). *Rag1*^{-/-} mice receiving WT CD4⁺ T cells started to develop clinical signs of EAE at 25.3 ± 0.6 d after MOG immunization, and reached a maximal disease score of 3.7 ± 0.3 . Remarkably, although *Rag1*^{-/-} mice received *Ikk α ^{AA}* CD4⁺ T cells developed EAE with only a slight delay relative to mice reconstituted with WT cells (28.3 ± 0.5 d; $P < 0.01$), the severity of the disease was significantly reduced (maximal

disease score: 0.7 ± 0.3 ; $P < 0.01$). All mice in this experiment developed EAE, but no mice died of the disease. Splenic T cell numbers of RAG1 knockout recipient mice were determined 22 d after the disease onset. No significant differences were observed between mice that received WT or IKK α ^{AA} cells. Collectively, these results establish that IKK α kinase activity in CD4⁺ T cells is essential for the development of EAE. It is to be noted that in addition to IKK α expressed by T cells, IKK α expressed by non-T cells (hematopoietic and non-hematopoietic cells) may also play a role in EAE. However, this issue is difficult to address because of the following: (a) IKK α expressed by nonhematopoietic cells may indirectly affect EAE by controlling lymphoid organogenesis, in addition to its potential direct effects in the CNS; (b) the difference in EAE between bone marrow chimeras (Fig. 1 b) and their parental mice (Fig. 1 a) is small. Thus, in this study we focused on the roles of IKK α in T cells.

To measure the anti-MOG response of transferred T cells, *Rag1*^{-/-} mice were sacrificed 22 d after disease onset, and their splenocytes were cultured in the presence of MOG peptide (Fig. 2). Total splenic T cell and splenocyte numbers were not significant different between mice received WT and *Ikk α ^{AA}* cells. Splenocytes isolated from mice receiving WT T

cells proliferated vigorously in response to MOG peptide and produced high levels of IL-2, IL-4, IL-17A, and IFN- γ . Strikingly, splenocytes from mice reconstituted with *Ikk α ^{AA}* T cells exhibited significantly reduced proliferation ($P < 0.05$) and cytokine production ($P < 0.01$; Fig. 2). Similar reduction in MOG-induced responses was observed in *Ikk α ^{AA}* splenocytes isolated from the mice tested in Fig. 1 (a and b; and not depicted). To determine




Figure 2. Reduced cytokine production by *Ikk α ^{AA}* CD4⁺ T cells primed in vivo. WT or *Ikk α ^{AA}* CD4⁺ T cells, 10^7 cells/mouse, were transferred into *Rag1*^{-/-} recipients through tail vein ($n = 3$). EAE was induced by immunizing mice with MOG peptide 24 h later. Mice were sacrificed on day 22 after disease onset, and their splenocytes, 0.5×10^6 /well, were cultured in complete DME with or without 5–50 μ g of MOG peptide. [³H]thymidine was added 48 h after the initiation of the cell culture. Cells were harvested 18 h later, and [³H]thymidine incorporation was measured (a). (b–e) Cytokine concentrations were measured by ELISA 40 h after the initiation of the culture. Results shown are means \pm SD of triplicate cultures. The differences between the two groups are statistically significant for all cultures with MOG peptide ($P < 0.01$). CPM, count per minute. Data are representative of two independent experiments.

whether IKK α mutation affects the survival of CD4⁺ T cells, we performed flow cytometry analysis of blood samples collected from mice reconstituted with T cells before they were immunized for EAE. We observed no significant differences in the number and frequency of T cells between WT and *Ikk α ^{AA}* groups. Additionally, WT and *Ikk α ^{AA}* T cells, when cultured in the presence of plate-bound anti-CD3 and soluble anti-CD28, did not have significant differences in survival as determined by flow cytometry after staining the cells with Annexin V. In contrast, IKK β -deficient T cells had a significantly increased rate of death under the same culture condition (unpublished data).

IKK α kinase activity is required for Th17 responses

The markedly reduced anti-MOG response of *Ikk α ^{AA}* CD4⁺ T cells indicates that the IKK α kinase activity may be required for T cell activation and/or differentiation. To test this theory, we isolated CD4⁺ T cells from naive WT and *Ikk α ^{AA}* mice and measured their responses to anti-CD3 and anti-CD28 stimulation (Fig. 3). We found that *Ikk α ^{AA}* mutant CD4⁺ T cells proliferated to the same extent as WT CD4⁺ T cells after anti-CD3 or anti-CD3 plus anti-CD28 stimulation (Fig. 3 a). Furthermore, upon stimulation with anti-CD3 and anti-CD28, *Ikk α ^{AA}* CD4⁺ T cells produced normal levels of IL-2, IL-4, and IFN- γ as compared with WT cells; however, when stimulated with anti-CD3 alone, they produced moderately less IL-4 and IFN- γ (Fig. 3). In contrast, *Ikk α ^{AA}* CD4⁺

T cells produced significantly less IL-17A than WT cells upon stimulation with anti-CD3 or anti-CD3 plus anti-CD28 (Fig. 3 e). IL-17A, but not IL-2, mRNA expression in *Ikk α ^{AA}* CD4⁺ T cells was also significantly reduced (Fig. S1). The viability of T cells in the two groups was not significantly different, as determined by trypan blue and/or Annexin V staining. Similar results were obtained when unfractionated WT and *Ikk α ^{AA}* splenocytes were stimulated with anti-CD3 and anti-CD28 (unpublished data). These results are in contrast to the global defect of the *Ikk α ^{AA}* CD4⁺ T cells to MOG restimulation as shown in Fig. 2. Because T cells tested in those experiments were isolated from mice with different degrees of EAE, the effect of the disease on T cell responsiveness could not be excluded. *Ikk α ^{AA}* mice did not show general defects in T cell activation, and similar percentages of CD4⁺CD44⁺ T cells ($15.5 \pm 1.3\%$ vs. $16.3 \pm 1.7\%$) and CD4⁺CD62L^{low} T cells ($22.6 \pm 2.1\%$ vs. $24.6 \pm 1.6\%$) were detected in spleens of naive WT and *Ikk α ^{AA}* mice, respectively.

These results indicate that IKK α may selectively regulate Th17 cell differentiation. To test this possibility, we compared the ability of WT and *Ikk α ^{AA}* CD4⁺ T cells to differentiate into Th17 cells in vitro. Differentiated Th17 cells were identified by flow cytometry after intracellular staining of IL-17A. When cultured under Th0 conditions, very few Th17 cells were spontaneously generated, but $\sim 14\%$ of WT or *Ikk α ^{AA}* CD4⁺ T cells produced IFN- γ (Fig. 4 a). In contrast, when cultured under Th17-inducing conditions, $9.9 \pm 0.6\%$ of WT CD4⁺ T cells produced IL-17; but only $5.6 \pm 0.8\%$ of *Ikk α ^{AA}* T cells were IL-17⁺ ($P < 0.01$; Fig. 4 b). Therefore, IKK α kinase activity is required for the development of the Th17 response.

IKK α specifically regulates *Il17a* gene expression

To determine whether IKK α regulates Th17 response at the transcriptional level, we examined the expression of a panel of Th17-related genes by quantitative real-time PCR. *Il17a* expression in *Ikk α ^{AA}* CD4⁺ T cells was significantly reduced as compared with WT cells when cultured under either Th0 or




Figure 3. Selective defect in cytokine production of *Ikk α ^{AA}* CD4⁺ T cells activated in vitro. CD4⁺ T cells isolated from naive WT ($n = 5$) and *Ikk α ^{AA}* ($n = 5$) mice were cultured in 96-well plates coated with 1 μ g/ml anti-CD3 in the presence or absence of 2 μ g/ml soluble anti-CD28. Proliferation was determined by [³H]thymidine incorporation at 48 h (a), and cytokine production was measured by ELISA at 40 h (b–e). Data are presented as means \pm SD of triplicate cultures and are representative of two independent experiments. *, $P < 0.05$; **, $P < 0.001$. CD3, cultured with anti-CD3; CD28, cultured with anti-CD28.




Figure 4. Defective IL-17 production and Th17 differentiation of *Ikk α ^{AA}* CD4⁺ T cells. CD4⁺ T cells from WT ($n = 4$) and *Ikk α ^{AA}* ($n = 4$) mice were cultured under either neutral (Th0) or Th17-inducing condition as indicated. (a) 72 h later, IL-17- and/or IFN- γ -producing cells were measured by flow cytometry after intracellular staining of cytokines. (b) Quantification of data shown in a. (c–g) 24 h after the initiation of the culture, total RNA was isolated and mRNA levels of the Th17 lineage genes were assessed by real-time RT-PCR. The lowest expression level of each gene was set to 1. The experiments were repeated at least three times with similar results. *, $P < 0.05$; **, $P < 0.001$.

IKK α selectively binds to the *Il17a* promoter

IKK α has been previously reported to be present in the nuclei of several nonhematopoietic cell types, such as keratinocytes, squamous epithelial cells, and fibroblasts (Anest et al., 2003; Yamamoto et al., 2003; Fernández-Majada et al., 2007). In the nucleus, IKK α can regulate target gene expression through a variety of mechanisms, some of which correlate with promoter-associated histone H3 phosphorylation (Anest et al., 2003; Yamamoto et al., 2003). Interestingly, we found that IKK α was not only present in the cytoplasm, but also constitutively expressed at high levels in the nuclei of resting CD4⁺ T cells (Fig. 5 a). The level of nuclear IKK α was $\sim 30\%$ of the cytoplasmic IKK α , and was not regulated by anti-CD3 or anti-CD28 stimulation. To test whether IKK α

regulates *Il17a* expression by a direct association with its promoter, we performed chromatin immunoprecipitation (ChIP) with anti-IKK α (Fig. 5). IKK α did not bind to the *Il17a* promoter in resting WT CD4⁺ T cells, but in cells that had been cultured in the Th17 differentiation medium for 1–2 d, IKK α was readily detected on the *Il17a* promoter, coinciding with the appearance of the IL-17A mRNA (Fig. 5, b and c). The *Il17a* promoter was selectively targeted by IKK α , as IKK α was not detected on promoters of *Il21*, *Il17f*, and *Il2* (Fig. 5 c). In WT T cells, IKK α binding to the *Il17a* promoter correlated with its histone H3 phosphorylation (Fig. 5 d). In contrast, in *Ikk α ^{AA}* T cells, mutant IKK α also bound to the *Il17a* promoter under Th17 differentiation conditions, but histone H3 phosphorylation did not occur (Fig. 5 d). On the other hand, histone H3 on *Il17f* promoter also underwent phosphorylation at serine10 position, 1 and 2 d after cells were cultured in Th17 differentiation medium, but this event was

Th17 condition (Fig. 4 c and Fig. S2). In contrast, *Il17f* (Fig. 4 d) and *Il23R* (not depicted) expressions were not affected by the loss of IKK α kinase activity. *Il21* expression was only marginally reduced in *Ikk α ^{AA}* CD4⁺ T cells (Fig. 4 e). Additionally, IL-17A heterogeneous nuclear RNA (hnRNA) could be readily detected in activated WT CD4⁺ T cells; in contrast, a much reduced signal was detected in *IKK α ^{AA}* T cells under the same condition (Fig. S3). The Th17 differentiation signals elicited by TGF- β and IL-6 eventually converge onto the induction of two Th17 lineage-specific transcription factors, ROR α and ROR γ T (Ivanov et al., 2006; Dong, 2008). However, the expression of *Rora* and *Rorgt* mRNAs and ROR γ and ROR γ T proteins was not affected by the IKK α mutation (Fig. 4, f and g, and Fig. S4). These data indicate that the effect of IKK α on Th17 lineage is *Il17a* specific and independent of ROR α and ROR γ T.

regulates *Il17a* expression by a direct association with its promoter, we performed chromatin immunoprecipitation (ChIP) with anti-IKK α (Fig. 5). IKK α did not bind to the *Il17a* promoter in resting WT CD4⁺ T cells, but in cells that had been cultured in the Th17 differentiation medium for 1–2 d, IKK α was readily detected on the *Il17a* promoter, coinciding with the appearance of the IL-17A mRNA (Fig. 5, b and c). The *Il17a* promoter was selectively targeted by IKK α , as IKK α was not detected on promoters of *Il21*, *Il17f*, and *Il2* (Fig. 5 c). In WT T cells, IKK α binding to the *Il17a* promoter correlated with its histone H3 phosphorylation (Fig. 5 d). In contrast, in *Ikk α ^{AA}* T cells, mutant IKK α also bound to the *Il17a* promoter under Th17 differentiation conditions, but histone H3 phosphorylation did not occur (Fig. 5 d). On the other hand, histone H3 on *Il17f* promoter also underwent phosphorylation at serine10 position, 1 and 2 d after cells were cultured in Th17 differentiation medium, but this event was




Figure 5. Epigenetic regulation of the *IL17a* locus by IKKα. (a) Purified CD4⁺ T cells from WT ($n = 4$) and *Ikkα^{ΔΔ}* ($n = 4$) mice were stimulated with 10 μg/ml plate-bound anti-CD3 and 10 μg/ml plate-bound anti-CD28 for the indicated times. Nuclear extract was blotted with anti-IKKα (mol wt 85 kD), anti-HDAC1 (anti-histone deacetylase-1, mol wt 60 kD), and anti-β-actin (mol wt 42kD). (b) Purified CD4⁺ T cells from WT mice ($n = 4$) were cultured under Th17-inducing condition. Total RNA was prepared at the indicated times. IL-17A mRNA levels were assessed by real-time RT-PCR. The expression level at day 0 was set to 1. (c) Purified WT CD4⁺ T cells were cultured as in b. ChIP was performed using anti-IKKα or control Ig at days (D) 0, 1, and 2, as indicated. Precipitated DNA was analyzed by PCR using *IL17a*, *IL2*, and *IL21* promoter-specific primers that gave rise to products of 167bp, 256bp, and 150bp, respectively. (d) Purified CD4⁺ T cells from WT ($n = 4$) and *Ikkα^{ΔΔ}* ($n = 4$) mice were cultured under the Th17-inducing condition. ChIP was performed using anti-IKKα, anti-pH3Ser10 (serine 10 phosphorylated histone H3), or control Ig, at the indicated times. Precipitated DNA was analyzed by PCR using *IL17a*, *IL2*, and *IL17f* promoter-specific primers, as in c. Data are representative of three independent experiments.

IKKα-independent (Fig. 5 d). As expected, IKKα mutation selectively affected PolIII binding to *IL17*, but not to *IL22* (Fig. S5). These results indicate that IKKα kinase activity is selectively required for driving the *IL17a* locus into an active state marked by H3 serine 10 phosphorylation.

IKKα regulates Th17 response independent from NF-κB

In addition to directly acting on the *IL17a* promoter, IKKα may also regulate Th17 response through NF-κB-dependent mechanisms. To test this possibility, we first examined IκBα degradation in WT and *Ikkα^{ΔΔ}* T cells. In WT CD4⁺ T cells, upon anti-CD3 and anti-CD28 stimulation, IκBα was quickly degraded followed by a recovery caused by resynthesis of IκBα (Fig. 6 a, left). The early degradation of IκBα in *Ikkα^{ΔΔ}* CD4⁺ T cells was not affected, but IκBα did not return to the prestimulation level at later time points. Similar results were obtained when cells were stimulated with PMA plus ionomycin (unpublished data). To test whether IKKα regulates the *IκBα* promoter as previously reported (Anest et al., 2003; Yamamoto et al., 2003), we performed ChIP-PCR analyses using CD4⁺ T cells stimulated with anti-CD3 and anti-CD28. We found that IKKα was recruited to *IκBα* promoter in both WT and *Ikkα^{ΔΔ}* CD4⁺ T cells after stimulation (Fig. 6 c). However, H3 Ser10 phosphorylation on the *IκBα* promoter was detected only in WT, but not in mutant, T cells. These results indicate that IKKα kinase activity is required for IκBα resynthesis, but not for its stimulation-induced degradation in T cells.

Next, we measured nuclear translocation of NF-κB subunits in T cells stimulated with anti-CD3 and anti-CD28. RelA, RelB, c-Rel, and p50 all migrated into the nucleus of WT CD4⁺ T cells upon activation (Fig. 6 a). As expected (Bonizzi et al., 2004), the IKKα mutation did not affect the

nuclear translocation of RelA, c-Rel, or p50, but completely blocked nuclear localization of RelB (Fig. 6 a), indicating that IKKα selectively controls RelB activation. The lack of an effect of IKKα mutation on nuclear RelA activity was also confirmed by RelA ELISA (Fig. S6). As the key kinase of the noncanonical (alternative) NF-κB pathway, IKKα mediates the processing of p100 and, consequently, the generation of the RelB-p52 heterodimer (Bonizzi et al., 2004). Defective RelB nuclear translocation in *Ikkα^{ΔΔ}* CD4⁺ T cells may be caused by impaired IKKα-regulated p100 processing. We therefore examined p100 processing and p52 nuclear translocation in anti-CD3- and anti-CD28-activated T cells (Fig. 6 b). In WT T cells, p100 and p52 were up-regulated in both the cytosolic and nuclear fractions upon stimulation. In *Ikkα^{ΔΔ}* T cells, however, p100 was also up-regulated in the cytosol and nucleus, but p52 expression was reduced in the cytosol and was undetectable in the nucleus after anti-CD3 and anti-CD28 stimulation. These data confirm that IKKα kinase activity is important for activation-induced p100 processing and is required for p52 and RelB nuclear translocation in T cells. Similar results were obtained when WT and *Ikkα^{ΔΔ}* T cells were stimulated with PMA and ionomycin (unpublished data).

These results raise the question of whether IKKα regulates Th17 cell differentiation through p52-RelB heterodimers. To test this, we examined whether RelB was required for Th cell differentiation by culturing WT and *Relb^{-/-}* CD4⁺ T cells under Th1- or Th17-inducing conditions (Fig. 6 d).




Figure 6. IKK α may not regulate Th17 gene expression through NF- κ B. (a) Purified CD4⁺ T cells from WT ($n = 4$) and *Ikk α ^{ΔΔ}* ($n = 4$) mice were stimulated with 10 μ g/ml plate-bound anti-CD3 and 10 μ g/ml plate-bound anti-CD28 for the indicated times. Nuclear and cytosolic protein lysates were blotted with anti-RelA (mol wt 65 kD), anti-RelB (mol wt 68 kD), anti-c-Rel (mol wt 75 kD), anti-p50 (mol wt 50 kD), anti-I κ B α (mol wt 36 kD), anti- β -actin (mol wt 42 kD), or anti-HDAC1 (mol wt 60 kD). (b) Purified CD4⁺ T cells from WT ($n = 4$) and *Ikk α ^{ΔΔ}* ($n = 4$) mice were stimulated for the indicated times, as in a. Nuclear and cytosolic protein lysates were blotted with anti-p100/p52 (mol wt of p100 and p52 are 100 and 52 kD, respectively), anti- β -actin (mol wt 42 kD), and anti-HDAC1 (mol wt 60 kD). (c) Purified CD4⁺ T cells from WT ($n = 4$) and *Ikk α ^{ΔΔ}* ($n = 4$) mice were stimulated for the indicated times, as in a. ChIP was performed using anti-IKK α , anti-pH3Ser10, or control Ig. Precipitated DNA was analyzed by PCR using *I κ B α* promoter-specific primers that gave rise to a 230-bp product. (d) Purified CD4⁺ T cells from WT ($n = 4$) and *Relb*^{-/-} ($n = 4$) mice were cultured under the Th1- or Th17-inducing condition. 72 h later, IL-17A- and IFN- γ -producing cells were measured by flow cytometry. Results are representative of two independent experiments.

Consistent with previously published data (Corn et al., 2005), *Relb*^{-/-} cells showed a moderately reduced capacity to differentiate into IFN- γ -producing Th1 cells (53.8 vs. 41.2%). Interestingly, similar percentages of IL-17-producing Th17 cells were generated in *Relb*^{-/-} and WT cultures (Fig. 6 d). Collectively, these results suggest that IKK α regulates the Th17 response through NF- κ B-independent mechanisms.

DISCUSSION

A novel finding from the current study is that IKK α selectively regulates the Th17 cell response. As a result, *Ikk α ^{ΔΔ}* mutant mice are more resistant to MOG-induced EAE. Although IKK α is important for the activation of the noncanonical NF- κ B pathway in CD4⁺ T cells, it does not seem to regulate Th17 response through NF- κ B-dependent mechanisms because *Relb*^{-/-} CD4⁺ T cells did not show any defect in Th17 differentiation. Using the TESS program, we were not able to identify any putative NF- κ B-binding site in either the murine or human *Il17a* promoter. In a promoter-transactivating assay, we didn't detect any effect of p50, p52, p65, p100, RelB, or c-Rel expression construct on the *Il17a* promoter reporter, whereas the ROR γ T construct activated the reporter significantly in the same assay (unpublished data). Expression of two Th17 lineage-specific transcription

factors, ROR α and ROR γ T was not changed in *Ikk α ^{ΔΔ}* CD4⁺ T cells, indicating that the defect may be attributed to another mechanism. This theory is supported by the observation that IKK α is recruited to the *Il17a* promoter during Th17 differentiation and a specific defect in IL-17A expression in *Ikk α ^{ΔΔ}* CD4⁺ T cells. Furthermore, phosphorylation of histone H3 on the *Il17a* promoter, an activation marker that correlates with Th17 differentiation, was absent in *Ikk α ^{ΔΔ}* T cells. Therefore, our study suggests a novel nuclear function for IKK α in Th17 differentiation.

In this study, we showed that RelB activation after anti-CD3 plus anti-CD28 stimulation was blocked in *Ikk α ^{ΔΔ}* CD4⁺ T cells. Although it has been reported that RelB is required for maximal T cell activation (Corn et al., 2005), we did not observe any defect in *Ikk α ^{ΔΔ}* CD4⁺ T cell proliferation and Th1 and Th2 cytokine production after anti-CD3 and anti-CD28 stimulation; this may be caused by additional effects of the IKK α mutation, such as reduced I κ B α resynthesis, which may compensate for the loss of RelB function. However, when stimulated *in vitro* with MOG peptide, splenocytes isolated from *Ikk α ^{ΔΔ}* mice that had developed EAE showed significantly reduced responses (Fig. 2). There are two possible reasons for this discrepancy. First, IKK α may be required for T cell activation induced by weak, but not strong,

ligands. Second, and more likely, the decreased anti-MOG T cell response may be secondary to the effect of IKK α on EAE. The increased severity of EAE in WT mice may help to activate and expand more MOG-specific CD4⁺ T cells than in *Ikk α ^{4A}* mice. As a consequence, there are likely more MOG-responsive T cells in the WT splenocyte culture than in the *Ikk α ^{4A}* culture (Fig. 2).

It should be emphasized that IKK α -mediated IL-17 regulation may be only one of the mechanisms whereby IKK α controls EAE, as other cytokines including Th1 cytokines may also be affected by IKK α (Figs. 2 and 3). With regard to IL-17, the most significant IKK α effect appears to be on its mRNA transcription (Figs. 3 and 4). However, because IKK α does not appear to affect the IL-17 protein levels of individual Th17 cells, but increases the frequency of IL-17-producing cells, we propose that IKK α may selectively affect IL-17 mRNA expression in a subpopulation of Th17 cells, but not all Th17 cells. This could be related to the stage of Th17 cell differentiation, the phase of cell cycle, and the microenvironment surrounding the Th17 cells. This may explain why many *Ikk α ^{4A}* cells do not produce any IL-17 protein, although others make normal amounts (Fig. 4).

Two subunits of the IKK complex, IKK α and IKK β , possess kinase activities (Häcker and Karin, 2006). Despite high sequence similarities, it is now evident that the two molecules possess distinct functions (Häcker and Karin, 2006). In T cells, IKK β mediates NF- κ B activation through the canonical pathway (Schmidt-Supprian et al., 2003). *Ikk β ^{-/-}* CD4⁺ T cells show significantly less I κ B α degradation with severely delayed kinetics (unpublished data). In contrast, IKK α was not required for the activation of the canonical NF- κ B pathway in CD4⁺ T cells. Instead, IKK α is required for the activation of the noncanonical NF- κ B pathway through p100 processing (Fig. 6 b). In the cytosol, p100 can still be processed, albeit at a markedly reduced rate in *Ikk α ^{4A}* cells. However, p52 nuclear translocation was completely abolished in these cells.

Gene expression is regulated by epigenetic mechanisms involving histone modifications. Dong et al. showed that similar to loci specific to Th1 and Th2 cells, Th17 loci encoding IL-17A and IL-17F were regulated by chromatin remodeling events (Akimzhanov et al., 2007). Transcriptionally permissive histone modifications, such as histone H3 acetylation and Lys-4 tri-methylation, were observed at the *Il17a* promoter in Th17 cells (Akimzhanov et al., 2007). We showed that during Th17 differentiation, histone H3 on both *Il17a* and *Il17f* promoters was phosphorylated at Ser10, a known prerequisite for histone H3 acetylation. However, IKK α was found to be selectively required for histone H3 phosphorylation at the *Il17a* promoter, but not *Il17f* promoter.

Recently, several groups have reported nuclear expression of IKK subunits (Anest et al., 2003; Yamamoto et al., 2003; Fernández-Majada et al., 2007; Lubin and Sweatt, 2007). Unlike IKK β and IKK γ (NEMO), which are exclusively cytoplasmic, IKK α contains a nuclear localization sequence. Furthermore, IKK α was found to be recruited to NF- κ B-dependent

(e.g., *Il8*, *Il6*, and *Ikb α*) and NF- κ B-independent (e.g., *c-fos*) gene promoters, where its presence correlated with H3 phosphorylation, and thus chromatin activation. However, nuclear IKK α was also found to be recruited to genes (e.g., *Maspin*) that undergo active repression (Luo et al., 2007). Although IKK α was proposed to directly phosphorylate histone H3 at Ser10 (Anest et al., 2003; Yamamoto et al., 2003), its association with gene repression and silencing suggest that IKK α -induced enhancement of H3 phosphorylation can also be mediated by indirect mechanisms. For example, IKK α may be responsible for the recruitment of identified H3 kinases to the promoters it activates. In addition, nuclear IKK α has also been shown to mediate RelA/p65 turnover in macrophages through its phosphorylation of Ser536 (Lawrence et al., 2005). This effect seems to be cell-type specific because we found that IKK α was not required for RelA/p65 phosphorylation on Ser536 in CD4⁺ T cells (unpublished data). This may partially explain the different roles of IKK α in T cells as compared with macrophages. Thus, the highly specific effect of IKK α on Th17 lineage and the resistance of *Ikk α ^{4A}* mice to MOG-induced EAE implicate IKK α as a potential therapeutic target for Th17-mediated autoimmune disorders.

MATERIALS AND METHODS

Mice and cell transfer. *Ikk α ^{4A}* mutant C57BL/6 mice were previously described (Senfleben et al., 2001). To generate bone marrow chimeric mice, C57BL/6 recipient mice were irradiated with 2 doses of 500 rads each. They were then intravenously injected with 10⁷ bone marrow cells collected from either WT or *Ikk α ^{4A}* mice. 8 wk after bone marrow reconstitution, mice were immunized for the induction of EAE as described below. For T cell transfer, 10⁷ CD4⁺ T cells isolated from WT or *Ikk α ^{4A}* mice were injected into *Rag1^{-/-}* C57BL/6 mice through the tail vein. 24 h after the transfer, mice were immunized with MOG for EAE induction. All procedures were preapproved by the Institutional Animal Care and Use Committee (IACUC) of the University of Pennsylvania.

Cell isolation, cell culture, and reagents. Naive CD4⁺CD25⁻CD44^{low}CD62L⁺ T cells were isolated by FACS, whereas total CD4⁺ T cells were isolated by MACS. The purity of the naive T cells isolated by FACS was >99%, whereas that of T cells isolated by MACS was >95%. Cells were cultured in complete DME containing 10% FBS, 100 U/ml penicillin, 100 μ g/ml streptomycin, 30 μ M β -mercaptoethanol, 1 mM sodium pyruvate, 10 mM HEPES, and 1 \times nonessential amino acids. Anti-IKK α , anti-RelA, anti-RelB, and anti-c-Rel were purchased from Santa Cruz Biotechnology, Inc. Anti- β -actin was purchased from Sigma-Aldrich. Anti-CD3 and anti-CD28 were purchased from eBioscience. Anti-I κ B α , anti-pRelA536, anti-p100/p52, and anti-pSer10H3 were purchased from Cell Signaling Technology. MOG 38–50 peptide was synthesized by Invitrogen. Pertussis toxin was purchased from List Biological Laboratories, Inc. CFA was purchased from DIFCO laboratories.

EAE induction and evaluation. EAE was induced as previously described (Hilliard et al., 2002). In brief, mice first received a subcutaneous immunization with 300 μ g MOG38–50 peptide emulsified in CFA and an intravenous injection of 200 ng pertussis toxin. A second injection of 200 ng pertussis toxin was given 48 h later. Mice were examined daily for clinical signs of EAE and scored as follows: 0, no disease; 1, tail paralysis; 2, hind limb weakness; 3, hind limb paralysis; 4, hind limb plus forelimb paralysis; 5, moribund or dead.

Th cell differentiation. CD4⁺ T cells were isolated using autoMACS automatic cell sorter (Miltenyi Biotec). Cells were cultured with 50 U/ml IL-2,

2 $\mu\text{g/ml}$ anti-CD28, and 2 $\mu\text{g/ml}$ anti-CD3 under either Th0- (no more addition of cytokines or antibodies), Th1- (10 ng/ml IL-12 and 20 ng/ml anti-IL-4), or Th17- (0.4 $\mu\text{g/ml}$ anti-IL-4, 10 $\mu\text{g/ml}$ anti-IFN- γ , 20 ng/ml IL-6, and 5 ng/ml TGF- β 1) inducing conditions. 5 d later, cells were washed and restimulated with anti-CD3 and anti-CD28 for 12 h, stained with anti-IL-17A and anti-IFN- γ , and examined by flow cytometry.

Real-time PCR. Total RNA was extracted using TRIzol reagent (Invitrogen). cDNA was synthesized using reverse transcription II (Invitrogen). Real-time quantitative PCR was performed using the SYBR Green master mix (Applied Biosystems). Data were normalized to β -actin mRNA. Primers used are as follows (forward and reverse): IL-17A, 5'-ACCTCACACGAGGCACAAGT-3' and 5'-CTTCATTGCGGTGGATC-3'; IL-17F, 5'-GAGGATAACACTGTGAGAGTTGAC-3' and 5'-GAGTTCATGGT-GCTGTCTTCC-3'; IL-21, 5'-TCATCATTGACCTCGTGGCC-3' and 5'-ATCGTACTTCTCCACTTGAATCCC-3'; ROR α , 5'-TCTCCCT-GCGCTCTCCGCAC-3' and 5'-TCCACAGATCTTGCATGGA-3'; ROR γ T, 5'-CCGCTGAGAGGGCTTAC-3' and 5'-TGCAGGAGTA-GGCCACATTACA-3'; IL-23R, 5'-GCCAAGAAGACCATTCCCGA-3' and 5'-TCAGTGCTACAACTTCTTCAGAGGACA-3'; p19, 5'-AAGT-TCTCTCCTTCCCTGTCGC-3' and 5'-TCTTGTGGAGCAGCA-GATGTGAG-3'; β -actin, 5'-GTGGGCCGCTCTAGGCACCAA-3' and 5'-CTCTTTGATGTCACGCACGATTTC-3'.

ChIP PCR. ChIP was performed using a Chromatin Immunoprecipitation kit (Millipore) according to the manufacturer's instructions. In brief, cells were first treated with formaldehyde and sonicated to break up the chromatin. Sonicated chromatin preparations were immunoprecipitated with specific antibodies. DNA was then eluted after extensive washing. PCR was performed using the following primers (forward and reverse): IL-17A promoter, 5'-GCAGCAGCTCAGATATGTCC-3' and 5'-TGAGGTCA-GCACAGAACCAC-3'; IL-17F promoter, 5'-CATGTGAATGGCA-CGATAGG-3' and 5'-TAATCCCCCACAAAGCAAC-3'; IL-2 promoter, 5'-CATAAGAAAGCGTTCATTG-3' and 5'-TACCTGTGTGGCAG-AAAGC-3'. A small portion of sonicated chromatin preparations was also amplified by PCR to ensure equal input.

ELISA. Antibodies used in ELISA were purchased from BD, including purified and biotinylated rat anti-mouse IL-2, IL-4, IL-6, IFN- γ , and IL-17A. Quantitative ELISA was performed using paired mAbs specific for corresponding cytokines according to the manufacturer's recommendations.

Western blot. Nuclear protein extract was prepared using a Nuclear Extract kit (Active Motif Inc.) per manufacturer's instructions. Total cell lysate was prepared by suspending cells in the RIPA buffer containing 150 mM NaCl, 10 mM Tris, pH 7.4, 0.1% SDS, 1% Triton X-100, 1% sodium deoxycholate, 100 μM Na₃VO₄, 5 mM EDTA, 1 mM PMSE and protease inhibitors cocktail. Proteins (20 μg per lane) were resolved by SDS-PAGE, transferred to a nitrocellulose membrane, and blotted with specific antibodies.

Statistical analyses. Disease severity, day of onset, and cytokine concentrations were analyzed by Student's *t* test. Disease scores were analyzed by Mann-Whitney *U* test.

Online supplemental material. Fig. S1 shows reduced IL-17A mRNA expression in *Ikk α ^{4/4}* CD4⁺ T cells. Fig. S2 shows defective IL-17A expression in *Ikk α ^{4/4}* CD4⁺ T cells during Th17 differentiation. Fig. S3 shows reduced IL-17A hnRNA expression in *Ikk α ^{4/4}* CD4⁺ T cells. Fig. S4 shows normal ROR γ /ROR γ T protein expression in *Ikk α ^{4/4}* CD4⁺ T cells. Fig. S5 shows the binding of RNA polymerase II to the *Il17*, *Il21*, and *Il22* promoters. Fig. S6 shows nuclear RelA/p65 activities as determined by ELISA. Online supplemental material is available at <http://www.jem.org/cgi/content/full/jem.20091346/DC1>.

The authors thank Jing Sun, Ruaidhri Carmody, and Christopher Hunter for their valuable discussions and reagents.

This work was supported by grants from the National Institutes of Health (AI50059, DK070691, and AI069289).

The authors have no competing financial interests related to this work.

Submitted: 19 June 2009

Accepted: 11 February 2011

REFERENCES

- Akimzhanov, A.M., X.O. Yang, and C. Dong. 2007. Chromatin remodeling of interleukin-17 (IL-17)-IL-17F cytokine gene locus during inflammatory helper T cell differentiation. *J. Biol. Chem.* 282:5969–5972. doi:10.1074/jbc.C600322200
- Anest, V., J.L. Hanson, P.C. Cogswell, K.A. Steinbrecher, B.D. Strahl, and A.S. Baldwin. 2003. A nucleosomal function for I κ B kinase- α in NF- κ B-dependent gene expression. *Nature*. 423:659–663. doi:10.1038/nature01648
- Bettelli, E., Y. Carrier, W. Gao, T. Korn, T.B. Strom, M. Oukka, H.L. Weiner, and V.K. Kuchroo. 2006. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature*. 441:235–238. doi:10.1038/nature04753
- Bettelli, E., T. Korn, and V.K. Kuchroo. 2007. Th17: the third member of the effector T cell trilogy. *Curr. Opin. Immunol.* 19:652–657. doi:10.1016/j.coi.2007.07.020
- Bonizzi, G., M. Bebién, D.C. Otero, K.E. Johnson-Vroom, Y. Cao, D. Vu, A.G. Jegga, B.J. Aronow, G. Ghosh, R.C. Rickert, and M. Karin. 2004. Activation of IKK α target genes depends on recognition of specific κ B binding sites by RelB:p52 dimers. *EMBO J.* 23:4202–4210. doi:10.1038/sj.emboj.7600391
- Corn, R.A., C. Hunter, H.C. Liou, U. Siebenlist, and M.R. Boothby. 2005. Opposing roles for RelB and Bcl-3 in regulation of T-box expressed in T cells, GATA-3, and Th effector differentiation. *J. Immunol.* 175:2102–2110.
- Cua, D.J., J. Sherlock, Y. Chen, C.A. Murphy, B. Joyce, B. Seymour, L. Lucian, W. To, S. Kwan, T. Churakova, et al. 2003. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature*. 421:744–748. doi:10.1038/nature01355
- Dong, C. 2008. TH17 cells in development: an updated view of their molecular identity and genetic programming. *Nat. Rev. Immunol.* 8:337–348. doi:10.1038/nri2295
- Fernández-Majada, V., C. Aguilera, A. Villanueva, F. Vilardell, A. Robert-Moreno, A. Aytés, F.X. Real, G. Capella, M.W. Mayo, L. Espinosa, and A. Bigas. 2007. Nuclear IKK activity leads to dysregulated notch-dependent gene expression in colorectal cancer. *Proc. Natl. Acad. Sci. USA.* 104:276–281. doi:10.1073/pnas.0606476104
- Häcker, H., and M. Karin. 2006. Regulation and function of IKK and IKK-related kinases. *Sci. STKE*. 2006:re13. doi:10.1126/stke.3572006re13
- Hilliard, B.A., N. Mason, L. Xu, J. Sun, S.-E. Lamhamed-Cherradi, H.-C. Liou, C. Hunter, and Y.H. Chen. 2002. Critical roles of c-Rel in autoimmune inflammation and helper T cell differentiation. *J. Clin. Invest.* 110:843–850.
- Hu, Y., V. Baud, T. Oga, K.I. Kim, K. Yoshida, and M. Karin. 2001. IKK α controls formation of the epidermis independently of NF- κ B. *Nature*. 410:710–714. doi:10.1038/35070605
- Ivanov, I.I., B.S. McKenzie, L. Zhou, C.E. Tadokoro, A. Lepelletier, J.J. Lafaille, D.J. Cua, and D.R. Littman. 2006. The orphan nuclear receptor ROR γ directs the differentiation program of proinflammatory IL-17⁺ T helper cells. *Cell*. 126:1121–1133. doi:10.1016/j.cell.2006.07.035
- Langrish, C.L., Y. Chen, W.M. Blumenschein, J. Mattson, B. Basham, J.D. Sedgwick, T. McClanahan, R.A. Kastelein, and D.J. Cua. 2005. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J. Exp. Med.* 201:233–240. doi:10.1084/jem.20041257
- Lawrence, T., M. Bebién, G.Y. Liu, V. Nizet, and M. Karin. 2005. IKK α limits macrophage NF- κ B activation and contributes to the resolution of inflammation. *Nature*. 434:1138–1143. doi:10.1038/nature03491
- Lubin, F.D., and J.D. Sweatt. 2007. The I κ B kinase regulates chromatin structure during reconsolidation of conditioned fear memories. *Neuron*. 55:942–957. doi:10.1016/j.neuron.2007.07.039
- Luo, J.L., W. Tan, J.M. Ricono, O. Korchynskiy, M. Zhang, S.L. Gonias, D.A. Cheresch, and M. Karin. 2007. Nuclear cytokine-activated IKK α

- controls prostate cancer metastasis by repressing Maspin. *Nature*. 446: 690–694. doi:10.1038/nature05656
- Romagnani, S. 1997. The Th1/Th2 paradigm. *Immunol. Today*. 18:263–266. doi:10.1016/S0167-5699(97)80019-9
- Schmidt-Supprian, M., G. Courtois, J. Tian, A.J. Coyle, A. Israël, K. Rajewsky, and M. Pasparakis. 2003. Mature T cells depend on signaling through the IKK complex. *Immunity*. 19:377–389. doi:10.1016/S1074-7613(03)00237-1
- Senftleben, U., Y. Cao, G. Xiao, F.R. Greten, G. Krähn, G. Bonizzi, Y. Chen, Y. Hu, A. Fong, S.C. Sun, and M. Karin. 2001. Activation by IKK α of a second, evolutionary conserved, NF- κ B signaling pathway. *Science*. 293:1495–1499. doi:10.1126/science.1062677
- Sil, A.K., S. Maeda, Y. Sano, D.R. Roop, and M. Karin. 2004. IkappaB kinase- α acts in the epidermis to control skeletal and craniofacial morphogenesis. *Nature*. 428:660–664. doi:10.1038/nature02421
- Sutton, C., C. Brereton, B. Keogh, K.H. Mills, and E.C. Lavelle. 2006. A crucial role for interleukin (IL)-1 in the induction of IL-17-producing T cells that mediate autoimmune encephalomyelitis. *J. Exp. Med.* 203:1685–1691. doi:10.1084/jem.20060285
- Veldhoen, M., and B. Stockinger. 2006. TGF β 1, a “Jack of all trades”: the link with pro-inflammatory IL-17-producing T cells. *Trends Immunol.* 27:358–361. doi:10.1016/j.it.2006.06.001
- Weaver, C.T., L.E. Harrington, P.R. Mangan, M. Gavrieli, and K.M. Murphy. 2006. Th17: an effector CD4 T cell lineage with regulatory T cell ties. *Immunity*. 24:677–688. doi:10.1016/j.immuni.2006.06.002
- Yamamoto, Y., U.N. Verma, S. Prajapati, Y.T. Kwak, and R.B. Gaynor. 2003. Histone H3 phosphorylation by IKK- α is critical for cytokine-induced gene expression. *Nature*. 423:655–659. doi:10.1038/nature01576
- Yang, X.O., A.D. Panopoulos, R. Nurieva, S.H. Chang, D. Wang, S.S. Watowich, and C. Dong. 2007. STAT3 regulates cytokine-mediated generation of inflammatory helper T cells. *J. Biol. Chem.* 282:9358–9363. doi:10.1074/jbc.C600321200
- Yang, X.O., B.P. Pappu, R. Nurieva, A. Akimzhanov, H.S. Kang, Y. Chung, L. Ma, B. Shah, A.D. Panopoulos, K.S. Schluns, et al. 2008. T helper 17 lineage differentiation is programmed by orphan nuclear receptors ROR α and ROR γ . *Immunity*. 28:29–39. doi:10.1016/j.immuni.2007.11.016