

RESEARCH ARTICLE

Open Access

Transcriptome analysis reveals *in vitro* cultured *Withania somnifera* leaf and root tissues as a promising source for targeted withanolide biosynthesis

Kalaiselvi Senthil¹, Murukarthick Jayakodi^{2†}, Pankajavalli Thirugnanasambantham^{1†}, Sang Choon Lee², Pradeepa Duraisamy¹, Preethi M Purushotham¹, Kalaiselvi Rajasekaran¹, Shobana Nancy Charles¹, Irene Mariam Roy¹, Arul Kumar Nagappan³, Gon Sup Kim³, Yun Sun Lee², Senthil Natesan⁴, Tae-Sun Min⁵ and Tae Jin Yang^{2*}

Abstract

Background: The production of metabolites via *in vitro* culture is promoted by the availability of fully defined metabolic pathways. Withanolides, the major bioactive phytochemicals of *Withania somnifera*, have been well studied for their pharmacological activities. However, only a few attempts have been made to identify key candidate genes involved in withanolide biosynthesis. Understanding the steps involved in withanolide biosynthesis is essential for metabolic engineering of this plant to increase withanolide production.

Results: Transcriptome sequencing was performed on *in vitro* adventitious root and leaf tissues using the Illumina platform. We obtained a total of 177,156 assembled transcripts with an average unigene length of 1,033 bp. About 13% of the transcripts were unique to *in vitro* adventitious roots but no unique transcripts were observed in *in vitro*-grown leaves. A putative withanolide biosynthetic pathway was deduced by mapping the assembled transcripts to the KEGG database, and the expression of candidate withanolide biosynthesis genes -were validated by qRT PCR. The accumulation pattern of withaferin A and withanolide A varied according to the type of tissue and the culture period. Further, we demonstrated that *in vitro* leaf extracts exhibit anticancer activity against human gastric adenocarcinoma cell lines at sub G1 phase.

Conclusions: We report here a validated large-scale transcriptome data set and the potential biological activity of *in vitro* cultures of *W. somnifera*. This study provides important information to enhance tissue-specific expression and accumulation of secondary metabolites, paving the way for industrialization of *in vitro* cultures of *W. somnifera*.

Keywords: Withanolides, Transcriptome, Illumina platform, Hiseq, *in vitro* adventitious roots

* Correspondence: tjyang@snu.ac.kr

†Equal contributors

²Department of Plant Science, Plant Genomics and Breeding Institute, Research Institute for Agriculture and Life Sciences, College of Agriculture and Life Sciences, Seoul National University, Seoul 151-921, Republic of Korea
Full list of author information is available at the end of the article

Background

Withania somnifera (L.) Dunal (Family, Solanaceae) commonly known as Ashwagandha or winter cherry is one of the top medicinal herbs used in ayurveda – the holistic system of Indian medicine. Most of its therapeutic properties are comparable to those of ginseng, and hence *W. somnifera* is also known as Indian ginseng. *W. somnifera* extract in various forms has been used as an adaptogen, aphrodisiac, liver tonic, anti-inflammatory agent, hepatoprotectant, astringent, immune booster [1], immunomodulator [2] and more recently to treat Alzheimer's disease [3], neurodegenerative disorders and stress [4] and also as an adjunct to chemotherapy and/or radiation therapy [5,6]. Wide variations in the type and content of secondary metabolites have been reported within chemotypes and parts of the *W. somnifera* plant [7,8]. The principle bioactive components, a group of secondary metabolites collectively called withanolides, are C-28 steroidal lactones [9]. Withanolides are biosynthesised through the isoprenoid pathway, probably via both the mevalonate and non-mevalonate pathways [8] wherein 24-methylene cholesterol is the first branching point towards the biosynthesis of different withanolides through a series of desaturation, hydroxylation, epoxidation, cyclization, chain elongation, and glycosylation steps [10,11].

W. somnifera has been identified by the National Medicinal Plant Board of India as one of the thirty-two priority medicinal plants that are in great demand in the domestic and international markets [12]. According to one estimate, the demand for withanolide production from dried plant material is 12,120 tonnes [13], whereas the annual production is 5,905 tonnes [14]. To meet the growing demand of the herbal industry, *in vitro* cultures could be an alternative to field-grown plants for consistent production of secondary metabolites within a short period [15]. Several reports indicate the potential of manipulating *in vitro* tissues to produce bioactive terpenoids, phenolics, saponins and anthocyanins [16–18], but any attempt to engineer efficient production of secondary metabolites requires understanding of their biosynthetic pathway(s), and our present knowledge of withanolide biosynthesis is limited to only a few genes involved in the pathway [10,19,20].

Next-generation sequencing (NGS) technology for transcriptomes (RNA-seq) provides a new approach for both obtaining gene sequences and quantifying transcriptomes of any organism. In recent years, RNA-seq has been a powerful method for identifying genes involved in important secondary metabolite pathways such as biosynthesis of ginsenosides in *Panax ginseng* [21–24], carotenoids in *Momordica cochinchinensis* [25], flavonoids, theanine and caffeine in tea (*Camellia sinensis*) [26], flavonoids in safflower (*Carthamus tinctorius* L.) [27], capsaicinoid in chili pepper (*Capsicum frutescens*) [28] and flavor components in radish (*Raphanus sativus* L.) [29]. Initial efforts have

been made to generate expressed sequence tags (ESTs) from *in vitro* tissues of *W. somnifera* [30]. In addition, leaf and root transcriptomes of field-grown tissues have been analysed to identify genes involved in withanolide biosynthesis [20], and very recently, analysis of the expression of pathogenesis-related genes has been carried out in leaf transcriptomes of *W. somnifera* [31]. However, no large-scale transcriptome information validated by expression profiling is available for *in vitro* tissues of *W. somnifera*.

Here, we used RNA-seq for large-scale transcriptome profiling and generated a comprehensive transcriptome for *W. somnifera* by assembling the transcriptomes of *in vitro* adventitious root and leaf tissues from the millions of short sequence reads generated by Illumina. Annotations including functional annotation, Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways - are also reported. In addition, we showed that *in vitro* tissue extracts have antioxidant activity and cytotoxicity against human adenocarcinoma gastric cancer (AGS) cell lines. Furthermore, we have developed a web database in which to access the *W. somnifera* transcriptome. Overall, this work represents the first large-scale transcriptome profiling of *in vitro* tissues for *W. somnifera* and provides comparative expression profiling of pathway genes involved in withanolide biosynthesis and their potential biological activity.

Results

Induction and maintenance of *in vitro* cultures

Direct adventitious roots were induced successfully from leaf explants of *W. somnifera*. Further induced roots were maintained in MS liquid medium in a bubble column bioreactor (Figure 1). At 45 days, adventitious roots showed maximal growth with increased branching. The roots were healthy and creamy white in colour. By 60 days, the roots turned brown and brittle. Leaf tissues from multiple shoots maintained *in vitro* were used for isolation of total RNA.

Illumina sequencing and *de novo* assembly

To establish a comprehensive overview of the *W. somnifera* transcriptome, we prepared RNA-seq libraries from total RNA isolated from 60-day-old *in vitro* cultures of adventitious root and leaf tissues and subjected them to paired-end (PE) sequencing with the Illumina platform. We obtained 135,186,223 and 113,849,837 cleaned high-quality reads from the *in vitro* adventitious root and leaf tissues, respectively (Table 1). We combined both sets of reads and assembled them using Trinity [32] to generate a unique transcript library. After read depth analysis, we obtained a total of 177,156 transcripts (including isoforms) with an average length of 1,033 bp (Table 1) that we used for subsequent analyses and also prepared a set




Figure 1 Adventitious root induction and proliferation in bioreactor. **a** -Induction of adventitious roots from the leaf explants of *W. somnifera*, **b** Propagation stage after 30 days of culture, **c** Mass production of root after 45 days in bioreactor.

of 105,662 non-redundant (Nr) transcripts by selecting only the longest sequence among isoforms.

Differentially expressed transcripts between *in vitro*-grown roots and leaves

High-quality reads from *in vitro* adventitious root and leaf samples were separately mapped to the assembled transcripts and the number of reads mapped was normalized by the FPKM method using RSEM [33]. We classified gene expression into four categories (low, moderate, high and very high), based on the FPKM values for each transcript in the respective tissue samples (Figure 2). The largest fraction of transcripts showed low expression (FPKM 0–2) in both tissue samples. A small fraction (14%–19%) was expressed moderately in both tissues and very few (1%–2%) had high expression (Figure 2). In addition, we identified transcripts expressed specifically in each tissue sample using tissue-by-tissue comparison between root and leaf. For this, we normalized the number of reads mapped to each transcript per million (TPM) mapped for root and leaf tissues and considered transcripts to be expressed specifically in a single tissue (root or leaf) if they showed at least 3 TPM in the tissue of interest and zero in the other. Among the total 177,156 transcripts analysed in this study, a large number (152,839) were present in both tissues and a total of 24,317 were preferentially expressed in root. By contrast, we did not find any transcripts expressed exclusively in leaf. To identify differentially-expressed transcripts, the edgeR bioconductor package was used with a threshold of greater than 2-fold change and a significant false discovery rate (FDR) value of ≤ 0.001 [34]. We found a total of 12,822 transcripts with different expression between root and leaf (Additional file 1: Figure S1). Among them, 8,013 transcripts were up-regulated and 4,809 transcripts were down-regulated.

Functional annotation of the assembled transcripts


For validation and annotation of assembled transcripts, sequence similarity searches were conducted against protein databases. BLAST searches indicated that 80,184 (45%) and 70,840 (40%) out of 177,156 transcripts showed significant similarity to known proteins in NCBI Nr and TAIR databases. As expected, the number of annotations we could assign increased when we annotated based on the closely related *Solanaceae* species of tomato and potato. Of all the transcripts, 83,204 (47%) had BLAST hits in tomato protein database, and 84,184 (47%) transcripts significantly matched proteins in the potato database.

Further, gene ontology (GO) annotation [35] was carried out BLAST2GO [35] and a total of 57,404 transcripts (32.40% of all the assembled transcripts) were assigned at least one GO term. Among the GO classifications, assignments to the cellular component class ranked highest (53,140), followed by biological process (50,786) and molecular function (44,018). Within the molecular function category, transcripts assigned to protein binding (11,436) and ATP binding (4,583) processes were the most common (Figure 3). Within the biological process category, the majority of the GO terms were assigned to response to salt stress (4,067) and response to cadmium ion (3,120). For cellular components, the assignments were mostly nucleus (15,104) and plasma membrane (13,613; Figure 3).

To identify the biological functions and interactions of genes, we used KEGG pathway [36] annotation. A total of 24,515 transcripts were assigned to 139 KEGG pathways. As shown in Figure 4, the KEGG metabolic pathways assigned most transcripts were involved in purine, starch and sucrose metabolism. Of the transcripts assigned to secondary metabolite biosynthetic pathways, a large pool with 505 members was mapped to phenylpropanoid biosynthesis (Table 2), with monoterpenoid biosynthesis

Table 1 Summary of *W. somnifera* transcriptome sequencing, assembly and annotation

Tissues	No. of raw reads	No. of trimmed reads	No. of assembled transcripts	Nr set (without isoforms)	Average transcript length (bp)	Longest transcript length (bp)
<i>in vitro</i> root	139,562,334	135,186,223	177,156	105,662	1,033	10,802
<i>in vitro</i> leaf	117,948,186	113,849,837				




(174) and terpenoid backbone biosynthesis (155) also including many transcripts. The overall annotation of the *W. somnifera* transcriptome provides a valuable resource for investigating specific processes, functional descriptions and pathways.

We next identified differentially expressed transcripts involved in the biosynthesis of major secondary metabolites (Table 2). Differentially expressed transcripts were observed for most of the pathways analysed, which was perhaps not surprising given the large number of transcripts that were unique to *in vitro* root tissues. We mapped the 24,317 *in vitro* root-specific transcripts to KEGG pathways and observed that only 3% of the transcripts were assigned to established pathways whereas the others were of unknown function. Interestingly, 13 transcripts for beta-amyrin synthase appear to be specific to

in vitro-grown roots as they were not found among field-grown root transcripts [20].

Expression of major withanolide biosynthesis genes in adventitious root and leaf tissues at different culture periods

To establish a complete expression profile for genes in the withanolide biosynthesis pathway, we extracted transcripts encoding enzymes involved in withanolide biosynthesis, including FPPS, SE, CAS, DXPS, GT, and HMGR (Table 3) from our assembly. We found expression of most transcripts in both types of *in vitro* tissues, along with some *in vitro* root-unique transcripts (Table 3). There were many transcripts for methyltransferase enzymes and very few for enzymes such as CECL, MECDPS, HMBPR, HMBPS and MVD (Table 3). Overall, our data clearly






Figure 4 KEGG pathway annotation of assembled *in vitro* tissue transcripts. Pathway assignment was summarized for two main categories: metabolism and secondary metabolite categories. A total of 24,515 transcripts were assigned to primary and secondary metabolic pathways.

showed that genes encoding putative enzymes involved in withanolide biosynthesis were expressed in both *in vitro* root and leaf tissues. For experimental confirmation, we chose major candidate genes *HMGR*, *FPPS*, *SE*, *CAS* and *GT* and performed qRT-PCR analysis (Figure 5). *FPPS*, *SE* and *CAS* genes exhibited higher expression at 45 days of growth in both leaf and root tissue. *HMGR* expression was high in 30-day-old *in vitro* adventitious root cultures, but declined sharply by the 45th day to levels observed in leaf tissues. In leaf tissue, the expression of *HMGR* was much lower, with maximal expression at 45 days of culture. *CAS* and *GT* shared similar expression levels in leaf tissues. In root tissue, *GT* levels were high until 60 days of culture, whereas in leaf tissue, *GT* exhibited similar expression during 30 and 45 days of growth but declined at 60 days. The expression of *SE* and *GT* seemed to follow a similar pattern with relatively higher expression in roots compared to leaves at all three time periods analysed, whereas *FPPS* and *CAS* shared a similar expression profile, with comparatively higher expression in leaves. The correlation matrix revealed significant linear correlation between the expression patterns of *FPPS* and *SE*, and the accumulation of major withanolides. *FPPS* and *SE* exhibited significant correlation ($r = 0.56$; $p = 0.01$) and ($r = 0.75$) respectively with the accumulation of withanolide A in 45-day-old root culture. In 45-day-old leaf culture, significant linear correlation ($r = 0.22$; $p = 0.01$) and a strong positive correlation ($r = 0.8$) were displayed by *FPPS* and *SE*, respectively, towards the accumulation of withaferin A. Among 60 day old root culture, significant correlation was exhibited by *SE* ($r = 0.602$) and *FPPS* ($r = 0.699$) towards the accumulation of withanolide A whereas, among 60 day old leaf culture, *FPPS* exhibited strong positive correlation ($r = 0.78$) followed by *SE*

($r = 0.73$) towards the accumulation of withaferin A. This shows that, though the roots turn brown by 60 days of culture, there is active metabolism taking place. Overall, the expression of the 5 tested genes in all time periods were comparable to the digital expression patterns observed in 60-day-old *in vitro* root and leaf tissue sequencing data based on FPKM values (Additional file 2: Figure S2).

Quantification of major withanolides in *in vitro* leaf and root tissues

The major secondary metabolites of *W. somnifera*, namely withanolide A and withaferin A, were quantified by high performance thin layer chromatography [37] (HPTLC; Figure 6). The contents of both components were significantly different in the two tissues tested (Additional file 3: Figure S3). Leaf tissues preferentially accumulated withaferin A whereas *in vitro* adventitious root tissues accumulated prodigious quantities of withanolide A. Significant differences were also observed in the accumulation patterns over time. *In vitro* adventitious roots contained high amounts of withanolide A (380 ± 0.36 $\mu\text{g/g}$ DW) at 45 days, but less at 60 days of culture. Similarly, withaferin A accumulation (980 ± 0.97 $\mu\text{g/g}$ DW) in leaf tissue was highest at 45 days and reduced on extended culturing period.

Antioxidant activity of extracts of *in vitro* leaf and root tissues

Complex antioxidant systems are essential for protecting cellular membranes and organelles from the damaging effects of reactive oxygen species (ROS), and plants have developed a series of enzymatic and non-enzymatic detoxification systems to counteract ROS [38]. The enzymatic and non-enzymatic antioxidant activities of the

Table 2 Differential expression of transcripts involved in major secondary metabolite biosynthetic pathways

Pathway	Pathway ID in KEGG DB	No. of transcripts assigned to pathway	Differentially expressed transcripts (more than 2 fold difference)	No. of unique transcripts	
				Adventitious roots	Leaves
Phenylpropanoid biosynthesis	map00940	505	185	104	0
Phenylalanine, tyrosine and tryptophan biosynthesis	map00400	209	49	15	0
Monoterpenoid biosynthesis	map00902	174	50	23	0
Terpenoid backbone biosynthesis	map00900	155	28	22	0
Pantothenate and CoA biosynthesis	map00770	141	52	16	0
Flavone and flavonol biosynthesis	map00944	129	57	22	0
Carotenoid biosynthesis	map00906	127	47	16	0
Zeatin biosynthesis	map00908	127	43	26	0
Flavonoid biosynthesis	map00944	121	49	15	0
Ubiquinone and other terpenoid-quinone biosynthesis	map00130	103	35	13	1
Isoquinoline alkaloid biosynthesis	map00950	101	34	11	0
Tropane, piperidine and pyridine alkaloid biosynthesis	map00960	100	35	12	0
Steroid hormone biosynthesis	map00140	90	33	16	0
Various types of N-glycan biosynthesis	map00513	73	10	6	0
Steroid biosynthesis	map00100	70	11	17	0
N-Glycan biosynthesis	map00513	61	10	5	0
Folate biosynthesis	map00790	59	11	7	0
Sesquiterpenoid and triterpenoid biosynthesis	map00909	45	15	4	0
Glucosinolate biosynthesis	map00966	43	18	11	0
Diterpenoid biosynthesis	map00904	38	8	14	0
Lipopolysaccharide biosynthesis	map00540	31	8	4	0
Phosphonate and phosphinate metabolism	map00440	28	4	6	0
Other types of O-glycan biosynthesis	map00514	11	0	1	0
Isoflavonoid biosynthesis	map00943	7	5	2	0
Anthocyanin biosynthesis	map00942	6	4	1	0

cultured *W. somnifera* tissues is represented in Table 4. Though both tissues were grown under controlled conditions, the antioxidant potentials differed between extracts of *in vitro* adventitious root (root extract) and leaf (leaf extract). The antioxidant capacity, as estimated from total antioxidants, of root extracts was much higher than that of leaf extracts (Table 4).

Inhibitory effect of methanolic extract of *in vitro* *W. somnifera* leaf tissue on cell proliferation

An initial cytotoxicity study based on the survival of adenocarcinoma gastric cancer (AGS) cell lines was performed using an MTT (3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide) assay. Cells were exposed to various concentrations (0–200 µg/mL) of methanolic extract of *in vitro*-grown leaf tissues (leaf extract) and the cell viability was determined. Cytotoxicity was

quantified as the percentage of viable treated cells relative to the viable cells in an untreated control. The half maximal inhibitory concentration (IC₅₀) for leaf extract was 118.85 µg/mL (Figure 7a). To determine the amount of sub G₀/G₁ cells present, the AGS cell lines were treated with leaf extract at the maximal IC₅₀ concentration (118.85 µg/mL). The cells were stained with propidium iodide (PI) and analyzed using flow cytometry. Statistically significant accumulation of cells with sub-G₁ DNA content and other remarkably detectable cell cycle changes were noted after 24 h incubation. Consistent with an increased sub-G₁ AGS cell population, there was a notable decrease of cells in G₁, S and G₂/M phases. The control cells that were treated with phosphate buffer saline (PBS) alone showed no changes in cell cycle regulation (Figure 7b). Annexin V/PI staining followed by flow cytometry was used to confirm the effect of apoptosis on

Table 3 Number of transcripts found in *in vitro* roots and leaves putatively encoding enzymes involved in withanolide biosynthesis


Enzymes	Abbreviation	Enzyme commission number	No. of transcripts		
			Total	Expressed in both adventitious roots and leaves	Uniquely expressed in adventitious roots
1-Deoxy-D-xylulose-5-phosphate synthase	DXPS	2.2.1.7	8	8	0
1-Deoxy-D-xylulose-5-phosphate reductoisomerase	DXPR	1.1.1.267	3	3	0
2-C-Methyl-D-erythritol 4-phosphate cytidyltransferase	MEP-CT	2.7.7.60	2	2	0
4-(Cytidine 5'-diphospho)-2-C-methyl-D-erythritol kinase	CDP-MEK	2.7.1.148	3	2	1
2-C-Methyl-D-erythritol 2,4-cyclodiphosphate synthase	MECDPS	4.6.1.12	1	1	0
(E)-4-Hydroxy-3-methylbut-2-enyl-diphosphate synthase	HMBPPS	1.17.7.1	1	1	0
4-Hydroxy-3-methylbut-2-enyl diphosphate reductase	HMBPPR	1.17.1.2	1	1	0
Isopentenyl-diphosphate delta-isomerase	IDI	5.3.3.2	4	4	0
Acetyl-CoA C-acetyltransferase	ACAT	2.3.1.9	19	12	7
Hydroxymethylglutaryl-CoA synthase	HMGCS	2.3.3.10	5	5	0
Hydroxymethylglutaryl-CoA reductase	HMGR	1.1.1.88	11	11	0
Hydroxymethylglutaryl-CoA reductase	HMGR	1.1.1.34	9	9	0
Mevalonate kinase	MK	2.7.1.36	3	3	0
Phosphomevalonatekinase	PVMK	2.7.4.2	8	8	0
Diphosphomevalonate decarboxylase	MVD	4.1.1.33	1	1	0
Geranyl-diphosphate synthase	GPPS	2.5.1.1	38	35	3
Farnesyl-diphosphate synthase	FPPS	2.5.1.10	19	19	0
Squalene synthase	SS	2.5.1.21	5	5	0
Squalene monooxygenase	SE	1.14.13.132	10	10	0
Cycloartenol synthase	CAS	5.4.99.8	28	15	13
Sterol 24-C-methyltransferase	SMT1	2.1.1.41	7	7	0
Methylsterol monooxygenase/Sterol-4a-methyl oxidase 2	SMO1/SMO2	1.14.13.72	7	6	1
Cycloeucalenol cycloisomerase	CEC1	5.5.1.9	1	1	0
Obtusifoliol 14-demethylase	CPY51G1	1.14.13.70	3	3	0
Delta 14-sterol reductase	FK	1.3.1.70	3	3	0
C-7,8 Sterol isomerase	HYD1	5.3.3.5	2	1	1
C-5 Sterol desaturase	STE1	1.14.21.6	12	10	2
7-Dehydro cholesterol reductase	DWF5	1.3.1.21	2	2	0
Sterol glycosyltransferases	SGT	2.4.1.173	53	42	11
Methyltransferases	MT		509	466	43

AGS cells. Treatment of cells with methanolic extract of *in vitro*-grown leaf tissue resulted in more apoptotic cells, and the percentage of cells in late apoptosis was much higher than that in early apoptosis (Figure 7c). These data suggest that leaf extract from *in vitro*-grown *W. somnifera* inhibits AGS cell proliferation by inducing cell cycle arrest and apoptosis.

Comparative analysis of *in vitro*- and field-grown leaf and root transcripts

To investigate tissue or culture method-unique gene expression, we compared our 60-day-old *in vitro* leaf

and root transcripts with one-year old field-grown root and leaf sequences that were downloaded from the NCBI SRA database (accession no. SRR520143) [20]. We selected high-quality reads from *in vitro* root and leaf samples and assembled them separately using Trinity. This assembly yielded a total of 160,323 transcripts with an average length of 1,028 bp for root and 173,009 transcripts with an average length of 1,060 for leaf. Then, we mapped field-grown root sequences onto our *in vitro* root transcripts with a minimum alignment length of 200 bp. Only 5.7% of the transcripts were not found in field-grown roots, and we consider these transcripts as



in vitro root-specific transcripts. Similarly, field-grown leaf sequences were aligned onto our *in vitro* leaf data set, revealing about 3.9% of *in vitro* leaf-specific transcripts.

Furthermore, we assembled field-grown leaves and root tissues separately using CLC assembler to identify field-

grown specific transcripts. We did not find a significant number of field grown tissue-specific transcripts. This implies that our assembly from cultured tissue contains a large enough number of contigs to avoid much loss of information.




Table 4 Antioxidant activity in methanolic extracts of adventitious root and leaves

Activity of antioxidants	Tissue	
	<i>In vitro</i> adventitious root	<i>In vitro</i> leaf
Enzymatic antioxidants (U/mg protein)		
SOD	5.66 \pm 0.35	3.38 \pm 0.25
Peroxidase	3.02 \pm 0.42	2.5 \pm 0.57
GST	4.0 \pm 0.25	2.4 \pm 0.09
Polyphenol oxidase	3.50 \pm 0.23	2.56 \pm 0.35
Non-Enzymatic antioxidants (mg/g)		
Vitamin A	3.54 \pm 0.35	2.56 \pm 0.25
Vitamin C	3.8 \pm 0.27	1.47 \pm 0.46
Vitamin E	10.2 \pm 0.27	8.3 \pm 0.57
Free radical scavenging activity (FRSA)	72.5 \pm 0.58%	55 \pm 0.47%

The values of Mean \pm SD of triplicates.
 SOD (U/mg): amount of SOD that causes 50% reduction in the extent of NBT oxidation.
 Peroxidase (U/mg): change in absorbance/min at 430 nm.
 GST (U/mg protein): moles of CDNB conjugated/min/mg of sample.
 Polyphenol oxidase: (U/mg protein): change in absorbance/minute at 495 nm.




Figure 7 Influence of *in vitro* leaf extract on AGS cell growth inhibition. **a** - Inhibitory effect of *in vitro* methanolic leaf extract on AGS cell proliferation was measured by MTT assay. IC-50 value of the extract on the cell viability was identified as 118.85 µg/mL. **b** - DNA cell cycle analysis of control and treated AGS cells. Cells were stained with PI to determine cell cycle phase distribution based on fluorescence of DNA. **c** - Apoptosis level was examined by flow cytometry for control and treated AGS cells. Cells were treated with methanolic extract of *in vitro* grown leaf tissue (IC₅₀ - 118 µg/mL) for 24 h. Data are representative of three similar experiments.

Identification of cDNA-derived EST-SSR markers

Transcript-based SSR markers are important resources for genetic diversity analysis, marker-assisted selection, and genetic mapping. For identification of SSRs, *W. somnifera* Nr set transcripts were searched with perl script MISA (<http://pgrc.ipk-gatersleben.de/misa/>). The following criteria for SSR identification were used in the MISA script: mono-nucleotides repeated more than ten times, di-nucleotides repeated more than six times, and tri-, tetra-, penta- or hexa-nucleotides repeated more than five times. A total of 16,283 potential EST-SSRs were identified in 13,944 transcripts, of which 1,951 sequences contained more than 1 EST-SSR and 678 EST-SSRs were present in compound form. The di-nucleotide SSRs represented the most abundant type (2,489) of SSRs identified followed by tri-nucleotide (1,681) SSRs. We also identified tetra (92)-

and penta (20)-nucleotide SSR motifs. Although only a small fraction of tetra- and penta-SSRs were identified in *W. somnifera*, the number is quite significant and will provide a cost-effective method for development of functional genetic markers in *W. somnifera*.

Withanomics database

We developed a public data resource, Withanomics, which provides a searchable interface for the *W. somnifera* transcriptome data. Withanomics is freely available at <http://im-crop.snu.ac.kr/withania/index.php>. The current version of this database provides the total transcriptome sequences (root + leaf) and leaf (*in vitro*) sequences and root (*in vitro*) sequences separately reported in this study, as well as EST sequences produced from our previous study [30]. Expression data for transcripts can be accessed

using unique transcript IDs which report the expression of the transcripts in root and leaf tissues. Functional searches for all transcripts can be carried out using functional keywords with the annotation database. Additionally, we included an *in vitro* root-specific function option to extract function of *in vitro* root-specific transcripts. The database uses the NCBI BLAST algorithm (version: 2.2.15) for sequence-based searches. From BLAST searches, users can match nucleotide or protein sequence(s) against the *W. somnifera* whole transcriptome data, separate tissue transcriptome data, and ESTs from root and shoot data at user-defined parameters. Furthermore, we developed a sequence extract tool to extract or download single or bulk sequences from our database. We anticipate that the database will be useful for accelerating genomic research in *Withania* and we will be updating this database annually or as new sequence data sets are available for *Withania*.

Discussion

Establishment of *in vitro* cultures and transcriptome sequences of *W. somnifera*

W. somnifera is a medicinal plant of high therapeutic value, and withanolides are the major secondary metabolites responsible for most of the pharmacological activity of this plant. The accumulation of secondary metabolites in field-grown plants can be influenced by various environmental factors as well as the age of the plant at harvest [39]. The growing demand in the pharmaceutical industry and the lack of homogeneity in field-grown material led us to search for an alternative approach for mass production of *W. somnifera* roots. Adventitious *W. somnifera* roots grown *in vitro* have been reported to produce the pharmaceutically important withanolides, especially withanolide A [8,40–42]. Root tips are auxin-accumulating regions where active cell multiplication takes place, and hence were chosen for establishment of suspension cultures. Leaves from shoots devoid of roots were chosen from *in vitro* multiplied plants. This was done to ensure that the transcriptome library we constructed would represent transcripts exclusively specific to root or leaf tissues. *In vitro* cultures have been successfully exploited for the production of paclitaxel, an anticancer agent from *Taxus* species, at industrial scale, and attempts are being made to increase production using elicitors [43]. However, only when bottlenecks in the biosynthetic pathways have been identified will it be possible to overexpress the genes that control the rate-limiting steps and thus boost the production of valuable secondary metabolites in *in vitro* cultures.

In our current study, we generated a total of 249 million high-quality reads from *in vitro*-grown leaf and root tissues, more than the previously reported ESTs and transcriptome data [20,30,31]. We obtained a total transcriptome size of

183 Mbps with average transcript length of 1,033 bp (Table 1) which is comparatively better than a previous *Withania* 454 transcriptome study [20]. Significant differences in the number of transcripts annotated previously [20] and in the present study were documented. Functional genomics studies require highly reliable reference sequences and our assembly data can be utilized as such for *W. somnifera* due to the high sequencing depth. GO annotation showed that many transcripts were assigned to regulation-related process such as response to stress and wounding and defence responses in the biological process category. Transcripts related to GO term nucleus were most abundant in the cellular component category, which is similar to the previously reported GO annotation of field-grown root and leaf tissues [20]. Further, pathway assignment for all transcripts was performed based on the KEGG database. Previously, Gupta *et al.* [20] identified a total of 124 pathways among one-year-old field-grown leaf and root tissue transcripts, while 139 pathways were identified in the present study. In both studies, the phenylpropanoid biosynthesis pathway was found to harbour the largest cluster of transcripts. In a recent study, Dasgupta *et al.* [31] reported 184 pathways from their *in vitro* leaf transcriptome library. The additional pathways may be due to pathogenesis-related genes expressed in response to salicylic acid signalling in the source tissue of their study.

Identification and expression of genes related to withanolide biosynthesis

With respect to the genes involved in withanolide biosynthesis, we observed all genes to be expressed in both roots and leaves, similar to the results of Gupta *et al.* [20]. As a part of our ongoing endeavour, we validated important withanolide pathway genes including *HMGR*, *FPPS*, *SE*, *CAS* and *GT* at different time periods of culture (Figure 5). *HMGR* catalyses the reduction of HMG-CoA to mevalonate, reported to be the first committed step in the isoprenoid pathway [44]. *FPPS* catalyses the biosynthesis of FPP, which serves as a substrate for the first committed reaction of several branched pathways [45]. *SE* and *CAS* genes comprise vital genetic components of biosynthetic pathways leading to production of phytosterols and a large cluster of triterpenoids [46]. *GT* performs the glycosylation of a wide variety of natural compounds, which is an important mechanism in detoxification of exogenous compounds [47].

Among these genes, the expression of *HMGR* was notable in that it showed very high expression during the 30th day of culture in *in vitro* roots. Differences in expression of *HMGR* could be essential to accommodate the changing demand for a cellular array of isoprenoids during normal growth and development of the plant, as is evidenced by the tissue-specific expression of members of

this multi-gene family in several plant species [48]. *HMGR* has been reported to be present in two copies in *Arabidopsis thaliana* [49] and three to four copies in *Solanum* species [48]. Expression of *HMGR* varies in a time-dependent manner in *in vitro* cultures of *Dioscorea zingiberensis* [50], which is similar to our observation. *FPPS* is another of key enzyme in the isoprenoid biosynthetic pathway, wherein it initiates triterpenoid precursor biosynthesis to support withanolide production [51]. Any change in the levels of *FPPS* would alter the flux of isoprenoids towards various branches of this pathway, playing a crucial role in isoprenoid metabolism. *CAS* plays essential roles in plant cell viability and in the regulation of triterpenoid biosynthesis. Upregulation of *CAS* results in enhanced accumulation of diosgenin in *in vitro* cultures of *D. zingiberensis* [50]. In general, the expression of *FPPS* and *CAS* were found to be higher in leaves at all time points in *in vitro* shoots compared to roots, which is in accordance with previous observations [51,8].

SE is a noncytochrome-P450 type monooxygenase that catalyses the formation of a hydroxyl group that is characteristic of sterols and triterpenols and represents a rate-limiting step in sterol biosynthesis [52-54]. A higher level of *SE* expression was observed in *in vitro* roots than in leaf tissues, consistent with a previous report of higher expression of *SE* in one-year-old field-grown roots than in shoot tissues of *W. somnifera* [8]. Glucosylation of sterols is catalyzed by sterol glucosyltransferases (SGTs), which are family 1 glycosyltransferases. The expression of *SE* and *GT* was found to be higher in *in vitro* roots at all time periods compared to leaves. Among the time periods analysed, all genes investigated except *HMGR* showed the highest expression at the 45th day of culture.

In the present study, the high expression of *SE* observed in root tissue was in corroboration with the data of Sabir *et al.* [8], who reported a correlation between high expression of the genes for squalene epoxidase and squalene synthase and relatively higher accumulation of withanolide in field-grown root tissues. *SE* was reported to be the rate-limiting enzyme in sterol biosynthesis [52-54], and this could explain the high withanolide accumulation in 45-day-old *in vitro* root culture. Though expression of *FPPS* was higher in *in vitro* root tissue at 45 days, *WsFPPS* may not modulate withanolide content [51] as *FPP* is also reported to be involved in biosynthesis of triterpenoids, carotenoids, plastoquinones and many other molecules. Next to *FPPS*, expression of *CAS* was highest in root tissues. After the formation of epoxy squalene, *CAS* catalyzes the next step, which involves the formation of 24-methylene cholesterol, a precursor molecule for withanolide synthesis. Hence, expression of *CAS* also plays a significant role in the biosynthesis of withanolides.

We can surmise that the key pathway genes are expressed to different levels in different parts of the plant thus showing tissue-specific expression. This variation may be because of the presence of specific trans-acting factors required for their functional activation. Expression analysis of other genes involved in the biosynthetic pathway along with inhibitor studies will aid in enhancing the specific accumulation of desirable secondary metabolites.

Pattern of withanolide accumulation

It has been hypothesised that withanolides are produced in different parts of the plant through expression of the metabolic pathway rather than being imported from leaves [41]. Our current results support this hypothesis in that the expression of important genes involved in withanogenesis showed a differential pattern along with the accumulation of important withanolides, namely withanolide A and withaferin A. The accumulation of both withanolide A and withaferin A was found to be high during 45 days of culture, but interestingly, accumulation of withanolide A in leaves was nearly absent as was that of withaferin A in roots.

This observation is contradictory to the previous reports of Sabir *et al.* [8], but is in accordance with the observation of Dasgupta *et al.* [31] who reported a low concentration of withanolide A in control leaf samples, which are comparable to the 45-day-old leaves in the present study. The accumulation of withanolide A in roots followed a similar pattern to the previous report [8], where withaferin A was absent in *in vitro* root cultures. Thus, withaferin A was found to be tightly linked to leaf tissue indicating that there was no coordination of the accumulation profile between root and leaf tissue. The results of the present study are particularly important as they indicate that the *in vitro*-cultured roots and shoots possess *de novo* withanogenic competence and that *in vitro* cultures can be manipulated for efficient production of specific withanolides.

Antioxidant efficacy of *in vitro* tissues

We found that the *in vitro*-cultured roots of *W. somnifera* are good source of both non-enzymatic and enzymatic antioxidant components. *In vitro*-cultured root tissue showed high SOD and GST activity in addition to polyphenol oxidase and peroxidase activity. Present in all sub-cellular organelles, SOD serves as the first line of defence against free radicals. GSTs are most often thought of as detoxification enzymes and can metabolize a wide range of toxic exogenous compounds via GSH conjugation [55]. Polyphenol oxidase is localized in the chloroplast and is a copper-associated enzyme with two binding sites for phenolic substrates. Previous research reported that polyphenol oxidase is activated only when the plant undergoes

physical damage [56]. This might explain the relatively low level of polyphenol oxidase activity in *in vitro* root and leaf tissue. Increased peroxidase activity is associated with environmental stress on plants [57]. The low peroxidase activity in *in vitro* cultures could result from a related lack of expression of the enzyme. Among the non-enzymatic antioxidants, vitamin E was found to be the most abundant in *in vitro* root cultures. Vitamin E is a lipid-soluble vitamin found in cell membranes and circulating lipoproteins that offers protection against oxidative damage by acting directly with a variety of oxygen radicals. Its antioxidant function is strongly supported by regeneration promoted by vitamin C [58]. Hence, these results support the view that *in vitro*-cultured roots are promising source of potential antioxidants and function similar to field-grown tissues.

Combining the promising evidence that the *in vitro* leaf tissues express higher quantities of withaferin A than the *in vitro* root tissues, we proceeded to analyse the cytotoxicity of *W. somnifera* leaf extract on AGS cell lines. We found that the samples had the ability to linearly produce 50% cell death with the maximally effective concentration being 100–120 µg/mL. The effect of *W. somnifera* leaf extract on AGS cancer cells establishes that tissues grown *in vitro* are capable of inducing apoptosis and regulating the cell cycle of DNA biosynthesis. It has been previously reported that withaferin A, a promising anticancer constituent of *W. somnifera*, inhibits growth of human breast cancer cells in culture and *in vivo* in association with apoptosis induction [59]. Overall, the apoptosis and cell cycle analysis in cancer cells demonstrates that the *in vitro* *W. somnifera* leaves have anticancer effects, especially on human gastric adenocarcinoma cell lines.

Conclusion

Here, transcriptome profiles for *in vitro*-cultured leaf and root tissues of *W. somnifera* provided substantial information with respect to expression of genes involved in withanolide biosynthesis and accumulation of major withanolides. We have also shown that accumulation of different metabolites is linked to tissue type and that the extracts of *in vitro* root and leaf tissues have antioxidant potential and anti-proliferative capacity against an AGS cancer cell line, respectively. Further, we developed a user-friendly public database to accelerate research. Altogether, our research provides valuable information on the *W. somnifera* *in vitro* transcriptome and withanolide biosynthetic pathway gene expression to promote effective engineering of the pathway genes for improved biosynthesis of withanolides.

Methods

Plant material and RNA extraction

Seeds of *W. somnifera* (L.) Dunal 'Jawahar Asgandh 20' were obtained from the Central Institute of Medicinal

and Aromatic Plants (Lucknow), and germinated *in vitro* on MS solid basal medium supplemented with 2% sucrose in the dark at 25°C. Shoots from *in vitro*-germinated seedlings were maintained on MS basal medium under standard culture conditions. Shoots were multiplied on MS medium supplemented with 1 mg/L BAP, with regular subcultures. Multiplied shoots were transferred to hormone-free medium and maintained. Leaves from these shoots devoid of roots were used for RNA extraction. Adventitious roots were induced from leaf explants following the procedure described by Wasnik et al. [60]. Direct adventitious roots were induced from leaf explants of *W. somnifera* cultivar 'Jawahar Asghand-20' on Murashige and Skoog (MS) medium supplemented with 30 g/L sucrose, 1 mg/L IBA and 0.25 mg/L IAA. For mass culture of adventitious roots, the root tips and branches from *in vitro*-induced adventitious roots were transferred to liquid MS basal medium (Figure 1) in a bubble column bioreactor (Biopia, Korea). RNA was extracted from adventitious root and leaf tissues grown on MS medium following standard protocols. The tissues were submerged in liquid nitrogen, snap frozen and crushed using a mortar and pestle. Total RNA was extracted from the frozen tissues using TRIzol reagent (Bangalore genei, India) according to the manufacturer's instructions. Briefly, 1.5 mL TRIzol was added to 200 mg tissue, homogenized and centrifuged. The supernatant was mixed with 0.2 mL chloroform and incubated at -20°C for 20 min. The tubes were then vortexed and centrifuged, and the aqueous phase was mixed well with 0.2 mL 2-propanol followed by centrifugation. All centrifugations were carried out at 4°C, 7200 × g for 20 min. Following centrifugation, the supernatant was discarded, and the RNA pellet was washed with 75% ethanol and vortexed. Samples were again centrifuged at 7200 × g for 10 min at 4°C. After removal of the supernatant, the remaining ethanol was allowed to evaporate. The total RNA was re-suspended in diethylpyrocarbonate (DEPC)-treated water.

Illumina sequencing and quality control

Raw reads (101 bp PE) were generated for 60-day-old *in vitro*-cultured adventitious root and leaf tissues without replication using the Illumina HiSeq2000. The library construction and sequencing was performed by the Macrogen (<http://dna.macrogen.com/>, Korea). The sequence data generated in this study have been deposited at NCBI in the Short Read Archive (SRA) with accession number SRP040231. The sequencing reads were cleaned by removing low-quality reads, primer adaptor sequence using NGS QC Toolkit [61]. De novo assembly was performed using Trinity [32] and to eliminate mis-assemblies and ensure a collection of informative transcripts, read depth analysis was carried out. The read depth percentage was determined by dividing the number of mapped reads for

each assembled transcript by the effective length (without 'N') of the transcript. Transcripts with <1% read-depth were removed as those were not considered to be assembled correctly from paired reads.

Functional annotation

Sequence similarity searches were conducted against the NCBI non-redundant (Nr) database, tomato ([http://solgenomics.net/organism/Solanum_lycopersicum/genome;version ITAG2.3](http://solgenomics.net/organism/Solanum_lycopersicum/genome;version_ITAG2.3)), potato (http://solgenomics.net/organism/Solanum_tuberosum/genome), and TAIR (The Arabidopsis Information Resource) protein databases using the BLASTx algorithm with an *E* value threshold of 10^{-5} . The BLAST2GO program was used to obtain GO terms for all assembled transcripts [35]. In many cases, multiple terms were assigned to the same transcripts, and the GO terms were classified into biological process, cellular component, and molecular function.

Quantitative reverse-transcription PCR (qRT-PCR) analysis

For expression analysis of pathway genes, adventitious root and leaf tissues were collected after 30, 45 and 60 days of culture for total RNA isolation using TRIzol reagent (Bangalore genei, India) [62]. Extracted RNA was treated enzymatically with DNase I to remove contaminant genomic DNA. Quality of RNA was examined in 1.2% denaturing formaldehyde agarose gels, and the concentration was determined using a spectrophotometer. cDNA synthesis was carried out from 2 µg RNA template using Superscript II reverse transcriptase (Invitrogen, USA), according to the manufacturer's instructions. qRT-PCR was performed in a realplex² Mastercycler (Eppendorf, Germany) in a 10-µL reaction volume containing 100 ng cDNA template with Luminoct SYBR[®] Green Master Mix (Sigma Aldrich, USA) and gene-specific primers. Specific primers were designed using the software primer3 [63] (Additional file 4: Table S1). The primers for *GAPDH* were used as a control to ensure that equal amounts of cDNA were used in all reactions. The PCR conditions were as follows: predenaturation at 95°C for 5 min, 40 cycles of 95°C for 30 s, 60°C for 30 s, 72°C for 30 s. The increase in fluorescence corresponding to the exponential increase in the product was used to determine the threshold cycle (C_t) in each reaction. Relative expression levels were calculated by the $2^{-\Delta\Delta C_t}$ method [64] with *GAPDH* as an internal control. All qRT-PCR reactions were performed in triplicate.

Quantification of major withanolides

Total withanolides from 1.0 g *in vitro* adventitious root and leaf tissues were extracted as described previously [65]. The final extraction was done in methanol, and the samples were evaporated to dryness, dissolved in 5.0 mL HPLC-grade methanol and subjected to HPTLC analysis.

The methanolic extracts of the *in vitro* adventitious root and *in vitro* leaf (root extract and leaf extract, respectively) tissues were applied to the plates as 6-mm bands, under a stream of nitrogen, by means of a Linomat V semiautomatic sample applicator (CAMAG, Switzerland) fitted with a 100-µL syringe. Linear ascending development to a distance of 8 cm was carried out on a twin-trough chamber saturated with the mobile phase, toluene: ethyl acetate: formic acid (5: 5: 1). Subsequent to development, the banding patterns were visualized in 254 nm, 366 nm and white light and the R_f values were calculated. Densitometric scanning was performed with a Scanner III (CAMAG, Switzerland) in the reflectance-absorbance mode at 234 nm for withanolide A and 223 nm for withaferin A [37]. Concentrations of the compound chromatographed were determined from the intensity of diffusely reflected light. Evaluation was carried out by comparing peak areas with linear regression [66]. Pearson's correlation coefficient (*r*) was used to correlate the expression levels with metabolite accumulation in 45- and 60 day-old culture in bivariate linear correlations.

Analysis of antioxidant activity

Total enzymatic and non-enzymatic antioxidant activity in methanolic extracts of 60-day-old adventitious root and leaf tissues were analysed following standard protocols. Total antioxidant activity was assayed using DPPH scavenging activity as described by Ansari *et al.* [67] and ascorbic acid was used as reference material. For the enzymatic antioxidants, the superoxide dismutase assay was performed following Beauchamp and Fedovich [68], that for polyphenol oxidase according to Ensimer [69], peroxidase according to Malik and Singh [70] and glutathione S transferase according to Habig *et al.* [71], and controls were run simultaneously without the plant extract. Among the non-enzymatic antioxidants, contents of ascorbic acid, vitamin A, and α -tocopherol were determined following the procedure of Roe and Kuether [72], Bayfield and Cole [73], and Backer *et al.* [74], respectively. The complete assay mixture without the plant extract served as the control to monitor non-specific binding of the substrates.

Cell proliferation analysis using MTT assay

The capacity of methanolic extracts of 60-day-old leaves, harvested from shoots maintained *in vitro*, to prevent proliferation of AGS cells was determined by performing 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) assay. AGS cells were seeded at 10×10^4 cells/mL in a 12-well plate and incubated for 24 h. The plant extracts were evaporated to dryness and dissolved in PBS before performing the experiment. The cells were treated with various concentrations of methanolic extracts of *in vitro*

grown leaf tissues and incubated for 24 h. The cells treated with PBS alone (0 µg/mL) were considered to be the control with 100% viability. After incubation, 100 µL MTT solution (5 mg/mL in 1× Phosphate Buffered Saline, PBS) was added to the wells and incubated for 3 h. Then 500 µL DMSO was added to each well after complete removal of the medium to dissolve the cellular crystalline deposits and kept for 20 minutes. The optical density was measured at 540 nm using an ELISA plate reader (EIA plate, USA).

Cell cycle analysis using propidium iodide

AGS cells were seeded in 5-mL plates at a concentration of 60×10^4 cells/well and incubated overnight at 37°C in an atmosphere of 5% CO₂. The cells were treated with various concentrations of methanolic extracts of *in vitro*-grown leaf tissues and incubated for 24 h. Cells were trypsinized, washed twice with cold PBS, and centrifuged. The pellet was fixed using cold 70% ethanol for 30 minutes at 4°C. The cells were washed once with PBS and resuspended in cold PI solution (50 µg/mL) containing RNase A (0.1 mg/mL) in PBS for 30 min in dark. In a Fluorescence Activated Cell Sorting (FACS) Calibur apparatus (Becton Dickinson, San Jose, CA, USA), approximately 10,000 counts were done for each sample. The relative proportions of G0/G1, S and G2/M cells were determined for cell cycle analysis. By determining the percentage of cell distribution in each phase, the effect of leaf extract on cell cycle was measured.

Cell apoptosis analysis

Annexin V/PI staining was employed to classify apoptosis levels in *in vitro* leaf extract-treated AGS cells using an Annexin V-fluorescein isothiocyanate (FITC) apoptosis detection kit (BDPharmingen, CA, USA). Approximately 5×10^5 cells were seeded into a 60-mL culture dish 24 h before being treated with extract. After 24 h, cells were trypsinized, washed with cold PBS and resuspended in 100 µL binding buffer. The cells were then stained with Annexin V-FITC and PI each in a Ca²⁺-enriched binding buffer. The cells were then incubated for 15 min in dark, 400 µL binding buffer was added and the assay was begun immediately using a FACScan flow cytometer (Becton Dickinson). The staining emissions were detected in the FL-1 and FL-3 channels. Approximately 10,000 counts were made for each sample. The distribution of live, early apoptosis, late apoptosis and necrotic cells were analysed.

Withanomics database

The Withanomics database was created to serve as a public web resource for *Withania* transcriptome data. This database was designed using PHP (v4.3.9) and MySQL (v4.1.20). The front-end language PHP was connected

with back-end MySQL by the Apache web server. The annotation, expression and other analysis data are stored in MySQL as tables, and this database is currently hosted on a CentOS (v5.8) Linux operating system. This database can be accessed at <http://im-crop.snu.ac.kr/withania/index.php>.

Additional files

Additional file 1: Figure S1. Differentially expressed transcripts between *in vitro* root and leaf tissues.

Additional file 2: Figure S2. The expression of the transcripts related HMGR, FPPS, SE, CAS, GT in *in vitro* root and leaf tissues are shown in heatmap.

Additional file 3: Figure S3. HPTLC chromatogram of standards, withaferin A and withanolide A and 45-day-old *in vitro* leaf and root samples. A - Chromatogram of standard withaferin A; B - Chromatogram of standard withanolide A; C - Chromatogram of 45-day-old *in vitro* leaf sample indicating the presence of withaferin A; D - Chromatogram of 45-day-old *in vitro* root sample indicating the presence of withanolide A.

Additional file 4: Table S1. Primers used for qRT-PCR analysis.

Abbreviations

HMGR: Hydroxymethylglutaryl-CoA reductase; FPPS: Farnesyl-diphosphate synthase; SE: Squalene monooxygenase; CAS: Cycloartenol synthase; GT: Glucosyltransferases; GAPDH: Glycerinaldehyde 3-phosphate dehydrogenase; SSR: Single sequence repeat.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KS, TSM and TJY conceived, designed and executed the study; PD and KR established and maintained *W. somnifera in vitro* cultures; SCL and YSL helped with sequencing; MJ carried out all bioinformatics analysis; PT performed expression studies, quantification and antioxidant analysis; PMP, SNC, IMR, AN, GSK worked on anti-proliferative activity and *in vitro* cytotoxicity. SN participated in design and drafting of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by a Major Research Project from the University Grants Commission GOI [F. No. 34-278/2008 (SR) dt. 2/01/2009], a RAPID Project from the Department of Biotechnology, GOI (No. BT/PR 11845/GBD/27/168/2009 dt. 4/02/2009), the WOS-A Program of the Department of Science and Technology [SR/WOS-A/LS-532/2011(G)], New Delhi, and the Next-Generation BioGreen 21 Program (No. PJ011030) of the Rural Development Administration, Republic of Korea.

Author details

¹Department of Biochemistry, Biotechnology and Bioinformatics, Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore 641043, India. ²Department of Plant Science, Plant Genomics and Breeding Institute, Research Institute for Agriculture and Life Sciences, College of Agriculture and Life Sciences, Seoul National University, Seoul 151-921, Republic of Korea. ³Lab of Biochemistry, School of Veterinary Medicine, Gyeongsang National University, Gyeongsangnam-do, Republic of Korea. ⁴Genomics and Proteomics Laboratory, Centre for Plant Molecular Biology & Biotechnology, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India. ⁵National Research Foundation, Seoul, Republic of Korea.

Received: 22 September 2014 Accepted: 2 January 2015
Published online: 22 January 2015

References

- Mirjalili MH, Moyano E, Bonfill M, Cusido RM, Palazón J. Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine. *Molecules*. 2009;14:2373–93.
- Sachdeva H, Sehgal R, Kaur. Studies on the protective and immunomodulatory efficacy of *Withania somnifera* along with cisplatin against experimental visceral leishmaniasis. *Parasitol Res*. 2013;112:2269–80.
- Sehgal N, Gupta A, Valli RK, Joshi SD, Mills JT, Hamel E, et al. *Withania somnifera* reverses Alzheimer's disease pathology by enhancing low-density lipoprotein receptor-related protein in liver. *Proc Natl Acad Sci*. 2012;109:3510–5.
- VenMurthy MR, Ranjekar PK, Ramassamy C, Deshpande M. Scientific basis for the use of Indian ayurvedic medicinal plants in the treatment of neurodegenerative disorders: ashwagandha. *Cent Nerv Syst Agents Med Chem*. 2010;10:238–46.
- Singh N, Verma P, Pandey BR, Gilca M. Role of *Withania somnifera* in prevention and treatment of cancer: an overview. *Int J Pharm Sci Drug Res*. 2011;3:274–9.
- Mansour HH, Hafez FH. Protective effect of *Withania somnifera* against radiation-induced hepatotoxicity in rats. *Ecotoxicol Environ Saf*. 2012;80:14–9.
- Chaurasiya ND, Sangwan RS, Misra LN, Tuli R, Sangwan NS. Metabolic clustering of a core collection of Indian ginseng *Withania somnifera* Dunal through DNA, isoenzyme, polypeptide and withanolide profile diversity. *Fitoterapia*. 2009;80:496–505.
- Sabir F, Mishra S, Sangwan RS, Jadaun JS, Sangwan NS. Qualitative and quantitative variations in withanolides and expression of some pathway genes during different stages of morphogenesis in *Withania somnifera* Dunal. *Protoplasma*. 2012;250:539–49.
- Misra L, Lal P, Sangwan RS, Sangwan NS, Uniyal GC, Tuli R. Unusually sulfated and oxygenated steroids from *Withania somnifera*. *Phytochemistry*. 2005;66:2702–7.
- Madina BR, Sharma LK, Chaturvedi P, Sangwan RS, Tuli. Purification and characterization of a novel glucosyltransferase specific to 27b-hydroxy steroidal lactones from *Withania somnifera* and its role in stress responses. *Biochem Biophys Acta*. 2007;1774:1199–207.
- Chaurasiya ND, Sangwan NS, Sabir F, Misra L, Sangwan RS. Withanolide biosynthesis recruits both mevalonate and DOXP pathways of isoprenogenesis in Ashwagandha *Withania somnifera* L. (Dunal). *Plant Cell Rep*. 2012;31:1889–97.
- Prajapati ND, Purohit SS, Sharma AK, Kumar. A handbook of medicinal plants. Jodhpur, India: Agrobios; 2007.
- Ramawat KG, Goyal S. The Indian herbal drugs scenario in global perspectives. In: Merillon JM, editor. *Bioactive molecules and medicinal plants*. Berlin, Heidelberg New York: Springer; 2008. p. 331.
- Sharada M, Ahuja A, Suri KA, Vij SP, Khajuria RK, Verma V, et al. Withanolide production by *in vitro* cultures of *Withania somnifera* (L.) and its association with differentiation. *Biol Plantarum*. 2007;51:161–4.
- Sivanandhan G, Arun M, Mayavan S, Rajesh M, Mariashibu TS, Manickavasagam M, et al. Chitosan enhances withanolides production in adventitious root cultures of *Withania somnifera* (L.) Dunal. *Ind Crop Prod*. 2012;37:124–9.
- Bernard F, Moghbel N, Hassannejad S. Treatment of licorice seeds with colchicine: changes in seedling DNA levels and anthocyanin and glycyrrhizic acid contents of derived callus cultures. *Nat Prod Commun*. 2012;7:1457–60.
- Sharma P, Yadav S, Srivastava A, Shrivastava N. Methyl jasmonate mediates upregulation of bacoside A production in shootcultures of *Bacopa monnieri*. *Biotechnol Lett*. 2013;35:1121–5.
- Marchev A, Christiane H, Sibylle S, Vasil G, Juliane S, Thomas B, et al. Sage *in vitro* cultures: a promising tool for the production of bioactive terpenes and phenolic substances. *Biotechnol Lett*. 2014;36:211.
- Sharma LK, Madina BR, Chaturvedi P, Sangwan RS, Tuli R. Molecular cloning and characterization of one member of 3b-hydroxy sterol glucosyltransferase gene family in *Withania somnifera*. *Arch Biochem Biophys*. 2007;460:48–55.
- Gupta P, Goel R, Pathak S, Srivastava A, Singh SP, Sangwan RS, et al. *De novo* assembly, functional annotation and comparative analysis of *Withania somnifera* leaf and root transcriptomes to identify putative genes involved in the withanolides biosynthesis. *PLoS One*. 2013;8:e62714.
- Jayakodi M, Lee SC, Park HS, Jang WJ, Lee YS, Choi BS, et al. Transcriptome Profiling and comparative analysis of *Panax ginseng* adventitious roots. *J Ginseng Res*. 2014. <http://dx.doi.org/10.1016/j.jgr.2014.05.008>.
- Luo H, Sun C, Sun Y, Wu Q, Li Y. Analysis of the transcriptome of *Panax notoginseng* root uncovers putative triterpene saponin-biosynthetic genes and genetic markers. *BMC Genomics*. 2011;12 Suppl 5:S5.
- Sun C, Li Y, Wu Q, Luo H, Sun Y. *De novo* sequencing and analysis of the American ginseng root transcriptome using a GS FLX Titanium platform to discover putative genes involved in ginsenoside biosynthesis. *BMC Genomics*. 2010;11:262.
- Li C, Zhu Y, Guo X, Sun C, Luo H. Transcriptome analysis reveals ginsenosides biosynthetic genes, microRNAs and simple sequence repeats in *Panax ginseng* C. A. Meyer. *BMC Genomics*. 2013;14:245.
- Hyun TK, Rim Y, Jang H-J, Kim CH, Park J, Kumar R, et al. *De novo* transcriptome sequencing of *Momordica cochinchinensis* to identify genes involved in the carotenoid biosynthesis. *Plant Mol Biol*. 2012;79:413–27.
- Shi CY, Yang H, Wei CL, Yu O, Zhang ZZ, Jiang CJ, et al. Deep sequencing of the *Camellia sinensis* transcriptome revealed candidate genes for major metabolic pathways of tea-specific compounds. *BMC Genomics*. 2011;12:131.
- Li H, Dong Y, Yang J, Liu X, Wang Y, Yao N, et al. *De novo* transcriptome of safflower and the identification of putative gene for oleosin and the biosynthesis of flavonoids. *PLoS One*. 2012;7:e30987.
- Liu S, Li W, Wu Y, Chen C, Lei J. *De novo* transcriptome assembly in chili pepper (*Capsicum frutescens*) to identify genes involved in the biosynthesis of capsaicinoids. *PLoS One*. 2013;8:e48156.
- Wang Y, Pan Y, Liu Z, Zhu X, Zhai L, Xu L, et al. *De novo* transcriptome sequencing of radish (*Raphanus sativus* L.) and analysis of major genes involved in glucosinolate metabolism. *BMC Genomics*. 2013;14:836.
- Senthil K, Wasnik NG, Kim YJ, Yang DC. Generation and analysis of expressed sequence tags from leaf and root of *Withania somnifera* (Ashwagandha). *Mol Biol Rep*. 2010;37:893–902.
- Dasgupta MG, George BS, Bhatia A, Sidhu OP. Characterization of *Withania somnifera* leaf transcriptome and expression analysis of pathogenesis – related genes during salicylic acid signaling. *PLoS One*. 2014;9:e94803.
- Grabherr MG, Haas BJ, Yassour M. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. 2011;15:644–52.
- Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics*. 2011;12:323.
- Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010;26:139–40.
- Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics*. 2005;21:3674–6.
- Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res*. 2000;28:27–30.
- Sharma V, Gupta AP, Bhandari P, Gupta RC, Singh B. A validated and quantitative HPTLC method for the quantification of Withaferin-A and Withanolide-A in different plant parts of two morphotypes of *Withania somnifera*. *Chromatographia*. 2007;66:801–4.
- Jaleel CA, Gopi R, Sankar B, Manivannan P, Kishorekumar A, Sridharan R, et al. Alterations in germination, seedling vigour, lipid peroxidation and proline metabolism in *Catharanthus roseus* seedlings under salt stress. *S Afr J Bot*. 2007;73:190–5.
- Awasthi D, Meitei KV, Misra R, Ali SA. Validation of harvesting period for obtaining optimum concentrations of Withanolides from *Withania somnifera* (L.) Dunal at different phenological stages of plant life. *Indan J Trop Biodiv*. 2009;16:129–32.
- Hahn EJ, Kim YS, Yu KW, Jeong CS, Paek KY. Adventitious root cultures of *Panax ginseng* c.v. meyer and ginsenoside production through large-scale bioreactor system. *J Plant Biotechnol*. 2003;5:1–6.
- Sangwan RS, Chaurasiya ND, Sangwan PL, Misra LN, Tuli R, Sangwan NS. Withanolide A is inherently *de novo* biosynthesized in roots of the medicinal plant Ashwagandha (*Withania somnifera*). *Physiol Plant*. 2008;133:278–87.
- Nagella P, Murthy HN. Establishment of cell suspension cultures of *Withania somnifera* for the production of withanolide A. *Bioresour Technol*. 2010;176:735–9.
- Onrubia M, Cusido RM, Ramirez K, Hernández-Vázquez L, Moyano E, Bonfill M, et al. Bioprocessing of plant *in vitro* systems for the mass production of pharmaceutically important metabolites: paclitaxel and its derivatives. *Curr Med Chem*. 2013;20:880–91.
- Abdin MZ, Kiran U, Aquil S. Molecular cloning and structural characterization of HMG-CoA reductase gene from *Catharanthus roseus* (L.) G Donn Cv Albus. *Indian J Biotech*. 2012;11:16–22.

45. Cunillera N, Arro M, Delourme D, Karst F, Boronat A, Ferrer A. *Arabidopsis thaliana* contains two differentially expressed farnesyl diphosphate synthase genes. *J Biol Chem*. 1996;271:7774–80.
46. Dhar N, Rana S, Bhat WW, Razdan S, Pandith SA, Khan S, et al. Dynamics of withanolide biosynthesis in relation to temporal expression pattern of metabolic genes in *Withania somnifera* (L.) Dunal: a comparative study in two morpho-chemovariants. *Mol Biol Rep*. 2013;40:7007–16.
47. Sakakibara KY. Functional genomics of family 1 glucosyltransferase in *Arabidopsis*. *Plant Biotech J*. 2009;26:267–74.
48. Dale S, Arro M, Becerra B, Morrice NG, Boronat A, Hardie DG, et al. Bacterial expression of the catalytic domain of 3-hydroxy-3-methylglutaryl-CoA reductase (isoform HMGR1) from *Arabidopsis thaliana*, and its inactivation by phosphorylation at Ser577 by *Brassica oleracea* 3-hydroxy-3-methylglutaryl-CoA reductase kinase. *Eur J Biochem*. 1995;233:506–13.
49. Choi D, Ward BL, Bostock RM. Differential induction and suppression of potato 3-hydroxy-3-methylglutaryl coenzyme A reductase genes in response to *Phytophthora infestans* and to its elicitor. *Plant Cell*. 1992;4:1333–44.
50. Diarra S, He J, Wang J, Li J. Ethylene treatment improves diosgenin accumulation in *in vitro* cultures of *Dioscorea zingiberensis* via up-regulation of CAS and HMGR gene expression. *Electron J Biotechnol*. 2013;16:6.
51. Gupta P, Akhtar N, Tewari SK, Sangwan RS, Trivedi PK. Differential expression of farnesyl diphosphate synthase gene from *Withania somnifera* in different chemotypes and in response to elicitors. *Plant Growth Regul*. 2011;65:93–100.
52. Hidaka Y, Satoh T, Kamei T. Regulation of squalene epoxidase in HepG2 cells. *J Lipid Res*. 1990;31:2087–94.
53. Abe I, Prestwich GD. Squalene epoxidase and oxidosqualene:lanosterol cyclase-key enzymes in cholesterol biosynthesis. In: Cane DE, editor. *Comprehensive natural products chemistry*. Volume 2. Oxford: Elsevier; 1999. p. 267–98.
54. Razdan S, Bhat WW, Rana S, Dhar N, Lattoo SK, Dhar RS, et al. Molecular characterization and promoter analysis of squalene epoxidase gene from *Withania somnifera* (L.) Dunal. *Mol Biol Rep*. 2012;40:905–16.
55. Khatun S, Ali MB, Hahn E, Paek K. Copper toxicity in *Withania somnifera*: Growth and antioxidant enzymes responses of *in vitro* grown plants. *Environ Exp Bot*. 2008;64:279–85.
56. Golbeck JH, Cammarata KV. Spinach thylakoid polyphenol oxidase isolation, activation properties of the native chloroplast enzyme. *Plant Physiol*. 1981;67:977–84.
57. Rabe R, Kreeb KH. Enzyme activities and chlorophyll and protein content in plants as indicators of air pollution. *Environ Pollut*. 1979;19:119–36.
58. Fusco D, Colloca G, Cesari M. Effects of antioxidant supplementation on the aging process. *Clin Interv Aging*. 2007;2:377–87.
59. Lee J, Hahn ER, Singh SV. Withaferin A inhibits activation of signal transducer and activation of transcription 3 in human breast cancer cell. *Carcinogenesis*. 2010;31:1991–8.
60. Wasnik NG, Muthusamy M, Chellappan S, Vaidhyanathan V, Pulla R, Senthil K, et al. Establishment of *in vitro* root cultures and analysis of secondary metabolites in Indian Ginseng - *Withania somnifera*. *Korean J Plant Res*. 2009;22:584–91.
61. Patel RK, Jain M. NGS QC Toolkit: a toolkit for quality control of next generation sequencing data. *PLoS One*. 2012;7:e30619.
62. Tank JG, Thaker VS. Changes in DNA and RNA level with endoreduplication can be determined using α -naphthyl red dye. *CIBTech J Bio Protocols*. 2012;1:23–31.
63. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. In: Misener S, Krawetz SA, editors. *Methods Mol Biol*. Volume 132. Totowa, NJ: Humana Press Inc.; 2000. p. 365–86.
64. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. *Methods*. 2001;25:402–8.
65. Patel JB, Lahiri SK, Shah MB. Development of a new method for identification and estimation of *Withania somnifera* root, and a method for quantitative analysis of withaferin A in young and old roots. *J Planar Chromatogr- Mod TLC*. 2009;22:283–6.
66. Jirge SS, Tatke PA, Gabhe SY. Development and validation of a novel HPTLC method for simultaneous estimation of betasitosteroldglucoside and withaferin A. *Int J Pharm Pharm Sci*. 2011;3:227–30.
67. Ansari AQ, Ahmed SA, Waheed MA, Juned S. Extraction and determination of antioxidant activity of *Withania somnifera* Dunal. *Eur J Exp Biol*. 2013;3:502–7.
68. Beauchamp BC, Fedovich. Superoxide dismutase assay and an assay applicable to acrylamide gel. *Anal Biochem*. 1976;10:276–87.
69. Enslinger K, Vamos – Vigyazo E. Inactivation of polyphenol oxidase and depletion of O-dihydroxyphenol content during enzymatic browning of fruit tissue. *J Amer*. 1995;12:1–9.
70. Malik CP, Singh MB. *Plant enzymology and histo-enzymology – A text manual*. New Delhi: Kalyani Publishers; 1980. p. 30–54.
71. Habig WH, Palist MJ, Jakoby WB. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem*. 1974;249:7130–9.
72. Roe JH, Kuether CA. Determination of ascorbic acid in whole blood and urine through the 2,4-dinitrophenyl hydrazine derivative of ascorbic acid. *J Biol Chem*. 1953;143:399–406.
73. Bayfield RF, Cole ER. Colorimetric estimation of vitamin A with trichloroacetic acid. *Meth Enzymol*. 1980;67:189–95.
74. Backer H, Frank O, De Angells B, Feingold S. Plasma tocopherol in man at various times after ingesting free or ocetylaned tocopherol. *Nutr Rep Int*. 1980;21:531–6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

