

NBER WORKING PAPER SERIES

TRANSFORM ANALYSIS AND
ASSET PRICING FOR AFFINE
JUMP-DIFFUSIONS

Darrell Duffie
Jun Pan
Kenneth Singleton

Working Paper 7105
<http://www.nber.org/papers/w7105>

NATIONAL BUREAU OF ECONOMIC RESEARCH
1050 Massachusetts Avenue
Cambridge, MA 02138
April 1999

We are grateful for extensive discussions with Jun Liu; conversations with Jean Jacod, Monika Piazzesi, Philip Protter, and Ruth Williams; and support from the Financial Research Initiative, The Stanford Program in Finance, and the Gifford Fong Associates Fund, at the Graduate School of Business, Stanford University. The views expressed herein are those of the authors and do not necessarily reflect the views of the National Bureau of Economic Research.

© 1999 by Darrell Duffie, Jun Pan, and Kenneth Singleton. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Transform Analysis and Asset Pricing for
Affine Jump-Diffusions
Darrell Duffie, Jun Pan, and Kenneth Singleton
NBER Working Paper No. 7105
April 1999
JEL No. G1

ABSTRACT

In the setting of “affine” jump-diffusion state processes, this paper provides an analytical treatment of a class of transforms, including various Laplace and Fourier transforms as special cases, that allow an analytical treatment of a range of valuation and econometric problems. Example applications include fixed-income pricing models, with a role for intensity-based models of default, as well as a wide range of option-pricing applications. An illustrative example examines the implications of stochastic volatility and jumps for option valuation. This example highlights the impact on option ‘smirks’ of the joint distribution of jumps in volatility and jumps in the underlying asset price, through both amplitude as well as jump timing.

Darrell Duffie
Graduate School of Business
Stanford University
Stanford, CA 93405
duffie@stanford.edu

Jun Pan
Graduate School of Business
Stanford University
Stanford, CA 93405
junpan@stanford.edu

Kenneth Singleton
Graduate School of Business
Stanford University
Stanford, CA 93405
and NBER
kenneths@future.stanford.edu

1 Introduction

In valuing financial securities in an arbitrage-free environment, one inevitably faces a trade-off between the analytical and computational tractability of pricing and estimation, and the complexity of the probability model for the state vector X . In the light of this trade-off, academics and practitioners alike have found it convenient to impose sufficient structure on the conditional distribution of X to give closed- or nearly closed-form expressions for securities prices. An assumption that has proved to be particularly fruitful in developing tractable, dynamic asset pricing models is that X follows an *affine jump-diffusion (AJD)*, which is, roughly speaking, a jump-diffusion process for which the drift vector, “instantaneous” covariance matrix, and jump intensities all have affine dependence on the state vector. Prominent among *AJD* models in the term-structure literature are the Gaussian and square-root diffusion models of Vasicek [1977] and Cox, Ingersoll, and Ross [1985]. In the case of option pricing, there is a substantial literature building on the particular affine stochastic-volatility model for currency and equity prices proposed by Heston [1993].

This paper synthesizes and significantly extends the extant literature on affine asset pricing models by deriving a closed-form expression for an “extended transform” of an *AJD* process X , and then showing that this transform leads to analytically tractable pricing relations for a wide variety of valuation problems. More precisely, fixing the current date t and a future payoff date T , suppose that the stochastic “discount rate” $R(X_t)$, for computing present values of future cash flows, is an affine function of X_t . Also, consider the generalized terminal payoff function $(v_0 + v_1 \cdot X_T) e^{u \cdot X(T)}$ of X_T , where v_0 is scalar and the n elements of each of the v_1 and u are scalars. These scalars may be real, or more generally, complex. We derive a closed-form expression for the transform

$$E_t \left(\exp \left(- \int_t^T R(X_s, s) ds \right) (v_0 + v_1 \cdot X_T) e^{u \cdot X_T} \right), \quad (1.1)$$

where E_t denotes expectation conditioned on the history of X up to t . Then, using this transform, we show that the tractability offered by extant, specialized affine pricing models extends to the entire family of *AJDs*. Additionally, by selectively choosing the payoff $(v_0 + v_1 \cdot X_T) e^{u \cdot X(T)}$, we significantly extend the set of pricing problems (security payoffs) that can be tractably addressed

with X following an AJD . To motivate the usefulness of our extended transform in theoretical and empirical analyses of affine models, we briefly outline three applications.

1.1 Affine, Defaultable Term Structure Models

There is a large literature on the term structure of default-free bond yields that presumes that the state vector underlying interest rate movements follows an AJD (see, e.g., Dai and Singleton [1999] and the references therein). Assuming that the instantaneous riskless short-term rate r_t is an affine function of an n -dimensional AJD process X_t (that is $r_t = \rho_0 + \rho_1 \cdot X_t$) Duffie and Kan [1996] show that the $(T - t)$ -period zero-coupon bond price,

$$E_t \left(\exp \left(- \int_t^T r_s ds \right) \right), \quad (1.2)$$

is known in closed form, where expectations are computed under the risk-neutral measure.¹

Recently, considerable attention has been focused on extending these models to allow for the possibility of default in order to price corporate bonds and other credit-sensitive instruments.² To illustrate the new pricing issues that may arise with the possibility of default, suppose that default is governed by a stochastic intensity λ_t and that, upon default, the holder recovers a constant fraction w of face value. Then, from results in Lando [1998], the price of a $(T - t)$ -period zero-coupon bond is given under technical integrability conditions by

$$E_t \left(\exp \left(- \int_t^T (r_s + \lambda_s) ds \right) \right) + w \int_t^T E_t \left(\lambda_s \exp \left(- \int_t^s (r_u + \lambda_u) du \right) \right) ds. \quad (1.3)$$

The first term in (1.3) is the value of a claim that pays \$1 contingent on survival to maturity T , while the second term is the value of the claim that pays w at date s should the issuer default at that date, and nothing otherwise.

Both the first term and, for each s , the expectation in the second term can be computed in closed form using our extended transform. Specifically,

¹The entire class of affine term structure models is obtained as the special case of (1.1) found by setting $R = r$, $u = 0$, $v_0 = 1$, and $v_1 = 0$.

²See, for example, Jarrow, Lando, and Turnbull [1997] and Duffie and Singleton [1999].

assuming that both r_t and λ_t are affine in an *AJD* process X_t , the first expectation in (1.3) is the special case of (1.1) that is obtained by letting $R(X_t, t) = r_t + \lambda_t$, $u = 0$, $v_0 = 1$ and $v_1 = 0$. Similarly, each expectation in (1.3) of the form $E_t(\lambda_s \exp(-\int_t^s r_u + \lambda_u du))$ is obtained as a special case of (1.1) by setting $u = 0$, $R(X_t, t) = r_t + \lambda_t$, and $v_0 + v_1 \cdot X_t = \lambda_t$. Thus, using our extended transform, the pricing of defaultable zero-coupon bonds with constant fractional recovery of par reduces to the computation of a one-dimensional integral of a known function. Similar reasoning can be used to derive closed-form expressions for zero prices in environments where the default arrival intensity is affine in X , and there is “gapping” risk associated with unpredictable transitions to different credit categories (see Lando [1998] for the case of $w = 0$).

A different application of the extended transform is pursued by Piazzesi [1998] who extends the *AJD* model in order to treat term-structure models with releases of macro-economic information and with central-bank interest-rate targeting. She considers jumps at both random and at deterministic times, and allows for an intensity process and interest-rate process that have linear-quadratic dependence on the underlying state vector, extending the basic results of this paper.

1.2 Estimation of Affine Asset Pricing Models

Another useful implication of (1.1) is that, by setting $R = 0$, $v_0 = 1$, and $v_1 = 0$, we obtain a closed-form expression for the conditional characteristic function ϕ of X_T given X_t , defined by $\phi(u, X_t, t, T) = E(e^{u \cdot X_T} | X_t)$. Because knowledge of ϕ is equivalent to knowledge of the joint conditional density function of X_T , this result is useful in estimation and all other applications involving the transition densities of an *AJD*.

For instance, Singleton [1998] exploits knowledge of ϕ to derive maximum likelihood estimators for *AJDs* based on the conditional density of X_{t+1} given X_t , obtained by Fourier inversion of ϕ ,

$$f(X_{t+1} | X_t; \gamma) = \frac{1}{(2\pi)^N} \int_{\mathbb{R}^N} e^{-iu \cdot X_{t+1}} \phi(u, X_t, t, t+1) du. \quad (1.4)$$

Das [1998] exploits (1.4) for the specific case of a Poisson-Gaussian *AJD* to compute method-of-moments estimators of a model of interest rates.

Method-of-moments estimators can also be constructed directly in terms

of the conditional characteristic function. By definition, ϕ satisfies

$$E_t [e^{iu \cdot X_{t+1}} - \phi(u, X_t, t, t+1)] = 0, \quad (1.5)$$

so any measurable function of X_t is orthogonal to the “error” ($e^{iu \cdot X_{t+1}} - \phi(u, X_t, t, t+1)$). Singleton [1998] uses this fact, together with the known functional form of ϕ , to construct generalized method-of-moments estimators of the parameters governing *AJDs* and, more generally, the parameters of asset pricing models in which the state follows an *AJD*. These estimators are computationally tractable and, in some cases, achieve the same asymptotic efficiency as the maximum likelihood estimator.³

1.3 Affine Option Pricing Models

In an influential paper in the option-pricing literature, Heston [1993] showed that the risk-neutral exercise probabilities appearing in the call option pricing formulas for bonds, currencies, and equities can be computed by Fourier inversion of the conditional characteristic function, which he showed is known in closed form for his particular affine, stochastic volatility model. Building on this insight,⁴ a variety of option-pricing models have been developed for state vectors having at most a single jump type (in the asset return), and whose behavior between jumps is that of a Gaussian or “square-root” diffusion.⁵

Knowing the extended transform (1.1) in closed-form, we can extend this option pricing literature to the case of general multi-dimensional *AJD* processes with much richer dynamic inter-relations among the state variables and much richer jump distributions. For example, we provide an analytically tractable method for pricing derivatives with payoffs at a future time T of the form $(e^{b \cdot X(T)} - c)^+$, where c is a constant strike price, $b \in \mathbb{R}^n$, X

³Liu and Pan [1997] and Liu [1997] propose alternative estimation strategies that exploit the special structure of affine diffusion models.

⁴Among the many recent papers examining option prices for the case of state variables following square-root diffusions are Bakshi, Cao, and Chen [1997], Bakshi and Madan [1999], Bates [1996], Bates [1997], Chen and Scott [1993], Chernov and Ghysels [1998], Pan [1998], Scott [1996], and Scott [1997], among others.

⁵More precisely, the short-term interest rate has been assumed to be an affine function of independent square-root diffusions and, in the case of equity and currency option pricing, spot-market returns have been assumed to follow stochastic-volatility models in which volatility processes are independent “square-root” diffusions that may be correlated with the spot-market return shock.

is an *AJD*, and $y^+ \equiv \max(y, 0)$. This leads directly to pricing formulas for plain-vanilla options on currencies and equities, quanto options (such as an option on a common stock or bond struck in a different currency), options on zero-coupon bonds, caps, floors, chooser options, and other related derivatives. Furthermore, we can price payoffs of the form $(b \cdot X(T) - c)^+$ and $e^{a \cdot X(T)} b \cdot X(T)$ and this allows us to price “slope-of-the-yield-curve” options and certain Asian options.⁶

In order to visualize our approach to option pricing, consider the price p at date 0 of a call option with payoff $(e^{d \cdot X(T)} - c)^+$ at date T , for given $d \in \mathbb{R}^n$ and strike c , where X is an n -dimensional *AJD*, with a short-term interest-rate process that is itself affine in X . For any real number y and any a and b in \mathbb{R}^n , let $G_{a,b}(y)$ denote the price of a security that pays $e^{a \cdot X(T)}$ at time T in the event that $b \cdot X(T) \leq y$. As the call option is in the money when $-d \cdot X(T) \leq -\ln c$, and in that case pays $e^{d \cdot X(T)} - ce^{0 \cdot X(T)}$, we have the option priced at

$$p = G_{d,-d}(-\ln c) - c G_{0,-d}(-\ln c). \quad (1.6)$$

Thus, it is enough to be able to compute the Fourier transform $\mathcal{G}_{a,b}(\cdot)$ of $G_{a,b}(\cdot)$ (treated as a measure), defined by

$$\mathcal{G}_{a,b}(z) = \int_{-\infty}^{+\infty} e^{izy} dG_{a,b}(y),$$

for then well-known Fourier-inversion methods can be used to compute terms of the form $G_{a,b}(y)$ in (1.6).

There are many cases in which the Fourier transform $\mathcal{G}_{a,b}(\cdot)$ of $G_{a,b}(\cdot)$ can be computed explicitly. We extend the range of solutions for the transform $\mathcal{G}_{a,b}(\cdot)$ from those already in the literature to include the entire class of *AJDs* by noting that $\mathcal{G}_{a,b}(z)$ is given by (1.1), for the complex coefficient vector $u = a + izb$, with $v_0 = 1$ and $v_1 = 0$. This, because of the affine structure, implies under regularity conditions, that

$$\mathcal{G}_{a,b}(z) = e^{\alpha^{(0)} + \beta^{(0)} \cdot X(0)}, \quad (1.7)$$

⁶In a complementary analysis of derivative security valuation, Bakshi and Madan [1999] show that knowledge of the special case of (1.1) with $v_0 + v_1 \cdot X_T = 1$ is sufficient to recover the prices of standard call options, but they do not provide explicit guidance as to how to compute this transform. Their applications to Asian and other options presumes that the state vector follows square-root or Heston-like stochastic-volatility models for which the relevant transforms had already been known in closed form.

where α and β solve known, complex-valued ordinary differential equations (*ODEs*) with boundary conditions at T determined by z . In some cases, these *ODEs* have explicit solutions. These include independent square-root diffusion models for the short-rate process, as in Chen and Scott [1995], and the stochastic-volatility models of asset prices studied by Bates [1997] and Bakshi, Cao, and Chen [1997]. Using our *ODE*-based approach, we derive other explicit examples, for instance stochastic-volatility models with correlated jumps in both returns and volatility. In other cases, one can easily solve the *ODEs* for α and β numerically, even for high-dimensional applications.

Similar transform analysis provides a price for an option with a payoff of the form $(d \cdot X_T - c)^+$, again for the general *AJD* setting. For this case, we provide in Appendix E an equally tractable method for computing the Fourier transform of $\tilde{G}_{a,b,a}(\cdot)$, where $\tilde{G}_{a,b,a}(y)$ is the price of a security that pays $e^{d \cdot X(T)} a \cdot X(T)$ at T in the event that $b \cdot X(T) \leq y$. This transform is again of the form (1.1), now with $v_1 = a$. Given this transform, we can invert to obtain $\tilde{G}_{a,b,a}(y)$ and the option price p' given by

$$p' = \tilde{G}_{a,-a,0}(-\ln c) - c G_{0,-a}(-\ln c). \quad (1.8)$$

As shown in Appendix E, these results can be used to price slope-of-the-yield-curve options and certain Asian options.

Our motivation for studying the general *AJD* setting is largely empirical. The *AJD* model takes the elements of the drift vector, “instantaneous” covariance matrix, and jump measure of X to be affine functions of X . This allows for conditional variances that depend on all of the state variables (unlike the Gaussian model), and for a variety of patterns of cross-correlations among the elements of the state vector (unlike the case of independent square-root diffusions). Dai and Singleton [1999], for instance, found that both time-varying conditional variances and negatively correlated state variables were essential ingredients to explaining the historical behavior of term structures of U.S. interest rates.

Furthermore, for the case of equity options, Bates [1997] and Bakshi, Cao, and Chen [1997] found that their affine stochastic-volatility models did not fully explain historical changes in the volatility smiles implied by *S&P500* index options. Within the affine family of models, one potential explanation for their findings is that they unnecessarily restricted the correlations between the state variables driving returns and volatility. Using the classification

scheme for affine models found in Dai and Singleton [1999], one may nest these previous stochastic-volatility specifications within an *AJD* model with the same number of state variables that allows for potentially much richer correlation among the return and volatility factors.

The empirical studies of Bates [1997] and Bakshi, Cao, and Chen [1997] also motivate, in part, our focus on multivariate jump processes. They concluded that their stochastic-volatility models (with jumps in spot-market returns only) do not allow for a degree of volatility of volatility sufficient to explain the substantial “smirk” in the implied volatilities of index option prices. Both papers conjectured that jumps in volatility, as well as in returns, may be necessary to explain option-volatility smirks. Our *AJD* setting allows for correlated jumps in both volatility and price. Jumps may be correlated because their amplitudes are drawn from correlated distributions, or because of correlation in the jump times. (The jump times may be simultaneous, or have correlated stochastic arrival intensities.)

In order to illustrate our approach, we provide an example of the pricing of plain-vanilla calls on the *S&P500* index. A cross-section of option prices for a given day are used to calibrate *AJDs* with simultaneous jumps in both returns and volatility. Then we compare the implied-volatility smiles to those observed in the market on the chosen day. In this manner we provide some preliminary evidence on the potential role of jumps in volatility for resolving the volatility puzzles identified by Bates [1997] and Bakshi, Cao, and Chen [1997].

The remainder of this paper is organized as follows. Section 2 reviews the class of affine jump-diffusions, and shows how to compute some relevant transforms, and how to invert them. Section 3 presents our basic option-pricing results. The example of the pricing of plain-vanilla calls on the *S&P500* index is presented in Section 4. Additional appendices provide various technical results and extensions.

2 Transform Analysis for *AJD* State-Vectors

This section presents the *AJD* state-process model and the basic transform calculations that will later be useful in option pricing. Technical details are presented in Appendix A.

2.1 The Affine Jump-Diffusion

We fix a probability space (Ω, \mathcal{F}, P) and an information filtration⁷ (\mathcal{F}_t) , and suppose that X is a Markov process in some state space $D \subset \mathbb{R}^n$, solving the stochastic differential equation

$$dX_t = \mu(X_t) dt + \sigma(X_t) dW_t + dZ_t, \quad (2.1)$$

where W is a Standard Brownian motion in \mathbb{R}^n ; $\mu : D \rightarrow \mathbb{R}^n$, $\sigma : D \rightarrow \mathbb{R}^{n \times n}$, and Z is a pure jump process whose jumps have a fixed probability distribution ν on \mathbb{R}^n and arrive with intensity $\{\lambda(X_t) : t \geq 0\}$, for some $\lambda : D \rightarrow [0, \infty)$. For notational convenience, we assume that X_0 is “known” (has a trivial distribution). Appendices provide additional technical details, as well as generalizations to multiple jump-types with different arrival intensities, and to time-dependent $(\mu, \sigma, \lambda, \nu)$.

We impose an “affine” structure on μ , $\sigma\sigma^\top$, and λ , in that all of these are assumed to be affine. In order for X to be well defined, there are joint restrictions on $(D, \mu, \sigma, \lambda, \nu)$. These restrictions are discussed in Duffie and Kan [1996] and Dai and Singleton [1999], and are reviewed briefly in Appendix A.

2.2 Transforms

First, we show that the Fourier transform of X_t and of certain related random variables is known in closed form up to the solution of an *ODE*. Then, we show how the distribution of X_t and the prices of options can be recovered by inverting this transform. Throughout this section, we specialize to the case of $v_0 = 1$ and $v_1 = 0$ in (1.1), and put our treatment of the extended transform in Appendix E.

We fix an affine discount-rate function $R : D \rightarrow \mathbb{R}$. The affine dependence of μ , $\sigma\sigma^\top$, λ , and R are determined by coefficients (K, H, l, ρ) defined by:

- $\mu(x) = K_0 + K_1 x$, for $K = (K_0, K_1) \in \mathbb{R}^n \times \mathbb{R}^{n \times n}$.
- $(\sigma(x)\sigma(x)^\top)_{ij} = (H_0)_{ij} + (H_1)_{ij} \cdot x$, for $H = (H_0, H_1) \in \mathbb{R}^{n \times n} \times \mathbb{R}^{n \times n \times n}$.
- $\lambda(x) = l_0 + l_1 \cdot x$, for $l = (l_0, l_1) \in \mathbb{R} \times \mathbb{R}^n$.
- $R(x) = \rho_0 + \rho_1 \cdot x$, for $\rho = (\rho_0, \rho_1) \in \mathbb{R} \times \mathbb{R}^n$.

⁷For technical details, see Appendix A.

For $c \in \mathbb{C}^n$, the set of n -tuples of complex numbers, we let $\theta(c) = \int_{\mathbb{R}^n} \exp(c \cdot z) d\nu(z)$ whenever the integral is well defined. This “jump transform” determines the jump-size distribution. The subsequent analysis suggests a practical advantage of choosing jump distributions with an explicitly known or easily computed jump transform θ .

The “coefficients” (K, H, l, θ) of X completely determine its distribution, given an initial condition $X(0)$. A “characteristic” $\chi = (K, H, l, \theta, \rho)$ captures both the distribution of X as well as the effects of any discounting, and determines a transform $\psi^\chi : \mathbb{C}^n \times D \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{C}$ of X_T conditional on \mathcal{F}_t , when well defined at $t \leq T$, by

$$\psi^\chi(u, X_t, t, T) = E^\chi \left(\exp \left(- \int_t^T R(X_s) ds \right) e^{u \cdot X(T)} \mid \mathcal{F}_t \right), \quad (2.2)$$

where E^χ denotes expectation under the distribution of X determined by χ . Here, ψ^χ differs from the familiar (conditional) characteristic function of the distribution of X_T because of the discounting at rate $R(X_t)$.

The key insight underlying our applications is that, under technical regularity conditions given in Appendix B, Proposition 1,

$$\psi^\chi(u, x, t, T) = e^{\alpha(t) + \beta(t) \cdot x}, \quad (2.3)$$

where β and α satisfy the complex-valued ODEs⁸

$$\dot{\beta}_t = \rho_1 - K_1^\top \beta_t - \frac{1}{2} \beta_t^\top H_1 \beta_t - l_1(\theta(\beta_t) - 1), \quad (2.4)$$

$$\dot{\alpha}_t = \rho_0 - K_0 \cdot \beta_t - \frac{1}{2} \beta_t^\top H_0 \beta_t - l_0(\theta(\beta_t) - 1), \quad (2.5)$$

with boundary conditions $\beta(T) = u$ and $\alpha(T) = 0$. The ODE (2.4)-(2.5) is easily conjectured from an application of Ito’s Formula to the candidate form (2.3) of ψ^χ .

Anticipating the application to option pricing, for each given $(d, c, T) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}_+$, our next goal is to compute (when well defined, as under conditions in Appendix B, Proposition 3) the “expected present value”

$$C(d, c, T, \chi) = E^\chi \left(\exp \left(- \int_0^T R(X_s) ds \right) (e^{d \cdot X(T)} - c)^+ \right). \quad (2.6)$$

⁸Here, $c^\top H_1 c$ denotes the vector in \mathbb{C}^n with k -th element $\sum_{i,j} c_i (H_1)_{ijk} c_j$.

We have

$$\begin{aligned} C(d, c, T, \chi) &= E^X \left(\exp \left(- \int_0^T R(X_s) ds \right) (e^{d \cdot X(T)} - c) \mathbf{1}_{d \cdot X(T) \geq \ln(c)} \right) \\ &= G_{d, -d}(-\ln(c); X_0, T, \chi) - c G_{0, -d}(-\ln(c); X_0, T, \chi), \end{aligned} \quad (2.7)$$

where, given some $(x, T, a, b) \in D \times [0, \infty) \times \mathbb{R}^n \times \mathbb{R}^n$, $G_{a,b}(\cdot; x, T, \chi) : \mathbb{R} \rightarrow \mathbb{R}_+$ is defined (under technical conditions provided in Appendix B) by

$$G_{a,b}(y; X_0, T, \chi) = E^X \left(\exp \left(- \int_0^T R(X_s) ds \right) e^{a \cdot X(T)} \mathbf{1}_{b \cdot X(T) \leq y} \right). \quad (2.8)$$

The Fourier-Stieltjes transform $\hat{G}_{a,b}(\cdot; X_0, T, \chi)$ of $G_{a,b}(\cdot; X_0, T, \chi)$, if well defined, is given by

$$\begin{aligned} \hat{G}_{a,b}(v; X_0, T, \chi) &= \int_{\mathbb{R}} e^{ivy} dG_{a,b}(y; X_0, T, \chi) \\ &= E^X \left(\exp \left(- \int_0^T R(X_s) ds \right) \exp[(a + ivb) \cdot X_T] \right) \\ &= \psi^X(a + ivb, X_0, 0, T). \end{aligned}$$

We may now extend the Lévy inversion formula⁹ (from the typical case of a proper cumulative distribution function) to obtain, under a technical integrability condition given in Appendix C, Proposition 2,

$$G_{a,b}(y; X_0, T, \chi) = \frac{\psi^X(a, X_0, 0, T)}{2} - \frac{1}{\pi} \int_0^\infty \frac{\text{Im}[\psi^X(a + ivb, X_0, 0, T)e^{-ivy}]}{v} dv, \quad (2.9)$$

where $\text{Im}(c)$ denotes the imaginary part of $c \in \mathbb{C}$. For $R = 0$, this gives the probability distribution function of $b \cdot X_T$. The associated transition density of X is obtained by differentiation of $G_{a,b}$. More generally, this provides the transition function of X with “killing” at rate¹⁰ R . Piazzesi [1998] extends this analysis to allow a limited degree of quadratic dependence of the short rate on the state vector.

⁹See, for example, Gil-Pelaez [1951] and Williams [1991] for a treatment of the Lévy inversion formula.

¹⁰A negative R is sometimes called a “creation” rate in Markov-process theory.

3 Option Pricing Theory

This section applies our basic transform analysis to the pricing of options. In all cases, we assume that the price process S of the asset underlying the option is of the exponential-affine form $S_t = e^{a(t)+b(t)\cdot X(t)}$. This is the case for many applications in affine settings, including underlying assets that are equities, currencies, and zero-coupon bonds.

Two traditional formulations¹¹ of the asset-pricing problem are:

1. Model the “risk-neutral” behavior of X under an equivalent martingale measure Q . That is, take X to be an affine jump-diffusion under Q with given characteristic χ_Q . Then apply (2.7) and (2.9).
2. Model the behavior of X as an affine jump-diffusion under the actual (that is, the “data-generating”) measure P . If one then supposes that the state-price density (also known as the “pricing kernel” or “marginal-rate-of-substitution” process) is an exponential-affine form in X , then X is also an affine jump-diffusion under Q , and one can either:
 - (a) calculate, as in Appendix D, the implied equivalent martingale measure Q and associated characteristic χ_Q of X under Q , and proceed as in the first alternative above, or
 - (b) simply apply the definition of the state-price density, which determines the price of an option directly in terms of $G_{a,b}$, computed using our transform analysis. This alternative is sketched in Section 3.2 below.

Of course the two approaches are consistent, and indeed the second formulation implies the first, as indicated. The second approach is more complete, and would be indicated for empirical time-series applications, for which the

¹¹A popular variant developed in a Gaussian setting by Jamshidian [1989]. In a setting in which X is an affine jump-diffusion under the equivalent martingale measure Q , one normalizes the underlying exponential-affine asset price by the price of a zero-coupon bond maturing on the option expiration date T . Then, in the new numeraire, the short-rate process is of course zero, and there is a new equivalent martingale measure $Q(T)$, often called the “forward measure,” under which prices are exponential affine. Application of Girsanov’s Theorem uncovers new affine behavior for the underlying state process X under $Q(T)$, and one can proceed as before. The change-of-measure calculations for this approach can be found in Appendix D.

“actual” distribution of the state process X as well as the parameters determining risk-premia must be specified and estimated, as in Pan [1998].

Applications of these approaches to call-option pricing are briefly sketched in the next two sub-sections. Other derivative pricing applications are provided in Section 3.3.

3.1 Risk-Neutral Pricing

Here, we take Q to be an equivalent martingale measure associated with a short-term interest rate process defined by $R(X_t) = \rho_0 + \rho_1 \cdot X_t$. This means that the market value at time t of any contingent claim that pays an \mathcal{F}_T -measurable random variable V at time T is, by definition,

$$E^Q \left(\exp \left(- \int_t^T R(X_s) ds \right) V \mid \mathcal{F}_t \right), \quad (3.1)$$

where, under Q , the state vector X is assumed to be an *AJD* with coefficients $(K^Q, H^Q, l^Q, \theta^Q)$. The relevant characteristic for risk-neutral pricing is then $\chi_Q = (K^Q, H^Q, l^Q, \theta^Q, \rho)$. It need not be the case that markets are complete. The existence of some equivalent martingale measure and the absence of arbitrage are in any case essentially equivalent properties, under technical conditions, as pointed out by Harrison and Kreps [1979]. For recent technical conditions, see for example Delbaen and Schachermayer [1994].

We let S denote the price process for the security underlying the option, and suppose for simplicity¹² that $\ln S_t = X_t^{(i)}$, an element of the state vector $X = (X^{(1)}, \dots, X^{(n)})$. Other components of the state process X may jointly specify the arrival intensity of jumps, the behavior of stochastic volatility, the behavior of other asset returns, interest-rate behavior, and so on. The given asset is assumed to have a dividend-yield process $\{\zeta(X_t) : t \geq 0\}$ defined by

$$\zeta(x) = q_0 + q_1 \cdot x, \quad (3.2)$$

for given $q_0 \in \mathbb{R}$ and $q_1 \in \mathbb{R}^n$. For example, if the asset is a foreign currency, then $\zeta(X_t)$ is the foreign short-term interest rate.

¹²The general case of $S_t = \exp(a_t + b_t \cdot X_t)$ can be similarly treated. Possibly after some innocuous affine change of variables in the state vector, possibly involving time dependencies in the characteristic χ , we can always reduce to the assumed case.

Because Q is an equivalent martingale measure, the coefficients $K_i^Q = ((K_0^Q)_i, (K_1^Q)_i)$ determining¹³ the “risk-neutral” drift of $X^{(i)} = \ln S$ are given by

$$(K_0^Q)_i = \rho_0 - q_0 - \frac{1}{2}(H_0^Q)_{ii} - l_0^Q (\theta^Q(\epsilon(i)) - 1) \quad (3.3)$$

$$(K_1^Q)_i = \rho_1 - q_1 - \frac{1}{2}(H_1^Q)_{ii} - l_1^Q (\theta^Q(\epsilon(i)) - 1), \quad (3.4)$$

where $\epsilon(i) \in \mathbb{R}^n$ has 1 as its i -th component, and any other component equal to 0.

Unless other security price processes are specified, the risk-neutral characteristic χ_Q is otherwise unrestricted by arbitrage considerations. There are analogous no-arbitrage restrictions on χ_Q for each additional specified security price process of the form $e^{a+b \cdot X(t)}$.

By the definition of an equivalent martingale measure and the results of Section 2.2, a plain-vanilla European call option with expiration time T and strike c has a price p at time 0 which, under the regularity in Appendix B, is given by (2.7) to be

$$p = G_{\epsilon(i), -\epsilon(i)}(-\ln(c); X_0, T, \chi_Q) - c G_{0, -\epsilon(i)}(-\ln(c); X_0, T, \chi_Q). \quad (3.5)$$

This extends Heston [1993], Bates [1996], Scott [1997], Bates [1997], Bakshi and Madan [1999], and Bakshi, Cao, and Chen [1997].

3.2 State-Price Density

Suppose the state vector X is an affine jump-diffusion with coefficients (K, H, l, θ) under the actual (data-generating) measure P . Let ξ be an (\mathcal{F}_t) -adapted

¹³Under (3.3)-(3.4), we have

$$\begin{aligned} S_t - S_0 &= \int_0^t S_u [R(X_u) - \zeta(X_u)] du + \int_0^t S_u \sigma^{(i)}(X_u)^\top dW_u^Q \\ &\quad + \sum_{0 < u \leq t} S_{u-} \left(\exp(\Delta X_u^{(i)}) - 1 \right) - \int_0^t S_u (\theta^Q(\epsilon(i)) - 1) (l_0^Q + l_1^Q \cdot X_u) du, \end{aligned}$$

where W^Q is an (\mathcal{F}_t) -standard Brownian motion in \mathbb{R}^n under Q . As the sum of the last 3 terms is a local Q -martingale, this indeed implies consistency with the definition of an equivalent martingale measure. (Here, $\Delta X(t) = X(t) - X(t-)$ denotes the jump of X at t .)

“state-price density,” defined by the property that the market value at time t of any security that pays an \mathcal{F}_T -measurable random variable V at time T is given by

$$\frac{1}{\xi(t)} E(V\xi(T) \mid \mathcal{F}_t).$$

We assume that $\xi_t = e^{a(t)+b(t)\cdot X(t)}$, for some bounded measurable $a : [0, \infty) \rightarrow \mathbb{R}$ and $b : [0, \infty) \rightarrow \mathbb{R}^n$. Without loss of generality, we take it that $\xi(0) = 1$.

Suppose the price of a given underlying security at time T is $e^{d\cdot X(T)}$, for some $d \in \mathbb{R}^n$. By the definition of a state-price density, a plain-vanilla European call option struck at c with exercise date T has a price at time 0 of

$$p = E[e^{a(T)+b(T)\cdot X(T)}(e^{d\cdot X(T)} - c)^+].$$

This leaves the option price

$$p = e^{a(T)} G_{b(T)+d, -d}(-\ln c; X_0, T, \chi^0) - ce^{a(T)} G_{b(T), -d}(-\ln c; X_0, T, \chi^0),$$

where $\chi^0 = (K, H, l, \theta, 0)$. (One notes that the short-rate process plays no role beyond that already captured by the state-price density.)

As mentioned at the beginning of this section, and detailed in Appendix E, an alternative is to translate the option-pricing problem to a “risk-neutral” setting.

3.3 Other Option-Pricing Applications

This section develops as illustrative examples several additional applications to option pricing. For convenience, we adopt the risk-neutral pricing formulation. That is, we suppose that the short rate is given by $R(X)$, where R is affine, and X is an affine jump-diffusion under an equivalent martingale measure Q . The associated characteristic χ_Q is fixed. While we treat the case of call options, put options can be treated by the same method, or by put-call parity.

3.3.1 Bond Derivatives

Consider a call option, struck at c with exercise date T , on a zero-coupon bond maturing at time $s > T$. Let $\Lambda(T, s)$ denote the time- T market price

of the underlying bond. From Duffie and Kan [1996], under the regularity conditions given in Appendix B,

$$\Lambda(T, s) = \exp(\alpha(T, s, 0) + \beta(T, s, 0) \cdot X_T),$$

where $\beta(T, s, 0)$ and $\alpha(T, s, 0)$ are defined by (B.3) and (B.4). At time T , the option pays

$$(\Lambda(T, s) - c)^+ = (e^{\alpha(T, s, 0) + \beta(T, s, 0) \cdot X(T)} - c)^+ \quad (3.6)$$

$$= e^{\alpha(T, s, 0)} (e^{\beta(T, s, 0) \cdot X(T)} - e^{-\alpha(T, s, 0)} c)^+. \quad (3.7)$$

The value of the bond option can therefore be obtained from (2.7) and (2.9). The same approach applies to caps and floors, which are simply portfolios of zero-coupon bond options with payment in arrears, as reviewed in Appendix G. This extends the results of Chen and Scott [1995] and Scott [1996]. Chacko and Das [1998] work out the valuation of asian interest-rate options for a large class of affine models. They provide numerical examples based on a multi-factor Cox-Ingersoll-Ross state vector.

3.3.2 Quantos

Consider a quanto of exercise date T and strike c on an underlying asset with price process $S = \exp(X^{(i)})$. The time- T payoff of the quanto is $(S_T M(X_T) - c)^+$, where $M(x) = e^{m \cdot x}$, for some $m \in \mathbb{R}^n$. The quanto scaling $M(X_T)$ could, for example, be the price at time T of a given asset, or the exchange rate between two currencies. The initial market value of the quanto option is then

$$G_{m+\epsilon(i), -\epsilon(i)}(-\ln(c); x, T, \chi_Q) - c G_{0, -\epsilon(i)}(-\ln(c); x, T, \chi_Q).$$

An alternative form of the quanto option pays $M(X_T)(S_T - c)^+$ at T , and has the price

$$G_{m+\epsilon(i), -\epsilon(i)}(-\ln(c); x, T, \chi_Q) - c G_{m, -\epsilon(i)}(-\ln(c); x, T, \chi_Q).$$

3.3.3 Foreign Bond Options

Let $\exp(X^{(i)})$ be a foreign-exchange rate, $R(X)$ be the domestic short interest rate, and $\zeta(X)$ be the foreign short rate, for affine ζ . Consider a

foreign zero-coupon bond maturing at time s , whose payoff at maturity, in domestic currency, is therefore $\exp(X^{(i)})$. The risk-neutral characteristic χ_Q is restricted by (3.3)-(3.4). From Proposition 1 in Appendix B, the domestic price at time t of the foreign bond is

$$\Lambda^f(t, s) = \exp(\alpha(t, s, \epsilon(i)) + \beta(t, s, \epsilon(i)) \cdot X_t).$$

We now consider an option on this bond with exercise date $T < s$ and domestic strike price c on the foreign s -year zero-coupon bond, paying $(\Lambda^f(T, s) - c)^+$ at time T , in domestic currency. The initial market value of this option can therefore be obtained as for a domestic bond option.

3.3.4 Chooser Options

Let $S^{(i)} = \exp(X^{(i)})$ and $S^{(j)} = \exp(X^{(j)})$ be two security price processes. An exchange, or “chooser,” option with exercise date T , pays $\max(S_T^{(i)}, S_T^{(j)})$. Depending on their respective dividend payout rates, the risk-neutral characteristic χ_Q is restricted by (3.3)-(3.4), applied to both i and j . The initial market value of this option is

$$G_{\epsilon(i), \epsilon(i) - \epsilon(j)}(0; x, T, \chi_Q) + G_{\epsilon(j), 0}(0, x, T, \chi_Q) - G_{\epsilon(j), \epsilon(j) - \epsilon(i)}(0; x, T, \chi_Q).$$

4 A “Double-Jump” Illustrative Model

As an illustration of the methodology, this section provides explicit transforms for a 2-dimensional affine jump-diffusion model. We suppose that S is the price process, strictly positive, of a security that pays dividends at a constant proportional rate $\bar{\zeta}$, and we let $Y = \ln(S)$. The state process is $X = (Y, V)^\top$, where V is the volatility process.

We suppose for simplicity that the short rate is a constant r , and that there exists an equivalent martingale measure Q under which¹⁴

$$d \begin{pmatrix} Y_t \\ V_t \end{pmatrix} = \begin{pmatrix} r - \bar{\zeta} - \bar{\lambda}\bar{\mu} - \frac{1}{2}V_t \\ \kappa_v(\bar{v} - V_t) \end{pmatrix} dt + \sqrt{V_t} \begin{pmatrix} 1 & 0 \\ \bar{\rho}\sigma_v & \sqrt{1 - \bar{\rho}^2}\sigma_v \end{pmatrix} dW_t^Q + dZ_t, \quad (4.1)$$

¹⁴Unless otherwise stated, the distributional properties of (Y, V) described in this section are in a “risk-neutral” sense, that is, under Q .

where W^Q is an (\mathcal{F}_t) -standard Brownian motion under Q in \mathbb{R}^2 , and Z is a pure jump process in \mathbb{R}^2 with constant mean jump-arrival rate $\bar{\lambda}$, whose bivariate jump-size distribution ν has the transform θ . A flexible range of distributions of jumps can be explored through the specification of θ . The risk-neutral coefficient restriction (3.3) is satisfied if and only if $\bar{\mu} = \theta(1, 0) - 1$.

Before we move on to special examples, we lay out the formulation for option pricing as a straightforward application of earlier results in the paper. At time t , the transform¹⁵ ψ of the log-price state variable Y_T can be calculated using the *ODE* approach in (2.5) as:

$$\psi(u, (y, v), t, T) = \exp(\alpha(T - t, u) + uy + \beta(T - t, u)v), \quad (4.2)$$

where, letting $b = \sigma_v \bar{\rho} u - \kappa_v$, $a = u(1 - u)$, and¹⁶ $\gamma = \sqrt{b^2 + a\sigma_v^2}$, we have

$$\beta(\tau, u) = -\frac{a(1 - e^{-\gamma\tau})}{2\gamma - (\gamma + b)(1 - e^{-\gamma\tau})}, \quad (4.3)$$

$$\alpha(\tau, u) = \alpha_0(\tau, u) - \bar{\lambda}\tau(1 + \bar{\mu}u) + \bar{\lambda} \int_0^\tau \theta(u, \beta(t, u)) dt, \quad (4.4)$$

where¹⁷

$$\alpha_0(\tau, u) = -r\tau + (r - \bar{\zeta})u\tau - \kappa_v \bar{v} \left(\frac{\gamma + b}{\sigma_v^2} \tau + \frac{2}{\sigma_v^2} \ln \left[1 - \frac{\gamma + b}{2\gamma} (1 - e^{-\gamma\tau}) \right] \right),$$

and where the term $\int_0^\tau \theta(u, \beta(t, u)) dt$ depends on the specific formulation of bivariate jump transform $\theta(\cdot, \cdot)$.

4.1 A Concrete Example

As a concrete example, consider the jump transform θ defined by

$$\theta(c_1, c_2) = \bar{\lambda}^{-1} (\lambda^y \theta^y(c_1) + \lambda^v \theta^v(c_2) + \lambda^c \theta^c(c_1, c_2)), \quad (4.5)$$

¹⁵That is, $\psi(u, (y, v), t, T) = \psi^\chi((u, 0)', (y, v)', t, T)$, where χ is the characteristic under Q of X associated with the short rate defined by $(\rho_0, \rho_1) = (r, 0)$.

¹⁶To be more precise, $\gamma = |\gamma^2|^{1/2} \exp\left(\frac{i \arg(\gamma^2)}{2}\right)$, where $\gamma^2 = b^2 + a\sigma_v^2$. Note that for any $z \in \mathbb{C}$, $\arg(z)$ is defined such that $z = |z| \exp(i \arg(z))$, with $-\pi < \arg(z) < \pi$.

¹⁷For any $z \in \mathbb{C}$, $\ln(z) = \ln|z| + i \arg(z)$.

where $\bar{\lambda} = \lambda^y + \lambda^v + \lambda^c$, and where

$$\begin{aligned}\theta^y(c) &= \exp\left(\mu_y c + \frac{1}{2}\sigma_y^2 c^2\right), \\ \theta^v(c) &= \frac{1}{1 - \mu_v c}, \\ \theta^c(c_1, c_2) &= \frac{\exp\left(\mu_{c,y} c_1 + \frac{1}{2}\sigma_{c,y}^2 c_1^2\right)}{1 - \mu_{c,v} c_2 - \rho_J \mu_{c,v} c_1}.\end{aligned}$$

What we incorporate in this example is in fact three types of jumps:

- Jumps in Y , with arrival intensity λ^y and normally distributed jump size with mean μ_y and variance σ_y^2 ,
- Jumps in V , with arrival intensity λ^v and exponentially distributed jump size with mean μ_v ,
- Simultaneous correlated jumps in Y and V , with arrival intensity λ^c . The marginal distribution of the jump size in V is exponential with mean $\mu_{c,v}$. Conditional on a realization, say z_v , of the jump size in V , the jump size in Y is normally distributed with mean $\mu_{c,y} + \rho_J z_v$, and variance $\sigma_{c,y}^2$.

In Bakshi, Cao, and Chen [1997] and Bates [1997], the *SVJ-Y* model, defined by $\lambda^v = \lambda^c = 0$, was studied using cross sections of options data to fit the “volatility smirk.” They find that allowing for negative jumps in Y is useful insofar as it increases the skewness of the distribution of Y_T , but that this does not generate the level of skewness implied by the volatility smirk observed in market data. They call for a model with jumps in volatility. Using this concrete “double-jump” example (4.5), we can address this issue, and provide some insights into what a richer specification of jumps may imply.

Before leaving this section to explore the implications of jumps for “volatility smiles,” we provide explicit option pricing through the transform formula (4.2), by exploiting the bivariate jump transform θ specified in (4.5). We have

$$\int_0^\tau \theta(u, \beta(t, u)) dt = \bar{\lambda}^{-1} (\lambda^y f^y(u, \tau) + \lambda^v f^v(u, \tau) + \lambda^c f^c(u, \tau)),$$

where

$$\begin{aligned}
f^y(u, \tau) &= \tau \exp\left(\mu_y u + \frac{1}{2}\sigma_y^2 u^2\right), \\
f^v(u, \tau) &= \frac{\gamma - b}{\gamma - b + \mu_v a} \tau - \frac{2\mu_v a}{\gamma^2 - (b - \mu_v a)^2} \ln\left(1 - \frac{(\gamma + b) - \mu_v a}{2\gamma}(1 - e^{-\gamma\tau})\right), \\
f^c(u, \tau) &= \exp\left(\mu_{c,y} u + \sigma_{c,y}^2 \frac{u^2}{2}\right) d,
\end{aligned}$$

where $a = u(1 - u)$, $b = \sigma_v \bar{\rho} u - \kappa_v$, $c = 1 - \rho_J \mu_{c,v} u$, and

$$d = \frac{\gamma - b}{(\gamma - b)c + \mu_{c,v} a} \tau - \frac{2\mu_{c,v} a}{(\gamma c)^2 - (bc - \mu_{c,v} a)^2} \ln\left[1 - \frac{(\gamma + b)c - \mu_{c,v} a}{2\gamma c}(1 - e^{-\gamma\tau})\right].$$

4.2 Jump Impact on “Volatility Smiles”

As an illustration of the implications of jumps for the volatility smirk, we first select three special cases of the “double-jump” example just specified,

SV: Stochastic volatility model with no jumps, obtained by letting $\bar{\lambda} = 0$.

SVJ-Y: Stochastic volatility model with jumps in price only, obtained by letting $\lambda^y \neq 0$, and $\lambda^v = \lambda^c = 0$.

SVJJ: Stochastic volatility with simultaneous and correlated jumps in price and volatility, obtained by letting $\lambda^c \neq 0$ and $\lambda^y = \lambda^v = 0$.

In order to choose plausible values for the parameters governing these three special cases, we calibrated these three benchmark models to the actual “market-implied” smiles on November 2, 1993, plotted in Figure 1.¹⁸ For each model, calibration was done by minimizing (by choice of the unrestricted parameters) the mean-squared pricing error (MSE), defined as the simple average of the squared differences between the observed and the modeled option prices across all strikes and maturities. The risk-free rate r is assumed to be 3.19%, and the dividend yield $\bar{\zeta}$ is assumed to be zero.

¹⁸The options data are downloaded from the home page of Yacine Ait-Sahalia. There is a total of 87 options with maturities (times to exercise date) ranging from 17 days to 318 days, and strikes prices ranging from 0.74 to 1.17 times the underlying futures price.




Figure 1: “Smile curves” implied by S&P 500 Index options of 6 different maturities. Option prices are obtained from market data of November 2, 1993.

Table 1 displays the calibrated parameters of the models. Interestingly, for this particular day, we see that adding a jump in volatility to the SVJ - Y model, leading to the model $SVJJ$ model, causes a substantial decline in the level of the parameter σ_v determining the volatility of the diffusion component of volatility. Thus, the volatility puzzle identified by Bates and Bakshi, Cao, and Chen, namely that the volatility of volatility in the diffusion component of V seems too high, is potentially explained by allowing for jumps in volatility. At the same time, the return jump variance σ_y^2 declines to approximately zero as we replace the SVJ - Y model with the $SVJJ$ model. A consequence of this is that the jump sizes of Y and of V are nearly perfectly anti-correlated. This jump distribution reinforces the negative skew typically




Figure 2: “Smile curves” implied by S&P 500 Index options with 17 days to maturity. Diamonds are observed Black-Scholes implied volatilities on November 2, 1993. *SV* is the Stochastic Volatility Model, *SVJ-Y* is the Stochastic Volatility Model with Jumps in Returns, and *SVJJ* is the Stochastic Volatility Model with Simultaneous and Correlated Jumps in Returns and Volatility. Model parameters were calibrated with options data of November 2, 1993.

found in estimation of the *SV* model for these data,¹⁹ as jumps down in return are associated with simultaneous jumps up in volatility.

In order to gain additional insight into the relative fit of the models to the option data used in our calibration, Figures 2 and 3 show the volatility smiles for the shortest (17-day) and longest (318-day) maturity options. For both maturities, there is a notable improvement of fit with the inclusion of jumps. Furthermore, the addition of a jump in volatility leads to a more pronounced smirk at both maturities and one that, based on the relative values of the

¹⁹In addition to the “calibration” results in the literature, see the time-series results of Chernov and Ghysels [1998] and Pan [1998]. For related work, see Poteshman [1998] and Benzoni [1998].

Table 1: Fitted Parameter Values for SV , $SVJ-Y$, and $SVJJ$ Models

	SV	$SVJ-Y$	$SVJJ$
$\bar{\rho}$	-0.70	-0.79	-0.82
\bar{v}	0.019	0.014	0.008
κ_v	6.21	3.99	3.46
σ_v	0.61	0.27	0.14
λ^c	0	0.11	0.47
$\bar{\mu}$	n/a	-0.12	-0.10
σ_y	n/a	0.15	0.0001
μ_v	n/a	0	0.05
ρ_J	n/a	n/a	-0.38
$\sqrt{V_0}$	10.1%	9.4%	8.7%
MSE	0.0124	0.0071	0.0041

The parameters are estimated by minimizing mean squared errors (MSE). A total of 87 options, observed on November 2, 1993, are used. $\sqrt{V_0}$ is the estimated value of stochastic volatility on the sample day. The risk-free rate is assumed to be fixed at $r = 3.19\%$, and the dividend yield at $\bar{\zeta} = 0$. From “risk neutrality,” $\bar{\mu} = \theta(1, 0) - 1$.

MSE in Table 1, produces a better overall fit on this day.

Next, we go beyond this fitting exercise, and study how the introduction of a volatility jump component to the SV and $SVJ-Y$ models might affect the “volatility smile,” and how correlation between jumps in Y and V affects the “volatility smirk.” We investigate the following three additional special cases:

1. The $SVJ-V$ model: We extend the fitted SV model by letting $\lambda^v = 0.1$ and $\lambda^y = \lambda^c = 0$. We measure the degree of contribution of the jump component of volatility by the fraction $\lambda^v \mu_v^2 / (\sigma_v^2 V_0 + \lambda^v \mu_v^2)$ of the initial instantaneous variance of the volatility process V that is due to the jump component. By varying μ_v , the mean of the volatility jumps, three levels of this volatility “jumpiness” fraction are considered: 0, 15%, and 30%. For each case, the time-0 instantaneous drift, variance, and correlation are fixed to those implied by the fitted SV model by varying σ_v , \bar{v} , and $\bar{\rho}$.




Figure 3: “Smile curves” implied by S&P 500 Index options with 318 days to maturity. Hexagrams are observed implied volatility of November 2, 1993. SV is the Stochastic Volatility Model, $SVJ-Y$ is the Stochastic Volatility Model with Jumps in Returns, and $SVJJ$ is the Stochastic Volatility Model with Simultaneous and Correlated Jumps in Returns and Volatility. Model parameters were calibrated with options data of November 2, 1993.

2. The $SVJ-Y-V$ model: We extend the fitted $SVJ-Y$ model by letting $\lambda^v = \lambda^y$, $\lambda^c = 0$, and λ^y be fixed as given in Table 1. Again, the volatility “jumpiness” is measured by the fraction of the instantaneous variance of V that is due to the jump component. Three jumpiness levels, 0, 15%, and 30% are again considered. For each case, the instantaneous drift, variance, and correlation are matched to the fitted $SVJ-Y$ model.
3. Finally, we modify the fitted $SVJJ$ model by varying the correlation between simultaneous jumps in Y and V . Five levels of correlation are considered: -1.0 , -0.5 , 0 , 0.5 , and 1.0 . For each case, the means and variances of jumps in V and Y are calibrated to the fitted $SVJJ$ model.

Table 2: “Instantaneous” Moments for the *SV* and *SVJ-V* Models

Model	Initial Instantaneous Moments		
	Drift(V)	Var(V)	Corr(Y, V)
<i>SV</i>	$\kappa_v(\bar{v} - V_0)$	$\sigma_v^2 V_0$	$\bar{\rho}$
<i>SVJ-V</i>	$\kappa_v(\bar{v} - \lambda^v \mu_v / \kappa_v - V_0)$	$\sigma_v^2 V_0 + \lambda^v \mu_v^2$	$\bar{\rho} \sigma_v \sqrt{V_0} (\sigma_v^2 V_0 + \lambda^v \mu_v^2)^{-1/2}$

Table 3: Jump Moments for the *SVJJ* Model

Variables	<i>SVJJ</i> Model: Jump Moments		
	Mean	Variance	Correlation
V	μ_v	μ_v^2	—
Y	$\mu_y + \rho_J \mu_v$	$\sigma_y^2 + \rho_J^2 \mu_v^2$	—
(V, Y)	—	—	$\mu_v \rho_J (\sigma_y^2 + \mu_v^2 \rho_J^2)^{-1/2}$

The implied 30-day “volatility smiles” for the above three variations are plotted in Figures 4, 5, and 6.

4.3 Multi-factor Volatility Specifications

Though our focus in this section has been on jump distributions, we are also interested in multi-factor models of the diffusion component of stochastic volatility. Bates [1997] has emphasized the potential importance of more than one volatility factor for explaining the “term structure” of return volatilities, and included two, independent volatility factors in his model. Similarly, the empirical analysis in Gallant, Hsu, and Tauchen [1998] of a non-affine, 3-factor model of asset returns, with two of the three state coordinates dedicated to volatility behavior, suggests that more than one volatility factor improves the goodness of fit for *S&P500* returns.

Our transform analysis applies directly to any affine formulation of multi-factor stochastic volatility models, including Bates’ model. Here, we also propose an examination of multi-factor volatility models in which there is a “long-term” stochastic trend component \bar{V}_t in volatility. For example, we




Figure 4: 30-day smile curve, varying volatility jumpiness, and no jumps in returns.

propose consideration of a three-factor model for $X = (Y, V, \bar{V})^\top$, given in its risk-neutral form by

$$d \begin{pmatrix} Y_t \\ V_t \\ \bar{V}_t \end{pmatrix} = \begin{pmatrix} r - \bar{\zeta} - \frac{1}{2}V_t \\ \kappa(\bar{V}_t - V_t) \\ \kappa_0(\bar{v} - \bar{V}_t) \end{pmatrix} dt + \begin{pmatrix} \sqrt{V_t} & 0 & 0 \\ \sigma\rho\sqrt{V_t} & \sigma\sqrt{1-\rho^2}\sqrt{V_t} & 0 \\ 0 & 0 & \sigma_0\sqrt{\bar{V}_t} \end{pmatrix} dW_t^Q, \quad (4.6)$$

where W^Q is an (\mathcal{F}_t) -standard Brownian motion in \mathbb{R}^3 under Q .

A one-factor volatility model, such as the SV model, may well oversimplify the term structure of volatility. In particular, the (SV) model has an auto-correlation of returns (over successive periods of length Δ) of $\exp(-\kappa_v\Delta)$, which decreases exponentially with Δ . This exponential decay is in direct contrast to a common empirical finding of a “long memory” in volatility. (See, for example, Bollerslev and Mikkelsen [1996] for findings




Figure 5: 30-day smile curve, varying volatility jumpiness. Independent arrivals of jumps in returns and volatility, with independent jump sizes.

based on spot-market data and Pan [1998] for results based on spot-market and options data.) The two-factor volatility model in (4.6), however, yields a more flexible volatility structure. The auto-correlation of Δ -period returns (with respect to the ergodic distribution of (V, \bar{V})) can be calculated for this model to be,

$$\text{corr}(V_t, V_{t+\Delta}) = e^{-\kappa\Delta} + (e^{-\kappa_0\Delta} - e^{-\kappa\Delta}) \frac{\kappa\sigma_0^2/(\kappa - \kappa_0)}{(\kappa + \kappa_0)\sigma^2/\kappa + \kappa\sigma_0^2/\kappa_0}.$$

In subsequent work, we plan to further investigate this or related multi-factor volatility specifications.




Figure 6: 30-day smile curve, varying the correlation between the sizes of simultaneous jumps in return and in volatility.

Appendices

A The Affine Jump-Diffusion

This appendix summarizes technical details for the basic *AJD* model, allowing for time-dependent coefficients.

We fix (Ω, \mathcal{F}, P) , a complete probability space, and $(\mathcal{F}_t)_{0 \leq t < \infty}$, a filtration of sub- σ -fields of \mathcal{F} satisfying the usual conditions.²⁰ We suppose that there is a strong Markov process X , with (X_t, t) in some $D \subset \mathbb{R}^n \times [0, \infty)$ for all

²⁰For technical definitions, see Protter [1990].

t , uniquely solving the stochastic differential equation

$$X_t = X_0 + \int_0^t \mu(X_s, s) ds + \int_0^t \sigma(X_s, s) dW_s + Z_t, \quad (\text{A.1})$$

where W is an (\mathcal{F}_t) -adapted Standard Brownian motion in \mathbb{R}^n ; $\mu : D \rightarrow \mathbb{R}^n$, $\sigma : D \rightarrow \mathbb{R}^{n \times n}$, where D is a subset of $\mathbb{R}^n \times [0, \infty)$ to be defined; Z is a pure jump process whose jump-counting process N has a stochastic intensity $\{\lambda(X_t, t) : t \geq 0\}$, for some $\lambda : D \rightarrow [0, \infty)$, and whose jump-size distribution is ν_t , a probability distribution on \mathbb{R}^n depending only on t . It is assumed that, for each t , $\{x : (x, t) \in D\}$ contains an open subset of \mathbb{R}^n .

We can equally well characterize the behavior of X in terms of the infinitesimal generator \mathcal{D} of its transition semigroup, defined by

$$\begin{aligned} \mathcal{D}f(x, t) &= f_t(x, t) + f_x(x, t)\mu(x, t) + \frac{1}{2}\text{tr}[f_{xx}(x, t)\sigma(x, t)\sigma(x, t)^\top] \\ &\quad + \lambda(x, t) \int_{\mathbb{R}^n} [f(x+z, t) - f(x, t)] d\nu_t(z), \end{aligned} \quad (\text{A.2})$$

for sufficiently regular $f : D \rightarrow \mathbb{R}$. The generator \mathcal{D} is defined by the property that, for any f in its domain, $\{f(X_t, t) - \int_0^t \mathcal{D}f(X_s, s) ds : t \geq 0\}$ is a martingale. (See Ethier and Kurtz [1986] for details.) In Appendix F, we consider more general jump behavior.

We impose an ‘‘affine’’ structure on μ , $\sigma\sigma^\top$, and λ , in that

$$\mu(x, t) = K_0(t) + K_1(t)x \quad (\text{A.3})$$

$$\sigma(x, t)\sigma(x, t)^\top = H_0(t) + \sum_{k=1}^n H_1^{(k)}(t)x_k \quad (\text{A.4})$$

$$\lambda(x, t) = l_0(t) + l_1(t) \cdot x, \quad (\text{A.5})$$

where for each $t \geq 0$, $K_0(t)$ is $n \times 1$, $K_1(t)$ is $n \times n$, $H_0(t)$ is $n \times n$ and symmetric, $H_1(t)$ is a tensor²¹ of dimension $n \times n \times n$, with symmetric $H^{(k)}(t)$ (for $k = 1, \dots, n$), $l_0(t)$ is a scalar, and $l_1(t)$ is $n \times 1$. The time-dependent coefficients $K = (K_0, K_1)$, $H = (H_0, H_1)$, and $l = (l_0, l_1)$ are bounded continuous functions on $[0, \infty)$. We further assume that, for each

²¹Let H be an $n \times n \times n$ tensor, fix its third index to k , the tensor is reduced to an $n \times n$ matrix $H^{(k)}$ with elements, $H_{ij}^{(k)} = H(i, j, k)$.

$t \geq 0$, $\int_0^t \lambda(X_s, s) ds < \infty$ P -a.s. This type of “affine jump-diffusion” process is introduced in Duffie and Kan [1996] for purposes of term-structure modeling.

We know that $\sigma(x, t)$ must be well defined for all (x, t) in D ; indeed one can define regularity conditions on μ , σ , λ , and ν such that a solution X exists for $D = \{(x, t) : \sigma(x, t)\sigma(x, t)^\top \text{ is positive semi-definite}\}$. See Duffie and Kan [1996] and Dai and Singleton [1999] for additional details. The conditions would include the requirement that for any $(x, t) \in D$, we have $(x + z, t) \in D$ for all z in the support of ν_t .

Letting \mathbb{C}^n denote the set of n -tuples of complex numbers, we let $\theta(c, t) = \int_{\mathbb{R}^n} \exp(c \cdot z) d\nu_t(z)$, for any $c \in \mathbb{C}^n$ such that the integral is well defined. This “jump transform” θ determines the probability distribution of each jump measure ν_t . We assume that θ is measurable.

B Transform Analysis

Fixing $T \in [0, \infty)$, the objective of this appendix is to compute the transform $\psi : \mathbb{C}^n \times D \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{C}$ of X_T conditional on \mathcal{F}_t , whenever well defined by

$$\psi(u, X_t, t, T) = E \left(\exp \left(- \int_t^T R(X_s, s) ds \right) \exp(u \cdot X_T) \mid \mathcal{F}_t \right), \quad (\text{B.1})$$

where $R(x, t) = \rho_0(t) + \rho_1(t) \cdot x$, for bounded measurable $\rho_0 : [0, T] \rightarrow \mathbb{R}$ and $\rho_1 : [0, T] \rightarrow \mathbb{R}^n$. The characteristic $\chi = (K, H, l, \theta, \rho)$ determines ψ . With technical regularity conditions, we can show that $\psi = \psi^\chi$, where

$$\psi^\chi(u, x, t, T) = \exp(\alpha(t, T, u) + \beta(t, T, u) \cdot x), \quad (\text{B.2})$$

where β and α satisfy the complex-valued ordinary differential equations

$$\frac{\partial}{\partial t} \beta(t, T, u) + \mathcal{B}(\beta(t, T, u), t) = 0, \quad \beta(T, T, u) = u, \quad (\text{B.3})$$

$$\frac{\partial}{\partial t} \alpha(t, T, u) + \mathcal{A}(\beta(t, T, u), t) = 0, \quad \alpha(T, T, u) = 0, \quad (\text{B.4})$$

and where, for any $c \in \mathbb{C}^n$,

$$\mathcal{B}(c, t) = K_1(t)^\top c + \frac{1}{2} c^\top H_1(t) c - \rho_1(t) + l_1(t) (\theta(c, t) - 1) \quad (\text{B.5})$$

$$\mathcal{A}(c, t) = K_0(t) \cdot c + \frac{1}{2} c^\top H_0(t) c - \rho_0(t) + l_0(t) (\theta(c, t) - 1), \quad (\text{B.6})$$

and where $(c^\top H_1(t)c)$ denotes the n -vector with k -th element $c^\top H_1^{(k)}(t)c$.

Our results will exploit the following technical conditions.

Definition 1: A characteristic (K, H, l, θ, ρ) is *well-behaved* at $(u, T) \in \mathbb{C}^n \times [0, \infty)$ if there is a unique solution X to (A.1) for $0 \leq t \leq T$ and for an initial condition $(X_0, 0) \in D$; if (B.3)-(B.4) are solved uniquely by β and α ; and if

$$(i) \ E \left(\int_0^T |\gamma_t| dt \right) < \infty, \text{ where } \gamma_t = \Psi_t (\theta(\beta(t, T, u), t) - 1) \lambda(X_t, t),$$

$$(ii) \ E \left[\left(\int_0^T \eta_t \cdot \eta_t dt \right)^{1/2} \right] < \infty, \text{ where } \eta_t = \Psi_t \beta(t, T, u)^\top \sigma(X_t, t), \text{ and}$$

$$(iii) \ E (|\Psi_T|) < \infty,$$

where, for each $t \leq T$,

$$\Psi_t = \exp \left(- \int_0^t R(X_s, s) ds \right) \exp (\alpha(t, T, u) + \beta(t, T, u) \cdot X_t). \quad (\text{B.7})$$

Proposition 1 (Transform of X): Suppose (K, H, l, θ, ρ) is well-behaved at (u, T) . Then Ψ is a martingale, and the transform ψ of X defined by (B.1) exists and is given by (B.2).

Proof: By Ito's formula,²²

$$\Psi_t = \Psi_0 + \int_0^t \Psi_s \mu_\Psi(s) ds + \int_0^t \eta_s dW_s + J_t, \quad (\text{B.8})$$

where

$$\mu_\Psi(t) = \frac{\partial}{\partial t} \alpha(t, T, u) + \mathcal{A}(\beta(t, T, u), t) + \left[\frac{\partial}{\partial t} \beta(t, T, u) + \mathcal{B}(\beta(t, T, u), t) \right] \cdot X_t,$$

and

$$J_t = \sum_{0 < \tau^{(i)} \leq t} (\Psi_{\tau^{(i)}} - \Psi_{\tau^{(i)-}}) - \int_0^t \gamma_s ds,$$

²²See Protter [1990] for a complex version of Ito's Formula.

where $\tau(i) = \inf\{t : N_t = i\}$ is the i -th jump time. Under condition (i), Lemma 1, to follow, shows that J is a martingale. Under condition (ii), $\int \eta dW$ is a martingale. Using (B.3) and (B.4), $\mu_\Psi \equiv 0$, and we are done. ■

Lemma 1: Under the assumptions of Proposition 1, J is a martingale.

Proof: Letting E_t denote \mathcal{F}_t -conditional expectation under P , for $0 \leq t \leq s \leq T$, we have

$$\begin{aligned} E_t \left(\sum_{t < \tau(i) \leq s} (\Psi_{\tau(i)} - \Psi_{\tau(i)-}) \right) &= E_t \left(\sum_{t < \tau(i) \leq s} E \left(\Psi_{\tau(i)} - \Psi_{\tau(i)-} \mid X_{\tau(i)-}, \tau(i) \right) \right) \\ &= E_t \left(\sum_{t < \tau(i) \leq s} \Psi_{\tau(i)-} (\theta(b(\tau(i)), \tau(i)) - 1) \right) \\ &= E_t \left(\sum_{t < \tau(i) \leq s} \int_{\tau(i-1)+}^{\tau(i)} \Psi_{u-} (\theta(b(u), u) - 1) dN_u \right) \\ &= E_t \left(\int_t^T \Psi_{u-} (\theta(b(u), u) - 1) dN_u \right). \end{aligned}$$

Because $\{\Psi_{t-} (\theta(b(t), t) - 1) : t \geq 0\}$ is an (\mathcal{F}_t) -predictable process, and the jump-counting process N has intensity $\{\lambda(X_t, t) : t \leq T\}$, integrability condition (i) implies that²³

$$E_t \left(\int_t^s \Psi_{u-} (\theta(b(u), u) - 1) dN_u \right) = E_t \left(\int_t^s \Psi_u (\theta(b(u), u) - 1) \lambda(X_u, u) du \right).$$

Hence J is a martingale. ■

C Transform Inversion

Proposition 2 (Transform Inversion): Suppose, for fixed $T \in [0, \infty)$, $a \in \mathbb{R}$, and $b \in \mathbb{R}^n$, that $\chi = (K, H, l, \theta, \rho)$ is well-behaved at $(a + ivb, T)$, for

²³See, for example, page 27 of Brémaud [1981]. We are applying the result for the real and imaginary components of the integrand, separately.

any $v \in \mathbb{R}$, and that

$$\int_{\mathbb{R}} |\psi^\chi(a + ivb, x, 0, T)| dv < \infty, \quad (\text{C.1})$$

where ψ^χ is defined by (B.2). Then $G_{a,b}(\cdot; x, T, \chi)$ is well defined by (2.8) and given by (2.9).

Proof: For $0 < \tau < \infty$, and a fixed $y \in \mathbb{R}$,

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\tau}^{\tau} \frac{e^{ivy} \psi^\chi(a - ivb, x, 0, T) - e^{-ivy} \psi^\chi(a + ivb, x, 0, T)}{iv} dv \\ &= \frac{1}{2\pi} \int_{-\tau}^{\tau} \int_{\mathbb{R}} \frac{e^{-iv(z-y)} - e^{iv(z-y)}}{iv} dG_{a,b}(z; x, T, \chi) dv \\ &= -\frac{1}{2\pi} \int_{\mathbb{R}} \int_{-\tau}^{\tau} \frac{e^{-iv(z-y)} - e^{iv(z-y)}}{iv} dv dG_{a,b}(z; x, T, \chi), \end{aligned}$$

where Fubini is applicable²⁴ because

$$\lim_{y \rightarrow +\infty} G_{a,b}(y; x, T, \chi) = \psi^\chi(a, x, 0, T) < \infty,$$

given that χ is well-behaved at (a, T) .

Next we note that, for $\tau > 0$,

$$\int_{-\tau}^{\tau} \frac{e^{-iv(z-y)} - e^{iv(z-y)}}{iv} dv = -\frac{\text{sgn}(z-y)}{\pi} \int_{-\tau}^{\tau} \frac{\sin(v|z-y|)}{v} dv$$

is bounded simultaneously in z and τ , for each fixed y .²⁵ By the bounded convergence theorem,

$$\begin{aligned} & \lim_{\tau \rightarrow \infty} \frac{1}{2\pi} \int_{-\tau}^{\tau} \frac{e^{ivy} \psi^\chi(a - ivb, x, 0, T) - e^{-ivy} \psi^\chi(a + ivb, x, 0, T)}{iv} dv \\ &= -\int_{\mathbb{R}} \text{sgn}(z-y) dG_{a,b}(z; x, T, \chi) \\ &= -\psi^\chi(a, x, 0, T) + (G_{a,b}(y; x, T, \chi) + G_{a,b}(y-; x, T, \chi)), \end{aligned}$$

²⁴Here, we also use the fact that, for any $u, v \in \mathbb{R}$, $|e^{iv} - e^{iu}| \leq |v - u|$.

²⁵We define $\text{sgn}(x)$ to be 1 if $x > 0$, 0 if $x = 0$, and -1 if $x < 0$.

where $G_{a,b}(y-; x, T, \chi) = \lim_{z \rightarrow y, z \leq y} G_{a,b}(z; x, T, \chi)$. Using the integrability condition (C.1), by the dominated convergence theorem we have

$$G_{a,b}(y; x, T, \chi) = \frac{\psi^\chi(a, x, 0, T)}{2} + \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{e^{ivy} \psi^\chi(a - ivb, x, 0, T) - e^{-ivy} \psi^\chi(a + ivb, x, 0, T)}{iv} dv.$$

Because $\psi^\chi(a - ivb, x, 0, T)$ is the complex conjugate of $\psi^\chi(a + ivb, x, 0, T)$, we have (2.9). \blacksquare

We summarize our main option-pricing tool as follows.

Proposition 3. The option-pricing formula (2.7) applies, where G is computed by (2.9), provided:

- (a) χ is well-behaved at $(d - ivd, T)$ and at $(-ivd, T)$, for all $v \in \mathbb{R}_+$, and
- (b) $\int_{\mathbb{R}} |\psi^\chi(d - ivd, x, 0, T)| dv < \infty$, and $\int_{\mathbb{R}} |\psi^\chi(-ivd, x, 0, T)| dv < \infty$.

D Change of Measure

This appendix provides the impact of a change of measure defined by a density process or a state-price-density process that is of the exponential-affine form in an affine jump-diffusion state process X .

Fixing $T > 0$, suppose, under the measure P , that a given characteristic $\chi = (K, H, l, \theta, \rho)$ is well-behaved at (b, T) for some $b \in \mathbb{R}^n$. Let

$$\xi_t = \exp \left(- \int_0^t R(X_s, s) ds \right) \exp (\alpha(t, T, b) + \beta(t, T, b) \cdot X_t). \quad (\text{D.1})$$

Under the conditions of Proposition 1, ξ is a positive martingale. We may then define an equivalent probability measure Q by $\frac{dQ}{dP} = \xi_T / \xi_0$.

In this section, we show how to compute the transform of X after a change of measure that arises from a normalization associated with ξ .

Proposition 4 (Transform under Change of Measure):

Let $\chi(Q) = (K^Q, H^Q, l^Q, \theta^Q)$ be defined by

$$K_0^Q(t) = K_0(t) + H_0(t)\beta(t, T, b), \quad K_1^Q(t) = K_1(t) + H_1(t)\beta(t, T, b), \quad (\text{D.2})$$

$$l_0^Q(t) = l_0(t)\theta(\beta(t, T, b), t), \quad l_1^Q(t) = l_1(t)\theta(\beta(t, T, b), t), \quad (\text{D.3})$$

$$\theta^Q(c, t) = \theta(c + \beta(t, T, b), t) / \theta(\beta(t, T, b), t), \quad H^Q(t) = H(t), \quad (\text{D.4})$$

where $H_1(t)b(t)$ denotes the $n \times n$ matrix with k -th column $H_1^{(k)}(t)b(t)$. Let $R^Q(x, t) = \rho_0^Q(t) + \rho_1^Q(t) \cdot x$, for some bounded measurable $\rho_0^Q : [0, \infty) \rightarrow \mathbb{R}$ and $\rho_1^Q : [0, \infty) \rightarrow \mathbb{R}^n$. Let $\rho^Q = (\rho_0^Q, \rho_1^Q)$ be such that $\chi(Q)$ is well-behaved at some (u, T) . Then, for $t \leq T$,

$$E^Q \left(\exp \left(- \int_0^T R^Q(X_s, s) ds \right) \exp(u \cdot X_T) \mid \mathcal{F}_t \right) = \psi^{\chi(Q)}(u, X_t, t, T), \quad (\text{D.5})$$

where $\psi^{\chi(Q)}$ is defined by (B.2).

Proof: Let

$$W_t^Q = W_t - \int_0^t \sigma(X_s, s)^\top \beta(s, T, b) ds, \quad t \geq 0. \quad (\text{D.6})$$

Lemma 2, below, shows that ξW^Q is a P -local martingale. It follows that W^Q is a Q -local martingale. Because $\int_0^t \sigma^\top(X_s, s) \beta(s, T, b) ds$ is a continuous finite-variation process, $[W_i^Q, W_j^Q]_t = [W_i^P, W_j^P]_t = \delta(i, j) t$, where $\delta(\cdot)$ is the kronecker delta. By Lévy's Theorem, W^Q is a standard Brownian motion in \mathbb{R}^n under Q .

Next, we let

$$M_t^Q = N_t - \int_0^t \theta(\beta(s, T, b)) \lambda(X_s, s) ds, \quad t \geq 0. \quad (\text{D.7})$$

Lemma 3, below, shows that ξM^Q is a P -local martingale. It follows that M^Q is a Q -local martingale. By the martingale characterization of intensity,²⁶ we conclude that, under Q , N is a counting process with the intensity $\{\lambda^Q(X_t, t) : t \geq 0\}$ defined by $\lambda^Q(x, t) = l_0^Q(t) + l_1^Q(t) \cdot x$.

Using the fact that, under Q , W^Q is a standard Brownian and the jump counting process N has intensity $\{\lambda^Q(X_t, t) : t \geq 0\}$, we may mimic the proof of Proposition 1, and obtain (D.5) replacing in the proof of Lemma 1 $E_t \left(\sum_{t < \tau^{(i)} \leq T} (\Psi_{\tau^{(i)}} - \Psi_{\tau^{(i)-}}) \right)$ with

$$E_t^Q \left(\sum_{t < \tau^{(i)} \leq T} (\Psi_{\tau^{(i)}} - \Psi_{\tau^{(i)-}}) \right) = \frac{1}{\xi_t} E_t \left(\sum_{t < \tau^{(i)} \leq T} \xi_{\tau^{(i)}} (\Psi_{\tau^{(i)}} - \Psi_{\tau^{(i)-}}) \right).$$

²⁶See, for example, page 28 of Brémaud [1981].

This completes the proof. \blacksquare

Lemma 2: Under the assumptions of Proposition 1, ξW^Q is a P -local martingale.

Proof: By Ito's Formula, with $0 \leq s \leq t \leq T$,

$$\begin{aligned}
\xi_t W_t^Q &= \xi_s W_s^Q + \int_s^t \xi_{u-} dW_u^Q + \int_s^t W_{u-}^Q d\xi_u \\
&\quad + \sum_{s < u \leq t} (\xi_u - \xi_{u-}) (W_t^Q - W_{t-}^Q) + \int_s^t d[\xi, W^Q]_u^c \\
&= \xi_s W_s^Q + \int_s^t \xi_{u-} (dW_u - \sigma^\top(X_u, u)b(u) du) \\
&\quad + \int_s^t W_u^Q d\xi_u + \int_s^t \xi_u \sigma^\top(X_u, u)b(u) du \\
&= \xi_s W_s^Q + \int_s^t \xi_{u-} dW_u + \int_s^t W_u^Q d\xi_u,
\end{aligned}$$

where $[\xi, W^Q]^c$ denotes the continuous part of the “square-brackets” process $[\xi, W^Q]$. As W and ξ are P -martingales, both $\left\{ \int_0^t \xi_{u-} dW_u : t \geq 0 \right\}$ and $\left\{ \int_0^t W_u^Q d\xi_u : t \geq 0 \right\}$ are P -local martingales. Hence, ξW^Q is a P -local martingale. \blacksquare

Lemma 3: Under the assumptions of Proposition 1, ξM^Q is a P -local martingale.

Proof: By Ito's Formula, with $0 \leq s \leq t \leq T$,

$$\begin{aligned}
\xi_t M_t^Q &= \xi_s M_s^Q + \int_s^t \xi_{u-} dM_u^Q + \int_s^t M_{u-}^Q d\xi_u + \sum_{s < u \leq t} (\xi_u - \xi_{u-})(N_u - N_{u-}) \\
&= \xi_s M_s^Q + \int_s^t \xi_{u-} dM_u + \int_s^t M_{u-}^Q d\xi_u + J^\xi,
\end{aligned}$$

where $M_t = N_t - \int_0^t \lambda(X_s, s) ds$, and where

$$J^\xi = \sum_{s < u \leq t} (\xi_u - \xi_{u-}) - \int_s^t \xi_u (\theta(\beta(u, T, b), u) - 1) \lambda(X_u, u) du.$$

As M and ξ are P -martingales, $\left\{ \int_0^t \xi_{u-} dM_u : t \geq 0 \right\}$ and $\left\{ \int_0^t M_{u-}^Q d\xi_u : t \geq 0 \right\}$ are P -local martingales. By a proof similar to that of Lemma 1, and using the Integration Theorem (γ) in Brémaud [1981], we can show that J^ξ is a P -local martingale. ■

For the remainder of this appendix, we denote Q by $Q(b)$, emphasizing the role of b in defining the change of probability measure given by (D.1). We let $\chi(b) = (K^{Q(b)}, H^{Q(b)}, l^{Q(b)}, \theta^{Q(b)}, \rho)$ denote the associated characteristic. The previous result shows in effect that, under $Q(b)$, the state vector X is still an affine jump-diffusion whose characteristics can be computed in terms of the characteristics of X under the measure P . This result provides us with an alternative approach to option pricing. We suppose that $Q(0)$ is an equivalent martingale measure. The price $\Gamma(X_0, a, d, c, T)$ of an option paying $(e^{a+d \cdot X_T} - c)^+$ at T is given by

$$\begin{aligned} \Gamma(X_0, a, d, c, T) &= E^{Q(0)} \left(\exp \left(- \int_0^T R(X_s, s) ds \right) (e^{a+d \cdot X_T} - c)^+ \right) \\ &= e^a E^{Q(0)} \left(\exp \left(- \int_0^T R(X_s, s) ds \right) e^{d \cdot X_T} \mathbf{1}_{d \cdot X_T \geq \ln(c) - a} \right) \\ &\quad - c E^{Q(0)} \left(\exp \left(- \int_0^T R(X_s, s) ds \right) \mathbf{1}_{d \cdot X_T \geq \ln(c) - a} \right). \end{aligned}$$

Provided the characteristic (K, H, l, θ, ρ) is well-behaved at (d, T) and $(0, T)$, we may introduce the equivalent probability measure $Q(d)$, and write

$$\begin{aligned} \Gamma(X_0, a, d, c, T) &= e^a \exp(\alpha(0, T, d) + \beta(0, T, d) \cdot X_0) E^{Q(d)} \left(\mathbf{1}_{d \cdot X_T \geq \ln(c) - a} \right) \\ &\quad - c \exp(\alpha(0, T, 0) + \beta(0, T, 0) \cdot X_0) E^{Q(0)} \left(\mathbf{1}_{d \cdot X_T \geq \ln(c) - a} \right). \end{aligned}$$

Let $\chi(1) = (K^{Q(d)}, H^{Q(d)}, l^{Q(d)}, \theta^{Q(d)}, 0)$ and $\chi(0) = (K^{Q(0)}, H^{Q(0)}, l^{Q(0)}, \theta^{Q(0)}, 0)$ be defined by (D.2)-(D.4) for $b = d$ and $b = 0$. We suppose that $\chi(1)$ and $\chi(0)$ are well behaved at (ivd, T) for any $v \in \mathbb{R}$. Then

$$\begin{aligned} E^{Q(d)} \left(\mathbf{1}_{d \cdot X_T \geq \ln(c) - a} \right) &= \frac{1}{2} + \frac{1}{\pi} \int_0^\infty \frac{\text{Im} [\psi^{\chi(1)}(ivd, x, 0, T) e^{-iv(\ln(c) - a)}]}{v} dv, \\ E^{Q(0)} \left(\mathbf{1}_{d \cdot X_T \geq \ln(c) - a} \right) &= \frac{1}{2} + \frac{1}{\pi} \int_0^\infty \frac{\text{Im} [\psi^{\chi(0)}(ivd, x, 0, T) e^{-iv(\ln(c) - a)}]}{v} dv, \end{aligned}$$

provided $\int_{\mathbb{R}} |\psi^{\chi(1)}(ivd, X_0, 0, T)| dv < \infty$ and $\int_{\mathbb{R}} |\psi^{\chi(0)}(ivd, X_0, 0, T)| dv < \infty$. These quantities may now be substituted into the previous relation in order to obtain the option price.

E “Extended” Transform Analysis

In this appendix, fixing a characteristic χ , we introduce an “extended” transform $\phi : \mathbb{R}^n \times \mathbb{C}^n \times D \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{C}$ of X_T conditional on \mathcal{F}_t , when well defined for $t \leq T$ by

$$\phi(v, u, X_t, t, T) = E \left(\exp \left(- \int_t^T R(X_s, s) ds \right) (v \cdot X_T) e^{u \cdot X_T} \mid \mathcal{F}_t \right). \quad (\text{E.1})$$

Under additional technical conditions, we can show that

$$\phi(v, u, x, t, T) = \psi^\chi(u, x, t, T) (A(t, T, v, u) + B(t, T, v, u) \cdot x), \quad (\text{E.2})$$

where ψ^χ is given by (B.2), and where B and A satisfy the linear ordinary differential equations

$$\begin{aligned} \frac{\partial}{\partial t} B(t, T, v, u) + K_1(t)^\top B(t, T, v, u) + \beta(t, T, u)^\top H_1(t) B(t, T, v, u) \\ + l_1(t) \Theta(\beta(t, T, u), t) \cdot B(t, T, v, u) = 0, \quad B(T, T, v, u) = v, \end{aligned} \quad (\text{E.3})$$

$$\begin{aligned} \frac{\partial}{\partial t} A(t, T, v, u) + K_0(t) \cdot B(t, T, v, u) + \beta(t, T, u)^\top H_0(t) B(t, T, v, u) \\ + l_0(t) \Theta(\beta(t, T, u), t) \cdot B(t, T, v, u) = 0, \quad A(T, T, v, u) = 0, \end{aligned} \quad (\text{E.4})$$

where $\Theta(c, t) = \int_{\mathbb{R}^n} \exp(c \cdot z) z d\nu_t(z)$.

Letting Ψ_t be defined by (B.7) and $\Phi_t = \Psi_t (A(t, T, v, u) + B(t, T, v, u) \cdot X_t)$, sufficient technical conditions are

(i) $E \left(\int_0^T |\tilde{\gamma}_t| dt \right) < \infty$, where

$$\tilde{\gamma}_t = \lambda(X_t, t) (\Phi_t (\theta(\beta(t, T, u), t) - 1) + \Psi_t B(t, T, v, u) \cdot \Theta(\beta(t, T, u), t)).$$

(ii) $E \left[\left(\int_0^T \tilde{\eta}_t \cdot \tilde{\eta}_t dt \right)^{1/2} \right] < \infty$, where

$$\tilde{\eta}_t = \Phi_t (\beta(t, T, u)^\top + B(t, T, v, u)^\top) \sigma(X_t, t).$$

(iii) $E (|\Phi_T|) < \infty$.

Definition E1: (K, H, l, θ, ρ) is “*extended well-behaved*” at (v, u, T) , if there is a unique solution X to (A.1) for $0 \leq t \leq T$, if (B.3)-(B.4) are solved uniquely by β and α , if (E.3)-(E.4) are solved uniquely by B and A , and if the above conditions (i)-(iii) are satisfied.

Proposition 5 (“Extended” Transform of X): Suppose $\chi = (K, H, l, \theta, \rho)$ is extended well-behaved at (v, u, T) . Then Φ is a martingale, and the transform ϕ of X defined by (E.1) is thus given by (E.2).

In principle, the extended transform ϕ can be computed by differentiation of the transform ψ , just as moments can be computed from a moment generating function. In practice, this may involve solving the same ODEs (E.3)-(E.4).

For fixed $a \in \mathbb{R}^n$, $b \in \mathbb{R}^n$, and $d \in \mathbb{R}^n$, we next define $\tilde{G}_{a,b,d}(\cdot; X_0, T, \chi)$ by

$$\tilde{G}_{a,b,d}(y; X_0, T, \chi) = E \left(\exp \left(- \int_0^T R(X_s, s) ds \right) (a \cdot X_T) e^{d \cdot X_T} \mathbf{1}_{b \cdot X_T \leq y} \right). \quad (\text{E.5})$$

Provided $\chi = (K, H, l, \theta, \rho)$ is extended well behaved at $(a, d + ivb, T)$, for any $v \in \mathbb{R}$, and that $\int_{\mathbb{R}} |\phi(a, d + ivb, x, 0, T)| dv < \infty$, $\tilde{G}_{a,b}$ can be obtained by the Fourier-inversion of ϕ , so that

$$\tilde{G}_{a,b,d}(y; x, T, \chi) = \frac{\phi(a, d, x, 0, T)}{2} - \frac{1}{\pi} \int_0^\infty \frac{\text{Im} [\phi(a, d + ivb, x, 0, T) \exp(-ivy)]}{v} dv. \quad (\text{E.6})$$

Now, anticipating the calculation of option prices, we consider, for given $c \in \mathbb{R}$ and $b \in \mathbb{R}^n$.

$$\tilde{C}^x(X_0, b, c, T) = E^x \left(\exp \left(- \int_0^T R(X_s, s) ds \right) (b \cdot X_T - c)^+ \right). \quad (\text{E.7})$$

We immediately obtain

$$\begin{aligned} \tilde{C}^x(X_t, b, c, T) &= E^x \left(\exp \left(- \int_0^T R(X_s, s) ds \right) (b \cdot X_T - c) \mathbf{1}_{b \cdot X_T \geq c} \right) \\ &= \tilde{G}_{b,-b,0}(-c; X_0, T, \chi) - c G_{0,-b}(-c; X_0, T, \chi), \end{aligned} \quad (\text{E.8})$$

where $G_{a,b}$ is given by (2.9) and $\tilde{G}_{a,b,0}$ is given by (E.6).

With this calculation, we could price a slope-of-the-yield-curve option, as yields in an *AJD* setting are themselves affine. Under the assumption of a deterministic short rate and dividend-yield process, that is, $\rho_1 = q_1 = 0$, we may also use this approach to price an asian option. For the latter, struck at c , at the expiration date T , the option pays $\left(\frac{1}{T} \int_0^T X_t^{(i)} dt - c\right)$, where $X^{(i)}$ is the price process of the underlying asset. If Q is an equivalent martingale measure, we must have

$$dX_t^{(i)} = (R(X_t, t) - \zeta(X_t, t)) X_t dt + dM_t^{(i)},$$

where M^i is a Q -martingale. For any $0 \leq t \leq T$, let $Y_t = \int_0^t X_s^{(i)} ds$. For short rate ρ_0 , we can let $\tilde{\rho}_0 = (\rho_0, 0)$ and $\tilde{\rho}_1 = (0, 0) = 0$, and see that $\tilde{X} = (X, Y)$ is an $(n+1)$ -dimensional affine jump diffusion with characteristic $\tilde{\chi} = \left(\tilde{K}, \tilde{H}, \tilde{l}, \tilde{\theta}, \tilde{\rho}\right)$ that can be easily derived from using the fact that $dY_t = X_t^{(i)} dt$. We may then use (E.8) and obtain the initial market value of the option as

$$\frac{1}{T} \tilde{C}^{\tilde{\chi}} \left(\tilde{X}_0, \epsilon^{(n+1)}, Tc, T \right).$$

F Extension to Multiple Jumps

We may easily relax the jump behavior of X to accommodate m types of jumps, with jump type i having jump-conditional distribution ν_t^i at time t , again depending only on t , and stochastic intensity $\{\lambda_i(X_t, t) : t \geq 0\}$, for $i \in \{1, \dots, m\}$, where $\lambda_i : D \rightarrow \mathbb{R}_+$ is defined by

$$\lambda_i(x, t) = l_0^i(t) + l_t^i(t) \cdot x,$$

for bounded measurable $l = ((l_0^1, l_1^1), \dots, (l_0^m, l_1^m))$. The jump transforms $\theta = (\theta^1, \dots, \theta^m)$ are defined by $\theta^i(c, t) = \int_{\mathbb{R}^n} \exp(c \cdot z) d\nu_t^i(z)$, $c \in \mathbb{C}^n$.

We can also characterize the behavior of X with multiple jumps in terms of the infinitesimal generator \mathcal{D} of its transition semigroup, with

$$\begin{aligned} \mathcal{D}f(x, t) &= f_t(x, t) + f_x(x, t)\mu(x, t) + \frac{1}{2} \text{tr} [f_{xx}(x, t)\sigma(x, t)\sigma(x, t)^\top] \\ &+ \sum_{i=1}^m \lambda^i(x, t) \int_{\mathbb{R}^n} [f(x+z, t) - f(x, t)] d\nu_t^i(z), \end{aligned} \quad (\text{F.1})$$

for sufficiently regular $f : D \rightarrow \mathbb{R}$.

In this general setting, Propositions 1, 2, and 3 apply after replacing the last terms in the right-hand sides of (B.5) and (B.6) with $\sum_{i=1}^m l_1^i(t) (\theta^i(c, t) - 1)$ and $\sum_{i=1}^m l_0^i(t) (\theta^i(c, t) - 1)$, respectively.

This can be extended to the case of an infinite number of jump types by allowing for a general Lévy jump measure that is affine in the state vector. (See Theorem 42, page 32, of Protter [1990].)

G Cap Pricing

A cap is a loan with face value, say 1, at a variable interest rate that is capped at some level \bar{r} . At time t , let $\tau, 2\tau, \dots, n\tau$ be the fixed dates for future interest payments. At each fixed date $k\tau$, the \bar{r} -capped interest payment, or “caplet,” is given by $\tau (\mathcal{R}((k-1)\tau, k\tau) - \bar{r})^+$, where $\mathcal{R}((k-1)\tau, k\tau)$ is the τ -year floating interest rate at time $(k-1)\tau$, defined by

$$\frac{1}{1 + \tau \mathcal{R}((k-1)\tau, k\tau)} = \Lambda((k-1)\tau, k\tau).$$

The market value at time 0 of the caplet paying at date $k\tau$ can be expressed as

$$\begin{aligned} \text{Caplet}(k) &= E^{\mathcal{Q}} \left[\exp \left(- \int_0^{k\tau} R(X_u, u) du \right) \tau (\mathcal{R}((k-1)\tau, k\tau) - \bar{r})^+ \right] \\ &= (1 + \tau \bar{r}) E^{\mathcal{Q}} \left[\exp \left(- \int_0^{(k-1)\tau} R(X_u, u) du \right) \left(\frac{1}{1 + \tau \bar{r}} - \Lambda((k-1)\tau, k\tau) \right)^+ \right]. \end{aligned}$$

Hence, the pricing of the k -th caplet is equivalent to the pricing of an in- $(k-1)\tau$ -for- τ put struck at $1/(1 + \tau \bar{r})$, which can be readily obtained by using Proposition 3 and put-call parity as $\text{Caplet}(k) = (1 + \tau \bar{r}) \bar{C}(k)$, where

$$\bar{C}(k) = \Gamma \left(X_0, \bar{\alpha}, \bar{\beta}, \frac{1}{1 + \tau \bar{r}}, (k-1)\tau \right) - \Lambda(0, k\tau) + \frac{\Lambda(0, (k-1)\tau)}{1 + \tau \bar{r}},$$

where $\Gamma(X_0, a, d, c, T)$ is the price of a claim to $(e^{a+d \cdot X(T)} - c)^+$ paid at T , and where $\bar{\alpha} = \alpha((k-1)\tau, k\tau, 0)$ and $\bar{\beta} = \beta((k-1)\tau, k\tau, 0)$.

References

- Bakshi, G., C. Cao, and Z. Chen (1997). Empirical Performance of Alternative Option Pricing Models. *Journal of Finance* 52, 2003–49.
- Bakshi, G. and D. Madan (1999). Spanning and Derivative-Security Valuation. *Journal of Financial Economics*, forthcoming.
- Bates, D. (1996). Jump and Stochastic Volatility: Exchange Rate Processes Implicit in Deutsche Mark Options. *The Review of Financial Studies* 9, 69–107.
- Bates, D. (1997). Post-'87 Crash fears in S&P 500 Futures Options. Working Paper 5894, National Bureau of Economic Research.
- Benzoni, L. (1998). Pricing Options under Stochastic Volatility: An Econometric Analysis. Working Paper, J.L. Kellogg Graduate School of Management, Northwestern University.
- Bollerslev, T. and H. O. Mikkelsen (1996). Modeling and Pricing Long Memory in Stock Market Volatility. *Journal of Econometrics* 73, 151–184.
- Brémaud, P. (1981). *Point Processes and Queues, Martingale Dynamics*. New York: Springer-Verlag.
- Chacko, G. and S. Das (1998). Pricing Average Interest Rate Options: A General Approach. Working Paper, Harvard Business School, Harvard University.
- Chen, R. and L. Scott (1993). Maximum Likelihood Estimation for a Multifactor Equilibrium Model of the Term Structure of Interest Rates. *The Journal of Fixed Income* 3, 14–31.
- Chen, R. and L. Scott (1995). Interest Rate Options in Multifactor Cox-Ingersoll-Ross Models of the Term Structure. *Journal of Derivatives* 3, 53–72.
- Chernov, M. and E. Ghysels (1998). What Data Should be Used to Price Options? Working Paper, Department of Finance, Pennsylvania State University.
- Cox, J. C., J. E. Ingersoll, and S. A. Ross (1985). A Theory of the Term Structure of Interest Rates. *Econometrica* 53(2), 385–407.

- Dai, Q. and K. Singleton (1999). Specification Analysis of Affine Term Structure Models. Working Paper, Graduate School of Business, Stanford University.
- Das, S. (1998). Poisson-Gaussian Processes and The Bond Markets. Working Paper, Harvard Business School, Harvard University.
- Delbaen, F. and W. Schachermayer (1994). A General Version of the Fundamental Theorem of Asset Pricing. *Math. Ann.* 300, 463–520.
- Duffie, D. and R. Kan (1996). A Yield-Factor Model of Interest Rates. *Mathematical Finance* 6(4), 379–406.
- Duffie, D. and K. Singleton (1999). Modeling Term Structures of Defaultable Bonds. *Review of Financial Studies*, forthcoming.
- Ethier, S. and T. Kurtz (1986). *Markov Processes, Characterization and Convergence*. New York: John Wiley & Sons.
- Gallant, R., C.-T. Hsu, and G. Tauchen (1998). Calibrating Volatility Diffusions and Extracting Integrated Volatility. Working Paper, Department of Economics, University of North Carolina.
- Gil-Pelaez, J. (1951). Note on the Inversion Theorem. *Biometrika* 38, 481–482.
- Harrison, M. and D. Kreps (1979). Martingales and Arbitrage in Multi-period Securities Markets. *Journal of Economic Theory* 20, 381–408.
- Heston, S. (1993). A Closed-Form Solution of Options with Stochastic Volatility with Applications to Bond and Currency Options. *The Review of Financial Studies* 6, 327–343.
- Jamshidian, F. (1989). An Exact Bond Option Formula. *Journal of Finance* 44, 205–209.
- Jarrow, R., D. Lando, and S. Turnbull (1997). A Markov Model for the Term Structure of Credit Spreads. *Journal of Finance* 10, 481–523.
- Lando, D. (1998). On Cox Processes and Credit Risky Securities. *Review of Derivatives Research* 2, 99–120.
- Liu, J. (1997). Generalized Method of Moments Estimation of Affine Diffusion Processes. Working Paper, Graduate School of Business, Stanford University.

- Liu, J. and J. Pan (1997). Asymptotic Maximum Likelihood Estimation of Affine Models. Working Paper, Graduate School of Business, Stanford University.
- Pan, J. (1998). “Integrated” Time-Series Analysis of Spot and Options Prices. Working Paper, Graduate School of Business, Stanford University.
- Piazzesi, M. (1998). Monetary Policy and Macroeconomic Variables in a Model of the Term Structure of Interest Rates. Working Paper, Stanford University.
- Poteshman, A. M. (1998). Estimating a General Stochastic Variance Model from Options Prices. Working Paper, Graduate School of Business, University of Chicago.
- Protter, P. (1990). *Stochastic Integration and Differential Equations*. New York: Springer-Verlag.
- Scott, L. (1996). The Valuation of Interest Rate Derivatives in a Multi-Factor Cox-Ingersoll-Ross Model that Matches the Initial Term Structure. Working Paper, University of Georgia.
- Scott, L. (1997). Pricing Stock Options in a Jump-diffusion Model with Stochastic Volatility and Interest Rates: Application of Fourier Inversion Methods. *Mathematical Finance* 7, 345–358.
- Singleton, K. (1998). Estimation of Affine Diffusion Models Based on the Empirical Characteristic Function. Working Paper, Graduate School of Business, Stanford University.
- Vasicek, O. (1977). An Equilibrium Characterization of the Term Structure. *Journal of Financial Economics* 5, 177–188.
- Williams, D. (1991). *Probability with Martingales*. Cambridge University Press.