

 Open access • Journal Article • DOI:10.1007/S11940-013-0277-Y

Treatment of primary CNS lymphoma. — [Source link](#)

Patrick Roth, Roger Stupp, Günter Eisele, Michael Weller

Institutions: University of Zurich

Published on: 01 Jan 2014 - Current Treatment Options in Neurology (Springer US)

Topics: Primary central nervous system lymphoma, Regimen, Salvage therapy, Radiation therapy and Rituximab

Related papers:

- [High-dose methotrexate with or without whole brain radiotherapy for primary CNS lymphoma \(G-PCNSL-SG-1\): a phase 3, randomised, non-inferiority trial](#)
- [Diagnosis and treatment of primary CNS](#)
- [Primary Central Nervous System Lymphoma.](#)
- [Primary central nervous system lymphoma: recent progress, many remaining questions.](#)
- [Pharmacotherapy for primary CNS lymphoma: progress beyond methotrexate?](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/treatment-of-primary-cns-lymphoma-bn8bsb4mo5>



Treatment of primary CNS lymphoma

Roth, P ; Stupp, R ; Eisele, G ; Weller, M

Abstract: OPINION STATEMENT: Primary central nervous system lymphoma is a particular challenge in clinical neuro-oncology. In contrast to most other malignant brain tumors, it may be considered a curable disease at least in younger patients who can tolerate intensive treatment regimens. Yet, therapeutic progress has been limited with little measurable improvement in outcome over the last two decades, mainly due to the low incidence of this tumor, which impedes the execution of large randomized clinical trials, and the failure of most large cooperative groups to conduct such trials. Whenever possible, high-dose methotrexate (HD-MTX) is the backbone of the therapeutic regimen. Response rates can be increased by the addition of second agents like ifosfamide or cytarabine, however, their impact on overall survival is less clear. Similarly, the use of the anti-CD20 antibody rituximab, commonly used in the treatment of B cell lymphomas outside the CNS, remains controversial and has not been examined in adequate clinical trials. The prognosis of patients, who do not qualify for HD-MTX-based chemotherapy, is considerably poorer. Radiation therapy is an active treatment with high response rates but does typically not result in long-lasting remissions. It remains an important therapeutic option as a salvage therapy in patients progressing on or no longer responding to HD-MTX-based treatment. The combination of HD-MTX and radiation therapy does not prolong overall survival. It is associated with significant neurotoxicity, and it should be avoided. Another matter of debate is whether consolidation therapy by other means, such as high-dose chemotherapy followed by stem cell support, is the most promising regimen. Given these numerous uncertainties, neuro-oncologists should strive for a treatment of PCNSL patients within clinical trials to allow for the development of improved therapeutic regimens.

DOI: <https://doi.org/10.1007/s11940-013-0277-y>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-95410>

Journal Article

Published Version

Originally published at:

Roth, P; Stupp, R; Eisele, G; Weller, M (2014). Treatment of primary CNS lymphoma. *Current Treatment Options in Neurology*, 16(1):277.

DOI: <https://doi.org/10.1007/s11940-013-0277-y>

Treatment of Primary CNS Lymphoma

Patrick Roth, MD^{1,3,*}

Roger Stupp, MD^{2,3}

Günter Eisele, MD^{1,3}

Michael Weller, MD^{1,3}

Address

*¹Department of Neurology, University Hospital Zurich, Frauenklinikstrasse 26, 8091 Zurich, Switzerland

E-mail: patrick.roth@usz.ch

²Department of Oncology, University Hospital Zurich, Zurich, Switzerland

³Brain Tumor Center Zurich, University Hospital Zurich, Zurich, Switzerland

Published online: 17 December 2013

© Springer Science+Business Media New York 2013

This article is part of the Topical Collection on *Neuro-oncology*

Keywords Primary CNS lymphoma · Chemotherapy · Methotrexate · Rituximab · Temozolomide · Irradiation · Neurotoxicity · Treatment · Steroids · Intrathecal therapy

Opinion statement

Primary central nervous system lymphoma is a particular challenge in clinical neuro-oncology. In contrast to most other malignant brain tumors, it may be considered a curable disease at least in younger patients who can tolerate intensive treatment regimens. Yet, therapeutic progress has been limited with little measurable improvement in outcome over the last two decades, mainly due to the low incidence of this tumor, which impedes the execution of large randomized clinical trials, and the failure of most large cooperative groups to conduct such trials. Whenever possible, high-dose methotrexate (HD-MTX) is the backbone of the therapeutic regimen. Response rates can be increased by the addition of second agents like ifosfamide or cytarabine, however, their impact on overall survival is less clear. Similarly, the use of the anti-CD20 antibody rituximab, commonly used in the treatment of B cell lymphomas outside the CNS, remains controversial and has not been examined in adequate clinical trials. The prognosis of patients, who do not qualify for HD-MTX-based chemotherapy, is considerably poorer. Radiation therapy is an active treatment with high response rates but does typically not result in long-lasting remissions. It remains an important therapeutic option as a salvage therapy in patients progressing on or no longer responding to HD-MTX-based treatment. The combination of HD-MTX and radiation therapy does not prolong overall survival. It is associated with significant neurotoxicity, and it should be avoided. Another matter of debate is whether consolidation therapy by other means, such as high-dose chemotherapy followed by stem cell support, is the most promising regimen. Given these numerous uncertainties, neuro-oncologists should strive for a treatment of PCNSL patients within clinical trials to allow for the development of improved therapeutic regimens.

Introduction

Primary central nervous system lymphoma (PCNSL) is a rare extranodal form of Non-Hodgkin lymphoma (NHL). Registry data from the United States report that only 2.2 % of all intracranial tumors are lymphomas [1]. Immunodeficiency (e.g., congenital or acquired such as human immunodeficiency virus [HIV] infection) is associated with an increased risk for the development of PCNSL. The improvement of antiretroviral therapy resulted in a decreased incidence of PCNSL in immunocompromised patients translating into fewer young patients affected with PCNSL. This contrasts with a continued increase in men and women aged 65 and older for reasons yet to be determined [2]. In immunocompetent patients, the median age at diagnosis is about 60 years.

PCNSL can involve all parts of the CNS including supratentorial and infratentorial compartments, spinal cord, meninges and cerebrospinal fluid (CSF). Finally, intraocular manifestation, either alone, then referred to as primary intraocular lymphoma (PIOL), or often associated with CNS involvement, can occur. Patients with PCNSL may present with various clinical symptoms and signs. The extent of personality changes, cognitive disturbances and focal neurological deficits will depend on tumor size, localization and extent of the peritumoral edema. Intraocular involvement may cause blurred vision or other complaints.

Computed tomography (CT) or preferentially magnetic resonance imaging (MRI) are used as imaging techniques with single or multiple supratentorial homogeneous contrast-enhancing tumor masses as a typical finding [3]. There are no specific features, which allow a reliable differentiation between PCNSL and other primary or secondary brain tumors as well as inflammatory lesions. The use of ultrasmall superparamagnetic iron oxide (USPIO) nanoparticles may improve the MRI diagnosis of PCNSL [4]. This technique, however, has not yet entered the clinical routine examination. Similarly, the value of positron emission tomography (PET) imaging using different tracers such as ^{18}F -fluorodeoxyglucose (^{18}F -FDG) or ^{11}C -methionine for the diagnosis and monitoring of PCNSL has remained questionable [5–7]. Upon diagnosis of a lymphoma in the CNS, the diagnosis of PCNSL can only be rendered after exclusion of tumor

manifestations outside the CNS. The detection of lymphoma cells in the CSF, also referred to as meningeal dissemination, in the eye and cranial nerves is still compatible with PCNSL. However, staging is recommended in every patient to exclude other lymphoma manifestations [8]. This is typically done by chest, abdomen and pelvis CT. Whether FDG-PET-based visualization results in increased sensitivity remains unclear [9]. Ocular infiltration must be ruled out by an appropriate ophthalmologic diagnostics. Vitrectomy followed by histopathological assessment of the aspirate is used to definitely verify or exclude lymphoma manifestations in the eye [10]. Bone marrow diagnostics may reveal the presence of tumor cells which, however, does not change the therapeutic approach in the absence of other systemic lymphoma manifestations [11]. Further recommendations of the International Extranodal Lymphoma Study Group (IELSG) include a lumbar puncture followed by CSF examination and testicular ultrasound in older men to exclude the presence of testicular lymphoma metastatic to brain [8].

Histopathological assessment of PCNSL tissue results in the diagnosis of a diffuse large B cell lymphoma (DLBCL) in about 90 % of all patients. These tumors typically express the B cell marker CD20, a prerequisite for consideration of treatment with the monoclonal antibody rituximab (see below). Standard cytology from body fluids such as vitreous or CSF may be broadened using flow cytometry analysis as well as PCR diagnostics of the rearranged immunoglobulin heavy-chain genes. Histopathological features as well as pitfalls in the diagnosis of PCNSL have been reviewed elsewhere [12]. CNS lymphoma with a histological diagnosis other than DLBCL is rare, including low-grade lymphomas and lymphomas of T cell origin as very uncommon findings [13, 14].

Because of the variable course of the disease, several attempts have been made to group patients into different clinical prognostic groups, which may help to estimate the clinical evolution. The “Memorial Sloan-Kettering Cancer Center prognostic model” published in 2006 is based on a large cohort of PCNSL patients [15]. Patients younger than 50 years had by far the

best outcome. In the group of patients aged 50 or older, the performance status allowed for further separation between two groups with patients being 50 years or older and a Karnofsky performance status (KPS) below 70 having the worst prognosis. A more comprehensive prognostic score was proposed by the IELSG using not only age and performance status but also elevated lactate dehydrogenase (LDH) and CSF protein levels as additional parameters which are associated with poor outcome [16]. Furthermore, the involvement of “deep brain structures” was determined as a negative factor and included in the prognostic assessment. This allowed for a separation of three patient groups, which differ in their prognosis.

Treatment

Similar to many other brain tumors, the available treatment modalities for PCNSL comprise surgery, irradiation and chemotherapy, with a clear focus on the latter two options. The agents or regimens (e.g., CHOP) commonly used to treat systemic lymphoma have no or only marginal activity in PCNSL [17]. There are probably several reasons including different biological properties of the tumors cells and the presence of the blood-brain barrier that impedes the penetration of many chemotherapeutics into the brain. Despite some progress within the last years, many important questions regarding the most effective treatment for PCNSL remain under debate. The rarity of PCNSL precludes the rapid execution of well-designed clinical trials with reasonable patient numbers and does not allow for rapid evaluation of different treatment regimens. Still, the publication of two larger randomized trials within the last years has shown that such studies are feasible and should be pursued within the next years [18, 19].

Surgery

The intention of surgery has been purely diagnostic for decades. However, an unplanned subgroup analysis of the G-PCNSL-SG1 dataset, comprising several hundred patients, revealed that complete resection may be associated with a better outcome [20•]. These findings are in contrast to very small and older retrospective series that suggested a biopsy-only approach. However, surgical techniques have improved and intra-operative imaging including ultrasound and MRI has advanced which might be the reason for this observation. Obviously, there will never be a randomized clinical trial comparing resection with biopsy in patients with lesions suspicious for a cerebral lymphoma. Therefore, based on the currently available evidence, complete resection should be considered in case of single lesions, which may be removed safely. In the case of multiple lesions or localization of the tumor mass in an eloquent region of the brain, open or stereotactic-guided biopsy will remain the gold standard because of the overall low risk of complications of such a procedure.

Radiation therapy

Radiation therapy has been used for decades for the treatment of PCNSL, but its role has been diminishing over the last years. Treatment of PCNSL has traditionally been considered to require the whole brain as the target volume

(WBRT). Standard WBRT doses are typically 40-50 Gray (Gy) with single fractions of not more than 2 Gy. Administration of higher doses does not result in prolonged disease control, but causes more neurotoxic side effects [21]. Trials, which assessed radiotherapy compared with supportive care, did not exist, but there is no doubt that WBRT is active against PCNSL. However, although high response rates can be achieved, they are commonly short-lasting. Furthermore, there has been increasing evidence that irradiation is associated with considerable neurotoxicity, particularly in combination with MTX-based chemotherapy and in elderly patients [22]. Major questions, which have been addressed, but only partially resolved within the last years, included the following:

- Is there any role for dose-reduced irradiation?
- Does consolidating radiation therapy after MTX-based chemotherapy prolong progression-free (PFS) and overall survival (OS)?
- When irradiation is deferred in patients who achieve a complete remission (CR) by chemotherapy alone, is it still active as a salvage treatment?

Several analyses indicate that the addition of WBRT to HD-MTX is associated with a considerable risk of cognitive impairment. In contrast, a profoundly lower incidence of neurotoxic effects is observed when HD-MTX was applied alone [23, 24••]. Because of these problems, the concept of dose-reduced WBRT was developed aiming at avoiding cognitive side effects, but still being active as consolidation. A report by Shaw and colleagues indicated a lack of neurocognitive decline when patients received consolidating WBRT with 23.4 Gy in 1.8 Gy/fraction after HD-MTX-based chemotherapy [25]. Owing to the lack of a control group, a benefit of the consolidation treatment could not be determined.

However, because of the futility of (higher) standard dose WBRT to prolong OS following MTX-based chemotherapy in the G-PCNSL-SG1 trial (see below), it can be anticipated that RT with a decreased dose will not provide more benefit in terms of survival unless an unusual dose response relationship of RT for PCNSL was postulated. When RT is deferred, it is still active in the situation of tumor recurrence. A retrospective case collection comprising 48 patients demonstrated an overall response rate of 79 % and a median OS of 16 months [26] suggesting that WBRT is an active treatment for patients with recurrent or refractory PCNSL.

Chemotherapy

Chemotherapy is regarded the treatment of first choice for patients newly diagnosed with PCNSL. Chemotherapy-only approaches can result in high response rates and, even more important, long-lasting remissions. There is a debate, however, on the most active compounds to be added to the established backbone of high-dose (HD)-MTX chemotherapy. The lack of benefit from many drugs which are used for the treatment of systemic lymphoma including the CHOP regimen is most likely due to their insufficient penetration of the blood-brain barrier [27].

Methotrexate alone or in combination

MTX doses that are commonly administered are in the range of 3 to 8 g/m², typically infused over 3 hours in 2 week-intervals. Doses less than 3 g/m² are

probably too low to reach sufficient concentrations in the CSF to kill lymphoma cells [28, 29]. A dose of 8 g/m² MTX was frequently associated with the need for a dose reduction during the course of the treatment, mostly due to renal toxicity [30, 31]. Therefore, most investigators chose a dose of MTX in the range of 3–4 g/m² for combined therapeutic approaches with additional drugs. Various combinations of MTX with other agents have been evaluated within the last three decades. However, the design of these corresponding reports, commonly either from retrospective case collections and limited size or both, and the data from few uncontrolled, but at least prospective series does not allow for final conclusions and precludes definition of a standard of care beyond HD-MTX (reviewed in [32]). In fact, there are only four randomized trials in PCNSL of which one was closed early for poor patient accrual. Drugs, which have been frequently used in combination with MTX, include cytarabine, ifosfamide, temozolomide, procarbazine, vincristine and rituximab. Various regimens have been assessed aiming at increasing response rates and duration of response. The only randomized trial comparing MTX alone (3.5 g/m²) with MTX (3.5 g/m²) plus high-dose cytarabine (four cycles of 2 g/m²) followed by WBRT as consolidation treatment in both arms demonstrated a superior response rate to the chemotherapy part in the MTX/cytarabine arm (CRR=46 %) compared to only 18 % in the cohort of patients receiving MTX alone [19]. The disappointingly and exceptionally poor results of the single agent MTX are likely because MTX was administered only every 3 weeks rather than the usual every 2 weeks, leading to a decreased and insufficient dose intensity. This is also reflected by the fact that only 15 % of the patients in the MTX only arm experienced grade 3/4 hematological toxicity. There was also a trend towards prolonged survival with the combination (3 year OS: 46 % vs. 32 %). An analysis of the G-PCNSL-SG1 trial indicates that increased response rates do not necessarily translate into prolonged OS. In this study, the addition of ifosfamide to HD-MTX resulted in increased CR rates but no superiority in terms of OS compared to HD-MTX alone was observed. In line with these findings, a comprehensive Cochrane database review concluded that the addition of other drugs to MTX may result in increased response rates as well as prolonged PFS whereas the benefit on OS remains unclear. In contrast, toxicity is significantly higher with combined treatment regimens [33]. Therefore, randomized trials with higher patient numbers and longer follow-up are required to determine ultimately the benefit from polychemotherapy as opposed to HD-MTX alone.

Chemotherapy and additional radiotherapy

Consolidating radiation therapy has been explored as an adjunct to chemotherapy for at least 20 years. There are numerous uncontrolled phase II studies exploring MTX-based regimens with additional WBRT resulting in variable response rates and survival data. These trials cannot be compared to each other because of their different designs [34–37]. However, it became clear that the addition of WBRT to chemotherapy is associated with considerable neurotoxicity, particularly in elderly patients. Only in 2010, the results of the randomized G-PCNSL-SG1 trial, which enrolled more than 500 patients with newly diagnosed PCNSL, were published [18]. The addition of WBRT to MTX-based chemotherapy resulted in prolonged PFS (18 months

for the WBRT group versus 12 months for patient who did not receive WBRT) whereas no difference in OS was observed (32 and 37 months, respectively). The trial was criticized for the long time needed for patient accrual, the high number of patients lost to follow-up and specifically the fact that approximately one in four patients did eventually not adhere to the randomization result, an inherent problem if patients are randomized several months prior to the time point where randomization becomes effective. However, the *per-protocol* as well as the *intention-to-treat* analysis yielded similar results for the major outcome measures, indicating that irradiation can be deferred until relapse after MTX-based first-line therapy. WBRT retains its role as a valuable treatment option in patients who fail to respond sufficiently to initial chemotherapy or do not qualify for MTX-based treatment, but outcome in relapsed PCNSL is commonly poor and long-term survival after WBRT would likely be long-term survival with cognitive impairment.

Other approaches for consolidation therapy

Because of the failure of WBRT to prolong OS when used as a consolidation treatment and the significant long-term toxicity associated with its administration, other strategies aiming at avoiding immediate irradiation have been evaluated. One of the most promising approaches is high-dose chemotherapy (HDC) with subsequent autologous stem cell transplantation (ASCT), while omitting WBRT. This concept that has emerged over the last 15 years may be a promising strategy, in particular for younger and fit patients. HDC followed by ASCT has been examined in relatively small and uncontrolled trials both in patients with newly diagnosed as well as refractory and recurrent PCNSL. Various conditioning therapies have been used including a combination of thiotepa together with busulfane or carmustine (BCNU) or the BEAM regimen (carmustine, etoposide, cytarabine, melphalan). Several smaller trials combined a HDC approach with additional irradiation yielding excellent survival results that, however, might be again compromised by late neurotoxicity [38–40]. Overall, MTX-based chemotherapy followed by high-dose therapy with carmustine and thiotepa and subsequent ASCT with or without WBRT resulted in promising long-term survival with a 5-year OS of 70 % [41]. Further studies reported promising results without significant neurotoxicity when HDC and subsequent ASCT were administered without additional WBRT [42–44].

Another approach assessed induction chemotherapy with MTX, temozolomide and rituximab followed by consolidation with etoposide and high-dose cytarabine for patients who achieved stable disease or better after induction. The 2-year OS was 58 % when all patients were analyzed and 93 % in the cohort of patients who had qualified for and received consolidation therapy with etoposide and cytarabine without evidence of neurotoxicity [45]. This regimen was further evaluated within a multicenter study resulting in survival curves, which display evidence of a plateau. The median OS had not yet been reached after a median follow-up of 4.9 years [46••]. These results indicate that chemotherapy alone can result in excellent long-term survival. However, the potential benefit of polychemotherapy and HDC/ASCT compared to standard chemotherapy alone needs to be confirmed in randomized trials.

Other drugs

In contrast to the unquestioned importance of MTX as the backbone of any first-line regimen, it is much less clear which other agents should be used in case of tumor recurrence, absence of response or for patients who cannot tolerate HD MTX. In the following section, we summarize some of the drugs that are frequently used either alone or as part of combined regimens.

Steroids

Similar to other brain tumors, the administration of corticosteroids results in a reduction of the peritumoral edema [47]. However, steroids also have a transient direct anti-tumor effect against lymphoma cells through induction of apoptosis, which can result in a rapid decrease of the size of the tumor mass. When steroids are administered before surgery, they may preclude histological confirmation of the diagnosis.

Rituximab

The putative beneficial role of rituximab, a monoclonal antibody targeting the cell surface antigen CD20, which is found in the majority of PCNSL, remains a matter of debate. Being part of most therapeutic regimens against systemic B cell lymphomas, its activity against lymphomas in the CNS is doubtful. CSF levels of rituximab after systemic application are approximately 0.1 % of serum levels, a phenomenon which is mainly caused by the presence of the blood-brain barrier which hampers the penetration of the antibody into the CSF [48]. Still, monotherapy with rituximab in the setting of recurrent PCNSL was reported to yield occasional radiographic responses in up to 33 % of patients without major toxicity concerns [49]. This may reflect the partially disrupted blood-brain barrier in PCNSL, which may still allow the antibody reaching some of brain lymphoma cells. Series looking at the efficacy of rituximab when added to a MTX-based regimen suggest a certain activity in PCNSL which is associated with increased CR rates that may eventually translate into prolonged OS without adding significant toxicity [50, 51]. However, these studies have limited value due to their retrospective nature. The results of the ongoing prospective HOVON 105 (EudraCT no. 2009-014722-42) and IELSG32 (NCT01011920) trials, which explore the activity of rituximab in a randomized fashion need to be awaited. Similarly, the role of intrathecal administration of rituximab has not yet been determined (see below).

Temozolomide

Temozolomide crosses the blood-barrier and has been included in several MTX-based regimens (see above). It has been mainly examined in patients with recurrent PCNSL who had previously been treated with MTX. Here,

single-agent temozolomide was typically well tolerated whereas its activity was overall limited with CR rates in the range of 20-30 % [52, 53]. Temozolomide alone, when used as first-line therapy in elderly patients who did not qualify for MTX-based chemotherapy, had also only modest activity (see below). The efficacy of temozolomide may depend on the activity of the DNA repair protein O⁶-methylguanine DNA methyltransferase (MGMT). In retrospective series, low MGMT protein expression examined by immunohistochemistry or high MGMT promoter methylation determined by methylation-sensitive high resolution melting (HS-HRM) analysis were predictive for response to temozolomide [54, 55]. However, larger series examining the predictive or prospective value of the MGMT status in PCNSL are lacking.

Topotecan

Topotecan is a topoisomerase I inhibitor that has been administered to patients with recurrent or MTX-refractory PCNSL. Treatment with topotecan can induce considerable response rates but may be associated with significant toxicity [56, 57].

Intrathecal therapy

Dissemination of lymphoma cells into the CSF is a common finding although the reported incidence rates are varying most likely due to different methods that were applied for the detection of meningeal affection. The presence of lymphoma cells in the CSF is probably not a negative prognostic marker [58]. Still, appropriate treatment of the CSF may be a prerequisite for long-lasting remissions. MTX administered at doses of 3 g/m² or higher will adequately penetrate through the blood-brain barrier to yield therapeutic CSF levels [59]. Therefore, intrathecal therapy is not considered as a standard of care which is supported by two retrospective analyses suggesting that additional intrathecal treatment does not result in prolonged OS compared to patient cohorts that were not treated intrathecally [60, 61]. In contrast, very promising results of a trial using a polychemotherapy approach including intraventricular therapy with MTX, prednisolone, and cytarabine were not reproduced when the intrathecal treatment was omitted [62–64]. Intraventricular administration of rituximab has been examined in a small cohort of PCNSL patients. The drug was well tolerated but the beneficial effects of this approach remain elusive [65]. Application of drugs by an Ommaya reservoir or a similar device are frequently associated with infectious complications and require an additional surgical procedure. This may be circumvented by the repeated application of drugs by lumbar puncture which, however, may not result in an equal distribution of the compound in the CSF. Whether the addition of lumbar or intraventricular treatment to MTX-based protocols results in

prolonged OS needs further investigation and should be assessed within appropriate clinical trials.

Elderly patients

Patients over age 60 are commonly referred to as “elderly”. However, some studies focusing on elderly patients only include patients who are at least 65 or even 70 years old. Owing to this variability, it becomes difficult to compare different studies enrolling elderly patients. As with younger patients, MTX should be the backbone of any treatment concept in elderly patients because of the lack of drugs with similar activity. In the elderly, HD-MTX is commonly well tolerated and can be safely administered in most patients [66]. Even in patients aged 80 and older, the application of MTX-containing regimens was feasible and resulted in long-term survival in some patients [67]. However, exceedingly high serum MTX levels, determined by area under the curve (AUC) calculations, are associated with reduced survival and should be avoided [68]. Promising CR rates can be achieved in elderly patients. However, the durability of these responses is reduced compared to younger patients indicating that consolidation is required in elderly patients in CR [66]. Again, various compounds have been added to MTX and assessed in smaller trials. The combination of MTX with procarbazine and lomustine resulted in a median OS of 15.4 months [69] which was only moderately changed to 17.5 months when rituximab was added in a subsequent trial [70]. A polychemotherapy regimen and additional intrathecal chemotherapy yielded a median OS of 14.3 months in a study with 50 PCNSL patients who had a median age of 72 years [71]. The combination of HD-MTX with temozolomide was reported to be well tolerated and resulted in an OS of 25 months [72]. However, a randomized phase II study comparing two different MTX-based regimens suggests higher activity of MPV-A (MTX, procarbazine, vincristine and cytarabine) compared to MTX combined with temozolomide [73•]. This trial, however, was not designed to prove superiority and these results must be confirmed in a phase III trial. No convincing treatment strategies are available for elderly patients who cannot be treated with HD-MTX. Radiation therapy is active but has several limitations as described above. Dose-intensified temozolomide alone has only limited efficacy [74]. In summary, similar to younger patients, HD-MTX should be used in the elderly whenever possible and considered appropriate. The benefit of further agents that are added to MTX remains to be determined.

Recurrent or refractory PCNSL

In the situation of recurrent or refractory PCNSL, no standard treatment has been defined. Several options are available and the parameters, which should be taken into account, include the performance status of the patient, benefit from prior therapies and the interval between last treatment and tumor recurrence. MTX re-challenge is generally considered promising in patients who

achieved a CR upon MTX-based treatment [75]. In contrast, patients whose tumors must be considered refractory to MTX should be offered an alternative treatment. Here, various approaches have been assessed. Similar to the situation of newly diagnosed PCNSL, the quality of the available studies is rather poor and randomized trials are lacking. WBRT is an active treatment in the setting of recurrent PCNSL with high response rates, which, however, frequently do not translate into long-lasting remissions [26, 76]. Temozolomide alone is well tolerated but has only modest activity [53]. The combination of temozolomide and rituximab was examined within a trial that was closed prematurely after a preliminary analysis had suggested disappointing activity [77]. Novel approaches include the administration of pemetrexed, a folate antimetabolite. In patients with relapsed or refractory PCNSL, it had single-agent activity but was also associated with considerable toxicity [78, 79]. Larger trials are required to determine the value of pemetrexed in PCNSL in more detail. Finally, HDC with subsequent ASCT may be a promising option for selected patients with recurrent PCNSL who qualify for such an intense and rather toxic approach [80, 81].

Compliance with Ethics Guidelines

Conflict of Interest

Patrick Roth has served as a consultant for Roche, Merck Sharp & Dohme, and Molecular Partners. Roger Stupp has served as a consultant for Merck KGaA, Merck Sharp & Dohme/Merck & Co., and Genentech/Roche. Michael Weller has served as a consultant for MagForce AG; has received grant support from Merck Serono, Roche, Antisense Pharma, and Bayer, and has received honoraria from Roche, Merck Serono, and Merck Sharp & Dohme. Günter Eisele declares that he has no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

References and Recommended Reading

Papers of particular interest, published recently, have been highlighted as:

- Of importance
 - Of major importance
1. Dolecek TA, Propp JM, Stroup NE, Kruchko C. CBTRUS statistical report: primary brain and central nervous system tumors diagnosed in the United States in 2005-2009. *Neuro Oncol.* 2012;14 Suppl 5:v1-49.
 2. O'Neill BP, Decker PA, Tieu C, Cerhan JR. The changing incidence of primary central nervous system lymphoma is driven primarily by the changing incidence in young and middle-aged men and differs from time trends in systemic diffuse large B-cell non-Hodgkin's lymphoma. *Am J Hematol* 2013 [epub ahead of print].
 3. Buehring U, Herrlinger U, Krings T, Thiex R, Weller M, Kuker W. MRI features of primary central nervous system lymphomas at presentation. *Neurology.* 2001;57:393-6.
 4. Farrell BT, Hamilton BE, Dosa E, Rimely E, Nasser M, Gahramanov S, et al. Using iron oxide nanoparticles to diagnose CNS inflammatory diseases and PCNSL. *Neurology.* 2013;81:256-63.

5. Maza S, Buchert R, Brenner W, Munz DL, Thiel E, Korfel A, et al. Brain and whole-body FDG-PET in diagnosis, treatment monitoring and long-term follow-up of primary CNS lymphoma. *Radiol Oncol*. 2013;47:103–10.
 6. Kawai N, Miyake K, Yamamoto Y, Nishiyama Y, Tamiya T. 18F-FDG PET in the diagnosis and treatment of primary central nervous system lymphoma. *BioMed Res Int*. 2013;2013:247152.
 7. Kawase Y, Yamamoto Y, Kameyama R, Kawai N, Kudomi N, Nishiyama Y. Comparison of 11C-methionine PET and 18F-FDG PET in patients with primary central nervous system lymphoma. *Mol Imaging Biol*. 2011;13:1284–9.
 8. Abrey LE, Batchelor TT, Ferreri AJ, Gospodarowicz M, Pulczynski EJ, Zucca E, et al. Report of an international workshop to standardize baseline evaluation and response criteria for primary CNS lymphoma. *J Clin Oncol*. 2005;23:5034–43.
 9. Mohile NA, Deangelis LM, Abrey LE. The utility of body FDG PET in staging primary central nervous system lymphoma. *Neuro Oncol*. 2008;10:223–8.
 10. Chan CC, Rubenstein JL, Coupland SE, Davis JL, Harbour JW, Johnston PB, et al. Primary vitreoretinal lymphoma: a report from an international primary central nervous system lymphoma collaborative group symposium. *Oncologist*. 2011;16:1589–99.
 11. Jahnke K, Hummel M, Korfel A, Burmeister T, Kiewe P, Klasen HA, et al. Detection of subclinical systemic disease in primary CNS lymphoma by polymerase chain reaction of the rearranged immunoglobulin heavy-chain genes. *J Clin Oncol*. 2006;24:4754–7.
 12. Deckert M, Brunn A, Montesinos-Rongen M, Terreni MR, Ponzoni M. Primary lymphoma of the central nervous system—a diagnostic challenge. *Hematol Oncol* 2013 [epub ahead of print].
 13. Jahnke K, Korfel A, O'Neill BP, Blay JY, Abrey LE, Martus P, et al. International study on low-grade primary central nervous system lymphoma. *Ann Neurol*. 2006;59:755–62.
 14. Shenker TN, Blay JY, O'Neill BP, Poortmans P, Thiel E, Jahnke K, et al. Primary CNS lymphoma of T-cell origin: a descriptive analysis from the international primary CNS lymphoma collaborative group. *J Clin Oncol*. 2005;23:2233–9.
 15. Abrey LE, Ben-Porat L, Panageas KS, Yahalom J, Berkey B, Curran W, et al. Primary central nervous system lymphoma: the Memorial Sloan-Kettering Cancer Center prognostic model. *J Clin Oncol*. 2006;24:5711–5.
 16. Ferreri AJ, Blay JY, Reni M, Pasini F, Spina M, Ambrosetti A, et al. Prognostic scoring system for primary CNS lymphomas: the International Extranodal Lymphoma Study Group experience. *J Clin Oncol*. 2003;21:266–72.
 17. Lachance DH, Brizel DM, Gockerman JP, Halperin EC, Burger PC, Boyko OB, et al. Cyclophosphamide, doxorubicin, vincristine, and prednisone for primary central nervous system lymphoma: short-duration response and multifocal intracerebral recurrence preceding radiotherapy. *Neurology*. 1994;44:1721–7.
 18. Thiel E, Korfel A, Martus P, Kanz L, Griesinger F, Rauch M, et al. High-dose methotrexate with or without whole brain radiotherapy for primary CNS lymphoma (G-PCNSL-SG-1): a phase 3, randomised, non-inferiority trial. *Lancet Oncol*. 2010;11:1036–47.
 19. Ferreri AJ, Reni M, Foppoli M, Martelli M, Pangalis GA, Frezzato M, et al. High-dose cytarabine plus high-dose methotrexate versus high-dose methotrexate alone in patients with primary CNS lymphoma: a randomised phase 2 trial. *Lancet*. 2009;374:1512–20.
 20. • Weller M, Martus P, Roth P, Thiel E, Korfel A, German PSG. Surgery for primary CNS lymphoma? Challenging a paradigm. *Neuro Oncol*. 2012;14:1481–4.
- Analysis of a large dataset suggesting that complete resection may be associated with prolonged survival in PCNSL patients.
21. Nelson DF. Radiotherapy in the treatment of primary central nervous system lymphoma (PCNSL). *J Neurooncol*. 1999;43:241–7.
 22. O'Brien PC, Roos DE, Pratt G, Liew KH, Barton MB, Poulsen MG, et al. Combined-modality therapy for primary central nervous system lymphoma: long-term data from a Phase II multicenter study (Trans-Tasman Radiation Oncology Group). *Int J Radiat Oncol Biol Phys*. 2006;64:408–13.
 23. Correa DD, Shi W, Abrey LE, Deangelis LM, Omuro AM, Deutsch MB, et al. Cognitive functions in primary CNS lymphoma after single or combined modality regimens. *Neuro Oncol*. 2012;14:101–8.
 24. •• Doolittle ND, Korfel A, Lubow MA, Schorb E, Schlegel U, Rogowski S, et al. Long-term cognitive function, neuroimaging, and quality of life in primary CNS lymphoma. *Neurology*. 2013;81:84–92.
- A comprehensive analysis of cognitive function and quality of life in PCNSL patients.
25. Shah GD, Yahalom J, Correa DD, Lai RK, Raizer JJ, Schiff D, et al. Combined immunochemotherapy with reduced whole-brain radiotherapy for newly diagnosed primary CNS lymphoma. *J Clin Oncol*. 2007;25:4730–5.
 26. Hottinger AF, DeAngelis LM, Yahalom J, Abrey LE. Salvage whole brain radiotherapy for recurrent or refractory primary CNS lymphoma. *Neurology*. 2007;69:1178–82.
 27. Mead GM, Bleehen NM, Gregor A, Bullimore J, Shirley D, Rampling RP, et al. A medical research

- council randomized trial in patients with primary cerebral non-Hodgkin lymphoma: cerebral radiotherapy with and without cyclophosphamide, doxorubicin, vincristine, and prednisone chemotherapy. *Cancer*. 2000;89:1359–70.
28. Shapiro WR, Young DF, Mehta BM. Methotrexate: distribution in cerebrospinal fluid after intravenous, ventricular and lumbar injections. *N Engl J Med*. 1975;293:161–6.
 29. Ferreri AJ, Guerra E, Regazzi M, Pasini F, Ambrosetti A, Pivnik A, et al. Area under the curve of methotrexate and creatinine clearance are outcome-determining factors in primary CNS lymphomas. *Br J Cancer*. 2004;90:353–8.
 30. Batchelor T, Carson K, O'Neill A, Grossman SA, Alavi J, New P, et al. Treatment of primary CNS lymphoma with methotrexate and deferred radiotherapy: a report of NABTT 96-07. *J Clin Oncol*. 2003;21:1044–9.
 31. Herrlinger U, Kuker W, Uhl M, Blaicher HP, Karnath HO, Kanz L, et al. NOA-03 trial of high-dose methotrexate in primary central nervous system lymphoma: final report. *Ann Neurol*. 2005;57:843–7.
 32. Korfel A, Schlegel U. Diagnosis and treatment of primary CNS lymphoma. *Nat Rev Neurol*. 2013;9:317–27.
 33. Bergner N, Monsef I, Illerhaus G, Engert A, Skoetz N. Role of chemotherapy additional to high-dose methotrexate for primary central nervous system lymphoma (PCNSL). *Cochrane Database Syst Rev*. 2012;11, CD009355.
 34. Bessell EM, Lopez-Guillermo A, Villa S, Verger E, Nomdedeu B, Petit J, et al. Importance of radiotherapy in the outcome of patients with primary CNS lymphoma: an analysis of the CHOD/BVAM regimen followed by two different radiotherapy treatments. *J Clin Oncol*. 2002;20:231–6.
 35. DeAngelis LM, Seiferheld W, Schold SC, Fisher B, Schultz CJ. Combination chemotherapy and radiotherapy for primary central nervous system lymphoma: Radiation Therapy Oncology Group Study 93-10. *J Clin Oncol*. 2002;20:4643–8.
 36. Poortmans PM, Kluin-Nelemans HC, Haaxma-Reiche H, Van't Veer M, Hansen M, Soubeyran P, et al. High-dose methotrexate-based chemotherapy followed by consolidating radiotherapy in non-AIDS-related primary central nervous system lymphoma: European Organization for Research and Treatment of Cancer Lymphoma Group Phase II Trial 20962. *J Clin Oncol*. 2003;21:4483–8.
 37. Ferreri AJ, Dell'Oro S, Foppoli M, Bernardi M, Brandes AA, Tosoni A, et al. MATILDE regimen followed by radiotherapy is an active strategy against primary CNS lymphomas. *Neurology*. 2006;66:1435–8.
 38. Illerhaus G, Marks R, Ihorst G, Guttenberger R, Ostertag C, Derigs G, et al. High-dose chemotherapy with autologous stem-cell transplantation and hyperfractionated radiotherapy as first-line treatment of primary CNS lymphoma. *J Clin Oncol*. 2006;24:3865–70.
 39. Colombat P, Lemevel A, Bertrand P, Delwail V, Rachieru P, Brion A, et al. High-dose chemotherapy with autologous stem cell transplantation as first-line therapy for primary CNS lymphoma in patients younger than 60 years: a multicenter phase II study of the GOELAMS group. *Bone Marrow Transplant*. 2006;38:417–20.
 40. Montemurro M, Kiefer T, Schuler F, Al-Ali HK, Wolf HH, Herbst R, et al. Primary central nervous system lymphoma treated with high-dose methotrexate, high-dose busulfan/thiotepa, autologous stem-cell transplantation and response-adapted whole-brain radiotherapy: results of the multicenter Ostdeutsche Studiengruppe Hamato-Onkologie OSHO-53 phase II study. *Ann Oncol*. 2007;18:665–71.
 41. Kasenda B, Schorb E, Fritsch K, Finke J, Illerhaus G. Prognosis after high-dose chemotherapy followed by autologous stem-cell transplantation as first-line treatment in primary CNS lymphoma—a long-term follow-up study. *Ann Oncol*. 2012;23:2670–5.
 42. Abrey LE, Moskowitz CH, Mason WP, Crump M, Stewart D, Forsyth P, et al. Intensive methotrexate and cytarabine followed by high-dose chemotherapy with autologous stem-cell rescue in patients with newly diagnosed primary CNS lymphoma: an intent-to-treat analysis. *J Clin Oncol*. 2003;21:4151–6.
 43. Kiefer T, Hirt C, Spath C, Schuler F, Al-Ali HK, Wolf HH, et al. Long-term follow-up of high-dose chemotherapy with autologous stem-cell transplantation and response-adapted whole-brain radiotherapy for newly diagnosed primary CNS lymphoma: results of the multicenter Ostdeutsche Studiengruppe Hamatologie und Onkologie OSHO-53 phase II study. *Ann Oncol*. 2012;23:1809–12.
 44. Alimohamed N, Daly A, Owen C, Duggan P, Stewart DA. Upfront thiotepa, busulfan, cyclophosphamide, and autologous stem cell transplantation for primary CNS lymphoma: a single centre experience. *Leuk Lymphoma*. 2012;53:862–7.
 45. Wieduwilt MJ, Valles F, Issa S, Behler CM, Hwang J, McDermott M, et al. Immunotherapy with intensive consolidation for primary CNS lymphoma: a pilot study and prognostic assessment by diffusion-weighted MRI. *Clin Cancer Res*. 2012;18:1146–55.
 - 46.●● Rubenstein JL, Hsi ED, Johnson JL, Jung SH, Nakashima MO, Grant B, et al. Intensive Chemotherapy and Immunotherapy in Patients With Newly Diagnosed Primary CNS Lymphoma: CALGB 50202 (Alliance 50202). *J Clin Oncol*. 2013;31:3061–8.
- This study demonstrates that long-term survival without disabling neurotoxicity can be achieved with a chemotherapy-alone regimen.

47. Roth P, Wick W, Weller M. Steroids in neurooncology: actions, indications, side-effects. *Curr Opin Neurol*. 2010;23:597–602.
48. Rubenstein JL, Combs D, Rosenberg J, Levy A, McDermott M, Damon L, et al. Rituximab therapy for CNS lymphomas: targeting the leptomeningeal compartment. *Blood*. 2003;101:466–8.
49. Batchelor TT, Grossman SA, Mikkelsen T, Ye X, Desideri S, Lesser GJ. Rituximab monotherapy for patients with recurrent primary CNS lymphoma. *Neurology*. 2011;76:929–30.
50. Birnbaum T, Stadler EA, von Baumgarten L, Straube A. Rituximab significantly improves complete response rate in patients with primary CNS lymphoma. *J Neurooncol*. 2012;109:285–91.
51. Chamberlain MC, Johnston SK. High-dose methotrexate and rituximab with deferred radiotherapy for newly diagnosed primary B-cell CNS lymphoma. *Neuro Oncol*. 2010;12:736–44.
52. Reni M, Mason W, Zaja F, Perry J, Franceschi E, Bernardi D, et al. Salvage chemotherapy with temozolomide in primary CNS lymphomas: preliminary results of a phase II trial. *Eur J Cancer*. 2004;40:1682–8.
53. Makino K, Nakamura H, Hide T, Kuratsu J. Salvage treatment with temozolomide in refractory or relapsed primary central nervous system lymphoma and assessment of the MGMT status. *J Neurooncol*. 2012;106:155–60.
54. Jiang X, Reardon DA, Desjardins A, Vredenburgh JJ, Quinn JA, Austin AD, et al. O(6)-methylguanine-DNA methyltransferase (MGMT) immunohistochemistry as a predictor of resistance to temozolomide in primary CNS lymphoma. *J Neurooncol*. 2013;114:135–40.
55. Adachi J, Mishima K, Wakiya K, Suzuki T, Fukuoka K, Yanagisawa T, et al. O(6)-methylguanine-DNA methyltransferase promoter methylation in 45 primary central nervous system lymphomas: quantitative assessment of methylation and response to temozolomide treatment. *J Neurooncol*. 2012;107:147–53.
56. Fischer L, Thiel E, Klasen HA, Birkmann J, Jahnke K, Martus P, et al. Prospective trial on topotecan salvage therapy in primary CNS lymphoma. *Ann Oncol*. 2006;17:1141–5.
57. Voloschin AD, Betensky R, Wen PY, Hochberg F, Batchelor T. Topotecan as salvage therapy for relapsed or refractory primary central nervous system lymphoma. *J Neurooncol*. 2008;86:211–5.
58. Korfel A, Weller M, Martus P, Roth P, Klasen HA, Roeth A, et al. Prognostic impact of meningeal dissemination in primary CNS lymphoma (PCNSL): experience from the G-PCNSL-SG1 trial. *Ann Oncol*. 2012;23:2374–80.
59. Abrey LE, Yahalom J, DeAngelis LM. Treatment for primary CNS lymphoma: the next step. *J Clin Oncol*. 2000;18:3144–50.
60. Khan RB, Shi W, Thaler HT, DeAngelis LM, Abrey LE. Is intrathecal methotrexate necessary in the treatment of primary CNS lymphoma? *J Neurooncol*. 2002;58:175–8.
61. Sierra Del Rio M, Ricard D, Houillier C, Navarro S, Gonzalez-Aguilar A, Idbaih A, et al. Prophylactic intrathecal chemotherapy in primary CNS lymphoma. *J Neurooncol*. 2012;106:143–6.
62. Pels H, Schmidt-Wolf IG, Glasmacher A, Schulz H, Engert A, Diehl V, et al. Primary central nervous system lymphoma: results of a pilot and phase II study of systemic and intraventricular chemotherapy with deferred radiotherapy. *J Clin Oncol*. 2003;21:4489–95.
63. Pels H, Juergens A, Glasmacher A, Schulz H, Engert A, Linnebank M, et al. Early relapses in primary CNS lymphoma after response to polychemotherapy without intraventricular treatment: results of a phase II study. *J Neurooncol*. 2009;91:299–305.
64. Juergens A, Pels H, Rogowski S, Fliessbach K, Glasmacher A, Engert A, et al. Long-term survival with favorable cognitive outcome after chemotherapy in primary central nervous system lymphoma. *Ann Neurol*. 2010;67:182–9.
65. Rubenstein JL, Li J, Chen L, Advani R, Drappatz J, Gerstner E, et al. Multicenter phase 1 trial of intraventricular immunochemotherapy in recurrent CNS lymphoma. *Blood*. 2013;121:745–51.
66. Roth P, Martus P, Kiewe P, Mohle R, Klasen H, Rauch M, et al. Outcome of elderly patients with primary CNS lymphoma in the G-PCNSL-SG-1 trial. *Neurology*. 2012;79:890–6.
67. Welch MR, Omuro A, DeAngelis LM. Outcomes of the oldest patients with primary CNS lymphoma treated at Memorial Sloan-Kettering Cancer Center. *Neuro Oncol*. 2012;14:1304–11.
68. Kasenda B, Rehberg M, Thurmann P, Franzem M, Veelken H, Fritsch K, et al. The prognostic value of serum methotrexate area under curve in elderly primary CNS lymphoma patients. *Ann Hematol*. 2012;91:1257–64.
69. Illerhaus G, Marks R, Muller F, Ihorst G, Feuerhake F, Deckert M, et al. High-dose methotrexate combined with procarbazine and CCNU for primary CNS lymphoma in the elderly: results of a prospective pilot and phase II study. *Ann Oncol*. 2009;20:319–25.
70. Fritsch K, Kasenda B, Hader C, Nikkhah G, Prinz M, Haug V, et al. Immunochemotherapy with rituximab, methotrexate, procarbazine, and lomustine for primary CNS lymphoma (PCNSL) in the elderly. *Ann Oncol*. 2011;22:2080–5.
71. Hoang-Xuan K, Taillandier L, Chinot O, Soubeyran P, Bogdhan U, Hildebrand J, et al. Chemotherapy

- alone as initial treatment for primary CNS lymphoma in patients older than 60 years: a multicenter phase II study (26952) of the European Organization for Research and Treatment of Cancer Brain Tumor Group. *J Clin Oncol*. 2003;21:2726–31.
72. Omuro AM, Taillandier L, Chinot O, Carin C, Barrie M, Hoang-Xuan K. Temozolomide and methotrexate for primary central nervous system lymphoma in the elderly. *J Neurooncol*. 2007;85:207–11.
73. • Omuro AM, Chinot OL, Taillandier L, Ghesquieres H, Soussain C, Delwail V, et al. Multicenter randomized phase II trial of methotrexate (MTX) and temozolomide (TMZ) versus MTX, procarbazine, vincristine, and cytarabine for primary CNS lymphoma (PCNSL) in the elderly: An Anocéf and Goelams Intergroup study. *J Clin Oncol*. 2013;31. No 15_suppl 2032.
- A randomized phase II trial suggesting superior activity of MTX, procarbazine, vincristine and cytarabine compared to MTX and temozolomide in elderly PCNSL patients.
74. Kurzwelly D, Glas M, Roth P, Weimann E, Lohner H, Waha A, et al. Primary CNS lymphoma in the elderly: temozolomide therapy and MGMT status. *J Neurooncol*. 2010;97:389–92.
75. Plotkin SR, Betensky RA, Hochberg FH, Grossman SA, Lesser GJ, Nabors LB, et al. Treatment of relapsed central nervous system lymphoma with high-dose methotrexate. *Clin Cancer Res*. 2004;10:5643–6.
76. Nguyen PL, Chakravarti A, Finkelstein DM, Hochberg FH, Batchelor TT, Loeffler JS. Results of whole-brain radiation as salvage of methotrexate failure for immunocompetent patients with primary CNS lymphoma. *J Clin Oncol*. 2005;23:1507–13.
77. Nayak L, Abrey LE, Drappatz J, Gilbert MR, Reardon DA, Wen PY, et al. North American Brain Tumor C. Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma*. 2013;54:58–61.
78. Raizer JJ, Rademaker A, Evens AM, Rice L, Schwartz M, Chandler JP, et al. Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. *Cancer*. 2012;118:3743–8.
79. Zhang JP, Lee EQ, Nayak L, Doherty L, Kesari S, Muzikansky A, et al. Retrospective study of pemetrexed as salvage therapy for central nervous system lymphoma. *J Neurooncol* 2013 [epub ahead of print].
80. Soussain C, Suzan F, Hoang-Xuan K, Cassoux N, Levy V, Azar N, et al. Results of intensive chemotherapy followed by hematopoietic stem-cell rescue in 22 patients with refractory or recurrent primary CNS lymphoma or intraocular lymphoma. *J Clin Oncol*. 2001;19:742–9.
81. Soussain C, Hoang-Xuan K, Taillandier L, Fourme E, Choquet S, Witz F, et al. Intensive chemotherapy followed by hematopoietic stem-cell rescue for refractory and recurrent primary CNS and intraocular lymphoma: Societe Francaise de Greffe de Moelle Osseuse-Therapie Cellulaire. *J Clin Oncol*. 2008;26:2512–8.