

# Tree-level soft emission of a quark pair in association with a gluon

Vittorio Del Duca,<sup>a,b,1</sup> Claude Duhr,<sup>c</sup> Rayan Haindl<sup>a</sup> and Zhengwen Liu<sup>d</sup>

<sup>a</sup>*Institute for Theoretical Physics, ETH Zürich,  
8093 Zürich, Switzerland*

<sup>b</sup>*Physik-Institut, Universität Zürich,  
Winterthurerstrasse 190, 8057 Zürich, Switzerland*

<sup>c</sup>*Bethe Center for Theoretical Physics, Universität Bonn,  
53115 Bonn, Germany*

<sup>d</sup>*Deutsches Elektronen-Synchrotron DESY,  
Notkestr. 85, 22607 Hamburg, Germany*

*E-mail:* [delducav@itp.phys.ethz.ch](mailto:delducav@itp.phys.ethz.ch), [cduhr@uni-bonn.de](mailto:cduhr@uni-bonn.de),  
[haindlr@phys.ethz.ch](mailto:haindlr@phys.ethz.ch), [zhengwen.liu@desy.de](mailto:zhengwen.liu@desy.de)

**ABSTRACT:** We compute the tree-level current for the emission of a soft quark-antiquark pair in association with a gluon. This soft current is the last missing ingredient to understand the infrared singularities that can arise in next-to-next-to-next-to-leading-order ( $N^3$ LO) computations in QCD. Its square allows us to understand for the first time the colour correlations induced by the soft emission of a quark pair and a gluon. We find that there are three types of correlations: besides dipole-type correlations that have already appeared in soft limits of tree-level amplitudes, we uncover for the first time also a three-parton correlation involving a totally symmetric structure constant. We also study the behaviour of collinear splitting amplitudes in the triple-soft limit, and we derive the corresponding factorisation formula.

**KEYWORDS:** Factorization, Renormalization Group, Higher-Order Perturbative Calculations

ARXIV EPRINT: [2206.01584](https://arxiv.org/abs/2206.01584)

<sup>1</sup>On leave from INFN, Laboratori Nazionali di Frascati, Italy.

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Tree-level soft currents</b>	<b>3</b>
2.1	The tree-level soft current for the emission of a single soft gluon	4
2.2	The tree-level soft current for the emission of a single quark pair	6
<b>3</b>	<b>Tree-level factorisation for soft <math>\bar{q}qg</math> emission</b>	<b>6</b>
3.1	The soft current	7
3.2	Strongly-ordered soft limits	12
<b>4</b>	<b>Soft limits of the splitting amplitudes</b>	<b>14</b>
4.1	Review of collinear factorisation	14
4.2	$m$ -parton soft limit of the $m$ -collinear splitting amplitudes	16
4.3	Soft $\bar{q}qg$ limit of tree-level splitting amplitudes	16
4.3.1	Soft $\bar{q}qg$ limit of the quadruple collinear splitting amplitudes	20
<b>5</b>	<b>Conclusions</b>	<b>21</b>
<b>A</b>	<b>Two-parton colour correlated soft function</b>	<b>22</b>
<b>B</b>	<b>Collinear limit of the two-parton colour correlation</b>	<b>25</b>

---

**1 Introduction**

The cornerstone of all methods to make predictions for modern collider experiments is perturbative Quantum Field Theory, where observables are expanded into a series in the coupling constants. The higher orders in the perturbative expansion capture, on the one hand, the effect of the exchange of virtual quanta, and, on the other hand, the emission of unobserved real particles in the final state. In theories featuring massless particles, both the real and virtual corrections are separately divergent (even after ultraviolet renormalisation), but their sum is finite for a scattering of colourless particles in the initial state (e.g., at an  $e^+e^-$  collider) due to the celebrated Kinoshita-Lee-Nauenberg theorem.<sup>1</sup> The cancellation of these infrared singularities, however, is very intricate, because it only happens after phase space integration, and the real and virtual corrections live in different phase spaces.

The infrared singularities stemming from real corrections arise from regions of phase space where massless particles become unresolved, i.e., they become either soft (meaning

---

<sup>1</sup>If the scattering features also hadrons in the initial state, the divergences only cancel after we perform mass factorisation.

that they have vanishing energies) or collinear to each other. The behaviour of scattering amplitudes in these unresolved limits is universal, in the sense that the amplitudes factorise into amplitudes without the unresolved particles, multiplied by a function that captures the divergence and does not depend on the underlying hard scattering. This universality of soft and collinear divergences is at the heart of so-called subtraction schemes, where, very loosely speaking, one subtracts the phase space divergences of real corrections at the integrand level, and adds them back in integrated form to cancel the infrared singularities of virtual corrections. Subtraction schemes at next-to-leading order (NLO) and next-to-next-to-leading order (NNLO) in the strong coupling constant are one of the cornerstones of modern precision computations in (massless) Quantum Chromodynamics (QCD). The construction and the success of subtraction schemes at NLO and NNLO rely crucially on the fact that the soft and collinear divergences describing tree-level amplitudes with up to two unresolved particles and one-loop amplitudes with one unresolved parton are well understood [1–12].

In order to reach the target precision for current and future collider experiments, like the Large Hadron Collider (LHC) at CERN and its potential successors, NNLO computations in QCD may not be sufficient, but also next-to-next-to-next-to-leading order ( $N^3$ LO) corrections will be required. While first examples of non-trivial two- and three-loop amplitudes relevant to  $N^3$ LO computations have recently become available [13–20], one of the major bottlenecks is that there is currently no general understanding of how to combine the real and virtual corrections.<sup>2</sup> Experience from NNLO shows that it is important to understand in detail all the unresolved limits that lead to singularities in an  $N^3$ LO computation. The relevant collinear singularities are by now completely understood, and they include the emission of four collinear partons at tree-level [28, 29], three collinear partons at one loop [30–32] or two collinear partons at two loops [33]. Soft singularities at  $N^3$ LO have also been studied. In particular, the emission of a soft gluon at two loops is well established [34–36], as is the emission of a pair of soft gluons or quarks at one loop [37, 38]. Tree-level soft emission of three partons, however, has so far only been studied for the emission of three soft gluons [39], but the case of a soft quark pair in addition to a gluon at tree-level has never been considered. The main goal of this paper is to provide for the first time this last soft limit needed to describe all infrared singularities that can arise in  $N^3$ LO computations.

The paper is organised as follows: in section 2, we review soft limits of tree-level scattering amplitudes, displaying explicitly the currents for a soft gluon and a soft  $\bar{q}q$  pair. In section 3, we present the main results of this paper, namely the tree-level current for the soft  $\bar{q}qg$  emission from QCD scattering amplitudes. In section 4, we consider kinematic sub-limits of the soft  $\bar{q}qg$  limit, which are useful for constructing subtraction schemes at  $N^3$ LO accuracy. In section 5 we draw our conclusions. We also include two appendices where we display analytic results which are too lengthy to be shown in the main text.

---

<sup>2</sup>For specific differential distributions for simple final states,  $N^3$ LO results are already available, cf. refs. [21–27].

## 2 Tree-level soft currents

The aim of this paper is to study the behavior of tree-level QCD amplitudes in the limit where a certain number of massless partons are *soft*, i.e., they have vanishing energies. To be more precise, consider the scattering of  $n$  particles with momenta  $p_i$ , and flavour, helicity and colour indices  $f_i$ ,  $s_i$  and  $c_i$ , respectively. If a subset of  $m$  massless partons become soft, the amplitude is divergent and the leading behaviour is captured by the factorisation formula,

$$\begin{aligned} \mathcal{S}_{1\dots m} \mathcal{M}_{f_1\dots f_n}^{c_1\dots c_n; s_1\dots s_n}(p_1, \dots, p_n) \\ = (\mu^\epsilon g_s)^m \mathbf{J}_{f_1\dots f_m}^{c_1\dots c_m; s_1\dots s_m}(p_1, \dots, p_m) \mathcal{M}_{f_{m+1}\dots f_n}^{c_{m+1}\dots c_n; s_{m+1}\dots s_n}(p_{m+1}, \dots, p_n), \end{aligned} \quad (2.1)$$

where  $g_s$  is the strong coupling constant and  $\mu$  is the scale introduced by dimensional regularisation. Throughout this paper we work in Conventional Dimensional Regularisation (CDR) in  $D = 4 - 2\epsilon$  dimensions. In particular, we assume that gluons have  $D - 2$  physical polarisations (quarks always have 2 polarisations). The symbol  $\mathcal{S}_{1\dots m}$  in eq. (2.1) denotes the operation of keeping only the leading divergent term in the soft limit. The scattering amplitude  $\mathcal{M}_{f_{m+1}\dots f_n}^{c_{m+1}\dots c_n; s_{m+1}\dots s_n}$  on the right-hand side is obtained from the amplitude on the left-hand side by simply removing the soft particles. The current  $\mathbf{J}_{f_1\dots f_m}^{c_1\dots c_m; s_1\dots s_m}$  describes the leading divergent behaviour of the amplitude in the soft limit, often referred to as the *eikonal* approximation in the literature.

The soft current depends on the colour, spin and flavour quantum numbers of the soft partons. In order to keep our notations compact, we find it useful to work with the colour-space formalism of refs. [40, 41], where scattering amplitudes are interpreted as vectors that can be expanded into an orthonormal basis in colour  $\otimes$  spin space,

$$|\mathcal{M}_{f_1\dots f_n}(p_1, \dots, p_n)\rangle = |c_1 \dots c_n\rangle \otimes |s_1 \dots s_n\rangle \mathcal{M}_{f_1\dots f_n}^{c_1\dots c_n; s_1\dots s_n}(p_1, \dots, p_n). \quad (2.2)$$

We will often suppress the dependence on the momenta. With this notation, the squared matrix element summed over spin and colour indices of the external particles can be written as

$$|\mathcal{M}_{f_1\dots f_n}|^2 \equiv \langle \mathcal{M}_{f_1\dots f_n} | \mathcal{M}_{f_1\dots f_n} \rangle = \sum_{\substack{(s_1, \dots, s_n) \\ (c_1, \dots, c_n)}} \left[ \mathcal{M}_{f_1\dots f_n}^{c_1\dots c_n; s_1\dots s_n} \right]^\dagger \mathcal{M}_{f_1\dots f_n}^{c_1\dots c_n; s_1\dots s_n}. \quad (2.3)$$

The soft current can then be interpreted as an operator  $\mathbf{J}_{f_1\dots f_m}(p_1, \dots, p_m)$  in this colour space, and the soft factorisation in eq. (2.1) takes the form,

$$\mathcal{S}_{1\dots m} |\mathcal{M}_{f_1\dots f_n}\rangle = (\mu^\epsilon g_s)^m \mathbf{J}_{f_1\dots f_m} |\mathcal{M}_{f_{m+1}\dots f_n}\rangle. \quad (2.4)$$

This operator acts on color space via the infinitesimal generators of the gauge transformations,

$$\mathbf{T}_i^a |c_1 \dots c_n\rangle = |c_1 \dots c'_i \dots c_n\rangle \mathbf{T}_{c'_i c_i}^a, \quad (2.5)$$

where we defined  $\mathbf{T}_{c'_i c_i}^a = t_{c'_i c_i}^a$  if parton  $i$  is a quark,  $\mathbf{T}_{c'_i c_i}^a = -t_{c_i c'_i}^a$  if it is an anti-quark (here  $t_{c'_i c_i}^a$  are the generators of the fundamental representation of  $SU(N_c)$ ), and  $\mathbf{T}_{c'_i c_i}^a = if^{c'_i a c_i}$  for

a gluon. Since colour must be conserved in every scattering process, the vector  $|\mathcal{M}_{f_{m+1}, \dots, f_n}\rangle$  must be a colour singlet, i.e., it must satisfy

$$\sum_{i=m+1}^n \mathbf{T}_i^a |\mathcal{M}_{f_{m+1}, \dots, f_n}\rangle = 0. \quad (2.6)$$

Henceforth we will simply use the shorthand,

$$\sum_{i=m+1}^n \mathbf{T}_i^a = 0, \quad (2.7)$$

where it is understood that this identity is only valid when we act on colour singlet states.

The soft current for the emission of a single soft gluon is known through two loops in the strong coupling constant [8, 10, 11, 34–36, 42, 43]. The double-soft current is known at tree-level and one-loop for the emission of a pair of soft gluons or quarks [1, 3, 6, 37, 38]. Triple-soft emission is currently only known at tree-level for three gluons [39].<sup>3</sup> The main aim of this paper is to compute for the first time the triple soft current for the emission of a soft quark pair in addition to a gluon. In the remainder of this section, we review the known results for the tree-level soft currents for the emission of a single soft gluon or quark pair, both in order to illustrate some general properties of soft currents and to define some quantities that will be useful when computing the soft current  $\mathbf{J}_{\bar{q}qg}$  in section 3.

## 2.1 The tree-level soft current for the emission of a single soft gluon

Let us start by discussing the case of the emission of a single soft gluon at tree-level. It will often be useful to consider a variant of the soft current operator where we have amputated the polarisation states, e.g., in the case of a single soft gluon, we define

$$\mathbf{J}_g(p_1) = |a\rangle \otimes |s\rangle \mathbf{J}_g^{a;s}(p_1) = |a\rangle \otimes |s\rangle \epsilon_\mu^s(p_1, n) \mathbf{J}_g^{a;\mu}(p_1). \quad (2.8)$$

Note that throughout this paper we always work in axial gauge for external gluons, where the polarisation vectors satisfy the constraints,

$$p_1^\mu \epsilon_\mu^s(p_1, n) = n^\mu \epsilon_\mu^s(p_1, n) = 0, \quad (2.9)$$

where  $n$  is a lightlike reference vector with  $p_1 \cdot n \neq 0$ . The soft current for the tree-level emission of a single gluon is given by [42, 43]

$$\mathbf{J}_g^{a;\mu}(p_1) = \sum_{i=2}^n \mathbf{J}_i^{a;\mu}(p_1) = \sum_{i=2}^n \mathcal{S}_i^\mu(p_1) \mathbf{T}_i^a, \quad (2.10)$$

with

$$\mathcal{S}_i^\mu(p_1) = \frac{p_i^\mu}{p_i \cdot p_1}. \quad (2.11)$$

---

<sup>3</sup>Quadruple-soft emission is known at tree-level for four gluons in the special case of the emission from two hard partons [39].

Note that, as a consequence of gauge invariance, the soft current is conserved,

$$p_{1\mu} \mathbf{J}_g^{a;\mu}(p_1) = \sum_{i=2}^n \mathbf{T}_i^a = 0 \quad \text{mod} \quad \sum_{i=2}^n \mathbf{T}_i^a = 0. \quad (2.12)$$

A similar relation also holds for multi-soft gluon emission, albeit, due to the non-abelian nature of the gauge interactions, only upon replacing *one* polarisation vector by its momentum. In ref. [39] a stronger version was shown to hold even if the remaining external polarisation vectors are amputated, although one needs to project onto the subspace spanned by the physical polarisations. Since our main interest is the current for the emission of a single soft gluon in addition to a soft quark pair, we will not pursue this further.

One is usually interested in the effect of the soft current and the color-correlations it induces after squaring the matrix element,

$$\begin{aligned} \mathcal{S}_1 |\mathcal{M}_{gf_2\dots f_n}|^2 &= (\mu^\epsilon g_s)^2 \langle \mathcal{M}_{f_2\dots f_n} | | \mathbf{J}_g(p_1) |^2 | \mathcal{M}_{f_2\dots f_n} \rangle \\ &= -(\mu^\epsilon g_s)^2 \sum_{i,j=2}^n \mathcal{S}_{ij}(p_1) |\mathcal{M}_{f_2\dots f_n}^{(ij)}|^2, \end{aligned} \quad (2.13)$$

where we have introduced the colour-correlated matrix element,

$$|\mathcal{M}_{f_2\dots f_n}^{(ij)}|^2 = \langle \mathcal{M}_{f_2\dots f_n} | \mathbf{T}_i \cdot \mathbf{T}_j | \mathcal{M}_{f_2\dots f_n} \rangle, \quad (2.14)$$

and the well-known eikonal function,

$$\mathcal{S}_{ij}(p_1) = \frac{2\chi_{ij}}{\chi_{1i}\chi_{1j}} \quad (2.15)$$

with  $\chi_{ij} = 2p_i \cdot p_j$  and  $\mathbf{T}_i \cdot \mathbf{T}_j = \mathbf{T}_j \cdot \mathbf{T}_i = \mathbf{T}_i^a \mathbf{T}_j^a$ . At this point we make some important comments. First, if we simply evaluate the square of the soft current in axial gauge, we find

$$\begin{aligned} |\mathbf{J}_g(p_1)|^2 &= [\mathbf{J}_g^{a;\mu}(p_1)]^\dagger d_{\mu\nu}(p_1, n) \mathbf{J}_g^{a;\nu}(p_1) \\ &= - \sum_{i,j=2}^n \mathbf{T}_i \cdot \mathbf{T}_j \mathcal{S}_{ij}(p_1) + \sum_{i,j=2}^n \mathbf{T}_i \cdot \mathbf{T}_j \left( \frac{n \cdot p_i}{(p_i \cdot p_1)(n \cdot p_1)} + \frac{n \cdot p_j}{(p_j \cdot p_1)(n \cdot p_1)} \right), \end{aligned} \quad (2.16)$$

where  $d_{\mu\nu}(p_1, n)$  denotes the sum over physical polarisations in axial gauge,

$$d_{\mu\nu}(p_1, n) = \sum_{s=1}^{D-2} \epsilon_\mu^s(p_1, n) \epsilon_\nu^s(p_1, n) = -g_{\mu\nu} + \frac{p_{1\mu} n_\nu + n_\mu p_{1\nu}}{p_1 \cdot n}. \quad (2.17)$$

We see that the terms dependent on  $n$  cancel due to colour conservation when acting on the amplitude in eq. (2.13). We will from now on drop terms in the squared current that vanish due to colour conservation in the hard amplitude. Second, we note that, as a consequence of  $\mathbf{T}_i \cdot \mathbf{T}_j = \mathbf{T}_j \cdot \mathbf{T}_i$ , the colour-correlated squared amplitude,  $|\mathcal{M}_{f_2\dots f_n}^{(ij)}|^2$ , features a symmetry under exchange of the emitters ( $i \leftrightarrow j$ ), to which we will refer as the *dipole symmetry*. The dipole symmetry of  $|\mathcal{M}_{f_2\dots f_n}^{(ij)}|^2$  implies that the kinematic factor (the eikonal function  $\mathcal{S}_{ij}(p_1)$ ) in eq. (2.13) is symmetric as well under the exchange of the emitters, ( $i \leftrightarrow j$ ). This is a general feature: the kinematic functions in the squared soft current inherit the symmetry properties of the colour operators that they multiply.

## 2.2 The tree-level soft current for the emission of a single quark pair

Let us also briefly review the features of the soft current for a single soft quark pair emission. It is again convenient to amputate the polarisation states,

$$\begin{aligned} \mathbf{J}_{\bar{q}q}(p_1, p_2) &= |\bar{i}j\rangle \otimes |s_1 s_2\rangle \mathbf{J}_{\bar{q}q}^{\bar{i}j; s_1 s_2}(p_1, p_2) \\ &= |\bar{i}j\rangle \otimes |s_1 s_2\rangle \bar{u}_{s_2}(p_2) \mathbf{J}_{\bar{q}q}^{\bar{i}j}(p_1, p_2) v_{s_1}(p_1), \end{aligned} \quad (2.18)$$

where  $\bar{i}$  and  $j$  denote the anti-fundamental and fundamental colour indices of the anti-quark and quark respectively. Note that  $\mathbf{J}_{\bar{q}q}^{\bar{i}j}$  is a matrix in Dirac spinor space, though we suppress the matrix indices for brevity. It is easy to see that the current for the emission of a soft quark pair is entirely determined by the soft current for single-gluon emission,

$$\mathbf{J}_{\bar{q}q}^{\bar{i}j}(p_1, p_2) = -\frac{1}{\chi_{12}} t_{j\bar{i}}^a \gamma_\mu \mathbf{J}_g^{a;\mu}(p_{12}), \quad p_{12} = p_1 + p_2, \quad (2.19)$$

where we work with a gluon propagator in Feynman gauge. We can square the soft current to obtain

$$\mathcal{S}_{12} |\mathcal{M}_{\bar{q}_1 q_2 f_3 \dots f_n}|^2 = (\mu^{2\epsilon} g_s^2)^2 T_F \sum_{i,j=3}^n Q_{ij}^{\bar{q}q}(p_1, p_2) |\mathcal{M}_{f_3 \dots f_n}^{(ij)}|^2, \quad (2.20)$$

where  $T_F = \frac{1}{2}$  is defined by  $\text{Tr}(t^a t^b) = T_F \delta^{ab}$ , and we defined the kinematic function,

$$Q_{ij}^{\bar{q}q}(p_1, p_2) = \frac{4}{\chi_{12}^2} \frac{\chi_{1i} \chi_{2j} + \chi_{1j} \chi_{2i} - \chi_{ij} \chi_{12}}{\chi_{i(12)} \chi_{j(12)}}, \quad (2.21)$$


and  $\chi_{i(jk)} = \chi_{ij} + \chi_{ik}$ . Note that, in addition to the dipole symmetry under exchange of the emitters ( $i \leftrightarrow j$ ), the soft  $\bar{q}q$  function features charge conjugation symmetry under the exchange of the quark and antiquark labels,

$$Q_{ij}^{\bar{q}q}(p_1, p_2) = Q_{ij}^{\bar{q}q}(p_2, p_1). \quad (2.22)$$

Finally, we recall that the eikonal currents, and so the kinematic functions of a single soft-gluon (2.15) or of a soft  $\bar{q}q$  pair (2.21), do not depend on the mass of the emitters, while the eikonal current of two soft gluons receives an extra contribution which is proportional to the squared mass of the emitters [6].

## 3 Tree-level factorisation for soft $\bar{q}qg$ emission

In this section we present the main result of this paper, namely our result for the tree-level soft current for soft  $\bar{q}qg$  emission. We start by quoting the results for the soft current, and then we discuss the analytic expressions for the squared current and the ensuing colour correlations.



**Figure 1.** Soft-gluon insertion diagrams for the emission of a soft gluon and  $q\bar{q}$  pair.

### 3.1 The soft current

Without loss of generality, we assume the anti-quark, quark and gluon to be partons 1, 2 and 3, respectively. The soft current is obtained by evaluating the diagrams in figure 1. In figure 1 the lines emanating from the hard amplitude (represented by the blue blob) represent Wilson lines in the representation of the hard partons. Soft gluons couple to Wilson lines using the well-known eikonal Feynman rules, cf., e.g., ref. [3].

We use axial gauge for the external gluon, but we evaluate all internal gluon propagators in Feynman gauge. We rescale the momenta of the three soft partons by  $\lambda$ ,  $p_i \rightarrow \lambda p_i$ ,  $i = 1, 2, 3$ , and we perform a Laurent expansion around  $\lambda = 0$ . The leading term in this Laurent expansion defines the soft current. We find it again convenient to amputate the polarisation vectors,

$$\mathbf{J}_{\bar{q}qg}^{\bar{i}j a; s_1 s_2 s_3}(p_1, p_2, p_3) = \epsilon_\mu^{s_3}(p_3, n) \bar{u}_{s_2}(p_2) \mathbf{J}_{\bar{q}qg}^{a; \mu}(p_1, p_2, p_3) v_{s_1}(p_1), \quad (3.1)$$

where for brevity we keep the colour indices of the quark pair implicit on the right-hand side.



We split the tree-level  $\bar{q}qg$  soft current into two contributions,

$$\mathbf{J}_{\bar{q}qg}^{a;\mu}(p_1, p_2, p_3) = \mathbf{K}^{a;\mu}(p_1, p_2, p_3) + \mathbf{W}^{a;\mu}(p_1, p_2, p_3). \quad (3.2)$$

The first term on the right-hand side is given by,

$$\begin{aligned} \mathbf{K}^{a;\mu}(p_1, p_2, p_3) &= \frac{1}{2} \{ \mathbf{J}_g^{a;\mu}(p_3), \mathbf{J}_{\bar{q}q}(p_1, p_2) \}, \\ &= -\frac{1}{2\chi_{12}} \sum_{i,j=4}^n \{ \mathbf{J}_i^{a;\mu}(p_3), \mathbf{J}_j^{b;\nu}(p_{12}) \} \gamma_\nu t^b. \end{aligned} \quad (3.3)$$

where the soft currents appearing on the right-hand side have been defined in the previous section. Equation (3.3) includes both uncorrelated ( $i \neq j$ ) soft emission from two different legs, figure 1IV, as well as contributions to the abelian part of figure 1III. The second term on the right-hand side of eq. (3.2) has the explicit form,

$$\begin{aligned} \mathbf{W}^{a;\mu}(p_1, p_2, p_3) &= \sum_{i=4}^n \mathbf{J}_i^{b;\nu}(p_{123}) \left\{ \frac{1}{\chi_{123}} \left[ \frac{1}{\chi_{13}} \gamma_\nu \not{p}_{13} \gamma^\mu (t^b t^a) - \frac{1}{\chi_{23}} \gamma^\mu \not{p}_{23} \gamma_\nu (t^a t^b) \right] \right. \\ &\quad \left. + \frac{i}{\chi_{12}} f^{abc} \left[ \frac{1}{2} \delta_\nu^\alpha (\mathcal{S}_i^\mu(p_3) - \mathcal{S}_i^\mu(p_{12})) + \frac{1}{\chi_{123}} U_\nu^{\alpha\mu}(p_{12}, p_3) \right] \gamma_\alpha t^c \right\}, \end{aligned} \quad (3.4)$$

where we have defined

$$U_\nu^{\alpha\mu}(p_{12}, p_3) = -2\delta_\nu^\alpha p_{12}^\mu + g^{\alpha\mu}(p_{12} - p_3)_\nu + 2\delta_\nu^\mu p_3^\alpha, \quad (3.5)$$

and

$$\chi_{1\dots k} = \sum_{i<j}^k \chi_{ij}. \quad (3.6)$$

Let us make two comments. First, we have seen that the soft current for the emission of a soft  $\bar{q}q$  pair can be expressed as the soft current for the emission of a single soft gluon contracted with the quark current, cf. eq. (2.19). From the Feynman diagrams in figure 1, it may appear that the  $\bar{q}qg$  soft current can similarly be factorised into the double soft gluon current  $\mathbf{J}_{gg}$  contracted with the quark current. This, however, is not so, because there are certain contributions to  $\mathbf{J}_{gg}$  that can be dropped due to the transversality condition in eq. (2.9), but they have to be kept if the gluon is off shell. Second, we note that, as expected, we find that the divergence of the triple-soft current defined in eq. (3.2) is proportional to the total colour charge of the hard partons, implying that the current is conserved,

$$p_{3,\mu} \mathbf{J}_{\bar{q}qg}^{a;\mu} = -\frac{1}{\chi_{12}} t^c \left[ \mathbf{J}^{c,\mu}(p_{12}) \gamma_\mu \delta^{ab} - i f^{abc} \frac{1}{\chi_{123}} \not{p}_3 \right] \sum_{i=4}^n \mathbf{T}_i^b = 0 \quad \text{mod} \quad \sum_{i=4}^n \mathbf{T}_i^b = 0, \quad (3.7)$$

which is reminiscent of eq. (2.12) in the case of single-gluon emission.

Next, we consider the structure of the squared soft current. We find that we can write the result as a sum over contributions from dipole, tripole and symmetrised colour correlations,

$$\mathcal{S}_{123} |\mathcal{M}_{\bar{q}_1 q_2 g_3 f_4 \dots f_n}|^2 = \left( \mathcal{S}_{123}^{(\text{dip})} + \mathcal{S}_{123}^{(\text{trip})} + \mathcal{S}_{123}^{(\text{sym})} \right) |\mathcal{M}_{\bar{q}_1 q_2 g_3 f_4 \dots f_n}|^2. \quad (3.8)$$

We now discuss the three contributions in turn. The dipole term reads

$$\begin{aligned} \mathcal{S}_{123}^{(\text{dip})} |\mathcal{M}_{\bar{q}_1 q_2 g_3 f_4 \dots f_n}|^2 &= (\mu^{2\epsilon} g_s^2)^3 T_F \sum_{i,j=4}^n \left[ C_F Q_{ij}^{\bar{q}qg(\text{ab})} + C_A \left( Q_{ij}^{\bar{q}qg(\text{nab})} + Q_{ij}^{\bar{q}qg(\text{mass})} \right) \right] |\mathcal{M}_{f_4 \dots f_n}^{(ij)}|^2, \end{aligned} \quad (3.9)$$

where  $C_F$  and  $C_A$  denote the Casimir operators in the fundamental and adjoint representations of  $\text{SU}(N_c)$ ,

$$C_F = \frac{N_c^2 - 1}{2N_c} \quad \text{and} \quad C_A = N_c. \quad (3.10)$$

The kinematic functions will be defined below. We suppress from now on the dependence of the kinematic functions on the momenta for readability. The dipole-correlated matrix element  $|\mathcal{M}_{f_4 \dots f_n}^{(ij)}|^2$  was already defined in eq. (2.14). The *sym* term comes from the symmetrised product of two dipole operators and is given by

$$\mathcal{S}_{123}^{(\text{sym})} |\mathcal{M}_{\bar{q}_1 q_2 g_3 f_4 \dots f_n}|^2 = (\mu^{2\epsilon} g_s^2)^3 T_F \sum_{i,j,k,\ell=4}^n Q_{ik;j\ell}^{\bar{q}qg} |\mathcal{M}_{f_4 \dots f_n}^{(ik;j\ell)}|^2, \quad (3.11)$$

where we introduced the symmetric four-parton correlation function, which also appears in the squared current for double-soft gluon emission at tree-level [3],

$$|\mathcal{M}_{f_4 \dots f_n}^{(ik;j\ell)}|^2 = \langle \mathcal{M}_{f_4 \dots f_n} | \{ \mathbf{T}_i \cdot \mathbf{T}_k, \mathbf{T}_j \cdot \mathbf{T}_l \} | \mathcal{M}_{f_4 \dots f_n} \rangle. \quad (3.12)$$

Besides the two- and four-parton correlations, which have already appeared in the context of single- and double-soft emission at tree-level, we also find a non-vanishing three-parton correlation,

$$\mathcal{S}_{123}^{(\text{trip})} |\mathcal{M}_{\bar{q}_1 q_2 g_3 f_4 \dots f_n}|^2 = (\mu^{2\epsilon} g_s^2)^3 \frac{1}{2} \sum_{i,j,k=4}^n Q_{ijk}^{\bar{q}qg} |\mathcal{M}_{f_4 \dots f_n}^{(ijk)}|^2, \quad (3.13)$$

where we defined the triple correlated tree-level squared amplitude by

$$|\mathcal{M}_{f_4 \dots f_n}^{(ijk)}|^2 = d^{abc} \langle \mathcal{M}_{f_4 \dots f_n} | \mathbf{T}_i^a \mathbf{T}_j^b \mathbf{T}_k^c | \mathcal{M}_{f_4 \dots f_n} \rangle, \quad (3.14)$$

and we have introduced the symmetric structure constant  $d^{abc}$ ,

$$d^{abc} = 2 \text{Tr} \left[ \left\{ t^a, t^b \right\} t^c \right]. \quad (3.15)$$

Colour correlations between three hard partons do not appear in the tree-level soft limits considered in refs. [1, 3, 39, 42, 43], and to the best of our knowledge it is the first time that they occur in tree-level currents. They did occur, however, already in the context of the one-loop soft-current for the emission of a soft  $q\bar{q}$  pair [38]. All other infrared limits, including the infrared poles of three-loop massless amplitudes, are free of  $d^{abc}$  terms. We therefore expect that the infrared singularities generated by the phase space integration of our  $d^{abc}$  term cancel against those with the same colour structure from the soft- $q\bar{q}$  emission at one loop.

Before we discuss the form of the kinematics functions, we outline how the colour correlations introduced above come about, and in particular how it occurs that only the totally symmetric structure constant  $d^{abc}$  enters the tripole correlations, but there is no correlation between three hard lines connected by the antisymmetric structure constant  $f^{abc}$ . In order to do so, we analyse the terms obtained from squaring the soft current in eq. (3.2). The square of the  $\mathbf{K}$  term in eq. (3.3) yields

$$|\mathbf{K}|^2 = -\frac{T_F}{16} \sum_{ijkl} \left[ \left\{ \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\}, \left\{ \mathbf{T}_k^a, \mathbf{T}_l^b \right\} \right\} + (j \leftrightarrow l) \right] \mathcal{S}_{jl}(p_3) Q_{ik}^{q\bar{q}}(p_1, p_2). \quad (3.16)$$

Using the identity [6],

$$\begin{aligned} \left\{ \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\}, \left\{ \mathbf{T}_k^a, \mathbf{T}_l^b \right\} \right\} + (j \leftrightarrow l) &= \left[ 2\{\mathbf{T}_i \cdot \mathbf{T}_k, \mathbf{T}_j \cdot \mathbf{T}_l\} \right. \\ &+ \left. \frac{1}{2} C_A (3\mathbf{T}_i \cdot \mathbf{T}_k (\delta_{il}\delta_{jk} + \delta_{ij}\delta_{kl}) - 4\mathbf{T}_i \cdot \mathbf{T}_j (\delta_{ik}\delta_{il} + \delta_{jk}\delta_{j\ell})) + (i \leftrightarrow k) \right] + (j \leftrightarrow l), \end{aligned} \quad (3.17)$$

eq. (3.16) is straightforwardly reduced to two- and four-parton colour correlations. Next, we examine the crossed term, obtained by contracting  $\mathbf{K}$  with  $\mathbf{W}$ . The product between  $\mathbf{K}$  and the second line of eq. (3.4) yields

$$-T_F \sum_{i,j,l} i f^{abc} \left[ \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\}, \mathbf{T}_l^c \right] \mathcal{Q}_{ijl}^{(1)} = -2T_F C_A \sum_{i,j} \mathbf{T}_i \cdot \mathbf{T}_j (\mathcal{Q}_{iji}^{(1)} - \mathcal{Q}_{ijj}^{(1)}), \quad (3.18)$$

where  $\mathcal{Q}_{ijl}^{(1)}$  is a function of kinematical invariants, and where in order to obtain the equality on the right-hand side, we used the identity [6],

$$i f^{abc} \left[ \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\}, \mathbf{T}_k^c \right] = 2C_A (\mathbf{T}_i \cdot \mathbf{T}_j) (\delta_{ik} - \delta_{jk}). \quad (3.19)$$

We see that eq. (3.18) is reduced to a two-parton colour correlation. The product between  $\mathbf{K}$  and the first line of eq. (3.4) contributes to two- and three-parton correlations. In particular, the product between  $\mathbf{K}$  and the second term in the first line of the  $\mathbf{W}$  term has the form,

$$\begin{aligned} \sum_{i,j,l} \left[ \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\} \mathbf{T}_l^c \text{Tr}(t^b t^a t^c) + \mathbf{T}_l^c \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\} \text{Tr}(t^c t^a t^b) \right] \mathcal{Q}_{ijl}^{(2)} \\ = \sum_{i,j,l} \left( -\frac{i}{2} T_F f^{abc} \left[ \left\{ \mathbf{T}_i^a, \mathbf{T}_j^b \right\}, \mathbf{T}_l^c \right] + d^{abc} \mathbf{T}_i^a \mathbf{T}_j^b \mathbf{T}_l^c \right) \mathcal{Q}_{ijl}^{(2)}, \end{aligned} \quad (3.20)$$

where, like  $\mathcal{Q}^{(1)}$  in eq. (3.18),  $\mathcal{Q}_{ijl}^{(2)}$  is also a function of kinematical invariants, and we used

$$\text{Tr}(t^a t^b t^c) = \frac{1}{4} d^{abc} + \frac{i}{2} T_F f^{abc}. \quad (3.21)$$

The contribution from the first term in the first line of  $\mathbf{W}$  has a similar form. Note, however, that the product between  $\mathbf{K}$  and the first term on the first line of  $\mathbf{W}$  has a relative sign difference with respect to the second term on the first line of  $\mathbf{W}$ . Therefore, the kinematic term of the three-parton correlation (3.24) inherits a relative minus sign

under the exchange of the quark and antiquark labels. Conversely, that sign is cancelled in the  $f^{abc}$  term since  $f^{abc}$  is antisymmetric. Finally, like in eq. (3.18), the  $f^{abc}$  term is reduced to dipole correlations using again the identity (3.19). This explains the absence of three-parton colour correlations involving an antisymmetric structure constant.

Let us now discuss the form of the kinematic functions. First we note that, due to the charge conjugation symmetry under exchange of the quark and anti-quark labels, the functions have a definite symmetry under an exchange of  $p_1$  and  $p_2$ . Further, while the four-parton and three-parton correlations in eqs. (3.11) and (3.13) and the abelian part of the dipole correlations in eq. (3.9) are insensitive to the mass of the emitters, the non-abelian dipole correlation contains additional terms proportional to the mass of emitters, which we have made explicit in eq. (3.9). This is in line with the dependence of the triple-soft gluon current on the mass [39].

The four-parton correlated term in eq. (3.11) only receives contributions from squaring the function  $\mathbf{K}^{a,\mu}$  in eq. (3.3). Therefore, the kinematic function associated to the four-parton correlations separates into the product of two eikonal functions, one associated to the soft gluon and the other to the soft  $\bar{q}q$  pair,

$$Q_{ik;j\ell}^{\bar{q}qg}(p_1, p_2, p_3) = -\frac{1}{2} \mathcal{S}_{j\ell}(p_3) Q_{ik}^{\bar{q}q}(p_1, p_2), \quad (3.22)$$

where the single-soft eikonal function and the soft- $\bar{q}q$  function are given in eqs. (2.15) and (2.21). From the symmetry of the colour operator in eq. (3.12), it follows that the kinematic factor in eq. (3.22) features dipole symmetries under the exchanges ( $i \leftrightarrow k$ ) or ( $j \leftrightarrow \ell$ ).

Let us now turn to the kinematic functions appearing in the dipole correlations. They must possess the dipole symmetry ( $i \leftrightarrow j$ ). The coefficient of  $C_F$  in eq. (3.9) is given by

$$\begin{aligned} Q_{ij}^{\bar{q}qg(ab)} = & \frac{8}{\chi_{123}^2 \chi_{i(123)} \chi_{j(123)}} \left\{ \frac{D}{2} \left[ \frac{\chi_{1j} \chi_{3i}}{\chi_{13}} - \frac{\chi_{12} \chi_{3i} \chi_{3j}}{\chi_{13} \chi_{23}} - \chi_{ij} \left( \frac{\chi_{13}}{\chi_{23}} + 1 \right) \right. \right. \\ & \left. \left. + \frac{\chi_{1i} \chi_{3j} + \chi_{1j} \chi_{3i} + \chi_{2i} \chi_{3j}}{\chi_{23}} \right] - \frac{\chi_{12} (\chi_{12} \chi_{ij} - \chi_{1i} \chi_{2j} - \chi_{1j} \chi_{2i})}{\chi_{13} \chi_{23}} \right. \\ & \left. + \frac{\chi_{12} (\chi_{3j} \chi_{1i} + \chi_{2j} \chi_{3i} + 2 \chi_{3j} \chi_{3i})}{\chi_{13} \chi_{23}} + \chi_{ij} \left( \frac{\chi_{13} - \chi_{12}}{\chi_{23}} - \frac{\chi_{12}}{\chi_{13}} + 2 \right) \right. \\ & \left. - \frac{\chi_{1j} (\chi_{1i} - \chi_{2i} + 2 \chi_{3i})}{\chi_{13}} - \frac{\chi_{3j} \chi_{i(12)} + \chi_{1j} (\chi_{3i} - \chi_{2i}) + \chi_{2i} \chi_{j(23)}}{\chi_{23}} \right\} \\ & + (1 \leftrightarrow 2), \end{aligned} \quad (3.23)$$

with  $\chi_{i(123)} = \chi_{1i} + \chi_{2i} + \chi_{3i}$ , which is manifestly symmetric under ( $i \leftrightarrow j$ ) once the ( $1 \leftrightarrow 2$ ) exchange has been added. The coefficients of  $C_A$  in eq. (3.9) are more lengthy, and we provide them in appendix A.

Finally, the kinematic factor appearing in the three-parton correlations can be written in an explicitly symmetric form as follows,

$$\begin{aligned}
 Q_{ijk}^{\bar{q}qg}(p_1, p_2, p_3) = & \frac{2}{3\chi_{123}} \left\{ \chi_{ij} \left[ \frac{u_{jk}}{\chi_{k(123)}\chi_{i(12)}} + \frac{1}{\chi_{j(123)}} \left( \frac{r_k}{\chi_{i(12)}} + \frac{t_{ik}}{\chi_{k(12)}} \right) \right] + \frac{u_{ij;k}}{\chi_{k(123)}} \right. \\
 & - \frac{2}{\chi_{12}\chi_{23}\chi_{k(123)}} \frac{\chi_{2j}(\chi_{1i}\chi_{2k} + \chi_{1k}\chi_{2i})}{\chi_{3j}\chi_{i(12)}} + \frac{2\chi_{ij}\chi_{2k}}{\chi_{23}\chi_{3k}} \frac{1}{\chi_{i(12)}\chi_{j(123)}} \\
 & \left. + 5 \text{ permutations of } (ijk) \right\} - (1 \leftrightarrow 2), \tag{3.24}
 \end{aligned}$$

with

$$\begin{aligned}
 r_k &= \frac{\chi_{13}\chi_{2k}}{\chi_{12}\chi_{23}\chi_{3k}} - \frac{\chi_{1k}}{\chi_{12}\chi_{3k}} + \frac{1}{\chi_{23}}, \\
 t_{ik} &= \frac{\chi_{13}\chi_{2k}}{\chi_{12}\chi_{23}\chi_{3i}} + \frac{\chi_{1k}}{\chi_{12}\chi_{3i}} - \frac{\chi_{3k}}{\chi_{23}\chi_{3i}}, \\
 u_{jk} &= -\frac{\chi_{13}\chi_{2k}}{\chi_{12}\chi_{23}\chi_{3j}} + \frac{\chi_{1k}}{\chi_{12}\chi_{3j}} + \frac{\chi_{3k}}{\chi_{23}\chi_{3j}}, \\
 u_{ij;k} &= \frac{-\chi_{1i}(\chi_{2j}\chi_{3k} + \chi_{2k}\chi_{3j}) + \chi_{1j}(\chi_{2k}\chi_{3i} - \chi_{2i}\chi_{3k}) - \chi_{1k}(\chi_{2i}\chi_{3j} + \chi_{2j}\chi_{3i})}{\chi_{12}\chi_{23}\chi_{3j}\chi_{i(12)}}. \tag{3.25}
 \end{aligned}$$

### 3.2 Strongly-ordered soft limits

In this section we examine the strongly-ordered soft limits, where one or more partons are much softer than the others. The strongly-ordered limits can be computed by rescaling the softer momenta by a parameter  $\lambda$  and keeping only the leading term in the Laurent expansion around  $\lambda = 0$ . Since the rescaling of the momentum components is uniform in  $\lambda$ , the ordering in the momentum components is equivalent to an ordering of the energies  $E_i$  of the soft particles. We will use the latter to label strongly-ordered limits. We note that the strongly-ordered soft limits where the quark is much softer than the anti-quark (or vice versa) is subleading, and we will not discuss those cases here. Hence, we discuss in the following only the two strongly-ordered soft limits where the energies of the quark and the anti-quark are of the same order.

The factorisation of the squared matrix element in any of the strongly-ordered limits can be written as in eq. (3.8), after replacing the kinematic factors with their expressions in the limit. We now discuss the two relevant strongly-ordered limits in turn.

Let us start with the case where the gluon is softer than the  $\bar{q}q$  pair,  $E_3 \ll E_1, E_2$ . The kinematic functions behave as

$$\begin{aligned}
 \mathcal{S}_3 Q_{ik;jl}^{\bar{q}qg}(p_1, p_2, p_3) &= Q_{ik;jl}^{\bar{q}qg}(p_1, p_2, p_3), \\
 \mathcal{S}_3 Q_{ijk}^{\bar{q}qg}(p_1, p_2, p_3) &= \frac{1}{6} Q_{ik}^{\bar{q}q}(p_1, p_2) [\mathcal{S}_{1j}(p_3) - \mathcal{S}_{2j}(p_3)] + 5 \text{ permutations of } (ijk), \\
 \mathcal{S}_3 Q_{ij}^{\bar{q}qg(\text{ab})}(p_1, p_2, p_3) &= 2 \mathcal{S}_{12}(p_3) Q_{ij}^{\bar{q}q}(p_1, p_2), \\
 \mathcal{S}_3 Q_{ij}^{\bar{q}qg(\text{nab})}(p_1, p_2, p_3) &= \left\{ \frac{1}{4} Q_{ij}^{\bar{q}q}(p_1, p_2) [\mathcal{S}_{1i}(p_3) - \mathcal{S}_{ij}(p_3) - \mathcal{S}_{12}(p_3)] \right. \\
 &\quad + \frac{\chi_{12}}{8} \mathcal{S}_{2j}(p_3) [\mathcal{S}_{1i}(p_{12}) \mathcal{S}_{2j}(p_{12}) - \mathcal{S}_{2i}(p_{12}) \mathcal{S}_{1j}(p_{12})] [\mathcal{S}_{1i}(p_{12}) - \mathcal{S}_{2i}(p_{12})] \\
 &\quad \left. - \frac{1}{2\chi_{12}} \mathcal{S}_{ij}(p_{12}) \mathcal{S}_{2i}(p_3) + \frac{1}{2} \mathcal{S}_{ij}(p_3) \mathcal{S}_{1i}(p_{12}) \mathcal{S}_{2i}(p_{12}) + (i \leftrightarrow j) \right\} + (1 \leftrightarrow 2). \\
 \mathcal{S}_3 Q_{ij}^{q\bar{q}g(\text{mass})}(p_1, p_2, p_3) &= m_i^2 \left( \frac{4\chi_{1j}}{\chi_{12}\chi_{13}\chi_{i(12)}^2\chi_{3j}} - \frac{4\chi_{ij}}{\chi_{12}\chi_{3i}\chi_{i(12)}^2\chi_{3j}} + (1 \leftrightarrow 2) \right) + (i \leftrightarrow j).
 \end{aligned} \tag{3.26}$$

We note that the function  $Q_{ik;jl}^{\bar{q}qg}$  in eq. (3.22) is exact in this strongly-ordered limit. In the case where the  $\bar{q}q$  pair is softer than the gluon,  $E_1, E_2 \ll E_3$ , the kinematic factors behave as

$$\begin{aligned}
 \mathcal{S}_{12} Q_{ik;jl}^{\bar{q}qg}(p_1, p_2, p_3) &= Q_{ik;jl}^{\bar{q}qg}(p_1, p_2, p_3), \\
 \mathcal{S}_{12} Q_{ijk}^{\bar{q}qg}(p_1, p_2, p_3) &= \mathcal{S}_{12} Q_{ij}^{\bar{q}qg(\text{ab})}(p_1, p_2, p_3) = 0, \\
 \mathcal{S}_{12} Q_{ij}^{\bar{q}qg(\text{nab})}(p_1, p_2, p_3) &= \left\{ \frac{\chi_{13}}{\chi_{12}\chi_{3(12)}} \mathcal{S}_{1i}(p_3) \left[ \frac{2\chi_{23}}{\chi_{12}\chi_{3(12)}} - \frac{1}{2} \mathcal{S}_{2i}(p_{12}) \right] \right. \\
 &\quad + \mathcal{S}_{ij}(p_3) \left[ \frac{\chi_{13}}{2\chi_{3(12)}} \mathcal{S}_{2i}(p_{12})^2 + \frac{\chi_{23}}{\chi_{12}\chi_{3(12)}} \mathcal{S}_{1i}(p_{12}) - \frac{2\chi_{13}\chi_{23}}{\chi_{12}^2\chi_{3(12)}^2} \right. \\
 &\quad \left. \left. + \frac{1}{4} (\mathcal{S}_{1i}(p_{12})\mathcal{S}_{2i}(p_{12}) - Q_{ij}^{\bar{q}q}(p_1, p_2)) \right] \right. \\
 &\quad \left. - \mathcal{S}_{ij}(p_{12}) \left[ \frac{\chi_{12}}{8\chi_{3(12)}} \mathcal{S}_{1j}(p_{12}) \mathcal{S}_{2i}(p_{12}) (\chi_{13} \mathcal{S}_{1j}(p_3) - \chi_{23} \mathcal{S}_{i2}(p_3)) \right. \right. \\
 &\quad \left. \left. + \frac{1}{2} \mathcal{S}_{i(12)}(p_3) \left( \mathcal{S}_{1j}(p_{12}) + \frac{1}{\chi_{12}} \right) \right] + (i \leftrightarrow j) \right\} + (1 \leftrightarrow 2), \\
 \mathcal{S}_{12} Q_{ij}^{q\bar{q}g(\text{mass})}(p_1, p_2, p_3) &= m_i^2 \left( \frac{4\chi_{3j}}{\chi_{12}\chi_{3(12)}\chi_{3i}^2\chi_{j(12)}} - \frac{8\chi_{23}\chi_{1j}}{\chi_{12}^2\chi_{3(12)}\chi_{3i}^2\chi_{j(12)}} \right. \\
 &\quad - \frac{4\chi_{ij}}{\chi_{12}\chi_{3i}^2\chi_{i(12)}\chi_{j(12)}} + \frac{8\chi_{1i}\chi_{2j}}{\chi_{12}^2\chi_{3i}^2\chi_{i(12)}\chi_{j(12)}} \\
 &\quad \left. + \frac{2}{\chi_{12}\chi_{3(12)}\chi_{3i}\chi_{i(12)}} + (1 \leftrightarrow 2) \right) + (i \leftrightarrow j),
 \end{aligned} \tag{3.27}$$

where we defined  $\mathcal{S}_{k(ij)}(p_l) = \mathcal{S}_{ik}(p_l) + \mathcal{S}_{jk}(p_l)$ . The function  $Q_{ik;jl}^{\bar{q}qg}$  in eq. (3.22) is again exact in this limit. The tripole contribution is power-suppressed and can therefore be neglected. The same is true for the coefficient of  $C_F$  in the dipole contribution.

Supplementary material for the terms in eqs. (3.8), (3.26) and (3.27) is provided in a computer-readable format with this published work. The content and conventions are appended as a short note at the start of each file.

## 4 Soft limits of the splitting amplitudes

So far we have only considered soft singularities of on-shell tree-level scattering amplitudes. Soft singularities, however, are not the only kinematic limits in which amplitudes become singular and factorise into universal building blocks, but they also exhibit collinear singularities when two or more massless particles become collinear. These limits need to be taken into account when constructing subtraction schemes for higher order computations. In particular, one is also interested in understanding iterated soft and collinear limits. In this section we discuss the behaviour of scattering amplitudes where a subset  $\bar{q}qg$  of collinear massless particles become soft. Before we do this, we give a short review of collinear factorisation in general.

### 4.1 Review of collinear factorisation

Throughout this section we follow closely the notations and conventions of section 2. Consider a tree-level scattering amplitude for  $n$  partons, and assume that a subset of  $m$  massless particles become collinear to a light-like direction  $P$ . The approach to the collinear limit is parametrised by a lightcone decomposition as follows,

$$p_i^\mu = z_i P^\mu + k_{\perp i}^\mu - \frac{k_{\perp i}^2}{2z_i} \frac{n^\mu}{P \cdot n}, \quad i = 1, \dots, m, \quad (4.1)$$

where  $p_i^2 = 0$  and  $P \cdot k_{\perp i} = 0$ . Above,  $n^\mu$  is an auxiliary light-like vector such that  $n \cdot k_{\perp i} = 0$  and  $n \cdot p_i \neq 0 \neq n \cdot P$ . The longitudinal momentum fractions  $z_i$  and the transverse momenta  $k_{\perp i}$  can be chosen to satisfy the following constraints,

$$\sum_{i=1}^m z_i = 1 \quad \text{and} \quad \sum_{i=1}^m k_{\perp i}^\mu = 0. \quad (4.2)$$

The collinear limit is performed by introducing a uniform scaling parameter  $\lambda_c$  as follows,

$$k_{\perp i} \rightarrow \lambda_c k_{\perp i}, \quad 1 \leq i \leq m. \quad (4.3)$$

Then, we expand the matrix element into a Laurent series around  $\lambda_c = 0$  and only keep the leading divergent term. In the collinear limit a scattering amplitude factorises<sup>4</sup> as [45–47],

$$\mathcal{C}_{1\dots m} |\mathcal{M}_{f_1\dots f_n}\rangle = \mathbf{Sp}_{ff_1\dots f_m} |\mathcal{M}_{ff_{m+1}\dots f_n}\rangle, \quad (4.4)$$

where  $f$  denotes the flavour of the parent particle (note that in QCD  $f$  can always be inferred from the flavours  $f_1 \dots f_m$  of the collinear partons). The symbol  $\mathcal{C}_{1\dots m}$  indicates

---

<sup>4</sup>The factorisation in eq. (4.4) is valid to all orders in perturbation theory if all the collinear particles are in the final state. For the case where at least one collinear parton is in the initial state, the factorisation in eq. (4.4) is known to hold only at tree-level [44].

that the equality only holds up to terms that are power-suppressed in the collinear limit. The scattering amplitude on the right-hand side of eq. (4.4) is obtained from the original amplitude by replacing the collinear particles with the light-like momentum  $P$ . The quantity  $\mathbf{Sp}$  is called the *splitting matrix* [30, 44] and only depends on the kinematics and the quantum numbers in the collinear set.

The factorisation in eq. (4.4) implies that also the squared matrix element must factorise,

$$\mathcal{C}_{1\dots m} |\mathcal{M}_{f_1\dots f_n}|^2 = \left( \frac{2\mu^{2\epsilon} g_s^2}{s_{1\dots m}} \right)^{m-1} \langle s' | \hat{\mathbf{P}}_{f_1\dots f_m} | s \rangle \langle s | \mathcal{T}_{ff_{m+1}\dots f_n} | s' \rangle, \quad (4.5)$$

where we defined the Mandelstam invariant  $s_{1\dots m} \equiv (p_1 + \dots + p_m)^2$ . The *helicity matrix*  $\mathcal{T}_{ff_{m+1}\dots f_n}$  denotes the square of the reduced amplitude by not summing over the spin of the parent parton,

$$\langle s | \mathcal{T}_{ff_{m+1}\dots f_n} | s' \rangle \equiv \langle \mathcal{M}_{ff_{m+1}\dots f_n} | s' \rangle \langle s | \mathcal{M}_{ff_{m+1}\dots f_n} \rangle. \quad (4.6)$$

The dependence of the helicity matrix on the lightcone direction  $P$  as well as the momenta of the hard partons is not written explicitly. The (*polarised*) *splitting amplitude*  $\hat{\mathbf{P}}$  is related to  $\mathbf{Sp}$  by

$$\left( \frac{2\mu^{2\epsilon} g_s^2}{s_{1\dots m}} \right)^{m-1} \hat{\mathbf{P}}_{f_1\dots f_m} = \frac{1}{\mathcal{C}_f} [\mathbf{Sp}_{ff_1\dots f_m}]^\dagger \mathbf{Sp}_{ff_1\dots f_m}, \quad (4.7)$$

where  $\mathcal{C}_f$  is the number of colour degrees of freedom of the parent parton with flavour  $f$ , i.e.,  $\mathcal{C}_g = N_c^2 - 1$  for a gluon and  $\mathcal{C}_q = N_c$  for a quark. In writing down eq. (4.7) we implicitly sum over the spin and colour indices of the collinear partons. Due to colour conservation in the hard amplitude, there are no non-trivial colour correlations, i.e., the operator  $\hat{\mathbf{P}}_{f_1\dots f_m}$  acts as the identity on colour space. However, there can be non-trivial spin correlations. When the parent parton is a quark, helicity conservation implies that the splitting amplitude is proportional to unity in spin space,

$$\langle s' | \hat{\mathbf{P}}_{f_1\dots f_m} | s \rangle = \delta^{ss'} \hat{P}_{f_1\dots f_m}, \quad (4.8)$$

where the quantity  $\hat{P}_{f_1\dots f_m}$  is a scalar. In the case where the parent parton is a gluon, it is possible to write the splitting amplitude in terms of Lorentz indices as [3],

$$\langle s | \hat{\mathbf{P}}_{f_1\dots f_m} | s' \rangle = \epsilon_\mu^s(P, n)^* \epsilon_\nu^{s'}(P, n) \hat{P}_{f_1\dots f_m}^{\mu\nu}, \quad (4.9)$$

with

$$\hat{P}_{f_1\dots f_m}^{\mu\nu} = g^{\mu\nu} A_{f_1\dots f_m} + \sum_{i,j=1}^m \frac{k_{\perp i}^\mu k_{\perp j}^\nu}{s_{1\dots m}} B_{ij, f_1\dots f_m}. \quad (4.10)$$

The splitting amplitudes for the squared matrix element have been computed at tree-level for the emission of up to four collinear partons [28, 29]. At one-loop level, they have been computed for the emission of up to three collinear partons [30–32].



So far all considerations were valid in the case where the collinear partons have non-zero energies. However, it can be interesting to understand how splitting amplitudes behave if a subset of collinear partons are soft. In ref. [48] the single- and double-soft limits of triple-collinear tree-level splitting amplitudes were considered. Those soft limits were generalised to a generic set of  $m$  collinear partons [28], where  $m \geq 2$  in the single-soft limit and  $m \geq 3$  in the double-soft limit, and applied to the single- and double-soft limits of quadruple-collinear splitting functions. It was shown in particular that the soft limits of splitting amplitudes involve universal quantities reminiscent of soft currents. In the remainder of this section we extend the analysis of refs. [28, 48] to certain classes of soft limits of more than two collinear partons.

### 4.2 $m$ -parton soft limit of the $m$ -collinear splitting amplitudes

We start by analysing the special case where all  $m$  collinear partons are soft. This implies that the parent parton must itself be soft, and therefore it must then be a gluon. The soft limit of the collinear factorisation in eq. (4.4) then becomes:

$$\mathcal{S}_{1\dots m} \mathcal{C}_{1\dots m} |\mathcal{M}_{f_1, \dots, f_n}\rangle = \mu^\epsilon g_s \mathbf{Sp}_{gf_1 \dots f_m} \mathbf{J}_g(P) |\mathcal{M}_{f_{m+1} \dots f_n}\rangle, \quad (4.11)$$

where the soft-gluon current is given in eq. (2.10). Squaring eq. (4.11) and summing over spin and colour indices, we obtain

$$\begin{aligned} & \mathcal{S}_{1\dots m} \mathcal{C}_{1\dots m} |\mathcal{M}_{f_1 \dots f_n}\rangle^2 \\ &= \left(\mu^{2\epsilon} g_s^2\right)^{m+1} \left(\frac{2}{s_{1\dots m}}\right)^{m-1} \sum_{i,j=m+1}^n \mathcal{S}_{ij\mu\nu}(P) \hat{P}_{f_1 \dots f_m}^{\mu\nu} |\mathcal{M}_{f_{m+1} \dots f_n}^{(ij)}|^2, \end{aligned} \quad (4.12)$$

where  $\hat{P}_{f_1 \dots f_m}^{\mu\nu}$  denotes the polarised splitting amplitude for a gluon to split into  $m$  partons  $f_1, \dots, f_m$  defined in eq. (4.9), and the sum runs over the hard partons. The soft factor  $\mathcal{S}_{ij}^{\mu\nu}$  is given by

$$\mathcal{S}_{ij}^{\mu\nu}(P) = \frac{p_i^\mu p_j^\nu}{(p_i \cdot P)(p_j \cdot P)}. \quad (4.13)$$

Equation (4.12) generalises straightforwardly the analogous derivation for  $m = 2$  collinear partons [6, 48].

### 4.3 Soft $\bar{q}qg$ limit of tree-level splitting amplitudes

In ref. [28], in the case of a tree-level amplitude of  $n$  massless partons, the behaviour of an  $m$ -parton splitting amplitude was analysed where a single gluon, or a  $\bar{q}q$  pair, or two gluons from the collinear set become soft.

We consider now the more general case of a tree amplitude with  $(m+r)$  massless and  $(n-m-r)$  massive partons. Firstly, we note that for an  $m$ -parton splitting amplitude where a single gluon or a  $\bar{q}q$  pair from the collinear set become soft, the kinematic coefficients are the same as the ones of ref. [28], because they are degree-zero homogeneous functions of the  $(n-m)$  momenta of the non-collinear partons. Thus, the kinematic coefficients do not depend on the momenta of the non-collinear partons. Likewise, for an  $m$ -parton

splitting amplitude where two gluons from the collinear set become soft, the massless pieces of the kinematic coefficients are the same as the ones of ref. [28], while the pieces which are proportional to the squared mass of the partons [6] are not singular in the collinear limit. Thus, we conclude that in the case of an  $m$ -parton splitting amplitude where a single gluon, or a  $\bar{q}q$  pair, or two gluons from the collinear set become soft, the factorisation formulae are the same as the ones of ref. [28] even in the more general case of a tree-level amplitude with massive partons.

We now focus on the behaviour of an  $m$ -parton splitting amplitude  $\hat{P}_{\bar{q}qg f_4 \dots f_m}^{ss'}$  in the limit where a  $\bar{q}q$  pair and a gluon from the collinear set become soft. The triple soft limit of the splitting amplitude is defined by performing a rescaling by a small parameter  $\lambda_s$  as follows,

$$z_i \rightarrow \lambda_s z_i, \quad k_{\perp i} \rightarrow \lambda_s k_{\perp i}, \quad \chi_{ij} \rightarrow \lambda_s^2 \chi_{ij}, \quad \chi_{ik} \rightarrow \lambda_s \chi_{ik}, \quad (4.14)$$

$$1 \leq i, j \leq 3, \quad 3 < k \leq m.$$

We expand the ensuing splitting amplitude in  $\lambda_s$ , and keep only the leading pole, of  $\mathcal{O}(\lambda_s^{-6})$ . We will argue that its coefficient is universal.

In order to obtain the factorisation of the splitting amplitude in the triple soft limit from eq. (4.14), we use the fact that the soft limit and the collinear limit are expected to commute, cf. ref. [3]. We then start from eq. (3.8) and take the collinear limit where partons 1 through  $m$  are collinear using colour conservation for the hard amplitude,

$$\sum_{j=m+1}^n \mathbf{T}_j^a = - \sum_{j=4}^m \mathbf{T}_j^a. \quad (4.15)$$

By doing so, we obtain the factorisation formula,

$$\begin{aligned} \mathcal{S}_{123} \left[ \left( \frac{2\mu^{2\epsilon} g_s^2}{s_{1\dots m}} \right)^{m-1} \hat{\mathbf{P}}_{\bar{q}qg f_4 \dots f_m} \right] &= (\mu^{2\epsilon} g_s^2)^3 \frac{T_F}{C_f} \left[ \sum_{i,j,k,\ell=4}^m U_{ijk\ell;123}^{\bar{q}qg} |\mathbf{Sp}_{ff_4 \dots f_m}^{(ik;j\ell)}|^2 \right. \\ &+ \sum_{i,j,k=4}^m U_{ijk;123}^{\bar{q}qg} |\mathbf{Sp}_{ff_4 \dots f_m}^{(ijk)}|^2 + \sum_{i,j=4}^m \left( C_F U_{ij;123}^{\bar{q}qg(ab)} + C_A U_{ij;123}^{\bar{q}qg(nab)} \right) |\mathbf{Sp}_{ff_4 \dots f_m}^{(ij)}|^2 \left. \right], \end{aligned} \quad (4.16)$$

with the factor  $C_f$  defined in eq. (4.7), and we introduced the two-, three- and four-parton colour-correlated splitting amplitudes,

$$\begin{aligned} |\mathbf{Sp}_{ff_4 \dots f_m}^{(ij)}|^2 &\equiv \left[ \mathbf{Sp}_{ff_4 \dots f_m} \right]^\dagger \mathbf{T}_i \cdot \mathbf{T}_j \mathbf{Sp}_{ff_4 \dots f_m}, \\ |\mathbf{Sp}_{ff_4 \dots f_m}^{(ijk)}|^2 &\equiv d^{abc} \left[ \mathbf{Sp}_{ff_4 \dots f_m} \right]^\dagger \mathbf{T}_i^a \mathbf{T}_j^b \mathbf{T}_k^c \mathbf{Sp}_{ff_4 \dots f_m}, \\ |\mathbf{Sp}_{ff_4 \dots f_m}^{(ik;j\ell)}|^2 &\equiv \left[ \mathbf{Sp}_{ff_4 \dots f_m} \right]^\dagger \{ \mathbf{T}_i \cdot \mathbf{T}_k, \mathbf{T}_j \cdot \mathbf{T}_\ell \} \mathbf{Sp}_{ff_4 \dots f_m}. \end{aligned} \quad (4.17)$$

The factorisation in eq. (4.16) follows straightforwardly from applying the collinear limit and imposing colour conservation as in eq. (4.15). A detailed derivation in the case of the single and double soft limits of splitting amplitudes can be found in ref. [28], where the two- and four-parton colour-correlated splitting amplitudes were introduced. The coefficients of colour-correlated splitting amplitudes are obtained by taking the collinear limit of the soft functions in eqs. (3.9), (3.11) and (3.13).

Applying colour conservation in the hard amplitude, eq. (4.15), we see that the hard matrix element completely factorises from the collinear limit of eqs. (3.9), (3.11) and (3.13). This is due to colour coherence: a cluster of collinear partons acts coherently as one single coloured object, i.e., the hard partons cannot resolve the individual collinear partons, and the colour correlations are in the space of the collinear partons only. The procedure to compute the contributions of the two- and four-parton colour correlations has been presented in detail in ref. [28] in the context of single- and double-soft limits of splitting amplitudes. The computation of the three-parton colour correlations is similar, and so here we content ourselves to simply state the results.

- The coefficient multiplying  $C_F$  in the two-parton colour-correlated term in eq. (4.16) is given by

$$U_{ij;123}^{\bar{q}qg(ab)} = Q_{ij}^{\bar{q}qg(ab)} + u_{(\text{dip})}^{(ab)}, \quad (4.18)$$

where  $Q_{ij}^{\bar{q}qg(ab)}$  is given in eq. (3.23) and  $u_{(\text{dip})}^{(ab)}$  collects contributions with at least one spectator outside of the collinear set,

$$\begin{aligned} u_{(\text{dip})}^{(ab)} = & \frac{8}{\chi_{123}^2 z_{123}^2} \left[ \frac{z_2(z_1 - z_2) - 2z_3 z_{12}}{\chi_{23}} + \frac{z_1 z_2 \chi_{12}}{\chi_{13} \chi_{23}} \left( \frac{z_3(z_3 + z_{123})}{z_1 z_2} + 2 \right) \right. \\ & \left. - \frac{z_1(z_1 - z_2 + 2z_3)}{\chi_{13}} + \frac{1}{2} D_{z_3} \frac{2z_2 \chi_{3(12)} - \chi_{12} z_3}{\chi_{13} \chi_{23}} \right] \\ & - \left\{ \frac{8}{\chi_{123}^2 z_{123} s_{i(123)}} \left[ z_i \left( \frac{\chi_{13}}{\chi_{23}} - \frac{\chi_{12} \chi_{123}}{\chi_{13} \chi_{23}} + 2 \right) + \frac{z_1(s_{2i} - s_{1i} - 2s_{3i})}{\chi_{13}} \right. \right. \\ & + \frac{D}{2} \frac{\chi_{13}(s_{i(12)} z_3 - \chi_{3(12)} z_i) + s_{3i}(\chi_{3(12)} z_1 - \chi_{12} z_3)}{\chi_{13} \chi_{23}} \\ & \left. \left. - \frac{s_{2i}(z_{23} + z_3 - z_1) + s_{3i} z_1 + s_{1i} z_3}{\chi_{23}} \right. \right. \\ & \left. \left. + \frac{\chi_{12}(z_3 s_{i(13)} + z_{23} s_{3i} + 2z_2 s_{1i})}{\chi_{13} \chi_{23}} \right] + (i \leftrightarrow j) \right\} \\ & + (1 \leftrightarrow 2), \end{aligned} \quad (4.19)$$

where we defined  $z_{1\dots k} = z_1 + \dots + z_k$ . The function multiplying  $C_A$  in the two-parton colour-correlated term in eq. (4.16) is

$$U_{ij;123}^{\bar{q}qg(\text{nab})} = Q_{ij}^{\bar{q}qg(\text{nab})} + u_{(\text{dip})}^{(\text{nab})}, \quad (4.20)$$

where  $Q_{ij}^{\bar{q}qg(\text{nab})}$  is given in eq. (A.1), while  $u_{(\text{dip})}^{(\text{nab})}$  is provided in appendix B. Note that there is no mass term in eq. (4.16), since the mass-dependent term  $Q_{ij}^{\bar{q}qg(\text{mass})}$  in eq. (A.2) is not singular in the collinear limit.

- The function multiplying the three-parton colour correlations in eq. (4.16) reads

$$U_{ijk;123}^{\bar{q}qg} = Q_{ijk}^{\bar{q}qg} + u_{(\text{trip})}, \quad (4.21)$$

where  $Q_{ijk}^{\bar{q}qg}$  is given in eq. (3.24), while  $u_{(\text{trip})}$  is

$$\begin{aligned}
 u_{(\text{trip})} = & \left\{ \frac{8z_1z_2z_{23}}{3\chi_{123}\chi_{12}\chi_{23}z_3z_{12}z_{123}} + \frac{2}{3\chi_{12}\chi_{123}\chi_{23}} \left\{ \frac{2z_2(\chi_{1(23)}z_i - \chi_{1i}z_{23} - z_1\chi_{i(23)})}{z_3z_{12}\chi_{i(123)}} \right. \right. \\
 & + \frac{2}{z_{123}} \left[ \frac{z_1z_2}{z_{12}} \left( \frac{\chi_{23}z_i - z_3\chi_{2i}}{z_2\chi_{3i}} - \frac{2\chi_{2i}}{\chi_{3i}} - 1 \right) - \frac{z_{23}(-\chi_{12}z_i + z_2\chi_{1i} + z_1\chi_{2i})}{z_3\chi_{i(12)}} \right] \\
 & - \chi_{ij} \left[ \frac{z_3\chi_{12} - z_2\chi_{13} + z_1\chi_{23}}{z_{123}\chi_{3i}\chi_{j(12)}} + \frac{1}{\chi_{i(123)}} \left( \frac{z_2\chi_{13} + z_1\chi_{23} - z_3\chi_{12}}{z_{12}\chi_{3j}} \right. \right. \\
 & + \left. \left. \frac{(2z_2 + z_3)\chi_{12} + z_2\chi_{13} - z_1\chi_{23}}{z_3\chi_{j(12)}} \right) \right] - z_i \left[ \frac{1}{\chi_{3i}} \left( \frac{u_{2;j}}{z_{12}\chi_{j(123)}} + \frac{u_{1;j}}{z_{123}\chi_{j(12)}} \right) \right. \\
 & + \left. \frac{1}{\chi_{i(12)}} \left( \frac{\chi_{12}\chi_{3j} + u_{3;j}}{z_{123}\chi_{3j}} + \frac{u_{2;j}}{z_3\chi_{j(123)}} \right) + \frac{1}{\chi_{i(123)}} \left( \frac{\chi_{12}\chi_{3j} + u_{3;j}}{z_{12}\chi_{3j}} + \frac{u_{1;j}}{z_3\chi_{j(12)}} \right) \right] \\
 & - \frac{1}{\chi_{i(123)}} \left[ \frac{\chi_{2i}(z_3\chi_{1j} - z_1(2\chi_{2j} + \chi_{3j})) + r_{ij}}{z_{12}\chi_{3j}} + \frac{\chi_{2i}(z_1\chi_{3j} - (2z_2 + z_3)\chi_{1j}) + r_{ij}}{z_3\chi_{j(12)}} \right] \\
 & - \frac{z_2\chi_{1j}\chi_{3i} - \chi_{3j}(z_2\chi_{1i} + z_1\chi_{2i}) - z_1\chi_{2j}\chi_{3i}}{z_{123}\chi_{3j}\chi_{i(12)}} \\
 & + \left. \frac{z_3\chi_{1i}\chi_{2j} + \chi_{2i}((2z_2 + z_3)\chi_{1j} + 2z_1\chi_{2j})}{z_{123}\chi_{3i}\chi_{j(12)}} \right\} \\
 & + 5 \text{ permutations of } (ijk) \Big\} - (1 \leftrightarrow 2), \tag{4.22}
 \end{aligned}$$

with

$$\begin{aligned}
 u_{1;j} &= -\chi_{12}\chi_{3j} + \chi_{13}\chi_{2j} + \chi_{1j}\chi_{23}, \\
 u_{2;j} &= \chi_{12}\chi_{3j} - \chi_{13}\chi_{2j} + \chi_{1j}\chi_{23}, \\
 u_{3;j} &= \chi_{2j}(2\chi_{12} + \chi_{13}) - \chi_{1j}\chi_{23}, \\
 r_{ij} &= -z_2(\chi_{1i}\chi_{3j} + \chi_{1j}\chi_{3i}) - \chi_{2j}((z_2 + z_{23})\chi_{1i} + z_1\chi_{3i}). \tag{4.23}
 \end{aligned}$$

- The function multiplying the symmetrised colour correlation in eq. (4.16) is given by

$$U_{ijkl;srt}^{\bar{q}qg} = \frac{1}{2} U_{ik;sr}^{q\bar{q}} U_{jl;t}, \tag{4.24}$$

with

$$U_{j\ell;t} = \frac{2z_j}{z_t\chi_{jt}} + \frac{2z_\ell}{z_t\chi_{lt}} - \mathcal{S}_{j\ell}(p_t), \tag{4.25}$$

$$\begin{aligned}
 U_{ik;sr}^{\bar{q}q} &= \frac{4}{\chi_{sr}^2} \left[ \frac{1}{4} \chi_{sr}^2 Q_{ik}^{\bar{q}q}(p_s, p_r) - \frac{z_s\chi_{ri} + z_r\chi_{si} - z_i\chi_{sr}}{z_{sr}\chi_{i(sr)}} \right. \\
 & \left. - \frac{z_s\chi_{rk} + z_r\chi_{sk} - z_k\chi_{sr}}{z_{sr}\chi_{k(sr)}} + \frac{2z_s z_r}{z_{sr}^2} \right]. \tag{4.26}
 \end{aligned}$$

We note that just like the function (3.22), eq. (4.24) is the product of two functions, one associated to the single-soft limit and the other to the soft- $\bar{q}q$  limit of splitting amplitudes introduced in ref. [28].

Finally, we note that, just like in the two-soft-gluon case, all the kinematic terms in the factorisation formula (4.16) of the splitting amplitude are independent of whether the tree amplitude describing the hard scattering contains massive partons or not.

Supplementary material for the coefficients of eq. (4.16) is provided in a computer-readable format with this published work.

### 4.3.1 Soft $\bar{q}qg$ limit of the quadruple collinear splitting amplitudes

So far all considerations were completely generic and hold for the soft- $\bar{q}qg$  limit of any splitting amplitude. We now focus on the soft- $\bar{q}qg$  limit within a quadruple collinear splitting  $f \rightarrow \bar{q}qgf$ , and we work out all the color factors explicitly. Equation (4.16) becomes

$$\begin{aligned} \mathcal{S}_{123} \left[ \left( \frac{2\mu^{2\epsilon} g_s^2}{s_{1234}} \right)^3 \hat{\mathbf{P}}_{\bar{q}qgf} \right] &= (\mu^{2\epsilon} g_s^2)^3 \frac{1}{\mathcal{C}_f} \left[ \left( C_F U_{44;123}^{\bar{q}qg(ab)} + C_A U_{44;123}^{\bar{q}qg(nab)} \right) |\mathbf{Sp}_{ff}^{(44)}|^2 \right. \\ &\quad \left. + U_{444;123}^{\bar{q}qg} |\mathbf{Sp}_{ff}^{(444)}|^2 + U_{4444;123}^{\bar{q}qg} |\mathbf{Sp}_{ff}^{(44;44)}|^2 \right], \end{aligned} \quad (4.27)$$

where the kinematic coefficients are given in eq. (4.17), and the colour-correlated splitting matrices are defined in eq. (4.17), with  $\mathbf{Sp}_{ff'} = \delta_{ff'}$  (note that  $\mathbf{Sp}_{ff'}$  acts as the identity on both colour and spin indices). We can now evaluate the colour correlations in eq. (4.17) explicitly. For the dipole and symmetric four-parton correlations, the colour algebra is trivial,

$$\frac{1}{\mathcal{C}_f} |\mathbf{Sp}_{ff}^{(44)}|^2 = C_4 \quad \text{and} \quad \frac{1}{\mathcal{C}_f} |\mathbf{Sp}_{ff}^{(44;44)}|^2 = 2 C_4^2, \quad (4.28)$$

where  $C_4$  denotes the Casimir in the representation of the fourth collinear parton, with flavour  $f$ . For the three-parton correlation we need to distinguish two cases, depending on the flavour  $f$ . If  $f = g$ , we have  $(\mathbf{T}_4^c)_{ab} = i f^{a_4 c b_4}$ , and using  $d^{abc} = 2 \text{Tr} \left( \{t^a, t^b\} t^c \right)$  and  $f^{abc} = -2i \text{Tr} \left( [t^a, t^b] t^c \right)$ , one can check that the three-parton correlation vanishes. If  $f$  is an (anti-)quark instead,  $(\mathbf{T}_4^c)_{ab} = t_{a_4 b_4}^c$ , we repeatedly use the identity,

$$t^a t^b = \frac{1}{2} \left[ \frac{1}{N_c} \delta^{ab} \mathbf{1} + \left( d^{abc} + i f^{abc} \right) t^c \right], \quad (4.29)$$

and we arrive at

$$d^{abc} t^a t^b t^c = \frac{(N_c^2 - 1)(N_c^2 - 4)}{4N_c}, \quad (4.30)$$

where we used that  $d^{aab} = 0$  and  $d^{abc} f^{abd} = 0$ , and  $d^{abc} d^{abd} = \frac{N_c^2 - 4}{N_c} \delta^{cd}$ . Hence, we obtain that

$$|\mathbf{Sp}_{ff}^{(444)}|^2 = \delta_{fq} \frac{(N_c^2 - 1)(N_c^2 - 4)}{4N_c}, \quad (4.31)$$

where

$$\delta_{fq} = \begin{cases} 0, & \text{if } f = g, \\ 1, & \text{otherwise.} \end{cases} \quad (4.32)$$

Putting it all together, we see that for  $m = 4$ , eq. (4.27) can be cast in the compact form,

$$\begin{aligned} \mathcal{S}_{123} \left[ \left( \frac{2\mu^{2\epsilon} g_s^2}{s_{1234}} \right)^3 \hat{\mathbf{P}}_{\bar{q}qgf} \right] &= (\mu^{2\epsilon} g_s^2)^3 \left[ C_A^2 \left( U_{44;123}^{\bar{q}qg(\text{nab})} + 2 U_{4444;123}^{q\bar{q}g} \right) \right. \\ &\quad \left. + C_F C_A U_{44;123}^{\bar{q}qg(\text{ab})} + \delta_{fq} \frac{(N_c^2 - 1)(N_c^2 - 4)}{4N_c} U_{444;123}^{q\bar{q}g} \right]. \end{aligned} \quad (4.33)$$

Equation (4.33) describes quadruple splitting amplitudes  $\hat{\mathbf{P}}_{\bar{q}qgf}$  in the triple soft limit. We stress that the derivation of eq. (4.33) only requires the knowledge of the triple soft current of section 3. In particular, we did not need any explicit results for the quadruple splitting amplitudes. We checked, however, that eq. (4.33) agrees with what we obtain by taking the soft limit of the quadruple collinear splitting amplitudes for  $f \in \{g, q, q'\}$  of refs. [28, 29].

## 5 Conclusions

In this work we have computed for the first time the tree-level current for the emission of a soft quark-antiquark pair in addition to a gluon. Our result is an operator in the colour space of the hard partons, and it is valid for any number of hard particles, irrespective of their flavour, spin, colour or mass.

We have also considered the square of the soft current and the colour correlations it induces on the squared matrix elements summed over colour and spin quantum numbers. Remarkably, we find that there are only three different types of colour correlations. These include dipole correlations, which correlate two hard partons and appear already for the emission of a single soft gluon, and symmetrised four-parton correlations that show up in two-gluon emissions. A novel feature of our result is the appearance of tripole correlations involving the totally symmetric structure constant  $d^{abc}$ . To the best of our knowledge, this is the first time that such colour correlations have been observed in soft limits of tree-level matrix elements.

If our soft current is used in the context of an  $N^3\text{LO}$  computation, e.g., to build a subtraction scheme at  $N^3\text{LO}$ , it is important to understand also kinematic sublimits, i.e., limits where a subset of unresolved particles develop additional singularities. We have in particular studied the strongly-ordered soft limits in which the gluon is softer than the quark pair (or vice-versa). We have worked out how collinear splitting amplitudes behave if a subset  $\bar{q}qg$  of collinear particles becomes soft. This leads to the same type of universal factorisation of splitting amplitudes as in refs. [28, 29], and we have derived the universal building blocks entering this factorisation in detail.

Our soft current was the last missing ingredient to describe all kinematic infrared singularities that can arise in  $N^3\text{LO}$  computations. Having a complete set of infrared limits was a crucial step towards building a subtraction scheme at NNLO. With the completion of this work, we have now a complete set of limits also at  $N^3\text{LO}$ , which opens the way to the development of subtraction schemes to combine real and virtual corrections at  $N^3\text{LO}$  in the future.

## Acknowledgments

We thank Einan Gardi for useful discussions and for a fruitful collaboration through the early stages of this work.

## A Two-parton colour correlated soft function

In this appendix we provide the explicit expression for the coefficients of  $C_A$  in eq. (3.9), with

$$Q_{ij}^{\bar{q}qg(\text{nab})} = \left[ \frac{1}{4} \mathcal{S}_{ij}(p_3) \left( \mathcal{S}_{1i}(p_{12}) \mathcal{S}_{2i}(p_{12}) - \frac{3}{4} Q_{ij}^{\bar{q}q}(p_1, p_2) \right) + \frac{1}{\chi_{123}^2 \chi_{i(123)} \chi_{j(123)}} \left( a_0 + a_1 \chi_{ij} + a_2 \chi_{ij}^2 \right) + (1 \leftrightarrow 2) \right] + (i \leftrightarrow j), \quad (\text{A.1})$$

and

$$Q_{ij}^{\bar{q}qg(\text{mass})} = m_i^2 \left[ \frac{1}{2\chi_{3i}^2} Q_{ij}^{\bar{q}q}(p_1, p_2) - \frac{1}{\chi_{12} \chi_{i(12)}} \mathcal{S}_{ij}(p_3) + \frac{2}{\chi_{123} \chi_{i(12)} \chi_{j(12)} \chi_{i(123)} \chi_{j(123)}} (c_0 + c_1 \chi_{ij}) + (1 \leftrightarrow 2) \right] + (i \leftrightarrow j). \quad (\text{A.2})$$

Like for the coefficient of  $C_F$  given in eq. (3.23), here we have exploited the dipole symmetry under the exchange of the indices labelling the hard emitters, as well as the charge conjugation symmetry of the  $q\bar{q}$  current. On the second lines of eqs. (A.1) and (A.2) we have extracted a pre-factor which captures the overall scaling of  $\mathcal{O}(\lambda^{-6})$  in the triple soft limit.

The coefficients  $a_0$ ,  $a_1$  and  $a_2$  on the right-hand side of eq. (A.1) are given by,

$$a_2 = \frac{\chi_{3(12)}}{\chi_{i(12)} \chi_{j(12)}} \left\{ \frac{\chi_{12}}{\chi_{3(12)}} \left( \frac{1}{2} - \frac{2\chi_{1i}}{\chi_{3i}} \right) + \frac{2\chi_{13}}{\chi_{3(12)}} + \frac{1}{\chi_{12}} \left[ \chi_{13} + \frac{\chi_{3(12)}}{\chi_{3i}} \left( \frac{\chi_{1j} \chi_{i(12)}}{\chi_{3j}} - 2\chi_{1i} \right) \right] + \frac{1}{\chi_{3i}} \left( \frac{2\chi_{1j} \chi_{i(12)}}{\chi_{3j}} - 4\chi_{1i} \right) \right\} + \frac{\chi_{12}}{2\chi_{3i} \chi_{3j}}, \quad (\text{A.3})$$

$$a_1 = D + \frac{2\chi_{12}^2}{\chi_{13} \chi_{23}} - \frac{2\chi_{12} \chi_{1i}}{\chi_{13} \chi_{3i}} - \frac{\chi_{1j} \chi_{2i}}{\chi_{3i} \chi_{3j}} + \frac{\chi_{3(12)}}{\chi_{i(12)} \chi_{j(12)}} \left\{ \frac{\chi_{12}}{\chi_{3(12)}} \left[ \chi_{1i} \left( \frac{10\chi_{1j}}{\chi_{13}} + \frac{8\chi_{1j}}{\chi_{23}} \right) + \chi_{2i} \left( \frac{9\chi_{1j}}{\chi_{13}} + \frac{9\chi_{1j}}{\chi_{23}} \right) \right] - \frac{6\chi_{1i}^2 \chi_{1j}}{\chi_{3(12)} \chi_{3i}} - \frac{4\chi_{1j} \chi_{3i}}{\chi_{3(12)}} - \frac{\chi_{2i}^2}{\chi_{3(12)} \chi_{3i}} \left( \frac{\chi_{13}^2 \chi_{1j}}{\chi_{12} \chi_{23}} + \frac{2\chi_{13} \chi_{1j}}{\chi_{23}} + 2\chi_{1j} \right) + \frac{\chi_{1j}}{\chi_{12}^2 \chi_{3(12)}} \left[ \chi_{2i} \left( \frac{\chi_{1j} (\chi_{13}^2 - \chi_{23}^2)}{\chi_{3j}} + (\chi_{13} - \chi_{23})^2 - 4\chi_{13} \chi_{23} \right) + 4\chi_{23} \chi_{1i} (\chi_{23} - \chi_{13}) + \frac{\chi_{2i}^2 (\chi_{23}^2 - \chi_{13}^2)}{\chi_{3i}} \right] - \frac{\chi_{1j}}{\chi_{12} \chi_{3(12)}} \left[ \frac{\chi_{2i}}{\chi_{3j}} \left( \frac{4\chi_{13} \chi_{1i} \chi_{2j}}{\chi_{3i}} + 3\chi_{1j} \chi_{23} \right) + \frac{3\chi_{13} \chi_{2i}^2 + \chi_{1i}^2 (4\chi_{13} + 10\chi_{23})}{\chi_{3i}} - \chi_{1i} (4\chi_{13} + 10\chi_{23}) \right] + \frac{\chi_{2i} \chi_{1j}}{\chi_{3(12)}} \left[ \frac{6\chi_{13}}{\chi_{23}} + \frac{\chi_{23}^2}{\chi_{12} \chi_{13}} - \frac{\chi_{1i}}{\chi_{3i}} \left( \frac{2\chi_{13}^2}{\chi_{12} \chi_{23}} + \frac{4\chi_{13}}{\chi_{23}} + 10 \right) \right] + \frac{6\chi_{23}}{\chi_{13}} - \frac{\chi_{1j}}{\chi_{3j}} \left( \frac{\chi_{23}^2}{\chi_{12} \chi_{13}} + \frac{2\chi_{23}}{\chi_{13}} + 2 \right) + \frac{\chi_{13}^2}{\chi_{12} \chi_{23}} + 9 \right] - \frac{2\chi_{1i} \chi_{1j}}{\chi_{12}^2 \chi_{3i}} \left( \frac{\chi_{13} \chi_{2i} \chi_{2j}}{\chi_{3j}} + 2\chi_{1i} \chi_{23} \right) + \frac{1}{\chi_{12}} \left[ \chi_{2i} \left( 5\chi_{1j} - \frac{\chi_{1i}}{\chi_{3i}} \left( \frac{4\chi_{1j}^2}{\chi_{3j}} + 10\chi_{1j} \right) \right) - \frac{2\chi_{1j}^2 \chi_{2i}^2}{\chi_{3i} \chi_{3j}} - 4\chi_{1j} \chi_{3i} \right]$$

$$\begin{aligned}
 & -\frac{\chi_{3(12)}\chi_{2i}}{\chi_{12}^2\chi_{3i}} \left[ \chi_{1i} \left( \frac{2\chi_{1j}^2}{\chi_{3j}} + 2\chi_{1j} \right) + \frac{\chi_{1j}^2\chi_{2i}}{\chi_{3j}} \right] + \frac{\chi_{1i}}{\chi_{3(12)}} \left( \frac{2\chi_{1j}\chi_{23}^2}{\chi_{12}\chi_{13}} + \frac{8\chi_{1j}\chi_{23}}{\chi_{13}} \right. \\
 & - \left. \frac{2\chi_{1j}\chi_{23}\chi_{i(12)}}{\chi_{3i}\chi_{13}} \left( \frac{\chi_{23}}{\chi_{12}} + 2 \right) + \frac{4\chi_{13}\chi_{1j}}{\chi_{23}} + 10\chi_{1j} \right) \left\} - \frac{\chi_{123}}{\chi_{i(12)}\chi_{j(12)}} \left\{ \frac{\chi_{1j}\chi_{2i}^2}{\chi_{23}\chi_{3i}} \right. \right. \\
 & + \frac{2\chi_{1i}\chi_{1j}\chi_{2i}\chi_{3(12)}}{\chi_{13}\chi_{23}\chi_{3i}} + \frac{2\chi_{1i}^2\chi_{1j}}{\chi_{13}\chi_{3i}} + \frac{\chi_{1j}^2\chi_{2i}}{\chi_{13}\chi_{3j}} - \frac{4\chi_{1i}\chi_{1j}\chi_{2i}\chi_{j(12)}}{\chi_{12}\chi_{3i}\chi_{3j}} \\
 & + \frac{\chi_{3(12)}}{\chi_{i(12)}\chi_{j(12)}} \left[ \frac{2\chi_{1j}^2(2\chi_{1i}\chi_{2i} + \chi_{i(12)}^2)}{\chi_{23}\chi_{3(12)}} + \frac{2\chi_{1j}^2}{\chi_{13}\chi_{3(12)}} \left( \frac{4\chi_{1i}\chi_{2i}\chi_{2j}}{\chi_{1j}} + 2\chi_{1i}\chi_{2i} + \chi_{i(12)}^2 \right) \right. \\
 & + \frac{\chi_{1j}\chi_{3i}}{\chi_{13}} \left( \frac{\chi_{1j}(2\chi_{1i} + \chi_{2i})}{\chi_{3j}} \left( \frac{\chi_{2j}}{\chi_{23}} + \frac{\chi_{j(12)}}{\chi_{3(12)}} \right) + \frac{2\chi_{1i}(\chi_{1j} + 2\chi_{2j}) + \chi_{1j}\chi_{2i}}{\chi_{23}} \right) \\
 & + \frac{\chi_{1j}\chi_{2i}^2\chi_{3j}}{\chi_{13}\chi_{23}} + \frac{\chi_{2i}^2\chi_{3j}}{\chi_{23}\chi_{3i}} \left( \frac{\chi_{1i}\chi_{1j} + \chi_{1j}\chi_{2i}}{\chi_{3(12)}} + \frac{\chi_{1i}\chi_{1j}}{\chi_{13}} \right) \\
 & - \frac{\chi_{1j}^2\chi_{2i}^2}{\chi_{12}^2} \left\{ 2 + \frac{4\chi_{1i}}{\chi_{3i}} \left[ \frac{\chi_{23}}{\chi_{3(12)}} \left( \frac{\chi_{1i}}{\chi_{2i}} \left( \frac{\chi_{13}}{\chi_{23}} + 4 \right) + \frac{\chi_{13}}{2\chi_{23}} + 2 + \frac{\chi_{1i}^2}{\chi_{2i}^2} + \frac{\chi_{2i}}{2\chi_{1i}} \right) \right. \right. \\
 & + \left. \frac{\chi_{2j}}{\chi_{3j}} \left( \frac{\chi_{1i}}{\chi_{2i}} \left( \frac{\chi_{1j}}{\chi_{2j}} + 3 \right) + \frac{\chi_{1j}}{2\chi_{2j}} + 3 + \frac{\chi_{2i}}{2\chi_{1i}} \right) + \frac{2\chi_{1i}\chi_{2j}}{\chi_{1j}\chi_{2i}} \right] \\
 & + \left. \frac{2\chi_{23}}{\chi_{3(12)}} \left( \frac{\chi_{13}\chi_{1j} + \chi_{2j}(4\chi_{13} + \chi_{23})}{\chi_{23}\chi_{3j}} + \frac{2\chi_{1i}^2}{\chi_{2i}^2} + \frac{4\chi_{1i}}{\chi_{2i}} \right) \right\} \\
 & + \frac{2\chi_{1j}^2}{\chi_{12}} \left\{ \frac{\chi_{3i}\chi_{1i}}{\chi_{3(12)}} \left( \frac{\chi_{1i}(2\chi_{1j} + 6\chi_{2j}) + \chi_{2i}(\chi_{1j} + 3\chi_{2j})}{\chi_{1i}\chi_{3j}} + 2 \right) \right. \\
 & + \left. \frac{\chi_{1i}^2\chi_{2i}}{\chi_{3(12)}\chi_{3i}} \left[ \frac{8\chi_{2j}}{\chi_{1j}} + 1 + \frac{\chi_{3j}}{\chi_{1j}} \left( \frac{3\chi_{2i}}{\chi_{1i}} + \frac{\chi_{2i}\chi_{i(23)}}{\chi_{1i}^2} \right) + \frac{\chi_{1i}}{\chi_{2i}} \left( \frac{4\chi_{2j}}{\chi_{1j}} + 1 \right) \right. \right. \\
 & + \left. \left. \frac{\chi_{2i}}{2\chi_{1i}} + \frac{\chi_{2i}^2}{2\chi_{1i}^2} \right] + \frac{\chi_{1i}\chi_{2i}}{\chi_{3(12)}} \left[ \frac{4\chi_{2j}\chi_{i(23)}}{\chi_{1j}\chi_{2i}} + 6 + \frac{\chi_{1i}}{\chi_{2i}} + \frac{\chi_{2i}\chi_{j(12)}}{2\chi_{3j}\chi_{1i}} + \frac{\chi_{i(23)}}{\chi_{1i}} \right] \right\} \left. \right\}, \tag{A.4} \\
 a_0 = & \frac{\chi_{3(12)}}{\chi_{i(12)}\chi_{j(12)}} \left\{ \frac{2\chi_{1i}^2\chi_{1j}^2}{\chi_{13}\chi_{3(12)}} \left( \frac{\chi_{23}}{\chi_{12}} + 2 \right) + \frac{\chi_{1j}\chi_{1i}^2\chi_{2i}^2}{\chi_{12}\chi_{3(12)}\chi_{3i}} \left( \frac{2\chi_{1i}\chi_{23}\chi_{2j}}{\chi_{13}\chi_{2i}^2} + \frac{\chi_{13}\chi_{1j}\chi_{2i}}{\chi_{23}\chi_{1i}^2} \right) \right. \\
 & + \frac{\chi_{1i}\chi_{1j}^2\chi_{23}\chi_{2i}}{\chi_{12}^2\chi_{3(12)}} \left( \frac{4\chi_{1i} + 4\chi_{3i}}{\chi_{2i}} - \frac{2\chi_{2i}\chi_{3j}}{\chi_{1i}\chi_{1j}} + \frac{2\chi_{3i}}{\chi_{1i}} - \frac{8\chi_{2j}}{\chi_{1j}} + 8 \right) + \frac{\chi_{2i}\chi_{3i}\chi_{1j}^2}{\chi_{13}\chi_{23}} \left( \frac{\chi_{2j}}{\chi_{3j}} - \frac{6\chi_{3j}}{\chi_{1j}} \right) \\
 & + \frac{2\chi_{1j}\chi_{2i}(4\chi_{1i}\chi_{2j} + \chi_{2i}\chi_{3j})}{\chi_{12}^2} + \frac{\chi_{1j}^2}{\chi_{3(12)}} \left\{ \frac{\chi_{2i}\chi_{1i}}{\chi_{12}} \left[ \frac{\chi_{1i}\chi_{23}}{\chi_{13}\chi_{3i}} \left( \frac{\chi_{2j}(2\chi_{13}^2 + 4\chi_{23}^2)}{\chi_{23}^2\chi_{1j}} + 2 \right) \right. \right. \\
 & - \left. \frac{4\chi_{13}}{\chi_{23}} \left( \frac{\chi_{13}^2 + \chi_{23}^2}{\chi_{13}^2} + \frac{\chi_{2j}}{\chi_{1j}} \right) + \frac{\chi_{3i}}{\chi_{13}\chi_{23}\chi_{1i}} \left( \frac{\chi_{2j}(\chi_{13}^2 + \chi_{23}^2)}{\chi_{3j}} + 3\chi_{23}^2 - \chi_{13}^2 \right) \right] \\
 & + \frac{2\chi_{1i}\chi_{2i}}{\chi_{13}} \left[ \frac{\chi_{3i}(3\chi_{23} - 2\chi_{13})}{2\chi_{23}\chi_{1i}} - \frac{5\chi_{13} + 3\chi_{23}}{\chi_{23}} - \frac{4\chi_{2j}}{\chi_{1j}} \right] \left. \right\} + \frac{2\chi_{1i}\chi_{1j}^2\chi_{3i}}{\chi_{13}\chi_{23}} \left( \frac{\chi_{2j}}{\chi_{3j}} - \frac{3\chi_{3j}}{\chi_{1j}} \right) \\
 & + \frac{2\chi_{1i}\chi_{1j}}{\chi_{3(12)}} \left\{ \frac{\chi_{1j}\chi_{3i}}{\chi_{12}} \left[ \frac{\chi_{2j}}{\chi_{3j}} \left( \frac{\chi_{13}}{\chi_{23}} + \frac{\chi_{23}}{\chi_{13}} \right) + \frac{3\chi_{2(13)}}{\chi_{13}} - \frac{\chi_{12}}{\chi_{23}} \right] + \frac{\chi_{2j}\chi_{3i}}{\chi_{12}} \left( \frac{2\chi_{13} + \chi_{12}}{\chi_{23}} + \frac{\chi_{23}}{\chi_{13}} \right) \right\} \\
 & + \frac{\chi_{1j}^2\chi_{2i}^2}{\chi_{13}\chi_{23}} \left( \frac{\chi_{1i}\chi_{3j}}{\chi_{1j}\chi_{3i}} - 6 \right) + \frac{\chi_{2i}^2}{\chi_{3(12)}} \left[ \frac{\chi_{1j}\chi_{3j}}{\chi_{12}} \left( \frac{3\chi_{1(23)}}{\chi_{23}} - \frac{\chi_{23}}{\chi_{13}} - \frac{2\chi_{12}}{\chi_{13}} \right) \right. \\
 & + \left. \frac{\chi_{1i}\chi_{1j}\chi_{j(13)}}{\chi_{12}\chi_{3i}} \left( \frac{\chi_{13}}{\chi_{23}} + \frac{\chi_{23}}{\chi_{13}} \right) + \frac{\chi_{1j}^2}{\chi_{12}} \left( \frac{\chi_{2j}}{\chi_{3j}} \left( \frac{\chi_{13}}{\chi_{23}} + \frac{\chi_{23}}{\chi_{13}} \right) - \frac{3\chi_{13}}{\chi_{23}} - \frac{3\chi_{23}}{\chi_{13}} \right) + \frac{\chi_{1j}^3\chi_{23}}{\chi_{12}\chi_{13}\chi_{3j}} \right] \\
 & + \frac{\chi_{1j}^2\chi_{2i}^2}{\chi_{12}^2} \left[ \frac{\chi_{2j}}{\chi_{2i}^2} \left( \frac{4\chi_{1i}^2\chi_{i(12)}}{\chi_{1j}\chi_{3i}} + \frac{4\chi_{1i}\chi_{3i}}{\chi_{3j}} \right) + \frac{2\chi_{1i}\chi_{j(13)}}{\chi_{1j}\chi_{3i}} + \frac{2\chi_{j(12)}}{\chi_{3j}} + \frac{2\chi_{2i}}{\chi_{3i}} + \frac{2\chi_{2j}\chi_{3i}}{\chi_{2i}\chi_{3j}} - 2 \right]
 \end{aligned}$$



$$\begin{aligned}
 & + \frac{\chi_{1j}^2 \chi_{2i}^2}{\chi_{12} \chi_{3(12)}} \left\{ \frac{4\chi_{1i} \chi_{j(13)}}{\chi_{1j} \chi_{3i}} + \frac{\chi_{1i} \chi_{3i}}{\chi_{2i}^2} \left( \frac{8\chi_{2j}}{\chi_{3j}} - \frac{6\chi_{j(23)}}{\chi_{1j}} + 2 + \frac{4\chi_{i(12)}}{\chi_{3i}} + \frac{6\chi_{1i}^2 \chi_{2j}}{\chi_{1j} \chi_{3i}^2} \right) + \frac{3\chi_{2i}}{\chi_{3i}} \right. \\
 & + \left. \frac{3\chi_{j(12)} + \chi_{2j}}{\chi_{3j}} + \frac{\chi_{1i}}{\chi_{2i}} \left[ \frac{\chi_{1i}}{\chi_{3i}} \left( \frac{10\chi_{2j}}{\chi_{1j}} + 2 \right) + \frac{1}{\chi_{1i}} \left( \frac{4\chi_{2j} \chi_{3i}}{\chi_{3j}} - \frac{6\chi_{3i} \chi_{3j}}{\chi_{1j}} - 8\chi_{i(12)} \right) \right] \right\} \\
 & - \frac{\chi_{12} \chi_{1j}^2 \chi_{2i}^2}{\chi_{13} \chi_{23} \chi_{3(12)}} \left[ \frac{4\chi_{1i} \chi_{3i}}{\chi_{2i}^2} \left( \frac{2\chi_{j(23)}}{\chi_{1j}} + 1 \right) + \frac{2\chi_{3i}}{\chi_{2i}} \left( \frac{4\chi_{3j}}{\chi_{1j}} + 1 \right) + \frac{2\chi_{3j}}{\chi_{1j}} \right] \\
 & + \frac{D}{2} \left[ \frac{\chi_{23} (2\chi_{1i} \chi_{1j} + \chi_{1j} \chi_{2i})}{\chi_{12} \chi_{13}} + \frac{\chi_{13} \chi_{1j} \chi_{2i}}{\chi_{12} \chi_{23}} + \chi_{3i} \chi_{3j} \left( \frac{2\chi_{12}}{\chi_{13} \chi_{23}} + \frac{1}{\chi_{12}} + \frac{2}{\chi_{13}} \right) \right. \\
 & - \left. \chi_{3i} \left( \frac{2\chi_{1j} \chi_{23}}{\chi_{12} \chi_{13}} + \frac{2\chi_{1j}}{\chi_{12}} + \frac{6\chi_{1j}}{\chi_{13}} + \frac{2\chi_{1j}}{\chi_{23}} \right) \right] - \frac{4\chi_{12} \chi_{1j} \chi_{2i}}{\chi_{13} \chi_{23}} + \frac{\chi_{1j}^2 \chi_{2i}}{\chi_{13} \chi_{3j}} \\
 & + \frac{1}{\chi_{3i}} \left( \frac{2\chi_{1i} \chi_{1j} \chi_{2i}}{\chi_{13}} + \frac{\chi_{1j} \chi_{2i}^2}{\chi_{23}} \right) + \frac{\chi_{123}}{\chi_{i(12)} \chi_{j(12)}} \left\{ \frac{\chi_{1j}^2 \chi_{2i}}{\chi_{12} \chi_{3i}} \left[ \frac{2\chi_{1i}^2}{\chi_{13}} \left( 1 - \frac{\chi_{2j} (\chi_{3(12)} + \chi_{23})}{\chi_{23} \chi_{1j}} \right) \right. \right. \\
 & + \left. \left. \frac{\chi_{2i}^2}{\chi_{23}} - \frac{2\chi_{1i}^3 \chi_{2j}}{\chi_{13} \chi_{1j} \chi_{2i}} + \frac{\chi_{1i} \chi_{2i} \chi_{3(12)}}{\chi_{13} \chi_{23}} \right] + \frac{\chi_{1j}^3 \chi_{2i}^2}{\chi_{12} \chi_{13} \chi_{3j}} \left( \frac{\chi_{2j} \chi_{3(12)}}{\chi_{1j} \chi_{23}} + 1 \right) \right. \\
 & + \frac{\chi_{3(12)}}{\chi_{i(12)} \chi_{j(12)}} \left\{ \frac{2\chi_{1j}^2 \chi_{2i}^2 (\chi_{1j} \chi_{2i} - \chi_{1i} \chi_{2j})}{\chi_{12} \chi_{13} \chi_{23}} + \frac{\chi_{1j}^2 \chi_{2i}}{\chi_{12} \chi_{3(12)}} \left( \frac{\chi_{1i} \chi_{1j} (\chi_{2i} - 2\chi_{1i}) + 3\chi_{2j} (\chi_{2i}^2 - 2\chi_{1i}^2)}{\chi_{13}} \right) \right. \\
 & + \left. \frac{\chi_{1i} \chi_{1j} (2\chi_{1i} + 3\chi_{2i}) + \chi_{2j} (\chi_{2i}^2 - 2\chi_{1i}^2)}{\chi_{23}} \right\} + \frac{4\chi_{1j}^2 \chi_{2i} \chi_{i(12)} (\chi_{2i} \chi_{j(12)} - 2\chi_{1i} \chi_{2j})}{\chi_{12}^2 \chi_{3(12)}} \\
 & + \frac{4\chi_{1i}^2 \chi_{1j}^2 \chi_{2i}^2}{\chi_{12}^2 \chi_{3(12)} \chi_{3i}} \left( \frac{\chi_{2i} (3\chi_{1j} - \chi_{2j})}{2\chi_{1i}} + \frac{\chi_{1i} (\chi_{1j} - 3\chi_{2j})}{\chi_{2i}} + (3\chi_{1j} - 3\chi_{2j}) \right) \\
 & + \frac{\chi_{1j} \chi_{2i}^2}{2\chi_{1i}^2} - \frac{\chi_{1i}^2 \chi_{2j}}{\chi_{2i}^2} + \frac{\chi_{1j}^2 \chi_{2i}^2 \chi_{3j}}{\chi_{12} \chi_{3(12)} \chi_{3i}} \left( \chi_{1i} \chi_{i(23)} \left( \frac{1}{\chi_{13}} - \frac{1}{\chi_{23}} \right) + \frac{\chi_{2i} \chi_{3i}}{\chi_{13}} \right) \\
 & + \frac{2\chi_{1j}^3 \chi_{2i}^2 (\chi_{2i} (\chi_{1j} + 3\chi_{2j}) - \chi_{1i} \chi_{2j})}{\chi_{12}^2 \chi_{3(12)} \chi_{3j}} + \frac{\chi_{1j}^2 \chi_{2j} \chi_{3i} (\chi_{13} - \chi_{23}) (\chi_{1j} \chi_{2i}^2 - 2\chi_{1i}^2 (\chi_{1j} + 2\chi_{2j}))}{\chi_{12} \chi_{13} \chi_{23} \chi_{3(12)} \chi_{3j}} \\
 & + \left. \frac{\chi_{1j}^2 \chi_{3i} \chi_{2i}}{\chi_{12} \chi_{3(12)}} \left[ \frac{\chi_{1i}}{\chi_{23}} \left( \frac{\chi_{2i} \chi_{j(12)}}{\chi_{1i}} - \frac{2\chi_{1i} \chi_{2j}}{\chi_{2i}} + 2\chi_{1j} \right) - \frac{\chi_{2j} (2\chi_{1i} + \chi_{2i})}{\chi_{13}} \right] \right\}. \tag{A.5}
 \end{aligned}$$

The coefficients  $c_0$  and  $c_1$  on the right-hand side of eq. (A.2) are given by,

$$\begin{aligned}
 c_1 = & \frac{\chi_{3(12)}}{\chi_{i(12)} \chi_{j(12)}} \left\{ \frac{\chi_{2i} \chi_{2j} \chi_{j(12)}}{\chi_{3(12)} \chi_{3i}} \left( \frac{\chi_{1i}^2 \chi_{3j}}{\chi_{2i} \chi_{2j} \chi_{3i}} + \frac{\chi_{2i} \chi_{3j}}{\chi_{2j} \chi_{3i}} + \frac{\chi_{2j} \chi_{3i}}{\chi_{2i} \chi_{3j}} + \frac{\chi_{3i}}{\chi_{2i}} - \frac{\chi_{3j}}{\chi_{2j}} - 2 \right) \right. \\
 & + \frac{\chi_{123}}{\chi_{12}} \frac{\chi_{j(12)}}{\chi_{3j}} \left[ \frac{\chi_{1j} \chi_{2j}}{\chi_{3(12)}} \left( \frac{2\chi_{1i} \chi_{2i} \chi_{3j}^2}{\chi_{1j} \chi_{2j} \chi_{3i}^2} - \frac{2\chi_{1i} \chi_{3j} \chi_{j(12)}}{\chi_{1j} \chi_{2j} \chi_{3i}} - \frac{\chi_{1i} \chi_{3j}^2}{\chi_{1j} \chi_{2j} \chi_{3i}} + \frac{\chi_{1j}}{\chi_{2j}} \right. \right. \\
 & - \left. \left. \frac{2\chi_{2i} \chi_{3j}}{\chi_{2j} \chi_{3i}} + \frac{\chi_{3j}}{\chi_{2j}} + 2 \right) + \frac{\chi_{1j} \chi_{2i} \chi_{3j}}{\chi_{123} \chi_{3i}} \left( \frac{\chi_{1i}^2 \chi_{j(123)}}{\chi_{1j} \chi_{2i} \chi_{3i}} - \frac{\chi_{1i} \chi_{j(12)}}{\chi_{1j} \chi_{2i} \chi_{3j}} + \frac{2\chi_{1i} \chi_{j(12)}}{\chi_{1j} \chi_{3i}} \right) \right. \\
 & + \left. \frac{\chi_{2j} \chi_{3i} \chi_{j(23)}}{\chi_{1j} \chi_{2i} \chi_{3j}} + \frac{\chi_{2i} \chi_{j(23)}}{\chi_{1j} \chi_{3i}} - \frac{\chi_{2j} \chi_{j(23)}}{\chi_{1j} \chi_{3j}} - \frac{\chi_{1j}}{\chi_{3j}} - \frac{\chi_{j(23)}}{\chi_{1j}} + \frac{\chi_{2i}}{\chi_{3i}} - \frac{2\chi_{2j}}{\chi_{3j}} \right) \left. \right\} \\
 & - \frac{\chi_{j(12)} (\chi_{3i} \chi_{j(12)} - \chi_{3j} \chi_{i(12)})}{\chi_{3i}^2 \chi_{3j}}, \tag{A.6}
 \end{aligned}$$

and

$$\begin{aligned}
 c_0 = & \frac{\chi_{3(12)}}{\chi_{i(12)}\chi_{j(12)}} \left\{ \frac{\chi_{1j}\chi_{2j}\chi_{i(12)}\chi_{j(12)}}{\chi_{23}\chi_{3(12)}} \left( \frac{\chi_{3j}^2\chi_{i(12)}}{\chi_{1j}\chi_{2j}\chi_{3i}} + \frac{\chi_{3j}\chi_{i(123)}\chi_{j(12)}}{\chi_{1j}\chi_{2j}\chi_{3i}} + \frac{3\chi_{2j}}{\chi_{1j}} + 4 \right) \right. \\
 & - \frac{\chi_{13}\chi_{23}\chi_{i(12)}\chi_{j(12)}}{\chi_{12}^2\chi_{3(12)}} \left[ \frac{\chi_{i(12)} \left( \chi_{123}\chi_{3j} + \chi_{3(12)}\chi_{j(12)} \right) (\chi_{1i}\chi_{2j} + \chi_{1j}\chi_{2i})}{\chi_{13}\chi_{23}\chi_{3i}^2} \right. \\
 & + \frac{\chi_{j(123)}}{\chi_{3i}} \left( \frac{\chi_{1j} (2\chi_{1i} + \chi_{2i}) - \chi_{1i}\chi_{2j}}{\chi_{13}} - \frac{\chi_{1j}\chi_{2i} - \chi_{2j} (\chi_{1i} + 2\chi_{2i})}{\chi_{23}} \right) \left. \right] \\
 & + \frac{\chi_{1j}\chi_{2j}\chi_{i(12)}\chi_{j(12)}}{\chi_{12}\chi_{3(12)}} \left[ \frac{\chi_{13}}{\chi_{23}} \left( \frac{\chi_{2j}}{\chi_{1j}} + 1 \right) + \frac{2\chi_{2j}}{\chi_{1j}} + 3 + \frac{\chi_{1j}\chi_{2(13)}}{\chi_{23}\chi_{2j}} \right. \\
 & + \frac{\chi_{j(12)}}{\chi_{1j}\chi_{3j}} \left( \frac{\chi_{13}}{\chi_{23}} + \frac{\chi_{1j}}{\chi_{2j}} + 2 \right) + \frac{\chi_{3j}}{\chi_{3i}} \left( \frac{3\chi_{1i}}{\chi_{1j}} + \frac{\chi_{1i}}{\chi_{2j}} - \frac{\chi_{13}}{\chi_{23}} \left( \frac{\chi_{1i}}{\chi_{1j}} + \frac{\chi_{2i}}{\chi_{1j}} \right) \right. \\
 & + \frac{2\chi_{2i}}{\chi_{2j}} + \frac{2\chi_{2i}}{\chi_{1j}} \left. \right) + \frac{\chi_{1i}\chi_{j(12)}}{\chi_{1j}\chi_{3i}} - \frac{\chi_{1i}\chi_{j(12)}}{\chi_{2j}\chi_{3i}} + \frac{2\chi_{3j}^2\chi_{i(12)}}{\chi_{1j}\chi_{2j}\chi_{3i}} - \frac{\chi_{13}\chi_{i(12)}\chi_{j(12)}}{\chi_{1j}\chi_{23}\chi_{3i}} \left. \right] \left. \right\} \\
 & + \frac{2\chi_{2j}\chi_{j(12)}^2}{\chi_{23}\chi_{3j}} + \frac{\chi_{j(12)} (\chi_{3i} - \chi_{i(12)}) (\chi_{2(13)}\chi_{3j} - \chi_{2j}\chi_{3(12)})}{\chi_{12}\chi_{23}\chi_{3i}} \\
 & - \frac{\chi_{i(12)}\chi_{j(12)} (\chi_{1i}\chi_{2j} + \chi_{1j}\chi_{2i})}{\chi_{12}\chi_{3i}^2}. \tag{A.7}
 \end{aligned}$$

## B Collinear limit of the two-parton colour correlation

In this appendix we consider the  $u_{(\text{dip})}^{(\text{nab})}$  term in eq. (4.20),

$$\begin{aligned}
 u_{(\text{dip})}^{(\text{nab})} = & \frac{4}{\chi_{123}^2 z_{123}^2} \left\{ \frac{D}{4} \left[ \frac{z_3 - z_{12}}{\chi_{12}} \left( \frac{4\chi_{12}z_3 - 2\chi_{13}z_2}{\chi_{23}} - z_{12} \left( 1 + \frac{z_3}{z_3 - z_{12}} \right) + 2z_1 \right) \right. \right. \\
 & + \frac{2z_3(z_1 - z_3)}{\chi_{23}} - \frac{2z_1z_3}{\chi_{13}} + \frac{2\chi_{12}z_3^2}{\chi_{13}\chi_{23}} \left. \right] + \frac{z_1(z_1 - z_2 + 2z_3)}{\chi_{13}} \\
 & - \frac{z_3\chi_{12}(z_{12} + 2z_3)}{\chi_{13}\chi_{23}} + \frac{2z_1z_2\chi_{23}}{\chi_{12}^2} \left[ \frac{z_{13}\chi_{13}}{z_1\chi_{23}} \left( \frac{z_{12}}{z_3} + 1 \right) - \frac{z_3z_1}{z_2z_{12}} - \frac{\chi_{3(12)}z_{123}}{2z_{12}\chi_{23}} \right] \\
 & + \frac{z_1z_2}{\chi_{23}} \left( \frac{z_1 - z_{23} - z_3}{z_{12}} + \frac{z_{23}}{z_1} - \frac{3z_3^2}{z_1z_2} - \frac{4z_2 + z_3}{z_2} \right) + \frac{z_1z_2}{\chi_{12}} \left[ \frac{z_{123} + 2z_2}{z_3} \right. \\
 & + \frac{z_3}{z_{12}} \left( 2z_1 \left( \frac{1}{z_3} - \frac{1}{z_2} \right) - 7 - \frac{2z_2}{z_1} \right) - \frac{z_2 - 2z_3}{4z_1} + \frac{9(z_1 + 2z_3)}{4z_2} - 3 \\
 & + \frac{\chi_{13}}{\chi_{23}} \left( \frac{z_1 - z_{23} - z_3}{z_{12}} + \frac{z_2}{z_1} + \frac{2z_2}{z_3} - 2 + \frac{3z_3}{z_1} \right) \left. \right] + \frac{2z_1z_2(z_2\chi_{13} - z_3\chi_{12})}{z_3\chi_{13}\chi_{23}} \\
 & + \frac{z_3^2}{\chi_{12}} \left( \frac{1}{4}(D - 6) + \frac{(z_2^2\chi_{13} + z_1^2\chi_{23})}{z_3z_{12}\chi_{12}} \right) \left. \right\} \\
 & - \left[ \frac{4}{\chi_{123}^2 z_{123} \chi_{i(123)}} (c_0 + c_1 z_i + c_2 z_i^2) + (i \leftrightarrow j) \right] + (1 \leftrightarrow 2), \tag{B.1}
 \end{aligned}$$

with

$$\begin{aligned}
 c_0 = & (D-6) \frac{z_3 \chi_{1i} \chi_{3i}}{2\chi_{12} \chi_{i(12)}} + \frac{z_3}{\chi_{i(12)}} \left( \frac{\chi_{1i}^2 \chi_{23}}{\chi_{12}^2} + \frac{z_1^2 \chi_{i(12)} \chi_{23} \chi_{3i}}{z_3 \chi_{12}^2 z_{12}} \right) + \frac{z_2 \chi_{1i}}{2\chi_{23}} \left( \frac{\chi_{2i}}{\chi_{3i}} + \frac{z_2}{z_3} \right) \\
 & + \frac{z_1 \chi_{2i}^2}{2\chi_{23} \chi_{3i}} + \frac{z_1 \chi_{2i}}{\chi_{23}} \left( \frac{z_2}{2z_3} - \frac{2\chi_{12}}{\chi_{13}} \right) - \frac{\chi_{123}}{\chi_{12}} \left\{ \frac{z_1 \chi_{2i}}{\chi_{12}} \left( \frac{z_3 \chi_{i(12)}}{2z_{12} \chi_{3i}} + \frac{z_{12} \chi_{3i}}{2z_3 \chi_{i(12)}} - 1 \right) \right. \\
 & - \frac{z_2 (z_2 \chi_{1i} \chi_{i(13)} + z_1 \chi_{2i} \chi_{i(23)})}{2z_3 \chi_{23} \chi_{i(12)}} + \frac{z_1 z_2 \chi_{1i} \chi_{2i}}{z_{12} \chi_{23} \chi_{i(12)}} \left( \frac{\chi_{2i}^2}{2\chi_{1i} \chi_{3i}} - \frac{z_{13} \chi_{2i}^2}{2z_2 \chi_{1i} \chi_{3i}} - \frac{z_{13} \chi_{2i}}{2z_2 \chi_{3i}} \right. \\
 & \left. \left. + \frac{\chi_{2i}}{\chi_{1i}} + \frac{\chi_{1i}}{2\chi_{3i}} - \frac{z_{23} \chi_{i(12)}}{2z_1 \chi_{3i}} + \frac{\chi_{2i}}{\chi_{3i}} + \frac{z_2}{z_1} + \frac{z_2^2}{2z_1 z_3} + \frac{z_1}{2z_3} + \frac{z_2}{z_3} + 2 \right) \right\} \\
 & + \frac{D}{4} \left( \frac{\chi_{13} (z_2 \chi_{i(12)} - z_2 \chi_{3i} + (z_{12} - z_3) \chi_{2i})}{\chi_{12} \chi_{23}} + \frac{2z_3 \chi_{12} \chi_{3i}}{\chi_{13} \chi_{23}} - \frac{2z_1 \chi_{3i}}{\chi_{13}} \right. \\
 & \left. - \frac{(z_{12} - 2z_3) \chi_{3i} + z_3 (\chi_{1i} + 3\chi_{2i})}{\chi_{23}} - \frac{z_1 \chi_{1i} + (z_3 - z_2) \chi_{2i} + z_2 \chi_{3i}}{\chi_{12}} \right) \\
 & + \frac{\chi_{1i} \chi_{2i}}{2\chi_{23}} \left[ \frac{z_1 z_2 \chi_{3i}}{z_3 \chi_{1i} \chi_{2i}} - \frac{(z_{12} + 6z_3) \chi_{3i}}{\chi_{1i} \chi_{2i}} + \frac{z_3}{\chi_{3i}} + \frac{2z_{23} + z_3 - 6z_1}{\chi_{1i}} - \frac{4z_2 + z_3}{\chi_{2i}} \right. \\
 & \left. + \frac{z_1}{z_{12}} \left( \frac{z_2}{\chi_{2i}} - \frac{z_2}{\chi_{1i}} \left( \frac{\chi_{3i}}{\chi_{2i}} + 1 \right) - \frac{z_3 z_2}{z_1 \chi_{2i}} \right) \right] + \frac{z_1 (\chi_{1i} - \chi_{2i} + 2\chi_{3i})}{\chi_{13}} \\
 & - \frac{\chi_{12} (z_3 \chi_{i(13)} + z_{23} \chi_{3i})}{\chi_{13} \chi_{23}} + \frac{\chi_{1i} \chi_{2i}}{2\chi_{12}} \left\{ \frac{\chi_{1i}}{\chi_{3i}} \left( \frac{2z_3 - z_1}{\chi_{1i}} + \frac{z_2}{\chi_{2i}} \right) + \frac{\chi_{13} z_{23}}{\chi_{23} \chi_{3i}} \right. \\
 & - \frac{(z_3 z_{12} + (z_1 - z_2)^2) \chi_{123} \chi_{2i}}{z_{12} \chi_{23} \chi_{3i} \chi_{i(12)}} + \frac{z_1}{\chi_{1i}} \left\{ 2 + \frac{z_2 \chi_{13}}{z_3 \chi_{23}} + \frac{z_2}{z_1} \left( \frac{2\chi_{13}}{\chi_{23}} - 1 \right) + \frac{z_3}{z_1} \left( \frac{3\chi_{13}}{\chi_{23}} + 2 \right) \right. \\
 & \left. - \frac{3\chi_{13}}{\chi_{23}} + \frac{\chi_{3i}}{\chi_{2i}} \left[ \frac{z_2}{z_1} \left( \frac{3\chi_{13}}{\chi_{23}} + 2 \right) + \frac{z_2}{z_3} \left( \frac{\chi_{13}}{\chi_{23}} + 2 \right) - \frac{z_2}{z_{12}} \left( \frac{\chi_{12} + 2\chi_{13}}{\chi_{23}} + 3 \right) - \frac{3z_3}{z_1} + 3 \right] \right. \\
 & \left. + \frac{\chi_{2i}}{\chi_{3i}} \left[ \frac{z_1}{z_{12}} \frac{\chi_{1(23)}}{\chi_{23}} \left( 1 - \frac{z_2}{z_1} + \frac{2\chi_{23}}{\chi_{1(23)}} \right) + \frac{\chi_{3(12)}}{\chi_{23}} \right] - \frac{z_2}{z_{12}} \left( \frac{2\chi_{12} + 3\chi_{13}}{\chi_{23}} - \frac{z_1 \chi_{123}}{z_2 \chi_{23}} + \frac{2z_{23}}{z_2} + \frac{2z_3}{z_1} \right) \right\} \\
 & + \frac{z_1 z_2}{z_{12} \chi_{3i}} \left( \frac{z_{23} \chi_{1(23)}}{z_1 \chi_{23}} + \frac{2z_2}{z_1} - \frac{z_{23} \chi_{1(23)}}{z_2 \chi_{23}} + \frac{z_3 (z_2 - z_1)}{z_1 z_2} + \frac{z_1}{z_2} + 1 \right) + \frac{z_2}{\chi_{2i}} \left( \frac{z_{23} \chi_{12}}{z_{12} \chi_{23}} + \frac{z_1}{z_{12}} \right. \\
 & + \frac{z_2}{z_3} \left( \frac{\chi_{13}}{\chi_{23}} + 1 \right) - \frac{z_1}{z_3} - \frac{2\chi_{13}}{\chi_{23}} + \frac{z_3 + 5z_1}{z_2} + 3 \left. \right) + \frac{z_1}{\chi_{i(12)}} \left\{ \frac{z_2 (z_1^2 - z_2^2) \chi_{1i}}{z_1^2 z_3 \chi_{2i}} \right. \\
 & + \frac{z_{12}^2}{z_1 z_3} + \frac{z_2 \chi_{1i} (\chi_{1(23)} - 5\chi_{23})}{z_1 \chi_{23} \chi_{2i}} - \frac{z_2}{z_1} \left( \frac{2\chi_{12} + 3\chi_{13}}{\chi_{23}} + 2 \right) - \frac{z_3}{z_1} \left( \frac{\chi_{12} + 2\chi_{13}}{\chi_{23}} + 3 \right) \\
 & + \frac{z_{12} \chi_{3i}}{z_1 \chi_{2i}} \left( \frac{3z_3}{z_{12}} - \frac{z_2}{z_3} \left( \frac{\chi_{1(23)}}{\chi_{23}} + 1 \right) - 2 \right) + \frac{2z_2}{z_{12}} \left( \frac{z_2}{z_1} \left( \frac{\chi_{1(23)}}{\chi_{23}} - 2 \right) + \frac{2\chi_{1(23)}}{\chi_{23}} - \frac{3z_1}{z_2} - 4 \right) \\
 & + \frac{z_2 \chi_{1i}}{z_1 \chi_{3i}} \left[ \frac{z_2 z_{12} \chi_{3i}}{z_1 z_3 \chi_{2i}} + \frac{z_1}{z_{12}} \left( \frac{\chi_{1(23)}}{\chi_{23}} + 1 \right) - \frac{z_{23}}{z_{12}} \left( \frac{\chi_{1(23)}}{\chi_{23}} + 1 \right) \right] + \frac{\chi_{2i}}{\chi_{1i}} \left[ \frac{2z_2}{z_{12}} \left( \frac{\chi_{1(23)}}{\chi_{23}} - 2 \right) \right. \\
 & + \frac{z_{13} \chi_{2i}}{z_{12} \chi_{3i}} \left( \frac{z_2}{z_{13}} \left( \frac{\chi_{1(23)}}{\chi_{23}} + 1 \right) - \frac{\chi_{1(23)}}{\chi_{23}} - 1 \right) - \frac{6z_1}{z_{12}} + \frac{\chi_{1(23)}}{\chi_{23}} + \frac{z_{123}}{z_3} \\
 & \left. \left. + \frac{\chi_{3i}}{\chi_{2i}} \left( \frac{\chi_{12}}{\chi_{23}} + \frac{3z_3 - 2z_{12}}{z_1} \right) \right] + \frac{\chi_{13}}{\chi_{23}} + 2 \right\} + \frac{z_1 z_2 \chi_{1i} \chi_{2i}}{2\chi_{12}^2} \left\{ \frac{2\chi_{13}}{z_1 \chi_{1i}} \left( \frac{\chi_{3i}}{\chi_{2i}} - 2 \right) \right. \\
 & + \frac{z_2}{z_{12}} \left[ \frac{4\chi_{13} \chi_{i(12)}}{z_1 \chi_{1i} \chi_{2i}} - \frac{2}{\chi_{i(12)}} \left( \frac{2z_{12} \chi_{3(12)} \chi_{1i}}{z_1 z_2 \chi_{2i}} + \frac{\chi_{13} \chi_{i(12)} \chi_{3i}}{z_1 \chi_{1i} \chi_{2i}} \right) + \frac{\chi_{3(12)} (3\chi_{i(12)} - \chi_{3i})}{z_2 \chi_{1i} \chi_{2i}} \right] \\
 & + \frac{2z_{12} \chi_{3(12)} \chi_{1i}}{z_3 z_1 \chi_{2i} \chi_{i(12)}} + \frac{z_3 z_{12}}{z_1 z_2} \left\{ \frac{1}{z_{12}} \left( \frac{2\chi_{13}}{\chi_{1i}} + \frac{\chi_{3(12)}}{\chi_{3i}} \right) + \frac{1}{\chi_{i(12)}} \left[ \frac{1}{z_3} \left( \frac{4\chi_{13} \chi_{2i}}{\chi_{1i}} + \chi_{3(12)} \right) \right. \right.
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{z_{12}\chi_{3(12)}}{z_3^2} - \frac{1}{z_{12}} \left( \frac{2\chi_{13}\chi_{2i}}{\chi_{1i}} + \chi_{3(12)} \right) \Bigg] \Bigg\} + \frac{2\chi_{3(12)}}{z_1 z_2} \left( \frac{z_1 \chi_{2i}}{\chi_{1i} \chi_{3i}} - \frac{z_2 z_{13}}{z_3 \chi_{2i}} + \frac{z_1 z_2 \chi_{3i}}{2z_3 \chi_{1i} \chi_{2i}} \right) \\
 & + \frac{\chi_{123}}{\chi_{i(12)}^2} \left\{ \frac{z_{12} \chi_{1i} \chi_{3i}}{z_1 z_3 \chi_{2i}} - \frac{2\chi_{i(12)}^2}{z_1 \chi_{2i}} + \frac{z_{123} \chi_{3i} \chi_{i(12)}}{z_3 \chi_{1i} \chi_{2i}} \left( \frac{\chi_{i(123)} \chi_{1i}}{z_1 \chi_{3i}} + \frac{\chi_{1i}^2}{z_1 \chi_{3i}} - \frac{\chi_{1i}}{z_1} \right) \right. \\
 & + \frac{z_{123} \chi_{i(12)}}{z_1 z_2} \left( \frac{\chi_{i(12)}}{2\chi_{3i}} - \frac{3z_{12}}{2z_3} + \frac{1}{2\chi_{i(12)} z_{123}} \left( \frac{z_{12}^2 \chi_{3i}}{z_3} + \frac{3z_3 \chi_{1i} \chi_{2i}}{\chi_{3i}} \right) \right) + \frac{\chi_{1i} \chi_{2i}}{z_{12} \chi_{3i}} \left[ \frac{z_3 \chi_{1i}^2}{z_1 \chi_{2i}^2} \right. \\
 & \left. \left. + \frac{z_3 \chi_{1i}}{\chi_{2i}} \left( \frac{1}{z_2} + \frac{3}{z_1} \right) + \frac{\chi_{i(12)}^2}{\chi_{1i} \chi_{2i}} \left( \frac{\chi_{i(123)}}{\chi_{2i}} \left( \frac{z_2 - z_3}{z_1} - 3 \right) + \frac{z_{123} \chi_{i(13)}}{z_1 \chi_{2i}} \right) \right] \right\} \\
 & - \frac{\chi_{123}}{4z_{12} \chi_{i(12)}} \frac{2z_{12} \chi_{2i} (\chi_{1i} (z_{23} - z_1) + z_1 \chi_{3i})}{\chi_{123} \chi_{23}}, \tag{B.2} \\
 c_1 = & \frac{D}{2} + \frac{\chi_{123}^2 z_{123} \chi_{i(123)}}{4\chi_{12} \chi_{3i} z_3} \left[ \frac{2}{\chi_{12}} \left( \frac{\chi_{1i} \chi_{2i}}{\chi_{i(12)}^2} + \frac{z_1 z_2}{z_{12}^2} \right) - \frac{3}{\chi_{i(12)} z_{12}} \frac{\chi_{2i} z_1 + \chi_{1i} z_2}{\chi_{12}} \right] \\
 & - \frac{z_{12}}{\chi_{i(12)}} \left[ \frac{\chi_{3(12)} \chi_{3i}}{2\chi_{12} z_{12}} - \frac{2\chi_{2i} z_1 z_2}{z_{12} \chi_{12}} \left( \frac{(\chi_{13} - \chi_{23})^2}{4z_{12} z_2 \chi_{12}} - \frac{\chi_{13} \chi_{23}}{z_1 z_2 \chi_{12}} + \frac{\chi_{13}^2}{z_1 z_{12} \chi_{12}} - \frac{z_3 \chi_{3(12)}}{2z_1 z_2 z_{12}} \right) \right] \\
 & + \frac{\chi_{12}^2}{\chi_{13} \chi_{23}} - \frac{\chi_{2i} \chi_{12}}{2\chi_{23} \chi_{3i}} - \frac{z_2 \chi_{12}}{2z_3 \chi_{23}} - \frac{z_2 \chi_{1i}}{2z_3 \chi_{3i}} - \frac{\chi_{123}}{\chi_{12}} \left\{ \frac{\chi_{i(12)}}{\chi_{3i}} \left( \frac{z_3}{4z_{12}} - \frac{z_1^2 \chi_{23}}{z_{12}^2 \chi_{12}} \right) \right. \\
 & \left. - \frac{\chi_{1i} \chi_{2i}}{\chi_{i(12)}^2} \left[ \frac{2z_1 \chi_{1i} \chi_{23}}{z_{12} \chi_{2i} \chi_{12}} - \frac{z_{123}}{z_{12}} \left( \frac{\chi_{2i}}{\chi_{1i}} + 1 \right) + \frac{z_1^2 \chi_{23}}{z_{12}^2 \chi_{12}} \left( 2 + \frac{\chi_{2i} \chi_{3(12)}}{\chi_{1i} \chi_{23}} \right) + \frac{z_{12} \chi_{13} \chi_{2i}}{z_3 \chi_{1i} \chi_{12}} \right] \right. \\
 & \left. + \frac{z_2 \chi_{12} \chi_{i(123)}}{2z_3 \chi_{23} \chi_{i(12)}} - \frac{\chi_{1i} \chi_{2i}}{2\chi_{3i} \chi_{i(12)}} - \frac{\chi_{1i} \chi_{2i}}{z_{12} \chi_{3i}} \left[ \frac{z_1 z_2}{z_3 \chi_{i(12)}} \left( \left( \frac{\chi_{2i}}{\chi_{1i}} + 2 \right) + \frac{z_1}{z_2} \right) - \frac{\chi_{12} z_{123}}{2\chi_{1i} \chi_{23}} \right] \right. \\
 & \left. - \frac{z_1 z_2}{2z_3 z_{12}} + \frac{(z_{123} + z_3) \chi_{3i}}{4z_3 \chi_{i(12)}} \right\} + \frac{\chi_{12}}{\chi_{13}} - \left( \frac{\chi_{13}}{\chi_{23}} + \frac{1}{2} \right) \left( \frac{\chi_{2i}}{\chi_{3i}} + \frac{z_2}{z_3} \right) + \frac{3\chi_{12} + 2\chi_{13}}{\chi_{23}} \\
 & - \frac{\chi_{i(12)}}{4\chi_{3i}} - 2 + \frac{z_1}{2z_{12}} \left( 1 - \frac{z_3}{z_1} + \frac{z_2 (\chi_{123} + \chi_{3(12)})}{z_1 \chi_{23}} - \frac{\chi_{1i}}{\chi_{3i}} \right) - \frac{\chi_{123} z_{123}}{2z_{12} \chi_{23}} \left( \frac{\chi_{2i}}{\chi_{3i}} + 1 \right) \\
 & + \frac{\chi_{1i} \chi_{2i}}{2\chi_{12}} \left\{ \frac{\chi_{13}}{\chi_{1i} \chi_{2i}} \left\{ \frac{1}{z_{12}} \left[ z_1 \left( \frac{2\chi_{23}}{\chi_{13}} + 1 \right) + z_2 \left( \frac{\chi_{13}}{\chi_{23}} + 2 \right) - \frac{\chi_{12} z_{23} + 2z_3 \chi_{3(12)}}{\chi_{13}} \right] \right. \right. \\
 & \left. \left. - \frac{z_2}{z_3} \left( \frac{\chi_{13}}{\chi_{23}} + \frac{z_1 \chi_{23}}{z_2 \chi_{13}} + 2 \right) - 4 \right\} - \frac{\chi_{13}}{\chi_{1i} \chi_{3i}} \left( \frac{z_2 \chi_{3(12)} + z_{23} \chi_{123}}{z_{12} \chi_{13}} + \frac{\chi_{13}}{\chi_{23}} \right. \right. \\
 & \left. \left. + \frac{z_1 \chi_{3(12)}}{z_3 \chi_{13}} + 2 \right) + \frac{\chi_{3(12)}}{\chi_{1i} \chi_{i(12)}} \left\{ \frac{\chi_{12}}{\chi_{3(12)}} \left[ \frac{\chi_{13}}{\chi_{12}} \left( \frac{\chi_{13}}{\chi_{23}} + 2 \right) - \frac{z_{12}}{z_3} - 1 \right] \right. \right. \\
 & \left. \left. + \frac{z_{123} + z_2}{z_{12}} - \frac{z_{12} + z_2}{z_3} \right\} + \frac{\chi_{3(12)}}{\chi_{2i} \chi_{i(12)}} \left( \frac{z_{123} + z_1}{z_{12}} + \frac{\chi_{23}}{\chi_{3(12)}} - \frac{z_1}{z_3} + 1 \right) \right. \\
 & \left. - \frac{\chi_{3(12)}}{\chi_{2i} \chi_{3i}} \left( \frac{z_{123}}{z_{12}} + \frac{z_2}{z_3} + \frac{z_{23} \chi_{12}}{z_{12} \chi_{3(12)}} + \frac{\chi_{23}}{\chi_{3(12)}} \right) + \frac{\chi_{123}}{z_{12} z_3} \left\{ \frac{z_1 z_2 (z_{12} - z_3) \chi_{i(123)}}{z_{12} \chi_{1i} \chi_{2i} \chi_{3i}} \right. \right. \\
 & \left. \left. - \frac{z_1 z_2}{\chi_{3i} \chi_{i(12)}} \left[ \frac{z_{12}^2}{z_1 z_2} \left( \frac{\chi_{3i}^2}{\chi_{1i} \chi_{2i}} - 1 \right) + \frac{\chi_{3i} \chi_{i(12)}}{\chi_{1i} \chi_{2i}} + \frac{\chi_{2i}}{\chi_{1i}} + \frac{\chi_{1i}}{\chi_{2i}} - \frac{3z_3^2}{2z_1 z_2} + \frac{z_2}{z_1} + \frac{z_1}{z_2} \right. \right. \\
 & \left. \left. + \frac{z_3 z_{12} \chi_{3i}^2}{z_1 z_2 \chi_{1i} \chi_{2i}} + 4 \right] + \frac{z_{123} \chi_{3i}}{\chi_{i(12)}^2} \left[ \frac{z_3^2 \chi_{1i}}{z_{123} \chi_{3i}^2} \left( \frac{\chi_{1i}}{2\chi_{2i}} + \frac{\chi_{2i}^2}{2\chi_{1i}^2} \right) + \frac{z_{12}^2}{z_{123} \chi_{3i}} \left( \frac{\chi_{i(12)} \chi_{3i}}{2\chi_{1i} \chi_{2i}} - 1 \right) \right. \right. \\
 & \left. \left. + \frac{z_{12} z_3}{z_{123} \chi_{3i}} \left( \frac{\chi_{i(12)} \chi_{3i}}{\chi_{1i} \chi_{2i}} - 1 \right) + \frac{z_3}{\chi_{3i}} \left( \frac{\chi_{1i}}{\chi_{2i}} + \frac{\chi_{2i}}{\chi_{1i}} + 2 \right) \right] \right\} \\
 & + \frac{z_1 z_2 \chi_{1i} \chi_{2i}}{2\chi_{12}^2} \left\{ \frac{\chi_{3(12)}}{z_1 z_2} \left[ \left( \frac{\chi_{13}}{\chi_{1i}} + \frac{\chi_{23}}{\chi_{2i}} \right) \left( \frac{1}{\chi_{i(12)}} - \frac{1}{\chi_{3i}} \right) - \frac{4\chi_{13} \chi_{23}}{\chi_{3(12)} \chi_{1i} \chi_{2i}} \right] \right\}
 \end{aligned}$$

$$\begin{aligned}
& + \frac{2\chi_{13}\chi_{3(12)}}{z_1\chi_{1i}\chi_{2i}} \left( \frac{1}{z_{12}} - \frac{1}{z_3} \right) + \frac{\chi_{3(12)}^2}{z_1\chi_{2i}} \left( \frac{1}{z_{12}\chi_{3i}} - \frac{1}{z_3\chi_{3i}} + \frac{1}{z_3\chi_{i(12)}} \right) \\
& + \frac{2(\chi_{13} - \chi_{23})}{z_{12}\chi_{2i}\chi_{i(12)}} \left( \frac{\chi_{23} - \chi_{13}}{z_1} + \frac{2\chi_{23}}{z_2} \right) + \frac{\chi_{123}}{\chi_{i(12)}^2} \left\{ \frac{2\chi_{13}}{z_{12}z_3} \left[ \frac{\chi_{i(12)}^2\chi_{i(123)}}{\chi_{1i}\chi_{2i}\chi_{3i}} \left( \frac{z_3}{z_1} + \frac{\chi_{3(12)}}{2\chi_{13}} \right) \right. \right. \\
& - \left. \left. \frac{2z_3\chi_{2i}}{z_1\chi_{1i}} - \frac{3\chi_{2i}}{\chi_{1i}} \right] - \frac{\chi_{3(12)}}{z_{12}^2} \left[ \frac{2z_1\chi_{2i}}{z_2\chi_{1i}} + \frac{\chi_{i(12)}^2\chi_{i(123)}}{\chi_{1i}\chi_{2i}\chi_{3i}} + \frac{6z_1\chi_{23}\chi_{i(12)}}{z_2\chi_{3(12)}\chi_{3i}} \right. \right. \\
& + \left. \left. \frac{2\chi_{23}}{\chi_{3(12)}} \left( \frac{z_1\chi_{1i}}{z_2\chi_{3i}} \left( \frac{\chi_{1i}}{\chi_{2i}} + \frac{\chi_{2i}^2}{\chi_{1i}^2} \right) + \frac{z_1^2\chi_{1i}}{z_2z_3\chi_{2i}} + \frac{z_2^2\chi_{1i}}{z_3z_1\chi_{2i}} + \frac{2z_1}{z_2} \right) \right] \right\} \\
& + \left. \frac{z_{123}\chi_{3(12)}}{z_1z_2z_3} \frac{\chi_{i(12)}}{\chi_{3i}} \left[ \frac{\chi_{3i}}{\chi_{3(12)}} \left( \frac{\chi_{13}}{\chi_{1i}} + \frac{\chi_{23}}{\chi_{2i}} \right) - \frac{\chi_{3i}}{\chi_{i(12)}} + 1 \right] \right\} \\
& + \frac{\chi_{123}}{4z_{12}\chi_{i(12)}} \left[ \frac{\chi_{1i}\chi_{2i}}{\chi_{123}} \left( \frac{30z_1 + 28z_2}{\chi_{2i}} + \frac{2z_{12}(\chi_{12} + 2\chi_{13})}{\chi_{1i}\chi_{23}} - \frac{2z_{12}\chi_{3i}}{\chi_{1i}\chi_{2i}} - \frac{2(2z_1 + z_2)z_{12}}{z_3\chi_{2i}} \right) \right. \\
& \left. + \frac{2z_{123}\chi_{2i}\chi_{i(12)}}{\chi_{23}\chi_{3i}} - \frac{2z_{12}\chi_{i(123)}}{\chi_{23}} \right], \tag{B.3}
\end{aligned}$$

$$\begin{aligned}
c_2 = & \frac{\chi_{12}}{4z_3\chi_{3i}} + \frac{\chi_{3(12)}^2}{4\chi_{12}z_{12}\chi_{i(12)}} \left( \frac{3\chi_{123}^2z_{123}\chi_{i(123)}}{z_3\chi_{3i}\chi_{3(12)}^2} + 1 \right) + \frac{\chi_{12}}{2z_{12}\chi_{3i}} \left[ \frac{\chi_{3(12)}}{\chi_{12}} \left( \frac{z_{12}}{z_3} - 1 \right) - \frac{1}{2} \right] \\
& + \frac{\chi_{1i}\chi_{2i}}{2\chi_{12}} \left\{ \frac{\chi_{3(12)}^2}{2z_{12}\chi_{1i}\chi_{2i}\chi_{i(12)}} \left[ \frac{z_{12}}{z_3} \left( \frac{\chi_{i(12)}}{\chi_{3i}} - 1 \right) - \frac{\chi_{i(12)}}{\chi_{3i}} \right] \right\} \\
& + \frac{\chi_{123}}{4z_{12}\chi_{i(12)}} \left( \frac{\chi_{3(12)}}{\chi_{123}} - \frac{z_{12}(\chi_{12} + 2\chi_{3(12)})}{z_3\chi_{123}} + 1 \right). \tag{B.4}
\end{aligned}$$

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP<sup>3</sup> supports the goals of the International Year of Basic Sciences for Sustainable Development.

## References

- [1] J.M. Campbell and E.W.N. Glover, *Double unresolved approximations to multiparton scattering amplitudes*, *Nucl. Phys. B* **527** (1998) 264 [[hep-ph/9710255](#)] [[INSPIRE](#)].
- [2] S. Catani and M. Grazzini, *Collinear factorization and splitting functions for next-to-next-to-leading order QCD calculations*, *Phys. Lett. B* **446** (1999) 143 [[hep-ph/9810389](#)] [[INSPIRE](#)].
- [3] S. Catani and M. Grazzini, *Infrared factorization of tree level QCD amplitudes at the next-to-next-to-leading order and beyond*, *Nucl. Phys. B* **570** (2000) 287 [[hep-ph/9908523](#)] [[INSPIRE](#)].
- [4] V. Del Duca, A. Frizzo and F. Maltoni, *Factorization of tree QCD amplitudes in the high-energy limit and in the collinear limit*, *Nucl. Phys. B* **568** (2000) 211 [[hep-ph/9909464](#)] [[INSPIRE](#)].
- [5] D.A. Kosower, *Multiple singular emission in gauge theories*, *Phys. Rev. D* **67** (2003) 116003 [[hep-ph/0212097](#)] [[INSPIRE](#)].
- [6] M. Czakon, *Double-real radiation in hadronic top quark pair production as a proof of a certain concept*, *Nucl. Phys. B* **849** (2011) 250 [[arXiv:1101.0642](#)] [[INSPIRE](#)].

- [7] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One loop  $n$  point gauge theory amplitudes, unitarity and collinear limits*, *Nucl. Phys. B* **425** (1994) 217 [[hep-ph/9403226](#)] [[INSPIRE](#)].
- [8] Z. Bern, V. Del Duca and C.R. Schmidt, *The Infrared behavior of one loop gluon amplitudes at next-to-next-to-leading order*, *Phys. Lett. B* **445** (1998) 168 [[hep-ph/9810409](#)] [[INSPIRE](#)].
- [9] D.A. Kosower and P. Uwer, *One loop splitting amplitudes in gauge theory*, *Nucl. Phys. B* **563** (1999) 477 [[hep-ph/9903515](#)] [[INSPIRE](#)].
- [10] Z. Bern, V. Del Duca, W.B. Kilgore and C.R. Schmidt, *The infrared behavior of one loop QCD amplitudes at next-to-next-to leading order*, *Phys. Rev. D* **60** (1999) 116001 [[hep-ph/9903516](#)] [[INSPIRE](#)].
- [11] S. Catani and M. Grazzini, *The soft gluon current at one loop order*, *Nucl. Phys. B* **591** (2000) 435 [[hep-ph/0007142](#)] [[INSPIRE](#)].
- [12] O. Braun-White and N. Glover, *Decomposition of triple collinear splitting functions*, *JHEP* **09** (2022) 059 [[arXiv:2204.10755](#)] [[INSPIRE](#)].
- [13] F. Caola, A. Chakraborty, G. Gambuti, A. von Manteuffel and L. Tancredi, *Three-loop helicity amplitudes for four-quark scattering in massless QCD*, *JHEP* **10** (2021) 206 [[arXiv:2108.00055](#)] [[INSPIRE](#)].
- [14] P. Bargiela, F. Caola, A. von Manteuffel and L. Tancredi, *Three-loop helicity amplitudes for diphoton production in gluon fusion*, *JHEP* **02** (2022) 153 [[arXiv:2111.13595](#)] [[INSPIRE](#)].
- [15] F. Caola, A. Chakraborty, G. Gambuti, A. von Manteuffel and L. Tancredi, *Three-Loop Gluon Scattering in QCD and the Gluon Regge Trajectory*, *Phys. Rev. Lett.* **128** (2022) 212001 [[arXiv:2112.11097](#)] [[INSPIRE](#)].
- [16] S. Abreu, F. Febres Cordero, H. Ita, B. Page and V. Sotnikov, *Leading-color two-loop QCD corrections for three-jet production at hadron colliders*, *JHEP* **07** (2021) 095 [[arXiv:2102.13609](#)] [[INSPIRE](#)].
- [17] S. Abreu, F. Febres Cordero, H. Ita, M. Klinkert, B. Page and V. Sotnikov, *Leading-color two-loop amplitudes for four partons and a  $W$  boson in QCD*, *JHEP* **04** (2022) 042 [[arXiv:2110.07541](#)] [[INSPIRE](#)].
- [18] S. Badger, H.B. Hartanto and S. Zoia, *Two-Loop QCD Corrections to  $Wb\bar{b}$  Production at Hadron Colliders*, *Phys. Rev. Lett.* **127** (2021) 012001 [[arXiv:2102.02516](#)] [[INSPIRE](#)].
- [19] S. Badger, H.B. Hartanto, J. Kryś and S. Zoia, *Two-loop leading-colour QCD helicity amplitudes for Higgs boson production in association with a bottom-quark pair at the LHC*, *JHEP* **11** (2021) 012 [[arXiv:2107.14733](#)] [[INSPIRE](#)].
- [20] S. Badger, H.B. Hartanto, J. Kryś and S. Zoia, *Two-loop leading colour helicity amplitudes for  $W^\pm\gamma + j$  production at the LHC*, *JHEP* **05** (2022) 035 [[arXiv:2201.04075](#)] [[INSPIRE](#)].
- [21] F.A. Dreyer and A. Karlberg, *Vector-Boson Fusion Higgs Production at Three Loops in QCD*, *Phys. Rev. Lett.* **117** (2016) 072001 [[arXiv:1606.00840](#)] [[INSPIRE](#)].
- [22] F.A. Dreyer and A. Karlberg, *Vector-Boson Fusion Higgs Pair Production at  $N^3$ LO*, *Phys. Rev. D* **98** (2018) 114016 [[arXiv:1811.07906](#)] [[INSPIRE](#)].
- [23] F. Dulat, B. Mistlberger and A. Pelloni, *Precision predictions at  $N^3$ LO for the Higgs boson rapidity distribution at the LHC*, *Phys. Rev. D* **99** (2019) 034004 [[arXiv:1810.09462](#)] [[INSPIRE](#)].

- [24] L.-B. Chen, H.T. Li, H.-S. Shao and J. Wang, *Higgs boson pair production via gluon fusion at  $N^3LO$  in QCD*, *Phys. Lett. B* **803** (2020) 135292 [[arXiv:1909.06808](#)] [[INSPIRE](#)].
- [25] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss, B. Mistlberger and A. Pelloni, *Fully Differential Higgs Boson Production to Third Order in QCD*, *Phys. Rev. Lett.* **127** (2021) 072002 [[arXiv:2102.07607](#)] [[INSPIRE](#)].
- [26] X. Chen, T. Gehrmann, N. Glover, A. Huss, T.-Z. Yang and H.X. Zhu, *Dilepton Rapidity Distribution in Drell-Yan Production to Third Order in QCD*, *Phys. Rev. Lett.* **128** (2022) 052001 [[arXiv:2107.09085](#)] [[INSPIRE](#)].
- [27] X. Chen et al., *Third-Order Fiducial Predictions for Drell-Yan Production at the LHC*, *Phys. Rev. Lett.* **128** (2022) 252001 [[arXiv:2203.01565](#)] [[INSPIRE](#)].
- [28] V. Del Duca, C. Duhr, R. Haindl, A. Lazopoulos and M. Michel, *Tree-level splitting amplitudes for a quark into four collinear partons*, *JHEP* **02** (2020) 189 [[arXiv:1912.06425](#)] [[INSPIRE](#)].
- [29] V. Del Duca, C. Duhr, R. Haindl, A. Lazopoulos and M. Michel, *Tree-level splitting amplitudes for a gluon into four collinear partons*, *JHEP* **10** (2020) 093 [[arXiv:2007.05345](#)] [[INSPIRE](#)].
- [30] S. Catani, D. de Florian and G. Rodrigo, *The Triple collinear limit of one loop QCD amplitudes*, *Phys. Lett. B* **586** (2004) 323 [[hep-ph/0312067](#)] [[INSPIRE](#)].
- [31] S. Badger, F. Buciuani and T. Peraro, *One-loop triple collinear splitting amplitudes in QCD*, *JHEP* **09** (2015) 188 [[arXiv:1507.05070](#)] [[INSPIRE](#)].
- [32] M. Czakon and S. Sapeta, *Complete collection of one-loop triple-collinear splitting operators for dimensionally-regulated QCD*, *JHEP* **07** (2022) 052 [[arXiv:2204.11801](#)] [[INSPIRE](#)].
- [33] C. Duhr, T. Gehrmann and M. Jaquier, *Two-loop splitting amplitudes and the single-real contribution to inclusive Higgs production at  $N^3LO$* , *JHEP* **02** (2015) 077 [[arXiv:1411.3587](#)] [[INSPIRE](#)].
- [34] Y. Li and H.X. Zhu, *Single soft gluon emission at two loops*, *JHEP* **11** (2013) 080 [[arXiv:1309.4391](#)] [[INSPIRE](#)].
- [35] C. Duhr and T. Gehrmann, *The two-loop soft current in dimensional regularization*, *Phys. Lett. B* **727** (2013) 452 [[arXiv:1309.4393](#)] [[INSPIRE](#)].
- [36] L.J. Dixon, E. Herrmann, K. Yan and H.X. Zhu, *Soft gluon emission at two loops in full color*, *JHEP* **05** (2020) 135 [[arXiv:1912.09370](#)] [[INSPIRE](#)].
- [37] Y.J. Zhu, *Double soft current at one-loop in QCD*, [arXiv:2009.08919](#) [[INSPIRE](#)].
- [38] S. Catani and L. Cieri, *Multiple soft radiation at one-loop order and the emission of a soft quark-antiquark pair*, *Eur. Phys. J. C* **82** (2022) 97 [[arXiv:2108.13309](#)] [[INSPIRE](#)].
- [39] S. Catani, D. Colferai and A. Torricelli, *Triple (and quadruple) soft-gluon radiation in QCD hard scattering*, *JHEP* **01** (2020) 118 [[arXiv:1908.01616](#)] [[INSPIRE](#)].
- [40] S. Catani and M.H. Seymour, *The Dipole formalism for the calculation of QCD jet cross-sections at next-to-leading order*, *Phys. Lett. B* **378** (1996) 287 [[hep-ph/9602277](#)] [[INSPIRE](#)].
- [41] S. Catani and M.H. Seymour, *A General algorithm for calculating jet cross-sections in NLO QCD*, *Nucl. Phys. B* **485** (1997) 291 [*Erratum ibid.* **510** (1998) 503] [[hep-ph/9605323](#)] [[INSPIRE](#)].

- [42] A. Bassetto, M. Ciafaloni and G. Marchesini, *Jet Structure and Infrared Sensitive Quantities in Perturbative QCD*, *Phys. Rept.* **100** (1983) 201 [[INSPIRE](#)].
- [43] S. Catani, *New Techniques for Calculating Higher-Order QCD Corrections*, in *Proceedings, Workshop on New Techniques for Calculating Higher Order QCD Corrections*, Zurich, Switzerland, 1992.
- [44] S. Catani, D. de Florian and G. Rodrigo, *Space-like (versus time-like) collinear limits in QCD: Is factorization violated?*, *JHEP* **07** (2012) 026 [[arXiv:1112.4405](#)] [[INSPIRE](#)].
- [45] D. Amati, R. Petronzio and G. Veneziano, *Relating Hard QCD Processes Through Universality of Mass Singularities. 2*, *Nucl. Phys. B* **146** (1978) 29 [[INSPIRE](#)].
- [46] D. Amati, R. Petronzio and G. Veneziano, *Relating Hard QCD Processes Through Universality of Mass Singularities*, *Nucl. Phys. B* **140** (1978) 54 [[INSPIRE](#)].
- [47] R.K. Ellis, H. Georgi, M. Machacek, H.D. Politzer and G.G. Ross, *Factorization and the Parton Model in QCD*, *Phys. Lett. B* **78** (1978) 281 [[INSPIRE](#)].
- [48] G. Somogyi, Z. Trócsányi and V. Del Duca, *Matching of singly- and doubly-unresolved limits of tree-level QCD squared matrix elements*, *JHEP* **06** (2005) 024 [[hep-ph/0502226](#)] [[INSPIRE](#)].