

RESEARCH PAPER

Troglitazone induces cytotoxicity in part by promoting the degradation of peroxisome proliferator-activated receptor γ co-activator-1 α protein

Correspondence

Chi-Wai Wong, Guangzhou Institute of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou Science City, Guangzhou 510663, China.
E-mail: wongcw123456@yahoo.com

Keywords

troglitazone; liver toxicity; mitochondrial mass; PGC-1 α

Received

13 January 2010

Revised

3 March 2010

Accepted

4 May 2010

Xuemei Liao, Yanfei Wang and Chi-Wai Wong

Guangzhou Institute of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China

BACKGROUND AND PURPOSE

Troglitazone (Tro), rosiglitazone (Rosi) and pioglitazone (Pio) are anti-diabetic thiazolidinediones that function as ligands for peroxisome proliferator-activated receptor γ (PPAR γ); however, Tro has been withdrawn from the market due to liver toxicity issues. Mitochondrial dysfunction induced by Tro has been suggested to be an important mechanism behind its cytotoxicity. Constitutively active nuclear hormone receptors, oestrogen-related receptor α and γ are thought to regulate mitochondrial mass and oxidative phosphorylation together with their co-activators PPAR γ co-activator-1 α and -1 β (PGC-1 α and PGC-1 β). Hence, in this study, we investigated whether Tro affects the expression and activity levels of these regulators.

EXPERIMENTAL APPROACH

Cellular viability was measured by an ATP-based assay. Mitochondrial mass and reactive oxygen species (ROS) were quantified by two different fluorogenic probes. Apoptosis was measured by an Annexin-V-based kit. Gene expression at the levels of mRNA and protein was measured by quantitative RT-PCR and Western analysis. Over-expression of PGC-1 α was mediated by an adenovirus.

KEY RESULTS

Tro, but not Rosi or Pio, selectively stimulated PGC-1 α protein degradation. As a result, Tro reduced mitochondrial mass, and superoxide dismutases 1 and 2 expressions, but induced ROS to initiate apoptosis. Using a ubiquitin-proteasome inhibitor MG132, it was established that blocking PGC-1 α degradation partially suppressed the reduction of mitochondrial mass. Importantly, over-expressing PGC-1 α partially restored the Tro-suppressed mitochondrial mass and attenuated the cytotoxic effects of Tro.

CONCLUSIONS AND IMPLICATIONS

Collectively, these results suggest that PGC-1 α degradation is an important mechanism behind the cytotoxic effects of Tro in the liver.

Abbreviations

$\Delta\psi_m$, mitochondrial membrane potential; ATP5b, ATP synthase; cyt-C, cytochrome C oxidase; ERR γ , oestrogen-related receptor γ ; PGC-1 α , peroxisome proliferator-activated receptor γ co-activator-1 α ; FBS, fetal bovine serum; JNK, c-Jun N-terminal protein kinase; PBS, phosphate-buffered saline; PGC-1 β , peroxisome proliferator-activated receptor γ co-activator-1 β ; Pio, pioglitazone; PPAR γ , peroxisome proliferator-activated receptor γ ; ERR α , oestrogen-related receptor α ; PPRE, peroxisome proliferator-activated receptor response element; ROS, reactive oxygen species; Rosi, rosiglitazone; SOD1, superoxide dismutase 1; SOD2, superoxide dismutase 2; Tfam, mitochondrial transcription factor A; Tro, troglitazone; TZD, thiazolidinedione

Introduction

Troglitazone (Tro), rosiglitazone (Rosi) and pioglitazone (Pio) are anti-diabetic thiazolidinediones (TZDs) that function as ligands for peroxisome proliferator-activated receptor γ (PPAR γ) (Desvergne and Wahli, 1999). TZDs bind to and enhance PPAR γ activity through its C-terminally located ligand binding domain (Lehmann *et al.*, 1995); thus, TZDs are thought to mediate their anti-diabetic effects by modulating the expression of PPAR γ target genes involved in regulating metabolism (Staels and Fruchart, 2005).

Despite their demonstrated efficacies in reducing blood glucose level and enhancing insulin sensitivity in patients, the risk-to-benefit ratios of this class of compounds are still hotly debated due to a range of side effects induced by TZDs (Scheen, 2004). All three TZDs induce oedema in patients and increase their body weights (Niemeyer and Janney, 2002; Tang *et al.*, 2003); Rosi and Pio are associated with increased risks of heart dysfunction (Home *et al.*, 2007; Lincoff *et al.*, 2007), and Tro has been withdrawn from the market due to liver toxicity issues (Watkins and Whitcomb, 1998). PPAR γ activation in the kidney collecting duct appears to be responsible for the oedema (Zhang *et al.*, 2005). However, although all three TZDs can activate PPAR γ in the liver, only Tro induced idiosyncratic hepatotoxicity, suggesting that activation of PPAR γ may not be the only mechanism responsible for Tro-induced liver toxicity.

Mitochondrial dysfunction induced by Tro has been suggested to be an important aspect behind hepatotoxicity (Masubuchi, 2006). Tro was found to decrease mitochondrial membrane potential ($\Delta\psi_m$), and increase permeability in both rat and human isolated hepatocytes, whereas Rosi had a minimal effect (Haskins *et al.*, 2001). The decrease in $\Delta\psi_m$ may be the result of inhibition of mitochondrial oxidative phosphorylation induced by Tro (Nadanaciva *et al.*, 2007). This defect is associated with the generation of reactive oxygen species (ROS) (Shishido *et al.*, 2003). As ROS is known to affect stress-activated protein kinases, Tro was demonstrated to activate apoptosis signal-regulating kinase 1 in human immortalized hepatocytes HC-04 (Lim *et al.*, 2008). In addition, Tro activated c-Jun N-terminal protein kinase (JNK), and a JNK inhibitor partially suppressed the Tro-mediated cell death in human HepG2 hepatocarcinoma cells (Bae and Song, 2003). Importantly, Tro caused liver injury in heterozygous superoxide dismutase 2 (*Sod2*[±]) mice, suggesting that an effect on the anti-oxidative capacity of mitochondria constitutes part of the mechanism behind its hepatotoxicity *in vivo* (Ong *et al.*, 2007).

The constitutively active nuclear hormone receptors, oestrogen-related receptor α and γ (ERR α and ERR γ), together with their co-activators, PPAR γ co-activator-1 α and -1 β (PGC-1 α and PGC-1 β), are thought to regulate mitochondrial mass and oxidative phosphorylation (Giguere, 2008). Because these factors govern many important aspects of mitochondrial function (Wu *et al.*, 1999; Kamei *et al.*, 2003), we hypothesized that Tro may affect the expression or activity of ERR α , ERR γ , PGC-1 α and PGC-1 β to induce cytotoxicity. In this study, we found that Tro, but not Rosi or Pio, selectively stimulated the degradation of PGC-1 α protein, leading to reductions in mitochondrial mass and the expression of superoxide dismutase 1 (SOD1) and SOD2, and the induction of ROS that then initiate apoptosis.

Methods

Cell culture and viability assay

Human hepatocarcinoma HepG2 cells were purchased from American Type Culture Collection and grown in MEM medium (Gibco, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS) (HyClone, Logan, UT, USA) and 1 \times penicillin/streptomycin (Gibco), and maintained at 37°C and 5% CO₂. For cell viability assay, cells were seeded in 96-well plates at a density of 30 000 cells per well with or without compounds added. After treatment of indicated time, viability was measured by using CellTiter-Glo Kit (Promega, Madison, WI, USA) and recorded by a VERITAS Microplate luminometer (Turner Biosystems, Sunnyvale, CA, USA).

Transfections

Transient transfections were performed using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. For luciferase reporter assays, cells at 85–95% confluency in 96-well plates were co-transfected with reporter plasmids (25 ng per well), *Renilla* luciferase (3 ng per well) as an internal control for transfection efficiency and the appropriate mammalian expression vectors (5 ng per well). Six hours after transfection, the cells were treated with drugs for 24 h. Luciferase activity was measured by using a VERITAS Microplate luminometer following the Dual-Glo Luciferase Assay System (Promega) technical manual.

Quantitative real-time PCR

Total RNA extraction, first-strand cDNA generation and quantitative real-time PCR analysis were performed as described previously (Nie and Wong, 2009). Relative gene expression was normalized to

β -actin levels. Sequences of primers used are listed in Supporting Information Table S1.

Western blot analysis

Cells were lysed using RIPA reagent (Shanghai Shenneng, Shanghai, China) according to the manufacturer's protocol, and protein extracts were analysed by 10% SDS-PAGE and blotted onto PVDF membrane. Membranes were incubated with rabbit anti-ERR α , -ERR γ , -PGC-1 α and -PGC-1 β antibodies all generated in-house, or mouse anti- β -actin antibody (Boster, Wuhan, China) followed by horseradish peroxidase-conjugated secondary antibody (Amersham, Buckinghamshire, UK) and developed with ECL reagent (Amersham).

Mitochondrial mass assay

Mitotracker green (Invitrogen) was added and quantified as described by Chen and Wong (2009). Briefly, cells were incubated in serum-free medium (pre-warmed to 37°C) with 150 nM Mitotracker Green FM for 20 min in the dark. After being stained, the cells were washed twice with cold phosphate-buffered saline (PBS) and suspended in 200 μ L PBS. Subsequently, the cells were analysed on a flow cytometer (FAC-SCalibur, BD Biosciences, San Jose, CA, USA) with excitation at 490 nm and emission at 516 nm. Data were processed by using the CellQuest program (BD Biosciences).

ROS assay

ROS production in mitochondria was measured by a cell-permeable fluorogenic probe MitoSOX Red (Invitrogen) that is selectively targeted to the mitochondria where it specifically reacts with superoxide anion. Drug-treated cells were loaded with 5 μ M MitoSOX Red for 10 min at 37°C, washed with PBS, and the fluorescence was detected with a flow cytometer (FAC-SCalibur, BD Biosciences) with excitation set at 510 nm and emission at 580 nm. Data were processed by using a CellQuest program (BD Biosciences).

Assessment of apoptosis

Annexin V-PE apoptosis detection kit I (BD Biosciences) was used for quantification as described previously (Wu *et al.*, 2009). Briefly, cells were washed twice with cold PBS and resuspended in 1 \times binding buffer at a concentration of 1 \times 10⁶ cells mL⁻¹, then 100 μ L of the solution (1 \times 10⁵ cells) was transferred to a 5 mL culture tube, and 5 μ L of Annexin V-PE and 5 μ L of 7-amino-actinomycin D were added. Subsequently, the cells were gently mixed and incubated for 15 min at room temperature (25°C) in the dark. Finally, the cells were suspended in 400 μ L PBS for analysis using a flow cytometer (FAC-

SCalibur, BD Biosciences), and data were processed by using the CellQuest program (BD Biosciences).

Adenovirus over-expressing human PGC-1 α

A recombinant adenovirus encoding the human PGC-1 α was prepared by transfecting a linearized pAd5vector containing 2.4 kb human PGC-1 α cDNA into Trex-293 cells. Purified virus at a multiplicity of 50 was used to infect HepG2 cells in media with 2.5% FBS. For mitochondrial mass assays, 2 \times 10⁵ HepG2 cells per well were plated in 24-well plates and incubated overnight before infection with the adenovirus. For cell viability assays, 2 \times 10⁴ HepG2 cells per well in 96-well plates were used. For mRNA collection, 4 \times 10⁵ HepG2 cells per well in six-well plates were used instead. Twenty-four hours after infection, cells were treated with dimethyl sulphoxide or 25 μ M Tro for another 24 h. Adenovirus without any additions was used as a control.

Statistical analysis

All assays were performed at least three independent times. Data are presented as mean \pm SD and analysed by Student's *t*-test or ANOVA (Figures 2C,D, 3A,D and 5). Asterisks indicate significant differences: **P* < 0.05, ***P* < 0.01, ****P* < 0.001.

Plasmids and chemicals

A pGL3-luciferase reporter (Promega) with the peroxisome proliferator-activated receptor response element (PPRE) from fatty acid-binding protein 4 promoter cloned in was kindly provided by Dr Brian Lavan. Expression vectors for PPAR γ 1 and PPAR γ 2 were cloned from cDNA generated from THP-1 macrophages. An expression vector for PGC-1 α has been described previously (Wang *et al.*, 2009). Tro and GW9662 were purchased from Cayman Chemical (Ann Arbor, MI, USA), while other chemical reagents were obtained from Sigma (St. Louis, MO, USA). Drug/molecular target nomenclature conforms to the *British Journal of Pharmacology's Guide to Receptors and Channels* (Alexander *et al.*, 2008).

Results

Specific effects of Tro on cell viability and apoptosis

Tro, Rosi and Pio share a common methoxy-phenylmethyl-1,3-thiazolidine-2,4-dione backbone structure with different substitutions (Figure 1A). All three glitazones enhanced the expression of a reporter luciferase driven by a PPRE in human hepatocarcinoma HepG2 cells, more specifically increasing the co-expression of the PPAR γ 1 isoform (Figure 1B). Consistent with the reported EC₅₀s of




Figure 1

Structure–activity relationships of glitazones with PPAR γ . (A) Chemical structures of Tro, Rosi and Pio. (B) HepG2 cells were transfected with expression and reporter plasmids as described in Methods. Relative activity was calculated and expressed as fold induction over that induced by dimethyl sulphoxide (vehicle control). Columns represent mean and vertical lines SD.

these glitazones, Rosi and Pio activated PPAR γ 1 at concentrations as low as 0.2 μ M, while Tro required higher concentrations to fully activate PPAR γ 1. Similar results were observed with the PPAR γ 2 isoform (Supporting Information Figure S1).

On the other hand, after the amount of FBS used for incubation had been reduced, only Tro dose-dependently reduced cell viability; Rosi and Pio did not significantly affect cell viability (Figure 2A–D). The reduction in cell viability mediated by Tro increased with time, whereas with Rosi and Pio viability was still preserved after a longer exposure time (Supporting Information Figure S2). In addition, this Tro-mediated reduction in cell viability was not blocked by the addition of a PPAR γ antagonist GW9662 (Figure 2E). Also, Tro, but not Rosi and Pio, increased apoptosis (Figure 2F). Collectively,

these data suggest that activation of PPAR γ is not involved in Tro-induced cytotoxicity.

Tro elevates mitochondrial ROS level and suppresses mitochondrial mass

Tro-induced hepatotoxicity has been linked to an increase in ROS (Shishido *et al.*, 2003). Hence, we investigated whether Tro-induced ROS is derived from the mitochondria. Using a fluorescence dye MitoSOX Red, which selectively targets the mitochondria and reacts with superoxide anion, we found that Tro dose-dependently elevated mitochondrial ROS level, but Rosi and Pio did not (Figure 3A). Next, it was found that the expression levels of SOD1 and SOD2, which are responsible for neutralizing superoxide anion, were specifically reduced by Tro (Figure 3B–C).




Figure 2

Differential effects of glitazones on cell viability and apoptosis. (A–D) HepG2 cells cultured in (A) 10%, (B) 5%, (C) 2.5% and (D) 0.5% FBS were treated with different doses of Tro, Rosi and Pio for 24 h before cell viability measurement. (E) HepG2 cells in 2.5% FBS were pretreated with 1 μ M of the PPAR γ antagonist GW9662 for 2 h before Tro addition. The relative viabilities were measured and expressed as in (A). (F) The % of apoptotic cells were measured after 24 h drug treatments in HepG2 cells cultured in 2.5% FBS. (A–E) The relative level under dimethyl sulphoxide treatment was set as 1. * P < 0.05, ** P < 0.01, *** P < 0.001.

Because Tro reduces mitochondrial antioxidative capacity, we next determined whether Tro also affects mitochondrial mass by using another fluorescence dye MitoTracker Green that stains the mitochondria independent of $\Delta\psi_m$. We found that Tro, but not Rosi or Pio, dose-dependently reduced mitochondrial mass (Figure 3D). Correspondingly, the expression levels of mitochondrial transcription factor A (Tfam), cytochrome C oxidase (cyt-C) and ATP synthase (ATP5b) that are involved in regulating mitochondrial mass, oxidative phosphorylation and ATP production, respectively, were specifically reduced by Tro (Figure 3E,F; Supporting Information Figure S3).

Tro selectively reduces PGC-1 α protein level

Tro suppressed the mRNA expression levels of SOD1, SOD2, Tfam, cyt-C and ATP5b; ERR α , ERR γ , PGC-1 α and PGC-1 β are responsible for governing the expression of these genes at the transcriptional level and controlling mitochondrial mass (Giguere, 2008). Hence, we next determined whether Tro affects the expression or activity levels of these mitochondrial regulators. We found that both Rosi and Pio did not significantly affect the mRNA expression levels of ERR α , ERR γ , PGC-1 α and PGC-1 β (Figure 4A–D), whereas Tro moderately increased the mRNA expression levels of ERR α , ERR γ and




Figure 3

Effects of Tro on mitochondrial ROS and mass. (A) The levels of mitochondrial ROS (B–C) mRNA expression levels (B, SOD1 and C, SOD2) and (D) the amounts of mitochondrial mass were measured after the drug treatments. (E–F) mRNA expression levels of (E) ATP5b and (F) Tfam were measured as in (B). (A–F) Drug treatments for 24 h were performed in HepG2 cells cultured in 2.5% FBS. The relative level under DMSO treatment was set as 1. $**P < 0.01$, $***P < 0.001$.

PGC-1 α (Figure 4A–C). In contrast, Western blot analysis revealed that of these three glitazones tested, only Tro specifically reduced the protein level of PGC-1 α , but not ERR α , ERR γ and PGC-1 β (Figure 4E), by strongly indicating that down-regulation of PGC-1 α expression at the protein level is correlated to the reductions in mitochondrial mass and target gene expression mediated by Tro.

Blocking PGC-1 α degradation partially reverses the Tro-altered mitochondrial function

Because the mRNA level of PGC-1 α was actually modestly increased by Tro, the reduction in the level of PGC-1 α protein is probably due to a reduced translation rate and/or an enhanced protein degra-

mentation rate. The activity of PGC-1 α is subjected to regulation by the ubiquitin–proteasome pathway (Sano *et al.*, 2007). In order to establish that PGC-1 α protein degradation is an important aspect of Tro-induced cytotoxicity, we used a proteasome inhibitor MG132 to suppress the activity of the ubiquitin–proteasome pathway. We found that MG132 stabilized the expression of PGC-1 α (Supporting Information Figure S4); meanwhile, it partially restored the loss of mitochondrial mass (Figure 5A) and the suppressed expression levels of Tfam, cyt-C and ATP5b (Figure 5B,C; Supporting Information Figure S5). However, MG132 alone strongly increased mitochondrial ROS level (data not shown), negating our ability to analyse its effect on Tro-mediated alteration of ROS production.




Figure 4

Effect of Tro on PGC-1 α protein levels. (A–D) mRNA expression levels of (A) ERR α , (B) ERR γ , (C) PGC-1 α and (D) PGC-1 β were measured after 24 h drug treatments in HepG2 cells cultured in 2.5% FBS. The relative level under dimethyl sulphoxide (DMSO) treatment was set as 1. * P < 0.05, ** P < 0.01. (E) Representative Western blots of ERR α , ERR γ , PGC-1 α , PGC-1 β , with β -actin as a control. (F) The reductions in PGC-1 α protein level from three independent experiments were quantified against DMSO as a control.

Over-expression of PGC-1 α partially alleviates the Tro-mediated cytotoxicity

Because MG132 blocks the degradation of not only PGC-1 α , but also other proteins, we next determined whether over-expressing PGC-1 α is able to restore the reduction of viability induced by Tro. We constructed an adenovirus encoding PGC-1 α , and found that it effectively enhanced the protein expression level of PGC-1 α compared to a control adenovirus (Supporting Information Figure S6). We found that increasing the expression of PGC-1 α was able to partially restore the Tro-mediated reductions of Tfam, cyt-C and ATP5b expression (Figure 6A;

data not shown) and mitochondrial mass (Figure 6B). Additionally, an increased expression of PGC-1 α was effective in alleviating the Tro-suppressed cell viability (Figure 6C), strongly suggesting that a decrease in the protein level of PGC-1 α through degradation is an important event behind the Tro-induced cytotoxicity.

Discussion

Tro, Rosi and Pio all function well as PPAR γ agonists. As members of the TZD class of drugs, these




Figure 5

The effect of MG132 on the Tro-induced alteration in mitochondrial function. (A) The amounts of mitochondrial mass with or without 5 μM MG132 pretreatment in the presence of increasing amounts of Tro were measured. (B–C) mRNA expression levels of (B) Tfam and (C) ATP5b were measured. (A–C) Drug treatments for 24 h were performed in HepG2 cells cultured in 2.5% FBS. (B–C) The relative level under dimethyl sulphoxide treatment was set as 1. $**P < 0.01$, $***P < 0.001$.

compounds were developed and marketed as anti-diabetic agents albeit target-dependent and compound-specific side effects. Significantly, Tro was withdrawn from the market due to liver toxicity issues. Tro was estimated to induce 10 liver-related deaths in one million (Plosker and Faulds, 1999). Tro treatments also inflicted less deadly, but more frequent, idiosyncratic liver injury character-

ized by elevations in serum alanine transaminase and aspartate transaminase levels (Plosker and Faulds, 1999).

The mechanism behind the Tro-mediated hepatotoxicity is likely to involve alterations in mitochondrial function (Masubuchi, 2006). PGC-1 α serves as a transcriptional co-activator for a number of transcription factors (Handschin, 2009). In particular, PGC-1 α functions with ERR α and ERR γ to guide the expression levels of mitochondrial regulators and enzymes (Giguere, 2008). In this study, we found that Tro, but not Rosi or Pio, selectively stimulated the degradation of PGC-1 α protein (Figure 4). As a result, Tro reduced mitochondrial mass and induced ROS to initiate apoptosis (Figures 2 and 3). Using a ubiquitin–proteasome inhibitor MG132, we established that blocking PGC-1 α degradation partially suppressed the reduction of mitochondrial mass (Figure 5). Importantly, increasing the expression of PGC-1 α was partly able to reverse the Tro-induced reduction in mitochondrial mass and its effects on cell viability (Figure 6). Collectively, these results suggest that PGC-1 α degradation is an important aspect behind Tro-induced liver toxicity.

Down-regulation of PGC-1 α at the protein level is likely to be a primary event preceding some of the mitochondrial dysfunction previously observed with Tro. Specifically, we found that the expression levels of ATP5b, cyt-C and Tfam, which are regulated by PGC-1 α with ERR α or ERR γ (Giguere, 2008), were suppressed by Tro, and that blocking PGC-1 α degradation by MG132 restored their expressions (Figure 5). Suppressing the expression levels of these oxidative phosphorylation enzymes (ATP5b and cyt-C) and mitochondrial biogenesis regulator (Tfam) is likely to result in reduced oxidative phosphorylation and ability to generate ATP, which may be the main mechanisms behind the decreased $\Delta\psi_m$ and cellular ATP levels previously reported (Tirmenstein *et al.*, 2002; Nadanaciva *et al.*, 2007).

Although we found that Tro induced the degradation of PGC-1 α protein and modestly increased its mRNA level, the mechanism leading to these effects is still not known. The modest increases in ERR α , ERR γ and PGC-1 α mRNA levels (Figure 4A–C) may actually reflect a compensatory mechanism to minimize the impact of the reduction in PGC-1 α protein induced by Tro (Figure 4E). Because the Tro-induced decrease in PGC-1 α protein was partially blocked by the proteasome inhibitor MG132, Tro may enhance the expression level of the ubiquitin E3 ligase, SCF^{Cdc4}, which has been shown to induce the degradation of PGC-1 α protein (Olson *et al.*, 2008). Tro may also enhance the activities of glycogen syn-




Figure 6

The effect of increasing the expression of PGC-1 α on the reduction in cell viability induced by Tro. (A–C) HepG2 cells cultured in 2.5% FBS were mock infected (blank) or infected with either a control adenovirus (vector) or an adenovirus encoding PGC-1 α (Ad5-PGC-1 α) for 24 h before drug treatments for another 24 h. (A) mRNA expression levels of Tfam (left panel) and ATP5b (right panel) were measured as in Figure 5. (B) The amounts of mitochondrial mass were measured as in Figure 3. (C) Cell viabilities were measured as in Figure 2. (A–C) The relative level under dimethyl sulfoxide treatment was set as 1. * $P < 0.05$, ** $P < 0.01$.

these kinase 3 β (GSK3 β) and p38 MAPK; both function to enhance the degradation of PGC-1 α mediated by SCF^{Cdc4} (Olson *et al.*, 2008). Alternatively, Tro may alter the expression levels of certain microRNAs that would target the 3'UTR of PGC-1 α mRNA suppressing its translation.

Regardless of the mechanism, the Tro-mediated effects are more clearly demonstrated in HepG2 cells supplemented with relatively lower amounts of FBS (Figure 2). This serum concentration-dependent effect may reflect growth factor and/or survival signalling pathways that impact on the stability of PGC-1 α protein. We speculate that reductions in growth factor and/or survival signalling pathways may contribute to the extent of Tro-induced liver toxicity in type II diabetics. Liver insulin resistance is a key aspect of type II diabetes (Leclercq *et al.*, 2007). Insulin resistance involves down-regulation of insulin-induced phosphorylation of insulin receptor substrates (IRS1/2) that play important roles in activating phosphoinositide 3-kinase (PI3K) and its downstream kinases, including Akt (Tanti and Jager, 2009). PI3K and Akt are responsible for communicating the

survival signal (Vara *et al.*, 2004). Intriguingly, Akt can phosphorylate and suppress the activity of GSK3 β (Cross *et al.*, 1995). It is possible that suppressing the survival pathway due to insulin resistance may exacerbate the Tro-induced liver toxicity, partly by attenuating the negative regulation by Akt of GSK3 β that promotes SCF^{Cdc4}-mediated PGC-1 α degradation. Other physiological stresses that activate p38MAPK may further impact on the extent of liver toxicity induced by Tro by stimulating GSK3 β phosphorylation of PGC-1 α , leading to its degradation by SCF^{Cdc4} (Olson *et al.*, 2008).

Acknowledgements

We would like to thank Mr Linbing Qu for generating the adenovirus. The work was supported by grants from the National Basic Research Program of China (973-Program) [2006CB50390] and the Knowledge Innovation Program of the Chinese Academy of Sciences [KSCX2-YW-R-085].

Conflict of interest

None to disclose.

References

- Alexander SPH, Mathie A, Peters JA (2008). *Guide to Receptors and Channels (GRAC)*, 3rd edition. Br J Pharmacol 153: S1–S209.
- Bae M-A, Song BJ (2003). Critical role of c-Jun N-terminal protein kinase activation in troglitazone-induced apoptosis of human HepG2 hepatoma cells. Mol Pharmacol 63: 401–408.
- Chen L, Wong C (2009). Estrogen-related receptor alpha inverse agonist enhances basal glucose uptake in myotubes through reactive oxygen species. Biol Pharm Bull 32: 1199–1203.
- Cross DA, Alessi DR, Cohen P, Andjelkovich M, Hemmings BA (1995). Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. Nature 378: 785–789.
- Desvergne B, Wahli W (1999). Peroxisome proliferator-activated receptors: nuclear control of metabolism 10.1210/er.20.5.649. Endocr Rev 20: 649–688.
- Giguere V (2008). Transcriptional control of energy homeostasis by the estrogen-related receptors. Endocr Rev 29: 677–696.
- Handschin C (2009). The biology of PGC-1[alpha] and its therapeutic potential. Trends Pharmacol Sci 30: 322–329.
- Haskins J, Rowse P, Rahbari R, de la Iglesia F (2001). Thiazolidinedione toxicity to isolated hepatocytes revealed by coherent multiprobe fluorescence microscopy and correlated with multiparameter flow cytometry of peripheral leukocytes. Arch Toxicol 75: 425–438.
- Home PD, Pocock SJ, Beck-Nielsen H, Gomis R, Hanefeld M, Jones NP *et al.* (2007). Rosiglitazone evaluated for cardiovascular outcomes – an interim analysis. N Engl J Med 357: 28–38.
- Kamei Y, Ohizumi H, Fujitani Y, Nemoto T, Tanaka T, Takahashi N *et al.* (2003). PPAR γ coactivator 1 β /ERR ligand 1 is an ERR protein ligand, whose expression induces a high-energy expenditure and antagonizes obesity. Proc Natl Acad Sci U S A 100: 12378–12383.
- Leclercq IA, Da Silva Morais A, Schroyen B, Van Hul N, Geerts A (2007). Insulin resistance in hepatocytes and sinusoidal liver cells: mechanisms and consequences. J Hepatol 47: 142–156.
- Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM, Kliewer SA (1995). An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor γ (PPAR γ). J Biol Chem 270: 12953–12956.
- Lim PLK, Liu J, Go ML, Boelsterli UA (2008). The mitochondrial superoxide/thioredoxin-2/Ask1 signaling pathway is critically involved in troglitazone-induced cell injury to human hepatocytes. Toxicol Sci 101: 341–349.
- Lincoff AM, Wolski K, Nicholls SJ, Nissen SE (2007). Pioglitazone and risk of cardiovascular events in patients with type 2 diabetes mellitus: a meta-analysis of randomized trials. JAMA 298: 1180–1188.
- Masubuchi Y (2006). Metabolic and non-metabolic factors determining troglitazone hepatotoxicity: a review. Drug Metab Pharmacokinet 21: 347–356.
- Nadanaciva S, Dykens JA, Bernal A, Capaldi RA, Will Y (2007). Mitochondrial impairment by PPAR agonists and statins identified via immunocaptured OXPHOS complex activities and respiration. Toxicol Appl Pharmacol 223: 277–287.
- Nie Y, Wong C (2009). Suppressing the activity of ERR α in 3T3-L1 adipocytes reduces mitochondrial biogenesis but enhances glycolysis and basal glucose uptake. J Cell Mol Med 13: 3051–3060.
- Niemeyer NV, Janney LM (2002). Thiazolidinedione-induced edema. Pharmacotherapy 22: 924–929.
- Olson BL, Hock MB, Ekholm-Reed S, Wohlschlegel JA, Dev KK, Kralli A *et al.* (2008). SCFCdc4 acts antagonistically to the PGC-1 transcriptional coactivator by targeting it for ubiquitin-mediated proteolysis. Genes Dev 22: 252–264.
- Ong MMK, Latchoumycandane C, Boelsterli UA (2007). Troglitazone-induced hepatic necrosis in an animal model of silent genetic mitochondrial abnormalities. Toxicol Sci 97: 205–213.
- Plosker GL, Faulds D (1999). Troglitazone: a review of its use in the management of type 2 diabetes mellitus. Drugs 57: 409–438.
- Sano M, Tokudome S, Shimizu N, Yoshikawa N, Ogawa C, Shirakawa K *et al.* (2007). Intramolecular control of protein stability, subnuclear compartmentalization, and coactivator function of peroxisome proliferator-activated receptor γ coactivator 1 α . J Biol Chem 282: 25970–25980.
- Scheen AJ (2004). Combined thiazolidinedione-insulin therapy: should we be concerned about safety? Drug Saf 27: 841–856.
- Shishido S, Koga H, Harada M, Kumemura H, Hanada S, Taniguchi E *et al.* (2003). Hydrogen peroxide overproduction in megamitochondria of troglitazone-treated human hepatocytes. Hepatology 37: 136–147.
- Staels B, Fruchart J-C (2005). Therapeutic roles of peroxisome proliferator-activated receptor agonists. Diabetes 54: 2460–2470.

- Tang WH, Francis GS, Hoogwerf BJ, Young JB (2003). Fluid retention after initiation of thiazolidinedione therapy in diabetic patients with established chronic heart failure. *J Am Coll Cardiol* 41: 1394–1398.
- Tanti J-F, Jager J (2009). Cellular mechanisms of insulin resistance: role of stress-regulated serine kinases and insulin receptor substrates (IRS) serine phosphorylation. *Curr Opin Pharmacol* 9: 753–762.
- Tirmenstein MA, Hu CX, Gales TL, Maleeff BE, Narayanan PK, Kurali E *et al.* (2002). Effects of troglitazone on HepG2 viability and mitochondrial function. *Toxicol Sci* 69: 131–138.
- Vara JAF, Casado E, de Castro J, Cejas P, Belda-Iniesta C, Gonzalez-Baron M (2004). PI3K/Akt signalling pathway and cancer. *Cancer Treat Rev* 30: 193–204.
- Wang J, Fang F, Huang Z, Wang Y, Wong C (2009). Kaempferol is an estrogen-related receptor [alpha] and [gamma] inverse agonist. *FEBS Lett* 583: 643–647.
- Watkins PB, Whitcomb RW (1998). Hepatic dysfunction associated with troglitazone. *N Engl J Med* 338: 916–917.
- Wu Z, Puigserver P, Andersson U, Zhang C, Adelmant G, Mootha V *et al.* (1999). Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell* 98: 115–124.
- Wu F, Wang J, Wang Y, Kwok T-T, Kong, S-K, Wong C (2009). Estrogen-related receptor [alpha] (ERR[alpha]) inverse agonist XCT-790 induces cell death in chemotherapeutic resistant cancer cells. *Chemico-Biol Interacts* 181: 236–242.
- Zhang H, Zhang A, Kohan DE, Nelson RD, Gonzalez FJ, Yang T (2005). Collecting duct-specific deletion of peroxisome proliferator-activated receptor γ blocks thiazolidinedione-induced fluid retention. *Proc Natl Acad Sci U S A* 102: 9406–9411.

Supporting information

Additional supporting information may be found in the online version of this article.

Figure S1 Transient transfections were performed as in 'Materials and Methods' except for using PPAR γ 2 expression vector. Six hours after transfection,

cells were treated with drugs for 24 h. Luciferase activity was measured and the relative activity was calculated and expressed as fold induction over dimethyl sulfoxide (DMSO) as a vehicle control.

Figure S2 HepG2 cells cultured in 2.5% fetal bovine serum (FBS) were treated dose-dependently with troglitazone, rosiglitazone and pioglitazone for (a) 24, (b) 48 and (c) 72 h before cell viability measurement as in 'Materials and Methods'. The relative level under dimethyl sulfoxide (DMSO) treatment was set as 1. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure S3 The mRNA expression levels of cyt-C in HepG2 cells cultured in 2.5% FBS after drug treatments for 24 hr were measured by quantitative RT-PCR. b-actin was used as a control and the relative level under DMSO treatment was set as 1. ** $P < 0.01$.

Figure S4 HepG2 cells cultured in 2.5% fetal bovine serum (FBS) were treated with drugs indicated. Twenty-four hours after treatment, the protein level of PGC-1 α and b-actin as a control were measured.

Figure S5 The mRNA expression levels of cyt-C in HepG2 cells cultured in 2.5% FBS after drug treatments for 24 h were measured by quantitative RT-PCR. b-actin was used as a control and the relative level under DMSO treatment was set as 1. *** $P < 0.001$.

Figure S6 HepG2 cells at 4×10^5 per well in 6-well plates were plated and incubated overnight before mock infection (blank) or infections with either a control adenovirus (vector) or an adenovirus encoding PGC-1 α (Ad5-PGC-1 α) at a multiplicity of infection of 50. Twenty-four hours after infection, the protein levels of PGC-1 α and b-actin as a control were measured.

Table S1 Sequences of primers used for RT-PCR are based on human genes and shown from 5' to 3'

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.