



Published in final edited form as:

J Occup Environ Med. 2015 October ; 57(10): 1031–1038. doi:10.1097/JOM.0000000000000534.

Trouble Sleeping Associated with Lower Work Performance and Greater Healthcare Costs: Longitudinal Data from Kansas State Employee Wellness Program

Siu-kuen Azor Hui, PhD, MSPH¹ and Michael A. Grandner, PhD, MTR²

¹Cancer Prevention and Control Program, Fox Chase Cancer Center, Philadelphia, PA

²Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA

Abstract

Objective—To examine the relationships between employees' trouble sleeping and absenteeism, work performance, and healthcare expenditures over a two year period.

Methods—Utilizing the Kansas State employee wellness program (EWP) dataset from 2008–2009, multinomial logistic regression analyses were conducted with trouble sleeping as the predictor and absenteeism, work performance, and healthcare costs as the outcomes.

Results—EWP participants (N=11,698 in 2008; 5,636 followed up in 2009) who had higher levels of sleep disturbance were more likely to be absent from work (all $p < 0.0005$), have lower work performance ratings (all $p < 0.0005$), and have higher healthcare costs ($p < 0.0005$). Longitudinally, more trouble sleeping was significantly related to negative changes in all outcomes.

Conclusions—Employees' trouble sleeping, even at a sub-clinical level, negatively impacts on work attendance, work performance, and healthcare costs.

INTRODUCTION

Sleep is an important area of focus in occupational medicine. Previous studies have shown associations between employees' sleep disturbances and a wide variety of negative occupational outcomes, including (1) absenteeism, (2) decreased productivity or presenteeism, (3) accidents and injuries, and (4) increased healthcare costs. According to a recent World Economic Forum report emphasizing chronic disease prevention at worksites as a strategy to enhance workforce wellness and performance, insufficient sleep is one of the eight major employee behaviors that employers should invest resources to address, to significantly reduce health care cost and increase productivity¹. In the US, employees' insufficient sleep caused an estimated \$150 billion in indirect costs (combined costs of absenteeism, presenteeism, and workplace accident or injuries)¹.

Address correspondences to: Siu-kuen Azor Hui, Fox Chase Cancer Center, Cancer Prevention and Control, 333 Cottman Ave., Young Pavilion 4171, Philadelphia, PA, 19111, USA, phone: 215-728-2453, fax: 215-214-2178, SKAzor.Hui@fccc.edu.

Conflict of interest statement: None declared from all authors.

Strong evidence from previous studies shows sleep disturbances are important factors of absenteeism; for instance, a study by Sivertsen and colleagues (2013)² examined data from the Hordaland Health Study and showed that insomnia and sleep apnea were both predictive of subsequent sick leave. These findings were echoed in the study by Bultmann and colleagues (2013)³, who found that in the Danish Work Environment Cohort Study, sleep disturbances and fatigue significantly predicted sickness absence. Rahkonen and colleagues (2012)⁴ examined data from employees of the City of Helsinki and found that frequent sleep problems were associated with increased sickness absences, both short and long in duration. Rajaratnam and colleagues (2011)⁵ found that police officers with probable sleep disorders were more likely to miss work as well. These and other studies suggest that poor sleep quality is associated with greater absenteeism.

Regarding decreased productivity at work (i.e. presenteeism) due to sleep problems, several studies have assessed these effects in varying ways. For example, Kessler and colleagues (2011)⁶ examined data from the American Insomnia Survey and found that poor sleep quality was significantly associated with lost work performance due to presenteeism. Swanson and colleagues (2011)⁷ found that self-reported symptoms of insomnia, sleep apnea, restless legs syndrome, and other sleep disorders were consistently associated with presenteeism. McKibben and colleagues (2010)⁸ found that sleep disturbances were associated with a 3-fold risk of impaired work performance and a 5-fold risk of limited day-to-day function among employees of the Florida Department of Health. In a landmark study, Rosekind and colleagues⁹ examined data from several US companies and found that for a typical good sleeper, the cost of decreased productivity due to insufficient sleep per year (based on salary) was \$1,293 per employee. This was increased to \$2,319 among those at risk for insufficient sleep, \$2,796 for those with insufficient sleep, and \$3,156 for those with insomnia. Also, this study found that the lost productivity was attributed to several domains, including impaired ability to meet time management demands, mental and interpersonal demands, output demands, and physical job demands.

Regarding increased accidents and injuries in occupational settings, many studies have showed that employees' sleep disturbances are significant risk factors. These findings have been reported across numerous professions, including physicians^{10,11}, nurses¹²⁻¹⁴, police officers⁵, truck drivers¹⁵⁻¹⁹, bus drivers²⁰, factory workers²¹, and others²². Further, Shahly and colleagues (2012)²³ found that self-reported poor sleep quality was associated with costly workplace accidents and errors. These studies focused on varying sleep factors (e.g., sleep apnea, sleep deprivation, shift work) but they show that, overall, trouble sleeping is an important risk factor for accidents and injuries.

In addition, employers incur substantial direct health care costs due to insufficient sleep of their employees. Strong evidence has shown that sleep deficiency or poor sleep quality is related to many chronic diseases such as coronary heart disease, diabetes, hypertension, overweight and obesity, and chronic stress and psychological problems²⁴.

Only a few studies to date have examined the potential impact of poor sleep on healthcare costs. This is an important consideration, given that healthcare costs are rising dramatically and this is a key expenditure for employers. Most of the studies in this domain have focused

on sleep apnea, showing that screening for, diagnosing, and/or treating sleep apnea can significantly reduce healthcare expenditures in occupational settings²⁵⁻²⁷. However, there have been a number of studies showing how ameliorating sleep problems can potentially reduce healthcare costs^{28,29}. These studies show that untreated sleep disorders, such as insomnia and sleep apnea, can profoundly increase healthcare expenditures. Despite these findings, previous studies have typically not examined the role of trouble sleeping in general, which may or may not meet criteria for a sleep disorder but may, still, impact health.

Moreover, not only poor sleep directly contribute to the chronic diseases, they may be indirectly contributing to their development through unhealthy behaviors²⁴, as previous studies indicated that sleep deficiency and/or poor sleep quality are associated with a number of behavioral risk factors of chronic diseases, such as smoking³⁰, alcohol abuse³¹, high stress³²⁻³⁴, low level of physical activities³⁵, and poor nutrition^{35,36}.

Despite these facts that sleep disturbances are contributing to numerous negative occupational outcomes and having sufficient and quality sleep are important health behaviors^{37,38}, the increasingly popular employee wellness programs (EWP) in US currently still lack a sleep improvement component to promote employee overall health. Among the employers offering a lifestyle management program in their EWP, most of them target nutrition/weight management (79%), smoking (77%), and fitness (72%)³⁹.

Taken together, the existing literature suggests that trouble sleeping may have an impact in a number of occupational demands, including absenteeism, presenteeism, accidents/injuries, and healthcare expenditures, but employers are not investing sufficient resources to alleviate these problems. Generalizability of previous studies on relations between sleep and occupational outcomes may be limited by a number of factors. For example, existing studies tended to focus on specific occupations (e.g., truck drivers), focus on specific sleep disorders (rather than general sleep problems), and focus on cross-sectional analyses (due to unavailability of longitudinal data). One possible avenue for an analysis that addresses some of these issues would be to examine data from an EWP in an organization large enough to include many different professions (increasing generalizability), using a measure of general sleep disturbance (to capture subclinical problems), and making use of follow-up data (to examine longitudinal relationships).

Accordingly, in order to bring greater awareness to employers about the significance of addressing sleep problems in the employee population, and establish a generalizable, quantified longitudinal effect of sleep disturbance on work performance and healthcare costs, the current study utilized a large Kansas State EWP dataset to examine the relationships between trouble sleeping and absenteeism, work performance, and healthcare expenditures over a two year period. This allowed us to investigate relationships between changes in trouble sleeping and changes in these important workplace-related outcomes.

METHODS

Data Source

The data for the current study were obtained through a data use agreement between the University of Kansas Medical Center and the Kansas Health Policy Authority in 2010. Data included basic personnel data of all Kansas state employees enrolled in the state health plans, as well as the complete individual-level responses of all HRA participants across 2008 and 2009. The personnel data in this dataset included the health plan members' age and total healthcare expenses (sum of expenses in medical care, prescription drugs and dental care) in the year for both years. These employees were eligible to participate in the Kansas State EWP, of which the online HRA was a major component. Each individual in these data had a unique alpha-numerical identifier. Because the coding of the numerical identifier was unknown to the authors, these data were not considered as personally identifiable, and it was deemed exempt by the Human Subjects Committee at the University of Kansas Medical Center.

Measures

All measures of sleep disturbance, absenteeism, and work performance were self-reported responses to the online health risk assessment (HRA) questionnaire in both 2008 and 2009. Online HRA is a gateway component of virtually all EWP, which collects information on employees' personal, familial, lifestyle, and emotional risk factors of common chronic diseases. Employees were given \$50 gift card to complete their online HRA and onsite biometric screening yearly.

Sleep disturbance was assessed with the question, "During the past 4 weeks, how often have you been bothered by any of the following problems?" with "Trouble Sleeping" as one item. The response choices were "Never," "Seldom," "Sometimes," "Often," and "Always".

Absenteeism was assessed by two questions: 1) In past 4 weeks, number of days you missed an entire work day because of problems with your physical or mental health (only include days missed for your own health, not someone else's health), and 2) In past 4 weeks, number of days you missed part of a work day because of problems with your physical or mental health (only include days missed for your own health, not someone else's health).

Self-rated work performance was assessed by the question, "On scale from 0 (worst) –10 (best), how would you rate your overall job performance on the days you worked during the past 4 weeks (28 days)?" Others' work performance was assessed by the question, "On scale from 0 (worst) –10 (best), how would you rate the usual performance of most workers in a job similar to yours?". Relative work performance score in our analysis was obtained by subtracting the others work performance rating from the self work performance rating.

Healthcare costs data were collected from the health services claims processed by the state employee health plans offered by the former Kansas Health Policy Authority (now subsumed in the Division of Health Care Finance, Kansas Department of Health and Environment). Covariates included age, sex, race/ethnicity, highest education level

achieved, total household income, and self-rated health. These were included since they are associated with both sleep quality and occupational factors in the literature.

Statistical Analyses

To examine relationships between trouble sleeping and baseline absenteeism, multinomial logistic regression analyses used absenteeism as outcome (0 days as reference, relative to 1–2 days, 3–6 days, and 7 or more days). Trouble sleeping was included as a categorical variable (reference = “never”). To investigate linear trends, the ordinal trouble sleeping variable was also assessed as a pseudo-continuous variable. To examine relationships between trouble sleeping and baseline self-rated performance, relative performance, and healthcare costs, these were input as continuous outcomes in multiple linear regression analyses. Trouble sleeping was again assessed as a categorical variable and a pseudo-continuous variable. Analyses were performed with and without covariates. To examine longitudinal changes in outcomes relative to longitudinal changes in trouble sleeping, change scores for all variables were computed by subtracting 2008 from 2009 data. (Thus, positive values mean an increase over 1 year.) Change scores for all variables were computed, including absenteeism variables which were treated as continuous for this calculation. Linear regression analyses examined trouble sleeping change scores as predictor of change scores for outcome variables, controlling for their baseline. All analyses were repeated after adjustment for covariates. *P* values <0.05 were considered significant. All analyses were performed using STATA 12.0 software (College Station, TX).

RESULTS

Sample Characteristics

The sample consisted of N=11,698 participants assessed in 2008 and N=5,636 who were followed up in 2009. The participation rates in the online HRA were 26% and 19% in the two years respectively. Characteristics of the sample are reported in Table 1, which displays demographic and socioeconomic covariates, health status, occupational outcome variables (absenteeism, work performance, and healthcare costs), and trouble sleeping.

Table 1 also displays differences between the complete sample and those that provided longitudinal data. Although only 48% of respondents provided follow-up data, this group did not differ from the full sample or those that only provided the first year data in any clinically meaningful way. For example, age, sex, race/ethnicity, income, and other factors were similarly distributed.

Absenteeism

Results of analyses assessing absenteeism at baseline are reported in Table 2. These include multinomial logistic regression analyses with absenteeism as outcome, (odds of 1–2, 3–6, and 7 or more days, relative to 0 days) and trouble sleeping as predictor. When trouble sleeping was assessed as a categorical variable, higher levels of trouble sleeping were associated with greater likelihood of absenteeism. This was consistent for missed full days, missed partial days, and total missed days. In addition, in all cases, a significant linear trend was found, demonstrating increased likelihood of each absenteeism category associated with

increasing levels of trouble sleeping. This pattern was maintained for both unadjusted and adjusted analyses.

Work Performance

Results of analyses assessing trouble sleeping associated with work performance measured at baseline are reported in Table 3. Regarding self-rated recent work performance, trouble sleeping was consistently associated with lower self-ratings of work performance. In addition trouble sleeping was consistently associated with a greater discrepancy between self-reported recent work performance and self-reported average performance of a worker in their job. Although workers typically rated themselves as above average, the degree to which they reported themselves to be above average depended on trouble sleeping. A linear trend between trouble sleeping and work productivity was also found. This was consistent for both unadjusted and adjusted analyses.

Healthcare Costs

Results of analyses assessing sleep disturbance at baseline with total healthcare costs for that year are also reported in Table 3. More trouble sleeping was, in general, associated with greater healthcare costs. For example, workers who report that they “always” experience trouble sleeping were associated with a mean \$5,206 in healthcare expenditures above those who “never” have problems; after adjusting for covariates, including overall health, this discrepancy was maintained but attenuated, representing an increased cost of \$3,461. In addition, a linear trend was found, such that in adjusted analyses, each category increase in the variable measuring trouble sleeping was associated with an additional \$725 cost.

Longitudinal Change

Table 4 describes relationships between change in trouble sleeping and change in absenteeism, work performance and healthcare costs over 1 year. Linear relationships between changes in trouble sleeping and changes in all outcomes were detected in both adjusted for baseline only and adjusted for baseline and covariates analyses. For example, in adjusted for baseline and covariates analysis, every 1-unit worsening in trouble sleeping over 1 year was associated with missing approximately 0.26 days (including 0.12 full and 0.14 partial days), a 8% decline in self-rated work performance, an 6% decline in relative work performance, and an increase of \$340 in healthcare expenditures.

DISCUSSION

The current study investigated the relationships between employees’ sleep disturbance and work attendance, work performance, health care costs over a two-year period, using a large Kansas State EWP participants’ HRA data. Our analyses found that cross-sectionally, higher levels of sleep disturbance were associated with greater likelihood of absenteeism (either full days or partial days), greater likelihood of lower self-ratings of work performance (either self only or relative to other workers). In terms of health care costs, our cross-sectional analyses also found significant association between more frequent trouble sleeping and higher health care costs. More importantly, in our longitudinal analyses, we found that worsening of sleep disturbance over one year was associated with further absenteeism, low

work productivity, and higher health care costs. These findings suggest that trouble sleeping of employees, even at a sub-clinical level, have significant negative impact on work performance and healthcare costs, which are important occupational outcomes to employers.

Previous studies on the relation between sleep disorders (e.g. insomnia, sleep apnea, etc.) or sleep disturbances (e.g. disrupted, insufficient sleep) variables and absenteeism reported that workers with these sleep problems are more likely to be absent from work^{7,40,41} and hence loss of productivity. This is detrimental to both the employee and employer. Our results are consistent with the literature with additional findings on the missed full days vs. partial days. After adjusting for potential covariates, we found that the employees who indicated even “seldom” had trouble sleeping were significantly associated with up to 6 missed days total. This association was stronger and the number of missed days total was higher as the level of trouble sleeping increased. This highlights the potential absenteeism consequence of even low frequency of trouble sleeping.

Perhaps more concerning to employers is the relation between trouble sleeping and presenteeism or lower work performance on the job. Several studies have found significant associations between sleep disturbances and lower work performance, more errors at work, more work disabilities, or more accidents at work^{6-9,42-46}. The presenteeism problem is more serious than the absenteeism one, as it is more costly to the employers. Not only the employers are paying the employees for being present at work, they are also more likely to pay for longer work time to complete a task or any compensation due to errors or disabilities caused by the fatigued (cognitively or physically) employees. Our results again confirmed the previous literature in this relation, in that even “seldom” had trouble sleeping was associated with lower subjective and relative work performance ratings. Our data added new insight on relative work performance, suggesting higher level of trouble sleeping is related to lesser degree of above average work performance. The correlation between trouble sleeping and lower productivity at work is alarming, and the fact that over half of our sample (56%) reported some level of sleep disturbance calls for development and implementation of effective intervention to monitor and improve employees’ sleep health.

Regarding the relation between trouble sleeping and health care expenditure, literature is limited in this area, but the available studies suggest the increased health care service utilization⁴² and increase medical and prescription costs⁴⁰ among employees with insomnia, as opposed to those without. Very few studies have been able to analyze actual healthcare expenses, rather they made estimated economic costs of workplace productivity loss associated with poor sleep^{6,9}. Our findings provided concrete evidence that each unit of increased sleep disturbance is associated with progressively higher total healthcare expense. This linear relation was true for employees whose sleep disturbance may not have met the diagnostic criteria of insomnia as well. The implications of this finding could be that as sleep disturbance increases, the employee either actually experienced more illnesses that need health care, or perceived to experience ill health and sought more health care services. These phenomena are likely due to the various negative physiological or mental health effects of sleep disturbances. Either way, the poor health status caused by trouble sleeping among employees directly cost employers’ business outcome, especially most employers are still paying for a large portion of their employees’ health insurance in the US.

A few strengths of the current study should be noted. Our HRA participants sample was large and occupationally diverse. It encompasses Kansas state employees from many industries (e.g., education, transportation, healthcare, administration etc.), the data from the present study is likely to be generalizable to multiple industries. Our dataset also included objective data on the healthcare expenses, which allowed us to examine the actual dollars spent associated with different degrees of trouble sleeping. Another unique aspect of our data is that we had the longitudinal HRA responses data across two years, which allowed us to examine the association between worsening trouble sleeping and absenteeism, work performance, and health care costs outcomes. Last but not least, our sample included employees who had sleep disturbances that might not have met diagnoses of sleep disorders, so we could examine the relation between sub-clinical sleep problems and important occupational outcomes.

Limitations

The single-item measure of sleep quality is problematic for several reasons. Most importantly, this question has not been specifically validated against any standard sleep measure; thus it is unclear to what degree the construct captured by this item represents better-validated measures of sleep. Second, self-reported, single-item, retrospective sleep items are not ideal for assessing sleep. Objective measures such as actigraphy and prospective measures such as sleep diary would be ideal. Nonetheless, single-item sleep quality measures have proven useful in many previous studies²⁴. The HRA responses data we used were self-reported, so we cannot know the actual respondents' absenteeism and work performance. The significant linear trends and associations found were from the cross-sectional data at baseline, and causal inference cannot be made. Our sample was also geographically limited to the Kansas State.

Finally, the low HRA participation rate could potentially have resulted in a sample biased on one of the measures of interest. This HRA participation rate (about 20%) is typical among EWP³⁹ and since the participation rate in the present study is in line with that of most other studies, the data are likely to be at least as representative as is the standard in the literature. Previous studies reported that when participation rates are lower than 30%, female workers are more likely to participate in worksite health promotion programs, though no other systematic demographic differences (e.g. age, race/ethnicity, marital status, education, income level) between participants and non-participants were consistently found^{47, 48} using Chi-square, *t*-tests or meta-analysis techniques (e.g. Cohen's *d*). This was also the case in our study population.

Further studies will be needed to address the weaknesses of this study, such as using more objective and standard subjective measure of sleep disturbance and objective measures of absenteeism, and work performance. It would also be more desirable to have longitudinal data with longer follow-up time to confirm the trend we found between the two years. The longer follow-up longitudinal data will also allow investigation of whether improved sleep over time may reverse the negative effect of absenteeism, work performance and health care costs.

Conclusions

The present study demonstrated that trouble sleeping was associated with a greater likelihood of missed work days, lower work performance (either subjective or relative), and higher overall health care costs. Longitudinal data analyses across two years also demonstrated that each unit of worsening trouble sleeping over time was associated with more missed work days, decline in work performance, and increased health care costs over time. These results indicate that it is important for employers to incorporate sleep improvement intervention as one of the essential lifestyle change interventions offered in EWP to promote health and productivity of the large employee population.

Acknowledgements

We thank Drs. Ellerbeck and Shireman at University of Kansas Medical Center, and Ms. Cheryl Miller (the former Program Administrator of the Kansas Employee Wellness Program) for facilitating the data acquisition for this study.

Sources of funding: Dr. Hui is supported by the National Cancer Institute (R03CA159903). Dr. Grandner is supported by the National Heart, Lung and Blood Institute (K23HL110216) and the National Institute of Environmental Health Sciences (R21ES022931).

REFERENCES

1. World Economic Forum. The new discipline of workforce wellness: Enhancing corporate performance by tackling chronic disease. Geneva: 2010.
2. Sivertsen B, Bjornsdottir E, Overland S, Bjorvatn B, Salo P. The joint contribution of insomnia and obstructive sleep apnoea on sickness absence. *J Sleep Res.* 2013 Apr; 22(2):223–230. [PubMed: 23043357]
3. Bultmann U, Nielsen MB, Madsen IE, Burr H, Rugulies R. Sleep disturbances and fatigue: independent predictors of sickness absence? A prospective study among 6538 employees. *European journal of public health.* 2013 Feb; 23(1):123–128. [PubMed: 22383478]
4. Rahkonen O, Lallukka T, Kronholm E, Vahtera J, Lahelma E, Laaksonen M. Sleep problems and sickness absence among middle-aged employees. *Scandinavian journal of work, environment & health.* 2012 Jan; 38(1):47–55.
5. Rajaratnam SM, Barger LK, Lockley SW, et al. Sleep disorders, health, and safety in police officers. *Jama.* 2011 Dec 21; 306(23):2567–2578. [PubMed: 22187276]
6. Kessler RC, Berglund PA, Coulouvrat C, et al. Insomnia and the performance of US workers: results from the America insomnia survey. *Sleep.* 2011 Sep; 34(9):1161–1171. [PubMed: 21886353]
7. Swanson LM, Arnedt JT, Rosekind MR, Belenky G, Balkin TJ, Drake C. Sleep disorders and work performance: findings from the 2008 National Sleep Foundation Sleep in America poll. *J Sleep Res.* 2011 Sep; 20(3):487–494. [PubMed: 20887396]
8. McKibben JB, Fullerton CS, Ursano RJ, et al. Sleep and arousal as risk factors for adverse health and work performance in public health workers involved in the 2004 Florida hurricane season. *Disaster medicine and public health preparedness.* 2010 Sep; 4(Suppl 1):S55–S62. [PubMed: 23105036]
9. Rosekind MR, Gregory KB, Mallis MM, Brandt SL, Seal B, Lerner D. The cost of poor sleep: workplace productivity loss and associated costs. *J Occup Environ Med.* 2010 Jan; 52(1):91–98. [PubMed: 20042880]
10. Mansukhani MP, Kolla BP, Surani S, Varon J, Ramar K. Sleep deprivation in resident physicians, work hour limitations, and related outcomes: a systematic review of the literature. *Postgraduate medicine.* 2012 Jul; 124(4):241–249. [PubMed: 22913912]

11. Reed DA, Fletcher KE, Arora VM. Systematic review: association of shift length, protected sleep time, and night float with patient care, residents' health, and education. *Ann Intern Med.* 2010 Dec 21; 153(12):829–842. [PubMed: 21173417]
12. Dorrian J, Paterson J, Dawson D, Pincombe J, Grech C, Rogers AE. Sleep, stress and compensatory behaviors in Australian nurses and midwives. *Revista de saude publica.* 2011 Oct; 45(5):922–930. [PubMed: 21845291]
13. Scott LD, Hwang WT, Rogers AE, Nysse T, Dean GE, Dinges DF. The relationship between nurse work schedules, sleep duration, and drowsy driving. *Sleep.* 2007 Dec; 30(12):1801–1807. [PubMed: 18246989]
14. Chang YS, Wu YH, Hsu CY, Tang SH, Yang LL, Su SF. Impairment of perceptual and motor abilities at the end of a night shift is greater in nurses working fast rotating shifts. *Sleep Med.* 2011 Oct; 12(9):866–869. [PubMed: 21925944]
15. Sharwood LN, Elkington J, Meuleners L, Ivers R, Boufous S, Stevenson M. Use of caffeinated substances and risk of crashes in long distance drivers of commercial vehicles: case-control study. *BMJ (Clinical research ed.).* 2013; 346:f1140.
16. Zhang C, Berger M, Malhotra A, Kales SN. Portable diagnostic devices for identifying obstructive sleep apnea among commercial motor vehicle drivers: considerations and unanswered questions. *Sleep.* 2012 Nov; 35(11):1481–1489. [PubMed: 23115397]
17. Verster JC, Taillard J, Sagaspe P, Olivier B, Philip P. Prolonged nocturnal driving can be as dangerous as severe alcohol-impaired driving. *J Sleep Res.* 2011 Dec; 20(4):585–588. [PubMed: 21226780]
18. Gurubhagavatula I, Nkwuo JE, Maislin G, Pack AI. Estimated cost of crashes in commercial drivers supports screening and treatment of obstructive sleep apnea. *Accident; analysis and prevention.* 2008 Jan; 40(1):104–115.
19. Pack AI, Maislin G, Staley B, et al. Impaired performance in commercial drivers: role of sleep apnea and short sleep duration. *American journal of respiratory and critical care medicine.* 2006 Aug 15; 174(4):446–454. [PubMed: 16690976]
20. Razmpa E, Sadegh Niat K, Saedi B. Urban bus drivers' sleep problems and crash accidents. *Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India.* 2011 Jul; 63(3):269–273.
21. Aderaw Z, Engdaw D, Tadesse T. Determinants of Occupational Injury: A Case Control Study among Textile Factory Workers in Amhara Regional State, Ethiopia. *Journal of tropical medicine.* 2011; 2011:657275. [PubMed: 22174723]
22. Salminen S, Oksanen T, Vahtera J, et al. Sleep disturbances as a predictor of occupational injuries among public sector workers. *J Sleep Res.* 2010 Mar; 19(1 Pt 2):207–213. [PubMed: 19840241]
23. Shahly V, Berglund PA, Coulovrat C, et al. The associations of insomnia with costly workplace accidents and errors: results from the America Insomnia Survey. *Archives of general psychiatry.* 2012 Oct; 69(10):1054–1063. [PubMed: 23026955]
24. Grandner MA. Addressing sleep disturbances: An opportunity to prevent cardiometabolic disease? *International review of psychiatry.* 2014 Apr; 26(2):155–176. [PubMed: 24892892]
25. Mar J, Rueda JR, Duran-Cantolla J, Schechter C, Chilcott J. The cost-effectiveness of nCPAP treatment in patients with moderate-to-severe obstructive sleep apnoea. *The European respiratory journal.* 2003 Mar; 21(3):515–522. [PubMed: 12662011]
26. Tan MC, Ayas NT, Mulgrew A, et al. Cost-effectiveness of continuous positive airway pressure therapy in patients with obstructive sleep apnea-hypopnea in British Columbia. *Canadian respiratory journal : journal of the Canadian Thoracic Society.* 2008 Apr; 15(3):159–165. [PubMed: 18437259]
27. Ayas NT, FitzGerald JM, Fleetham JA, et al. Cost-effectiveness of continuous positive airway pressure therapy for moderate to severe obstructive sleep apnea/hypopnea. *Arch Intern Med.* 2006 May 8; 166(9):977–984. [PubMed: 16682570]
28. Wade AG. The societal costs of insomnia. *Neuropsychiatric disease and treatment.* 2010; 7:1–18. [PubMed: 21326650]
29. Leger D, Bayon V. Societal costs of insomnia. *Sleep Med Rev.* 2010 Dec; 14(6):379–389. [PubMed: 20359916]

30. de Leeuw R, Eisenlohr-Moul T, Bertrand P. The association of smoking status with sleep disturbance, psychological functioning, and pain severity in patients with temporomandibular disorders. *Journal of orofacial pain*. 2013 Winter;27(1):32–41. [PubMed: 23424718]
31. Chaput JP, McNeil J, Despres JP, Bouchard C, Tremblay A. Short sleep duration is associated with greater alcohol consumption in adults. *Appetite*. 2012 Dec; 59(3):650–655. [PubMed: 22841812]
32. Okamura H, Tsuda A, Yajima J, et al. Short sleeping time and psychobiological responses to acute stress. *International journal of psychophysiology : official journal of the International Organization of Psychophysiology*. 2010 Dec; 78(3):209–214. [PubMed: 20692301]
33. Lauterbach D, Behnke C, McSweeney LB. Sleep problems among persons with a lifetime history of posttraumatic stress disorder alone and in combination with a lifetime history of other psychiatric disorders: a replication and extension. *Comprehensive psychiatry*. 2011 Nov-Dec; 52(6):580–586. [PubMed: 21489420]
34. Kompier MA, Taris TW, van Veldhoven M. Tossing and turning--insomnia in relation to occupational stress, rumination, fatigue, and well-being. *Scandinavian journal of work, environment & health*. 2012 May; 38(3):238–246.
35. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Dietary nutrients associated with short and long sleep duration. Data from a nationally representative sample. *Appetite*. 2013 May;64:71–80. [PubMed: 23339991]
36. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Sleep symptoms associated with intake of specific dietary nutrients. *J Sleep Res*. 2014 Feb; 23(1):22–34. [PubMed: 23992533]
37. Colten, HR.; Altevogt, BM. *Sleep disorders and sleep deprivation : an unmet public health problem*. Washington, DC: Institute of Medicine : National Academies Press; 2006. Institute of Medicine Committee on Sleep Medicine and Research.
38. Zee PC, Badr MS, Kushida C, et al. Strategic opportunities in sleep and circadian research: report of the joint task force of the sleep research society and american academy of sleep medicine. *Sleep*. 2014; 37(2):219–227. [PubMed: 24501434]
39. Mattke, S.; Liu, H.; Caloyeras, JP., et al. *Workplace Wellness Programs Study: Final Report*. Santa Monica, CA: RAND Corporation; 2013.
40. Kleinman NL, Brook RA, Doan JF, Melkonian AK, Baran RW. Health benefit costs and absenteeism due to insomnia from the employer's perspective: a retrospective, case-control, database study. *The Journal of clinical psychiatry*. 2009 Aug; 70(8):1098–1104. [PubMed: 19758521]
41. Godet-Cayre V, Pelletier-Fleury N, Le Vaillant M, Dinet J, Massuel MA, Leger D. Insomnia and absenteeism at work. Who pays the cost? *Sleep*. 2006 Feb; 29(2):179–184. [PubMed: 16494085]
42. Rosekind MR, Gregory KB. Insomnia risks and costs: health, safety, and quality of life. *The American journal of managed care*. 2010 Aug; 16(8):617–626. [PubMed: 20712395]
43. Yumang-Ross DJ, Burns C. Shift work and employee fatigue: implications for occupational health nursing. *Workplace health & safety*. 2014 Jun; 62(6):256–261. quiz 262. [PubMed: 24971821]
44. Omachi TA, Claman DM, Blanc PD, Eisner MD. Obstructive sleep apnea: a risk factor for work disability. *Sleep*. 2009 Jun; 32(6):791–798. [PubMed: 19544756]
45. Douglas NJ. Sleep, performance and the European Working Time Directive. *Clinical medicine (London, England)*. 2005 Mar-Apr;5(2):95–96.
46. Niu SF, Chung MH, Chen CH, Hegney D, O'Brien A, Chou KR. The effect of shift rotation on employee cortisol profile, sleep quality, fatigue, and attention level: a systematic review. *The journal of nursing research : JNR*. 2011 Mar; 19(1):68–81. [PubMed: 21350389]
47. Lewis RJ, Huebner WW, Yarborough CM III. Characteristics of participants and nonparticipants in worksite health promotion. *American Journal of Health Promotion*. 1996; 11:99–106. [PubMed: 10163602]
48. Robroek SJ, van Lenthe FJ, van Empelen P, Burdorf A. Determinants of participation in worksite health promotion programmes: a systematic review. *Int J Behav Nutr Phys Act*. 2009; 6:26. [PubMed: 19457246]

Table 1

Characteristics of the Baseline Sample (N=11,698)

Variable	Category	Total Sample	2008 Data Only	2008 and 2009 Data
N		11,698	6,062	5,636
Age	Mean ± SD	44.60 ± 11.50	43.87 ± 12.02	45.39 ± 10.86
Sex	Female	63.93%	63.43%	64.46%
Race/Ethnicity	Non-Hispanic White	86.53%	85.43%	87.70%
	Black/African-American	4.51%	5.51%	3.44%
	Hispanic/Latino	3.42%	3.70%	3.12%
	Native American	2.62%	3.10%	2.11%
	Asian/Other	2.92%	2.26%	3.62%
Education	Post-graduate	25.58%	25.26%	25.92%
	College graduate	33.63%	31.92%	35.47%
	Some college	27.00%	27.52%	26.44%
	High school	13.17%	14.42%	11.83%
	Less than high school	0.62%	0.89%	0.34%
	\$100,000+	2.14%	2.46%	1.79%
Income	\$85,001 – \$100,000	1.89%	2.14%	1.61%
	\$55,001 – \$85,000	14.17%	13.30%	15.12%
	\$35,001 – \$55,000	39.61%	36.75%	42.67%
	\$20,001 – \$35,000	34.90%	36.19%	33.52%
	\$0–\$20,000	7.29%	9.16%	5.29%
	Excellent	11.56%	11.38%	11.75%
Health	Very Good	42.02%	40.91%	43.20%
	Good	38.74%	38.98%	38.48%
	Fair	7.27%	8.23%	6.23%
	Poor	0.42%	0.49%	0.34%
	0 Days	72.61%	71.61%	73.69%
Absenteeism (Missed Full Days in the past 4 weeks)	1–2 Days	22.06%	22.29%	21.82%
	3–6 Days	4.50%	5.13%	3.81%

Variable	Category	Total Sample	2008 Data Only	2008 and 2009 Data
Absenteeism (Missed Part Days)	7 or More Days	0.83%	0.97%	0.67%
	0 Days	76.39%	77.17%	75.55%
	1-2 Days	20.28%	19.28%	21.34%
	3-6 Days	2.80%	3.04%	2.54%
	7 or More Days	0.54%	0.51%	0.57%
Absenteeism (Missed Total Days)	0 Days	61.17%	61.33%	61.00%
	1-2 Days	27.42%	26.71%	28.19%
	3-6 Days	9.39%	9.72%	9.03%
	7 or More Days	2.02%	2.24%	1.77%
Work Performance (Subjective)	perf_r	8.32 ± 1.35	8.30 ± 1.38	8.33 ± 1.32
Work Performance (Relative)	perf_d1	0.89 ± 1.57	0.89 ± 1.60	0.88 ± 1.55
Healthcare Costs	Mean ± SD	5,016.65 ± 11,691.27	5,199.46 ± 13,689.51	4,820.03 ± 9,060.05
	Never	44.05%	43.60%	44.54%
Trouble Sleeping	Seldom	22.00%	20.69%	23.40%
	Sometimes	22.11%	23.00%	21.15%
	Often	8.60%	8.91%	8.27%
	Always	3.25%	3.81%	2.64%
	Mean ± SD	-0.09 ± 1.98		
Absenteeism (Missed Full Days) Change	Mean ± SD	-0.01 ± 2.03		
Absenteeism (Missed Part Days) Change	Mean ± SD	-0.10 ± 3.23		
Work Performance (Subjective) Change	Mean ± SD	-0.07 ± 1.41		
Work Performance (Relative) Change	Mean ± SD	0.06 ± 1.76		
Trouble Sleeping Category Change	4 Categories Improved	15.83%		
	3 Categories Improved	4.61%		
	2 Categories Improved	0.77%		
	1 Category Improved	0.14%		
	No Change	51.42%		
	1 Category Worse	19.83%		
2 Categories Worse	6.24%			

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Variable	Category	Total Sample	2008 Data Only	2008 and 2009 Data
	3 Categories Worse	1.00%		
	4 Categories Worse	0.16%		

Table 2

Results of Multinomial Logistic Regression Analyses Investigating Associations between Trouble Sleeping (Reference = Never) and Absenteeism (Reference = 0 Days)

Trouble Sleeping Category	1–2 Days			3–6 Days			7 Or More Days		
	OR	95% CI	P	OR	95% CI	P	OR	95% CI	P
UNADJUSTED									
<i>Missed Full Days</i>									
Never	Reference			Reference			Reference		
Seldom	1.40	1.25–1.58	<0.0005	1.30	0.99–1.70	0.054	1.41	0.73–2.71	0.301
Sometimes	1.66	1.48–1.87	<0.0005	2.33	1.84–2.95	<0.0005	2.52	1.42–4.44	0.001
Often	2.55	2.18–2.97	<0.0005	4.50	3.42–5.91	<0.0005	6.53	3.59–11.88	<0.0005
Always	2.71	2.13–3.43	<0.0005	6.48	4.53–9.26	<0.0005	11.52	5.75–23.09	<0.0005
<i>Linear Trend^a</i>	1.31	1.26–1.36	<0.0005	1.63	1.52–1.75	<0.0005	1.88	1.61–2.20	<0.0005
<i>Missed Partial Days</i>									
Never	Reference			Reference			Reference		
Seldom	1.45	1.29–1.64	<0.0005	1.64	1.19–2.26	0.002	2.90	1.22–6.90	0.016
Sometimes	1.74	1.54–1.96	<0.0005	2.18	1.61–2.94	<0.0005	5.32	2.43–11.65	<0.0005
Often	2.20	1.88–2.58	<0.0005	3.94	2.49–5.56	<0.0005	11.46	5.04–26.04	<0.0005
Always	2.76	2.19–3.49	<0.0005	5.81	3.73–9.07	<0.0005	10.38	3.45–31.23	<0.0005
<i>Linear Trend^a</i>	1.30	1.25–1.35	<0.0005	1.55	1.42–1.70	<0.0005	1.93	1.59–2.34	<0.0005
<i>Missed Days Total</i>									
Never	Reference			Reference			Reference		
Seldom	1.37	1.23–1.53	<0.0005	1.64	1.37–1.97	<0.0005	1.68	1.09–2.58	0.018
Sometimes	1.60	1.44–1.78	<0.0005	2.36	1.99–2.80	<0.0005	3.43	2.37–4.97	<0.0005
Often	2.12	1.81–2.47	<0.0005	4.54	3.68–5.59	<0.0005	8.43	5.66–12.55	<0.0005
Always	2.15	1.67–2.78	<0.0005	7.17	5.40–9.52	<0.0005	12.91	7.86–21–19	<0.0005
<i>Linear Trend^a</i>	1.25	1.21–1.30	<0.0005	1.63	1.55–1.72	<0.0005	1.98	1.79–2.20	<0.0005
ADJUSTED									
<i>Missed Full Days</i>									

Trouble Sleeping Category	1–2 Days			3–6 Days			7 Or More Days		
	OR	95% CI	P	OR	95% CI	P	OR	95% CI	P
Never	<i>Reference</i>								
Seldom	1.30	1.16–1.47	<0.0005	1.16	0.88–1.52	0.296	1.20	0.62–2.32	0.583
Sometimes	1.45	1.29–1.64	<0.0005	1.78	1.39–2.26	<0.0005	1.78	0.99–3.18	0.053
Often	1.95	1.66–2.29	<0.0005	1.76	2.07–3.68	<0.0005	3.70	1.99–6.90	<0.0005
Always	1.92	1.50–2.45	<0.0005	3.40	2.33–4.96	<0.0005	5.58	2.69–11.59	<0.0005
<i>Linear Trend</i> ^a	1.21	1.16–1.26	<0.0005	1.39	1.29–1.50	<0.0005	1.56	1.32–1.85	<0.0005
<i>Missed Partial Days</i>									
Never	<i>Reference</i>								
Seldom	1.36	1.20–1.54	<0.0005	1.49	1.08–2.05	0.016	2.38	1.00–5.69	0.051
Sometimes	1.54	1.36–1.74	<0.0005	1.78	1.31–2.42	<0.0005	3.86	1.74–8.55	0.001
Often	1.74	1.47–2.05	<0.0005	2.62	1.83–3.76	<0.0005	6.76	2.90–15.76	<0.0005
Always	2.03	1.59–2.58	<0.0005	3.42	2.15–5.47	<0.0005	5.37	1.72–16.70	0.004
<i>Linear Trend</i> ^a	1.21	1.16–1.26	<0.0005	1.36	1.24–1.50	<0.0005	1.64	1.33–2.01	<0.0005
<i>Missed Days Total</i>									
Never	<i>Reference</i>								
Seldom	1.28	1.14–1.43	<0.0005	1.46	1.22–1.76	<0.0005	1.44	0.94–2.23	0.097
Sometimes	1.42	1.27–1.58	<0.0005	1.87	1.57–2.23	<0.0005	2.48	1.69–3.62	<0.0005
Often	1.67	1.42–1.97	<0.0005	2.94	2.36–3.66	<0.0005	4.69	3.09–7.12	<0.0005
Always	1.60	1.24–2.08	<0.0005	4.08	3.03–5.50	<0.0005	6.02	3.57–10.14	<0.0005
<i>Linear Trend</i> ^a	1.17	1.12–1.21	<0.0005	1.42	1.35–1.50	<0.0005	1.64	1.47–1.82	<0.0005

^aEvaluating trouble sleeping as a pseudo-continuous ordinal variable; effects for 1-category increase

* Adjusted for age, sex, race/ethnicity, education, income, and overall health

Table 3

Associations Trouble Sleeping between, Work Performance, and Healthcare Costs

Trouble Sleeping Category	Subjective Performance (0 to 10)			Relative Performance (Self - Other)			Healthcare Costs (\$)		
	B	95% CI	P	B	95% CI	P	B	95% CI	P
UNADJUSTED									
Never	Reference			Reference					
Seldom	-0.16	-0.23 - -0.10	<0.0005	-0.11	-0.18 - -0.03	0.005	\$551.05	\$1.52-\$1100.58	0.049
Sometimes	-0.23	-0.29 - -0.16	<0.0005	-0.12	-0.19 - -0.04	0.002	\$1943.71	\$1395.11-\$2492.32	<0.0005
Often	-0.51	-0.60 - -0.42	<0.0005	-0.31	-0.41 - -0.20	<0.0005	\$3639.55	\$2854.87-\$4424.22	<0.0005
Always	-0.48	-0.62 - -0.34	<0.0005	-0.31	-0.47 - -0.14	<0.0005	\$5206.07	\$3995.97-\$6416.17	<0.0005
Linear Trend ^a	-0.14	-0.16 - -0.12	<0.0005	-0.08	-0.11 - -0.06	<0.0005	\$1166.70	\$981.73-\$1351.66	<0.0005
ADJUSTED*									
Never	Reference			Reference					
Seldom	-0.15	-0.21 - -0.09	<0.0005	-0.10	-0.17 - -0.02	0.011	-\$13.38	-\$560.54-\$533.77	0.962
Sometimes	-0.21	-0.28 - -0.15	<0.0005	-0.11	-0.18 - -0.03	0.005	\$1027.34	\$473.18-\$1581.50	<0.0005
Often	-0.42	-0.51 - -0.33	<0.0005	-0.25	-0.36 - -0.15	<0.0005	\$2337.19	\$1541.31-\$3133.07	<0.0005
Always	-0.36	-0.50 - -0.22	<0.0005	-0.24	-0.41 - -0.08	0.004	\$3461.89	\$2242.13-\$4681.64	<0.0005
Linear Trend ^a	-0.11	-0.14 - -0.09	<0.0005	-0.07	-0.09 - -0.04	<0.0005	\$725.15	\$532.98-\$917.33	<0.0005

^a Evaluating trouble sleeping as a pseudo-continuous ordinal variable; effects for 1-category increase

* Adjusted for age, sex, race/ethnicity, education, income, and overall health

Associations between Changes in Trouble Sleeping and Changes in Absenteeism, Work Performance, and Healthcare Costs

Table 4

	Adjusted for Baseline		Adjusted for Baseline and Covariates	
Absenteeism (Full Days)	0.14	0.09-0.19	<0.0005	0.12
Absenteeism (Partial Days)	0.15	0.10-0.21	<0.0005	0.14
Absenteeism (Total)	0.28	0.21-0.36	<0.0005	0.26
Subjective Performance	-0.08	-0.12 - -0.05	<0.0005	-0.08
Relative Performance	-0.08	-0.13 - -0.04	<0.0005	-0.06
Healthcare Costs	\$411.15	\$224.02-\$598.29	<0.0005	\$340.45
				\$152.60 - \$528.30
				0.001
				<0.0005