

UC Berkeley

UC Berkeley Previously Published Works

Title

Tuberculosis among health-care workers in low- and middle-income countries: a systematic review.

Permalink

<https://escholarship.org/uc/item/4sg6m98m>

Journal

PLoS medicine, 3(12)

ISSN

1549-1277

Authors

Joshi, Rajnish
Reingold, Arthur L
Menzies, Dick
et al.

Publication Date

2006-12-01

DOI

10.1371/journal.pmed.0030494

Peer reviewed

Tuberculosis among Health-Care Workers in Low- and Middle-Income Countries: A Systematic Review

Rajnish Joshi^{1,2}, Arthur L. Reingold¹, Dick Menzies³, Madhukar Pai^{3*}

1 Division of Epidemiology, School of Public Health, University of California Berkeley, Berkeley, California, United States of America, **2** Department of Medicine, Mahatma Gandhi Institute of Medical Sciences, Sevagram, Maharashtra, India, **3** Montreal Chest Institute, McGill University, Montreal, Canada

Funding: RJ is supported by a training fellowship from the Fogarty AIDS International Training Program (AITRP grant 1-D-43-TW00003–17). DM and MP receive research support from the Canadian Institutes of Health Research grants MOP-81362 and RRD153877. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Academic Editor: Mushtaque Chowdhury, BRAC, Bangladesh

Citation: Joshi R, Reingold AL, Menzies D, Pai M (2006) Tuberculosis among health-care workers in low- and middle-income countries: A systematic review. *PLoS Med* 3(12): e494. doi:10.1371/journal.pmed.0030494

Received: March 20, 2006

Accepted: October 11, 2006

Published: December 26, 2006

Copyright: © 2006 Joshi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: CI, confidence interval; HCW, health-care worker; IGRA, interferon-gamma release assay; IRR, incidence rate ratio; LMIC, low- and middle-income country; LTBI, latent TB infection; MDR-TB, multidrug-resistant tuberculosis; NTM, non-tuberculous mycobacteria; OR, odds ratio; PPD, purified protein derivative of tuberculin; TB, tuberculosis; TST, tuberculin skin test; TU, tuberculin unit; WHO, World Health Organization; XDR-TB, extensively drug-resistant tuberculosis

* To whom correspondence should be addressed. E-mail: madhukar.pai@mcgill.ca

ABSTRACT

Background

The risk of transmission of *Mycobacterium tuberculosis* from patients to health-care workers (HCWs) is a neglected problem in many low- and middle-income countries (LMICs). Most health-care facilities in these countries lack resources to prevent nosocomial transmission of tuberculosis (TB).

Methods and Findings

We conducted a systematic review to summarize the evidence on the incidence and prevalence of latent TB infection (LTBI) and disease among HCWs in LMICs, and to evaluate the impact of various preventive strategies that have been attempted. To identify relevant studies, we searched electronic databases and journals, and contacted experts in the field. We identified 42 articles, consisting of 51 studies, and extracted data on incidence, prevalence, and risk factors for LTBI and disease among HCWs. The prevalence of LTBI among HCWs was, on average, 54% (range 33% to 79%). Estimates of the annual risk of LTBI ranged from 0.5% to 14.3%, and the annual incidence of TB disease in HCWs ranged from 69 to 5,780 per 100,000. The attributable risk for TB disease in HCWs, compared to the risk in the general population, ranged from 25 to 5,361 per 100,000 per year. A higher risk of acquiring TB disease was associated with certain work locations (inpatient TB facility, laboratory, internal medicine, and emergency facilities) and occupational categories (radiology technicians, patient attendants, nurses, ward attendants, paramedics, and clinical officers).

Conclusions

In summary, our review demonstrates that TB is a significant occupational problem among HCWs in LMICs. Available evidence reinforces the need to design and implement simple, effective, and affordable TB infection-control programs in health-care facilities in these countries.

The Editors' Summary of this article follows the references.

Introduction

The risk of transmission of *Mycobacterium tuberculosis* from individuals with tuberculosis (TB) to other patients and to health-care workers (HCWs) has been recognized for many years [1,2]. This risk is greater when larger numbers of infectious (smear-positive) TB patients are managed at a health-care facility, and can be reduced with implementation of effective infection-control measures [3,4]. In the United States and other high-income countries, the risk of nosocomial transmission of TB was high in the pre-chemotherapy era, but declined with the reduction in incidence of TB disease in the population [3]. This trend, however, changed between 1985 and 1993, when several outbreaks of multidrug-resistant tuberculosis (MDR-TB) were reported in nosocomial and congregate settings in the United States [5,6]. This led to recommendations for a comprehensive set of infection-control practices to protect HCWs and reduce nosocomial transmission [1,4]. In the years following the publication of these recommendations, there was a dramatic decline in the burden of TB among HCWs [1,4,7]. In a recent review of TB among HCWs in high-income countries, the overall incidence of TB disease in the general population and native-born HCWs was less than 10 and 25 per 100,000 per year, respectively [8].

The situation is very different in low- and middle-income countries (LMICs), which account for more than 90% of the global TB burden [9,10]. Because these countries have high TB rates and limited resources [11,12], they focus largely on case detection and treatment using the DOTS strategy [1,9]. In these countries, even low-cost strategies to reduce TB transmission in health-care facilities are seldom implemented [3,13].

We conducted a systematic review to summarize the evidence on the incidence and prevalence of latent TB infection (LTBI) and TB disease among HCWs in LMICs. We specifically addressed the following questions: (1) What is the prevalence of LTBI and what are the risk factors for LTBI in HCWs? (2) What is the incidence of LTBI in HCWs and what risk factors are associated with higher incidence rates? (3) What is the incidence of TB disease in HCWs and how does it compare with the incidence in the population? (4) Are certain occupations, or some work locations within a health-care facility, at higher risk of TB than others? (5) How effective are various strategies in reducing the incidence of LTBI and/or disease among HCWs in LMICs?

Methods

Search Strategy

We searched the following electronic databases for primary studies: PubMed (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?DB=pubmed>, 1950 to December 2005), BIOSIS (<http://scientific.thomson.com/products/bp>, 1969 to December 2005), Embase (<http://www.embase.com>, 1974 to November 2005), and Web of Science (<http://isiknowledge.com>, 1945 to December 2005). Our search strategy included terms such as “tuberculosis,” “health personnel,” “health-care worker,” “nosocomial,” “infection control,” “disease transmission,” “occupational exposure,” and “nosocomial tuberculosis.” (Details of the complete search strategy are provided in Table S1.) We hand-searched the indices of the *International*

Journal of Tuberculosis and Lung Disease; Tuberculosis; and Tubercle and Lung Disease for relevant articles not already captured by the electronic searches. We identified additional studies by contacting experts in the field and by searching reference lists of primary studies, review articles, and textbook chapters.

Study Selection

Our search strategy aimed to identify all the available published studies in the English language that reported data on the incidence and prevalence of LTBI and TB disease in HCWs. Although non-English studies were excluded, we extracted data from studies that had English abstracts, and these limited data are included in Tables S1 and S2. This included cross-sectional tuberculin surveys, cohort studies on tuberculin-conversion rates, retrospective or prospective studies on the incidence of TB disease, studies on risk factors for acquiring LTBI or TB disease, and studies documenting the effect of preventive strategies aimed at reducing nosocomial transmission. We restricted the review to studies conducted in countries classified by the World Bank as low or middle income [14]. We excluded case reports or case series of nosocomial transmission or outbreaks, as well as conference abstracts. Two reviewers (RJ and MP) independently screened the citations (titles and abstracts) identified from all sources. Subsequently, full-text articles of the studies selected in the initial screen of titles and abstracts were reviewed to identify the final set of eligible studies.

Data Extraction

Two reviewers (RJ and MP) independently extracted data from a subset of eligible studies. The inter-rater agreement on TB outcomes (such as LTBI incidence and prevalence, and TB disease incidence, etc.) was 100% in this pilot study. Subsequently, data from the full set of included studies were extracted by one reviewer (RJ). Data extracted included: country, survey year, type of health-care facility, number of TB patients managed in the facility, infection-control practices (such as personal protection, administrative measures, engineering controls, etc.) in the facility at the time of the study, prevalence and incidence of LTBI, prevalence and incidence of TB disease, risk factors for LTBI or TB disease, infection-control interventions (personal, administrative, and engineering controls), and evaluations of their effectiveness, delays in diagnosis at the facility, and demographic and other relevant details about HCWs included in the studies. We used the following definitions to standardize the data-extraction process.

Health-care facility: All facilities where patients seek health care, including hospitals, clinics, dispensaries, health centers, and imaging and laboratory facilities. We did not include prisons, nursing homes, correctional facilities, and other congregate settings.

HCW: Any individual who works in a health-care setting including, but not restricted to, physicians, nurses, allied health personnel (nursing assistants, operation theater technicians, etc.), health educators, social workers, midwives, community health workers based in hospitals, laboratory personnel, pharmacists, radiographers, volunteers, orderlies, and health-facility administrators.

LTBI: A positive tuberculin skin test (TST) done by any standard method using 1TU (tuberculin unit) or 2TU of purified protein derivative of tuberculin (PPD) RT23 or 5TU




Figure 1. Study Flow Chart
doi:10.1371/journal.pmed.0030494.g001

of PPD-S, with induration size ≥ 10 mm on a single test [15]. Owing to the high prevalence of BCG vaccination and non-tuberculous mycobacteria (NTM) in LMICs [16], a positive result after the second TST in a two-step TST (i.e., boosting) was excluded from the calculation of LTBI prevalence.

Tuberculin conversion: Defined as a newly positive TST after a documented negative-baseline TST (at any time after a negative two-step baseline, or more than 1 y after a negative single TST). An increase of 10 mm over the baseline was defined as conversion [4].

TB disease: Included all forms of pulmonary, as well as extra-pulmonary, TB where a definitive (microbiologically confirmed), or presumptive (based on clinical, imaging, or pathology criteria) diagnosis was made. The definition included self-reported past treatment for TB disease.

Income category definitions: The countries were grouped according to 2004 gross national income per capita criteria as suggested by the World Bank, which classifies LMICs as those with per capita income value of less than US\$10,066 [14].

Infection-control interventions were defined as any personal protection (including, but not limited to, respirators), administrative measures (including, but not limited to, early diagnosis and isolation policy, reducing time for which TB patients would be hospitalized, and reducing waiting times for infectious patients in outpatient and radiology facilities), and environmental controls (including, but not limited to, negative-pressure isolation rooms, HEPA filters, etc.)

Data on the estimated incidence of all forms of TB disease in the general population were obtained from the World Health Organization's (WHO) global TB database [17]. All risk estimates among HCWs were calculated with respect to these country incidence rates. Since there may be considerable variations in TB incidence rates within a specific country, we also used the authors' estimates of regional/local incidence rates, if reported.




Figure 2. Prevalence of TB Infection in HCWs as Determined by TST Surveys

The circles and the lines represent the point estimates and 95% CIs, respectively. The size of the circle indicates the study size, and the diamond indicates the pooled estimate (weighted average) and its 95% CI. The heterogeneity chi-square statistics are 142.6 ($p < 0.01$) for the medical students and 600.9 ($p < 0.01$), for all HCWs, indicating significant heterogeneity across studies. (a) Indicates estimate of TST positivity in nursing students at entry into nursing school. (b) Indicates estimate of TST positivity after 3 y in nursing school.
doi:10.1371/journal.pmed.0030494.g002

Data Synthesis

Studies were heterogeneous in many respects, including baseline TB incidence in the population, institutional TB case loads, types of tests used to detect TB, job descriptions and classifications of HCWs, and preventive measures used at health-care facilities. Hence we analyzed the studies in prespecified subgroups. Studies of medical or nursing students were analyzed separately, as their risk of exposure may be different from other HCWs. Studies on TB disease were analyzed separately from studies on LTBI.

The incidence rates of LTBI and TB disease among HCWs, and corresponding estimates in the general population, were used to determine the excess risk among HCWs attributable to nosocomial exposure. We calculated the risk estimates for the incidence of TB disease in HCWs for various occupations and work locations, with incidence of TB disease in the general population as a reference. Data were analyzed using Stata (Version 9) and Meta-DiSc (Version 1.2) software.

In meta-analyses, heterogeneity refers to a substantial degree of variability in study results. Such heterogeneity can be due to differences in methodological quality, study design, sampling variability, and study populations across studies. In the presence of significant heterogeneity, pooled or summary estimates from meta-analyses are difficult to interpret. We addressed heterogeneity using subgroup (stratified) analyses. Because the studies estimating the

Table 1. Prevalence of LTBI in Medical or Nursing Students as Determined by TST Surveys in LMICs

Study; Country; (Number Tested)	TB Disease Incidence in the Country ^a ; (Year)	PPD Used; Cut Off; Reading Interval	TST Positivity; Positive-Tested (95% CI)	Proportion with BCG Scars (%)	Significant Risk Factors for Positive TST after Multivariate Analysis; OR (95% CI)
Keyanja 2005 [26]; Uganda; (53 medical students) ^b	403; (2004)	5TU Tubersol; 10 mm; 72 h+	21/53; 40% (26 to 54)	66	ND
Pai 2005 [28]; India; (226 medical students, 126 nursing students) ^b	168; (2004)	1TU RT23; 10 mm; 48–72 h	88/352; 25% (21 to 30)	71	Age (reference 18–20 y); 21–23 y; 1.5 (0.8 to 2.6)
Maciel 2005 [18]; Brazil; (178 nursing students)	62; (2003)	2TU RT23; 10 mm; 48–72 h	36/178; 20% (14 to 27)	100	Age (reference 18–20 y); 23 y+; 4.2 (1.9 to 9.3)
Silva 2000 [27]; Brazil; (455 medical students)	62; (2003)	2TU RT23; 10 mm; 48–72 h	33/455; 7% (5 to 10)	67.8	Training level (reference early or intermediate); senior year; 2.9 (1.1 to 7.6)
Teixeira 2005 [24]; Brazil; (1,032 medical students)	62; (2003) and 28–114 ^c ; (2002)	2TU RT23; 10 mm; 48–72 h	72/1,032; 7% (5 to 9)	88	Male sex; 1.8 (1.1 to 3.0)
Golchin 2005 [25]; Iran; (160 nursing students)	28; (2003)	5TU RT23; 10 mm; 48–72 h	3/160; 2% ^c (0.4 to 5) and 18/123; 15% ^d (9 to 22)	80.6	Training level (reference prediclinical); early clinical; 1.4 (0.7 to 2.6) Training level (reference prediclinical); late clinical; 1.9 (1.0 to 3.5) TB incidence in area ^e (reference low); high; 5.1 (1.6 to 16.8) TB incidence in area ^e (reference low); intermediate; 4.3 (1.3 to 14.6)
Levy 2005 [20]; Brazil; (84 nursing students, 86 medical students)	62; (2003)	5TU Tubersol; 10 mm; 48–72 h	119/170; 70% (62 to 77)	All were BCG-vaccinated 6 mo prior to TST	ND

^aAll figures represent WHO estimates [17] for incidence of all forms of TB, in the general population, in the country per 100,000 per year.

^bSubgroup analysis.

^cAt entry of nursing education.

^dOn completion of nursing education.

^eThis study was done in three regions with TB incidence in the general population of 114 (high), 63 (intermediate), and 28 (low) per 100,000.

ND, not determined.

doi:10.1371/journal.pmed.0030494.t001



Table 2. Prevalence of LTBI in All HCWs as Determined by TST Surveys

Study; Country; Number Tested	TB Disease; Incidence in the Country ^a ; (Survey Year)	PPD Used; Cut Off; Reading Interval	TST; Positive-Tested (95% CI)
Kassim 2000 [30]; Ivory Coast; (n = 512)	268; (1996)	5TU Tubersol; 10 mm; 48–72 h	405/512; 79% (75% to 82%)
Keskiner 2004 [33]; Turkey; (n = 491)	31; (2000)	5TU; 10 mm; 72 h+	355/491; 72% (68% to 76%)
Molina-Gamboa 1994 [34]; Mexico; (n = 175)	50; (1990)	5TU Tubersol; 10 mm; 72 h+	123/175; 70% (62% to 77%)
Do 1999 [35]; Thailand; (n = 911)	142; (2000)	5TU Tubersol; 10 mm; 48–72 h	623/911; 68% (65% to 71%)
Yanai 2003 [22]; Thailand; (n = 1,202)	142; (2000)	5TU Tubersol; 10 mm; 48–72 h	764/1,202; 63% (61% to 67%)
Alonso-Echanove 2001 [23]; Peru; (n = 270)	250; (1998)	2TU RT23; 10 mm; 48–72 h	170/270; 63% (57% to 69%)
Bonifacio 2002 [19]; Peru; (n = 97)	211; (2001)	5TU; 10 mm; 48–72 h	58/97; 60% (50% to 70%)
Kayanja 2005 [26]; Uganda; (n = 396)	403; (2004)	2TU RT23; 10 mm; 48–72 h	225/396; 57% (52% to 62%)
Roth 2005 [21]; Brazil; (n = 4,419)	62; (2003)	5TU Tubersol; 10 mm; 48–72 h	2,181/4,419; 49% (48% to 51%)
Orrett 2000 [32]; Trinidad; (n = 182)	11 ^b ; (1997)	5TU Tubersol; 10 mm; 72 h+	81/182; 44% (37% to 52%)
Pai 2005 [28]; India; (n = 719)	168; (2004)	1TU RT23; 10 mm; 48–72 h	298/720; 41% (38% to 45%)
Garcia-Garcia 2001 [31]; Mexico; (n = 823)	36 ^c ; (2000)	2TU RT23; 10 mm; 72 h+	280/823; 34% (31% to 37%)
Naidoo 2002 [29]; South Africa; (n = 78)	561; (2001)	NR; 10 mm; 24–48 h	26/78; 33% (23% to 45%)

^aAll figures represent WHO estimates [17] for incidence of all forms of TB, in the general population, in the country per 100,000 per year.

^bThe incidence in the general population of all forms of TB in Trinidad, as reported in [32], is 270 per 100,000.

^cThe study was conducted in Chihuahua, Mexico. The estimated incidence of all forms of TB in the general population in the city was 13.5 per 100,000 in the same year.

^dTB incidence (number of cases treated at each facility). High: TB clinic, infectious diseases, pulmonary medicine. Low: obstetrics, administrative.

^eAge is considered as a continuous variable in this model. OR represents risk for yearly increment in age.

NR, not reported.

ND, not determined.

doi:10.1371/journal.pmed.0030494.t002

prevalence of LTBI had comparable methodologies, we generated pooled (summary) estimates by using a fixed-effects model, where studies were weighed by the overall sample size, and we corrected for over-dispersion to allow for heterogeneity that was due to between-study variability.

Results

Description of Included Studies

Of the 1,901 unique citations identified in the literature search, 42 articles describing the results of 51 studies met our eligibility criteria (Figure 1). The prevalence of LTBI was determined by tuberculin surveys in 18 studies (35.3%) [18–35]. Seven studies (11%) [18–22,36,37] determined the incidence of LTBI using tuberculin conversions. Twenty

(39.2%) studies [23,38–56] determined the incidence of TB disease in HCWs. Six studies (11.7%) [21,22,24,44,48,57] reported the use of infection-control measures to reduce nosocomial transmission. Environmental attributes such as type of ventilation in the health-care facility, area devoted to patient care, patient and HCW crowding, and cumulative hospital stay of infectious patients were not reported by any study.

Prevalence of LTBI and Risk Factors for LTBI

Medical and nursing students. As seen in Figure 2 and Table 1, the prevalence of LTBI in medical or nursing students varied widely from 2% (Iran [25]) to 40% (Uganda [26]). The prevalence estimates correlated well (correlation coefficient $[R] = 0.91$, $p = 0.01$) with TB disease incidence rates in the

Table 2. Extended.

HCWs with BCG Scar (%)	Significant Risk Factors for Positive TST after Multivariate Analysis; OR (95% CI)	
	Nonoccupational	Occupational
83	Education level (reference no education): secondary or more 3.0 (1.3 to 6.9) Education level (reference no education): primary 2.9 (1.3 to 6.8)	Years in profession >1 y; 1.9 (1.1 to 3.3) Work location: high versus low TB incidence ^d ; 3.4 (1.8 to 6.2)
93	No. of BCG scars 1.1 (1.1 to 1.5)	Occupation (compared with female physicians): male physician 1.5 (1.23 to 1.69) Occupation (compared with female physicians): nurse 1.5 (1.29 to 1.66) Occupation (compared with female physicians): radiology technician 1.7 (1.35 to 1.73) Occupation (compared with female physicians): laboratory technician 1.6 (1.30 to 1.74) Occupation (compared with female physicians): male housekeeper 1.6 (1.38 to 1.40)
80	ND	ND
77	Male sex 2.3 (1.6 to 3.3) BCG scar present 1.9 (1.3 to 2.6)	Years in profession >1 y: 2.3 (1.2 to 4.3) Occupation (reference office clerks): physicians and nurses 1.7 (1.3 to 2.4) Occupation (reference office clerks): maintenance personal 1.8 (1.1 to 3.0)
82	Age > 30 y: 1.6 Male sex 1.6 BCG scar present 1.6	Years in profession >1 y: 1.5
NR		Years in profession >1: 1.5 (1.0 to 2.2) Work in medical ward: 2.1 (1.5 to 2.9); history of contact: 3.2 (1.9 to 5.3) Collected sputum: 1.5 (1.2 to 1.9) Use of common staff areas (laboratories): 2.7 (1.6 to 4.5)
NR	ND	ND
66	Age ^e 1.04 (1.02 to 1.07)	Work location (reference administrative work): medical wards 2.26 (1.17 to 4.38) Work location (reference administrative work): surgical/obstetrics 2.43 (1.30 to 4.55)
67	Age ^e 1.02 (1.02 to 1.03) Male sex 1.5 (1.2 to 1.7) BCG scar 1.9 (1.6 to 2.2)	>6 mo in profession 1.4 (1.2 to 1.7) Nurse 1.3 (1.1 to 1.5) Involved in patient care 1.2 (1.0 to 1.4)
4	Age (reference <25 y): 40 y+ 2.2 (0.6 to 18.5) Age (reference <25 y): 35–39 y 1.6 (0.3 to 15.9) Age (reference <25 y): 30–34 y 2.1 (0.5 to 19.2) Age (reference <25 y): male sex 1.2 (0.7 to 2.3)	Employment >1 y: 2.4 (1.1 to 5.0)
71	Age (reference 18–20 y): 41 y+ 2.9 (0.9 to 9.0) Age (reference 18–20 y): 31–40 y 1.7 (0.6 to 4.8) Age (reference 18–20 y): 21–30 y 1.9 (1.1 to 3.3)	Years in profession (reference ≤ 1 y work) >10: 3.2 (1.1 to 9.4) Years in profession (reference ≤ 1 y work) 6–10: 2.8 (1.2 to 6.2)
84	Age ^e 1.04 (1.02 to 1.07) BCG scar 2.1 (1.2 to 3.4)	Reference (other occupations combined): participation in autopsies 9.3 (2.1 to 40.5) Reference (other occupations combined): emergency/radiology 2.0 (1.0 to 3.8) Reference (other occupations combined): physicians and nurses 1.5 (1.0 to 2.1)
NR	ND	ND

general population (e.g., ranging from 28 per 100,000 in Iran to 403 per 100,000 in Uganda). As shown in Table 1, levels of training and age were associated with the prevalence of LTBI in most studies. The prevalence of LTBI in senior years was two to three times higher compared with junior years in two studies from Brazil [24,27]. A study from India [28] reported a 4-fold higher prevalence in medical students who were more than 23 y of age than in medical students aged 18–20 y (corresponding to an additional 3–5 y spent in training). The overall pooled prevalence of LTBI was 12% (95% confidence interval [CI] 10 to 13) among medical or nursing students, excluding the study by Levy et al. [20], where all medical students had received BCG vaccination within the previous 6 mo, and 70% of them were TST-positive. In other studies, where BCG was usually given in childhood, the presence of a BCG scar was not significantly associated with LTBI.

All HCWs. As shown in Figure 2 and Table 2, the prevalence of LTBI in all HCWs ranged from 33% (95% CI

23 to 45) [29] to 79% (95% CI 75 to 82) [30] in various studies, with a pooled prevalence estimate of 54% (95% CI 53 to 55). Increasing age and duration of employment in the health-care facility (indicating longer cumulative exposure), were associated with higher prevalence of LTBI in most studies (Table 2). The prevalence of LTBI in HCWs increased by 1.04 times (95% CI 1.02 to 1.07) with each additional year of increase in age [26,31], and by 1.5 (95% CI 1.0 to 2.2) [23] to 2.4 (95% CI 1.1 to 5.0) [31] times with employment duration of more than 1 y. The prevalence of LTBI was 3-fold higher [28] with ≥10 y of employment. Working in medical wards, participation in procedures such as sputum collection and autopsies, and a history of contact with TB patients were independent occupational risk factors for LTBI.

Several studies reported a prevalence of LTBI in nurses, a subgroup with a high level of patient contact, and thus potential exposure to TB cases. As seen in Table 3, the prevalence of LTBI among nurses ranged from 43% to 87%.

Table 3. Prevalence of LTBI in Nurses

Study; Country	Number of Nurses Tested	TB Disease Incidence in the Country ^a (Year)	Tuberculin Used; Cut Off; Reading Interval	TST Positivity (%) in Nurses (Nu)	TST Positivity in HCWs (All)	Difference in Prevalence (Excess Prevalence in Nurses) (Nu – All, 95% CI)
García-García 2001 [31]; Mexico	296	36 ^b ; (2000)	2TU RT23; 10 mm; 72 h+	69.6	34	35.6 (25.2 to 45.8)
Pai 2005 [28]; India	161	168; (2004)	1TU RT23; 10 mm; 48–72 h	65	41.4	23.6 (10.4 to 35.1)
Keskiner 2004 [33]; Turkey	75	31; (2000)	5TU RT23; 10 mm; 72 h+	86	72	14 (–9.1 to 35.2)
Kassim 2000 [30]; Ivory Coast	189 ^d	268; (1996)	5TU Tubersol; 10 mm; 48–72 h	87	79	8 (–7.6 to 23.0)
Kayanja 2005 [26]; Uganda	153	403; (2004)	5TU Tubersol; 10 mm; 72 h+	64	57	7 (–8.0 to 21.2)
Roth 2005 [21]; Brazil	1,597	62; (2003)	5TU Tubersol; 10 mm; 48–72 h	68	63	5 (14.0 to 23.1)
Do 1999 [35]; Thailand	361 ^d	142; (2000)	5TU Tubersol; 10 mm; 48–72 h	72	68	4 (–6.6 to 13.9)
Yanai 2003 [22]; Thailand	542 ^d	142; (2000)	5TU Tubersol; 10 mm; 48–72 h	64.8	63.5	1.3 (–6.9 to 9.3)
Orrert 2000 [32]; Trinidad	132	11 ^c ; (1997)	5TU Tubersol; 10 mm; 72 h+	43	44.5	–1.5 (–16.8 to 12.6)

^aAll figures represent WHO estimates [17] for incidence of all forms of TB, in the general population, in the country per 100,000 per year.

^bThe study was conducted in Chihuahua, Mexico. The estimated incidence of all forms of TB in the general population in the city was 13.5 per 100,000 in the same year.

^cThe incidence of all forms of TB in the general population in Trinidad, as reported in [32] is 270 per 100,000.

^dPhysicians and nurses combined.

The weighted average for TST positivity in all studies is 68 (95% CI 67 to 70). The weighted average for attributable risk is 11.5 (95% CI 9 to 14).
doi:10.1371/journal.pmed.0030494.t003

It was reported in eight studies that the prevalence of LTBI in nurses was higher than that in other HCWs (ranging from being 1.3% higher [22] to 35.6% higher [31]). One study, however [32], reported a lower prevalence in nurses, compared to other HCWs).

Incidence of LTBI (Tuberculin Conversion) and Risk Factors

In six out of the seven studies covered in Table 4, the annual risk of TB infection ranged from 3.9% to 14.3%, after accounting for the incidence of TB infection in the general population, and the risk attributable to occupational exposure ranged from 2.6% to 11.3%. (A study in medical students by Levy et al. [20], where all participants were BCG-vaccinated 6 mo prior to the first TST was excluded, as this vaccination program could be responsible for a low LTBI incidence of 0.5% in this study). There was marked, although non-significant, correlation between the incidence of LTBI and TB hospital admissions per year ($R = 0.86$, $p = 0.13$). A higher level of clinical training (odds ratio [OR] = 4.77, 95% CI 1.01 to 22.46) [36], BCG vaccination after baseline TST (OR 2.9, 95% CI 1.1 to 7.6) [21], nursing occupation (OR 1.7, 95% CI 1.1 to 2.7) [21], and recent exposure to TB (OR 1.6, 95% CI 1.0 to 2.6) [21] were independent risk factors for TST conversion in these studies.

Incidence of TB Disease

Results of studies that estimated the incidence of TB disease are shown in Table 5. Two studies (one from Russia [41] and another from South Africa [39]) reported lower TB disease incidence in HCWs than in the general population. In South Africa, the incidence of TB in the general population increased rapidly (from 321 per 100,000 per year in 1991 to 1,250 per 100,000 per year in 1996), with 44% of these cases being attributed to HIV infection [54]. The incidence of TB disease in Russia in the general population (113 per 100,000) was much higher than in the Samara Oblast (74.9 per 100,000), where a study by Dimitrova et al. [41] was carried out, which could be responsible for the low attributable risk estimates in the Dimitrova study [41]. In the remaining studies, the risk to HCWs of TB disease attributable to nosocomial exposure ranged from 25 to 5,361 per 100,000 per year. HCWs in facilities with fewer HCWs for every TB patient seen (HCW-to-patient ratio less than 50 per 100 TB patients), had a higher incidence of TB disease in HCWs ($R = -0.45$, $p = 0.18$).

Association between Work Location and Job Categories with Risk of TB Disease

Studies that reported rates of TB disease by work location and occupational categories are shown in Table 6. We calculated incidence rate ratios (IRRs) by using an estimated general population incidence rate for all TB cases in the country as a comparison. Workers in TB inpatient facilities (IRR ranged from 14.6 to 99.0), laboratories (IRR = 78.6), general medicine wards (IRR ranged from 3.9 to 36.6), and emergency rooms (IRR ranged from 26.6 to 31.9) had a higher risk for TB disease compared with the general population. Workers in outpatient medical facilities had an intermediate risk (IRR ranged from 4.2 to 11.6), and workers in surgery, obstetrics and gynecology, administration, and operating theaters had a lower risk. There was considerable heterogeneity in the risk of TB disease between different



Table 4. Incidence of LTBI or Tuberculin Conversion in HCWs

Study; Country; TB Incidence in Population ^a (Year)	Type of HCWs; Number and Period of Follow-up	PPD Used; Testing Protocol; Conversion Increment	Exposure to TB in Hospital TB Admissions per Year	Infection-Control Measures Used	Number Converted/Retested	Annual Risk of TB Infection in HCWs (95% CI)	Annual Risk of TB Infection in the Population (%)	Annual Risk of Infection Attributable to Nosocomial Transmission
Bonifacio 2002 [19]; Peru; 211; (2001)	Residents, interns; n = 35; 12 mo	5TU; one-step; 10 mm	1,220	Only 7.4% self-reported consistent use of N95 mask.	5/35	14.3% (4.8% to 30.2%)	3 ^b	11.3
Roth 2005 [21]; Brazil; 62; (2003)	All HCWs; n = 1,209; 6–12 mo	5TU; two-step; 10 mm	960 ^b	Two out of four hospitals had administrative, personnel, and engineering control	105/1,209	8.7% (7.2% to 10.4%)	0.5 ^c	8.2
Yanai 2003 [22]; Thailand; 142; (2000)	All HCWs; n = 164; 331.6 py	5TU; two-step; 10 mm	356	Respiratory-isolation facilities; 49% used N95 masks	24/331.6 py	7.2% (1.9% to 12.5%)	1.4 ^d	5.8
Maciel 2005 [18]; Brazil; 62; (2003)	Nursing students; n = 76; 2 y	2TU; two-step; 10 mm	NR	Isolation; PFR 95 masks; surgical masks used	8/76	5.3% (1.4% to 13%)	0.5 ^c	4.8
Silva 2002 [36]; Brazil; 62; (2003)	Medical students; n = 414; 12 mo	2TU; two-step; 10 mm	270	Respiratory isolation and respirator masks used	16/414	3.9% (1.1% to 12.1%)	0.5 ^c	3.4
Pai 2005; India [37]; 168; (2004)	Medical, nursing students; n = 147; 18 mo	1TU; one-step; 10 mm	NR	None	6/147	4.1% ^g (1.5% to 8.7%)	1.5 ^f	2.6
Levy 2005 [20]; Brazil; 62; (2003)	Medical, nursing students; n = 46 ^c ; 48 mo	5TU; two-step; 10 mm	NR	All were BCG-vaccinated 6 mo prior to TST	1/46	0.5% (0.4% to 0.8%)	0.5 ^e	0

^aAll figures represent WHO estimates [17] for incidence of all forms of TB, in the general population, in the country per 100,000 per year.

^bThis study was conducted in four hospitals.

^cThese HCWs were administered BCG vaccine 6 mo prior to baseline TST and were still negative.

^dEstimated ARTI for Thailand in 1997 [75].

^eNational estimates for Brazil [76].

^fNational estimates for India [77].

^gThis Indian study reported a higher ARTI estimate of 5% when a whole-blood interferon-gamma assay was used to detect conversions. If that estimate is used, the risk attributable to nosocomial exposure would be 3.5%.

^hAnnual risk was estimated in urban shantytowns.

ARTI, annual risk of TB infection; NR, not reported; py, person-year.

doi:10.1371/journal.pmed.0030494t004



Table 5. Incidence of TB Disease (All Forms) in HCWs

Study; Country	Type of Facility and Study	HCWs Exposed/ 100 TB Patients Seeking Care	HCWs per 100 TB Patients	HCWs with TB Disease/ Person-Year Exposure	TB Disease Incidence among HCWs ^a	TB Disease Incidence in the Country ^b ; (Year)	IRR (95% CI)	Risk of TB Disease Attributable to Nosocomial Transmission
Kanyere 2003 [50]; Malawi	Central hospital; retrospective cohort	571/3,607	16	33/571	5,780	419 ^d ; (2001)	13.8 (9.4 to 19.6)	5,361
Eyob 2002 [42]; Ethiopia	TB referral center; retrospective cohort	90/4,862	2	5/90 ^c	5,556	277 ^g ; (1998)	20.0 (6.5 to 47.3)	5,279
Harris 1999 [46]; Malawi	Across 40 hospitals; retrospective cohort	3,042/14,532	21	108/3,042	3,550	389 ^d ; (1996)	9.1 (7.3 to 11.3)	3,161
Harris 2002 [44]; Malawi	Across 40 hospitals; retrospective cohort	2,979/NR	—	96/2,979	3,222	389 ^d ; (1996)	8.3 (6.5 to 10.3)	2,833
Harris 1997 [45]; Malawi	Central hospital; retrospective cohort	310/1,112	28	12/620	1,935	389 ^d ; (1996)	4.9 (2.5 to 8.8)	1,546
Rao 2004 [55]; India	Tertiary-care hospital; prospective and retrospective cohorts	NR	—	13/1,032	1,260	168; (2002)	7.5 (3.9 to 13.2)	1,092
Babus 1997 [38]; Croatia	Nurses in two TB-care and one tertiary-care hospital; retrospective cohort	170/610	28	18/2,560	703	74 ^h ; (1990)	9.5 (5.3 to 16.1)	629
Naidoo 2006 [56]; South Africa	Across eight hospitals in province; retrospective cohort	49,392/NR	—	583/49,392	1,180	718 ⁱ ; (2004)	1.6 (1.5 to 1.8)	462
Alonso-Echanove 2001 [23]; Peru	Tertiary-care hospital; retrospective cohort	2,300/NR	—	36/9,200	391 ^e	111 ^g ; (1998)	3.5 (2.4 to 5.1)	280
Skodric 2000 [53]; Serbia	Pulmonary facility; retrospective cohort	267/600	44	9/3,204	287	59 ^f ; (1998)	4.8 (2.1 to 9.7)	228
Hosoglu 2005 [47]; Turkey	Tertiary-care facility; retrospective cohort	734/NR	—	22/11,010	199.9	40 ^f ; (1995)	5.0 (2.8 to 8.6)	160
Jelip 2004 [48]; Malaysia	All HCWs in the state; retrospective case control	7,312/NR	—	205/73,120	280.4	104 ^g ; (2003)	2.69 (2.1 to 3.4)	176.4
Wilkinson 1998 [54]; South Africa	Secondary-care hospital; retrospective cohort	725/760	95	22/4,350	506	379 ^g ; (1997)	1.33 (0.8 to 2.0)	127
Cuhadaroglu 2002 [40]; Turkey	Tertiary-care hospital; retrospective cohort	3,359/NR	—	31/33,590	96	29 ^f ; (2001)	3.2 (1.8 to 5.5)	67
Jiamrajasangi 2005 [49]; Thailand	Tertiary-care hospital; prospective cohort	3,959/NR	—	78/41,462	188	142 ^g ; (2000)	1.32 (1.0 to 1.7)	46
Gopinath 2004 [43]; India	Tertiary-care hospital; retrospective cohort	6,016/1,174	512	125/60,163	208	168; (2000)	1.2 (1.0 to 1.6)	40
Kilinc 2002 [51]; Turkey	Two secondary- and two tertiary-care hospitals each; retrospective cohort	6,156/NR	—	59/80,028	69	41 ^f ; (1998)	1.8 (1.2 to 2.7)	28
Kruuner 2001 [52]; Estonia	All HCWs in the country; retrospective cohort	14,730/806	1827	67/73,650	91	66; (1998)	1.4 (1.0 to 2.0)	25
Dimitrova 2005 [41]; Russia	All TB and general health facilities in Samara Oblast; retrospective cohort	64,855/2,445	2652	474/583,695	81.2	103 ^m ; (2002)	0.7 (0.6–0.9)	–32
Balt 1998 [39]; South Africa	Four TB-care hospitals; retrospective cohort	NR	—	2/726	275	379 ^g ; (1997)	0.7 (0.08 to 2.6)	–104

^aAll figures represent incidence of TB per 100,000 HCWs.

^bAll figures represent WHO estimates [17] for incidence of all forms of TB, in the general population, in the country per 100,000 population per year.

^cHCW figures are for 1998 only.

^dAuthors have used case-notification rates in the country concerned for comparison in their study.

^eThese figures represent sputum-positive pulmonary TB cases in both HCWs and the general population.

^fAuthors used average incidence rate over the entire study period for comparison.

^gThe regional TB incidence rates in the different studies are as follows:

^hThe incidence in the general population was 777 per 100,000 in 1997.

ⁱThe incidence in the general population was 48 per 100,000 (women aged 20–49 y).

^jThe incidence in the general population was between 386 and 782 per 100,000 from 1999 to 2003.

^kThe incidence in the general population was 55.4 per 100,000 in 2000 (women aged 15–64 y).

^lThe incidence in the general population was 154 per 100,000 in 2003.

^mThe incidence in the general population was 36 per 100,000 in 1998.

ⁿThe incidence in the general population was 74.9 per 100,000 in 2002.

^oThe incidence in the general population was 286 per 100,000 in 1997.

^pThe incidence in the general population was 1,543 per 100,000 in 1997 in individuals aged 15–43-y.

NR, not reported.

doi:10.1371/journal.pmed.0030494.t005



Table 6. Occupational Risk Factors for TB Disease in HCWs

Occupational Risk Factor	Occupational Risk Factor for TB Disease	IRR ^b	CI (95%)	Study Reference
Location of work	Inpatient TB facility (TB ward)	9.5	5.6 to 15.0	Babus 1997 [38]
		10.7	10.1 to 11.3	Dimitrova 2005 [41]
		28.6	0.6 to 206.1	Harries 1997 [45]
	Laboratory facility (general laboratory)	86.9	57.4 to 126.7	Kilinc 2002 [51]
		78.6	42.5 to 135.3	Alonso-Echanove 2001 [23]
		1.5	0.8 to 2.6	Jiamrajarasangi 2005 [49]
	Inpatient general medicine facility	14.0	9.5 to 20.0	Kilinc 2002 [51]
		18.9	8.4 to 37.2	Alonso-Echanove 2001 [23]
		35.4	9.4 to 100.9	Harries 1997 [32]
	Emergency facility/ICU	10.3	4.9 to 19.1	Jiamrajarasangi 2005 [49]
		31.9	10.2 to 76.8	Alonso-Echanove 2001 [23]
	Outpatient medicine/TB facility	2.8	2.5 to 3.1	Dimitrova 2005 [41]
		2.8	1.1 to 5.8	Jiamrajarasangi 2005 [49]
	Combined inpatient and outpatient TB facility	11.2	0.3 to 68.8	Harries 1997 [45]
		8.5	6.2 to 12.1	Dimitrova 2005 [41]
		0.8	2.8 to 1.7	Jiamrajarasangi 2005 [49]
	Surgical facility	7.8	3.0 to 16.5	Alonso-Echanove 2001 [23]
		8.6	1.7 to 25.8	Harries 1997 [45]
	Obstetrics and gynecology	0.2	0.006 to 1.4	Jiamrajarasangi 2005 [49]
		2.8	0.06 to 16.0	Harries 1997 [32]
Administrative facility	1.1	0.3 to 2.8	Jiamrajarasangi 2005 [49]	
	0.6	0.07 to 2.07	Jiamrajarasangi 2005 [49]	
Job title	Nurses	1.2 ^a	1.0 to 1.5	Kruuner 2001 [52]
		1.3	1.0 to 1.8	Jiamrajarasangi 2005 [49]
		2.7	1.2 to 5.4	Cuhadaroglu 2002 [40]
		3.0	0.07 to 17.1	Eyob 2002 [42]
		3.1	0.8 to 8.1	Hosoglu 2005 [47]
		7.3	4.8 to 10.7	Harries 1999 [46]
		10.4	5.1 to 18.9	Kanyerere 2003 [50]
		27.9	16.8 to 43.8	Kilinc 2002 [51]
		0.5	0.01 to 2.7	Cuhadaroglu 2002 [40]
	Doctors	2.2	1.8 to 2.6	Kruuner 2001 [52]
		6.3	0.1 to 36.7	Eyob 2002 [42]
		6.8	3.6 to 11.7	Hosoglu 2005 [47]
		10.9	5.9 to 20.7	Kilinc 2002 [51]
		4.2	2.6 to 4.3	Cuhadaroglu 2002 [40]
		4.0	1.3 to 9.3	Hosoglu 2005 [47]
	Allied health staff	25.0	14.0 to 41.5	Kilinc 2002 [51]
		7.9	0.2 to 47.8	Kanyerere 2003 [50]
		19.0	9.6 to 34.2	Harries 1999 [46]
	Paramedics	7.3	1.5 to 21.6	Eyob 2002 [42]
		7.9	2.1 to 20.9	Harries 1999 [46]
		16.4	3.3 to 50.0	Eyob 2002 [42]
	Clinical officers	5.5	0.1 to 32.0	Harries 1999 [46]
		53.0	5.6 to 256.2	Kanyerere 2003 [50]
		12.4	7.4 to 20.9	Harries 1999 [46]
	Laboratory assistants	52.2	19.4 to 120.4	Kanyerere 2003 [50]
		9.1	6.3 to 12.6	Harries 1999 [46]
		13.2	6.7 to 23.5	Kanyerere 2003 [50]
	Laboratory technicians	0.7	0.3 to 1.7	Jiamrajarasangi 2005 [49]
		8.0	1.6 to 24.1	Eyob 2002 [42]
		5.7	0.7 to 21.1	Eyob 2002 [42]
Radiology technicians	25.4	8.0 to 62.2	Eyob 2002 [42]	
	0.5	0.1 to 1.4	Jiamrajarasangi 2005 [49]	
	5.6	1.5 to 14.5	Eyob 2002 [42]	
Patient attendants				
Ward attendants				
Custodial workers				
Cleaners				
Guards, drivers				
Administrative staff				

^aIncludes nurses and laboratory staff.

^bAll risk estimates have been calculated using WHO estimates [17] for incidence of all forms of TB in the country concerned as a baseline reference.
doi:10.1371/journal.pmed.0030494.t006

occupations: radiology technicians, patient attendants, nurses, ward attendants, paramedics, clinical officers, laboratory personnel, and physicians had a high incidence of TB disease, while the incidence of TB disease was lowest in administrative staff.

Impact of Infection-Control Strategies on the Incidence of TB Infection or Disease

Most authors reported that no specific TB infection-control programs were being used in the health-care facilities where the studies were carried out. Only three studies

Table 7. Impact of Multiple Administrative, Personal, and Engineering Control Measures on Nosocomial Transmission of TB

Study: (Medical Facilities and Year Strategy Initiated); (Number of HCWs)	Preventive Strategy Used		Impact
	Epidemiological Measure in Absence of Preventive Measure	Epidemiological Measure in Presence of Preventive Measure	
	Administrative	Personal	Engineering
Harries 2002 [44], Malawi; 40 TB-care facilities established plus TB-control guidelines (1998); (n = 2,697)	(1) Priority to patients with chronic cough in OPD. (2) Rapid sputum collection, transport, and reporting. (3) Visitors kept to a minimum. (4) CXR at quiet times in the day.	(1) Proper cough hygiene. (2) Mask worn by TB patients and when undergoing surgical procedures. (3) Visitors kept to a minimum. (4) CXR at quiet times in the day.	(1) More ventilated TB wards. (2) TB patients spend more day-time outside wards when possible. (3) Windows left open most of the time.
			Incidence of TB disease before prevention ^a (1996): clinical officer 7,407 Incidence of TB disease after prevention ^a (1999): patient attendant 4,348 Incidence of TB disease after prevention ^a (1999): ward attendant 3,954 Incidence of TB disease after prevention ^a (1999): TB officer 1,785 Incidence of TB disease after prevention ^a (1999): nurses 2,060 Incidence of TB disease after prevention ^a (1999): overall 3,222
Yanal 2003 [22], Thailand; provincial referral hospital (1997–1998); (n = 1,202)	(1) Early suspicion of TB. (2) Early sputum collection and reporting. (3) Early initiation of treatments. (4) Isolation of patients with TB. (5) One-stop OPD TB service.	(1) N95 mask use by HCWs. (2) HEPA filter in TB laboratory areas.	(1) TB-isolation room in wards. (2) Maximizing ventilation in wards. (3) Class II safety cabinets in laboratory. (4) Ultra-violet germicidal irradiation system in laboratory.
			Incidence of TB disease before prevention ^a (1995–1997): all HCWs 179.21 Incidence of LTI per year before prevention (1995–1997): 9.3% (3.3 to 15.3)
Roth 2005 [21], Brazil; two hospitals with preventive measures and two without preventive measures (1998); (n = 4,868)	(1) Rapid diagnosis and treatment of TB patients. (2) Isolation of TB patients in private rooms.	(1) N95 mask use by HCWs. (2) HEPA filter in laboratory areas.	Incidence of LTI in hospitals without prevention measures (1998–1999): 16 per 1,000 person-months ^b Incidence of LTI in hospitals with prevention measures in place (1998–1999): 7.75 per 1,000 person-months ^b

^aPer 100,000 of general population per year.^bAverage value computed.NS, not significant; OPD, outpatient department.
doi:10.1371/journal.pmed.0030494.t007



Figure 3. A Young Medical Trainee Examines a Patient with Pulmonary TB at a Rural Hospital in India

In such low-income countries, more years of clinical training and greater exposure to TB patients are important risk factors for acquiring new TB infection.

doi:10.1371/journal.pmed.0030494.g003

[21,22,44] evaluated the impact of multiple infection-control strategies on the risk of TB infection or disease. Another two studies [24,48] analyzed whether a lack of personal-protection measures was associated with a risk of TB infection. One study [57] evaluated the knowledge, attitudes, and use of TB infection-control measures by HCWs.

As seen in Table 7, Harries et al. [44] evaluated the impact of multiple administrative control measures which were implemented in 40 district and mission hospitals in Malawi, following adoption of infection-control guidelines. The data were collected by interviewing HCWs and by screening the TB registers at these facilities. The study revealed that the infection-control guidelines were not uniformly implemented, and the median compliance with various measures was 76% (range 3% to 100%). There was a non-significant decrease in TB disease incidence after 1 y of implementing these measures.

The introduction of multiple administrative, personal, and engineering controls in a single hospital in Thailand [22] resulted in a significant drop in the annual incidence of LTBI in HCWs from 9.3% to 2.2%. However, the incidence of TB disease in HCWs showed a non-significant increase (from 179 to 252 per 100,000) 1–2 y after initiation of these control measures. During the course of this study [22], the proportion of HIV-positive TB patients treated at this facility increased from 3% to 57%; if there was a similar increase in HIV among HCWs, the incidence could have increased despite a fall in new infections.

In another study from Brazil [21], a cross-sectional tuberculin survey determined the baseline LTBI prevalence in four hospitals. Hospital A initiated administrative controls and provided N95 respirators for all HCWs required to enter a TB-isolation room. Hospital B had initiated administrative controls 3 mo before the baseline TST testing and, at the onset of the study, had introduced N95 respirators and had begun construction of negative-pressure isolation rooms. Hospitals C and D had no TB-control measures in place

throughout the study. Baseline TST positivity was significantly different in the four hospitals (46.7%, 69.6%, 65.8%, and 62.2% in hospitals A, B, C, and D, respectively). After 1 y, the incidence of LTBI (in initially tuberculin-negative workers) was significantly lower in hospitals A and B, which had implemented multiple infection-control measures, compared with the other two hospitals.

In a case-control study by Jelip et al. [48], HCWs with TB disease were 5.9 times (95% CI 0.76 to 46.4) more likely to have poor knowledge about TB transmission, and 4.3 times (95% CI 0.95 to 19.8) more likely to be unaware of the need for respiratory protection. In a study among medical students [24], although 90% were aware of the risk of TB transmission, only 46% reported the use of personal-protection measures. In a study from Thailand [57], although 97% of HCWs were aware of TB infection-control policies, only 52% used personal-protection measures (e.g., respirators), and only 72% implemented respiratory isolation for TB cases. Failure to use personal protection was associated with a 2.6-fold (95% CI 1.06 to 6.64) increased risk of TB disease in HCWs [46].

Discussion

Principal Findings

Our systematic review of 51 studies demonstrates that the prevalence (range 33% to 79%) and incidence (range 0.5% to 14.3% per year) of LTBI, and the attributable risk of TB disease due to nosocomial exposure (from 25 to 5,361 per 100,000 per year), were high among HCWs in LMICs. The attributable risk was higher in health-care facilities that had more TB patients per HCW. Certain work locations (in-patient TB facility, laboratory, general medicine, and emergency facilities) and occupational categories (radiology technicians, patient attendants, nurses, ward attendants, paramedics, and clinical officers) were associated with a higher risk of TB disease. Finally, there is little published evidence on the effects of infection-control measures in LMIC.

In a recent review of TB among HCWs in high-income settings, the prevalence of LTBI ranged from 5% to 55%, in different occupations [8]. The annual risk of TB infection ranged from 0.1% to 10% in studies published prior to 1995 [56], but only from 0.1% to 1.2% after the widespread introduction of infection-control measures [8]. The annual incidence of TB disease in the general population was much lower (2–55 per 100,000) in HCWs in high-income countries (excluding foreign-born HCWs) [8], than among HCWs in LMICs (69–5,780 per 100,000). Health-care facilities in LMICs had a median of 36 (range 2–2,652) HCWs per 100 TB patients treated at the facility, which is much lower than facilities in high-income countries, which have a median of 6,450 (range 100–77,000) HCWs per 100 TB patients [58]. Thus, HCWs in low-income countries are likely to experience significantly higher TB exposure (Figure 3), and it is therefore not surprising that the epidemiology of TB among HCWs is very different in high-income countries versus LMICs.

Strengths and Limitations of the Review

Our systematic review had several strengths. We used a comprehensive search strategy using multiple sources and databases to retrieve relevant studies. Two reviewers (RJ and

MP) independently selected and extracted data from the included studies. Subgroup analyses were done to minimize heterogeneity across studies. Data were pooled only when studies were reasonably consistent in their methods. However, our review had certain limitations. First, despite the comprehensive literature search, a few eligible studies were missed, because we included only English-language studies. Our literature search had identified ten non-English articles [59–68] which potentially could have been included in the final review. Of these ten papers, seven provided an abstract in English, and the summary results are presented in Table S2. As seen in Table S2, apart from one study [63], the overall results are fairly similar to the English studies included in our review. This finding, although reassuring, does not completely rule out a language bias in our review. Second, publication bias is a known problem in systematic reviews and meta-analyses. If studies with high TB rates among HCWs were more likely to have been published, this may over-estimate the TB burden among HCWs. Thirdly, there were limited data on the magnitude of TB exposure in HCWs. Most studies did not report even simple indicators of exposure, such as the number of TB patients cared for at the facility, the number of inpatient days of TB patients, or the number of TB patients per HCW. Hospital infection-control policies and programs are important, yet often unmeasured, potential confounders—and these were not described in most studies. Finally, although we pooled prevalence estimates because of methodological similarity between studies on LTBI prevalence, the pooled averages will need to be interpreted with caution because of heterogeneity in study results.

Limitations of Primary Studies Included in the Review

In addition to the limitations of the review, the primary studies included within its scope had several limitations. These limitations are discussed separately for each of the four main outcomes in our review: (1) the prevalence of LTBI; (2) the incidence of LTBI; (3) the incidence of TB disease; and (4) the impact of infection-control measures.

Prevalence of LTBI in HCWs

Studies that reported LTBI prevalence among HCWs had several limitations. The first limitation pertains to the use of the TST. The prevalence of occupational LTBI could have been over-estimated because it was based on TST. The TST detects lifetime cumulative occupational plus nonoccupational exposure to *M. tuberculosis*, as well as the effects of NTM exposure and BCG vaccination. Prevalence of NTM and its effect on TST is difficult to estimate because studies from several countries were included, and there are no data on NTM prevalence in each setting.

The practice, timing, and frequency of BCG vaccination vary widely across countries, which complicates the analyses as BCG can be an important cause of false-positive TST. The results of TST are also influenced by the type of test material (PPD), technique of reading, and definition of a positive test. Although all studies used the definition of 10 mm or more induration after 48–72 h for TST to be positive, the PPD used varied, which could reduce comparability of studies. These limitations may affect the prevalence of LTBI, but should not affect the analysis of risk factors associated with LTBI.

Recently, interferon-gamma release assays (IGRAs) have become available for the diagnosis of LTBI. In contrast to the

TST, IGRAs use antigens that are significantly more specific than PPD. Thus, IGRAs are highly specific and are therefore less likely to be affected by previous BCG vaccination and NTM exposure [69,70]. Only one study in our review used an IGRA for estimation of prevalence among HCWs. This study showed comparable LTBI prevalence using TST and IGRA, and high agreement between the test results [28].

Another major limitation of prevalence studies is the lack of concurrent data on LTBI prevalence in the population. Thus, it is not easy to determine whether HCWs had a significantly higher LTBI prevalence than the community. However, despite this limitation, our review shows that the prevalence of LTBI was lower among young medical or nursing students newly entering the health-care profession, but increased with each year of training (an index of cumulative exposure). Similarly, the prevalence of LTBI among other HCWs increased with duration of employment, again reflecting cumulative exposure. HCWs whose occupation involves closer patient contact (such as nurses) also had higher LTBI prevalence. These results indirectly suggest that nosocomial TB contributes to the burden of LTBI among HCWs.

Incidence of LTBI in HCWs

Almost all studies that estimated the incidence of new TB infections used serial tuberculin skin testing. In addition to the known limitations of TST, serial TST has additional problems such as boosting, choice between a single-step or a two-step baseline protocol, and the definitions used for conversion. Most studies followed the two-step testing protocol so as not to overestimate true LTBI incidence due to boosting. Only one study used an IGRA for estimating the rate of new TB infection [37]. This study found a higher conversion rate when an IGRA was used, raising the possibility that IGRAs may be more sensitive for recent infection than the TST [37]. This hypothesis deserves further study.

Despite the above limitations, the results suggest that HCWs have a higher risk of TB infection than the estimates of risk in the general population. The high attributable risk estimates for LTBI incidence provide the most convincing evidence for nosocomial transmission of TB in health-care settings. In these studies, more years of clinical training and greater exposure to TB patients were risk factors for new infection, and this provides additional support for nosocomial transmission.

Incidence of TB Disease in HCWs

The incidence of TB disease in HCWs was generally higher than the estimated TB rates in the general population. However, several methodological problems may affect the interpretation of these studies. HCWs may be more likely to seek medical care, and hence case-detection rates may be higher than in the general population. Our review included probable and self-reported TB cases, which also could have inflated the incidence of TB disease among HCWs. On the other hand, HCWs are less likely to develop TB because HCWs have a higher average socio-economic status, and are younger and healthier, than the general population in LMIC (i.e., healthy-worker effect) [71]. We used WHO estimates for the incidence of TB in the relevant country to ensure comparability of results across studies. Using data from a

single source ensures uniformity in determining attributable risk for different studies. However, there may be substantial regional variation of incidence within countries, and it would seem that regional estimates of TB incidence would be more valid to compare with rates of TB in a particular institution. However, the estimates of relative and attributable risk were not markedly different when we analyzed the studies using WHO data, or using local estimates provided by the authors of the study concerned.

Despite the above limitations, most studies reported higher estimates of TB disease among HCWs than in the general population, and this is suggestive of nosocomial transmission. The high rates of TB disease among young HCWs are particularly worrisome. Some of this may be explained by coinfection with HIV—particularly in countries with a very high prevalence of HIV, such as in Sub-Saharan Africa. Few studies reported the prevalence of HIV infection among HCWs, and thus the impact of HIV on TB disease among HCWs could not be addressed.

Molecular studies involving DNA fingerprinting could provide confirmatory evidence of nosocomial transmission, as they have in high-income countries [72]. However, our literature search revealed only a single case report [73] from a LMIC in which molecular methods were used to confirm nosocomial transmission of *M. tuberculosis* to a nurse.

Impact of Infection-Control Measures on TB in HCWs

Only three studies [21,22,44] in this review assessed the impact of infection-control measures on reducing the risk of TB in HCWs. Thus, it is difficult to make any inferences regarding effectiveness of control measures. One study determined that administrative measures had little impact on the development of TB disease, but the study considered a period of only 1 y—a relatively short interval to detect changes in this outcome [44]. Two studies that measured the incidence of TB infection detected significant reduction within 1 y following the introduction of multiple infection-control measures [22,31]. In one of these studies [22], the incidence of active disease in HCWs actually increased over the study period, but this could have been due to an increasing HIV prevalence in HCWs, which was not measured. The other study [31] compared TB incidence rates in hospitals with different infection-control policies, and other differences identified between the hospitals could have confounded the estimates. Taken together, the limited available evidence suggests that a reduction in the risk of TB infection is possible with simple administrative controls, but this needs to be evaluated in larger, better-controlled studies.

In summary, there is consistent epidemiologic evidence that TB is an important occupational disease in HCWs. There is clear evidence of heavy exposure, with little or no infection-control measures in place. Thus, it is not surprising that there is consistent evidence of excess prevalence and incidence of TB infection, as well as a higher incidence of TB disease among HCWs than in the general populations in the same LMICs. Although estimates vary widely, infection and disease are roughly correlated with indicators of exposure—including more years of work, or clinical training, and work that has been identified as high risk among HCWs in high-income countries. Finally, there is evidence, albeit limited and weak, that the incidence of infection drops after the implementation of infection-control measures. This epi-

demiological evidence implies that a substantial proportion of LTBI and TB disease in the HCWs in LMICs is the result of nosocomial TB transmission.

Conclusions

Our review presents fairly strong evidence that nosocomial TB is an important occupational problem among HCWs in LMICs, and reduction of that risk should be a priority. Currently available evidence is limited, but it suggests that relatively simple interventions, such as early diagnosis of TB, segregation of infectious TB patients, or education and training of HCWs, might be effective. Additional low-cost measures could include engineering controls such as exhaust ventilation, improved natural ventilation, or sunlight [74]. However, well-designed field studies evaluating the cost, feasibility, and effectiveness of these interventions in resource-limited settings are urgently needed.

There are several important reasons as to why nosocomial transmission of TB should be addressed in LMICs. First, occupational TB can lead to the loss of skilled workers, and this can adversely impact health-care services in the long run. Second, transmission of TB can have serious, and even fatal, consequences for patients and HCWs. This is particularly true with MDR-TB strains, and in patient populations with high HIV seroprevalence. Hospitals have been shown to be important focal points of MDR-TB transmission, with explosive outbreaks, and associated with high mortality. Third, implementation of effective TB infection control can promote awareness of TB, and the adoption of improved practices for the diagnosis and treatment of TB, particularly in the private health sector. Low-cost administrative interventions are feasible and, if implemented, should require minimal resources. Given the evidence summarized in this review, national TB-control programs and public health agencies in LMICs must begin to address nosocomial TB transmission as an integral part of their TB-control efforts. HCWs are essential in the fight against TB, and their health needs to be protected as well as that of patients with TB. With the recent emergence of extensively drug-resistant tuberculosis (XDR-TB), the need to implement infection-control measures has been reemphasized by global agencies such as the WHO and the Stop TB Partnership [78]. Efforts are ongoing to update existing infection-control guidelines in the wake of XDR-TB, and to develop programs that are suitable for resource-limited countries. We strongly support these initiatives and call for more resources and partnerships to tackle the chronically neglected problem of nosocomial TB in low-income countries.

Supporting Information

Table S1. Strategy Used to Search Pubmed to Identify Studies for This Systematic Review

Found at doi:10.1371/journal.pmed.0030494.st001 (38 KB DOC).

Table S2. Summary of Non-English Studies That Were Excluded from the Review (Seven Studies where an English Abstract Was Available)

Found at doi:10.1371/journal.pmed.0030494.st002 (47 KB DOC).

Acknowledgments

Author contributions. RJ, AR, DM, and MP conceived and designed the study. RJ and MP collected, analyzed, and interpreted the data. RJ and MP drafted the manuscript. RJ, AR, DM, and MP performed critical revision of the manuscript.

References

- Fennelly KP, Iseman MD (1999) Health care workers and tuberculosis: The battle of a century. *Int J Tuberc Lung Dis* 3: 363–364.
- Sepkowitz KA (1996) Tuberculin skin testing and the health care worker: Lessons of the Prophit Survey. *Tuberc Lung Dis* 77: 81–85.
- Harries AD, Maher D, Nunn P (1997) Practical and affordable measures for the protection of health care workers from tuberculosis in low-income countries. *Bull World Health Organ* 75: 477–489.
- Jensen PA, Lambert LA, Iademarco MF, Ridzon R (2005) Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health-care settings, 2005. *MMWR Recomm Rep* 54: 1–141.
- Pearson ML, Jereb JA, Frieden TR, et al. (1992) Nosocomial transmission of multidrug-resistant *Mycobacterium tuberculosis*. A risk to patients and health care workers. *Ann Intern Med* 117: 191–196.
- Sague C, Dooley S, Hutton M, et al. (1991) Nosocomial transmission of multidrug-resistant tuberculosis MDR-TB in an urban hospital. 31st Interscience Conference on Antimicrobial Agents and Chemotherapy; 29 September to 2 October 1991; Chicago, Illinois, United States. 140 p.
- Fella P, Rivera P, Hale M, Squires K, Sepkowitz K (1995) Dramatic decrease in tuberculin skin test conversion rate among employees at a hospital in New York City. *Am J Infect Control* 23: 352–356.
- Seidler A, Nienhaus A, Diel R (2005) Review of epidemiological studies on the occupational risk of tuberculosis in low-incidence areas. *Respiration* 2: 431–446.
- Jones-Lopez EC, Ellner JJ (2005) Tuberculosis infection among HCWs. *Int J Tuberc Lung Dis* 9: 591.
- van Gorkom J (1999) Nosocomial transmission in low-income countries. *Int J Tuberc Lung Dis* 3: 1144–1145.
- Whitehead M, Dahlgren G, Evans T (2001) Equity and health sector reforms: Can low-income countries escape the medical poverty trap? *Lancet* 358: 833.
- Makinen M, Waters H, Rauch M, et al. (2000) Inequalities in health care use and expenditures: Empirical data from eight developing countries and countries in transition. *Bull World Health Organ* 78: 55.
- Pai M, Kalantri SP, Aggarwal AN, Menzies D, Blumberg HM (2006) Nosocomial tuberculosis in India. *Emerg Infect Dis* 12: 1311–1318.
- World Bank Country Classification (2005). Available: <http://www.worldbank.org/data/countryclass/classgroups.htm>. Accessed 18 November 2006.
- Targeted tuberculin testing and treatment of latent tuberculosis infection (2000) American Thoracic Society. *MMWR Recomm Rep* 49: 1–51.
- Menzies D (1999) Interpretation of repeated tuberculin tests. Boosting, conversion, and reversion. *Am J Respir Crit Care Med* 159: 15–21.
- Global Tuberculosis Database 2006. Available: http://www.who.int/tb/country/global_tb_database/en/index.html. Accessed 18 November 2006.
- Maciel ELN, Viana MC, Zeitoun RCG, Ferreira I, Fregona G, et al. (2005) Prevalence and incidence of *Mycobacterium tuberculosis* infection in nursing students in Vitoria, Espirito Santo. *Rev Soc Bras Med Trop* 38: 469–472.
- Bonifacio N, Saito M, Gilman RH, et al. (2002) High risk for tuberculosis in hospital physicians, Peru. *Emerg Infect Dis* 8: 747–748.
- Levy MZ, Medeiros EA, Shang N, et al. (2005) TST reversion in a BCG-revaccinated population of nursing and medical students, Sao Paulo, Brazil, 1997–2000. *Int J Tuberc Lung Dis* 9: 771–776.
- Roth VR, Garrett DO, Laserson KF, et al. (2005) A multicenter evaluation of tuberculin skin test positivity and conversion among health care workers in Brazilian hospitals. *Int J Tuberc Lung Dis* 9: 1335–1342.
- Yanai H, Limpakarnjanarat K, Uthairoravit W, Mastro TD, Mori T, et al. (2003) Risk of *Mycobacterium tuberculosis* infection and disease among health care workers, Chiang Rai, Thailand. *Int J Tuberc Lung Dis* 7: 36–45.
- Alonso-Echanove J, Granich RM, Laszlo A, et al. (2001) Occupational transmission of *Mycobacterium tuberculosis* to health care workers in a university hospital in Lima, Peru. *Clin Infect Dis* 33: 589–596.
- Teixeira EG, Menzies D, Comstock GW, et al. (2005) Latent tuberculosis infection among undergraduate medical students in Rio de Janeiro State, Brazil. *Int J Tuberc Lung Dis* 9: 841–847.
- Golchin M, Rostami M (2005) Tuberculin test in nursing and human-sciences students. *J Res Med Sci* 10: 172–176.
- Kayanja HK, Debanne S, King C, Whalen CC (2005) Tuberculosis infection among health care workers in Kampala, Uganda. *Int J Tuberc Lung Dis* 9: 686–688.
- Silva VM, Cunha AJ, Oliveira JR, et al. (2000) Medical students at risk of nosocomial transmission of *Mycobacterium tuberculosis*. *Int J Tuberc Lung Dis* 4: 420–426.
- Pai M, Gokhale K, Joshi R, et al. (2005) *Mycobacterium tuberculosis* infection in health care workers in rural India—Comparison of a whole-blood interferon gamma assay with tuberculin skin testing. *JAMA* 293: 2746–2750.
- Naidoo S, Mahommed A (2002) Knowledge, attitudes, behaviour and prevalence of TB infection among dentists in the western Cape. *SADJ* 57: 476–478.
- Kassim S, Zuber P, Wiktor SZ, et al. (2000) Tuberculin skin testing to assess the occupational risk of *Mycobacterium tuberculosis* infection among health care workers in Abidjan, Cote d'Ivoire. *Int J Tuberc Lung Dis* 4: 321–326.
- Garcia-Garcia ML, Jimenez-Corona A, Jimenez-Corona ME, et al. (2001) Factors associated with tuberculin reactivity in two general hospitals in Mexico. *Infect Control Hosp Epidemiol* 22: 88–93.
- Orrett FA (2000) Prevalence of tuberculin skin test reactivity among health care workers at a teaching hospital in Trinidad. *Clin Microbiol Infect* 6: 45–48.
- Keskiner R, Ergonul O, Demiroglu Z, Eren S, Baykam N, et al. (2004) Risk of tuberculous infection among healthcare workers in a tertiary-care hospital in Ankara, Turkey. *Infect Control Hosp Epidemiol* 25: 1067–1071.
- Molina-Gamboa J, Fivera-Morales I, Ponce-de-Leon-Rosales S (1994) Prevalence of tuberculin reactivity among healthcare workers from a Mexican hospital. *Infect Control Hosp Epidemiol* 15: 319–320.
- Do AN, Limpakarnjarat K, Uthairoravit W, et al. (1999) Increased risk of *Mycobacterium tuberculosis* infection related to the occupational exposures of health care workers in Chiang Rai, Thailand. *Int J Tuberc Lung Dis* 3: 377–381.
- Silva VM, Cunha AJ, Kritski AL (2002) Tuberculin skin test conversion among medical students at a teaching hospital in Rio de Janeiro, Brazil. *Infect Control Hosp Epidemiol* 23: 591–594.
- Pai M, Joshi R, Dogra S, Mendiratta DK, Narang P, et al. (2006) Serial testing of health care workers for tuberculosis infection using interferon- γ assay. *Am J Respir Crit Care Med* 174: 349–355.
- Babus V (1997) Tuberculosis morbidity risk in medical nurses in specialized institutions for the treatment of lung diseases in Zagreb. *Int J Tuberc Lung Dis* 1: 254–258.
- Balt E, Durrheim DN, Weyer K (1998) Nosocomial transmission of tuberculosis to health care workers in Mpumalanga. *S Afr Med J* 88: 1363–1366.
- Cuhadaroglu C, Erelel M, Tabak L, Kilicaslan Z (2002) Increased risk of tuberculosis in health care workers: A retrospective survey at a teaching hospital in Istanbul, Turkey. *BMC Infect Dis* 2: 14.
- Dimitrova B, Hutchings A, Atun R, et al. (2005) Increased risk of tuberculosis among health care workers in Samara Oblast, Russia: Analysis of notification data. *Int J Tuberc Lung Dis* 9: 43–48.
- Eyob G, Gebeyhu M, Goshu S, Girma M, Lemma E, et al. (2002) Increase in tuberculosis incidence among the staff working at the Tuberculosis Demonstration and Training Centre in Addis Ababa, Ethiopia: A retrospective cohort study (1989–1998). *Int J Tuberc Lung Dis* 6: 85–88.
- Gopinath KG, Siddique S, Kirubakaran H, Shanmugam A, Mathai E, et al. (2004) Tuberculosis among healthcare workers in a tertiary-care hospital in South India. *J Hosp Infect* 57: 339–342.
- Harries AD, Hargreaves NJ, Gausi F, Kwanjana JH, Salaniponi FM (2002) Preventing tuberculosis among health workers in Malawi. *Bull World Health Organ* 80: 526–531.
- Harries AD, Kamenya A, Namarika D, et al. (1997) Delays in diagnosis and treatment of smear-positive tuberculosis and the incidence of tuberculosis in hospital nurses in Blantyre, Malawi. *Trans R Soc Trop Med Hyg* 91: 15–17.
- Harries AD, Nyirenda TE, Banerjee A, Boeree MJ, Salaniponi FM (1999) Tuberculosis in health care workers in Malawi. *Trans R Soc Trop Med Hyg* 93: 32–35.
- Hosoglu S, Tanrikulu AC, Dagli C, Akalin S (2005) Tuberculosis among health care workers in a short working period. *Am J Infect Control* 33: 23–26.
- Jelip J, Mathew GG, Yusin Y, et al. (2004) Risk factors of tuberculosis among health care workers in Sabah, Malaysia. *Tuberculosis (Edinb)* 84: 19–23.
- Jiamjarasrangi W, Hirunsuthikul N, Kamolratanakul P (2005) Tuberculosis among health care workers at King Chulalongkorn Memorial Hospital, 1988–2002. *Int J Tuberc Lung Dis* 9: 1253–1258.
- Kanyerere HS, Salaniponi FM (2003) Tuberculosis in health care workers in a central hospital in Malawi. *Int J Tuberc Lung Dis* 7: 489–492.
- Kilinc O, Ucan ES, Cakan MD, et al. (2002) Risk of tuberculosis among healthcare workers: Can tuberculosis be considered as an occupational disease? *Respir Med* 96: 506–510.
- Kruuner A, Danilovitch M, Pehme L, Laisaar T, Hoffner SE, et al. (2001) Tuberculosis as an occupational hazard for health care workers in Estonia. *Int J Tuberc Lung Dis* 5: 170–176.
- Skodric V, Savic B, Jovanovic M, et al. (2000) Occupational risk of tuberculosis among health care workers at the Institute for Pulmonary Diseases of Serbia. *Int J Tuberc Lung Dis* 4: 827–831.
- Wilkinson D, Gilks CF (1998) Increasing frequency of tuberculosis among staff in a South African district hospital: Impact of the HIV epidemic on the supply side of health care. *Trans R Soc Trop Med Hyg* 92: 500–502.
- Rao KG, Aggarwal AM, Behera D (2004) Tuberculosis among physicians in training. *Int J Tuberc Lung Dis* 8: 1392–1394.
- Naidoo S, Jinabhai CC (2006) TB in health care workers in KwaZulu Natal, South Africa. *Int J Tuberc Lung Dis* 10: 676–682.
- Luksamijarulkul P, Supapvanit C, Loosereewanich P, Aiumlaor P (2004) Risk assessment towards tuberculosis among hospital personnel: Administrative control, risk exposure, use of protective barriers and microbial air quality. *Southeast Asian J Trop Med Public Health* 35: 1005–1011.
- Menzies D, Fanning A, Yuan L, Fitzgerald M (1995) Current concepts—Tuberculosis among health-care workers. *N Engl J Med* 332: 92–98.
- Aksytina LP, Leonov IV (1998) Tuberculosis as hospital acquired infection. *Probl Tuberk* 1: 5–7.
- Priimak AA, Plotnikova LM (1992) Tuberculosis morbidity in medical workers and the measures for their social protection. *Probl Tuberk* (11–12): 24–26.

61. Protsiuk RG, Serdechnyi VI (1996) The tuberculosis incidence among medical workers in Donetsk Province. *Lik Sprava* (1–2): 191–193.
62. Olona M, Campins M, Oltra C, Vague J (1989) Prevalence of tuberculosis infections among hospital staff. *Rev Clin Esp* 185: 44–45.
63. Ostrosky-Zeichner L, Rangel-Frausto MS, Garcia-Romero E, Vazquez A, Ibarra MJ, et al. (2000) Tuberculosis in health personnel: Importance of surveillance and control programs. *Salud Publica Mex* 42: 48–52.
64. Fimenez Martinez I, Sanchez Mozo I, Cembero Gil C, Dominguez Carmona M (1976) Study of the Mantoux reaction in a hospital institution. *Rev Med Univ Navarra* 20: 83–97.
65. Houssine El, Laraqui C, Laraqui O, El Kabouss Y, et al. (2004) Assessment of the risk of tuberculosis among health care personnel in Meknes. *Sante* 14: 167–171.
66. Laraqui CH, Ottmani S, Hammou MA, Bencheikh N, Mahjour J (2001) Tuberculosis in public health workers in Morocco. *Int J Tuberc Lung Dis* 5: 939–945.
67. Kohout J (1999) Occupational tuberculosis in health professionals of the West Bohemian Region. *Prakt Lek* 79: 646–648.
68. Kuklova D, Kneidlova M, Mencik M, Naus A (1989) The incidence of occupational diseases in the health services within the catchment area of the Clinic for Occupational Diseases of the School of Medicine in Prague. *Vnitr Lek* 35: 472–478.
69. Pai M, Kalantri S, Dheda K (2006) New tools and emerging technologies for the diagnosis of tuberculosis: Part I. Latent tuberculosis. *Expert Rev Mol Diagn* 6: 413–422.
70. Pai M, Riley LW, Colford JM Jr (2004) Interferon-gamma assays in the immunodiagnosis of tuberculosis: A systematic review. *Lancet Infect Dis* 4: 761–776.
71. Li CY, Sung FC (1999) A review of the healthy worker effect in occupational epidemiology. *Occup Med (Lond)* 49: 225–229.
72. Seidler A, Nienhaus A, Diel R (2004) The transmission of tuberculosis in the light of new molecular biological approaches. *J Occup Environ Med* 61: 96–102.
73. Wilkinson D, Crump J, Pillay M, Sturm AW (1997) Nosocomial transmission of tuberculosis in Africa documented by restriction fragment length polymorphism. *Trans R Soc Trop Med Hyg* 91: 318.
74. Granich R, Binkin NJ, Jarvis WR, Simone PM, Rieder HL, et al. (1999) Guidelines for the prevention of tuberculosis in health care facilities in resource-limited settings. WHO/CDS/TB/99.269. Geneva, Switzerland: World Health Organization. 51 p. Available: <http://www.who.int/docstore/gtb/publications/healthcare/PDF/WHO99-269.pdf>. Accessed 23 November 2006.
75. Palwatwichai A (2001) Tuberculosis in Thailand. *Respirology* 6: 65–70.
76. Ministério de Saude (1994) Control da tuberculose: Uma proposta de integração ensino-servico, 4a edicao. Available: http://www.sbp.com.br/img/Nota%20tecnica2_%20Dose_BCG_26%2005%202006%20FINAL%202006.pdf. Accessed 23 November 2006.
77. Chadha VK, Kumar P, Jagannatha PS, et al. (2005) Average annual risk of tuberculosis infection in India. *Int J Tuberc Lung Dis* 9: 116–118.
78. Ravighione M (2006) XDR-TB. Entering the post-antibiotic era? *Int J Tuberc Lung Dis* 10: 1185–1187.

Editors' Summary

Background. One third of the world's population is infected with *Mycobacterium tuberculosis*, the bacterium that causes tuberculosis (TB). In many people, the bug causes no health problems—it remains latent. But about 10% of infected people develop active, potentially fatal TB, often in their lungs. People with active pulmonary TB readily spread the infection to other people, including health-care workers (HCWs), in small airborne droplets produced when they cough or sneeze. In high-income countries such as the US, guidelines are in place to minimize the transmission of TB in health-care facilities. Administrative controls (for example, standard treatment plans for people with suspected or confirmed TB) aim to reduce the exposure of HCWs to people with TB. Environmental controls (for example, the use of special isolation rooms) aim to prevent the spread and to reduce the concentration of infectious droplets in the air. Finally, respiratory-protection controls (for example, personal respirators for nursing staff) aim to reduce the risk of infection when exposure to *M. tuberculosis* is unavoidably high. Together, these three layers of control have reduced the incidence of TB in HCWs (the number who catch TB annually) in high-income countries.

Why Was This Study Done? But what about low- and middle-income countries (LMICs) where more than 90% of the world's cases of TB occur? Here, there is little money available to implement even low-cost strategies to reduce TB transmission in health-care facilities—so how important an occupational disease is TB in HCWs in these countries? In this study, the researchers have systematically reviewed published papers to find out the incidence and prevalence (how many people in a population have a specific disease) of active TB and latent TB infections (LTBIs) in HCWs in LMICs. They have also investigated whether any of the preventative strategies used in high-income countries have been shown to reduce the TB burden in HCWs in poorer countries.

What Did the Researchers Do and Find? To identify studies on TB transmission to HCWs in LMICs, the researchers searched electronic databases and journals, and also contacted experts on TB transmission. They then extracted and analyzed the relevant data on TB incidence, prevalence, risk factors, and control measures. Averaged-out over the 51 identified studies, 54% of HCWs had LTBI. In most of the studies, increasing age and duration of employment in health-care facilities, indicating a longer cumulative exposure to infection, was associated with a higher prevalence of LTBI. The same trend was seen in a subgroup of medical and nursing students. After accounting for the incidence of TB

in the relevant general population, the excess incidence of TB in the different studies that was attributable to being a HCW ranged from 25 to 5,361 cases per 100,000 people per year. In addition, a higher risk of acquiring TB was associated with working in specific locations (for example, inpatient TB facilities or diagnostic laboratories) and with specific occupations, including nurses and radiology attendants; most of the health-care facilities examined in the published studies had no specific TB infection-control programs in place.

What Do These Findings Mean? As with all systematic reviews, the accuracy of these findings may be limited by some aspects of the original studies, such as how the incidence of LTBI was measured. In addition, the possibility that the researchers missed some relevant published studies, or that only studies where there was a high incidence of TB in HCWs were published, may also affect the findings of this study. Nevertheless, they suggest that TB is an important occupational disease in HCWs in LMICs and that the HCWs most at risk of TB are those exposed to the most patients with TB. Reduction of that risk should be a high priority because occupational TB leads to the loss of essential, skilled HCWs. Unfortunately, there are few data available to indicate how this should be done. Thus, the researchers conclude, well-designed field studies are urgently needed to evaluate whether the TB-control measures that have reduced TB transmission to HCWs in high-income countries will work and be affordable in LMICs.

Additional Information. Please access these Web sites via the online version of this summary at <http://dx.doi.org/10.1371/journal.pmed.0030494>.

- US National Institute of Allergy and Infectious Diseases patient fact sheet on tuberculosis
- US Centers for Disease Control and Prevention information for patients and professionals on tuberculosis
- MedlinePlus encyclopedia entry on tuberculosis
- NHS Direct Online, from the UK National Health Service, patient information on tuberculosis
- US National Institute for Occupational Health and Safety, information about tuberculosis for health-care workers
- American Lung Association information on tuberculosis and health-care workers