

## REVIEW ARTICLE OPEN

## Tumor evolution and chemoresistance in ovarian cancer

Soochi Kim<sup>1,2</sup>, Youngjin Han<sup>2,3</sup>, Se Ik Kim<sup>4</sup>, Hee-Seung Kim<sup>4</sup>, Seong Jin Kim<sup>5,6</sup> and Yong Sang Song<sup>2,3,4,7</sup>

Development of novel strategies to overcome chemoresistance is central goal in ovarian cancer research. Natural history of the cancer development and progression is being reconstructed by genomic datasets to understand the evolutionary pattern and direction. Recent studies suggest that intra-tumor heterogeneity (ITH) is the main cause of treatment failure by chemoresistance in many types of cancers including ovarian cancer. ITH increases the fitness of tumor to adapt to incompatible microenvironment. Understanding ITH in relation to the evolutionary pattern may result in the development of the innovative approach based on individual variability in the genetic, environment, and life style. Thus, we can reach the new big stage conquering the cancer. In this review, we will discuss the recent advances in understanding ovarian cancer biology through the use of next generation sequencing (NGS) and highlight areas of recent progress to improve precision medicine in ovarian cancer.

*npj Precision Oncology* (2018)2:20 ; doi:10.1038/s41698-018-0063-0

## INTRODUCTION

Most ovarian cancer arises from the epithelium of the ovary and fallopian tube. Cancers developing from the germ cells (eggs) or stromal cells are less common. The epithelial ovarian cancer (EOC) is not a single disease entity, rather composed of several histological subtypes, with each subtype characterized by different microscopic appearances and biological and genetic backgrounds.<sup>1</sup> This diversity extends to various clinical outcomes of the disease, where patients with different histological subtypes respond differently to the same treatments and also have different prognoses. Ovarian cancer has long been classified into four representative histological subtypes, serous, endometrioid, mucinous, and clear cell adenocarcinomas.<sup>2</sup> The WHO classification was recently revised and is valid since 2014.<sup>3</sup> In addition, the rate of these histological subtypes of ovarian cancer are also different across racial and ethnic groups.<sup>4</sup> Thus, stratification of the ovarian cancer according to their histological subtypes and tumor stage, as well as the ethnicity, are essential consideration for the decision of treatment methods.

Stage of the disease, which is determined surgically, is the critical determinant of ovarian cancer prognosis.<sup>1</sup> The 5-year overall survival rate is significantly different between International Federation of Gynaecological Oncology (FIGO) stage I and stage III/IV cancers, nearly 90% and around 10–40%, respectively.<sup>5</sup> There are three broad classification of prognostic factors in ovarian cancer. The tumor, the patient and lastly, the clinical interventions. The tumor itself can be sub-classified by the stage, grade, and histological subtypes. The patient with ovarian cancer are sub-classified by their age, physical, and socioeconomic status. The clinical interventions can be sub-classified by the quality of surgically removed tumor and also whether the patient have received the post-operative adjuvant chemotherapy.<sup>5</sup>

The standard treatment of advanced EOC is based on the maximum debulking surgery, followed by platinum-based and taxane-based chemotherapy, which remained the same over the past three decades.<sup>6</sup> Meanwhile, many anticancer agents, including molecular-targeted agents, and combination therapies have been developed and validated clinically. However, the overall survival rate has not been improved significantly due to chemoresistance.<sup>7</sup> Therefore, understanding the underlying molecular mechanisms associated with the chemoresistance is the critical step to improve treatment results in ovarian cancer. In this review, we will focus on the current treatment and prognosis of EOC, firstly. Next, we will explore novel strategies to overcome chemoresistance in ovarian cancer, focusing on anticancer strategy targeting tumoral evolution and intratumor heterogeneity. The recent study results of next generation sequencing (NGS) techniques will be reviewed. We will discuss the possible changes in care to pave the path towards precision medicine.

## CURRENT TREATMENT IN EOC

In 1976, the report by Wiltshaw and Kroner on the efficacy of cisplatin in ovarian cancer opened the modern era of platinum-based combination chemotherapy.<sup>8</sup> In the 1990s, Paclitaxel was introduced into platinum-based treatment, and significantly improved the progression-free survival and overall survival of the patients with advanced-stage ovarian cancer.<sup>9</sup> Current standard of care for the patients with advanced-stage ovarian cancer involves primary debulking surgery followed by the platinum-based and taxane-based combination chemotherapy including Taxol and carboplatin.<sup>6,10</sup> The concept of primary debulking surgery is diminishing the residual tumor to the minimum where adjuvant chemotherapy will be optimally effective.<sup>6</sup> Cytoreductive surgery is initially recommended for

<sup>1</sup>Seoul National University Hospital Biomedical Research Institute, Seoul 03080, Republic of Korea; <sup>2</sup>Cancer Research Institute, Seoul National University College of Medicine, Seoul 03080, Republic of Korea; <sup>3</sup>WCU Biomodulation, Department of Agricultural Biotechnology, Seoul National University, Seoul 03080, Republic of Korea; <sup>4</sup>Department of Obstetrics and Gynecology, Seoul National University College of Medicine, Seoul 03080, Republic of Korea; <sup>5</sup>Precision Medicine Research Center, Advanced Institutes of Convergence Technology, Seoul National University, Suwon, Gyeonggi-do 16229, Republic of Korea; <sup>6</sup>Department of transdisciplinary Studies, Graduate School of Convergence Science and Technology, Seoul National University, Suwon, Gyeonggi-do 16229, Republic of Korea and <sup>7</sup>Interdisciplinary Program in Cancer Biology, Seoul National University College of Medicine, Seoul 03080, Republic of Korea

Correspondence: Yong Sang. Song (ysson@snu.ac.kr)

Received: 3 November 2017 Revised: 6 August 2018 Accepted: 9 August 2018

Published online: 17 September 2018

the patients with clinical stage II–IV disease.<sup>11</sup> An alternative option is neoadjuvant chemotherapy followed by interval debulking surgery, which has been shown to be safer and better tolerated than primary debulking surgery for the patients with more advanced disease.<sup>10</sup> Currently, neoadjuvant chemotherapy is considered for the patients with advanced-stage disease expected in surgery or physically poor surgical candidates.<sup>10,12–14</sup> However, it remains still controversial in selection of the patients, who will benefit from a neoadjuvant chemotherapy.<sup>15,16</sup> Further prospective studies on this issue are warranted.

Platinum compound is the commonly selected chemo-agent for the primary treatment in ovarian cancer. Cisplatin is the very first platinum compound used as the primary treatment in ovarian cancer; however, it has a dose-limiting toxicity including minor symptoms like nausea and serious injuries on kidney and peripheral neuropathy. To overcome the toxicity associated with the use of cisplatin, the organo-platinum analogues of cisplatin, such as carboplatin has been developed and replaced cisplatin for primary chemotherapy in ovarian cancer.<sup>17</sup> The following regimen was recommended by Gynecologic Cancer Inter-Group (GCIg) since 2004, and paclitaxel and carboplatin are intravenously administered every 3 weeks for six cycles in most of patients with advanced-stage ovarian cancer.<sup>18,19</sup> Drug response or resistance to the chemotherapy is partly related to histological subtypes in EOC patients. High grade serous ovarian cancer patients respond well to platinum-based chemotherapy, whereas clear cells and mucinous types are remarkably platinum resistant.<sup>20–22</sup> After the first round of chemotherapy, ovarian cancer patients can be sub-categorized based on the time period in which the disease progresses. NCCN guidelines in ovarian cancer defines platinum-sensitive recurrence (PSR) and platinum-resistant recurrence (PRR) based on the cut-off value of 6 months from the last day of first round chemotherapy. If patients relapse 6 months or more after the first round of platinum-based chemotherapy, they are considered PSR.<sup>11</sup> Many researchers designed ovarian cancer nomogram to help both patients and clinicians to predict platinum sensitivity and overall patient survival.<sup>23–25</sup> However, current limitation is that the ovarian cancer nomogram is based on data from clinicopathological parameters including age, disease stage, grade, histology, and residual disease. Future studies and development of nomograms including NGS data would improve both prediction and treatment.

Dualistic model of ovarian carcinogenesis based on morphology, molecular studies distinguishing histological subtypes of ovarian cancer by specific genetic alterations, and the mouse model of ovarian carcinogenesis attributed to our current understanding of ovarian cancer origin.<sup>26,27</sup> However, the origins of ovarian cancer are still on debate, primarily because the precursor lesions of ovarian cancer are largely unknown. Fallopian tube has been suggested as origin of high-grade serous ovarian cancer (HGSOC), the most common histologic subtype of EOCs with poor outcomes. Previous studies about the prevalence of occult ovarian and fallopian tube cancers in women with *BRCA1/2* germline mutations who received risk-reducing salpingo-oophorectomy indicated the fimbriae of the fallopian tube as the potential precursor lesions of HGSOCs.<sup>28–30</sup> Single cell epithelial layer with *TP53* mutation and serous tubal intraepithelial carcinoma (STIC) have been observed in patients with HGSOC.<sup>31,32</sup> Adding to the immunohistochemistry, targeted sequencing also revealed that the fallopian tubal lesions harbor the same *TP53* mutations as surrounding invasive carcinomas.<sup>33,34</sup> Mouse model of ovarian cancer further support this notion that HGSOCs arise from the fallopian tube.<sup>35</sup> Recent evolutionary analyses using NGS technique revealed that both *TP53* mutation and STICs are precursors of ovarian cancer.<sup>36</sup>

## Clinico-pathological and molecular heterogeneity in ovarian cancer


Kobel et al. reported clinico-pathological and molecular heterogeneity of ovarian cancer in 2008. They analyzed the expression of potential molecular markers in 500 ovarian cancer patients across all histological subtypes. Interestingly, analysis of the entire cohort identified 10 potential molecular markers which were differentially expressed between early and advanced stage ovarian cancer. However, no markers remained to be different when the analysis was restricted to each individual subtypes.<sup>37</sup> The most predominant histological subtype in ovarian cancer, is serous adenocarcinoma.<sup>4</sup> It has been documented that there are disparities among racial and ethnic groups with respect to the rate of histological subtypes.<sup>38</sup> Particularly, the prevalence of clear cell adenocarcinoma is higher in Japan (Fig. 1a). Takenaka and his colleagues examined the clinico-pathological heterogeneity in 72 Japanese ovarian cancer patients, using targeted deep sequencing.<sup>39</sup> Interestingly, the most commonly mutated genes in the entire cohort were *TP53*, *PIK3CA*, and *KRAS*, but there were disparities among histological subtypes with respect to the frequency of these genetic mutations. This is consistent with previous report, which demonstrated that *TP53* and *BRCA1/2* mutations are molecular and genetic signatures of serous histological subtype, and *PIK3CA* and *KRAS* are dominant mutations in clear cell and mucinous subtypes, respectively.<sup>7,40–42</sup> Endometrioid histological subtype of ovarian cancer are histologically and molecularly similar to those of endometrial cancer. They share mutations in *PTEN*, *PIK3CA*, *ARID1A*, *KRAS*, *PPP2R1*, and *CTNNB1*<sup>43–45</sup> but are genomically different as the frequency of *PTEN* and *CTNNB1* mutations are significantly different between the two malignancies.<sup>46</sup>

Recent exome sequencing study in mucinous ovarian cancer further highlighted that ovarian cancer is a heterogeneous group of tumors, and its mutation profiles are distinct across histological subtypes.<sup>47</sup> This study supports that molecular features of histological subtypes are distinct (Fig. 1b) and should be considered as a group of diseases sharing the same tumor site, the ovary or fallopian tube.

Recently, our group explored the molecular heterogeneity in ovarian cancer in a number of methods. Primary culture of cellular fraction of ascites derived from ovarian cancer patients were heterogeneous and presented morphologically distinct spheres.<sup>48</sup> This is consistent with data from cancer cell lines, in vivo xenograft model and patient-derived ascites.<sup>49,50</sup> The clinical significance of morphologically heterogeneous tumor spheres is unclear in ovarian cancer model. However, tumor spheres have been increasingly studied in a number of cancer models, including breast cancer,<sup>51</sup> glioblastoma,<sup>52</sup> and pancreatic cancer.<sup>53</sup> Heterogeneity in the tumor microenvironment is also known to be an important factor contributing to tumoral heterogeneity. Ascites in ovarian cancer patients created heterogeneous pro-inflammatory tumor microenvironment. At the same time, ascites increased invasiveness in a selective subset of ovarian cancer cell lines, expressing membrane-bound IL-6 receptor.<sup>54</sup> In addition, pro-inflammatory M1 macrophage but not M2 macrophage increased ovarian cancer metastasis through NF $\kappa$ B activation.<sup>55</sup> These findings support the presence of clinico-pathological and molecular heterogeneity in ovarian cancer caused by stromal and inflammatory cells in tumor microenvironment. Understanding clinico-pathological and molecular heterogeneity in ovarian cancer are important to improve precision medicine.<sup>56</sup>

## Spatial and temporal heterogeneity in ovarian cancer

Intra-tumor heterogeneity (ITH) has been documented for many decades, initially in the late 1800s by Rudolf Virchow, from a morphological aspect.<sup>57</sup> The development of karyotyping and cytogenetic technologies in the 1980s and microarray technologies in the late 1990s led to numerous studies showing tumoral



**Fig. 1** Histological and molecular heterogeneity in ovarian cancer. **a** The rate of histological subtypes in Caucasian (above) and Japanese (below). **b** Most frequently mutated genes identified by NGS technology according to histological subtypes in ovarian cancer

Table 1. Potential druggable targets identified by NGS technology in HGSOc				
Samples	Druggable targets		Genomic analysis	Reference
	Dominant	Sample-specific		
Primary samples 25 serous	<i>TP53</i>	<i>PTEN; CDKN2A</i>	Targeted sequencing	39
31 spatially separated samples 6 serous	<i>TP53</i>	<i>PIK3CA; TNNB1; PDGFR; NF1; SH3GL1; RBM15</i>	Exome sequencing,	58
Tumor cells from ascites 3 serous	<i>TP53</i>	<i>BRCA1; CSMD3; MACF1; CAPN7; DMD; ORS1A; PREX2; AP1b1</i>	Exome sequencing	59
11 spatially separated samples 1 serous	<i>TP53</i>	<i>KIF13A; CTSG; SLCO3A1; SPIC</i>	Exome sequencing	65
92 serous Primary 80 Ascites 12	<i>TP53</i>	<i>BRCA1; BRCA2; RB1; NF1; PTEN; RAD51B</i>	Whole genome sequencing	80
135 spatially and temporally separated samples 4 serous	<i>TP53</i>	<i>BRCA1; BRCA2; APC; NF1</i>	Exome sequencing Targeted sequencing	81
62 temporally separated samples 31 serous	<i>TP53</i>	<i>MECOM; ERBB2; CCNE1; ERCC2; ERCC1; TERT; BRCA1</i>	Exome sequencing	82
38 serous Primary 20 Recurrent 18	<i>TP53</i>	<i>MYCC; NF1; KRAS; BRCA1</i>	Targeted sequencing	112

heterogeneity in qualitative ways. The development of NGS technologies around 2005, led to a paradigm shift in the field of genomics, away from the qualitative studies based on single markers, and to the large-scale quantitative datasets. The subsequent application of NGS technologies to human tumors revealed that ITH is common in many types of human cancers including ovarian cancer.<sup>58–62</sup>

NGS has been applied to the molecular characterization of tumors, and to identify new druggable targets, as well as to select appropriate patients for clinical trials.<sup>63</sup> Potential druggable targets recently identified by NGS in serous and non-serous subtypes of ovarian cancer are listed in Tables 1 and 2. The lack of successful treatment strategies in ovarian cancer made the Cancer Genome Atlas (TCGA) researchers to integrate genomic analyses

**Table 2.** Potential druggable targets identified by NGS technology in non-serous ovarian cancer

Samples	Druggable targets		Genomic Analysis	Reference
	Dominant	Sample-specific		
Primary samples 27 clear cell	<i>PIK3CA</i>	<i>KRAS; AKT1; CTNNB1; TP53</i>	Targeted sequencing	39
Primary samples 10 endometrioid	None	<i>PIK3CA; KRAS; PTEN; ERBB2; NRAS; TP53</i>	Targeted sequencing	39
Primary samples 3 mucinous	<i>ERBB2; TP53</i>	<i>PIK3CA; CDKN2A</i>	Targeted sequencing	39
Primary samples 6 endometrioid	None	<i>PIK3CA; PTEN; PIK3R1; ARID1A; CTNNB1</i>	Exome sequencing Targeted sequencing	42
Primary samples 3 mucinous	None	<i>TP53, KRAS</i>	Exome sequencing Targeted sequencing	42
Primary samples 11 mucinous	<i>KRAS</i>	<i>TP53; BRAF; CDKN2A</i>	Exome sequencing	47
5 endometrioid Primary 3 Recurrent 2		<i>TP53; MYCC</i>	Targeted sequencing	112
3 clear cell Primary 2 Recurrent 1	None	<i>TP53; BRCA1; MET</i>	Targeted sequencing	112
Primary sample 1 mucinous	None	<i>TP53; AURKA; NOTCH1; FGF1R</i>	Targeted sequencing	112
Primary samples 48 clear cell	<i>ARID1A</i>	<i>PIK3CA; PPP2R1A; KRAS BRAF, ERBB2, PDGFRB, PGR</i>	Exome sequencing	113
Primary samples 105 clear cell	<i>PIK3CA</i>	<i>TP53, KRAS</i>	Targeted sequencing	114
Primary samples 39 clear cell	<i>ARID1A; PIK3A</i>	<i>KRAS; PPP2R1A; PTEN; MLL3; ARID1B; PIK3R1</i>	Exome sequencing	115,116

of ovarian cancer into molecular abnormalities related to the pathophysiology, clinical outcome, and potential druggable targets in HGSOC.<sup>64</sup> The TCGA study provides a large-scale integrative view of the genomic aberrations in HGSOC with extensive heterogeneity between individual tumors. However, ITH of ovarian cancer in primary and metastatic lesions has remained largely unexplored. Recent study of our group explored the tumoral evolution during the metastasis through analysis of genetic mutations by NGS.<sup>65</sup> There was a negligible accumulation of new mutations during metastasis. Interestingly, only 6% of somatic mutations were common mutations present in all samples, and the majority of somatic mutations detected in the metastatic samples were also present in the primary tumor samples. This study supports that peritoneal seeding arises with little accumulation of somatic mutations and copy number alteration, which might suggest that metastatic potential may have developed at an early stage of ovarian cancer development.

The idea that primary and metastases are clonally related is not new. In 1992, Jacob et al. performed several analyses to determine the clonal origin of metastases in ovarian cancer. This study supports that most metastases in ovarian cancer are monoclonal based on the pattern of genetic alterations including the loss of heterozygosity, p53 gene mutation and X-chromosome inactivation.<sup>66</sup> Numerous studies also reported that the primary and its corresponding metastases show the same pattern of genomic alterations, which support the idea that the major gross genetic changes in ovarian cancer take place in the primary tumor.<sup>67–70</sup> It has also been shown that the malignant cells presented in ascites reflect its corresponding primary tumor.<sup>71</sup> Moreover, Yin and

colleagues reported that synchronous bilateral ovarian cancer are monoclonal; however, primary and metastatic tumors further evolved upon dissociation.<sup>72</sup>

Unlike other solid tumors, the biological behavior of ovarian cancer has been thought to be unique and rather simple. Most patients are diagnosed in late-stage disease and disease is predominantly confined to the peritoneal cavity and there is a clear sign of predilected metastasis to the omentum. These clinical observations lead to a misleading conclusion in the past that the metastasis in ovarian cancer is easier and they lack hematogenous metastasis.<sup>72,73</sup> Although there is a clear sign of predilected metastasis to the omentum, recent studies by Anil Sood and Buckanovich demonstrated an alternative route of metastasis in ovarian cancer, involving hematogenous pattern of metastasis in ovarian cancer.<sup>74,75</sup> Using a parabiosis model, they showed that intraperitoneal injection of ovarian cancer cells in host mouse could hematogenously metastasized to the omentum of guest mouse. ErbB3-neuregulin 1 (NRG1) axis was shown to be a dominant pathway responsible for this route of metastasis.<sup>74</sup> ErbB3 is overexpressed in a number of cancers including ovarian cancer and is associated with chemoresistance.<sup>76,77</sup> More recently, intravenous injection of ovarian cancer cell lines generated the intra-ovarian tumor in 80% of mice and development of ascites. And also in both murine and primary human ovarian tumor cell models, subcutaneous injection of ovarian tumor cells resulted in metastasis to the ovary, supporting hematogenous spread of ovarian cancer in other metastatic sites possibly.<sup>75</sup> These results support that the ovary plays a critical role in ovarian cancer cell metastasis within the peritoneal cavity.



Ovarian cancer cells leave the ovary as single cells or clusters, it is thought that they move through a passive mechanism, carried by the physiological movement of peritoneal fluid, the malignant ascites.<sup>73</sup> The malignant ascites creates tumor-friendly microenvironment thereby contributing to tumor heterogeneity in ovarian cancer (reviewed in ref. <sup>78</sup>). Interestingly, the degree of spatial genetic diversity varied widely across the patients, reflecting extensive intra-patient and inter-patient variability. The difference in the tumor microenvironment may explain these heterogeneity, at least in part.<sup>58</sup> However, our understanding in intra-patient and inter-patient variability is limited. Larger studies are warranted to understand the pattern or direction of tumor evolution in ovarian cancer.

#### Tumoral heterogeneity and chemoresistance

After cytoreductive surgery and adjuvant chemotherapy, the majority of the ovarian cancer patients achieve a clinical complete remission. However, ~50–70% of the patients will experience the recurrent associated with occult chemoresistance. Overcoming chemoresistance is one of the major challenges faced in ovarian cancer research. ITH has been proposed as the main cause of treatment failure and drug resistance in ovarian cancer and other primary cancer.<sup>79</sup>

Most studies of tumor evolution and heterogeneity handled a single time-point samples, providing very indirect information. This is primarily due to difficulties in collection of longitudinal samples from cancer patients and the high costs of genomic profiling. In 2013, Bashashati et al. obtained longitudinal samples from the same patient at primary debulking surgery and after 42 months, with 21 cycles of multi-agent chemotherapy in between. Highly conserved mutations were observed between primary samples and relapsed samples, exhibiting near-identical genomic landscapes.<sup>58</sup> In the same year, Castellarin et al. performed whole exome sequencing (WES) on cancer cells harvested from ascites at multiple time points. Ascites were collected at primary, at the first recurrence, and the second recurrence from three HGSOC patients receiving standard treatment to determine the clonal origin and mutational adaptations across the recurrence.<sup>59</sup> From both studies, genetically distinct heterogeneous clones are considered to be present prior to treatment and most of the mutations in primary tumors persisted despite treatment, suggesting that most clones are able to evade current chemotherapy.<sup>58,59</sup>

In contrast, whole genome sequencing of HGSOC patients found few differences between primary tumor and ascites samples. The number of non-silent coding single nucleotide variants (SNVs) increased with the number of platinum-based chemotherapy cycles, suggesting that primary tumor further evolved during treatment. However, there were no additional platinum drug induced mutations in driver genes. Moreover, the genetic alterations were personalized rather than generalized.<sup>80</sup> More recent study reported in 2015, analyzed the patterns of clonal evolution of relapsed samples in HGSOC cases using phylogenetic analysis. The phylogenetic analysis of relapsed samples showed that there were small changes in heterogeneity during neoadjuvant therapy and the prevalent clonal population at clinical recurrence arose from early divergence events.<sup>81</sup> In another study, 31 paired tumor biopsies from HGSOC patient from primary debulking surgery to disease relapse were analyzed. Although the relative number of primary-specific and relapse-specific mutations vary substantially across the patients, in general, this study supported that recurrent tumors originate from a single clone that escapes platinum therapy.<sup>82</sup>









#### Cellular heterogeneity and chemoresistance

It is well established fact that tumor tissues are comprised of diverse cellular components, including cancer cells and stromal

cells.<sup>78</sup> Recent advances in single-cell technology further expanded our understanding on the cellular heterogeneity by identifying cellular subsets within the tumor and stromal compartments. Proteome analysis of HGSOC at the single-cell level identified clinically relevant subsets, which were bearing mesenchymal traits, based on high vimentin and low E-cadherin expression. Furthermore, these cells co-expressed cMyc and HE4, and these phenotypes were implicated in tumor development and carboplatin resistance.<sup>83</sup> Epithelial to mesenchymal transition (EMT) is the feature of cancer stem cell (CSC) and implicated in ovarian cancer progression including chemoresistance. A number of molecules are associated with EMT and chemoresistance in ovarian cancer. Notch is well-known stem cell-signaling pathway, and of those the activation of Notch3 and its downstream gene SUSD2 expression is associated with EMT through upregulation of EpCAM and conferred chemoresistance to cisplatin in HGSOC.<sup>84</sup> Moreover, cellular heterogeneity is more pronounced in primary tumors and cellular subsets at a transitory epithelial/mesenchymal hybrid stage that are characterized by low membrane E-cadherin, high cytoplasmic E-cadherin, high CD133, high CD44 and low Tie2 expression displayed CSC features.<sup>85</sup> Other recently reported stemness and EMT regulatory genes in ovarian cancer include, KDM5A,<sup>86</sup> splice isoform of CD44s (standard)<sup>87</sup> and CD73.<sup>88</sup> KDM5A is histone demethylase and its expression is associated with drug resistance in breast cancer<sup>89</sup> glioblastoma multiforme,<sup>90</sup> and ovarian cancer.<sup>86</sup> CD44 variant containing exons v8-10 (CD44v8-10) recently have been identified as new CSC marker in gastric cancer<sup>91</sup> and cisplatin resistance in urothelial cancer.<sup>92</sup> In ovarian cancer, CD44v8-10 expression in primary ovarian cancer is associated with epithelial morphology and better prognosis, whereas expression of its soluble form in ascites is associated with poor prognosis.<sup>93</sup> Upregulation of transforming growth factor  $\beta$  (TGF $\beta$ ) mediated left-right determination factor (LEFTY) was initially reported in pancreatic cancer cell lines as a novel tumor suppressor, based on suppressive effect on pancreatic cancer cell proliferation.<sup>94</sup> LEFTY is identified as a new CSC marker and associated with EMT in clear cell subtype.<sup>95</sup>

Cellular diversity in tumor tissues and its stromal microenvironment can induce chemoresistance. Recent study showed that mesenchymal stromal cells (MSC), especially cancer-associated fibroblasts (CAFs) could induce chemoresistance in trans manner by secreting CCL2 and CCL5, which in turn promote IL-6 production in ovarian cancer cells and subsequent chemoresistance.<sup>96</sup> CAFs are also main source of IL-6 secretion in ovarian tumor microenvironment and induce chemoresistance to paclitaxel.<sup>97</sup> Role of IL-6 in ovarian cancer chemoresistance is noted in a number of studies and downstream molecular mechanisms implicated in chemoresistance includes PYK2,<sup>96</sup> JAK2/STAT3,<sup>97</sup> Ras/MEK/ERK and PI3K/Akt signaling.<sup>98</sup> A heparin-binding growth factor called midkine from CAFs induced cisplatin resistance in a number of cancer cells including ovarian cancer by increasing the expression levels of lncRNA ANRIL in cancer cells.<sup>99</sup> How CAFs promote chemoresistance in ovarian cancer is not fully understood yet. Recent report demonstrated crosstalk between CAFs and endothelial cells. It has been shown that CAFs regulate lipoma preferred partner (LPP) gene in endothelial cells via MFAP5 thereby increasing resistance to paclitaxel in vivo.<sup>100</sup>

Irrespective of germ-line and somatic mutations, BRCA1/2 mutations are known to be related with development and progression of ovarian cancer, especially for those of the high-grade serous subtype.<sup>101–103</sup> Poly ADP-ribose polymerase (PARP) inhibitors, such as olaparib, rucaparib, and veliparib have recently been recognized as promising therapeutics for ovarian cancer patients with inactivating BRCA1/2 mutations as a single agent or a combination with other anticancer agents.<sup>103–105</sup> To gain an insight into the selection of patients who may benefit from the treatment with PARP inhibitors, rapid and precise sequencing methods would be required. Recent study has analyzed both

Patterns of clonal evolution		Clinical challenge	
Linear			<b>Simple,</b> Low ITH, single biopsy represent entire tumor
Branching			<b>Difficult,</b> Extensive ITH, spatial-temporal biopsy are required
Neutral			<b>Difficult,</b> Extensive ITH, spatial-temporal biopsy are required
Punctuated			<b>Simple,</b> ITH generated early Single biopsy samples represent entire tumor Subclonal expansion are rare

**Fig. 2** Patterns of clonal evolution and clinical challenges. Linear evolution occurs in the presence of clonal selection over time but can generate intra-tumoral and inter-tumoral heterogeneity if the selective sweep is incomplete or in a different microenvironment. Branching evolution occurs in the presence of multiple clonal selection over time, thus generate extensive ITH. Neutral evolution occurs in the absence of selective sweep but accumulation of random mutation over time result in extensive ITH. Punctuated evolution occurs in the absence of selective sweep, ITH occurs in the early stage of the tumor development and there is no further subclonal selection and expansion. Color dots indicate clones with different genotypes

germline and somatic mutations in BRCA1/2 genes. The study showed detection of mutational status of BRCA1/2 by newly developed technology called ‘single-molecule molecular inversion probe (smMIP)-based targeted NGS’ using paraffin-embedded tissue.<sup>106</sup> Although PARP inhibitors are effective at targeting ovarian cancer with BRCA1/2 deficiency, explained by the concept of synthetic lethality, having a good grasp of those who would benefit from the therapy is extremely important in order to potentiate therapeutic outcomes. Recent clinical trial studies have shown differential response rate of PARP inhibitor between platinum-sensitive and platinum-resistant HGSOC groups.<sup>107</sup>

#### Tumoral heterogeneity and clinical implication (limitation)

ITH is a result of the evolutionary forces of selection pressures.<sup>108</sup> During ongoing evolution, subclones are selected according to their fitness to survive in divergent microenvironmental conditions. Selection is executed based on the phenotypes or genotypes that have survival advantage in a given environment and allow further clonal expansion and evolution. Therefore, the degree of ITH in a given patient correlates with the rate of recurrent disease. This raises the question whether we could improve therapeutic responses and reduce recurrent tumor in ovarian cancer patients by alternating treatment styles.

NGS data can further be used to reconstruct clonal lineages and expansion to understand tumor evolution. A number of algorithms are available to construct phylogenetic trees from ITH datasets (reviewed in ref.<sup>109</sup>). Thus, the history of a given tumor can be tracked retrospectively by estimating the order in which mutations occurred. There are four models of tumor evolution,

linear, branching, neutral, and punctuated evolution.<sup>110,116</sup> Models of tumor evolution have different implications for the clinical diagnosis, prognosis, and treatment of cancer patients (Fig. 2).

Accumulating NGS studies in ovarian cancer suggest that it is possible to improve therapeutic response and could potentially prevent the development of recurrences by targeting multiple mutated pathways present in primary tumors. Although their primary tumor is often completely removed during debulking surgery, the majority of the patients with ovarian cancer die as a consequence of metastatic disease. Studies showed a rather consistent picture of HGSOC, showing a dynamic entity composed of multiple populations of genetically and phenotypically distinct subclones evolved from a single ancestral clone with patient-specific patterns of branched evolution.<sup>58,65,81,111</sup>

Linear and punctuated evolution imply limited ITH, simplifying diagnostic assays, because single biopsy samples represent entire tumor. In contrast both the branching and neutral evolution suggest that ITH is extensive and would require multi-sampling approaches from different spatial regions to detect all of the clinically actionable mutations in the tumor. For example, longitudinal samples collected from primary to recurrent disease showed that a mutation which took up small portion in the primary tumor sample, became dominant close to 100% in recurrent tumor samples.<sup>59</sup> These results suggest that the primary and recurrent diseases shared similar genetic alterations, but current therapy fails to destroy all the clones in the primary case. Larger studies are needed to establish whether the mutational conservation between primary and relapsed tumor samples is a feature of long survivorship in ovarian cancer patients. A number of studies support that patients with tumors with high clonal expansion show short survival and resistant relapse.<sup>58,81</sup>

Lambrechts et al. reconstructed tumor evolution from 31 paired surgical biopsies collected at primary and subsequent relapse. Authors failed to observe a dominant platinum-induced mutation signature but every tumor pair showed branched evolution pattern.<sup>82</sup> This suggest that for HGSOC, biopsies to detect novel mutations are needed at every disease relapse for personalized anticancer therapy.

## CONCLUSION

The current treatment protocols for women with ovarian cancer are not subtype specific, while each histologic subtype contains unique mutation patterns. Treatment for cancer is moving towards the personalized therapy. Advances in genomics increased our understanding of the tumor genomes. We know that the cancer cells in a tumor are not all identical, composed of different clones, defined as sets of cancer cells that share a common genotype. Evolution theory successfully guided us to understand the cancer progression model. The interaction of cancer cells with micro-environment modulates tumor heterogeneity, affecting the response to therapy. Therefore, understanding tumoral heterogeneity caused by genetic alterations of cancer cells and their interaction with tumor microenvironment will help the researchers to design new therapeutic approaches towards precision medicine. Further understanding of tumor evolution with molecular, histopathological, and clinical characterizations will enlighten the precision medicine in ovarian cancer.

## ACKNOWLEDGEMENTS

This research was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number: HI16C2037 and HA17C0037).

## AUTHOR CONTRIBUTIONS

S.K. authored majority of the manuscript, others contributed to various sections of this review and overall editing. Y.S.S. supervised the idea generation and revised it critically.

## ADDITIONAL INFORMATION

**Competing interests:** The authors declare no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## REFERENCES

- Meinhold-Heerlein, I. & Hauptmann, S. The heterogeneity of ovarian cancer. *Arch. Gynecol. Obstet.* **289**, 237–239 (2014).
- Gilks, C. B. et al. Tumor cell type can be reproducibly diagnosed and is of independent prognostic significance in patients with maximally debulked ovarian carcinoma. *Hum. Pathol.* **39**, 1239–1251 (2008).
- Meinhold-Heerlein, I. et al. The new WHO classification of ovarian, fallopian tube, and primary peritoneal cancer and its clinical implications. *Arch. Gynecol. Obstet.* **293**, 695–700 (2016).
- Koshiyama, M., Matsumura, N. & Konishi, I. Clinical efficacy of ovarian cancer screening. *J. Cancer* **7**, 1311–1316 (2016).
- Heintz, A. P. et al. Carcinoma of the ovary. FIGO 26th annual report on the results of treatment in gynecological cancer. *Int. J. Gynaecol. Obstet.* **95**(Suppl. 1), S161–S192 (2006).
- Kim, A., Ueda, Y., Naka, T. & Enomoto, T. Therapeutic strategies in epithelial ovarian cancer. *J. Exp. Clin. Cancer Res* **31**, 14 (2012).
- Davis, A., Tinker, A. V. & Friedlander, M. "Platinum resistant" ovarian cancer: what is it, who to treat and how to measure benefit? *Gynecol. Oncol.* **133**, 624–631 (2014).
- Wiltshaw, E. & Kroner, T. Phase II study of cis-dichlorodiammineplatinum(II) (NSC-119875) in advanced adenocarcinoma of the ovary. *Cancer Treat. Rep.* **60**, 55–60 (1976).

- McGuire, W. P. et al. Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer. *N. Engl. J. Med.* **334**, 1–6 (1996).
- Vergote, I. et al. Neoadjuvant chemotherapy or primary surgery in stage IIIC or IV ovarian cancer. *N. Engl. J. Med.* **363**, 943–953 (2010).
- Morgan, R. J. Jr et al. Ovarian cancer, version 1.2016, NCCN clinical practice guidelines in oncology. *J. Natl Compr. Cancer Netw.* **14**, 1134–1163 (2016).
- Morrison, J., Haldar, K., Kehoe, S. & Lawrie, T. A. Chemotherapy versus surgery for initial treatment in advanced ovarian epithelial cancer. *Cochrane Database Syst. Rev.*, CD005343, <https://doi.org/10.1002/14651858.CD005343.pub3> (2012).
- Vandenput, I. et al. Neoadjuvant chemotherapy followed by interval debulking surgery in patients with serous endometrial cancer with transperitoneal spread (stage IV): a new preferred treatment? *Br. J. Cancer* **101**, 244–249 (2009).
- Chi, D. S. et al. An analysis of patients with bulky advanced stage ovarian, tubal, and peritoneal carcinoma treated with primary debulking surgery (PDS) during an identical time period as the randomized EORTC-NCIC trial of PDS vs neoadjuvant chemotherapy (NACT). *Gynecol. Oncol.* **124**, 10–14 (2012).
- Leary, A., Cowan, R., Chi, D., Kehoe, S., & Nankivell, M. Primary surgery or neoadjuvant chemotherapy in advanced ovarian cancer: the debate continues. *Am. Soc. Clin. Oncol. Educ. Book* **35**, 153–162 (2016).
- Vergote, I. et al. Neoadjuvant chemotherapy is the better treatment option in some patients with stage IIIC to IV ovarian cancer. *J. Clin. Oncol.* **29**, 4076–4078 (2011).
- Bookman, M. A. First-line chemotherapy in epithelial ovarian cancer. *Clin. Obstet. Gynecol.* **55**, 96–113 (2012).
- Stuart, G. C. et al. 2010 Gynecologic Cancer InterGroup (GIG) consensus statement on clinical trials in ovarian cancer: report from the Fourth Ovarian Cancer Consensus Conference. *Int. J. Gynecol. Cancer* **21**, 750–755 (2011).
- du Bois, A. et al. 2004 consensus statements on the management of ovarian cancer: final document of the 3rd International Gynecologic Cancer InterGroup Ovarian Cancer Consensus Conference (GIG OCC 2004). *Ann. Oncol.* **16**(Suppl. 8), viii7–viii12 (2005).
- Sugiyama, T. et al. Clinical characteristics of clear cell carcinoma of the ovary: a distinct histologic type with poor prognosis and resistance to platinum-based chemotherapy. *Cancer* **88**, 2584–2589 (2000).
- Itamochi, H. et al. Sensitivity to anticancer agents and resistance mechanisms in clear cell carcinoma of the ovary. *Jpn. J. Cancer Res.* **93**, 723–728 (2002).
- Mabuchi, S., Sugiyama, T. & Kimura, T. Clear cell carcinoma of the ovary: molecular insights and future therapeutic perspectives. *J. Gynecol. Oncol.* **27**, e31 (2016).
- Lee, C. K. et al. Prognostic nomogram to predict progression-free survival in patients with platinum-sensitive recurrent ovarian cancer. *Br. J. Cancer* **105**, 1144–1150 (2011).
- Previs, R. A. et al. A prognostic nomogram to predict overall survival in women with recurrent ovarian cancer treated with bevacizumab and chemotherapy. *Gynecol. Oncol.* **132**, 531–536 (2014).
- Xu, X. L. et al. A novel nomogram based on LODDS to predict the prognosis of epithelial ovarian cancer. *Oncotarget* **8**, 8120–8130 (2017).
- Cho, K. R. Ovarian cancer update: lessons from morphology, molecules, and mice. *Arch. Pathol. Lab. Med.* **133**, 1775–1781 (2009).
- Karnezis, A. N. & Cho, K. R. Of mice and women—non-ovarian origins of "ovarian" cancer. *Gynecol. Oncol.* **144**, 5–7 (2017).
- Cass, I. et al. BRCA-mutation-associated fallopian tube carcinoma: a distinct clinical phenotype? *Obstet. Gynecol.* **106**, 1327–1334 (2005).
- Piek, J. M. et al. BRCA1/2-related ovarian cancers are of tubal origin: a hypothesis. *Gynecol. Oncol.* **90**, 491 (2003).
- Medeiros, F. et al. The tubal fimbria is a preferred site for early adenocarcinoma in women with familial ovarian cancer syndrome. *Am. J. Surg. Pathol.* **30**, 230–236 (2006).
- Lee, Y. et al. A candidate precursor to serous carcinoma that originates in the distal fallopian tube. *J. Pathol.* **211**, 26–35 (2007).
- Kindelberger, D. W. et al. Intraepithelial carcinoma of the fimbria and pelvic serous carcinoma: evidence for a causal relationship. *Am. J. Surg. Pathol.* **31**, 161–169 (2007).
- Kuhn, E. et al. TP53 mutations in serous tubal intraepithelial carcinoma and concurrent pelvic high-grade serous carcinoma—evidence supporting the clonal relationship of the two lesions. *J. Pathol.* **226**, 421–426 (2012).
- McDaniel, A. S. et al. Next-generation sequencing of tubal intraepithelial carcinomas. *JAMA Oncol.* **1**, 1128–1132 (2015).
- Kim, J. et al. High-grade serous ovarian cancer arises from fallopian tube in a mouse model. *Proc. Natl Acad. Sci. USA* **109**, 3921–3926 (2012).
- Labidi-Galy, S. I. et al. High grade serous ovarian carcinomas originate in the fallopian tube. *Nat. Commun.* **8**, 1093 (2017).
- Kobel, M. et al. Ovarian carcinoma subtypes are different diseases: implications for biomarker studies. *PLoS Med.* **5**, e232 (2008).



38. Haruta, S. et al. Molecular genetics and epidemiology of epithelial ovarian cancer (Review). *Oncol. Rep.* **26**, 1347–1356 (2011).
39. Takenaka, M. et al. Profiling of actionable gene alterations in ovarian cancer by targeted deep sequencing. *Int. J. Oncol.* **46**, 2389–2398 (2015).
40. Kurman, R. J. & Shih, Ie, M. The origin and pathogenesis of epithelial ovarian cancer: a proposed unifying theory. *Am. J. Surg. Pathol.* **34**, 433–443 (2010).
41. Kurman, R. J. & Shih, Ie, M. Pathogenesis of ovarian cancer: lessons from morphology and molecular biology and their clinical implications. *Int. J. Gynecol. Pathol.* **27**, 151–160 (2008).
42. Teer, J. K. et al. Mutational heterogeneity in non-serous ovarian cancers. *Sci. Rep.* **7**, 9728 (2017).
43. Kurman, R. J. & Shih, Ie, M. Molecular pathogenesis and extraovarian origin of epithelial ovarian cancer—shifting the paradigm. *Hum. Pathol.* **42**, 918–931 (2011).
44. Schultheis, A. M. et al. Massively parallel sequencing-based clonality analysis of synchronous endometrioid endometrial and ovarian carcinomas. *J. Natl Cancer Inst.* **108**, djv427 (2016).
45. Valtcheva, N. et al. Tracking the origin of simultaneous endometrial and ovarian cancer by next-generation sequencing—a case report. *BMC Cancer* **17**, 66 (2017).
46. McConechy, M. K. et al. Ovarian and endometrial endometrioid carcinomas have distinct CTNBN1 and PTEN mutation profiles. *Mod. Pathol.* **27**, 128–134 (2014).
47. Ryland, G. L. et al. Mutational landscape of mucinous ovarian carcinoma and its neoplastic precursors. *Genome Med.* **7**, 87 (2015).
48. Kim, B. et al. PGC1alpha induced by reactive oxygen species contributes to chemoresistance of ovarian cancer cells. *Oncotarget* **8**, 60299–60311 (2017).
49. Ning, Y., Luo, C., Ren, K., Quan, M. & Cao, J. FOXO3a-mediated suppression of the self-renewal capacity of sphere-forming cells derived from the ovarian cancer SKOV3 cell line by 7-difluoromethoxy-5,4'-di-n-octyl genistein. *Mol. Med. Rep.* **9**, 1982–1988 (2014).
50. Yin, M. et al. Tumor-associated macrophages drive spheroid formation during early transcoelomic metastasis of ovarian cancer. *J. Clin. Invest.* **126**, 4157–4173 (2016).
51. Smart, C. E. et al. In vitro analysis of breast cancer cell line tumourspheres and primary human breast epithelia mammospheres demonstrates inter- and intrasphere heterogeneity. *PLoS One* **8**, e64388 (2013).
52. Soeda, A. et al. CORRIGENDUM: the evidence of glioblastoma heterogeneity. *Sci. Rep.* **5**, 9630 (2015).
53. Feng, H. et al. Homogeneous pancreatic cancer spheroids mimic growth pattern of circulating tumor cell clusters and macrometastases: displaying heterogeneity and crater-like structure on inner layer. *J. Cancer Res. Clin. Oncol.*, <https://doi.org/10.1007/s00432-017-2434-2> (2017).
54. Kim, S. et al. Malignant ascites enhances migratory and invasive properties of ovarian cancer cells with membrane bound IL-6R in vitro. *Oncotarget* **7**, 83148–83159 (2016).
55. Cho, U., Kim, B., Kim, S., Han, Y. & Song, Y. S. Pro-inflammatory M1 macrophage enhances metastatic potential of ovarian cancer cells through NF-kappaB activation. *Mol. Carcinog.*, <https://doi.org/10.1002/mc.22750> (2017).
56. Lawrenson, K. & Gayther, S. A. Ovarian cancer: a clinical challenge that needs some basic answers. *PLoS Med.* **6**, e25 (2009).
57. Brown, T. M. & Fee, E. Rudolf Carl Virchow: medical scientist, social reformer, role model. *Am. J. Public Health* **96**, 2104–2105 (2006).
58. Bashashati, A. et al. Distinct evolutionary trajectories of primary high-grade serous ovarian cancers revealed through spatial mutational profiling. *J. Pathol.* **231**, 21–34 (2013).
59. Castellarin, M. et al. Clonal evolution of high-grade serous ovarian carcinoma from primary to recurrent disease. *J. Pathol.* **229**, 515–524 (2013).
60. Ding, L. et al. Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* **481**, 506–510 (2012).
61. Gerlinger, M. et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N. Engl. J. Med.* **366**, 883–892 (2012).
62. Mackenzie, R. et al. Targeted deep sequencing of mucinous ovarian tumors reveals multiple overlapping RAS-pathway activating mutations in borderline and cancerous neoplasms. *BMC Cancer* **15**, 415 (2015).
63. Takeda, M. et al. Clinical application of amplicon-based next-generation sequencing to therapeutic decision making in lung cancer. *Ann. Oncol.* **26**, 2477–2482 (2015).
64. Cancer Genome Atlas Research, N. Integrated genomic analyses of ovarian carcinoma. *Nature* **474**, 609–615 (2011).
65. Lee, J. Y. et al. Tumor evolution and intratumor heterogeneity of an epithelial ovarian cancer investigated using next-generation sequencing. *BMC Cancer* **15**, 85 (2015).
66. Jacobs, I. J. et al. Clonal origin of epithelial ovarian carcinoma: analysis by loss of heterozygosity, p53 mutation, and X-chromosome inactivation. *J. Natl Cancer Inst.* **84**, 1793–1798 (1992).
67. Adib, T. R. et al. Predicting biomarkers for ovarian cancer using gene-expression microarrays. *Br. J. Cancer* **90**, 686–692 (2004).
68. Hibbs, K. et al. Differential gene expression in ovarian carcinoma: identification of potential biomarkers. *Am. J. Pathol.* **165**, 397–414 (2004).
69. Israeli, O. et al. Genomic analyses of primary and metastatic serous epithelial ovarian cancer. *Cancer Genet. Cytogen.* **154**, 16–21 (2004).
70. Khalique, L. et al. The clonal evolution of metastases from primary serous epithelial ovarian cancers. *Int. J. Cancer* **124**, 1579–1586 (2009).
71. Choi, Y. J. et al. Intraindividual genomic heterogeneity of high-grade serous carcinoma of the ovary and clinical utility of ascitic cancer cells for mutation profiling. *J. Pathol.* **241**, 57–66 (2017).
72. Yin, X. et al. Clonality, heterogeneity and evolution of synchronous bilateral ovarian cancer. *Cancer Res.* <https://doi.org/10.1158/0008-5472.CAN-17-1461> (2017).
73. Lengyel, E. Ovarian cancer development and metastasis. *Am. J. Pathol.* **177**, 1053–1064 (2010).
74. Pradeep, S. et al. Hematogenous metastasis of ovarian cancer: rethinking mode of spread. *Cancer Cell* **26**, 77–91 (2014).
75. Coffman, L. G. et al. New models of hematogenous ovarian cancer metastasis demonstrate preferential spread to the ovary and a requirement for the ovary for abdominal dissemination. *Transl. Res.* **175**, 92–102 e102 (2016).
76. Davies, S. et al. High incidence of ErbB3, ErbB4, and MET expression in ovarian cancer. *Int. J. Gynecol. Pathol.* **33**, 402–410 (2014).
77. Meetze, K. et al. Neuregulin 1 expression is a predictive biomarker for response to AV-203, an ERBB3 inhibitory antibody, in human tumor models. *Clin. Cancer Res.* **21**, 1106–1114 (2015).
78. Kim, S., Kim, B. & Song, Y. S. Ascites modulates cancer cell behavior, contributing to tumor heterogeneity in ovarian cancer. *Cancer Sci.* **107**, 1173–1178 (2016).
79. Swanton, C. Intratumor heterogeneity: evolution through space and time. *Cancer Res.* **72**, 4875–4882 (2012).
80. Patch, A. M. et al. Whole-genome characterization of chemoresistant ovarian cancer. *Nature* **521**, 489–494 (2015).
81. Schwarz, R. F. et al. Spatial and temporal heterogeneity in high-grade serous ovarian cancer: a phylogenetic analysis. *PLoS Med.* **12**, e1001789 (2015).
82. Lambrechts, S. et al. Genetic heterogeneity after first-line chemotherapy in high-grade serous ovarian cancer. *Eur. J. Cancer* **53**, 51–64 (2016).
83. Gonzalez, V. D. et al. Commonly occurring cell subsets in high-grade serous ovarian tumors identified by single-cell mass cytometry. *Cell Rep.* **22**, 1875–1888 (2018).
84. Xu, Y. et al. SUSD2 promotes cancer metastasis and confers cisplatin resistance in high grade serous ovarian cancer. *Exp. Cell Res.* **363**, 160–170 (2018).
85. Strauss, R. et al. Analysis of epithelial and mesenchymal markers in ovarian cancer reveals phenotypic heterogeneity and plasticity. *PLoS One* **6**, e16186 (2011).
86. Feng, T., Wang, Y., Lang, Y. & Zhang, Y. KDM5A promotes proliferation and EMT in ovarian cancer and closely correlates with PTX resistance. *Mol. Med. Rep.* **16**, 3573–3580 (2017).
87. Bhattacharya, R., Mitra, T., Ray Chaudhuri, S. & Roy, S. S. Mesenchymal splice isoform of CD44 (CD44s) promotes EMT/invasion and imparts stem-like properties to ovarian cancer cells. *J. Cell. Biochem.* **119**, 3373–3383 (2018).
88. Lupia, M. et al. CD73 regulates stemness and epithelial-mesenchymal transition in ovarian cancer-initiating cells. *Stem Cell Rep.* **10**, 1412–1425 (2018).
89. Hou, J. et al. Genomic amplification and a role in drug-resistance for the KDM5A histone demethylase in breast cancer. *Am. J. Transl. Res.* **4**, 247–256 (2012).
90. Banelli, B. et al. The histone demethylase KDM5A is a key factor for the resistance to temozolomide in glioblastoma. *Cell Cycle* **14**, 3418–3429 (2015).
91. Lau, W. M. et al. CD44v8-10 is a cancer-specific marker for gastric cancer stem cells. *Cancer Res.* **74**, 2630–2641 (2014).
92. Hagiwara, M. et al. Variant isoforms of CD44 involves acquisition of chemoresistance to cisplatin and has potential as a novel indicator for identifying a cisplatin-resistant population in urothelial cancer. *BMC Cancer* **18**, 113 (2018).
93. Sosulski, A. et al. CD44 splice variant v8-10 as a marker of serous ovarian cancer prognosis. *PLoS One* **11**, e0156595 (2016).
94. Miyata, N. et al. Transforming growth factor beta and Ras/MEK/ERK signaling regulate the expression level of a novel tumor suppressor Lefty. *Pancreas* **41**, 745–752 (2012).
95. Matsumoto, T. et al. TGF-beta-mediated LEFTY/Akt/GSK-3beta/Snail axis modulates epithelial-mesenchymal transition and cancer stem cell properties in ovarian clear cell carcinomas. *Mol. Carcinog.* <https://doi.org/10.1002/mc.22816> (2018).
96. Pasquier, J. et al. CCL2/CCL5 secreted by the stroma induce IL-6/PYK2 dependent chemoresistance in ovarian cancer. *Mol. Cancer* **17**, 47 (2018).
97. Wang, L. et al. CAFs enhance paclitaxel resistance by inducing EMT through the IL6/JAK2/STAT3 pathway. *Oncol. Rep.* **39**, 2081–2090 (2018).
98. Wang, Y. et al. Autocrine production of interleukin-6 confers cisplatin and paclitaxel resistance in ovarian cancer cells. *Cancer Lett.* **295**, 110–123 (2010).



99. Zhang, D. et al. Midkine derived from cancer-associated fibroblasts promotes cisplatin-resistance via up-regulation of the expression of lncRNA ANRIL in tumour cells. *Sci. Rep.* **7**, 16231 (2017).
100. Leung, C. S. et al. Cancer-associated fibroblasts regulate endothelial adhesion protein LPP to promote ovarian cancer chemoresistance. *J. Clin. Invest.* **128**, 589–606 (2018).
101. Schrader, K. A. et al. Germline BRCA1 and BRCA2 mutations in ovarian cancer: utility of a histology-based referral strategy. *Obstet. Gynecol.* **120**, 235–240 (2012).
102. Takaoka, M. & Miki, Y. BRCA1 gene: function and deficiency. *Int. J. Clin. Oncol.* **23**, 36–44 (2018).
103. Dougherty, B. A. et al. Biological and clinical evidence for somatic mutations in BRCA1 and BRCA2 as predictive markers for olaparib response in high-grade serous ovarian cancers in the maintenance setting. *Oncotarget* **8**, 43653–43661 (2017).
104. Musella, A. et al. Rucaparib: an emerging parp inhibitor for treatment of recurrent ovarian cancer. *Cancer Treat. Rev.* **66**, 7–14 (2018).
105. Gray, H. J. et al. Phase I combination study of the PARP inhibitor veliparib plus carboplatin and gemcitabine in patients with advanced ovarian cancer and other solid malignancies. *Gynecol. Oncol.* **148**, 507–514 (2018).
106. Weren, R. D. et al. Novel BRCA1 and BRCA2 tumor test as basis for treatment decisions and referral for genetic counselling of patients with ovarian carcinomas. *Hum. Mutat.* **38**, 226–235 (2017).
107. Morgan, R. D., Clamp, A. R., Evans, D. G. R., Edmondson, R. J. & Jayson, G. C. PARP inhibitors in platinum-sensitive high-grade serous ovarian cancer. *Cancer Chemother. Pharmacol.* **81**, 647–658 (2018).
108. Stratton, M. R. Exploring the genomes of cancer cells: progress and promise. *Science* **331**, 1553–1558 (2011).
109. Schwartz, R. & Schaffer, A. A. The evolution of tumour phylogenetics: principles and practice. *Nat. Rev. Genet.* **18**, 213–229 (2017).
110. Davis, A., Gao, R. & Navin, N. Tumor evolution: linear, branching, neutral or punctuated? *Biochim. Biophys. Acta* **1867**, 151–161 (2017).
111. Hoogstraat, M. et al. Genomic and transcriptomic plasticity in treatment-naive ovarian cancer. *Genome Res.* **24**, 200–211 (2014).
112. Ross, J. S. et al. Comprehensive genomic profiling of epithelial ovarian cancer by next generation sequencing-based diagnostic assay reveals new routes to targeted therapies. *Gynecol. Oncol.* **130**, 554–559 (2013).
113. Shibuya, Y. et al. Identification of somatic genetic alterations in ovarian clear cell carcinoma with next generation sequencing. *Genes Chromosomes Cancer.* <https://doi.org/10.1002/gcc.22507> (2017).
114. Friedlander, M. L. et al. Molecular profiling of clear cell ovarian cancers: identifying potential treatment targets for clinical trials. *Int. J. Gynecol. Cancer* **26**, 648–654 (2016).
115. Murakami, R. et al. Exome sequencing landscape analysis in ovarian clear cell carcinoma shed light on key chromosomal regions and mutation gene networks. *Am. J. Pathol.* **187**, 2246–2258 (2017).
116. Burrell, R. A. & Swanton, C. Tumour heterogeneity and the evolution of polyclonal drug resistance. *Mol. Oncol.* **8**, 1095–1111 (2014).



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018