



Published in final edited form as:

J Am Chem Soc. 2010 June 9; 132(22): 7784–7793. doi:10.1021/ja102337n.

The Thermochemical and Kinetic Properties of Ascorbate are Tuned by its Local Environment: Solution Chemistry and Biochemical Implications

Jeffrey J. Warren* and James M. Mayer*

Department of Chemistry, University of Washington, Box 351700, Seattle, WA 98195-1700

Abstract

Ascorbate (Vitamin C) is a ubiquitous biological cofactor. While its aqueous solution chemistry has long been studied, many *in vivo* reactions of ascorbate occur in enzyme active sites or at membrane interfaces, which have varying local environments. This report shows that the rate and driving force of oxidations of two ascorbate derivatives by the TEMPO radical (2,2'-6,6'-tetramethylpiperidine-1-oxyl) in acetonitrile are very sensitive to the presence of various additives. These reactions proceed by the transfer of a proton and an electron (a hydrogen atom), as is typical of biological ascorbate reactions. The measured rate and equilibrium constants vary substantially with added water or other polar solutes in acetonitrile solutions, indicating large shifts in the reducing power of ascorbate. The correlation of rate and equilibrium constants indicates that this effect has a thermochemical origin rather than being a purely kinetic effect. This contrasts with previous examples of solvent effects on hydrogen atom transfer reactions. Potential biological implications of this apparently unique effect are discussed.

Introduction

Ascorbic acid (Vitamin C) is a key biological cofactor, necessary for human health,¹ that has a wide range of biochemical roles. It is important as an antioxidant, for instance reducing α -tocopheroxyl radical or oxidized glutathione,² and as a cofactor for metalloenzymes such as ascorbate oxidase³ and cytochrome *b*₅₆₁.⁴ At physiological pH, it exists predominantly as the ascorbate monoanion, AscH^- (Scheme 1). Oxidation of ascorbate forms the ascorbyl radical $\text{Asc}^{\bullet-}$, which has one proton and one electron less than AscH^- , and further oxidation gives dehydroascorbate, which can hydrate to the corresponding hemiketal in aqueous solution.⁵

Ascorbate has traditionally been considered a one-electron reductant, but it almost always reacts with loss of both an electron and a proton (Scheme 1) because the neutral ascorbyl radical is a high energy, highly acidic species [$\text{p}K_{\text{a}}(\text{AscH}^{\bullet}) = -0.45$].⁶ In many cases, ascorbate reacts to transfer the e^- and H^+ in a single kinetic step, in effect as a hydrogen atom. Njus and co-workers have shown that ascorbate reacts as a hydrogen atom donor to oxyl radicals, cytochrome *b*₅₆₁, and in other systems.⁷⁸ In such cases, the thermochemistry is in some ways better described by the O–H bond dissociation free energy (BDFE), $73.6 \pm 1.0 \text{ kcal mol}^{-1}$ for aqueous ascorbate (see below), rather than the redox potential.⁹ The standard 0.282 mV potential for ascorbate at pH 7 is mechanistically less informative because it is for the more complex $2e^-/1\text{H}^+$ transfer to produce dehydroascorbate.⁶


We recently described oxidations of ascorbate or 5,6-isopropylidene ascorbate ($i\text{AscH}^-$, Scheme 1) by oxyl radicals¹⁰ or porphyrin-iron-imidazole complexes,¹¹ and showed that these reactions proceed via a concerted transfer of $\text{H}^+ + e^-$ (H^\bullet) rather than by a stepwise mechanism with initial proton or electron transfer.¹⁰ The ascorbyl radical ($\text{Asc}^{\bullet-}$ or $i\text{Asc}^{\bullet-}$) is remarkably persistent in 'dry' MeCN solvent, decaying over hours at ambient temperatures.¹⁰ This contrasts with the rapid disproportionation of ascorbyl radicals in water (milliseconds at 10^{-4} M concentrations).¹² The surprisingly long lifetime has allowed us to directly measure equilibrium constants, as described below, and thereby determine the O–H BDFE of AscH^- in MeCN as 67.7 ± 0.8 kcal mol⁻¹. Remarkably, this is 5.9 ± 1.3 kcal mol⁻¹ weaker than the aqueous value.

Such a large change in X–H homolytic bond strength with solvent is, to our knowledge, unprecedented. The studies described here show that this change is not due the bulk properties of the medium such as dielectric (ϵ). Instead, the dramatic shift in ascorbate homolytic bond strength is predominantly due to the local environment about the ascorbate, especially the presence of hydrogen bond donors. Ascorbate reactions often occur within enzyme active sites or at membrane interfaces that are different from bulk water, so it seems likely that Nature has taken advantage of this unusual property of ascorbate; the last section below explores three possible examples.

Results

I. Ascorbate + TEMPO rate and equilibrium constants in pure solvents

To examine the intrinsic reactivity of ascorbate derivatives, we have examined their oxidation by the stable TEMPO radical (eq 1, TEMPO = 2,2',6,6'-tetramethylpiperidine-1-oxyl). These are representative and easily studied reactions, which occur rapidly by hydrogen atom transfer to form TEMPOH and the corresponding ascorbyl radical (eq 1) as previously described.¹⁰ The studies presented here used the acetonitrile-soluble ascorbate salt with potassium 18-crown-6 ($[\text{K18C6}^+]\text{AscH}^-$) and the DBU- H^+ salt of the isopropylidene derivative $i\text{AscH}^-$ (DBU = 1,2-diazabicyclo[5.4.0]undec-7-ene), avoiding the low melting and more difficult to handle ${}^n\text{Bu}_4\text{N}^+$ compounds used in the previous study. $[\text{K18C6}^+]\text{AscH}^-$ was prepared from potassium ascorbate¹³ + 18-crown-6 in MeCN, and $[\text{DBU-H}^+]\text{AscH}^-$ was generated *in situ* from solutions of $i\text{AscH}_2$ by adding 1 equiv of DBU. Both of these salts are spectroscopically and kinetically identical to their ${}^n\text{Bu}_4\text{N}^+$ analogs.



(1)

Rate constants for the oxidation of $[\text{K18C6}^+]\text{AscH}^-$ by TEMPO (eq 1) have been determined in MeCN, DMSO and water (and in MeCN with various additives, see below). The reactions in MeCN and DMSO were performed under pseudo-first-order conditions of excess TEMPO (15 to 180 equiv), observing the growth of the ascorbyl radical optical band at 377 nm by using UV/Vis stopped-flow spectrophotometry (Figure 1). SPECFIT™ global analysis software was used to fit the spectra over the entire observed spectral range (320–550 nm).¹⁴ The data follow a first order kinetic model over >4 half-lives. A plot of the

derived pseudo-first-order rate constants as a function of [TEMPO] is linear (see Supporting Information), with slopes that are independent of starting [AscH⁻]. These slopes give bimolecular rate constants $k_{1,\text{MeCN}} = 500 \pm 50 \text{ M}^{-1} \text{ s}^{-1}$ and $k_{1,\text{DMSO}} = 100 \pm 10 \text{ M}^{-1} \text{ s}^{-1}$ (see Tables 1 and 2, below). The rate constants (and equilibrium constants, see below) are unaffected by the presence or absence of 0.1 M ⁿBu₄NPF₆. Unless specifically noted, all rate and equilibrium constants were determined at 298 K without added electrolyte. Rate constants for oxidations of isopropylidene ascorbate (*i*AscH⁻) by TEMPO have been measured similarly, under a wide range of conditions, extending the previous report¹⁰ (see Tables 1 and 3, below).

In MeCN and DMSO, with <10 equivalents of TEMPO, reaction 1 proceeds to an equilibrium mixture of the four species. The position of equilibrium can be measured by using the stopped-flow instrument to do a spectrophotometric titration. Reactions are complete after a few seconds, and the position of the equilibrium can be determined before there is significant decay of the ascorbyl radical product. Equilibrium constants were calculated by plotting [TEMPOH][Asc^{•-}]/[AscH⁻] vs. [TEMPO]. These plots show good linearity, validating the mass balance assumption and the equilibrium treatment (see Supporting Information). The derived equilibrium constant for [K18C6⁺]AscH⁻ + TEMPO, $K_{1,\text{MeCN}}$, is 0.14 ± 0.02 ($\Delta G^\circ_{1,\text{MeCN}} = 1.3 \pm 0.1 \text{ kcal mol}^{-1}$), which is within error of the previously measured K for ⁿBu₄N⁺AscH⁻, 0.12 ± 0.04 .¹⁰ Equilibrium constants for analogous reactions were measured similarly and are given in Tables 2 and 3.

Rate constants have also been measured in water (pH 7.1, 100 mM phosphate buffer), for both AscH⁻ and *i*AscH⁻. In water, the ascorbyl radical optical band is not observed because disproportionation of Asc^{•-} is fast¹⁵ compared to the initial reaction of AscH⁻ + TEMPO. Therefore these reactions were done with excess ascorbate (10 – 42 mM) and monitored using the disappearance of the optical band for TEMPO (4.3 mM). The data were fit using SPECTFITTM to a model with an initial bimolecular hydrogen atom transfer step followed by Asc^{•-} disproportionation ($k_{\text{disp}} = 1 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ in 0.045 M phosphate buffer, pH 7¹⁵), yielding $k_{1,\text{H}_2\text{O}} = 2.4 \pm 0.3 \text{ M}^{-1} \text{ s}^{-1}$ and $k_{1,i,\text{H}_2\text{O}} = 5.5 \pm 0.5 \text{ M}^{-1} \text{ s}^{-1}$. The former rate constant is in excellent agreement with a recent determination at more acidic pH for the reduction of TEMPO by ascorbate.¹⁶ The corresponding equilibrium constants cannot be determined in water, again due to the rapid ascorbyl radical disproportionation.

II. Ascorbate bond dissociation free energies in pure solvents

The free energy change of eq 1 in pure solvents (ΔG°_1) is the difference in O–H BDFEs of the ascorbate and TEMPOH in that solvent. Thus, the measured values for K_1 ($= e^{-\Delta G^\circ_1/RT}$) and the well known TEMPO-H BDFEs ($66.5 \pm 0.5 \text{ kcal mol}^{-1}$ in MeCN,¹⁷ $67.5 \pm 1.0 \text{ kcal mol}^{-1}$ in DMSO¹⁸) provide the ascorbate O–H BDFEs in those solvents (Table 1). The bond strength of ascorbate in water and MeCN can also be derived using a well established procedure^{18,19} from the known $\text{p}K_a$ and $E_{1/2}$ values:⁶ $\text{BDFE}(X\text{-H}) = 1.37\text{p}K_a + 23.06E_{1/2} + C_G$ [$C_G(\text{H}_2\text{O}) = 57.6 \text{ kcal mol}^{-1}$ with $E_{1/2}$ vs. NHE^{18b}]. The ascorbate BDFE in MeCN derived from this equation is within error and less precise than that derived from K_1 , so the latter is given in Table 1.²⁰

For both ascorbate and isopropylidene ascorbate, the O–H bond dissociation free energies are the same in MeCN and DMSO, with that for AscH⁻ about 1 kcal mol⁻¹ stronger than *i*AscH⁻. The O–H BDFE of ascorbate is about 6 kcal mol⁻¹ larger in water than in the polar organic solvents, as noted above. These solution BDFEs are related to the gas-phase BDFE of ascorbate by the free energies of solution of AscH⁻, Asc[•] and H[•], and about half of the difference between the organic solvents and water can be ascribed to differences in $\Delta G^\circ(\text{H}^\bullet)_{\text{solv}}$.²¹ Still, the 6 kcal mol⁻¹ strengthening of the ascorbate O–H bond between MeCN/DMSO vs. H₂O is significantly larger than would be expected. For substrates with one

hydroxylic moiety the change in BDFE from MeCN to water is typically +2–4 kcal mol⁻¹ and from DMSO to water is about +1–3 kcal mol⁻¹ (see Supporting Information).

III. Rate and equilibrium constants in mixed MeCN/H₂O solvent

To probe the substantial difference in BDFEs for ascorbate in acetonitrile versus water, reaction 1 has been studied in acetonitrile/water mixtures. The effects of other ‘additives’ are explored in the next section.

Rate constants for ascorbate (AscH⁻) oxidation by TEMPO were measured under pseudo-first-order conditions, as described above, in solutions with the mole fraction of water (χ_{water}) from $\sim 3 \times 10^{-5}$ (“dry” acetonitrile, see Experimental Section) to 0.012. These values correspond to concentrations of water up to 200 mM, or 0.1 to 720 equivalents of water per ascorbate. Qualitatively, the reaction becomes substantially slower in the presence of even quite small amounts of water. Second-order rate constants for reactions with $\chi_{\text{water}} < 0.003$ were determined by simple pseudo-first-order kinetic analysis, as above. However, in reactions with higher water concentrations, the decay of the ascorbyl radical is more appreciable and had to be treated explicitly when fitting the data. Above $\chi_{\text{water}} = 0.012$, the decay is too rapid to extract meaningful kinetic results. Fortunately, the isopropylidene derivative *i*Asc^{•-} decays more slowly so that experiments up to $\chi_{\text{water}} = 0.028$ could be analyzed. The kinetic data were globally fit with a model of a second-order approach-to-equilibrium step followed by the second-order decay of the ascorbyl radical (Scheme 2).²² In all cases, the spectrum of ascorbyl radical was fixed in SPECFITTM to the known spectrum in pure acetonitrile,¹⁰ which is reasonable since the λ_{max} and ϵ of the radicals are similar in pure MeCN (377 nm/3900 ± 200 M⁻¹ cm⁻¹)¹⁰ and H₂O (360 nm/3500 ± 200 M⁻¹ cm⁻¹).²³

Equilibrium constants for these reactions with added water are also determined using Scheme 2, with K_{eq} given by the ratio of the rate constants k_1/k_{-1} . This is best done for reactions where a limited amount of TEMPO is used. For reactions with $\chi_{\text{water}} < 0.003$, analysis with the model in Scheme 2 gives the same values of k_1 and K_1 as those from the pseudo-first-order plots and plots of [TEMPOH][Asc^{•-}]/[AscH⁻] versus [TEMPO] in MeCN (Table 1). The derived rate and equilibrium constants for reactions with added water are given in Tables 2 and 3 and shown graphically in Figure 2. To pick one example, the rate constant for the *i*AscH⁻ reaction is a remarkable three times slower in the presence of just $\chi_{\text{water}} = 0.014$, and the equilibrium constant is 8 times smaller.

IV. Effects of other additives on rate and equilibrium constants

Additives other than water also have a large effect on the rate and equilibrium constants for reaction 1 in MeCN. These experiments were done with *i*AscH⁻ for experimental convenience (see above). We have examined the effects of adding: ethylene glycol (HOCH₂CH₂OH), glycerol [HOCH₂CH(OH)CH₂OH], dimethoxyethane (DME, MeOCH₂CH₂OMe), diethanolamine [(HOCH₂CH₂)₂NH], *N*-acetyl-leucine-methyl ester [MeC(O)NHCH(*i*Bu)C(O)OMe], DMSO, choline chloride (HOCH₂CH₂NMe₃⁺Cl⁻), tetramethylammonium chloride (Me₄N⁺Cl⁻), and, as noted above, ^{*n*}Bu₄N⁺PF₆⁻. These were chosen to include both hydrogen bond donors and acceptors, molecules with biological-type functional groups, and charged species. Acidic compounds, such as NH₄⁺Cl⁻, cannot be used because they would protonate the ascorbate substrate under these conditions. For each additive, k_{1i} and K_{1i} were determined as described above.

Most of the additives cause a decrease rate and equilibrium constants, consistent with the effect of added water (Table 4 and Figure 3). The exceptions are dimethoxyethane, diethanolamine, Me₄N⁺Cl⁻, and ^{*n*}Bu₄N⁺PF₆⁻, which have little or no effect on the reaction.

The lack of any change with 100 mM ${}^n\text{Bu}_4\text{N}^+\text{PF}_6^-$ is a striking contrast to the effect of only 6.5 mM choline chloride ($\chi = 3.4 \times 10^{-4}$), which decreases k_{1i} and K_{1i} by factors of 1.6 and 2.9, respectively. DMSO has a much weaker effect than other additives: at $\chi = 7.7 \times 10^{-3}$, DMSO lowers k_{1i} by a factor of 1.3 while ethylene glycol causes a 5.7-fold rate decrease.

These shifts in k_{1i} and K_{1i} presumably result from the additive binding to or in some way differentially solvating the ascorbate and/or ascorbyl radical. To probe such a binding, optical spectra of ascorbate in the presence of ethylene glycol or choline chloride have been examined, but no changes were observed. ${}^1\text{H}$ NMR spectra, however, show downfield shifts of the resonance for the proton at C4 for both AscH^- and $i\text{AscH}^-$ (see Scheme 1 for numbering), with either ethylene glycol or choline chloride. The addition of choline chloride has a qualitatively larger effect on the chemical shift at lower concentrations than ethylene glycol, consistent with the above results that choline chloride has the larger effect on k_1 and K_1 .²⁴

V. Other reactions of TEMPO, TEMPOH and ascorbate

In principle, the changes in k_1 and K_1 with increasing water or other additives could be due primarily to changes in the ascorbate/ascorbyl-radical couple, to changes in the TEMPO/TEMPOH couple, or to changes in both couples. To resolve this question, hydrogen atom transfer reactions of TEMPO, TEMPOH, and ascorbate with other reagents have been studied in MeCN as a function of added water.

1. Reactions with phenoxyl radicals— $i\text{AscH}^-$ and TEMPOH each rapidly and completely transfer H^\bullet to the 2,6-di-*tert*-butyl-4-methoxyphenoxyl radical (${}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$). In pure MeCN, both reactions have $K_{\text{eq}} = 3 \times 10^5$ based on the relevant BDFEs.²⁵ The rate constants in MeCN with various amounts of added water are shown in Figure 4 and given in Supporting Information Table S1. For TEMPOH + ${}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$, the rate constant at $\chi_{\text{water}} = 0.028$ is slower than that in pure MeCN by only 13%. Similarly, the rate constant for TEMPOH + ${}^t\text{Bu}_3\text{PhO}^\bullet$ ($K_{\text{eq}} = 5 \times 10^7$)¹⁷ is reduced by only 15% at $\chi_{\text{water}} = 0.028$. In contrast, k_{1i} for TEMPO + $i\text{AscH}^-$ decreases by a factor of 7 at this water concentration (Table 3). The mild dependence of the TEMPOH rate constants on small amounts of water is the expected behavior for a hydrogen atom transfer reaction (see below). These reactions do not show the unusual solvent effect.

In contrast, reactions of $i\text{AscH}^-$ with ${}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$ are substantially slowed by added water, as shown in Figure 4. For instance, this reaction is slower by 38% at $\chi_{\text{water}} = 0.0038$ vs. pure MeCN. This reduction is remarkably similar to the decrease of 35% for TEMPO + $i\text{AscH}^-$ at the same χ_{water} . This water concentration is *seven times less* than amount that causes only 13–15% in the TEMPOH reactions. In sum, the reaction of ascorbate with ${}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$ shows the same unusual slowing with added water but the reactions of TEMPOH do not. The similarity between the water effects for the two reactions of $i\text{AscH}^-$, with ${}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$ and with TEMPO, suggests that this effect is due to the ascorbate rather than the hydrogen abstractor.

2. Hydrogen atom transfer from a ruthenium complex to TEMPO—Added water could also potentially slow k_1 and k_{1i} by forming a hydrogen bond to the $\text{R}_2\text{N}-\text{O}^\bullet$ moiety of TEMPO, which is a moderately strong hydrogen bond acceptor.²⁶ To test this possibility, we have explored the effect of added water on the ability of TEMPO to remove H^\bullet from $(\text{acac})_2\text{Ru}^{\text{II}}(\text{py-imH})$ ²⁷ ($\text{acac} = 2,4\text{-pentanedioato}$, $\text{py-imH} = 2\text{-(2'-pyridyl)imidazole}$). This reaction has been previously examined in pure MeCN, where it has $\Delta G^\circ = -4.4 \pm 0.1$ kcal mol^{-1} ($K_{\text{eq}} = 1800 \pm 200$) and $k = (1.4 \pm 0.1) \times 10^3$ $\text{M}^{-1} \text{s}^{-1}$.^{27b} Between $\chi_{\text{water}} = 3 \times 10^{-5}$ and 7.2×10^{-3} the measured rate constant is within the 7% error bars of the value in pure

MeCN (see Supporting Information). At $\chi_{\text{water}} = 2.8 \times 10^{-2}$, there is a decrease in k of a factor of 1.4. In contrast, this concentration of water reduces k_i by a factor of 7.2 (Table 3).

3. Hydrogen atom transfer from ${}^t\text{Bu}_2\text{NOH}$ to TEMPO—The effect of added water on the reaction of TEMPO with ${}^t\text{Bu}_2\text{NOH}$ has also been explored. Experimentally, this reaction is most conveniently studied in the reverse direction, ${}^t\text{Bu}_2\text{NO}^\bullet + \text{TEMPOH}$, but since the reaction is close to isoergic the approach-to-equilibrium kinetics yield the forward and reverse rate constants as well as the equilibrium constant. In pure MeCN, based on the reported results,²⁸ TEMPO + ${}^t\text{Bu}_2\text{NOH}$ has $\Delta G^\circ = -1.3 \pm 0.1 \text{ kcal mol}^{-1}$ ($K_{\text{eq}} = 9.0 \pm 1.5$) and $k = 17.3 \pm 3.6$. We find that addition of water to $\chi = 1.4 \times 10^{-2}$ has essentially no effect on either the forward or the reverse rate constants, or the equilibrium constant. At $\chi_{\text{water}} = 2.8 \times 10^{-2}$ there is no change rate constant for TEMPO + ${}^t\text{Bu}_2\text{NO}^\bullet$, although k for the reverse reaction and the equilibrium constant both decrease by about 15%. Thus neither the reaction of TEMPO with $(\text{acac})_2\text{Ru}^{\text{II}}(\text{py-imH})$ nor TEMPO + ${}^t\text{Bu}_2\text{NOH}$ show the substantial effect of added water that is observed for TEMPO + ascorbates. This effect of additives is thus indicated to be a property of ascorbate rather than TEMPO.

Discussion

I. Overview

Most biochemical reactions of ascorbate, as is known^{4,7} but not always appreciated, initially form the ascorbyl radical and therefore involve loss of e^- and H^+ or, equivalently, H^\bullet . These reactions are best described as hydrogen atom transfer (or concerted proton-electron transfer²⁵). The results reported here show that small changes in the reaction medium cause surprisingly large changes in the kinetics and thermodynamics of the H-atom transfer reactions of ascorbates with TEMPO and with ${}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$. The reaction of the potassium-18-crown-6 salt of ascorbate with TEMPO is 1.9 times slower and 4.0 times less favorable in MeCN with *ca.* 1 mol% water (~200 mM water), versus the reaction in pure MeCN. With the more experimentally tractable isopropylidene ascorbate derivative, $[\text{DBU-H}]^+i\text{AscH}^-$, the presence of this amount of water decreases k_i and K_{1i} by factors of 3.3 and 9.6, respectively. This effect is not limited to water: the addition of *ca.* 1 mol% ethylene glycol, choline chloride, or *N*-acetyl-leucine-methyl ester to the MeCN solvent all cause k_i to fall by more than a factor of two and K_{1i} to drop by at least a factor of eight.

The results reported show that the changes observed in rate and equilibrium constants with various additives are due to changes in the reactivity of ascorbate, not TEMPO or TEMPOH. The reactions of $\text{AscH}^- + \text{TEMPO}$, $i\text{AscH}^- + \text{TEMPO}$, and $\text{AscH}^- + {}^t\text{Bu}_2(\text{MeO})\text{ArO}^\bullet$ all show this remarkable effect. In contrast, none of the reactions of TEMPO or TEMPOH show substantial changes in their rate or equilibrium constants upon addition of water, except for the reactions of TEMPO with the ascorbates. The TEMPO/TEMPOH couple appears to behave like a typical oxyl radical, while the ascorbate/ascorbyl radical couple is anomalous. This conclusion is supported by the unusual solvent dependence of its bond dissociation free energy of ascorbate, as discussed above and in the Supporting Information).

II. The kinetic and thermodynamic results indicate that ascorbate is strongly affected by local solvation

These large variations in rate and equilibrium constants for reaction 1 cannot be explained by variation of the bulk properties of the solvent, such as the dielectric constant ϵ , nor by a common kind of kinetic solvent effect observed for other HAT reactions (see below). The presence of water in the small amounts used here ($\chi_{\text{water}} < 0.03$) has virtually no effect on ϵ .²⁹ Furthermore, the effects of the other additives do not follow that expected for changes ϵ .

For example, the dielectric constants of pure MeCN (37.5) and ethylene glycol (37.7) are nearly identical,³⁰ yet a small amount of ethylene glycol in the MeCN solvent ($\chi = 7.0 \times 10^{-4}$, 13.5 mM) reduces k_{1i} by 32% and K_{1i} by 48%. With the ionic material choline chloride ($\text{HOCH}_2\text{CH}_2\text{NMe}_3^+\text{Cl}^-$), addition of only 13 equivalents ($\chi = 3.4 \times 10^{-4}$, 6.5 mM) reduces k_{1i} and K_{1i} by factors of 1.6 and 2.9. This is not due to an ionic strength or ion pairing effect, since a much larger concentration of an inert salt, 100 mM $^n\text{Bu}_4\text{N}^+\text{PF}_6^-$, has no effect on the rate or equilibrium constants. The equilibrium constant for $i\text{AscH}^- + \text{TEMPO}$ decreases by 1,800% as water concentration is raised to $\chi_{\text{water}} = 0.028$. This decrease is vastly larger than would be predicted from ideal solution behavior, since the reaction is 2.5 kcal mol⁻¹ less favorable in pure H₂O vs. pure MeCN (see below) and 2.8% of 2.5 kcal mol⁻¹ would imply a barely noticeable change in K_{1i} of 12%.

If the large variations in rate and equilibrium constants with small amounts of additives are not due to changes in the bulk properties of the solvent, they must be due to differences in the local environment that the H-transfer reaction occurs in. As shown below, it is the presence of H-bond donors in the local environment that are causing the observed effect.

Solvent effects on hydrogen atom transfer reactions have been examined in many systems, particularly the effects of solvent polarity and viscosity on selectivity.³¹ For HAT reactions involving C-H bonds, solvent effects are usually very small³² (which is why gas phase C-H bond strengths can be used to rationalize reactions in solution). For hydroxylic substrates (ROH), Ingold, Litwinienko and co-workers have found substantial kinetic solvent effects (KSEs) due to hydrogen bonding from the substrate to the solvent, $\text{ROH}\cdots\text{Solv}$.³³ Their 'KSE model' indicates that only the non-hydrogen bonded substrate is reactive toward HAT, so reactions are slower in solvents that are better hydrogen bond acceptors. They gauge the H-bond accepting ability of solvent using the hydrogen bond basicity parameters (β_2^{H}) developed by Abraham.³⁴ This model does predict the slower rates for reaction 1 in the better hydrogen bond accepting DMSO ($\beta_2^{\text{H}} = 0.78$) versus MeCN ($\beta_2^{\text{H}} = 0.44$) (Tables 2, 3). However, the large rate deceleration with the addition of small amounts of water is not consistent with the KSE model. Since MeCN is a slightly better H-bond acceptor than H₂O ($\beta_2^{\text{H}}(\text{MeCN}) = 0.44 > \beta_2^{\text{H}}(\text{H}_2\text{O}) = 0.38$), the KSE model predicts that H-transfer rate constants should be unaffected by small amounts of water. These results show that the major effects of additives on the reactions of ascorbate are not due to a hydrogen bonding interaction that blocks the transfer of the ascorbate hydrogen. If this were the case, ethylene glycol and dimethoxyethane would be expected to show very similar effects on the rate constants (k_1). Thus, ascorbate shows a different type of solvent effect on its HAT reactivity that is distinct from the Ingold kinetic solvent effect model for hydrogen atom transfer reactions.

It is striking that small amounts of water and other additives affect *both* the rates of reaction and the equilibrium constants. In fact, the equilibrium constants change substantially more than the rate constants. This contrasts with the KSE model, which does not correlate with the (ground state) reaction thermodynamics because it involves hydrogen bonding only with the reactants and not with the products. Brønsted plots of $\ln(k)$ vs. $\ln(K)$ for the ascorbate and isopropylidene ascorbate reactions at different water concentrations are linear (Figure 5). Figure 5b also includes values for all of the other additives, showing that water and the other materials all behave very similarly – a remarkable and unexpected result. The slopes of the plots are the Brønsted α values, or equivalently, $\Delta\Delta G^\ddagger/\Delta\Delta G^\circ$. For AscH^- , $\alpha = 0.48 \pm 0.03$. For $i\text{AscH}^-$, the slope is 0.61 ± 0.04 including all additives, and a very similar 0.60 ± 0.05 for just the MeCN + H₂O reactions of $i\text{AscH}^-$. A combined plot with all the data shows a very strong linear trend among all the data ($i\text{AscH}^-$ and AscH^-), with a slope of 0.58 ± 0.04 and a correlation coefficient of 0.96 (Figure 5c).

Rate/driving force relationships in organic H-atom transfer reactions have long been studied,³⁵ although we are unaware of an example such as this with small amounts of additives. Traditionally these correlations have used reaction enthalpies, but we have found that free energies are more appropriate.⁹ In addition, we have found that a Marcus theory formalism connecting k and K is very valuable for understanding HAT reactions.^{36,37} For reactions such as these with small driving forces ($\Delta G^\circ \ll \lambda/2$), Marcus theory predicts a linear Brønsted plot with a slope of 0.5,³⁸ which is in excellent agreement with the observed results. *Thus, the observed changes in rate constant stem predominantly from the changes in the driving force of the reactions.*

III. Local solvation or binding shifts the thermochemistry of ascorbate

Small amounts of water and other materials strongly shift the equilibrium constant for the reaction of ascorbate and isopropylidene ascorbate with TEMPO. As shown in the previous section, this is due to the local solvation or binding of the additive to the ascorbate and/or ascorbyl radical. The diverse range of additives gives insight into the nature of this binding or solvation. Ethylene glycol (HOCH₂CH₂OH) at 0.8% markedly lowers both k_{1i} and K_{1i} (factors of 5.7 and 20, respectively) while dimethoxyethane (DME, MeOCH₂CH₂OMe) has essentially no effect at a similar mole fraction. Ethylene glycol and DME are similar in size are comparable hydrogen bond acceptors,³⁹ but only the former can be a hydrogen bond donor. Thus, it is the hydrogen bond *donating* character of the material that is critical. This is again distinct from Ingold's kinetic solvent effect observations, where the hydrogen bond *accepting* character of the solvent is the key contributor to the rate constant for hydrogen atom transfer. The importance of hydrogen bond donation is also indicated by the very small effect of DMSO, a good hydrogen bond acceptor, and by the much larger effect of choline chloride (HOCH₂CH₂NMe₃⁺Cl⁻) vs. Me₄N⁺Cl⁻. All of the results reported here are consistent with hydrogen bond donation from the additive to ascorbate playing the key role in the observed changes in k and K .

The effect of additives on the equilibrium constants can be analyzed using a thermochemical cycle such as Scheme 3, although this is probably a simplification because the binding is probably not well described by a 1:1 model. In this model, the difference in the measured equilibrium constants, K_{1A}/K_1 , is equal to the difference in binding constants for the additive A with AscH⁻ and Asc^{•-} ($K_{\text{Asc}^\bullet}/K_{\text{AscH}^-}$). The binding constants K_{AscH^-} and K_{Asc^\bullet} in this analysis are analogous to the K_d values for AscH⁻ and Asc^{•-} binding to a protein (see next section). Qualitatively, when an additive (or enzyme binding pocket) binds ascorbate more tightly than the ascorbyl radical, then the hydrogen atom transfer step will be less favorable ($K_{1A} < K_1$). Since these equilibrium constants are related to the O–H bond dissociation free energy (BDFE) of the ascorbate, hydrogen bond donating environments effectively increase the BDFE. A more detailed discussion of BDFEs is given in the Supporting Information.

The difference in binding to AscH⁻ and Asc^{•-} is presumably due to the difference in charge distribution in the two anions, as indicated in the drawings in Scheme 1 above. Ascorbate has a single enolate oxygen that formally carries the negative charge, while the ascorbyl radical has a more delocalized triketone-like structure. This is reflected in the difference in basicity of AscH⁻ and Asc^{•-} (pK_a of the conjugate acids 4.0 and -0.45 , respectively^{6,8}). This issue has been addressed by Costanzo *et al.* using DFT calculations and molecular dynamics simulations. They show that all three exocyclic oxygen atoms carry significant negative charge in both AscH⁻ and Asc^{•-}, with all the atomic charges being smaller in the radical.⁴⁰ Molecular dynamics simulations indicate that in aqueous solution, the enolate oxygen of AscH⁻ is the preferred hydrogen bonding site. Oxidation to the radical causes the loss of 0.6 of a hydrogen bond at this oxygen, consistent with the change in calculated charges. Thus, these calculations appear to support the conclusion that hydrogen bond donors bind more strongly to AscH⁻ than to Asc^{•-}.

Two other reports have described unusual solvent effects on hydrogen transfer reactions of ascorbate.^{16,41} The reduction of PhNO by ascorbate is 40% slower in 1:1 water:dioxane than that in pure water, which is twice as large as the maximum KSE-predicted shift in k from pure water to pure dioxane.⁴¹ In addition, the H₂O/D₂O kinetic isotope effect increases from 2.4 to 8.5 from water to water/dioxane. While these effects could be a result of bulk polarity changes, it seems likely that the issues of local solvation implicated here are also playing a role. A very recent study of the reduction of TEMPO by ascorbate in water:dioxane mixtures reported large kinetic isotope effects indicative of hydrogen tunneling.¹⁶ Tunneling has also been implicated in other HAT reactions of TEMPO.^{27b,28} It is not clear how tunneling is contributing to the local solvent effect discussed here. However, it seems unlikely that tunneling effects play a dominant role given the very strong Brønsted correlations between rate and equilibrium constants. We also note that Amorati *et al.* have also reported a solvent effect that is distinct from Ingold's KSE model.⁴² They found that the kinetics and thermodynamics of HAT from hydroquinones to 2,2'-diphenylpicrylhydrazyl radical (DPPH) or peroxy radicals are affected by hydrogen bonding to the non-transferring hydrogen of the hydroquinone. Such a "remote" hydrogen bonding effect may be analogous to what is reported here, but their study involved hydrogen bond acceptors while the ascorbate reactions above are modulated by hydrogen bond donors.

IV. Biological Implications

The results reported here show that changes in the hydrogen-bond donating environment can have a substantial impact on the reactivity of ascorbate. In particular, hydrogen bond donating environments increase the ascorbate O–H bond strength, making ascorbate a poorer reductant. Given the wide range of biological processes involving ascorbate, it seems likely that Nature has taken advantage of this apparently unique feature. For example, do ascorbate binding pockets in certain enzymes have hydrogen bond donating or hydrogen bond accepting character that would thermodynamically favor formation of ascorbate or ascorbyl, respectively? Below we briefly discuss three *in vivo* processes in which modulation of the thermochemistry of ascorbate by the local H-bonding environment could play an important role.

Cytochrome b_{561} is a transmembrane protein, found in both plants and animals, that is thought to mediate ascorbate recycling across the membrane (Figure 6a).^{43,44} *In vitro* studies show that it can catalyze the reduction of the ascorbyl radical to ascorbate. In mammalian chromaffin b_{561} , the putative Asc^{•−} binding site on the intravesicular side of the membrane is fairly hydrogen bond donating, with the sequence −xYSLHSWxGx− (x is a hydrophobic residue).⁴³ On the extravesicular side of the lipid bilayer, the putative AscH[−] binding site has fewer hydrogen bond donating groups (−ALLVYRVFR−), as well as a well conserved lysine residue (Lys85), proposed to play an electrostatic role in binding.⁴³ Based on the results presented here, the H-bonding environment of the intravesicular Asc^{•−} binding pocket thermodynamically favors formation of AscH[−], and the extravesicular AscH[−] binding pocket favors the reverse reaction. While the mechanism of AscH[−]/Asc^{•−} interconversion may be more complicated than the simple HAT reactions studied here,⁴⁵ it is likely that the kinetic barriers reflect the energetics, so this effect should facilitate the reduction of Asc^{•−} on the intravesicular side of the membrane and the oxidation of AscH[−] on the extravesicular side.

Ascorbate peroxidase (APX) has an ascorbate binding pocket that is well characterized.⁴⁶ It contains several hydrogen bonding interactions (*e.g.* Lys30, Arg172 and the heme-propionate), and is fairly exposed to solvent. This pocket thus appears to hold ascorbate in an environment closer to that of aqueous solution, with a high bond strength (high one electron/one-proton redox potential). The aqueous 1H⁺/1e[−] ascorbate potential is 0.33 V at

pH 7,⁸ somewhat above the outer-sphere $\text{Fe}^{\text{II/III}}$ potential of the heme in APX of +0.206 V.⁴⁷ Since the catalytic cycle of APX requires an iron(III) resting state,⁴⁸ it is important that the bound ascorbate not be any more reducing than aqueous ascorbate. Thus, the enzyme has apparently evolved to have a strongly hydrogen bond donating binding site for ascorbate to avoid reduction of the resting ferric state.

The ascorbate reduction of the α -tocopheroxyl radical, to give the ascorbyl radical and α -tocopherol (vitamin E), is perhaps one of the most discussed and studied reactions of ascorbate. This simple HAT reaction is very well documented *in vitro*,^{49,50} though its *in vivo* importance is debated.⁵¹ α -Tocopherol is membrane bound, likely with its phenolic moiety near the water-lipid interface.⁴⁹ Conversely, ascorbate is hydrophilic and does not significantly partition into lipids, although there is evidence that it can localize in the polar lipid head groups at the surface of the membrane.⁵² This membrane surface region is polar, but less hydrated and contains fewer H-bond donors than the cellular bulk (Figure 6b).⁵³ Based upon our results, ascorbate at the surface of the membrane is predicted to act as a stronger H-atom donor (reductant) to tocopheroxyl radical than ascorbate in aqueous solution.⁵⁴ In addition, the O–H bond in α -tocopherol should be a little stronger in the polar head group region than in the bulk lipid.⁵⁵ Thus the ‘intermediate’ region at lipid-water interfaces appears to be an ideal location for Nature to maximize the driving force for, and therefore rate of, H-transfer from AscH^- to α -tocopheroxyl. Rapid reduction of α -tocopheroxyl is valuable to minimize its activity as a pro-oxidant.⁵⁶

Conclusions

Rate and equilibrium constants for the oxidation of ascorbate by the aminoxyl radical TEMPO (eq 1) in acetonitrile are very sensitive to small amounts of various additives. The results are similar for the potassium 18-crown-6 salt of ascorbate (AscH^-) and the more soluble DBU- H^+ -isopropylidene ascorbate ($i\text{AscH}^-$). Addition of water up to 1–3 mole% causes decreases of factors of 2–7 in k_1 and almost 4–20 in K_1 . For AscH^- , the 4.5 fold decrease in K_1 at only 1.2 mole% water is a quarter of the total change from pure MeCN to pure water. Other hydrogen bond donating additives such as glycerol, choline chloride, and *N*-acetyl-leucine-methyl ester have similar effects. On the other hand, the hydrogen bond acceptors dimethoxyethane and DMSO do not significantly affect the reaction outside of normal kinetic solvent effects. Thus, the changes in rate and equilibrium constants are much larger than can be explained by changes in bulk solvent properties such as the dielectric constant or ionic strength. In addition, since the changes in k and K_{eq} are due to hydrogen bond donors not acceptors, this effect is clearly distinct from Ingold’s kinetic solvent effect model for hydrogen atom transfer reactions.

The changes in rate constant directly parallel the changes in equilibrium constant, with Brønsted plots, $\ln(k)$ vs. $\ln(K)$, being linear with slopes close to the theoretical value of 0.5 (0.48 for AscH^- and 0.61 for $i\text{AscH}^-$). Therefore the changes in rate are predominantly a consequence of the changes in the driving force for the reactions. This contrasts with previously described kinetic solvent effects in oxyl radical reactions, which are purely kinetic effects due to hydrogen bonding to the transferring hydrogen atom. The results show that ascorbate is a stronger reductant – better able to donate a proton and an electron (or H^\bullet)^{4,7,8} – in local environments with less hydrogen bond donors. The change in reducing power of ascorbate is also indicated by the change in bond strength (BDFE) in pure solvents (Table 1). The unusually large solvation/binding effect of ascorbate is suggested to derive from the substantial differences in charge distribution between ascorbate and the more delocalized ascorbyl radical. This study also indicates that the reactivity of ascorbate can, in many cases, be best understood using the free energy for loss of both a proton and an electron (H^\bullet).

The tuning of the properties of ascorbate by its local environment is likely to be of importance in many of the biological actions of ascorbate. Polar, protic environments favor ascorbate while environments without hydrogen bond donors favor the ascorbyl radical. Three examples illustrate different ways in which Nature may have utilized this unique property, to facilitate ascorbate recycling by cytochrome *b*₅₆₁, to avoid unwanted side effects in ascorbate peroxidase, and in the regeneration of α -tocopherol by ascorbate. As new reactivity and structural information emerges about ascorbate-utilizing proteins, specific hydrogen bonding interactions should be considered with respect to the reactivity of the enzyme.

Methods

Physical Techniques and Instrumentation

UV-Vis stopped flow measurements were obtained using an OLIS RSM-1000. ¹H NMR spectra were recorded on a Bruker Avance 300 MHz or 500 MHz spectrometers. All reactions were performed in the absence of air using standard glovebox and high vacuum techniques.

Materials

Reagent grade solvents were purchased from Fischer Scientific, unless otherwise noted. All other chemicals were purchased from Aldrich. Anhydrous acetonitrile was purchased from Honeywell Burdick and Jackson, sparged with Ar and plumbed directly into an N₂-filled glovebox with stainless steel tubing. The supplier specification for water content in the MeCN is 5 ppm ($\chi_{\text{water}} = 1.5 \times 10^{-5}$ M); our independent measurement by Karl Fischer titration was 10 ppm. Benzene and DMSO were dried using a Grubbs-type catalyst in a Seca solvent system. Deuterated solvents were obtained from Cambridge Isotope Laboratories. TEMPOH, 2,4,6-tri-*tert*-butyl phenoxyl, and 2,6-di-*tert*-butyl-4-methoxy phenoxyl were prepared according to the literature.²⁵ (acac)₂Ru^{II}(py-imH) was prepared as previously described.^{27a} Deionized water was distilled before use in kinetic experiments. 18-crown-6 was recrystallized from acetonitrile and dried under vacuum. Choline chloride was recrystallized from absolute ethanol and dried under vacuum. Glyme was freshly distilled from Na/benzophenone ketyl. *N*-acetyl-leucine-methyl ester was prepared from leucine-methyl ester (Aldrich) by the method of Applewhite,⁵⁷ recrystallized from ethanol/pentane and stored in an N₂-filled glovebox. Ethylene glycol was distilled from MgSO₄, then a small amount of Na⁰ was added and the solvent stirred overnight, and finally was distilled freshly before use. Glycerol was added to an equal volume of *n*-butanol, and rotated slowly in an ice bath until crystals formed. The crystals were collected by vacuum filtration, washed with ice-cold acetone and dried under vacuum. Diethanolamine was distilled and crystallized from its melt.

Preparation of potassium 18-crown-6 ascorbate

Potassium ascorbate was prepared according to the literature.¹³ A swivel-frit assembly was charged with 0.5 g (2.3 mmol) potassium ascorbate and 0.6 g (2.3 mmol) 18-crown-6. The system was evacuated and approximately 20 mL CH₃CN was freshly vacuum transferred from CaH₂. The resulting yellow solution was stirred for 1 hour and filtered. The solvent was removed under vacuum to give a yellow solid, which was recrystallized from CH₃CN/Et₂O and washed once with ~10 mL Et₂O. ¹H NMR (DMSO-*d*₆) 3.24 (m, 1H), 3.34 (m, 1H), 3.45 (m, 1H), 3.55 (s, 24 H), 3.85 (d, 1H).

Stopped-Flow Kinetics

In a typical procedure, solutions of *i*AscH₂ + DBU (0.68 mM each) and TEMPO (1.3–71 mM) in MeCN were prepared and loaded into gastight syringes inside a N₂-filled glovebox. Mixtures containing additives, except for water, were also prepared in the glovebox. The syringes were removed from the glovebox individually for each stopped-flow run. When necessary, freshly distilled and de-gassed water was quickly transferred from an N₂-filled 2-neck flask to both syringes (*i*AscH⁻ and TEMPO) via a gas tight μL syringe. The stopped-flow syringes were capped and inverted twice to mix thoroughly and immediately placed onto the stopped-flow. A minimum of 5 kinetic runs were collected at each [TEMPO]. The data were analyzed using SPECFIT™ software as described in the text.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The U.S. National Institutes of Health (GM50422) and the University of Washington, Department of Chemistry are gratefully acknowledged for financial support.

References

1. Packer, L.; Fuchs, J., editors. Vitamin C in Health and Disease. Vol. 5. Marcel Dekker; New York: 1997.
2. Reed, DJ. Interaction of vitamin E, ascorbic acid and glutathione in protection against oxidative damage. In: Packer, L.; Fuchs, J., editors. Vitamin E in Health and Disease. Marcel-Dekker; New York: 1993. p. 269-281.
3. Kroneck, PMH.; Armstrong, FA.; Merkle, H.; Marchesini, A. Ascorbate oxidase: molecular properties and catalytic activity. In: Seib, PA.; Tolbert, BM., editors. Ascorbic Acid: Chemistry, Metabolism and Uses. American Chemical Society; Washington, D.C: 1982. p. 223-248. *Advances in Chemistry Series 200*
4. Njus D, Wigle M, Kelley PM, Kipp BH, Schlegel HB. *Biochemistry*. 2001; 40:11905–11911. [PubMed: 11570891]
5. Kerber RC. *J Chem Ed*. 2008; 85:1237–1242.
6. Creutz C. *Inorg Chem*. 1981; 20:4449–4452.
7. Njus D, Kelley PM. *Biochim Biophys Acta*. 1993; 1144:235–248. [PubMed: 8399278]
8. Njus D, Kelley PM. *FEBS Lett*. 1991; 283:147–151. [PubMed: 1647978]
9. The BDFE is a better predictor of reactivity than the more commonly used bond dissociation enthalpy, BDE; see: Mader EA, Davidson ER, Mayer JM. *J Am Chem Soc*. 2007; 129:5153–5166. [PubMed: 17402735]
10. Warren JJ, Mayer JM. *J Am Chem Soc*. 2008; 130:7546–7547. [PubMed: 18505256]
11. Warren JJ, Mayer JM. *J Am Chem Soc*. 2008; 130:2774–2776. [PubMed: 18257574]
12. Bielski BHJ, Allen AO. *J Am Chem Soc*. 1970; 92:3793–3794. [PubMed: 5422775]
13. Tajmir-Riahi HA, Boghai DM. *J Inorg Biochem*. 1992; 45:73–84.
14. Binstead, RA.; Zuberbühler, AD.; Jung, B. Specfit, Version 3.0.38 (32-bit Windows). Spectrum Software Associates; Chapel Hill, NC: 2006.
15. Bielski BHJ, Allen AO, Schwarz HA. *J Am Chem Soc*. 1981; 103:3516–3518.
16. Sajenko I, Pilepić V, Brala CJ, Sajenko I, Uršić. *J Phys Chem A*. 2010; 114:3423–3430. [PubMed: 20151699]
17. Mader EA, Manner VW, Markle TF, Wu A, Franz JA, Mayer JM. *J Am Chem Soc*. 2009; 131:4335–4345. [PubMed: 19275235]
18. The BDFE in DMSO is derived from the known ^{18a}*E*_{1/2} and p*K*_a values using BDFE(X–H) = 1.37p*K*_a + 23.06*E*_{1/2} + *C*_G, with *C*_G = 71.1 kcal mol⁻¹ (ferrocene⁺⁰ electrochemical reference).

- 9,^{18b}(a) Bordwell FG, Liu W-Z. *J Am Chem Soc.* 1996; 118:10819–10823.(b) Mader, EA. Ph.D. Thesis. University of Washington; 2007. Hydrogen atom transfer reactions of iron and cobalt alpha-diimines: A study of intrinsic and thermodynamic effects.
19. Tilset, M. The thermodynamics of systems involving electron-transfer paths. In: Balzani, V., editor. *Electron Transfer in Chemistry*. Vol. 2. Wiley-VHC; New York: 2001. p. 677-713.
20. As shown in reference ¹⁰, the BDFE(*i*AscH⁻)_{MeCN} derived from p*K*_a and *E*_{1/2} values is 64.4 ± 2.0 kcal mol⁻¹. The large difference in aqueous and MeCN BDFEs is therefore not an artifact of the method of BDFE determination (from *K*₁ vs. p*K*_a and *E*_{1/2}).
21. A more detailed discussion of Δ*G*_{solv}^o(H[•]) in different solvents can be found in reference ^{18b}. Briefly, following Rounder,^{21a} Δ*G*_{solv}^o(H[•]) is well approximated by Δ*G*_{solv}^o(H₂), which is (in kcal mol⁻¹), 5.12 in MeCN, 5.61 in DMSO, and 8.98 in water.^{18b} Thus, the solvation of H[•] contributes +3.9 kcal mol⁻¹ to the difference in BDFEs between MeCN and H₂O (3.4 kcal mol⁻¹ for DMSO vs. H₂O). This is only one aspect of the solvation, however, and Δ*G*_{solv}^o(XH) is not necessarily close to Δ*G*_{solv}^o(X[•]) for hydroxylic substrates ROH such as ascorbate which can form strong hydrogen bonds with H-bond accepting solvents.
22. The second step, Asc^{-•} disproportionation, was fit in the kinetic model as bimolecular process but could mechanistically be more complicated. The derived values for *k*₁ and *k*₋₁ do not appear to be very sensitive to the value of the disproportionation rate constant.
23. The *λ*ε given for aqueous ascorbate is the average of the values reported in reference ¹² and in: Schuler RH. *Radiat Res.* 1977; 69:417–433. [PubMed: 847094]
24. Similar NMR shifts are observed for *i*AscH⁻ + choline chloride. A plausible quantitative analysis of the AscH⁻ ¹H NMR shifts is given in the Supporting Information.
25. Waidmann CR, Zhou X, Tsai EA, Kaminsky W, Hrovat DA, Borden WT, Mayer JM. *J Am Chem Soc.* 2009; 131:4729–4743. [PubMed: 19292442]
26. Frachi P, Lucarini M, Pedrielli P, Pedulli GF. *CHEM PHYS CHEM.* 2002; 3:789–793. [PubMed: 12436906]
27. (a) Wu A, Masland J, Swartz RD, Kaminsky W, Mayer JM. *Inorg Chem.* 2007; 46:11190–11201. [PubMed: 18052056] (b) Wu A, Mayer JM. *J Am Chem Soc.* 2008; 130:14745–14754. [PubMed: 18841973]
28. Wu A, Mader EA, Datta A, Hrovat DA, Borden WT, Mayer JM. *J Am Chem Soc.* 2009; 131:11985–11997. [PubMed: 19618933]
29. Jellema R, Bulthuis J, van der Zwan G. *J Mol Liq.* 1997; 73–74:179–193.
30. Riddick, JA.; Bunger, WB. *Organic Solvents: Physical Properties and Methods of Purification*. In: Weissberger, A., editor. *Techniques of Chemistry Series*. Vol. 2. Wiley-Interscience; New York: 1970.
31. Tanko, JM.; Suleman, NK. *Solvent Effects in the Reactions of Neutral Free Radicals*. In: Martinho Simões, JA.; Greenburg, A.; Liebman, JF., editors. *Energetics of Organic Free Radicals*. Vol. 4. Blackie; Glasgow: 1996. p. 244-293. SEARCH series
32. Avila DV, Brown CE, Ingold KU, Luszyk J. *J Am Chem Soc.* 1993; 115:466–470.
33. Litwinienko G, Ingold KU. *Acc Chem Res.* 2007; 40:222–230. and references therein. [PubMed: 17370994]
34. Abraham MH, Grellier PL, Prior DV, Morris JJ, Taylor PJ. *J Chem Soc Perkin Trans 2.* 1990; 1990:521–529.
35. (a) Ingold, KU. *Free Radicals*. Kochi, JK., editor. Vol. 1. Wiley-Interscience; New York: 1973. p. 69-70.(b) Russell, GA. *Free Radicals*. Kochi, JK., editor. Vol. 1. Wiley-Interscience; New York: 1973. p. 283-293.
36. Mayer JM. *Annu Rev Phys Chem.* 2004; 55:363–90. [PubMed: 15117257]
37. Roth JP, Yoder JC, Won T-J, Mayer JM. *Science.* 2001; 294:2524–2526. [PubMed: 11752572]
38. Marcus RA. *J Phys Chem.* 1968; 72:891–899.
39. Alkyl ethers and primary alcohols have essentially identical hydrogen bond basicities (β₂^H). See reference ³⁴.
40. Costanzo F, Sulpizi M, Vandevondele J, Della Valle RG, Sprik M. *Molec Phys.* 2007; 105:17–23.
41. Vuina D, Pilepić V, Ljubas D, Sanković K, Sajenko I, Uršić S. *Tet Lett.* 2007; 48:3633–3637.

42. Amorati R, Franchi P, Pedulli GF. *Angew Chem Int Ed*. 2007; 46:6336–6338.
43. Verelst W, Asard H. *Genome Biol*. 2003; 4:R38.1–R38.12. [PubMed: 12801412]
44. Nakanishi N, Takeuchi F, Tsubaki M. *J Biochem*. 2007; 142:553–560. [PubMed: 17905810]
45. It has been suggested that the ascorbate oxidation reaction at the extravesicular active site utilizes an essential histidine residue that acts as a proton acceptor from ascorbate.^{4,45} If this histidine is bound to the iron that accepts the electron,^{4,45} we would still call this H-atom transfer,¹⁰ but it is possible that it could have different properties than the classical HAT from ascorbate to TEMPO described here. More detailed structural data would aid in understanding the nature of the ascorbate/ascorbyl binding pockets and the potential proton acceptors and hydrogen bonding interactions. (a) Bashtovyy D, Bérczi A, Asard H, Páli R. *Protoplasma*. 2003; 221:31–40. [PubMed: 12768339] (b) Bérczi A, Su D, Lakshminarasimhan N, Vargas A, Asard H. *Arch Biochem Biophys*. 2005; 443:82–92. [PubMed: 16256064]
46. Sharp KH, Mewies M, Moody PCE, Raven EL. *Nat Struct Biol*. 2003; 10:303–307. [PubMed: 12640445]
47. Efimov I, Papadopoulou ND, McLean KJ, Badyal SK, Macdonald IK, Munro AW, Moody PCE, Raven EL. *Biochemistry*. 2007; 46:8017–8023. [PubMed: 17580972]
48. Raven EL. *Nat Prod Rep*. 2003; 20:367–381. [PubMed: 12964833]
49. Burton GW, Ingold KU. *Acc Chem Res*. 1986; 19:194–201.
50. May JM, Qu Z, Mendiratta S. *Arch Biochem Biophys*. 1998; 349:281–289. [PubMed: 9448716]
51. Hamilton ITJ, Gilmore WS, Benzie IFF, Mulholland CW, Strain JJ. *Brit J Nutr*. 2000; 84:261–267. [PubMed: 10967604]
52. Crans DC, Baruah B, Gaidamauskas E, Lemons BG, Lorenz BB, Johnson MD. *J Inorg Biochem*. 2008; 102:1334–1347. [PubMed: 18331759]
53. Nagle JF, Tristram-Nagle S. *Biochim Biophys Acta*. 2000; 1469:159–195. [PubMed: 11063882]
54. This prediction assumes that α -tocopherol is not effected by H-bond donors in the same unusual way that ascorbate is, that the α -tocopherol BDFE is not unusually shifted by local solvation. This assumption is supported by the BDFE data in the Supporting Information, and by reports of similar solvent effects on α -tocopherol and phenol reactions in a variety of solvents: cf. Valgimigli L, Banks JT, Ingold KU, Lusztyk J. *J Am Chem Soc*. 1995; 117:9966–9971.
55. In general, homolytic bond strengths increase from non-polar to polar media. This is in part due to changes in the free energy of solvation of H^\bullet (see Supporting Information).
56. Bowry VW, Stocker R. *J Am Chem Soc*. 1993; 115:6029–6044.
57. Applewhite TH, Waite H, Neimann C. *J Am Chem Soc*. 1958; 80:1465–1469.




Figure 1. The kinetic data (\circ) for reaction of 0.28 mM AsCH^- with 8.7 mM TEMPO in DMSO and a first-order global fit (—) from SPECFITTM. Inset: Stopped-flow UV-Vis spectra over 1.1 s for this reaction; the broad absorption centered at 460 nm is due to TEMPO.




Figure 2. Plots of rate constants [red solid symbols, left axes] and equilibrium constants [blue open symbols, right axes], as a function of mole fraction of added distilled water in MeCN for the reactions of (a) ascorbate with TEMPO (k_1 , K_1) and (b) isopropylidene ascorbate with TEMPO (k_{1i} , K_{1i}).




Figure 3. Plots of rate constants [k_{1i} , plot (a)] and equilibrium constants [K_{1i} , plot (b)] for the reaction of isopropylidene ascorbate with TEMPO (eq 1) in MeCN, as a function of mole fraction of various additives. Error bars have been omitted for clarity, see Tables 2, 3, and 4.




Figure 4. Relative rate constants, $k_{\text{rel}} = k(\chi_{\text{water}})/k(\chi_{\text{water}}=0)$, as a function of mole fraction water in MeCN for TEMPOH + 2,4,6-tri-*tert*-butyl phenoxyl (○), TEMPOH + 2,4-di-*tert*-butyl-4-methoxyphenoxyl (●) and *i*AsCH⁻ + 2,4-di-*tert*-butyl-4-methoxyphenoxyl (■).




Figure 5. Plots of rate constants $\ln(k_1)$ versus equilibrium constants $\ln(K_1)$ for the reaction of (a) AsCH^- in MeCN with added water, \blacktriangle ($R^2 = 0.99$), (b) $i\text{AsCH}^-$ with TEMPO in MeCN with different additives: \blacksquare , reactions with added water ($R^2 = 0.93$); \circ , reactions with other additives, and (c) the aggregate data set from ($R^2 = 0.96$).







Figure 6. (a) Proposed structural model for bovine cytochrome b_{561} , modified from reference ⁴⁴, showing the ascorbate/ascorbyl radical conversions on both sides of the membrane. Fe = iron-heme and H = histidine. (b) Depiction of a local environment for the ascorbate reduction of α -tocopheroxyl in a lipid bilayer, as adapted from references ⁴⁹ and ⁵²; the bars at the right delineate the regions having different levels of hydration (H-bond donating water)

**Scheme 1.**

Redox chemistry of ascorbate (with the carbon numbering is shown for AscH⁻).

**Scheme 2.**

Kinetic model used to fit reaction 1 in the presence of water or other additives.

**Scheme 3.**

The effect of additives *A* on the reaction of AscH^- with TEMPO.

Table 1

Ascorbate bond dissociation free energies (BDFEs) and rate and equilibrium constants for their reactions with TEMPO (eq 1) in pure solvents.^a

Reagent	Solvent	k (M ⁻¹ s ⁻¹)	K_1	BDFE ^b
AscH ⁻	MeCN	500 ± 50	0.14 ± 0.02	67.7 ± 0.8
AscH ⁻	DMSO	100 ± 10	0.9 ± 0.1	67.6 ± 1.0
AscH ⁻	H ₂ O	2.4 ± 0.3	~1.2 × 10 ⁻³ ^c	73.6 ± 1.0
<i>i</i> AscH ⁻	MeCN	1720 ± 150	1.2 ± 0.2	66.5 ± 0.8
<i>i</i> AscH ⁻	DMSO	250 ± 20	0.85 ± 0.10	67.5 ± 1.0

^a At 298 K.

^b BDFE values in kcal mol⁻¹. Determined for equilibrium measurements with TEMPO (eq 1) except as noted.

^c Calculated from BDFE_{water}(AscH⁻) and BDFE_{water}(TEMPOH) (see below).

Table 2Rate and equilibrium constants for ascorbate (AscH^-) + TEMPO (eq 1).^a

solvent	χ_{water}	k_1	K_1
MeCN	$3 \times 10^{-5} b$	500 ± 50	0.14 ± 0.02
MeCN/H ₂ O	2.2×10^{-3}	410 ± 50	0.10 ± 0.01
MeCN/H ₂ O	3.6×10^{-3}	370 ± 40	0.080 ± 0.008
MeCN/H ₂ O	7.2×10^{-3}	280 ± 70	0.048 ± 0.005
MeCN/H ₂ O	7.2×10^{-3}	300 ± 60	0.052 ± 0.005
MeCN/H ₂ O	1.2×10^{-2}	260 ± 50	0.035 ± 0.004
H ₂ O	1	2.4 ± 0.3	ND
DMSO	$2 \times 10^{-4} b$	100 ± 10	0.9 ± 0.1

^a At 298 K. Rate constants in $\text{M}^{-1} \text{s}^{-1}$.^b χ_{water} in "dry" solvent from Karl Fischer titration.

Table 3Rate and equilibrium constants for isopropylidene ascorbate ($i\text{AscH}^-$) + TEMPO (eq 1).^a

solvent	χ_{water}	k_{1i}	K_{1i}
MeCN	3×10^{-5} ^b	1720 ± 150 ^c	1.2 ± 0.2 ^c
MeCN/H ₂ O	1.5×10^{-3}	1290 ± 100	0.65 ± 0.07
MeCN/H ₂ O	3.6×10^{-3}	1020 ± 100	0.43 ± 0.06
MeCN/H ₂ O	7.2×10^{-3}	780 ± 80	0.38 ± 0.04
MeCN/H ₂ O	7.2×10^{-3}	745 ± 80	0.25 ± 0.03
MeCN/H ₂ O	1.2×10^{-2}	525 ± 70	0.13 ± 0.02
MeCN/H ₂ O	1.4×10^{-2}	560 ± 70	0.15 ± 0.02
MeCN/H ₂ O	2.8×10^{-2}	240 ± 50	0.065 ± 0.006
H ₂ O	1	5.5 ± 0.4	ND
DMSO	2×10^{-4} ^b	250 ± 20	0.85 ± 0.10

^a At 298 K. Rate constants in $\text{M}^{-1} \text{s}^{-1}$.^b χ_{water} in "dry" solvent from Karl Fischer titration.^c Reference 10.

Table 4Rate and equilibrium constants for $i\text{AscH}^- + \text{TEMPO}$ (eq 1) in MeCN with additives.^a

Additive	χ in MeCN	k_{ti}	K_{ti}
– none – ^b	<i>b</i>	1720 ± 150	1.2 ± 0.2
ethylene glycol	7.0×10^{-4}	1170 ± 80	0.65 ± 0.09
“	7.7×10^{-3}	300 ± 40	0.085 ± 0.010
glycerol	5.1×10^{-5}	1500 ± 100	0.87 ± 0.10
dimethoxyethane	4.8×10^{-3}	1680 ± 130	1.2 ± 0.2
diethanolamine	2.7×10^{-4}	1600 ± 120	0.90 ± 0.10
<i>N</i> -AcO-Leu-OMe ^c	3.0×10^{-3}	1150 ± 100	0.35 ± 0.05
DMSO	7.7×10^{-3}	1360 ± 100	0.50 ± 0.05
“	2.8×10^{-2}	1100 ± 100	0.40 ± 0.05
choline chloride ^d	5.5×10^{-5}	1410 ± 120	1.1 ± 0.2
“	3.4×10^{-4}	1100 ± 100	0.46 ± 0.06
Me ₄ N ⁺ Cl ⁻	9.5×10^{-4}	1550 ± 150	0.9 ± 0.1
ⁿ Bu ₄ N ⁺ PF ₆ ⁻	5.2×10^{-3}	1720 ± 150	1.2 ± 0.2

^a At 298 K. Rate constants in M⁻¹ s⁻¹.^b $\chi_{\text{H}_2\text{O}} \cong 3 \times 10^{-5}$ present in ‘dry’ MeCN.^c *N*-acetyl-leucine-methyl ester (MeC(O)NHCH(ⁱBu)C(O)OMe).^d HOCH₂CH₂NMe₃⁺Cl⁻.