

SCIENTIFIC REPORTS

OPEN

Two new cellulolytic fungal species isolated from a 19th-century art collection

Carolina Coronado-Ruiz^{1,2}, Roberto Avendaño¹, Efraín Escudero-Leyva^{2,3}, Geraldine Conejo-Barboza^{4,5}, Priscila Chaverri^{2,3,6} & Max Chavarría^{1,2,4}

The archive of the Universidad de Costa Rica maintains a nineteenth-century French collection of drawings and lithographs in which the biodeterioration by fungi is rampant. Because of nutritional conditions in which these fungi grew, we suspected that they possessed an ability to degrade cellulose. In this work our goal was to isolate and identify the fungal species responsible for the biodegradation of a nineteenth-century art collection and determine their cellulolytic activity. Fungi were isolated using potato-dextrose-agar (PDA) and water-agar with carboxymethyl cellulose (CMC). The identification of the fungi was assessed through DNA sequencing (nrDNA ITS and α -actin regions) complemented with morphological analyses. Assays for cellulolytic activity were conducted with Gram's iodine as dye. Nineteen isolates were obtained, of which seventeen were identified through DNA sequencing to species level, belonging mainly to genera *Arthrinium*, *Aspergillus*, *Chaetomium*, *Cladosporium*, *Colletotrichum*, *Penicillium* and *Trichoderma*. For two samples that could not be identified through their ITS and α -actin sequences, a morphological analysis was conducted; they were identified as new species, named *Periconia epilithographicola* sp. nov. and *Coniochaeta cipronana* sp. nov. Qualitative tests showed that the fungal collection presents important cellulolytic activity.

Variations in the composition and appearance of a material as a consequence of the action of microorganisms is known as biodeterioration¹. This phenomenon becomes evident with the presence of reddish-brown or yellowish-brown patches, microfungus structures and textural changes, which are commonly found in ancient documents². These conditions apply to a nineteenth-century French collection of drawings and lithographs by Bernard Romain Julien (1802–1871) that is held in the archive of the School of Plastic Arts of Universidad de Costa Rica. The damage due to the microbial proliferation in these works of art is related to the storage conditions, especially to the damp and warm environments³. To design an effective and specific treatment according to the species growing in the laminae led to the isolation and identification of the fungal species responsible for the foxing of the lithographs.

Previous investigations of the microbiota in antique documents reported the presence of fungi that belong mainly to genera *Alternaria*, *Aspergillus*, *Chaetomium*, *Cladosporium*, *Penicillium*, and *Trichoderma*^{1,2,4–7}. For instance, El Bergadi *et al.* (2013) isolated, identified and characterized the microbiota of manuscripts from an ancient collection of the Medina of Fez and found *Aspergillus niger*, *Aspergillus oryzae*, *Mucor racemosus*, and *Penicillium chrysogenum*, as the most frequent species from a total of 31 fungal isolates⁸.

Because of nutritional limitations in which these fungi grew and where cellulose of the laminae was the only source of carbon, the species responsible for the biodeterioration were believed to possess cellulolytic activity^{8,9}. This cellulolytic activity is of interest for multiple biotechnological processes, such as treatment of agroindustrial residues^{10,11} or production of cellulases¹². This condition was first deduced and published in 1903 by van Ieterson in “La décomposition de la cellulose par les microorganismes”⁷. An investigation of the microbial diversity in a nineteenth-century Islamic and Koranic book led to the discovery of nine fungal species with the ability to

¹Centro Nacional de Innovaciones Biotecnológicas (CENIBiot), CeNAT-CONARE, 1174-1200, San Jose, Costa Rica.

²Centro de Investigaciones en Productos Naturales (CIPRONA), Universidad de Costa Rica, 11501-2060, San Jose, Costa Rica. ³Escuela de Biología, Universidad de Costa Rica, 11501-2060, San Jose, Costa Rica. ⁴Escuela de Química, Universidad de Costa Rica, 11501-2060, San Jose, Costa Rica. ⁵Instituto de Investigaciones en Arte (II Arte), 11501-2060, San Jose, Costa Rica. ⁶Department of Plant Science and Landscape Architecture, University of Maryland, College Park, MD, 20742, Maryland, USA. Correspondence and requests for materials should be addressed to M.C. (email: max.chavarría@ucr.ac.cr)

degrade carboxymethylcellulose (CMC), including *Aspergillus niger*, *A. oryzae* and *Hypocrea lixii*⁸. Michaelsen *et al.* (2009) and Pinzari *et al.* (2006) described *Aspergillus versicolor*, *A. nidulans*, *A. terreus* and *Chaetomium globosum* as agents in the microbiological damage of old documents^{5,6}.

Given the interest in the developing methods for protecting and preserving ancient documents from microbial degraders¹³ and the importance of obtaining microorganisms or enzymes with the capacity to degrade ligno-cellulosic wastes¹⁴, the aim of the present work was to isolate and identify the fungal species responsible for the biodegradation of a nineteenth-century art collection and to determine their cellulolytic activity. We found 19 fungal isolates belonging mainly to genera *Arthrimum*, *Aspergillus*, *Chaetomium*, *Cladosporium*, *Colletotrichum*, *Penicillium* and *Trichoderma*. Two samples not identified through their DNA sequences were identified through morphological analysis as new fungal species, namely *Periconia epilithographicola* sp. nov. and *Coniochaeta cipronana* sp. nov. Qualitative tests showed that the fungus collection presents important cellulolytic activity.

Methods

Sampling and isolation of cellulolytic fungi. A total of 13 laminae from a nineteenth-century French collection of lithographs belonging to Universidad de Costa Rica with signs of biodeterioration were sampled in areas of critical damage (colored or discolored areas, microfungus structures or other observable textural changes in the paper) with sterile cotton swabs, which were subsequently submerged in Phosphate Buffered Saline (PBS, 100 µL). Samples (50 µL) were cultured onto potato dextrose agar (PDA; Difco Potato Dextrose agar, BD company, France), and onto water agar with carboxymethyl cellulose (CMC, 1%, Sigma-Aldrich) with kanamycin (km, 50 µg/mL, Sigma-Aldrich). Morphologically distinct colonies were isolated and purified onto plates with the same culture media^{2,15–17}.

Molecular identification. To identify the various fungal isolates, DNA extraction was performed using the method described by Lodhi *et al.* (1994) with modifications¹⁸. First, two disks (diameter 0.8 cm) from each fungal colony were introduced into Eppendorf tubes (1.5 mL). An extraction buffer (750 µL) was added, followed by the vortex of the sample and an incubation period (30 min at 67 °C). DNA was then precipitated with the addition of a CHCl₃:octanol mixture (24:1, 750 µL), separation of the supernatant, and addition of isopropanol (600 µL, Sigma-Aldrich) and ethanol (500 µL, 70% v/v, Sigma-Aldrich). The DNA was eventually resuspended in AE buffer (50 µL, Qiagen) containing RNase (1 µL, Fermentas). PCR reactions were performed to amplify the ITS (ITS 4 and ITS 5) and actin (Act-512F and Act-783R) regions using a reaction mix (PCR Master Mix, 10 µL, 2X, Thermo Scientific), water (7 µL), primers (0.5 µL, 10 µM) and DNA (2 µL, 50 ng/µL)^{19,20}. All PCR reactions were performed in a thermocycler (Applied Biosystems 9902, Norwalk, USA) according to conditions described by Carbone & Kohn (1999) and White *et al.* (1990) for actin and ITS primers, respectively^{19,20}.

The amplified products were purified with a clean-up kit (EXO-AP, Thermo Scientific, USA) and sequenced with a genetic analyzer (ABI 3130xl) and a reaction kit (Big Dye v.3 Terminator Cycle Sequencing Ready Reaction Kit, Applied Biosystems, USA), using ITS and actin primers (1 µM). Sequences were analyzed with software (MEGA 7), and were run through a Standard Nucleotide BLAST (Genbank, NCBI nucleotide database) to assess the similarity with reported sequences of fungal species. The BLAST searches were run excluding uncultured/environmental samples in the database. To corroborate the results, the BLAST search was repeated limiting the search to sequences only from type material. All sequences have been deposited in the GenBank database under the accession numbers that appear in Supplementary Table S1.

Morphological identification. Two species that did not have a close match to anything in Genbank, were examined in more detail to determine their morphological characteristics. Morphological analyses followed recommendations and techniques described by Ellis (1971) for hyphomycetous fungi and common methods in mycology^{21,22}. Fungal isolates were cultured in CMD (BBL Corn Meal Dextrose agar, BD Company, France) and PDA (Difco Potato Dextrose agar, BD company, France) for 7 to 10 days near 25 °C. An optical microscope (Olympus BX-40, Japan) was used with an attached camera (18 megapixels, OMAX, Korea); software (ToupView, ToupTek Photonics, China) was used to measure structures.

Screening of cellulolytic activity. Cellulase-producing microorganisms were screened on agar plates enriched with only CMC as a source of carbon, with Gram's iodine as indicator (Prelab)^{23–26}. This qualitative determination is based on the interaction of iodine with cellulose and its components in its degraded form, such that the integral biopolymer holds Gram's iodine dye; whereas areas with cellulose hydrolyzed by enzymes result in clear zones or the appearance of a pale halo^{15,27}. The halo was measured for the subsequent calculation of the enzymatic index (EI), a semi-quantitative estimate of the enzyme activities, according to this formula¹⁵.

$$EI = \frac{\text{Diameter of hydrolysis zone}}{\text{Diameter of colony}} \quad (1)$$

For this purpose, fungal discs (diameter 0.8 cm) were grown in a solid medium composed of water agar (1.6%), CMC (1%) and kanamycin (km, 50 µg/mL). After cultures were incubated (7 days, 30 °C), plates were flooded with Gram's iodine stain (10 mL, 10 min) and washed with water to enable the observation, photographing and measurement of the clear zone around the fungal growth^{23–26}. Software (ImageJ, version 1.51j8) was used to measure the diameters²⁸. The experiment was repeated twice (on separate days) with duplicates of each isolate. *Pleurotus ostreatus* served as a positive control²⁹.




Figure 1. Fungal diversity in ancient lithographs. Several fungi were isolated from stained and degraded areas from nineteenth-century drawing laminae. On the top row from left to right are isolates #5, #10, #19, #9, #8, and #15. On the bottom row from left to right are isolates #16, #22, #21, #11, #23 and #26. Samples in the image were grown in PDA during 6 days. We thank Dr. Salomón Chaves (Instituto de Investigaciones en Arte) for authorizing the use of images from the collection of drawings by Bernard Romain Julien in this manuscript.

Results and Discussion

Isolation and identification of fungi isolated from drawings and lithographs. Through the screening of the lithographs, the total count of fungi isolated was 19, of which eight grew directly in water agar with CMC-km and eleven were first isolated from PDA and then recultivated in water agar with CMC as the sole source of carbon. The proliferation of fungi in the latter culture medium is in accordance with the environment in which they were isolated (limited sources of carbon, with cellulose as sole nutrient). Laminae #5 was the most contaminated, with ten isolations; followed by laminae #7 and #10, with 3 isolations each (see Supplementary Figure S1). The fungal isolates showed diverse forms, sizes, elevations, borders, surfaces, opacity, color and growth rates, as shown in Fig. 1.

BLAST searches in GenBank database resulted in the classification of nineteen isolates into fifteen species and nine genera (Table 1). These nineteen isolates had at least a 98% similarity with known species. The most prevalent genus was *Cladosporium*. Of the nine identified genera, *Aspergillus*, *Chaetomium*, *Cladosporium*, *Penicillium*, and *Trichoderma* are reported as common microbiota in ancient works of art^{1,2,17,30,31}. The actin region was sequenced to confirm the results obtained with the ITS region, and to classify to species level some samples that could not be done with ITS. For all cases in which both ITS and actin sequences were obtained, the fungi were classified within the same species, except isolate #9 in which the actin region denied conclusive results obtained with ITS.

Two isolates were only identified to genus or class levels using both ITS and actin regions. Specifically, isolate #19 was classified within the genus *Periconia*, and isolate #21 was classified within the class Sordariomycetes, both in the phylum Ascomycota. Since these two isolates did not have a close match to any sequence in the Genbank, traditional morphological analyses and descriptions (e.g. microscopy and use of taxonomic literature) were done to elucidate the identity of these isolates.

Description of two new fungal species. *Periconia epilithographicola* C. Coronado-Ruiz, R. Avendaño, E. Escudero-Leyva, G. Conejo-Barboza, P. Chaverri & M. Chavarría *sp. nov.* Fig. 2. **Mycobank:** MB825093 **GenBank:** MF422162 (ITS) & MF422179 (actin). **Etymology:** *epilithographicola*, because it was found growing over art lithographs. **Holotype:** Costa Rica, San José, San Pedro de Montes de Oca, Universidad de Costa Rica; on art lithographs; May 19th, 2016; collected by Avendaño R.; extype culture CBS 144017, a permanently preserved, metabolically inactive culture (= #19). **Diagnosis:** *Periconia* species producing a pinkish to reddish pigment. Straight conidiophores; globose, echinulated, golden-brown conidia. **Colonies:** At 25 °C after three weeks, on CMD, attaining 25 mm diam., colony white, cottony. On PDA, attaining 60 mm diam., colony effuse, pinkish (similar to OAC486), with creeping hyphae; conidiophores visible, forming small agglutinated black sticky drop-like structures. **Conidiophores:** macronematous, with creeping hyphae forming stipes 251.6–270 · 3.6–6.1 μm,

Isolate#	Identification	ITS and closest accession number			Actin and closest accession number		
		Accession	Identity	Coverage	Accession	Identity	Coverage
4	<i>Cladosporium sphaerospermum</i>	KP701988.1	100%	100%	EU570272.1	98%	99%
5	<i>Penicillium chrysogenum</i> ^a	KC009774.1	100%	100%	AM920435.1	97%	100%
6	<i>Penicillium westlingii</i> ^b	JN617668.1	100%	100%	AM920435.1	83%	55%
7	<i>Cladosporium tenuissimum</i> ^c	KP701937.1* KJ596320.1*	100%	100%	LN834582.1	100%	99%
8	<i>Aspergillus niger</i>	KJ365316.1	100%	100%	AM270331.1	99%	99%
9	<i>Cladosporium</i> sp.	KP701937.1* KJ596320.1*	100%	100%	—	—	—
10	<i>Arthrimum arundinis</i> ^b	KF144889.1	100%	100%	AY951865.1	76%	76%
11	<i>Cladosporium angustisporum</i> ^c	MG250413.1* MG199960.1* KP701978.1* KP701964.1* KP701938.1* KP701935.1* KP701930.1* KP701908.1*	100%	100%	LN834540.1	100%	100%
12	<i>Aspergillus versicolor</i>	NR_131277.1	100%	95%	—	—	—
13	<i>Chaetomium</i> cf. <i>subglobosum</i> ^b	NR_144826.1	96%	99%	KF545191.1	99%	100%
15	<i>Cladosporium angustisporum</i> ^c	MG250413.1* MG199960.1* KP701978.1* KP701964.1* KP701938.1* KP701935.1* KP701930.1* KP701908.1*	100%	100%	LN834540.1	100%	100%
16	<i>Cladosporium cladosporioides</i> ^c	MG250413.1* MG199960.1* KP701978.1* KP701964.1* KP701938.1* KP701935.1* KP701930.1* KP701908.1*	100%	100%	KT600582.1	99%	96%
17	<i>Chaetomium</i> cf. <i>subglobosum</i> ^b	NR_144826.1	100%	99%	KF545191.1	99%	100%
19	<i>Periconia</i> sp. ^d	HQ608027.1	99%	100%	KP184118.1	83%	95%
20	<i>Chaetomium</i> cf. <i>subglobosum</i> ^b	NR_144826.1	99%	99%	KF545191.1	99%	100%
21	<i>Coniochaeta</i> sp. ^d	KX869958.1	99%	100%	AY579255.1	71%	100%
22	<i>Aspergillus niger</i>	KJ365316.1	100%	100%	AM270331.1	99%	99%
23	<i>Trichoderma</i> cf. <i>Longibrachiatum</i> ^b	KT336509.1	100%	100%	JQ238613.1	98%	99%
26	<i>Colletotrichum kahawae</i> ^c	NR_144787.1	100%	98%	JX009431.1	99%	100%

Table 1. Molecular identification of isolated fungi using ITS and actin regions. ^aIsolates had homology with two fungi of different species with the ITS region analysis, but through sequencing of the actin region it was possible to confirm the identification. ^bNo register in the NCBI GenBank database for actin sequencing regions. ^cITS region sequencing allowed to identify only the isolates at genus level; the actin region enabled an identification at specie level. ^dNo register in the NCBI GenBank database for either ITS or actin sequencing regions. *Accessions with same similarity.

straight, branched singly near the base, seven or more septate, grayish to black. **Conidiogenous cells:** holoblastic, (5.1–) 7 · 10 (–11.5) μm (n = 15), sub-globose to ellipsoid, finely roughened, yellowish to brown (slightly more brilliant than OAC757). **Conidia:** globose, (7.8–) 9.2 (–10.7) μm diam. (n = 30), golden to brown (similar to OAC705), echinulated, catenated, sometimes forming long chains. **Habitat:** Growing on aged lithographs of Instituto de Investigaciones en Arte (Universidad de Costa Rica). **Notes:** Several *Periconia* species share similar characteristics of the conidiophore, differing mainly in the conidia size. *Periconia pseudobyssoides* conidia are larger, (12–) 15–17 (–20) μm diam. and brown-reddish³²; *P. byssoides* conidia are 10–15 μm in diam²¹; *P. saraswatiurensis* conidia are 9–12 μm diam., also secreting dark green to purple pigments in culture²¹. The only species with a similar conidial size is *P. jabalpurensis* but it lacks septa in the conidiophores; *P. macrospinoso* shows conidia of up to 35 μm diam. with long spines (<2 μm)²¹, which does not fit *Periconia epilithographicola*.

Coniochaeta cipronana C. Coronado-Ruiz, R. Avendaño, E. Escudero-Leyva, G. Conejo-Barboza, P. Chaverri & M. Chavarría sp. nov. Fig. 3. **Mycobank:** MB825094 **GenBank:** MF422164 (ITS) & MF422181 (actin). **Etymology:** as a reference to Centro de Investigaciones en Productos Naturales (CIPRONA, Universidad de Costa Rica) for the impact and transcendence of the research in the field of natural products over 38 years. **Holotype:** Costa Rica, San José, San Pedro de Montes de Oca, Universidad de Costa Rica; from art lithograph; May 19th, 2016; collected by Avendaño R.; extype culture CBS 144016, a permanently preserved, metabolically inactive culture (= #21).




Figure 2. *Periconia epigraphicola*. (A) PDA Culture ca 6 days. (B), PDA Culture ca 20 days. (C) Conidiogenous cell forming conidia. (D) Conidiogenous cell. (E) Catenated spinulose conidia. (F) Spinulose conidia.

Diagnosis: *Nodulisporium*-like conidiophore, with macro- and microconidia, hyaline, macroconidia 5–7-septate slightly curved, fusiform, microconidia cylindrical 1–2-septate. **Colonies:** At 25 °C after 3 weeks on CMD, reaching 20 mm diam., hyaline to white. On PDA attaining 25 mm diam., colony white, then turning purple (similar to OAC555), cracking and turning the media dull orange (lighter than OAC789). **Conidiophores:** *Nodulisporium*-like. **Conidiogenous cells:** Simple, mainly straight or sometimes curled, cylindrical, (5.8–) 21.8 (–28.8) · (2.2–) 2.8 (–3) µm (n = 15), arising directly from hyphae and stretching toward the apex, sometimes dichotomously branched, dimorphic, without collarete, hyaline. Short conidiophores (3–) 4.8 (–5) · 2 µm (n = 15). **Conidia:** macroconidia fusiform, (40.1–) 60.6 (–74.6) · (3–) 3.5 (–5) µm (n = 30), 5–7-septate, slightly curved, hyaline, smooth; microconidia cylindrical (11.6–) 16.5 (–25) · (2.2–) 2.8 (–3.6) µm (n = 30), 1–2-septate, hyaline, smooth. **Habitat:** Growing on aged lithographs of Instituto de Investigaciones en Arte (Universidad de Costa Rica). **Notes:** This species, because of the *Nodulisporium*-like conidiophore, is similar to *Coniochaeta ershadii*, especially in the size of the conidiogenous cells. The conidia produced by *C. ershadii* are prominently smaller³³ than those present in *C. cipronana*; the presence of macro- and microconidia is also a distinguishing character.

The new fungal species described belong to *Periconia* Tode and *Coniochaeta* (Sacc.) Cooke genera (see Supplementary Figures S2 and S3). *Periconia* is a polyphyletic genus Pleosporales (Dothideomycetes, Ascomycota), with a complicated taxonomy and a poorly understood phylogeny³². This genus has been widely reported as a common endophyte from the roots of several plants, like a *Periconia* species isolated from *Piper longum* producing metabolites with a high pharmacological potential³⁴ and the melanized hyphae are believed to protect the fungi from environmental oxidation³⁵. Some species have been reported as parasites in leaves of *Xanthium strumarium* and *Ipomea muricaria* in India and others as decomposers in bamboo statches³⁶. *Coniochaeta* (Coniochaetaceae, Coniochaetales, Sordariomycetes, Ascomycota) was introduced as a subgenus of *Rosellinia* De Not. for species with hairy perithecia but differing by the absence of amyloid asci in their sexual stages³³. Many *Coniochaeta* conidiophores produce *Lecytophora*-like structures. Like *Periconia*, *Coniochaeta* requires further taxonomic and phylogenetic studies³⁷. About 70 species and six synonyms are included in the genus *Coniochaeta* and most of the isolates are reported from dung, necrotic wood, soil and plant surfaces³⁸.




Figure 3. *Coniochaeta cipronana*. (A) PDA Culture ca 6 days. (B) PDA Culture ca 20 days. (C–E) Conidiogenous cell with small conidia. (F) Dicotomic conidiophore. (G) Macroconidia 5–7 septate.

Cellulase activity of the fungal isolates. Assay of the cellulase activity showed that 95% of the samples produce extracellular enzymes that break down cellulose into smaller oligosaccharides or monosaccharides, as evident from the clear zone observed after staining the plates with Gram's iodine (see Table 2 and Supplementary Figure S4). This fraction that includes the two new species (sample #19: *Periconia epilithographicola* and sample #21: *Coniochaeta cipronana*) also comprehends species of *Arthrinium*, *Aspergillus*, *Chaetomium*, *Cladosporium*, *Colletotrichum*, *Penicillium*, and *Trichoderma*, being the first four commonly reported with cellulolytic activity^{31,39–45}. These observations are congruent with the habitat of restricted carbon sources, in which sheets or laminae made of fibers of cellulose pulp were the support material for the growth of microorganisms.

Importantly, 32% of the total isolates had a significantly superior enzymatic index relative to a positive control (*P. ostreatus*), i.e., isolates #5 (*Penicillium chrysogenum*), #7 (*Cladosporium tenuissimum*), #11 (*Cladosporium angustisporum*) and #23 (*Trichoderma* cf. *longibrachiatum*). Other studies have characterized these species as effective cellulase producers^{16,40}. Isolates #5 (*Penicillium chrysogenum*) and #23 (*Trichoderma* cf. *longibrachiatum*) had an outstanding performance relative to the positive control and the rest of the isolated fungi. Specifically, isolate #5 presented an enzymatic index for cellulase activity almost twice of that of the positive control. Isolates of these species not only have presented important cellulase activity but also have been the object of study for their capacity to produce xylanases⁴⁶, or tanases⁴⁷.

The case of isolate # 23 (*Trichoderma* cf. *longibrachiatum*) was even more striking. For this fungus, EI is reported for 24 h (see Table 2) because after 7 days (the period in which the other isolates were measured) the microorganism had covered the entire Petri plate, evidence of an accelerated growth and a large capacity to use the CMC as the sole source of carbon. The result (1.39 ± 0.03) was slightly smaller than the positive control (1.8 ± 0.1 , measured after seven days). However, as previously mentioned, isolate # 23 was measured at 24 h. This result implies a large rate of enzymatic (cellulase) production from fungus #23 in a medium rich in cellulose, relative to the rest of the fungi studied, which is important for the development of biotechnological applications

Isolate#	Enzymatic index
4	No activity
5	3,3 ± 0,2
6	1,91 ± 0,07
7	2,87 ± 0,05
8	0,92 ± 0,02
9	1,6 ± 0,1
10	1,62 ± 0,08
11	2,74 ± 0,02
12	1,47 ± 0,09
13	1,243 ± 0,005
15	2,87 ± 0,03
16	1,47 ± 0,03
17	1,199 ± 0,004
19	1,861 ± 0,002
20	1,17 ± 0,08
21	1,57 ± 0,06
22	1,086 ± 0,006
23	1,39 ± 0,03*
26	0,80 ± 0,06
Control: <i>P. ostreatus</i>	1,8 ± 0,1

Table 2. Enzymatic indices of the isolates on CMC agar stained with Gram Iodine after incubation for seven days. *EI was measured after 24 h.

and industry. Many studies have featured this species as a fungus with great cellulase activity^{48–51}. Many commercial cellulases can be purchased in purified form after production with this species (e.g. C9748 Sigma-Aldrich or E-CELTR from Megazyme). Investigations with isolation # 23 will continue to evaluate its potential to degrade lignocellulosic residues from agricultural activity in Costa Rica (e.g., wastes from pineapple production).

In summary, in isolating, identifying and characterizing the cellulolytic activity of the fungi responsible for the biodegradation of a nineteenth-century collection, several species of fungi were found to have the ability to produce cellulases. In addition, two new species of fungi were identified and named *Periconia epilithographicola* sp. nov. and *Coniochaeta cipronana* sp. nov., which also have cellulolytic activity. A knowledge of the microorganisms that colonized the Bernard Romain Julien collection belonging to Universidad de Costa Rica will allow the development of strategies directed to the conservation of these ancient lithographs. This work also contributes to the knowledge of new species with cellulolytic activity, which is a topic of perennial interest for biotechnology because of the important role of fungal cellulolytic enzymes in commercial food processing, performing the hydrolysis of cellulose during drying of beans, in the textile industry and laundry detergents, in the conversion of biomass into industrially important solvents or fuels, and their potential application for the bioremediation of wastes.

References

- Sterflinger, K. & Piñar, G. Microbial deterioration of cultural heritage and works of art — tilting at windmills. *Appl. Microbiol. Biotechnol.* **97**, 9637–9646 (2013).
- Mesquita, N. *et al.* Fungal diversity in ancient documents. a case study on the archive of the University of Coimbra. *Int. Biodeter. Biodegr.* **63**, 626–629 (2009).
- Rakotonirainy, M. S., Heude, E. & Lave, B. Isolation and attempts of biomolecular characterization of fungal strains associated to foxing on a 19th century book. *J. Cult. Herit.* **8**, 126–133 (2007).
- Garg, K. L., Kamal, K. & Mishra, A. K. Role of fungi in the deterioration of wall paintings. *Sci. Total Environ.* **167**, 255–271 (1995).
- Michaelsen, A., Piñar, G., Montanari, M. & Pinzari, F. Biodeterioration and restoration of a 16th-century book using a combination of conventional and molecular techniques: A case study. *Int. Biodeter. Biodegr.* **63**, 161–168 (2009).
- Pinzari, F., Pasquariello, G. & De Mico, A. Biodeterioration of paper: a SEM study of fungal spoilage reproduced under controlled conditions. *Macromol. Symp.* **238**, 57–66 (2006).
- Sterflinger, K. & Pinzari, F. The revenge of time: fungal deterioration of cultural heritage with particular reference to books, paper and parchment. *Environ. Microbiol.* **14**, 559–566 (2012).
- El Bergadi, F., Laachari, F., Elabed, S., Mohammed, I. & Ibsouda, S. Cellulolytic potential and filter paper activity of fungi isolated from ancients' manuscripts from the Medina of Fez. *Ann. Microbiol.* **64**, 815–822 (2013).
- Montegut, D., Indictor, N. & Koestler, R. J. Fungal deterioration of cellulosic textiles: a review. *Int. Biodeterior.* **28**, 209–226 (1991).
- Anwar, Z., Gulfaz, M. & Irshad, M. Agro-industrial lignocellulosic biomass a key to unlock the future bio-energy: A brief review. *J. Radiat. Res. Appl. Sci.* **7**, 163–173 (2014).
- Berlemont, R. Distribution and diversity of enzymes for polysaccharide degradation in fungi. *Sci. Rep.* **7**, 222 (2017).
- Chiranjeevi, T. *et al.* Optimization of holocellulolytic enzymes production by *Cladosporium Cladosporioides* using Taguchi-L16 orthogonal array. *J. Biobased Mater. Bioenergy.* **6**, 1–10 (2012).
- Sequeira, S., Phillips, A., Cabrita, E. & Macedo, M. Antifungal treatment of paper with calcium propionate and parabens: Short-term and long-term effects. *Int. Biodeterior. Biodegradation.* **120**, 203–215 (2017).
- Montella, S. *et al.* Discovery of genes coding for carbohydrate-active enzyme by metagenomic analysis of lignocellulosic biomasses. *Sci. Rep.* **7**, 42623 (2017).
- Florencio, C., Couri, S. & Farinas, C. Correlation between agar plate screening and solid-state fermentation for the prediction of cellulase production by *Trichoderma* strains. *Enzyme Res.* **2012**, 793708–793715 (2012).

16. Makeshkumar, V. & Mahalingam, P. U. Isolation and characterization of rapid cellulose degrading fungal pathogens from compost of agro wastes. *Int. J. Pharm. Biol. Arch.* **2**, 1695–1698 (2011).
17. Sanmartín, P., DeAraujo, A. & Vasanthakumar, A. Melding the old with the new: trends in methods used to identify, monitor, and control microorganisms on cultural heritage materials. *Microb. Ecol.* **74**, 1–17 (2016).
18. Lodhi, A., Ye, G., Weeden, N. & Reisch, B. A simple and efficient method for DNA extraction from Grapevine Cultivars and *Vitis* species. *Plant Mol. Biol. Rep.* **12**, 6–13 (1994).
19. White, T., Bruns, S., Lee, S. & Taylor, J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*. 315–322 (1990).
20. Carbone, I. & Kohn, L. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycology*. **91**, 553–556 (1999).
21. Ellis, M. *Dematiaceous hyphomycetes* (CABI Publishing, 1971).
22. Sutton, B.C. The Coelomycetes. Fungi Imperfecti with Pycnidia, Acervuli and Stromata (CABI Publishing, 1980).
23. Bakar, N., Abd-Aziz, S., Hassan, M. & Ghazali, F. Isolation and selection of appropriate cellulolytic mixed microbial cultures for cellulases production from oil palm empty fruit bunch. *Biotech.* **9**, 73–78 (2010).
24. Gohel, H. R., Contractor, C. N., Ghosh, S. K. & Braganza, V. J. A comparative study of various staining techniques for determination of extra cellular cellulase activity on Carboxy MethylCellulose (CMC) agar plates. *Int. J. Curr. Microbiol. App. Sci.* **3**, 261–266 (2014).
25. Johnsen, H. R. & Krause, K. Cellulase activity screening using pure carboxymethylcellulose: Application to soluble cellulolytic samples and to plant tissue prints. *I. J. Mol. Sci.* **15**, 830–838 (2014).
26. Meddeb-Mouelhi, F., Moisan, J. K. & Beaugerard, M. A comparison of plate assay methods for detecting extracellular cellulase and xylanase activity. *Enzyme Microb. Technol.* **66**, 16–19 (2014).
27. Wang, Y. & Easteal, A. Interaction between iodine and ethyl cellulose. *J. Appl. Polym. Sci.* **71**, 1303–1314 (1999).
28. Schneider, C., Rasband, W. & Eliceiri, K. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*. **9**, 671–675 (2012).
29. Valášková, V. & Baldrian, P. Estimation of bound and free fractions of lignocellulose-degrading enzymes of wood-rotting fungi *Pleurotus ostreatus*, *Trametes versicolor* and *Piptoporus betulinus*. *Research in Microbiology*. **157**, 119–124 (2006).
30. Dunca, S. *et al.* Study of the contaminating microbiota of old paper supports. *Eur. Sci. J.* **3**, 237–251 (2014).
31. Naji, K., Abdullah, Q., Al-Zaqri, A. & Alghalibi, S. Evaluating the biodeterioration enzymatic activities of fungal contamination isolated from some ancient Yemeni mummies preserved in the national museum. *Biochem. Res. Int.* **2014**, 1–9 (2014).
32. Markovskaja, S. & Kacergius, A. Morphological and Molecular Characterization of *Periconia pseudobyssoides* sp. nov. and a closely related *P. byssoides*. *Mycol. Progress.* **13**, 291–302 (2014).
33. Asgari, B., Zare, R. & Gams, W. *Coniochaeta ershadii*, a new species from Iran and a key to well-documented *Coniochaeta* species. *Nova Hedwigia*. **84**, 175–187 (2007).
34. Verma, V. C., Lobkovsky, E., Gange, A. C., Singh, S. K. & Prakash, S. Piperine production by endophytic fungus *Periconia* sp. isolated from *Piper longum* L. *J. Antibiot.* **64**, 427–431 (2011).
35. Yuan, Z., Zhang, C., Lin & Kubicek, C. Identity, diversity, and molecular phylogeny of the endophytic mycobiota in the roots of rare wild rice (*Oryza granulata*) from a nature reserve in Yunnan, China. *Appl. Environ. Microbiol.* **76**, 1642–1652 (2010).
36. Rao, R. & Rao, D. The genus *Periconia* from India. *Mycopathol. Mycol. Appl.* **22**, 285–310 (1964).
37. Khan, Z. *et al.* *Coniochaeta polymorpha*, a new species from endotracheal aspirate of a preterm neonate, and transfer of *Lecytophora* species to *Coniochaeta*. *Antonie van Leeuwenhoek*. **104**, 243–252 (2013).
38. Xie, J. *et al.* An endophytic *Coniochaeta velutina* producing broad spectrum antimycotics. *J. Microbiol.* **53**, 390–397 (2015).
39. Li, Y. *et al.* The different roles of *Penicillium oxalicum* LaeA in the production of extracellular cellulase and β -xylosidase. *Front. Microbiol.* **7**, 1–14 (2016).
40. Abe, C. *et al.* Fungi isolated from maize (*Zea mays* L.) grains and production of associated enzyme activities. *Int. J. Mol. Sci.* **16**, 15328–15346 (2015).
41. Ciferri, O. Microbial degradation of paintings. *Appl. Environ. Microbiol.* **65**, 879–885 (1999).
42. El-Said, A., Saleem, A., Maghraby, T. & Hussein, M. Cellulase activity of some phytopathogenic fungi isolated from diseased leaves of broad bean. *Int. J. Curr. Microbiol. App. Sci.* **3**, 883–900 (2014).
43. Ghose, T. K. Measurement of cellulase activities. *Pure Appl. Chem.* **59**, 257–268 (1987).
44. Gopinath, S., Anbu, P. & Hilda, A. Extracellular enzymatic activity profiles in fungi isolated from oil-rich environments. *Mycoscience*. **46**, 119–126 (2005).
45. Wang, M. & Lu, X. Exploring the synergy between cellobiose dehydrogenase from *Phanerochaete chrysosporium* and cellulase from *Trichoderma reesei*. *Front. Microbiol.* **7**, 1–10 (2016).
46. Haas, H., Herfurth, E., Stoffler, G. & Redl, B. Purification, characterization and partial amino acid sequences of a xylanase produced by *Penicillium chrysogenum*. *Biochim. Biophys. Acta*. **1117**, 279–286 (1992).
47. Rajakumar, G. & Nandy, S. Isolation, purification, and some properties of *Penicillium chrysogenum* Tannase. *Appl. Environ. Microbiol.* **46**, 525–527 (1983).
48. Sandhu, D. & Kalra, M. Production of cellulase, xylanase and pectinase by *Trichoderma longibrachiatum* on different substrates. *T. Brit. Mycol. Soc.* **79**, 409–413 (1982).
49. Roger, J. & Nacas, J. Interrelationship of Xylanase Induction and Cellulase Induction of *Trichoderma longibrachiatum*. *Appl. Environ. Microbiol.* **56**, 2535–2539 (1990).
50. Kalra, M., Sidhu, M. & Sandhu, D. Partial purification, characterization and regulation of cellulolytic enzymes from *Trichoderma longibrachiatum*. *J. Appl. Bacteriol.* **61**, 73–80 (1986).
51. Chutani, P. & Sharma, K. Concomitant production of xylanases and cellulases from *Trichoderma longibrachiatum* MDU-6 selected for the deinking of paper waste. *Bioprocess Biosyst. Eng.* **39**, 747–758 (2016).

Acknowledgements

We are grateful to Instituto de Investigaciones en Arte (II Arte) of Universidad de Costa Rica and Dr. Salomon Chaves Badilla for the access and support for the study and selection of the collection of drawings. This work was supported by resources of the Vice-rectory of Research of Universidad de Costa Rica (project number 726-B6-115), Centro Nacional de Innovaciones Biotecnológicas (CENIBiot) and Reintegra (Red en Ciencia y Conservación del Patrimonio Cultural Nacional).

Author Contributions

M.C. conceived and designed the experiments; C.C.-R., R.A., G.C.-B., E.E.-L. performed the experiments; C.C.-R., R.A., E.E.-L., P.C., M.C. analyzed the data; P.C., M.C. contributed reagents or materials or analytical tools; C.C.-R., E.E.-L., P.C., M.C. wrote the paper. All authors reviewed and approved the final version of the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-24934-7>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018