



NOTE

Two new xanthenes and cytotoxicity from the bark of *Garcinia schomburgkiana*

Sutin Kaennakam¹ · Kamonwan Mudsing¹ · Kitiya Rassamee² · Pongpun Siripong² · Santi Tip-pyang¹

Received: 25 May 2018 / Accepted: 17 August 2018 / Published online: 31 August 2018
© The Author(s) 2018

Abstract

Two new xanthone derivatives, named schomburgones A (**1**) and B (**2**), along with eight known compounds, including xanthenes (**3–8**) and anthraquinones (**9–10**) were isolated from the bark of *Garcinia schomburgkiana*. Their structures were determined by spectroscopic analysis especially 1D and 2D NMR spectroscopies. All isolated compounds were evaluated for their cytotoxicity against five cancer cell lines (KB, HeLa S-3, HT-29, MCF-7 and HepG-2). Compounds **3–6** and **8** showed good cytotoxicity against all the five cancer cell lines with IC₅₀ values in the range of 1.45–9.46 μM.

Keywords *Garcinia schomburgkiana* · Clusiaceae · Xanthone · Cytotoxicity

Introduction

Garcinia schomburgkiana Pierre (family Clusiaceae) is a medium-sized tree distributed in Thailand, Laos, Vietnam, and Cambodia. In folk medicine in these countries, its leaves, roots, and fruits are used for the treatment of cough, menstrual disturbances, expectorant, laxative and diabetes [1]. Previous chemical and biological studies on the chemical constituents of *G. schomburgkiana* showed the presence of xanthenes, depsidones, biphenyls, flavonoids, triterpenoids, and phloroglucinols, some of which exhibited anti-malarial activity and cytotoxicity [2], [3]. Here, we reported two new xanthone derivatives, named schomburgones A (**1**) and B (**2**), along with six known xanthenes (**3–8**) and two known anthraquinones (**9–10**) from the bark of this plant. The structures of all isolated compounds were elucidated

using spectroscopic methods especially 1D and 2D NMR spectroscopies and compared with their ¹H and ¹³C NMR spectroscopic data from the literature. The cytotoxicity of all isolated compounds was evaluated using the MTT method against five cancer cell lines.

Results and discussion

Phytochemical investigation of CH₂Cl₂ crude extract from the bark of *G. schomburgkiana* led to the isolation of two new xanthone derivatives, named schomburgones A (**1**) and B (**2**), along with eight known compounds (Fig. 1), including isocudranixanthone B (**3**) [4], gerontoxanthone I (**4**) [5], nigrolineaxanthone E (**5**) [6], isojacareubin (**6**) [7], dulxanthone A (**7**) [8], macluraxanthone (**8**) [9], vismiaquinone A (**9**) [10], and 3-geranylemodin (**10**) [11]. The structures of all isolated compounds were elucidated using spectroscopic methods especially NMR spectroscopies and compared with their ¹H and ¹³C NMR spectroscopic data from the literature.

Schomburgone A (**1**) was obtained as yellow oil. Its molecular formula was determined as C₂₄H₂₄O₆ by the negative HRESIMS measurement through the ion peak at *m/z* 407.1527 [M–H]⁺ (calcd. for C₂₄H₂₃O₆, 407.1495). The UV spectrum displayed absorption bands at λ_{max} 395, 315 and 243 nm, which is typical of the xanthone chromophore [12]. The IR spectrum showed phenolic hydroxyl groups and a hydrogen bonded carbonyl group at 3422 and 1632 cm⁻¹. The ¹H NMR spectrum showed the presence

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11418-018-1240-8>) contains supplementary material, which is available to authorized users.

✉ Sutin Kaennakam
n-s-k@hotmail.com

✉ Santi Tip-pyang
santi.Ti@chula.ac.th

¹ Center of Excellence in Natural Products Chemistry, Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

² Natural Products Research Section, Research Division, National Cancer Institute, Bangkok 10400, Thailand




Fig. 1 Chemical structures of **1–10**


of a 3,3-dimethylallyl substituent, which was confirmed by two singlets at δ_{H} 1.67 (3H, s, H-4') and 1.85 (3H, s, H-5') for the vinyl methyls, a triplet at δ_{H} 5.20 (1H, t, $J=7.23$ Hz, H-2') for the vinylic proton and a doublet at δ_{H} 3.44 (2H, d, $J=7.23$ Hz, H-1') for the allylic proton of prenyl group. In addition, the methoxy signal, a hydroxyl signal, two aromatic proton signals and a hydrogen bonded hydroxyl signal appeared as five singlets at δ_{H} 3.89 (3H, s, OCH₃-3), 6.26 (1H, s, OH-6), 6.33 (1H, s, H-2), 6.85 (1H, s, H-5) and 13.38 (1H, s, OH-1), respectively. The signals at δ_{H} 1.50 (6H, s, H-4'' and H-5''), 5.83 (1H, d, $J=10.23$ Hz, H-2'') and 8.02 (1H, d, $J=10.23$ Hz, H-1'') in the spectrum were indicative of a dimethylchromene ring. The angular fusion of the chromene ring at C-8 was deduced from the low field shift of H-1'' (δ_{H} 8.02), which was located in the deshielding area of the carbonyl group. The ¹H and ¹³C NMR spectroscopic data

(Table 1) were shown to be similar to those of the known xanthone, paxanthone B [13], except that the hydroxyl group at C-3 of paxanthone B was replaced by a methoxy group. In the HMBC correlations of **1** (Fig. 2), the methoxy proton at δ_{H} 3.89 showed a cross-peak with δ_{C} 163.5 (C-3). In addition, a methine proton at δ_{H} 8.02 showed cross-peaks with δ_{C} 77.1 (C-3''), 108.5 (C-8a), and 136.9 (C-7), confirming that a dimethylchromene ring was located at C-8, and a methylene proton at δ_{H} 3.44 showed cross-peaks with δ_{C} 131.7 (C-3'), 153.7 (C-4a), and 163.5 (C-3), indicating that a prenyl group was attached to C-4. Thus, the complete assignment of schomburgone A was determined as **1**.

Schomburgone B (**2**) was obtained as yellow oil. Its molecular formula was determined as C₁₉H₁₈O₅ by the negative HRESIMS measurement through the ion peak at m/z 325.1098 [M-H]⁺ (calcd. for C₁₉H₁₇O₅, 325.1076). The

Table 1 NMR spectroscopic data (400 MHz, CDCl₃) for **1** and **2**

Position	1		2	
	δ_{H} (J in Hz)	δ_{C}	δ_{H} (J in Hz)	δ_{C}
1		162.0		163.0
2	6.33 (s)	94.1	6.43 (s)	96.2
3		163.5		166.1
4		107.2		113.9
4a		153.7		154.0
5	6.85 (s)	102.6		145.9
6		151.1	7.26 (d, 7.62)	120.3
7		136.9	7.23 (t, 7.62)	124.6
8		119.9	7.71 (d, 7.62)	116.6
8a		108.5		121.0
9		183.0		182.0
9a		103.9		104.1
10a		153.5		144.8
1'	3.44 (d, 7.23)	21.7		42.0
2'	5.20 (t, 7.23)	122.4	6.70 (dd, 10.63, 17.67)	156.3
3'		131.7	5.22 (d, 17.67), 5.07 (d, 10.63)	104.5
4'	1.67 (s)	25.9	1.61 (s)	28.4
5'	1.85 (s)	18.0	1.61 (s)	28.4
1''	8.02 (d, 10.23)	121.2		
2''	5.83 (d, 10.23)	132.5		
3''		77.1		
4''	1.50 (s)	27.5		
5''	1.50 (s)	27.5		
1-OH	13.38 (s)		13.25 (s)	
5-OH			6.42 (s)	
6-OH	6.26 (s)			
3-OCH ₃	3.89 (s)	56.1	3.91 (s)	56.2

**Fig. 2** Key HMBC (arrow curves) and COSY (bold lines) correlations of **1** and **2**

UV spectrum displayed absorption bands at λ_{max} 394, 315 and 244 nm. The IR spectrum showed phenolic hydroxyl groups and a hydrogen bonded carbonyl group at 3432 and 1642 cm^{-1} . The ¹H NMR spectrum showed the presence of a 1,1-dimethylallyl group, which was confirmed by a singlets at δ_{H} 1.61 (6H, s, H-4' and H-5') for two methyls, a doublet of doublet at δ_{H} 6.70 (1H, dd, $J = 10.63$ Hz, 17.67,

H-2') for the methine proton and two doublets at δ_{H} 5.07 (1H, d, $J = 10.63$ Hz, H-3') and 5.22 (1H, d, $J = 17.67$ Hz, H-3') for the methylene protons. Moreover, the methoxy signal, hydroxyl signal, aromatic proton signal and hydrogen bonded hydroxyl signal appeared as four singlets at δ_{H} 3.91 (3H, s, OCH₃-3), 6.42 (1H, s, OH-5), 6.43 (1H, s, H-2) and 13.25 (1H, s, OH-1), respectively. The ABC-type aromatic protons were assigned at δ_{H} 7.23 (1H, t, $J = 7.62$ Hz, H-7), 7.26 (1H, d, $J = 7.62$ Hz, H-6) and 7.71 (1H, d, $J = 7.62$ Hz, H-8). The ¹H and ¹³C NMR spectroscopic data (Table 1) were shown to be similar to those of the known xanthone, pancixanthone A [14], except that the hydroxyl group at C-3 of pancixanthone-A was substituted by a methoxy group. In the HMBC correlations of **2** (Fig. 2), the methoxy proton at δ_{H} 3.91 showed a cross-peak with δ_{C} 166.1 (C-3). Moreover, two methyl protons at δ_{H} 1.61 showed cross-peaks with δ_{C} 113.9 (C-4) and 156.3 (C-2'), confirming that a 1,1-dimethylallyl group was connected at C-4. Thus, the completed assignment of schomburgone B was determined as **2**.

In previous research many xanthenes showed cytotoxicity [15]. Therefore, all isolated compounds were evaluated in vitro for their cytotoxicity against five cancer cell lines (KB, HeLa S-3, HT-29, MCF-7 and HepG-2) (Table 2). Compounds **3–6** and **8** showed good cytotoxicity against all five cancer cell lines with IC₅₀ values in the range of 1.45–9.46 μM . Compounds **1** and **7** showed weak cytotoxicity against all five cancer cell lines with IC₅₀ values in the range of 34.69–73.10 μM . Compounds **2**, **9** and **10** showed inactive cytotoxicity against all five cancer cell lines with IC₅₀ values >100 μM . The SAR studied data (Fig. 1; Table 2) of xanthenes suggest that the ortho hydroxy group at C-5 and C-6 and the 1,1-dimethylallyl group at C-4 might improve the cytotoxicity as inferred from the comparison of their cytotoxicity of compounds **1–10**.

Experimental

General experimental procedures

NMR spectra were recorded on Bruker 400 AVANCE spectrometer. HRESIMS spectra were obtained using a Bruker MICROTOF model mass spectrometer. The UV–visible absorption spectra were recorded on a UV-2550 UV–Vis spectrometer (Shimadzu, Kyoto, Japan). The IR spectra were measured on a Nicolet 6700 FT-IR spectrometer using KBr discs.

Plant material

The bark of *G. schomburgkiana* was collected from Bang Ramat Road, Khwaeng Bang Ramat, Khet Taling Chan, Bangkok Thailand (13°45'42"N, 100°24'56"E), in June

Table 2 In vitro cytotoxicity of compounds **1–10** against five cancer cell lines

Compounds	IC ₅₀ (μM) ± SD				
	KB	HeLa S-3	HT-29	MCF-7	HepG-2
1	45.05 ± 2.08	69.22 ± 4.02	61.92 ± 2.40	52.21 ± 1.71	73.19 ± 1.14
2	> 100	> 100	> 100	> 100	> 100
3	5.23 ± 0.19	7.95 ± 0.25	7.87 ± 0.30	6.70 ± 0.81	5.93 ± 0.94
4	4.69 ± 0.21	7.57 ± 0.26	9.18 ± 0.38	5.26 ± 0.55	4.89 ± 0.83
5	5.08 ± 0.36	5.82 ± 0.15	4.17 ± 0.07	7.19 ± 0.36	9.46 ± 0.45
6	4.30 ± 0.12	6.60 ± 0.24	5.92 ± 0.40	3.21 ± 0.71	3.19 ± 0.14
7	38.17 ± 6.83	65.26 ± 3.89	34.69 ± 2.29	46.03 ± 1.29	54.80 ± 1.18
8	1.45 ± 0.09	1.62 ± 0.20	1.87 ± 0.30	1.70 ± 0.81	1.93 ± 0.94
9	> 100	> 100	> 100	> 100	> 100
10	> 100	> 100	> 100	> 100	> 100
Doxorubicin	0.13 ± 0.006	0.03 ± 0.001	0.31 ± 0.07	0.42 ± 0.14	1.23 ± 0.02

IC₅₀ ≤ 10 = good activity, 10 < IC₅₀ ≤ 30 = moderate activity, IC₅₀ > 100 = inactive

2017. The plant material was identified by Dr. Suttira Sedlak, a botanist at the Walai Rukhavej Botanical Research Institute, Mahasarakham University, and a specimen retained as a reference (Khumkratok no. 92-08).

Extraction and isolation

The air-dried bark of *G. schomburgkiana* (2.0 kg) was extracted with CH₂Cl₂ at room temperature for 7 days (2 × 25 L). The CH₂Cl₂ crude extract (91.0 g) was further separated by column chromatography (CC) over silica gel CC and eluted with a gradient of Hexane–EtOAc (90, 70, 50 and 30% Hexane–EtOAc each 5 L) to give six fractions (A–F). Fraction A (4.0 g) was purified by Sephadex LH-20 column eluted with 80% CH₂Cl₂–MeOH (2 L) and further applied to a radial chromatography (chromatotron) with 95% hexane–EtOAc (200 mL) to afford compound **9** (3.2 mg). Fraction B (10.5 g) was purified by Sephadex LH-20 column eluted with 80% CH₂Cl₂–MeOH (2 L) and further applied to a chromatotron with 50% hexane–CH₂Cl₂ (200 mL) to obtain compounds **2** (4.2 mg), **4** (2.5 mg) and **7** (2.3 mg). Compound **1** (7.2 mg) was separated by Sephadex LH-20 column eluted with 50% CH₂Cl₂–MeOH (2 L) from fraction C (2.0 g). Fraction D (6.5 g) was purified by Sephadex LH-20 column eluted with 50% CH₂Cl₂–MeOH (2 L) to give compounds **5** (8.5 mg) and **8** (4.6 mg). Fraction E (8.5 g) was purified by Sephadex LH-20 column eluted with 50% CH₂Cl₂–MeOH (2 L) and further applied to a chromatotron with 70% hexane–EtOAc (200 mL) to obtain compounds **3** (5.2 mg) and **10** (5.5 mg). Finally, fraction F (1.2 g) was subjected to silica gel CC eluted with 100% CH₂Cl₂ and further purified by Sephadex LH-20 column eluted with 80% CH₂Cl₂–MeOH (2 L) to yield compound **6** (6.5 mg).

Schomburgone A (**1**): yellow oil; UV (CHCl₃) λ_{max} (log ε): 395 (3.6), 315 (4.2) and 243 (4.4) nm., IR ν_{max} (KBr): 3422 and 1632 cm⁻¹; ¹H (400 MHz, CDCl₃) and ¹³C

NMR (100 MHz, CDCl₃) spectroscopic data, see Table 1; HRESIMS *m/z* 407.1527 [M–H]⁺ (calcd. for C₂₄H₂₃O₆, 407.1495).

Schomburgone B (**2**): yellow oil; UV (CHCl₃) λ_{max} (log ε): 394 (3.5), 315 (4.0) and 244 (4.2) nm., IR ν_{max} (KBr): 3432 and 1642 cm⁻¹; ¹H (400 MHz, CDCl₃) and ¹³C NMR (100 MHz, CDCl₃) spectroscopic data, see Table 1; HRESIMS *m/z* 325.1098 [M–H]⁺ (calcd. for C₁₉H₁₇O₅, 325.1076).

Cytotoxicity assay

All isolated compounds (**1–10**) were subjected to cytotoxic evaluation against KB (human epidermoid carcinoma), HeLa S-3 (human cervical carcinoma), HT-29 (human colon adenocarcinoma), MCF-7 (human breast adenocarcinoma) and HepG-2 (human liver carcinoma) cell lines employing the colorimetric method [16]. Doxorubicin was used as the reference substance which exhibits activity against five cancer cell lines. The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (Sigma Chemical Co., USA) was dissolved in saline to make a 5 mg/mL stock solution. Cancer cells (3 × 10³ cells) suspended in 100 μg/wells of MEM medium containing 10% fetal calf serum (Gibco BRL, Life Technologies, NY, USA) were seeded onto a 96-well culture plate (Costar, Corning Incorporated, NY, USA). After 24 h pre-incubation at 37 °C in a humidified atmosphere of 5% CO₂/95% air to allow cellular attachment, various concentrations of test solution (0.1, 0.3, 1.0, 3.0, 10.0, 30.0, and 100.0 μM, each 10 μL/well) were added and these were then incubated for 48 h under the above conditions. At the end of the incubation, 10 μL of tetrazolium reagent was added into each well followed by further incubation at 37 °C for 4 h. The supernatant was decanted, and DMSO (100 μL/well) was added to allow formosan solubilization. The optical density of each well was detected using a Microplate

reader at 550 nm and for correction at 595 nm. Each determination represented the average mean of six replicates. The 50% inhibition concentration (IC₅₀ value) was determined by curve fitting.

Acknowledgments The authors are grateful to the Graduate School of Chulalongkorn University for a Postdoctoral Fellowship (Ratchadaphiseksomphot Endowment Fund) to SK. We also thank Dr. Suttira Sedlak, Walai Rukhavej Botanical Research Institute, Mahasarakham University, Mahasarakham 44000, Thailand for identification and deposition of the plant material.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Mungmee C, Sitthigool S, Suttisri R, Buakeaw A (2012) Xanthenes and biphenyls from *Garcinia schomburgkiana* wood and their cytotoxicity. *Thai J Pharm Sci* 36:6–9
- Le DH, Nishimura K, Takenaka Y, Mizushima Y, Tanahashi T (2016) Polyprenylated Benzoylphloroglucinols with DNA polymerase inhibitory activity from the fruits of *Garcinia schomburgkiana*. *J Nat Prod* 79:1798–1807
- Sukandar ER, Siripong P, Khumkratok S, Tip-Pyang S (2016) New depsidones and xanthone from the roots of *Garcinia schomburgkiana*. *Fitoterapia* 111:73–77
- Kobayashi M, Mahmud T, Yoshioka N, Shibuya H, Kitagawa I (1997) Indonesian medicinal plants. XXI. Inhibitors of Na⁺/H⁺ exchanger from the bark of *Erythrina variegata* and the roots of *Maclura cochinchinensis*. *Chem Pharm Bull* 45:1615–1619
- Chang CH, Lin CC, Kawata Y, Hattori M, Namba T (1989) Prenylated xanthenes from *Cudrania cochinchinensis*. *Phytochemistry* 28:2823–2826
- Rukachaisirikul V, Ritthiwigrom T, Pinsa A, Sawangchote P, Taylor WC (2003) Xanthenes from the stem bark of *Garcinia nigrolineata*. *Phytochemistry* 64:1149–1156
- Helboe P, Arends P (1973) Xanthone studies. VI. Synthesis of jacareubin, isojacareubin, and some hydroxyxanthenes with allylic substituents. *Arch Pharm Chemi Sci Ed* 1:549–555
- Ito C, Miyamoto Y, Nakayama M, Kawai Y, Rao KS, Furukawa H (1997) A novel depsidone and some new xanthenes from *Garcinia species*. *Chem Pharm Bull* 45:1403–1413
- Wolfrom ML, Komitsky FJ, Fraenkel G, Looker JH, Dickey EE, McWain P (1963) Macluraxanthone and two accompanying pigments from the root bark of the osage orange. *Tetrahedron Lett* 4:749–755
- Delle Monache F, Ferrari F, Marini-Bettolo GB, Maxfield P, Cerini S, Fedeli W (1979) Vismiones from *Vismia baccifera* var. dealdata (H. B. K.): chemistry and X-ray structure determination. *Gazz Chim Ital* 109:301–310
- Botta B, Delle Monache F, Delle Monache G, Marini Bettolo GB, Oguakwa JU (1983) 3-Geranyloxy-6-methyl-1,8-dihydroxy-anthraquinone and vismiones C, D, and E, from *Psorospermum febrifugum*. *Phytochemistry* 22:539–542
- Kaennakam S, Siripong P, Tip-Pyang S (2015) Kaennacowanols A-C, three new xanthenes and their cytotoxicity from the roots of *Garcinia cowa*. *Fitoterapia* 102:171–176
- Ishiguro K, Nakajima M, Fukumoto H, Isoi K (1995) Co-occurrence of prenylated xanthenes and their cyclization products in cell suspension cultures of *Hypericum patulum*. *Phytochemistry* 38:867–869
- Ito C, Miyamoto Y, Rao KS, Furukawa H (1996) A novel dibenzofuran and two new xanthenes from *Calophyllum paniciflorum*. *Chem Pharm Bull* 44:441–443
- Vo HT, Nguyen NTT, Nguyen HT, Do KQ, Connolly JD, Maas G (2012) Cytotoxic tetraoxygenated xanthenes from the bark of *Garcinia schomburgkiana*. *Phytochem Lett* 5:553–557
- Kongkathip N, Kongkathip B, Siripong P, Sangma C, Luangkamin S, Niyomdech M (2003) Potent antitumor activity of synthetic 1,2-naphthoquinones and 1,4-naphthoquinones. *Bioorganic Med Chem* 11:3179–3191