

**TWO-WEIGHT, WEAK-TYPE NORM INEQUALITIES FOR  
FRACTIONAL INTEGRALS, CALDERÓN-ZYGMUND  
OPERATORS AND COMMUTATORS**

D. CRUZ-URIBE, SFO AND C. PÉREZ

ABSTRACT. We give  $A_p$ -type conditions which are sufficient for the two-weight, weak-type  $(p, p)$  inequalities for fractional integral operators, Calderón-Zygmund operators and commutators. For fractional integral operators, this solves a problem posed by Sawyer and Wheeden [28]. At the heart of all of our proofs is an inequality relating the Hardy-Littlewood maximal function and the sharp maximal function which is strongly reminiscent of the good- $\lambda$  inequality of Fefferman and Stein [13].

1. INTRODUCTION

Let  $M$  be the Hardy-Littlewood maximal operator. Given a pair of weights  $(u, v)$  and  $p$ ,  $1 < p < \infty$ , it is well known that the weak-type inequality

$$(1.1) \quad u(\{x \in \mathbb{R}^n : Mf(x) > t\}) \leq \frac{C}{t^p} \int_{\mathbb{R}^n} |f|^p v \, dx$$

holds if and only if  $(u, v) \in A_p$ : there exists a positive constant  $K$  such that for all cubes  $Q$ ,

$$(1.2) \quad \left( \frac{1}{|Q|} \int_Q u \, dx \right) \left( \frac{1}{|Q|} \int_Q v^{-p'/p} \, dx \right)^{p/p'} \leq K.$$

For other classical operators, however, the  $A_p$  condition is not sufficient for the weak  $(p, p)$  inequality. In fact, of the operators we are interested in, a necessary and sufficient condition for the weak  $(p, p)$  inequality is known only for fractional integral

---

1991 *Mathematics Subject Classification.* 42B20, 42B25.

*Key words and phrases.* weights, weak-type inequalities, fractional integral operators, Calderón-Zygmund operators, commutators.

We would like to thank E. Sawyer for sharing with the second author an unpublished manuscript which suggested our approach. We would also like to thank the referee for pointing out an error in Theorem 1.6.

The first author was supported by a Ford Foundation fellowship; the second author by DGICYT Grant PB40192, Spain. The second author is also grateful for the invitation and hospitality of the Centre de Recerca Matemàtica, Barcelona, where this work was completed.

operators. (See Sawyer [27].) This result is interesting and important, but it has the drawback that the condition involves the fractional integral operator.

Sufficient,  $A_p$ -type conditions can also be gotten from sufficient conditions for the strong  $(p, p)$  inequality. Neugebauer [18] showed that

$$(1.3) \quad \left( \frac{1}{|Q|} \int_Q u^r dx \right)^{1/rp} \left( \frac{1}{|Q|} \int_Q v^{-rp'/p} dx \right)^{1/rp'} \leq C, \quad r > 1,$$

is sufficient for the strong  $(p, p)$  inequality for the maximal operator, for Calderón-Zygmund operators and commutators. Sawyer and Wheeden [28] showed that for  $0 < \alpha < n$ ,

$$(1.4) \quad |Q|^{\alpha/n} \left( \frac{1}{|Q|} \int_Q u^r dx \right)^{1/rp} \left( \frac{1}{|Q|} \int_Q v^{-rp'/p} dx \right)^{1/rp'} \leq C, \quad r > 1,$$

is sufficient for the strong-type  $(p, p)$  inequality for fractional integral operators. (Additional sufficient conditions are found in [20], [21] and [24]. We give precise definitions of these operators in Section 2 below.)

In general, sufficient conditions for the weak  $(p, p)$  inequality which are derived from strong  $(p, p)$  conditions are not sharp. The purpose of this paper is to show that for the operators we consider, there are conditions which are weaker than (1.3) and (1.4) which are sufficient for the weak-type inequality. Roughly, it suffices to strengthen the  $A_p$  condition (1.2) by introducing a “power bump” on the left-hand term alone, rather than on both terms as in (1.3) and (1.4).

Our first result is for fractional integral operators. It solves a problem posed by Sawyer and Wheeden [28].

**Theorem 1.1.** *Given a pair of weights  $(u, v)$ ,  $p$ ,  $1 < p < \infty$ , and  $\alpha$ ,  $0 < \alpha < n$ , suppose that for some  $r > 1$  and for all cubes  $Q$ ,*

$$(1.5) \quad |Q|^{\alpha/n} \left( \frac{1}{|Q|} \int_Q u^r dx \right)^{1/rp} \left( \frac{1}{|Q|} \int_Q v^{-p'/p} dx \right)^{1/p'} \leq C < \infty.$$

*Then the fractional integral operator  $I_\alpha$  satisfies the weak  $(p, p)$  inequality*

$$(1.6) \quad u(\{x \in \mathbb{R}^n : |I_\alpha f(x)| > t\}) \leq \frac{C}{t^p} \int_{\mathbb{R}^n} |f|^p v dx.$$

Our second result is for Calderón-Zygmund operators.

**Theorem 1.2.** *Let  $T$  be a Calderón-Zygmund operator. Given a pair of weights  $(u, v)$  and  $p$ ,  $1 < p < \infty$ , suppose that for some  $r > 1$  and for all cubes  $Q$ ,*

$$(1.7) \quad \left( \frac{1}{|Q|} \int_Q u^r dx \right)^{1/rp} \left( \frac{1}{|Q|} \int_Q v^{-p'/p} dx \right)^{1/p'} \leq C < \infty.$$

*Then  $T$  satisfies the weak  $(p, p)$  inequality*

$$(1.8) \quad u(\{x \in \mathbb{R}^n : |Tf(x)| > t\}) \leq \frac{C}{t^p} \int_{\mathbb{R}^n} |f|^p v dx.$$

*Remark 1.3.* Though for clarity we have stated Theorem 1.2 for Calderón-Zygmund operators, it is true for a much larger class of operators. To be precise: if there exists some  $\delta$ ,  $0 < \delta < 1$ , and a constant  $C_\delta$  such that for every  $f \in C_0^\infty(\mathbb{R}^n)$ ,

$$(1.9) \quad M^\#(|Tf|^\delta)(x)^{1/\delta} \leq C_\delta Mf(x),$$

then (1.7) implies (1.8).

Alvarez and Pérez [3] showed that inequality (1.9) holds for Calderón-Zygmund operators. In this case it can be thought of as extending the classical estimate

$$(1.10) \quad M^\#(Tf)(x) \leq C_r M(|f|^r)(x)^{1/r},$$

where  $T$  is a regular singular integral operator and  $r > 1$ , (see García-Cuerva and Rubio de Francia [14, p. 204].) In some sense, (1.9) contains more information than (1.10) since the latter does not suffice to prove Theorem 1.2.

Alvarez and Pérez also showed that inequality (1.9), and so Theorem 1.2, hold for the following operators: weakly strongly singular integral operators (see C. Fefferman [12]), some pseudo-differential operators in the Hörmander class (see Hörmander [15]), and a class of oscillatory integral operators related to those introduced by Phong and Stein [25]. They used (1.9) to generalize Coifman's theorem [7] relating the  $L^p$  norm of singular integral operators and the maximal function.

*Remark 1.4.* For Calderón-Zygmund operators we have been able to prove stronger results; see [11]. By different methods we showed that we may replace the “power bump” in (1.7) by a “bump” in the scale of Orlicz spaces. More precisely, we replace the  $L^r$  norm by the  $L(\log L)^{p-1+\delta}$  norm with  $\delta > 0$ . However we are unable to extend these results to the broader class of operators discussed in the previous remark.

*Remark 1.5.* Conditions (1.5) and (1.7) are sufficient for the fractional maximal operator and the Hardy-Littlewood maximal operator to be bounded from  $L^{p'}(u^{-p'/p})$  to  $L^{p'}(v^{-p'/p})$ . (See [21], [22].) We conjecture that the boundedness of the corresponding maximal operator is itself sufficient for inequalities (1.6) and (1.8) to hold. In particular we believe that the Orlicz space conditions given in [21] and [22] are sufficient.

Our last result is about (linear) commutators. These operators are defined by

$$C_b^k f(x) = \int (b(x) - b(y))^k K(x, y) f(y) dy,$$

where  $K$  is a kernel satisfying the standard estimates and  $b$  is a locally integrable function. (See Section 2 for a precise definition.)

Since commutators have a greater degree of “singularity” than the corresponding Calderón-Zygmund operators, we need a slightly stronger condition. Roughly, we need to “bump” the right-hand term as well, but it suffices to do so in the scale of Orlicz spaces. Recall that if  $B$  is an increasing Young function and if  $Q$  is any cube, we define the mean Luxemburg norm of a measurable function  $f$  with respect to  $B$  by

$$\|f\|_{B,Q} = \inf \left\{ \lambda > 0 : \frac{1}{|Q|} \int_Q B \left( \frac{|f|}{\lambda} \right) dx \leq 1 \right\}.$$

(For more information on Orlicz spaces, see Section 2 below.)

**Theorem 1.6.** *Let  $T$  be a Calderón-Zygmund operator and  $b$  a function in BMO. Given a pair of weights  $(u, v)$ ,  $p$ ,  $1 < p < \infty$ , and  $k \geq 0$ , suppose that for some  $r > 1$  and for all cubes  $Q$ ,*

$$(1.11) \quad \left( \frac{1}{|Q|} \int_Q u^r dx \right)^{1/rp} \|v^{-1/p}\|_{C_k, Q} \leq C < \infty,$$

where  $C_k(t) = t^{p'} \log(e + t)^{kp'}$ . Then the commutator  $C_b^k$  satisfies the weak  $(p, p)$  inequality

$$(1.12) \quad u(\{x \in \mathbb{R}^n : |C_b^k f(x)| > t\}) \leq \frac{C}{t^p} \int_{\mathbb{R}^n} |f|^p v dx.$$

When  $k = 0$ ,  $C_b^0 = T$ , and so in this case Theorem 1.6 reduces to Theorem 1.2.

*Remark 1.7.* As a corollary to Theorem 1.6 we get a new proof of the one-weight, strong  $(p, p)$  norm inequality for commutators, which was first proved in a more general form by Alvarez, Bagby, Kurtz and Pérez [2] and Segovia and Torrea [29]. If  $w \in A_p$  then  $w$  and  $w^{-p'/p}$  both satisfy the reverse Hölder inequality and so inequality (1.11) holds for some  $r > 1$  and for  $p \pm \epsilon$ . The strong-type inequality follows by interpolation.

The proofs of Theorems 1.1, 1.2 and 1.6 all follow the same outline. Each relies on our so-called principal lemma, Theorem 3.4 below, which relates the Hardy-Littlewood maximal operator and the Fefferman-Stein sharp maximal operator via an inequality

strongly reminiscent of a good- $\lambda$  inequality. To apply Theorem 3.4 we use three results which relate the given operator, the sharp maximal operator and the maximal operator. For Calderón-Zygmund operators this is inequality (1.9). Similar inequalities hold for fractional integral operators and commutators: see Lemmas 4.4 and 6.1.

The remainder of this paper is organized as follows: in Section 2 we give a number of definitions and lemmas needed in later sections. The heart of the paper is Section 3, where we prove Theorem 3.4. Finally, in Sections 4, 5 and 6 we prove Theorems 1.1, 1.2 and 1.6.

Throughout this paper all notation is standard or will be defined as needed. All cubes are assumed to have their sides parallel to the coordinate axes. Given a cube  $Q$ ,  $l(Q)$  will denote the length of its sides and for any  $r > 0$ ,  $rQ$  will denote the cube with the same center as  $Q$  and such that  $l(rQ) = rl(Q)$ . We will denote the collection of all dyadic cubes by  $\Delta$  and by  $\Delta(Q)$  the collection of all dyadic subcubes relative to the (not necessarily dyadic) cube  $Q$ . By weights we will always mean non-negative, locally integrable functions which are positive on a set of positive measure. Given a Lebesgue measurable set  $E$  and a weight  $w$ ,  $|E|$  will denote the Lebesgue measure of  $E$  and  $w(E) = \int_E w dx$ . Given  $1 < p < \infty$ ,  $p' = p/(p-1)$  will denote the conjugate exponent of  $p$ . Finally,  $C$  will denote a positive constant whose value may change at each appearance.

## 2. PRELIMINARY IDEAS

In this section we give a number of definitions and lemmas needed in later sections.

**The main operators.** First we define the operators in Theorems 1.1, 1.2 and 1.6.

*Fractional integral operators:* Given  $\alpha$ ,  $0 < \alpha < n$ , define the fractional integral operator of order  $\alpha$  by

$$I_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}} dy.$$

For more information, see Stein [31, pp. 117-120].

*Calderón-Zygmund operators:* Given a kernel  $K$  on  $\mathbb{R}^n \times \mathbb{R}^n$ —i.e. a locally integrable, complex-valued function defined off the diagonal—we say that it satisfies the standard estimates if there exist  $\delta$ ,  $0 < \delta \leq 1$ , and  $C$  finite such that for all distinct points  $x$  and  $y$  in  $\mathbb{R}^n$ , and all  $z$  such that  $|x-z| < \frac{1}{2}|x-y|$ :

- (1)  $|K(x, y)| \leq C|x-y|^{-n}$ ;
- (2)  $|K(x, y) - K(z, y)| \leq C|x-z|^\delta/|x-y|^{n+\delta}$ ;
- (3)  $|K(y, x) - K(y, z)| \leq C|x-z|^\delta/|x-y|^{n+\delta}$ .

A bounded linear operator  $T : C_0^\infty(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}^n)$  (here  $\mathcal{D}'$  is the space of distributions) is said to be associated with a kernel  $K$  if

$$\langle Tf, g \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} K(x, y)g(x)f(y) dx dy$$

for all  $f$  and  $g$  in  $C_0^\infty(\mathbb{R}^n)$  with  $\text{supp}(f) \cap \text{supp}(g) = \emptyset$ .  $T$  is said to be a Calderón-Zygmund operator if its associated kernel satisfies the standard estimates and it extends to a bounded linear operator on  $L^2$ . For more information, see Coifman and Meyer [8] and Christ [6].

Important examples of such operators are the Calderón-Zygmund singular integral operators:

$$Tf(x) = \text{p. v.} \int_{\mathbb{R}^n} k(x-y)f(y) dy,$$

where  $k \in L_{loc}^1(\mathbb{R}^n \setminus \{0\})$  and  $K(x, y) = k(x-y)$  satisfies the standard estimates. For more information see García-Cuerva and Rubio de Francia [14, p. 192].

*Commutators:* Given a Calderón-Zygmund operator  $T$  and a function  $b$  in BMO, let  $M_b$  denote multiplication by  $b$ . We define the linear operators  $C_b^k$  by  $C_b^0 = T$ ,  $C_b^1 = [M_b, T] = M_bT - TM_b$ , and for  $k > 1$ ,  $C_b^k = [M_b, C_b^{k-1}]$ . If  $f \in C_0^\infty(\mathbb{R}^n)$  then

$$C_b^k f(x) = \int (b(x) - b(y))^k K(x, y)f(y) dy, \quad x \notin \text{supp}(f).$$

Commutators were introduced by Coifman, Rochberg and Weiss [9], who showed they are bounded on  $L^p$ ,  $1 < p < \infty$ .

**Maximal operators.** Key to the proofs of our results are a number of maximal operators. For completeness we give their definitions here.

*The maximal operator:* Given a locally integrable function  $f$  and  $\alpha$ ,  $0 \leq \alpha < n$ , define

$$M_\alpha f(x) = \sup_{Q \ni x} \frac{1}{|Q|^{1-\alpha/n}} \int_Q |f| dy.$$

If  $\alpha = 0$  this is the Hardy-Littlewood maximal operator and we write  $Mf$  for  $M_0f$ ; if  $0 < \alpha < n$  this is the fractional maximal operator of order  $\alpha$ . We use the Hardy-Littlewood maximal operator to control Calderón-Zygmund operators and commutators, and the fractional maximal operator to control fractional integral operators. (See inequality (1.9) and Lemmas 4.4 and 6.1.)

We define the dyadic maximal and fractional maximal operators  $M^d$  and  $M_\alpha^d$  similarly except the supremums are restricted to dyadic cubes containing  $x$ . Given  $\delta > 0$  we define the  $\delta$ -maximal operator by  $M_\delta f(x) = M(|f|^\delta)(x)^{1/\delta}$ . We define  $M_\delta^d$  similarly. From the context there should be no confusion between the fractional maximal operator and the  $\delta$ -maximal operator.

*The sharp maximal operator:* Given a locally integrable function  $f$  and a cube  $Q$ , let  $f_Q$  denote the average of  $f$  over  $Q$ :

$$f_Q = \frac{1}{|Q|} \int_Q f \, dx.$$

Define the sharp maximal function of  $f$  by

$$M^\# f(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y) - f_Q| \, dy.$$

The sharp maximal function was introduced by Fefferman and Stein [13]. Again, define the dyadic sharp maximal function  $M^{\#,d}$  by restricting the supremum to dyadic cubes. Given  $\delta > 0$ , define the sharp  $\delta$ -maximal function by  $M_\delta^\# f(x) = M^\#(|f|^\delta)(x)^{1/\delta}$ , and define  $M_\delta^{\#,d}$  similarly.

**Orlicz spaces.** In Section 6 we will need the following facts about Orlicz spaces. (For further information see Bennett and Sharpley [4] or Rao and Ren [26].) A function  $B : [0, \infty) \rightarrow [0, \infty)$  is a Young function if it is convex and increasing, and if  $B(0) = 0$  and  $B(t) \rightarrow \infty$  as  $t \rightarrow \infty$ .

Given a Young function  $B$ , define the mean Luxemburg norm of  $f$  on a cube  $Q$  by

$$\|f\|_{B,Q} = \inf \left\{ \lambda > 0 : \frac{1}{|Q|} \int_Q B \left( \frac{|f|}{\lambda} \right) \, dy \leq 1 \right\}.$$

When  $B(t) = t^p$ ,  $1 \leq p < \infty$ ,

$$\|f\|_{B,Q} = \left( \frac{1}{|Q|} \int_Q |f|^p \, dx \right)^{1/p};$$

that is, the Luxemburg norm coincides with the (normalized)  $L^p$  norm. There is another characterization of the Luxemburg norm, due to Krasnosel'skiĭ and Rutickiĭ [17, p. 92] (also see Rao and Ren [26, p. 69]) which we will need:

$$(2.1) \quad \|f\|_{B,Q} \leq \inf_{s>0} \left\{ s + \frac{s}{|Q|} \int_Q B \left( \frac{|f|}{s} \right) \, dx \right\} \leq 2\|f\|_{B,Q}.$$

Given three Young functions  $A$ ,  $B$  and  $C$  such that for all  $t > 0$ ,

$$(2.2) \quad A^{-1}(t)C^{-1}(t) \leq B^{-1}(t),$$

then we have the following generalized Hölder's inequality due to O'Neil [19]: for any cube  $Q$  and all functions  $f$  and  $g$ ,

$$(2.3) \quad \|fg\|_{B,Q} \leq 2\|f\|_{A,Q}\|g\|_{C,Q}.$$

Define the maximal operator  $M_B$  by

$$M_B f(x) = \sup_{Q \ni x} \|f\|_{B,Q}.$$

The dyadic maximal operator  $M_B^d$  is defined in similarly, except the supremum is restricted to dyadic cubes containing  $x$ . It follows from an inequality due to Stein [30] that for  $k \geq 1$ , if  $B_k(t) = t \log(e+t)^{k-1}$ , then  $M^k f \approx M_{B_k} f$ , where  $M^k = M \cdot M \cdots M$  is the  $k$ -th iterate of the maximal function. (See Carozza and Passarelli di Napoli [5] and the references given there.)

**The Calderón-Zygmund Decomposition.** Our proofs depend heavily on the Calderón-Zygmund decomposition and a generalization of it to Orlicz space norms. To be precise and to establish notation, we state the result here. For a proof see [22]; this is an adaptation of the classical proof given in García-Cuerva and Rubio de Francia [14, p. 137].

**Lemma 2.1.** *Given a Young function  $B$ , suppose  $f$  is a non-negative function such that  $\|f\|_{B,Q}$  tends to zero as  $l(Q)$  tends to infinity. Then for each  $t > 0$  there exists a disjoint collection of dyadic cubes  $\{C_i^t\}$  such that for each  $i$ ,  $t < \|f\|_{B,C_i^t} \leq 2^nt$ ,*

$$\{x \in \mathbb{R}^n : M_B^d f(x) > t\} = \bigcup_i C_i^t,$$

and

$$\{x \in \mathbb{R}^n : M_B f(x) > 4^nt\} \subset \bigcup_i 3C_i^t.$$

Moreover, the cubes are maximal: if  $Q$  is a dyadic cube such that  $Q \subset \{M_B^d f(x) > t\}$ , then  $Q \subset C_i^t$  for some  $i$ .

To recapture the classical lemma, let  $B(t) = t$  and note that if  $f \in L^q$  for some  $q$ ,  $1 \leq q < \infty$ , then

$$\|f\|_{B,Q} = \frac{1}{|Q|} \int_Q f \, dx \rightarrow 0 \quad \text{as} \quad |Q| \rightarrow \infty.$$

More generally, to apply Lemma 2.1 it suffices to assume that  $f$  is bounded and has compact support.

### 3. THE PRINCIPAL LEMMA

In this section we prove our principal lemma: an inequality linking the sharp maximal function and the Hardy-Littlewood maximal function. In spirit, though not in detail it resembles the good- $\lambda$  inequality of Fefferman and Stein [13]. (Also see García-Cuerva and Rubio de Francia [14, pp. 161-3] and Journé [16, p. 41].)



To state the principal lemma we first need a definition and a lemma.

**Definition 3.1.** Given  $r > 1$  and a weight  $u$ , define the set function  $A_u^r$  on measurable sets  $E \subset \mathbb{R}^n$  by

$$A_u^r(E) = |E|^{1/r'} \left( \int_E u^r dx \right)^{1/r} = |E| \left( \frac{1}{|E|} \int_E u^r dx \right)^{1/r}.$$

(The second equality holds provided  $|E| > 0$ .)

**Lemma 3.2.** For any  $r > 1$  and weight  $u$ , the set function  $A_u^r$  has the following properties:

- (1) If  $E \subset F$  then  $A_u^r(E) \leq \left( \frac{|E|}{|F|} \right)^{1/r'} A_u^r(F)$ ;
- (2)  $u(E) \leq A_u^r(E)$ ;
- (3) If  $\{E_j\}$  is a sequence of disjoint sets and  $\bigcup_j E_j = E$  then

$$\sum_j A_u^r(E_j) \leq A_u^r(E).$$

*Proof.* Condition (1) follows immediately from Definition 3.1, and Condition (2) is just Hölder's inequality. Condition (3) also follows from Hölder's inequality:

$$\begin{aligned} \sum_j |E_j|^{1/r'} \left( \int_{E_j} u^r dx \right)^{1/r} &\leq \left( \sum_j |E_j| \right)^{1/r'} \left( \sum_j \int_{E_j} u^r dx \right)^{1/r} \\ &= |E|^{1/r'} \left( \int_E u^r dx \right)^{1/r}. \end{aligned}$$

□

*Remark 3.3.* The key property is Condition (1), which plays the same role that the  $A_\infty$  condition plays in the proof of weighted good- $\lambda$  inequalities. (See, for example, Journé [16, p. 41].) If  $A_u$  were another set function which satisfied Conditions (2) and (3) of Lemma 3.2, satisfied

$$(3.1) \quad A_u(E) \leq \phi(|E|/|F|)A_u(F), \quad \phi(t) \rightarrow 0 \text{ as } t \rightarrow 0,$$

and for some  $r > 1$  satisfied (for technical reasons in the proof)  $A_u(E) \leq CA_u^r(E)$ , we could immediately derive corresponding conditions governing weak-type norm inequalities for the operators we are interested in.

Originally, we had hoped to replace the “power bumps” in (1.5), (1.7) and (1.11) by Orlicz space conditions. Intuitively, the appropriate set function would be  $A(E) = |E| \|u\|_{B,E}$ , where  $B$  is some Young function—for example,  $B(t) = t \log(e+t)^\delta$ ,  $\delta > 0$ .

For such  $B$ , Conditions (2) and (3) hold; we will show this in the course of proving Lemma 5.1 below. However, Condition (1) fails.

*Remark added in proof.* The first author and A. Fiorenza have characterized the class of Young functions  $B$  for which  $A(E) = |E| \|u\|_{B,E}$  satisfies (3.1). This class includes Orlicz functions which grow slower than  $t^r$  for any  $r > 1$ . These results will appear in [10].

We can now state and prove our principal lemma.

**Theorem 3.4.** *Given a non-negative function  $f \in L^q$  for some  $q$ ,  $1 \leq q < \infty$ ,  $r$ ,  $1 < r \leq q'$ , a weight  $u$ , and  $\delta > 0$ , then there exists  $\epsilon > 0$  such that for each  $t > 0$  there exists a subcollection  $\{Q_j^t\}$  of dyadic cubes from the Calderón-Zygmund decomposition of  $f^\delta$  at height  $t^\delta$ ,  $\{C_i^{t^\delta}\}$ , with the property that*

$$\left( \frac{1}{|Q_j^t|} \int_{Q_j^t} |f^\delta - (f^\delta)_{Q_j^t}| dx \right)^{1/\delta} > \epsilon^{1/\delta} t,$$

and such that for all  $p \geq q/r'$ ,

$$(3.2) \quad \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : M_\delta^d f(x) > t\}) \leq C \sup_{t>0} t^p \sum_j A_u^r(Q_j^t).$$

The constants  $\epsilon$  and  $C$  depend only on  $r$ ,  $p$  and  $n$ .

As a corollary to the proof we have the following stronger inequality.

**Corollary 3.5.** *With the same hypotheses and notation as Theorem 3.4, we have that*

$$(3.3) \quad \sup_{t>0} t^p \sum_i A_u^r(C_i^{t^\delta}) \leq C \sup_{t>0} t^p \sum_j A_u^r(Q_j^t).$$

*Remark 3.6.* In our applications of these results we always have  $f \in L^q$  for any  $q > 1$ , so we can get any value of  $p \geq 1$ . If  $r$  can be taken close to 1 then we can get any  $p > 0$ .

*Proof.* First note that it will suffice to prove this result for  $\delta = 1$ . For arbitrary  $\delta > 0$ ,  $M_\delta^d f(x) > t$  is equivalent to  $M^d(f^\delta)(x) > t^\delta$ , so the general case follows if we replace  $f$  by  $f^\delta$  and  $t$  by  $t^\delta$ .

Second, we may assume that  $u$  is bounded and has compact support. To see that the general case follows, fix a weight  $u$  and let  $u_k = \min(u, k)\chi_{B(0,k)}$ . Since  $u_k$  is bounded, inequalities (3.2) and (3.3) hold with  $u$  replaced by  $u_k$ . Since  $\lim_n u_k = \sup_n u_k = u$ ,

if we take the limit as  $n$  tends to infinity we may exchange limit and supremum and apply the monotone convergence theorem to get the desired result.

Fix  $p, q/r' \leq p < \infty$ , and fix  $f$ . For each  $t > 0$ , let  $\Omega_t = \{x \in \mathbb{R}^n : M^d f(x) > t\}$ . Now fix  $N = 2^n + 1$  (the reason for this choice will be clear below); by the Calderón-Zygmund decomposition, Lemma 2.1,  $\Omega_{Nt} = \bigcup_k C_k^{Nt}$  and  $\Omega_t = \bigcup_i C_i^t$ . By maximality, for each  $k$ ,  $C_k^{Nt} \subset C_i^t$  for some  $i$ . By Lemma 3.2, Conditions (2) and (3),

$$\begin{aligned} t^p u(\Omega_{Nt}) &= t^p \sum_k u(C_k^{Nt}) \\ &\leq t^p \sum_k A_u^r(C_k^{Nt}) \\ &= t^p \sum_i \sum_{C_k^{Nt} \subset C_i^t} A_u^r(C_k^{Nt}) \\ &\leq t^p \sum_i A_u^r(\Omega_{Nt} \cap C_i^t). \end{aligned}$$

Fix  $\epsilon < N^{-pr'}$ ; again the reason for this choice will be clear below. Divide the indices  $i$  into two sets:  $i \in F$  if

$$\frac{1}{|C_i^t|} \int_{C_i^t} |f - f_{C_i^t}| dx \leq \epsilon t,$$

and  $i \in G$  if the opposite inequality holds. The cubes  $\{C_i^t : i \in G\}$  are the cubes in the conclusion of the theorem, and we relabel them  $\{Q_j^t\}$ .

If  $i \in F$  then we claim that

$$A_u^r(\Omega_{Nt} \cap C_i^t) \leq \epsilon^{1/r'} A_u^r(C_i^t).$$

By Lemma 3.2, Condition (1), it will suffice to show that

$$|\Omega_{Nt} \cap C_i^t| \leq \epsilon |C_i^t|.$$

By the maximality of the Calderón-Zygmund decomposition, if  $x \in \Omega_{Nt} \cap C_i^t$  then

$$M^d f(x) = M^d(f\chi_{C_i^t})(x).$$

Hence,

$$\begin{aligned} \Omega_{Nt} \cap C_i^t &= \{x \in C_i^t : M^d(f\chi_{C_i^t})(x) > Nt\} \\ &= \{x \in C_i^t : M^d(f\chi_{C_i^t})(x) - f_{C_i^t} > Nt - f_{C_i^t}\} \\ &\subset \{x \in C_i^t : M^d(|f - f_{C_i^t}|\chi_{C_i^t})(x) > t\}. \end{aligned}$$

Since the dyadic maximal operator is weak-type  $(1, 1)$  with constant 1, (see Journé [16, p. 10]), and since  $i \in F$ ,

$$|\Omega_{Nt} \cap C_i^t| \leq \frac{1}{t} \int_{C_i^t} |f - f_{C_i^t}| dx \leq \epsilon |C_i^t|.$$

Therefore, we have shown that

$$(3.4) \quad \begin{aligned} t^p \sum_k A_u^r(C_k^{Nt}) &\leq t^p \sum_i A_u^r(\Omega_{Nt} \cap C_i^t) \\ &\leq t^p \sum_{i \in F} A_u^r(C_i^t) + t^p \sum_{i \in G} A_u^r(C_i^t) \end{aligned}$$

$$(3.5) \quad \leq \epsilon^{1/r'} t^p \sum_i A_u^r(C_i^t) + t^p \sum_j A_u^r(Q_j^t).$$

Therefore, if we take the supremum of (3.5) over  $0 < t < M$  and the supremum of (3.4) over  $0 < t < M/N$ , we get

$$\sup_{0 < t < M/N} t^p \sum_k A_u^r(C_k^{Nt}) \leq \sup_{0 < t < M} \epsilon^{1/r'} t^p \sum_i A_u^r(C_i^t) + \sup_{0 < t < M} t^p \sum_j A_u^r(Q_j^t);$$

equivalently,

$$\sup_{0 < t < M} t^p \sum_i A_u^r(C_i^t) \leq \epsilon^{1/r'} N^p \sup_{0 < t < M} t^p \sum_i A_u^r(C_i^t) + N^p \sup_{0 < t < M} t^p \sum_j A_u^r(Q_j^t).$$

To get the desired inequality we need to re-arrange terms; to do this we need to show that for each  $M > 0$ ,

$$\sup_{0 < t < M} t^p \sum_i A_u^r(C_i^t) < \infty.$$

But for fixed  $t$ , by Condition (3) of Lemma 3.2 and the definition of  $A_u^r$ ,

$$\begin{aligned} t^p \sum_i A_u^r(C_i^t) &\leq t^p A_u^r(\Omega_t) \\ &= t^p |\Omega_t|^{1/r'} \left( \int_{\Omega_t} u^r dx \right)^{1/r}. \end{aligned}$$

Let  $B$  be the support of  $u$ ; by assumption  $|B| < \infty$ . Further,  $u$  is bounded. Therefore,

$$\leq |B|^{1/r} \|u\|_\infty t^p |\Omega_t|^{1/r'}.$$

Since  $f \in L^q$ , by the weak  $(q, q)$  inequality for the dyadic maximal operator,

$$\leq |B|^{1/r} \|u\|_\infty t^{p-q/r'} \|f\|_q^{q/r'}.$$

Therefore, since  $p - q/r' \geq 0$ ,

$$\sup_{0 < t < M} t^p \sum_i A_u^r(C_i^t) < |B|^{1/r} \|u\|_\infty M^{p-q/r'} \|f\|_q^{1/r'} < \infty.$$

Thus we can re-arrange terms; since  $\epsilon < N^{-pr'}$ , we get

$$\begin{aligned} \sup_{0 < t < M} t^p u(\{x \in \mathbb{R}^n : M^d f(x) > t\}) &\leq \sup_{0 < t < M} t^p \sum_i A_u^r(C_i^t) \\ &\leq \frac{N^p}{1 - \epsilon^{1/r'} N^p} \sup_{0 < t < M} t^p \sum_j A_u^r(Q_j^t). \end{aligned}$$

Since this holds for all  $M > 0$ , if we take the limit as  $M$  tends to infinity we get inequalities (3.2) and (3.3).  $\square$

#### 4. FRACTIONAL INTEGRAL OPERATORS

In this section we prove Theorem 1.1. The proof depends on three lemmas; the first two are due to Sawyer and Wheeden [28].

**Lemma 4.1.** *Given a non-negative function  $f$  and  $\alpha$ ,  $0 < \alpha < n$ , there exists a constant  $C_\alpha$ , depending only on  $\alpha$  and  $n$ , such that for any cube  $Q_0$ ,*

$$\sum_{Q \in \Delta(Q_0)} |Q|^{\alpha/n} \int_Q f \, dx \leq C_\alpha |Q_0|^{\alpha/n} \int_{Q_0} f \, dx.$$

**Definition 4.2.** Given  $\alpha$ ,  $0 < \alpha < n$ , and  $z \in \mathbb{R}^n$ , define the translated dyadic fractional integral operator  $I_{\alpha,z}^d$  by

$$I_{\alpha,z}^d f(x) = \sum_{\substack{Q+z \in \Delta \\ Q \ni x}} |Q|^{\alpha/n-1} \int_Q f \, dy.$$

If  $z = 0$  we write  $I_\alpha^d$  for  $I_{\alpha,0}^d$ .

**Lemma 4.3.** *Given a weight  $u$ ,  $\alpha$ ,  $0 < \alpha < n$ , and  $p$ ,  $1 < p < \infty$ , then there exists a constant  $C$  such that for every function  $f$ ,*

$$\sup_{t>0} t^p u(\{x \in \mathbb{R}^n : |I_\alpha f(x)| > t\}) \leq C \sup_{z \in \mathbb{R}^n} \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : |I_{\alpha,z}^d f(x)| > t\}).$$

**Lemma 4.4.** *Given  $\alpha$ ,  $0 < \alpha < n$ , there exists a constant  $D_\alpha$  such that for any function  $f$ , dyadic cube  $Q_0$  and  $x \in Q_0$ ,*

$$\frac{1}{|Q_0|} \int_{Q_0} |I_\alpha^d f - (I_\alpha^d f)_{Q_0}| \, dx \leq D_\alpha M_\alpha^d f(x).$$

*Proof.* By the definition of  $I_\alpha^d$ , for  $x \in Q_0$ ,

$$I_\alpha^d f(x) = \sum_{\substack{x \in Q \in \Delta \\ Q \subset Q_0}} |Q|^{\alpha/n-1} \int_Q f dx + \sum_{\substack{Q \in \Delta \\ Q_0 \subseteq Q}} |Q|^{\alpha/n-1} \int_Q f dx;$$

hence,

$$\frac{1}{|Q_0|} \int_{Q_0} I_\alpha^d f dx = \frac{1}{|Q_0|} \sum_{\substack{Q \in \Delta \\ Q \subset Q_0}} |Q|^{\alpha/n} \int_Q f dx + \sum_{\substack{Q \in \Delta \\ Q_0 \subseteq Q}} |Q|^{\alpha/n-1} \int_Q f dx.$$

Therefore, by Lemma 4.1,

$$\begin{aligned} \frac{1}{|Q_0|} \int_{Q_0} |I_\alpha^d f - (I_\alpha^d f)_{Q_0}| dx &\leq \frac{2}{|Q_0|} \sum_{\substack{Q \in \Delta \\ Q \subset Q_0}} |Q|^{\alpha/n} \int_Q |f| dx \\ &\leq 2C_\alpha |Q_0|^{\alpha/n-1} \int_{Q_0} |f| dx \\ &\leq 2C_\alpha M_\alpha^d f(x). \end{aligned}$$

□

*Proof of Theorem 1.1.* By Lemma 4.3 it will suffice to prove inequality (1.6) with  $I_\alpha$  replaced by  $I_{\alpha,z}^d$  and with a constant independent of  $z$ . In the proof that follows it will be clear that all the constants are independent of  $z$ , so in fact it will suffice to prove inequality (1.6) for  $I_\alpha^d$ .

Since  $I_\alpha^d$  is a positive operator, by a standard argument we may assume that  $f$  is non-negative, bounded and has compact support. Fix  $p$ ,  $1 < p < \infty$ ; then  $I_\alpha^d f \in L^q$ , where  $q > 1$  is such that  $p \geq q/r'$ , so we can apply Theorem 3.4 to it. Let  $\delta = 1$ . Then there exists  $\epsilon > 0$  such that for each  $t > 0$  there exists a sequence of disjoint dyadic cubes  $\{Q_j^t\}$  such that

$$(4.1) \quad \frac{1}{|Q_j^t|} \int_{Q_j^t} |I_\alpha^d f - (I_\alpha^d f)_{Q_j^t}| dx > \epsilon t$$

and (by the Lebesgue differentiation theorem)

$$\begin{aligned} \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : |I_\alpha^d f(x)| > t\}) &\leq \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : M^d(I_\alpha f)(x) > t\}) \\ &\leq C \sup_{t>0} t^p \sum_j A_u^r(Q_j^t). \end{aligned}$$

Fix  $t$ ; then by Lemma 4.4, for each  $j$ ,

$$Q_j^t \subset \{x \in \mathbb{R}^n : M_\alpha^d f(x) > \epsilon D_\alpha^{-1} t\}.$$

By an argument analogous to that for the dyadic maximal operator (cf. Lemma 2.1), we can write the right-hand side as the union of disjoint dyadic cubes  $\{P_k^t\}$  such that for each  $k$ ,

$$|P_k^t|^{\alpha/n-1} \int_{P_k^t} f \, dx > \epsilon D_\alpha^{-1} t.$$

Further, the  $P_k^t$ 's are maximal with this property; in particular, for each  $j$  there exists  $k$  such that  $Q_j^t \subset P_k^t$ . Therefore, by Lemma 3.2, Condition (3),

$$\begin{aligned} t^p \sum_j A_u^r(Q_j^t) &= t^p \sum_k \sum_{Q_j^t \subset P_k^t} A_u^r(Q_j^t) \\ &\leq t^p \sum_k A_u^r(P_k^t) \\ &\leq (\epsilon^{-1} D_\alpha)^p \sum_k |P_k^t| \left( \frac{1}{|P_k^t|} \int_{P_k^t} u^r \, dx \right)^{1/r} \left( |P_k^t|^{\alpha/n-1} \int_{P_k^t} f \, dx \right)^p. \end{aligned}$$

By Hölder's inequality and inequality (1.5),

$$\begin{aligned} &\leq C \sum_k |P_k^t|^{\alpha p/n} \left( \frac{1}{|P_k^t|} \int_{P_k^t} u^r \, dx \right)^{1/r} \left( \frac{1}{|P_k^t|} \int_{P_k^t} v^{-p'/p} \, dx \right)^{p/p'} \int_{P_k^t} f^p v \, dx \\ &\leq C \sum_k \int_{P_k^t} f^p v \, dx \\ &\leq C \int_{\mathbb{R}^n} f^p v \, dx. \end{aligned}$$

The constant is independent of  $t$ , so if we take the supremum over all  $t > 0$  we get inequality (1.6). □

*Remark 4.5.* At the cost of a more complex argument similar to that for Calderón-Zygmund operators (cf. Lemma 5.1 below) we could dispense with the dyadic fractional integral operator and prove Theorem 1.1 directly for  $I_\alpha$ . The key inequality is the non-dyadic analogue of Lemma 4.4 due to Adams [1]:  $M^\#(I_\alpha f)(x) \leq C M_\alpha f(x)$ .

## 5. CALDERÓN-ZYGMUND OPERATORS

In this section we prove Theorem 1.2. The proof is similar to that of Theorem 1.1, but is complicated by the fact that we cannot pass to an equivalent dyadic operator. To compensate we need the following lemma which is also needed in the proof of Theorem 1.6.

**Lemma 5.1.** *Let  $B$  be a Young function. Suppose that for some function  $f \in L^q$ ,  $1 \leq q < \infty$ , and for some  $t > 0$  there exists a constant  $\mu$ ,  $0 < \mu \leq 1$ , and a collection of dyadic cubes  $\{Q_j\}$  such that for each  $j$ ,*

$$|Q_j \cap \{x \in \mathbb{R}^n : M_B f(x) > t\}| \geq \mu |Q_j|.$$

*Then there exists a constant  $\nu > 0$ , depending on  $n$  and  $\mu$ , and a subcollection  $\{P_k\}$  of the Calderón-Zygmund decomposition with respect to  $B$  of  $f$  at height  $\nu t$ ,  $\{C_i^{\nu t}\}$ , such that for each  $j$ ,  $Q_j \subset 3P_k$  for some  $k$ .*

*If we replace  $M_B$  by  $M_B^d$  in the hypothesis then we can strengthen the conclusion by finding  $P_k$ 's such that  $Q_j \subset P_k$  and by letting  $\mu = \nu$ .*

*Proof.* We first consider the non-dyadic case. By Lemma 2.1,

$$E_t = \{x \in \mathbb{R}^n : M_B f(x) > t\} \subset \bigcup_i 3C_i^{\gamma t},$$

where  $\gamma = 4^{-n}$ . If we had  $Q_j \subset 3C_i^{\gamma t}$  for some  $i$  we would be done, but this need not be the case, even if  $\mu = 1$ . However, for each  $j$  there is a collection of indices  $A_j$  such that

$$Q_j \cap E_t \subset \bigcup_{i \in A_j} 3C_i^{\gamma t} \quad \text{and} \quad 3C_i^{\gamma t} \cap Q_j \neq \emptyset, \quad i \in A_j.$$

There are two possibilities: first, there exists  $i \in A_j$  such that  $l(Q_j) \leq l(3C_i^{\gamma t})$ . Then  $Q_j \subset 9C_i^{\gamma t}$  and by inequality (2.1),

$$\begin{aligned} 2\|f\|_{B,9C_i^{\gamma t}} &\geq \inf_{s>0} \left\{ s + \frac{s}{|9C_i^{\gamma t}|} \int_{9C_i^{\gamma t}} B\left(\frac{|f|}{s}\right) dx \right\} \\ &\geq 9^{-n} \inf_{s>0} \left\{ s + \frac{s}{|C_i^{\gamma t}|} \int_{C_i^{\gamma t}} B\left(\frac{|f|}{s}\right) dx \right\} \\ &= 9^{-n} \|f\|_{B,C_i^{\gamma t}} \\ &> 9^{-n} \gamma t. \end{aligned}$$



Alternatively,  $l(Q_j) > l(3C_i^{\gamma t})$  for all  $i \in A_j$ . But then for each  $i \in A_j$ ,  $3C_i^{\gamma t} \subset 3Q_j$ , and so

$$\begin{aligned}
2|3Q_j| \|f\|_{B,3Q_j} &\geq \inf_{s>0} \left\{ s|3Q_j| + s \int_{3Q_j} B\left(\frac{|f|}{s}\right) dx \right\} \\
&\geq \sum_{i \in A_j} \inf_{s>0} \left\{ s|C_i^{\gamma t}| + s \int_{C_i^{\gamma t}} B\left(\frac{|f|}{s}\right) dx \right\} \\
&= \sum_{i \in A_j} |C_i^{\gamma t}| \|f\|_{B,C_i^{\gamma t}} \\
&> 3^{-n} \gamma t \sum_{i \in A_j} |3C_i^{\gamma t}| \\
&\geq 3^{-n} \gamma t |Q_j \cap E_t| \\
&\geq 9^{-n} \mu \gamma t |3Q_j|.
\end{aligned}$$

So in either case, for each  $j$  there exists a cube  $\bar{Q}_j$  containing  $Q_j$  such that

$$\|f\|_{B,\bar{Q}_j} > \frac{\mu \gamma t}{2 \cdot 9^n}.$$

Now by the same argument that is used to prove the Calderón-Zygmund decomposition, Lemma 2.1, we can show that there exists a subcollection  $\{P_k\}$  of  $\{C_i^{\nu t}\}$ ,  $\nu = \frac{1}{2} \mu \gamma 36^{-n} = \frac{1}{2} \mu 144^{-n}$ , such that for each  $j$ ,  $Q_j \subset \bar{Q}_j \subset 3P_k$  for some  $k$ . This completes the proof for  $M_B$ .

The proof in the dyadic case is very similar, but is simplified considerably by the fact that if two dyadic cubes intersect then one is contained in the other.  $\square$

*Proof of Theorem 1.2.* By a standard argument, we may assume that  $f \in C^\infty(\mathbb{R}^n)$  and has compact support. Fix  $p$ ,  $1 < p < \infty$ ; then  $Tf \in L^q$ , where  $q > 1$  is such that  $p \geq q/r'$ . Hence, we may apply Theorem 3.4 to it. Fix  $\delta < 1$ . Then there exists  $\epsilon > 0$  such that for each  $t > 0$  there exists a sequence of disjoint dyadic cubes  $\{Q_j^t\}$  such that

$$\left( \frac{1}{|Q_j^t|} \int_{Q_j^t} ||Tf|^\delta - (|Tf|^\delta)_{Q_j^t}| dx \right)^{1/\delta} > \epsilon^{1/\delta} t$$

and

$$\begin{aligned}
\sup_{t>0} t^p u(\{x \in \mathbb{R}^n : |Tf(x)| > t\}) &\leq \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : M_\delta^d(Tf)(x) > t\}) \\
&\leq C \sup_{t>0} t^p \sum_j A_u^r(Q_j^t).
\end{aligned}$$

As we noted in the Introduction,  $T$  satisfies inequality (1.9). Therefore, for each  $j$ ,

$$Q_j^t \subset \{x \in \mathbb{R}^n : M_\delta^\#(Tf)(x) > \epsilon^{1/\delta}t\} \subset \{x \in \mathbb{R}^n : Mf(x) > \beta t\},$$

where  $\beta = C_\delta^{-1}\epsilon^{1/\delta}$ .

By Lemma 5.1 (with  $\mu = 1$ ), for each  $t > 0$  there exists a sequence of disjoint dyadic cubes  $\{P_k^t\}$  such that for each  $j$ ,  $Q_j^t \subset 3P_k^t$  for some  $k$ , and such that

$$\frac{1}{|P_k^t|} \int_{P_k^t} |f| dx > \rho t,$$

where  $\rho > 0$  depends only on  $\beta$  and  $n$ . Then by Lemma 3.2, Condition (3), for each  $t > 0$ ,

$$\begin{aligned} t^p \sum_j A_u^r(Q_j^t) &= t^p \sum_k \sum_{Q_j^t \subset 3P_k^t} A_u^r(Q_j^t) \\ &\leq t^p \sum_k A_u^r(3P_k^t) \\ &\leq \rho^{-p} \sum_k |3P_k^t| \left( \frac{1}{|3P_k^t|} \int_{3P_k^t} u^r dx \right)^{1/r} \left( \frac{1}{|P_k^t|} \int_{P_k^t} |f| dx \right)^p. \end{aligned}$$

By Hölder's inequality and inequality (1.7),

$$\begin{aligned} &\leq C \sum_k \left( \frac{1}{|3P_k^t|} \int_{3P_k^t} u^r dx \right)^{1/r} \left( \frac{1}{|3P_k^t|} \int_{3P_k^t} v^{-p'/p} dx \right)^{p/p'} \int_{P_k^t} |f|^p v dx \\ &\leq C \sum_k \int_{P_k^t} |f|^p v dx \\ &\leq C \int_{\mathbb{R}^n} |f|^p v dx. \end{aligned}$$

The constant is independent of  $t$ , so if we take the supremum over all  $t > 0$  we get inequality (1.8). This completes our proof.  $\square$

## 6. COMMUTATORS

In this section we prove Theorem 1.6. The proof depends on Theorem 1.2 and the following analogue of inequality (1.9) for commutators.

**Lemma 6.1.** *Given a Calderón-Zygmund operator  $T$ , a function  $b$  in BMO, constants  $\delta_0$  and  $\delta_1$ ,  $0 < \delta_0 < \delta_1 < 1$ , and  $k \geq 1$ , there exists a constant  $K$ , depending on the BMO norm of  $b$ , such that for every function  $f \in C_0^\infty(\mathbb{R}^n)$  and any  $x \in \mathbb{R}^n$ ,*

$$M_{\delta_0}^{\#,d}(C_b^k f)(x) \leq K \sum_{i=0}^{k-1} M_{\delta_1}^d(C_b^i f)(x) + K M^{k+1} f(x).$$

This result is found in [23, 24]. As given there, the non-dyadic maximal operator appears in the first term on the right-hand side, but it is immediate from the proof that it is still true with the dyadic maximal operator there.

*Proof of Theorem 1.6.* When  $k = 0$ , Theorem 1.6 reduces to Theorem 1.2, so we may fix  $k \geq 1$ . By a standard argument we may assume that  $f \in C^\infty(\mathbb{R}^n)$  and has compact support. Fix  $p$ ,  $1 < p < \infty$ ; then  $C_b^i f \in L^q$ , where  $0 \leq i \leq k$  and  $q > 1$  is such that  $p \geq q/r'$ . Hence, we may apply Theorem 3.4 to  $C_b^k f$ . Fix  $\delta_0$  and  $\delta_1$ ,  $0 < \delta_0 < \delta_1 < 1$ . There there exists  $\epsilon > 0$  such that for each  $t > 0$  there exists a sequence of disjoint dyadic cubes  $\{Q_j^t\}$  such that

$$\left( \frac{1}{|Q_j^t|} \int_{Q_j^t} \|C_b^k f\|^{\delta_0} - (|C_b^k f|^{\delta_0})_{Q_j^t} dx \right)^{1/\delta_0} > \epsilon^{1/\delta_0} t$$

and

$$\begin{aligned} \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : |C_b^k f(x)| > t\}) &\leq \sup_{t>0} t^p u(\{x \in \mathbb{R}^n : M_{\delta_0}^d(C_b^k f)(x) > t\}) \\ &\leq C \sup_{t>0} t^p \sum_j A_u^r(Q_j^t). \end{aligned}$$

By Lemma 6.1, for each  $j$  and  $t$ ,

$$\begin{aligned} Q_j^t &\subset \bigcup_{i=1}^{k-1} \{x \in \mathbb{R}^n : M_{\delta_1}^d(C_b^i f)(x) > \beta t\} \\ &\quad \cup \{x \in \mathbb{R}^n : M_{\delta_1}^d(Tf)(x) > \beta t\} \\ &\quad \cup \{x \in \mathbb{R}^n : M^{k+1} f(x) > \beta t\} \\ &\equiv \left( \bigcup_{i=1}^{k-1} F_i^{\beta t} \right) \cup F_0^{\beta t} \cup F_k^{\beta t}, \end{aligned}$$

where  $\beta = \epsilon^{1/\delta_0} K^{-1}(k+1)^{-1}$ . For each  $j$  and  $t$  we cannot have that  $|Q_j^t \cap F_i^{\beta t}| < (k+1)^{-1}|Q_j^t|$  for all  $i$ . Hence, for some  $i$ ,  $|Q_j^t \cap F_i^{\beta t}| \geq (k+1)^{-1}|Q_j^t|$ ; if this is the

case we write  $Q_j^t \in \mathcal{F}_i^{\beta t}$ . Thus,

$$\sup_{t>0} t^p \sum_j A_u^r(Q_j^t) \leq \sum_{i=0}^k \sup_{t>0} t^p \sum_{Q_j^t \in \mathcal{F}_i^{\beta t}} A_u^r(Q_j^t).$$

To complete the proof we will show that each term of the outer sum on the right-hand side is dominated by  $C \int_{\mathbb{R}^n} |f|^p v dx$ . There are three cases.

*Case 1: cubes in  $\mathcal{F}_k^{\beta t}$ .* As we noted in Section 2, there exists a constant  $\beta' > 0$  such that

$$\{x \in \mathbb{R}^n : M^{k+1} f(x) > \beta t\} \subset \{x \in \mathbb{R}^n : M_B f(x) > \beta' t\},$$

where  $B(t) = t \log(e+t)^k$ . Therefore, by Lemma 5.1 (with  $\mu = (k+1)^{-1}$ ), there exists a constant  $\nu > 0$  such that, for each  $t > 0$  there exists a collection of disjoint dyadic cubes  $\{P_l^t\}$  such that for each  $j$ ,  $Q_j^t \subset 3P_l^t$  for some  $l$  and such that  $\|f\|_{B, P_l^t} > \nu t$ . We now proceed exactly as we did at the end of the proof of Theorem 1.2. Since  $C_k(t) = t^{p'} \log(e+t)^{kp'}$ ,  $C_k^{-1}(t) \approx t^{1/p'} \log(e+t)^{-k}$ , and so  $t^{1/p} C_k^{-1}(t) \leq B^{-1}(t)$ . Then, by Lemma 3.2, Conditions (2) and (3), the generalized Hölder's inequality (2.3) and inequality (1.11),

$$\begin{aligned} \sup_{t>0} t^p \sum_{Q_j^t \in \mathcal{F}_k^{\beta t}} A_u^r(Q_j^t) &\leq \sup_{t>0} t^p \sum_l A_u^r(3P_l^t) \\ &\leq C \sup_{t>0} \sum_l |3P_l^t| \left( \frac{1}{|3P_l^t|} \int_{3P_l^t} u^r dx \right)^{1/r} \|f\|_{B, P_l^t}^p \\ &\leq C \sup_{t>0} \sum_l \left( \frac{1}{|3P_l^t|} \int_{3P_l^t} u^r dx \right)^{1/r} \|v^{-1/p}\|_{C_k, P_l^t}^p \int_{P_l^t} |f|^p v dx \\ &\leq C \sup_{t>0} \sum_l \left( \frac{1}{|3P_l^t|} \int_{3P_l^t} u^r dx \right)^{1/r} \|v^{-1/p}\|_{C_k, 3P_l^t}^p \int_{P_l^t} |f|^p v dx \\ &\leq C \sup_{t>0} \sum_l \int_{P_l^t} |f|^p v dx \\ &\leq C \int_{\mathbb{R}^n} |f|^p v dx. \end{aligned}$$

*Case 2: cubes in  $\mathcal{F}_0^{\beta t}$ .* Given  $t > 0$ , let  $s = (\beta t)^{\delta_1}$ . Again by Lemma 5.1 (the dyadic case), if  $Q_j^t \in \mathcal{F}_0^{\beta t}$  then for some  $i$ ,  $Q_j^t \subset C_i^s$ , where  $\{C_i^s\}$  is the Calderón-Zygmund

decomposition of  $|Tf|^{\delta_1}$  at height  $s$ . Hence, by Lemma 3.2, Conditions (2) and (3),

$$\sup_{t>0} t^p \sum_{Q_j^t \in \mathcal{F}_0^{\beta t}} A_u^r(Q_j^t) \leq \sup_{t>0} t^p \sum_i A_u^r(C_i^s).$$

By Corollary 3.5 there exists  $\epsilon > 0$  and a subcollection  $\{\bar{Q}_j^t\}$  of  $\{C_i^s\}$  such that if  $x \in \bar{Q}_j^t$  then  $M_{\delta_1}^{\#,d}(Tf)(x) > \beta''t$ , where  $\beta'' = \epsilon^{1/\delta_1}\beta$ , and such that

$$\sup_{t>0} t^p \sum_i A_u^r(C_i^s) \leq C \sup_{t>0} t^p \sum_j A_u^r(\bar{Q}_j^t).$$

We can now argue exactly as we did in the proof of Theorem 1.2 to get

$$\sup_{t>0} t^p \sum_{Q_j^t \in \mathcal{F}_0^{\beta t}} A_u^r(Q_j^t) \leq C \int_{\mathbb{R}^n} |f|^p v \, dx.$$

*Case 3: cubes in  $\mathcal{F}_i^{\beta t}$ ,  $1 \leq i \leq k-1$ .* Fix  $i$ ; then arguing exactly as we did in Case 2, by Corollary 3.5 there exists  $\epsilon > 0$  and a collection of disjoint dyadic cubes  $\{\bar{Q}_j^t\}$  such that if  $x \in \bar{Q}_j^t$  then  $M_{\delta_1}^{\#}(C_b^i f)(x) > \beta''t$ , where  $\beta'' = \epsilon^{1/\delta_1}\beta$ , and such that

$$\sup_{t>0} t^p \sum_{Q_j^t \in \mathcal{F}_i^{\beta t}} A_u^r(Q_j^t) \leq C \sup_{t>0} t^p \sum_j A_u^r(\bar{Q}_j^t).$$

We now apply Lemma 6.1 and repeat the argument at the beginning of this proof. When we do so we reduce the degree of the highest order commutator appearing from  $i$  to  $i-1$ . Therefore, after repeating our argument a finite number of times, we will reduce to collections of cubes satisfying conditions such as those in Case 1 and Case 2. Repeating those arguments will then give us the desired inequality.  $\square$

## REFERENCES

- [1] D.R. Adams, *A note on Riesz potentials*, Duke Math. J. 42 (1975), 765-778.
- [2] J. Alvarez, R. Bagby, D. Kurtz and C. Pérez, *Weighted estimates for commutators of linear operators*, Studia Math. 104 (1993), 195-209.
- [3] J. Alvarez and C. Pérez, *Estimates with  $A_\infty$  weights for various singular integral operators*, Bollettino U.M.I. (7) 8-A (1994), 123-133.
- [4] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, New York, 1988.
- [5] M. Carozza and A. Passarelli di Napoli, *Composition of maximal operators*, Publ. Mat. 40 (1996), 397-409.
- [6] M. Christ, *Lectures on Singular Integral Operators*, CBMS Regional Conference Series in Mathematics, 77, Amer. Math. Soc., Providence, 1990.
- [7] R. Coifman, *Distribution function inequalities for singular integrals*, Proc. Acad. Sci. U.S.A. 69 (1972), 2838-2839.

- [8] R. Coifman and Y. Meyer, *Au delà des opérateurs pseudo-différentiels*, Astérisque 57 (1979), 1-84.
- [9] R. Coifman, R. Rochberg and G. Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math. 103 (1976), 611-635.
- [10] D. Cruz-Uribe, SFO, and A. Fiorenza, *The  $A_\infty$  property for Young functions and weighted norm inequalities*, preprint.
- [11] D. Cruz-Uribe, SFO, and C. Pérez, *Sharp two-weight, weak-type norm inequalities for singular integral operators*, Math. Res. Let. 6 (1999), 417-428.
- [12] C. Fefferman, *Inequalities for strongly singular convolution operators*, Acta Math. 124 (1970), 9-36.
- [13] C. Fefferman and E.M. Stein,  *$H^p$  spaces in several variables*, Acta Math. 129 (1972), 137-193.
- [14] J. García-Cuerva and J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North Holland Math. Studies 116, North Holland, Amsterdam, 1985.
- [15] L. Hörmander, *Pseudo-differential operators and hypoelliptic equations*, Proc. Symp. Pure Math 10 (1967), 138-183.
- [16] J.-L. Journé, *Zygmund Operators, Pseudo-Differential Operators and the Cauchy Integral of Calderón*, Lecture Notes in Mathematics, 994, Springer Verlag, New York, 1983.
- [17] M.A. Krasnosel'skiĭ and Ya.B. Rutickiĭ, *Convex functions and Orlicz spaces*, P. Noordhoff, Groningen, 1961.
- [18] C.J. Neugebauer, *Inserting  $A_p$ -weights*, Proc. Amer. Math. Soc. 87 (1983), 644-648.
- [19] R. O'Neil, *Fractional integration in Orlicz spaces*, Trans. Amer. Math. Soc. 115 (1963), 300-328.
- [20] C. Pérez, *Weighted norm inequalities for singular integral operators*, J. London Math. Soc. 49 (1994), 296-308.
- [21] C. Pérez, *Two weighted inequalities for potential and fractional type maximal operators*, Indiana Math. J. 43 (1994), 663-683.
- [22] C. Pérez, *On sufficient conditions for the boundedness of the Hardy-Littlewood maximal operator between weighted  $L^p$ -spaces with different weights*, Proc. London Math. Soc. 71 (1995), 135-57.
- [23] C. Pérez, *Endpoint estimates for commutators of singular integral operators*, J. Func. Anal. 128 (1995), 163-185.
- [24] C. Pérez, *Sharp estimates for commutators of singular integrals via iterations of the Hardy-Littlewood maximal function*, J. Fourier Anal. Appl. 3 (1997), 743-756.
- [25] D.H. Phong and E.M. Stein, *Hilbert integrals, singular integrals and Radon transforms*, Acta Math. 157 (1985), 99-157.
- [26] M.M. Rao and Z.D. Ren, *Theory of Orlicz Spaces*, Marcel Dekker, New York, 1991.
- [27] E.T. Sawyer, *A two weight weak type inequality for fractional integrals*, Trans. Amer. Math. Soc. 281 (1984), 339-345.
- [28] E.T. Sawyer and R. Wheeden, *Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces*, Amer. J. Math. 114 (1992), 813-874.
- [29] C. Segovia and J.L. Torrea, *Weighted inequalities for commutators of fractional and singular integrals*, Publ. Mat. 35 (1991), 209-235.
- [30] E.M. Stein, *Note on the class  $L \log L$* , Studia Math. 32 (1969), 305-310.
- [31] E.M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, 1970.

DEPT. OF MATHEMATICS, TRINITY COLLEGE, HARTFORD, CT 06106-3100, USA

*E-mail address:* david.cruzuribe@mail.trincoll.edu

DEP. DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN

*E-mail address:* carlos.perez@uam.es