

Pacific Journal of Mathematics

UNBOUNDED REPRESENTATIONS OF *-ALGEBRAS

STANLEY P. GUDDER AND W. SCRUGGS

UNBOUNDED REPRESENTATIONS OF *-ALGEBRAS

S. GUDDER AND W. SCRUGGS

Basic results on unbounded operator algebras are given, a general class of representations, called adjointable representations is introduced and irreducibility of representations is considered. A characterization of self-adjointness for closed, strongly cyclic *-representations is presented.

1. Introduction. Algebras of unbounded operators and unbounded representations of *-algebras have been important in quantum field theory [1, 3, 9, 10] and certain studies of Lie algebras [5, 7]. The present paper proceeds along the lines initiated and developed by Robert Powers [6, 7] and much of the notation and definitions follow [6]. In §2, we present some basic results concerning unbounded operator algebras, introduce a class of representations called adjointable representations, and consider irreducibility of representations. Section 3 characterizes the self-adjointness of closed, strongly cyclic *-representations.

2. Adjointable representations. Let M and N be subspaces (linear manifolds) in a Hilbert space H . Let $L(M, N)$ and $L_c(M, N)$ denote the collection of linear operators and closable linear operators, respectively with domain M and range in N . For simplicity we use the notation $L(M) = L(M, M)$ and $L_c(M) = L_c(M, M)$. Notice that $L_c(H)$ is the set of bounded linear operators on H . We denote the domain of an operator A by $D(A)$ and if A is closable we denote the closure of A by \bar{A} . A collection of operators \mathcal{B} is an *op-algebra* if there exists a subspace M such that $\mathcal{B} \subseteq L(M)$ and $A, B \in \mathcal{B}$ implies $AB, (\alpha A + B) \in \mathcal{B}$ for all $\alpha \in \mathbb{C}$. A set $\mathcal{B} \subseteq L(M)$ is *symmetric* if M is dense and $A \in \mathcal{B}$ implies $D(A^*) \supseteq M$ and $A^*|M \in \mathcal{B}$. A symmetric op-algebra $\mathcal{B} \subseteq L(M)$ that contains $I|M$ is called an *op*-algebra*. It is easy to see that if $\mathcal{B} \subseteq L(M)$ is an op*-algebra, then the map $A \rightarrow A^*|M$ is an involution so \mathcal{B} is a *-algebra. Also, if π is a representation of a *-algebra \mathcal{A} , then $\pi(\mathcal{A}) = \{\pi(A): A \in \mathcal{A}\}$ is an op-algebra and if π is a *-representation of \mathcal{A} , then $\pi(\mathcal{A})$ is an op*-algebra (we always assume that a *-algebra contains an identity I).

A set $\mathcal{B} \subseteq L(M, N)$ is *directed* if for any $B_1, B_2 \in \mathcal{B}$ there exists a $B_3 \in \mathcal{B}$ such that $\|B_1x\|, \|B_2x\| \leq \|B_3x\|$ for all $x \in M$. For example, if $\mathcal{B} \subseteq L_c(H)$ and $\{\lambda I: \lambda \geq 0\} \subseteq \mathcal{B}$, then \mathcal{B} is directed. Indeed, just let $B_3 = (\|B_1\| + \|B_2\|)I$. For an example of an unbounded directed set, let $\mathcal{B} \subseteq L(M, H)$ and suppose $B_1, B_2 \in \mathcal{B}$ implies $B_3 =$

$I \mid M + B_1^* B_1 + B_2^* B_2 \in \mathcal{B}$. Then for any $x \in M$ we have

$$\begin{aligned} \|B_3 x\|^2 &= \|x\|^2 + \|(B_1^* B_1 + B_2^* B_2)x\|^2 + 2\langle(B_1^* B_1 + B_2^* B_2)x, x\rangle \\ &\geq 2(\|B_1 x\|^2 + \|B_2 x\|^2) \\ &\geq \|B_1 x\|^2, \|B_2 x\|^2. \end{aligned}$$

In particular, any op *-algebra is directed.

An *extension* \mathcal{B}_1 of $\mathcal{B} \subseteq L(M, N)$ is a set of operators $\mathcal{B}_1 \subseteq L(M_1, N_1)$ where $M \subseteq M_1$, $N \subseteq N_1$ and for which there exists a bijection $\phi: \mathcal{B} \rightarrow \mathcal{B}_1$ such that $\phi(B)|M = B$ for every $B \in \mathcal{B}$. If $\mathcal{B} \subseteq L(M, N)$, the \mathcal{B} -topology on M is the topology generated by the set of seminorms $\{\|x\|, \|Bx\|: B \in \mathcal{B}\}$. The completion of M in the \mathcal{B} -topology is denoted by $\hat{M}_{\mathcal{B}}$ or simply \hat{M} if no confusion can arise. We say that $\mathcal{B} \subseteq L(M, N)$ is *collectively closed* if for any net $x_\alpha \in M$ satisfying $x_\alpha \rightarrow x \in H$, $Bx_\alpha \rightarrow y(B) \in H$ for every $B \in \mathcal{B}$, then $x \in M$ and $Bx = y(B)$. Clearly if all $B \in \mathcal{B}$ are closed then \mathcal{B} is collectively closed; the converse need not hold.

THEOREM 1.

- (1) $\mathcal{B} \subseteq L(M, N)$ is collectively closed if and only if $M = \hat{M}_{\mathcal{B}}$.
- (2) If $\mathcal{B} \subseteq L_c(M, N)$, then the set $\mathcal{B}_1 = \{\bar{B} \mid M_1: B \in \mathcal{B}\}$ where $M_1 = \cap \{D(\bar{B}): B \in \mathcal{B}\}$ is collectively closed.
- (3) If $\mathcal{B} \subseteq L_c(M, N)$, then the set $\bar{\mathcal{B}} = \{\bar{B} \mid \hat{M}_{\mathcal{B}}: B \in \mathcal{B}\}$ is the minimal collectively closed extension of \mathcal{B} . Moreover, if $\mathcal{B} \subseteq L_c(M)$ and $A, B \in \mathcal{B}$ implies $AB \in \mathcal{B}$, then $\bar{\mathcal{B}} \subseteq L_c(\hat{M}_{\mathcal{B}})$.
- (4) If $\mathcal{B} \subseteq L_c(M, N)$ is directed, then $\hat{M}_{\mathcal{B}} = \cap \{D(\bar{B}): B \in \mathcal{B}\}$.
- (5) If $\mathcal{B} \subseteq L_c(M)$ is an op-algebra, then $\bar{\mathcal{B}}$ is an op-algebra. If $\mathcal{B} \subseteq L_c(M)$ is an op *-algebra, then $\bar{\mathcal{B}}$ is an op *-algebra and $\hat{M}_{\mathcal{B}} = \cap \{D(\bar{B}): B \in \mathcal{B}\}$.

Proof.

- (1) Suppose $\mathcal{B} \subseteq L(M, N)$ is collectively closed and $x_\alpha \in M$ is a Cauchy net in the \mathcal{B} -topology. Then x_α and Bx_α are Cauchy in H so there exist $x, y(B) \in H$ such that $x_\alpha \rightarrow x$, $Bx_\alpha \rightarrow y(B)$ in H for every $B \in \mathcal{B}$. Since \mathcal{B} is collectively closed, $x \in M$ and $Bx_\alpha \rightarrow Bx$, so $x_\alpha \rightarrow x$ in the \mathcal{B} -topology and M is complete in the \mathcal{B} -topology. Hence $M = \hat{M}_{\mathcal{B}}$. Conversely, suppose $M = \hat{M}_{\mathcal{B}}$, and x_α is a net in M such that $x_\alpha \rightarrow x$ and $Bx_\alpha \rightarrow y(B)$ in H for every $B \in \mathcal{B}$. Then x_α is Cauchy in the \mathcal{B} -topology. Since M is complete in the \mathcal{B} -topology there exists an $x' \in M$ such that $x_\alpha \rightarrow x'$ and $Bx_\alpha \rightarrow Bx'$ in H for every $B \in \mathcal{B}$. Hence $x = x' \in M$ and $Bx = Bx' = y(B)$.

(2) This is straightforward.

(3) It is clear that

$\hat{M}_{\mathcal{B}} = \{x \in \cap \{D(\bar{B}): B \in \mathcal{B}\}: M \ni x_\alpha \rightarrow x, Bx_\alpha \rightarrow \bar{B}x \text{ for all } B \in \mathcal{B}\}$. We now show that $\hat{M}_{\mathcal{B}}$ is complete in the $\bar{\mathcal{B}} \mid \hat{M}_{\mathcal{B}}$ -topology. Suppose $x_\alpha \in \hat{M}_{\mathcal{B}}$ is Cauchy in the $\bar{\mathcal{B}} \mid \hat{M}_{\mathcal{B}}$ -topology. Then x_α and Bx_α are Cauchy in H for every $B \in \mathcal{B}$. Hence there exists an $x \in H$ such that $x_\alpha \rightarrow x$ and $Bx_\alpha \rightarrow \bar{B}x$ for every $B \in \mathcal{B}$. Since $x_\alpha \in \hat{M}_{\mathcal{B}}$ there exists a net $x_{\alpha\beta} \in M$ such that $x_{\alpha\beta} \rightarrow x_\alpha$ and $Bx_{\alpha\beta} \rightarrow \bar{B}x_\alpha$ in H for every $B \in \mathcal{B}$. Now $x_{\alpha\beta}$ is a net in M and $x_{\alpha\beta} \rightarrow x, Bx_{\alpha\beta} \rightarrow \bar{B}x$ in H for every $B \in \mathcal{B}$. Hence $x \in \hat{M}_{\mathcal{B}}$. It follows from (1) that $\bar{\mathcal{B}}$ is collectively closed. Clearly $\bar{\mathcal{B}}$ is an extension of \mathcal{B} . Moreover, $\bar{\mathcal{B}}$ is a minimal collectively closed extension since any collectively closed extension of \mathcal{B} must contain $\hat{M}_{\mathcal{B}}$ in its domain. Now suppose $\mathcal{B} \subseteq L_c(M)$ and $A, B \in \mathcal{B}$. If $x \in \hat{M}_{\mathcal{B}}$, then there exists a net $x_\alpha \in M$ such that $x_\alpha \rightarrow x$ and $Bx_\alpha \rightarrow \bar{B}x$ for every $B \in \mathcal{B}$. For fixed $A \in \mathcal{B}$ we have $Ax_\alpha \in M$ and $Ax_\alpha \rightarrow \bar{A}x$ and for every $B \in \mathcal{B}$, since $BA \in \mathcal{B}$, $BAx_\alpha \rightarrow \bar{BA}x = \bar{B}\bar{A}x$. Hence $\bar{A}x \in \hat{M}_{\mathcal{B}}$ and $\bar{\mathcal{B}} \subseteq L_c(\hat{M}_{\mathcal{B}})$.

(4) Suppose that $\mathcal{B} \subseteq L_c(M, N)$ is directed. We have seen that $\hat{M}_{\mathcal{B}} \subseteq \cap \{D(B): B \in \mathcal{B}\}$. If $x \in \cap \{D(\bar{B}): B \in \mathcal{B}\}$, then for each $B \in \mathcal{B}$ there exists a sequence $x(B, i) \in M$ such that $x(B, i) \rightarrow x$ and $Bx(B, i) \rightarrow \bar{B}x$. For each $B \in \mathcal{B}$ and for each integer $n > 0$ there exists an integer $n_B > 0$ such that $\|x(B, n_B) - x\| < n^{-1}$ and $\|Bx(B, n_B) - \bar{B}x\| < n^{-1}$. For $A, B \in \mathcal{B}$, define the order $(A, n_A) < (B, m_B)$ if $\|Az\| \leq \|Bz\|$ for every $z \in M$ and $n < m$. Since \mathcal{B} is directed, $\{(B, n_B)\}$ is a directed partially ordered set and $x(B, n_B)$ is a net. Notice that if $\|Az\| \leq \|Bz\|$ for every $z \in M$ then $\|\bar{A}y\| \leq \|\bar{B}y\|$ for every $y \in \cap \{D(\bar{B}): B \in \mathcal{B}\}$. Indeed let $z_i \in M$ be a sequence such that $z_i \rightarrow y$ and $Bz_i \rightarrow \bar{B}y$. Since $\|Az_i - Az_j\| \leq \|Bz_i - Bz_j\|$, Az_i is Cauchy and hence $Az_i \rightarrow \bar{A}y$. Therefore,

$$\|\bar{A}y\| = \lim \|Az_i\| \leq \lim \|Bz_i\| = \|\bar{B}y\|.$$

Clearly, $x(B, m_B) \rightarrow x$ and to show that $Ax(B, m_B) \rightarrow \bar{A}x$ let $\epsilon > 0$ and let $n > 0$ be an integer such that $n^{-1} < \epsilon$. Then for $(B, m_B) > (A, n_A)$ we have

$$\begin{aligned} \|Ax(B, m_B) - \bar{A}x\| &= \|\bar{A}x(B, m_B) - \bar{A}x\| \\ &\leq \|\bar{B}x(B, m_B) - \bar{B}x\| \\ &< m^{-1} < n^{-1} < \epsilon. \end{aligned}$$

It follows that $x \in \hat{M}_{\mathcal{B}}$.

(5) This is a straightforward consequence of (2) and (3).

In the work of R. Powers [6] only hermitian representations are considered. But there are important representations that are not hermitian. For example, even if π is hermitian, π^* need not be. We therefore treat a larger class of representations, which we call adjointable, that includes π^* whenever π is hermitian.

Let \mathcal{A} be a $*$ -algebra and let π, π_1 be two representations of \mathcal{A} with domains $D(\pi), D(\pi_1) \subseteq H$. We say that π and π_1 are *adjoint* and write $\pi a \pi_1$, if $\langle \pi(A)x, y \rangle = \langle x, \pi_1(A^*)y \rangle$ for every $A \in \mathcal{A}$ and $x \in D(\pi)$, $y \in D(\pi_1)$. Notice that a is a symmetric relation; that is $\pi a \pi_1$ if and only if $\pi_1 a \pi$. Also, $\pi a \pi$ if and only if π is hermitian. Furthermore, if $\pi a \pi_1$ and $\pi_1 a \pi_2$ then $\pi(A) = \pi_2(A)$ on $D(\pi) \cap D(\pi_2)$ for every $A \in \mathcal{A}$ and if $D(\pi) = D(\pi_2)$ then $\pi = \pi_2$. We say that a representation π is *adjointable* if there exists a representation π_1 such that $\pi a \pi_1$.

If π is a representation of a $*$ -algebra \mathcal{A} , we define $D(\pi^*) = \cap\{D(\pi(A)^*): A \in \mathcal{A}\}$ and $\pi^*(A) = \pi(A^*)^*|D(\pi^*)$ for all $A \in \mathcal{A}$. (To save parentheses we use the notation $\pi(A)^* = [\pi(A)]^*$.) In general, π^* need not be a representation since, for one thing, $D(\pi^*)$ need not be dense. If π is hermitian, then π^* is a representation [6]. Hence, if π is hermitian, then

$$\langle \pi(A)x, y \rangle = \langle x, \pi(A)^*y \rangle = \langle x, \pi^*(A^*)y \rangle$$

for every $A \in \mathcal{A}$ and $x \in D(\pi)$, $y \in D(\pi^*)$ so $\pi a \pi^*$ and each is adjointable.

THEOREM 2.

- (1) π is adjointable if and only if $D(\pi^*)$ is dense.
- (2) If π is adjointable, then π^* is a closed representation and is the largest representation adjoint to π .
- (3) Suppose $\pi \subset \pi_1$. If $\pi_1 a \pi_2$, then $\pi a \pi_2$. If π_1 is adjointable, then so is π and $\pi_1^* \subset \pi^*$.
- (4) If π is adjointable, then there exists a smallest closed representation $\bar{\pi}$ which extends π . If $\pi a \pi_1$, then $\bar{\pi} a \pi_1$.
- (5) If π is adjointable, then $\pi^*, \bar{\pi}$ are adjointable, π^{**} is a closed representation and $\pi \subset \bar{\pi} \subset \pi^{**}$, $\pi^{***} = \pi^*$, $\bar{\pi}^* = \pi^*$.
- (6) If π is hermitian and π_1 is an hermitian extension of π , then $\pi \subset \pi_1 \subset \pi^*$.
- (7) If π is hermitian, then π^{**} and $\bar{\pi}$ are hermitian and $\pi \subset \bar{\pi} \subset \pi^{**} \subset \pi^*$.

Proof.

- (1) If π is adjointable and $\pi a \pi_1$ then $\pi_1(A^*) \subset \pi(A)^*$ for every $A \in \mathcal{A}$ so $D(\pi_1) \subseteq D(\pi^*)$ and $D(\pi^*)$ is dense. Conversely, suppose

$D(\pi^*)$ is dense. For $x \in D(\pi)$, $y \in D(\pi^*)$ we have

$$\begin{aligned}\langle \pi(A^*)x, \pi^*(B)y \rangle &= \langle \pi(A^*)x, \pi(B^*)^*y \rangle \\ &= \langle \pi(B^*)\pi(A^*)x, y \rangle \\ &= \langle \pi(B^*A^*)x, y \rangle \\ &= \langle x, \pi(B^*A^*)^*y \rangle.\end{aligned}$$

Hence $\pi^*(B)y \in D(\pi(A^*)^*)$ and $\pi(A^*)^*\pi^*(B)y = \pi(B^*A^*)^*y$ for every $A, B \in \mathcal{A}$. It follows that $\pi^*(B): D(\pi^*) \rightarrow D(\pi^*)$ and $\pi^*(A)\pi^*(B) = \pi((AB)^*)^* = \pi^*(AB)$. Moreover, π^* is linear since for $x \in D(\pi)$, $y \in D(\pi^*)$ we have

$$\begin{aligned}\langle \pi^*(\alpha A + B)y, x \rangle &= \langle \pi(\bar{\alpha}A^* + B^*)^*y, x \rangle \\ &= \langle y, \bar{\alpha}\pi(A^*)x \rangle + \langle y, \pi(B^*)x \rangle \\ &= \langle [\alpha\pi^*(A) + \pi^*(B)]y, x \rangle.\end{aligned}$$

It follows that π^* is a representation and $\pi a \pi^*$.

(2) It was shown in (1) that π^* is a representation if π is adjointable. It follows from Theorem 1 (2) that π^* is closed. If $\pi a \pi_1$ then $\langle \pi(A)x, y \rangle = \langle x, \pi_1(A^*)y \rangle$ for all $x \in D(\pi)$, $y \in D(\pi_1)$. Hence, $D(\pi_1) \subseteq D(\pi^*)$ and $\pi_1(A^*) \subset \pi(A)^* = \pi^*(A^*)$ for every $A \in \mathcal{A}$ so $\pi_1 \subset \pi^*$.

(3) Suppose $\pi \subset \pi_1$ and $\pi_1 a \pi_2$. Then for every $x \in D(\pi)$, $y \in D(\pi_2)$ we have $\langle \pi(A)x, y \rangle = \langle \pi_1(A)x, y \rangle = \langle x, \pi_2(A^*)y \rangle$. Hence $\pi a \pi_2$. For all $x \in D(\pi)$, $y \in D(\pi_1^*)$ we have $\langle \pi(A)x, y \rangle = \langle x, \pi_1^*(A^*)y \rangle$. Hence $\pi a \pi_1^*$ and by (2) we have $\pi_1^* \subset \pi^*$.

(4) If π is adjointable, then by (1), $D(\pi^*)$ is dense. Then $D(\pi(A)^*)$ is dense so $\pi(A)$ is closable for every $A \in \mathcal{A}$. Define $D(\bar{\pi}) = \bar{D}(\pi)_B$ where $B = \{\pi(A): A \in \mathcal{A}\}$ and $\bar{\pi}(A) = \pi(A)|D(\bar{\pi})$. It follows from Theorem 1 (3) that $\{\bar{\pi}(A): A \in \mathcal{A}\}$ is the minimal collectively closed extension of B . It is straightforward to show that $\bar{\pi}$ is a representation and that $\pi a \pi_1$ implies $\bar{\pi} a \pi_1$.

(5) If π is adjointable then so is π^* and from (2) π^{**} is a closed representation. If $x \in D(\pi)$, $y \in D(\pi^*)$ then for all $A \in \mathcal{A}$ we have $\langle \pi^*(A^*)y, x \rangle = \langle \pi(A)^*y, x \rangle = \langle y, \pi(A)x \rangle$. Hence $x \in \cap\{D[\pi^*(A^*)]: A \in \mathcal{A}\} = D(\pi^{**})$ and $\pi^{**}(A)x = \pi^*(A^*)^*x = \pi(A)x$ so $\pi \subset \pi^{**}$. Since $\pi \subset \bar{\pi}$ we have by (3) that $\bar{\pi}^* \subset \pi^*$. Since $\pi a \pi^*$ from (4) we have $\bar{\pi} a \pi^*$. Hence by (2) $\pi^* \subset \bar{\pi}^*$ so $\pi^* = \bar{\pi}^*$. By (3) $\pi^{***} \subset \pi^*$. Since $\pi^* a \pi^{**}$, by (2) we have $\pi^* \subset \pi^{***}$ so $\pi^{***} = \pi^*$.

(6) For all $x \in D(\pi)$, $y \in D(\pi_1)$, $A \in \mathcal{A}$ we have

$$\langle \pi(A)x, y \rangle = \langle \pi_1(A)x, y \rangle = \langle x, \pi_1(A^*)y \rangle.$$

Hence $\pi_1 \alpha \pi$ and by (2) $\pi_1 \subset \pi^*$.

(7) It is shown in [6] that $\bar{\pi}$ is hermitian if π is hermitian. Since π is hermitian we have $\pi \subset \pi^*$. Applying (3) twice gives $\pi^{**} \subset \pi^{***}$ so π^{**} is hermitian. Since π^{**} is closed we have from (2) that $\pi \subset \bar{\pi} \subset \pi^{**}$ and from (6) $\pi^{**} \subset \pi^*$.

We now show that the extensions in (7) can be distinct. Let \mathcal{A} be the free commutative $*$ -algebra on one hermitian generator A . Define the representation π of \mathcal{A} on the Hilbert space $H = L^2[0, 1]$ as follows: $D(\pi) = \{f \in C^\infty[0, 1]: f^{(n)}(0) = f^{(n)}(1) = 0, n = 0, 1, 2, \dots\}$ $\pi(A) = -i d/dt$. It is straightforward to show that π is hermitian and that $\pi = \bar{\pi} = \pi^{**} \subsetneq \pi^*$ [8]. Now let π_1 be the representation of \mathcal{A} on H defined by:

$$\begin{aligned} D(\pi_1) &= \{f \in C^\infty[0, 1]: f(0) = f(1), f^{(n)}(0) = f^{(n)}(1), n = 1, 2, \dots\} \\ \pi_1(A) &= -i d/dt. \end{aligned}$$

It is straightforward to show that π_1 is hermitian and that $\pi_1 = \bar{\pi}_1 \subsetneq \pi_1^{**} = \pi_1^*$ [8].

We now consider commutants and irreducibility. If $\pi \alpha \pi_1$, define $C(\pi, \pi_1)$ to be the set of operators $C \in L_c(H)$ satisfying $\langle C\pi(A)x, y \rangle = \langle Cx, \pi_1(A^*)y \rangle$ for every $x \in D(\pi)$, $y \in D(\pi_1)$, $A \in \mathcal{A}$. The proof of the following lemma is straightforward.

LEMMA 3.

- (1) $C(\pi, \pi_1)$ is a weakly closed subspace of $L_c(H)$ containing I .
- (2) $C(\pi, \pi_1) = \{C \in L_c(H): C: D(\pi) \rightarrow D(\pi_1^*), C\pi(A) = \pi_1^*(A)C | D(\pi)\}$.
- (3) $C \in C(\pi, \pi_1)$ if and only if $C^* \in C(\pi_1, \pi)$.

The *commutant* of a $*$ -representation π is defined as $\pi(\mathcal{A})' = C(\pi, \pi)$. It follows from Lemma 3 that $\pi(\mathcal{A})'$ is a weakly closed, symmetric subspace of $L_c(H)$ containing I . However, $\pi(\mathcal{A})'$ need not be a von Neumann algebra [6]. If π is self-adjoint then $\pi(\mathcal{A})'$ is a von Neumann algebra [6]. If π is a $*$ -representation, the *strong commutant* is defined by

$$\pi(\mathcal{A})'_s = \{C \in \pi(\mathcal{A})': C: D(\pi) \rightarrow D(\pi)\}.$$

Hence

$$\begin{aligned}\pi(\mathcal{A})'_s &= \{C \in L_c(H): C: D(\pi) \rightarrow D(\pi), C\pi(A) \\ &= \pi(A)C | D(\pi), \forall A \in \mathcal{A}\}.\end{aligned}$$

It is easy to see that $\pi(\mathcal{A})'_s$ is an op-algebra in $L_c(H)$ containing I and if π is closed, then $\pi(\mathcal{A})'_s$ is weakly closed [1]. Again $\pi(\mathcal{A})'_s$ need not be a von Neumann algebra but if π is self-adjoint, then $\pi(\mathcal{A})'_s$ is a von Neumann algebra and $\pi(\mathcal{A})'_s = \pi(\mathcal{A})'$.

LEMMA 4. *A *-representation π is self-adjoint if and only if $\pi(\mathcal{A})' = \pi(\mathcal{A})'_s$ and $D(\pi^*) = \cup\{Cx: x \in D(\pi), C \in \pi(\mathcal{A})'\}$.*

Proof. Necessity follows from our previous observations. For sufficiency, if $\pi(\mathcal{A})' = \pi(\mathcal{A})'_s$ then $C: D(\pi) \rightarrow D(\pi)$ for all $C \in \pi(\mathcal{A})'$. Hence $D(\pi^*) = \cup\{Cx: x \in D(\pi), C \in \pi(\mathcal{A})'\} \subset D(\pi)$.

For a bounded *-representation π of a *-algebra \mathcal{A} on a Hilbert space H the following conditions are equivalent [2, 4].

- (i) $\pi(\mathcal{A})' = \{\lambda I: \lambda \in C\}$.
- (ii) The only invariant closed subspaces of H are $\{0\}$ and H .
- (iii) Every nonzero vector in $H = D(\pi)$ is cyclic.

A bounded *-representation π is said to be *irreducible* if π satisfies any one (and hence all) of these three conditions.

For unbounded self-adjoint representations one can give examples [6, 8] which show that no two of the above conditions are equivalent. Also, there is more than one natural way to extend some of the above conditions for unbounded self-adjoint representations. Let π be a self-adjoint representation. We say that a subspace M is a *self-adjoint invariant subspace* for π if M is invariant and $\pi|_M$ is self-adjoint. The following are natural conditions that one might use to define irreducibility for a self-adjoint representation π of a *-algebra \mathcal{A} with domain $D(\pi) \subseteq H$.

- (1) $\pi(\mathcal{A})' = \{\lambda I: \lambda \in C\}$.
- (2) The only invariant subspaces for π which are complete in the $\pi(\mathcal{A})$ -topology are $\{0\}$ and $D(\pi)$.
- (2') The only self-adjoint invariant subspaces for π are $\{0\}$ and $D(\pi)$.
- (3) Every nonzero vector in $D(\pi)$ is strongly cyclic.
- (3') Every nonzero vector in $D(\pi)$ is cyclic.

THEOREM 5. *If π is self-adjoint representation of the *-algebra \mathcal{A} on the Hilbert space H , then*

$$(2') \leftrightarrow (1) \leftarrow (2)$$

$$\uparrow \qquad \downarrow$$

$$(3') \longleftarrow (3)$$

Proof. (2) \rightarrow (3). Suppose (2) holds, $0 \neq \phi \in D(\pi)$ and $M = \{\pi(A)\phi : A \in \mathcal{A}\}$. Clearly, $M \neq \{0\}$ and M is an invariant subspace of H for π . Let \hat{M} be the completion of M in the $\pi(\mathcal{A})$ -topology. Since π is closed, $\hat{M} \subseteq D(\pi)$ and clearly \hat{M} is a subspace of H . We now show that \hat{M} is invariant under π . If $x \in \hat{M}$, then there exists a net $x_\alpha \in M$ such that $x_\alpha \rightarrow x$ in the $\pi(\mathcal{A})$ -topology. Fix an $A \in \mathcal{A}$. Then for every $B \in \mathcal{A}$ we have

$$\pi(B)\pi(A)x_\alpha = \pi(BA)x_\alpha \rightarrow \pi(BA)x = \pi(B)\pi(A)x.$$

Hence $\pi(A)x_\alpha \rightarrow \pi(A)x$ in the $\pi(\mathcal{A})$ -topology so $\pi(A)x \in \hat{M}$ and $\pi(A)\hat{M} \subseteq \hat{M}$. Since (2) holds, $\hat{M} = D(\pi)$. Hence M is dense in $D(\pi)$ in the $\pi(\mathcal{A})$ -topology so ϕ is a strongly cyclic vector for π .

(3) \rightarrow (2). Suppose (2) does not hold. Then there exists a $\pi(\mathcal{A})$ -complete invariant subspace M of H with $M \neq \{0\}, D(\pi)$. If $0 \neq \phi \in M$, then clearly ϕ is not a strongly cyclic vector for π .

(1) \rightarrow (2'). Suppose (2') does not hold. Then there exists a nontrivial self-adjoint invariant subspace M for π . Now M is not dense in H since otherwise $\pi|_M$ is a *-representation of \mathcal{A} on $\bar{M} = H$ with domain $M \subseteq D(\pi)$. Then $\pi|_M \subset \pi = \pi^* \subset (\pi|_M)^*$. Since $\pi|_M$ is self-adjoint, $\pi|_M = \pi$ and $D(\pi) = M$ which is a contradiction. By Theorem 4.7 [6] the projection E on \bar{M} satisfies $E \in \pi(\mathcal{A})'$. Since $E \neq 0, I$, (1) does not hold.

(2') \rightarrow (1). Suppose (1) does not hold. Since π is self-adjoint, $\pi(\mathcal{A})'$ is a von Neumann algebra so there exists a nontrivial projection $E \in \pi(\mathcal{A})'$. By Theorem 4.7 [6], $ED(\pi)$ is a nontrivial self-adjoint invariant subspace for π . Thus (2') does not hold.

(3') \rightarrow (1). Suppose (3') holds. Let $0 \neq E \in \pi(\mathcal{A})'$ be a projection. By Theorem 4.7 [6], $ED(\pi) = M$ is a self-adjoint invariant subspace for π . Let $0 \neq \phi \in M$. Since ϕ is cyclic and $\{\pi(A)\phi : A \in \mathcal{A}\} \subseteq M$, M is dense in H . As in (1) \rightarrow (2') above, $M = D(\pi)$ and hence $E = I$. Since 0 and I are the only projections in $\pi(\mathcal{A})'$, we have $\pi(\mathcal{A})' = \{\lambda I : \lambda \in C\}$.

(3) \rightarrow (3'). This is trivial. (2) \rightarrow (1). Since (2) \rightarrow (2') trivially, this follows from (2') \rightarrow (1) above.

3. Closed strongly cyclic *-representations. In this section we shall mainly be concerned with characterizing self-

adjointness for closed strongly cyclic *-representations. Let π be a *-representation of a *-algebra \mathcal{A} with domain $D(\pi) \subseteq H$. The *unbounded commutant* $\pi(\mathcal{A})^c$ of π is defined as the set of operators $C \in L(D(\pi), H)$ such that $\langle C\pi(A)x, y \rangle = \langle Cx, \pi(A^*)y \rangle$ for all $x, y \in D(\pi)$ and $A \in \mathcal{A}$. The *strong unbounded commutant* is defined by $\pi(\mathcal{A})_s^c = \{C \in \pi(\mathcal{A})^c : CD(\pi) \rightarrow D(\pi)\}$. Notice that $\pi(\mathcal{A})' | D(\pi) \subset \pi(\mathcal{A})^c$ and $\pi(\mathcal{A})_s' | D(\pi) \subseteq \pi(\mathcal{A})_s^c$. In fact,

$$\begin{aligned}\pi(\mathcal{A})' &= \{\bar{C} : C \in \pi(\mathcal{A})^c, C \text{ bounded}\} \\ \pi(\mathcal{A})_s' &= \{\bar{C} : C \in \pi(\mathcal{A})_s^c, C \text{ bounded}\}.\end{aligned}$$

We say that a net $B_\alpha \in L(M, N)$ converges weakly to $B \in L(M, N)$ if $\langle B_\alpha x, y \rangle \rightarrow \langle Bx, y \rangle$ for every $x, y \in M$. Moreover, $\mathcal{B} \subseteq L(M, N)$ is *weakly closed* if for any net $B_\alpha \in \mathcal{B}$ which converges weakly to some $B \in L(M, N)$ we have $B \in \mathcal{B}$. The proof of the next lemma is straightforward.

LEMMA 6.

- (1) If π is self-adjoint, then $\pi(\mathcal{A})^c = \pi(\mathcal{A})_s^c$.
- (2) $\pi(\mathcal{A})^c = \{C \in L(D(\pi), D(\pi^*)) : C\pi(A) = \pi^*(A)C, \forall A \in \mathcal{A}\}$.
- (3) $\pi(\mathcal{A})_s^c = C \in L(D(\pi)) : C\pi(A) = \pi(A)C, \forall A \in \mathcal{A}\}$.
- (4) $\pi(\mathcal{A})^c$ is a weakly closed subspace of $L(D(\pi), D(\pi^*))$ containing $I | D(\pi)$.
- (5) $\pi(\mathcal{A})_s^c$ is an op-algebra in $L(D(\pi))$.
- (6) $\pi(\mathcal{A})^c = \pi(\mathcal{A})_s^c$ if and only if $\pi(\mathcal{A})^c$ is an op-algebra.

Let \mathcal{A} be a *-algebra and let π, π_1 be *-representations of \mathcal{A} on Hilbert spaces H, H_1 , respectively. We say that π and π_1 are *equivalent*, and write $\pi \cong \pi_1$, if there exists a unitary transformation V from H onto H_1 such that $VD(\pi) = D(\pi_1)$ and $\pi(A) = V^*\pi_1(A)V$ for every $A \in \mathcal{A}$.

Let ω be a state on \mathcal{A} . Then by the *GNS* construction for *-algebras [6], there exists a closed, strongly cyclic *-representation π_ω of \mathcal{A} with strongly cyclic vector x_0 such that $\omega(A) = \langle \pi_\omega(A)x_0, x_0 \rangle$ for every $A \in \mathcal{A}$. Moreover, if π is any closed, strongly cyclic *-representation of \mathcal{A} with strongly cyclic vector y_0 such that $\langle \pi(A)y_0, y_0 \rangle = \omega(A)$ for every $A \in \mathcal{A}$ then $\pi \cong \pi_\omega$ [6].

We now characterize states ω such that π_ω is self-adjoint. A linear functional $F: \mathcal{A} \rightarrow C$ is ω -*bounded* if for every $B \in \mathcal{A}$ there exists an $M_B \geq 0$ such that $|F(BA)| \leq M_B \omega(A^*A)^{1/2}$ for every $A \in \mathcal{A}$. For example, if $A_\alpha \in \mathcal{A}$ is a net such that $\omega(A_\alpha^*A^*AA_\alpha)$ is Cauchy for every $A \in \mathcal{A}$, then the functional $F(A) = \lim \omega(A_\alpha^*A)$ is ω -bounded. Indeed, for every $B \in \mathcal{A}$ we have

$$\begin{aligned} |F(BA)| &= \lim |\omega(A^*BA)| = \lim |\omega(A^*B^*A_\alpha)| \\ &\leq \omega(A^*A)^{1/2} \lim \omega(A_\alpha^*BB^*A_\alpha)^{1/2}. \end{aligned}$$

If every ω -bounded linear functional has the above form, then we call ω a *Riesz state*.

THEOREM 7. *Let ω be a state on the $*$ -algebra \mathcal{A} . Then π_ω is self-adjoint if and only if ω is a Riesz state.*

Proof. Recall that π_ω is constructed as follows. Let \mathcal{I} be the left ideal $\{A \in \mathcal{A} : \omega(A^*A) = 0\}$ and let H_0 be the inner product space consisting of equivalence classes $[A]$ in \mathcal{A}/\mathcal{I} with inner product $\langle [A], [B] \rangle = \omega(B^*A)$. Let H be the Hilbert space completion of H_0 . Define a $*$ -representation π_0 of \mathcal{A} with domain $D(\pi_0) = H_0$ by $\pi_0(A)[B] = [AB]$. If $\pi_\omega = \bar{\pi}_0$, then π_ω is a closed, strongly cyclic $*$ -representation with domain $D(\pi_\omega) = \hat{H}_{0\pi_0(\mathcal{A})}$ and strongly cyclic vector $[I]$. Now suppose π_ω is self-adjoint and $F: \mathcal{A} \rightarrow C$ is ω -bounded. If $\omega(A^*A) = 0$, then $F(A) = 0$ so $F: \mathcal{I} \rightarrow 0$. Hence F can be considered as a linear functional on H_0 . Since $|F([A])| \leq M_I \| [A] \|$, F is a continuous linear functional on H_0 and by the Riesz theorem there exists a $z \in H$ such that $F([A]) = \langle [A], z \rangle$ for every $[A] \in H_0$. Now for every $B \in \mathcal{A}$ we have

$$\begin{aligned} |\langle \pi_0(B)[A], z \rangle| &= |\langle [BA], z \rangle| = |F([BA])| \\ &= |F(BA)| \leq M_B \| [A] \|. \end{aligned}$$

Hence $z \in D(\pi_0^*) = D(\pi_\omega^*) = D(\pi_\omega)$, so there exists a net $[A_\alpha] \in H_0$ which converges to z in the $\pi_0(\mathcal{A})$ -topology. Thus $[AA_\alpha]$ is Cauchy for every $A \in \mathcal{A}$. Finally, for every $A \in \mathcal{A}$ we have

$$F(A) = \lim \langle [A], [A_\alpha] \rangle = \lim \omega(A_\alpha^*A).$$

Conversely, suppose ω is a Riesz state and $x \in D(\pi_\omega^*)$. Define the linear functional $F: \mathcal{A} \rightarrow C$ by $F(A) = \langle [A], x \rangle$. Then for every $A, B \in \mathcal{A}$ we have

$$|F(BA)| = |\langle \pi_0(B)[A], x \rangle| \leq M_B \| [A] \| = M_B \omega(A^*A)^{1/2}$$

so F is ω -bounded. Hence there exists a net $A_\alpha \in \mathcal{A}$ such that $\omega(A_\alpha^*A^*AA_\alpha)$ is Cauchy for every $A \in \mathcal{A}$ and $F(A) = \lim \omega(A_\alpha^*A)$ for every $A \in \mathcal{A}$. It follows that $[A_\alpha]$ is Cauchy in the $\pi_0(\mathcal{A})$ -topology and hence there exists a $y \in D(\pi_\omega)$ such that $[A_\alpha] \rightarrow y$. Furthermore, for every $A \in \mathcal{A}$ we have $F(A) = \lim \omega(A_\alpha^*A) = \lim \langle [A], [A_\alpha] \rangle = \langle [A], y \rangle$. Hence $x = y \in D(\pi_\omega)$ and π_ω is self-adjoint.

COROLLARY. A closed, strongly cyclic *-representation π with strongly cyclic vector x_0 is self-adjoint if and only if the state $A \rightarrow \langle \pi(A)x_0, x_0 \rangle$ is a Riesz state.

A state ω is *faithful* if $\omega(A^*A) = 0$ implies $A = 0$. A vector $x_0 \in D(\pi)$ is *separating* if $\pi(A)x_0 = 0$ implies $\pi(A) = 0$. If ω is faithful then the strongly cyclic vector x_0 for π_ω is separating. Conversely, if x_0 is separating, then $\omega(A^*A) = 0$ implies $\pi_\omega(A) = 0$. A representation π of \mathcal{A} is *ultra-cyclic* if there exists an $x_0 \in D(\pi)$ such that $D(\pi) = \{\pi(A)x_0 : A \in \mathcal{A}\}$. We then call x_0 an *ultra-cyclic vector*. Ultra-cyclic representations are important because of the following result.

LEMMA 8. π is a closed, strongly cyclic *-representation if and only if π is the closure of an ultra-cyclic *-representation π^0 .

Proof. Suppose π is a closed, strongly cyclic *-representation of \mathcal{A} with strongly cyclic vector x_0 . Define $D(\pi^0) = \{\pi(A)x_0 : A \in \mathcal{A}\}$ and $\pi^0(B)\pi(A)x_0 = \pi(BA)x_0$. Then π^0 is an ultra-cyclic *-representation and $\bar{\pi}^0 = \pi$. Conversely, if π is the closure of an ultra-cyclic *-representation π^0 with ultra-cyclic vector x_0 , then π is a closed *-representation. Moreover, since $D(\pi)$ is the completion of $D(\pi^0)$ in the $\pi(\mathcal{A})$ -topology, x_0 is a strongly cyclic vector for π .

We call π^0 in the proof of Lemma 8 the *underlying* ultra-cyclic *-representation for π . We can obtain information about π by studying the simpler representation π^0 . For example, a condition characterizing the essential self-adjointness of π^0 characterizes the self-adjointness of π . Moreover, $\pi^{0*} = \pi^*$ and $\pi^0(\mathcal{A}') = \pi(\mathcal{A})'$.

Let π be an arbitrary ultra-cyclic *-representation of \mathcal{A} with a separating ultra-cyclic vector x_0 . For $x \in D(\pi^*)$ define $\pi^c(x) \in L(D(\pi), D(\pi^*))$ by $\pi^c(x)\pi(A)x_0 = \pi^*(A)x$. This is a well-defined operator since $\pi(A)x_0 = \pi(B)x_0$ implies $\pi(A) = \pi(B)$. Then for every $y, z \in D(\pi)$ we have

$$\langle \pi(A^*)y, z \rangle = \langle y, \pi(A)z \rangle = \langle y, \pi(B)z \rangle = \langle \pi(B^*)y, z \rangle.$$

Hence $\pi(A^*) = \pi(B^*)$, so $\pi(A^*)^* = \pi(B^*)^*$ and finally

$$\pi^*(A) = \pi(A^*)^*|D(\pi^*) = \pi(B^*)^*|D(\pi^*) = \pi^*(B).$$

It is straightforward to see that $D(\pi)$ is a *-algebra with identity x_0 under the product $(\pi(A)x_0) \circ (\pi(B)x_0) = \pi(AB)x_0$ and involution $(\pi(A)x_0)^* =$

$\pi(A^*)x_0$. Moreover, for every $x, y, z \in D(\pi)$ we have $\langle x \circ y, z \rangle = \langle y, x^* \circ z \rangle$.

THEOREM 9. *Let π be an ultra-cyclic *-representation of \mathcal{A} with a separating, ultra-cyclic vector x_0 .*

- (1) π^c is a weakly continuous linear bijection from $D(\pi^*)$ into $\pi(\mathcal{A})^c$.
- (2) The following statements are equivalent.
 - (a) $Cx_0 \in D(\pi)$ for every $C \in \pi(\mathcal{A})^c$.
 - (b) $\pi(\mathcal{A})^c$ is an op-algebra.
 - (c) π is self-adjoint.
- (3) $\pi(\mathcal{A})^c$ is an op*-algebra if and only if π is self-adjoint and there exists an involution b on the *-algebra $D(\pi)$ satisfying

$$(3.1) \quad \langle x^*, y \rangle = \langle y^b, x \rangle$$

for every $x, y \in D(\pi)$.

- (4) If $\pi(\mathcal{A})^c$ is an op*-algebra, then π^c is a weakly continuous b -anti-isomorphism of $D(\pi)$ onto $\pi(\mathcal{A})^c$.

Proof.

- (1) Clearly, π^c is linear. To show that π^c maps $D(\pi^*)$ into $\pi(\mathcal{A})^c$, for $x \in D(\pi^*)$, $A \in \mathcal{A}$, $z \in D(\pi)$ and $y = \pi(B)x_0 \in D(\pi)$ we have

$$\begin{aligned} \langle \pi^c(x)\pi(A)y, z \rangle &= \langle \pi^c(x)\pi(AB)x_0, z \rangle \\ &= \langle \pi^*(AB)x, z \rangle = \langle \pi^*(B)x, \pi(A^*)z \rangle \\ &= \langle \pi^c(x)\pi(B)x_0, \pi(A^*)z \rangle = \langle \pi^c(x)y, \pi(A^*)z \rangle. \end{aligned}$$

To show that π^c is surjective, let $C \in \pi(\mathcal{A})^c$. Then $Cx_0 \in D(\pi^*)$ and for any $y = \pi(A)x_0 \in D(\pi)$ we have

$$\begin{aligned} \pi^c(Cx_0)y &= \pi^c(Cx_0)\pi(A)x_0 = \pi^*(A)Cx_0 \\ &= C\pi(A)x_0 = Cy. \end{aligned}$$

To show that π^c is injective, suppose that $x, x_1 \in D(\pi^*)$ and $\pi^c(x) = \pi^c(x_1)$. Then

$$x = \pi^*(1)x = \pi^c(x)x_0 = \pi^c(x_1)x_0 = \pi^*(1)x_1 = x_1.$$

To show that π^c is weakly continuous, suppose that $x_i, x \in D(\pi^*)$ and

$x_i \rightarrow x$ in norm. Then for any $y = \pi(B)x_0 \in D(\pi)$ and $z \in D(\pi)$ we have

$$\begin{aligned}\lim \langle \pi^c(x_i)y, z \rangle &= \lim \langle \pi^*(B)x_i, z \rangle \\ &= \lim \langle x_i, \pi(B^*)z \rangle = \langle x, \pi(B^*)z \rangle \\ &= \langle \pi^*(B)x, z \rangle = \langle \pi^c(x)\pi(B)x_0, z \rangle \\ &= \langle \pi^c(x)y, z \rangle.\end{aligned}$$

(2) (a) \rightarrow (b). Suppose that (a) holds and $C \in \pi(\mathcal{A})^c$, $y = \pi(A)x_0 \in D(\pi)$. We then have $Cy = C\pi(A)x_0 = \pi^*(A)Cx_0 = \pi(A)Cx_0 \in D(\pi)$. Hence, by Lemma 6(6), $\pi(\mathcal{A})^c$ is an op-algebra.

(b) \rightarrow (c). If $x \in D(\pi^*)$, then by (1) $\pi^c(x) \in \pi(\mathcal{A})^c$. If $\pi(\mathcal{A})^c$ is an op-algebra, then $x = \pi^*(1)x = \pi^c(x)x_0 \in D(\pi)$. Hence $D(\pi^*) = D(\pi)$ and π is self-adjoint.

(c) \rightarrow (a). If π is self-adjoint, then $\pi(\mathcal{A})^c \subseteq L(D(\pi))$.

(3) Suppose $\pi(\mathcal{A})^c$ is an op*-algebra. Then, by (2), π is self-adjoint. If $C \in \pi(\mathcal{A})^c$, then $D(\pi) \subseteq D(C^*)$ and $C^*|D(\pi) \in \pi(\mathcal{A})^c$ so $C^*: D(\pi) \rightarrow D(\pi)$. For $x \in D(\pi)$, by (1) $\pi^c(x) \in \pi(\mathcal{A})^c$ so $x^b \equiv \pi^c(x)^*x_0 \in D(\pi)$. For $x = \pi(A)x_0 \in D(\pi)$ and $y \in D(\pi)$ we have

$$\begin{aligned}\langle y^b, x \rangle &= \langle \pi^c(y)^*x_0, x \rangle = \langle x_0, \pi^c(y)\pi(A)x_0 \rangle \\ &= \langle x_0, \pi(A)y \rangle = \langle \pi(A^*)x_0, y \rangle = \langle x^*, y \rangle\end{aligned}$$

so (3.1) holds. That b is an involution now follows from (3.1). For example,

$$\begin{aligned}\langle (y \circ z)^b, x \rangle &= \langle x^*, y \circ z \rangle = \langle y^* \circ x^*, z \rangle \\ &= \langle z^b, x \circ y \rangle = \langle x^* \circ z^b, y \rangle = \langle y^b, z^{b*} \circ x \rangle \\ &= \langle z^b \circ y^b, x \rangle.\end{aligned}$$

The other properties of an involution follow in a similar way. Conversely, suppose π is self-adjoint and there exists an involution b on $D(\pi)$ satisfying (3.1). Then by (2), $\pi(\mathcal{A})^c$ is an op-algebra. If $C \in \pi(\mathcal{A})^c$, then for any $x = \pi(B)x_0 \in D(\pi)$ and $y = \pi(A)x_0 \in D(\pi)$ we have

$$\begin{aligned}\langle Cy, x \rangle &= \langle C\pi(A)x_0, x \rangle = \langle Cx_0, \pi(A^*)x \rangle \\ &= \langle Cx_0, \pi(A^*B)x_0 \rangle = \langle Cx_0, [\pi(B^*A)x_0]^* \rangle \\ &= \langle \pi(B^*A)x_0, (Cx_0)^b \rangle = \langle \pi(A)x_0, \pi(B)[(Cx_0)^b] \rangle \\ &= \langle y, \pi^c[(Cx_0)^b]x \rangle.\end{aligned}$$

Hence $D(\pi) \subseteq D(C^*)$, $C^*|D(\pi) = \pi^c[(Cx_0)^b] \in \pi(\mathcal{A})^c$ and so $\pi(\mathcal{A})^c$ is an op*-algebra.

(4) Suppose $\pi^c(\mathcal{A})^c$ is an op*-algebra. It follows from (1) that $\pi^c: D(\pi) \rightarrow \pi^c(\mathcal{A})^c$ is a weakly continuous linear bijection. For $x = \pi(A)x_0 \in D(\pi)$ and $y = \pi(B)x_0 \in D(\pi)$ we have

$$\pi^c(x)y = \pi^c(x)\pi(B)x_0 = \pi(B)x = \pi(B)x_0 \circ \pi(A)x_0 = y \circ x.$$

It is now clear that π^c is an anti-isomorphism. To show that π^c is a b -anti-isomorphism, for $x \in D(\pi)$ and $y = \pi(A)x_0 \in D(\pi)$ we have

$$\begin{aligned} \pi^c(x^b)y &= y \circ x^b = y \circ [\pi^c(x)^*x_0] = \pi(A)\pi^c(x^*)x_0 \\ &= \pi^c(x)^*\pi(A)x_0 = \pi^c(x)^*y. \end{aligned}$$

COROLLARY. *Let π be a closed, strongly cyclic *-representation of \mathcal{A} with separating, strongly cyclic vector x_0 and let π^0 be the underlying ultra-cyclic representation. Then π is self-adjoint if and only if $Cx_0 \in D(\pi)$ for every $C \in \pi^0(\mathcal{A})^c$.*

Proof. If π is self-adjoint and $C \in \pi^0(\mathcal{A})^c$, then $Cx_0 \in D(\pi^{0*}) = D(\pi^*) = D(\pi)$. Conversely, suppose $Cx_0 \in D(\pi)$ for every $C \in \pi^0(\mathcal{A})^c$. If $x \in D(\pi^*)$, then $x \in D(\pi^{0*})$ so by Theorem 9(1), $\pi^c(x) \in \pi^0(\mathcal{A})^c$. Hence $x = \pi^c(x)x_0 \in D(\pi)$, so $D(\pi) = D(\pi^*)$ and π is self-adjoint.

REFERENCES

1. H.J. Borchers and J. Yngvason, *On the algebra of field operators. The weak commutant and integral decomposition of states*, Commun. Math. Phys. **42** (1975), 231-252.
2. G.G. Emch, *Algebraic Methods in Statistical Mechanics and Quantum Field Theory*, John Wiley & Sons, Inc., New York (1972).
3. G. Lassner, *Topological algebras of operators*, Rep. Math. Phys., **3** (1972), 279-293.
4. M.A. Naimark, *Normed Rings*, Noordhoff Publishers, Groningen, The Netherlands (1970).
5. E. Nelson, *Analytic vectors*, Ann. Math., **70** (1959), 572-615.
6. R.T. Powers, *Self-adjoint algebras of unbounded operators*, Commun. Math. Phys., **21** (1971), 85-124.
7. ———, *Self-adjoint algebras of unbounded operators II*, Trans. Amer. Math. Soc., **187** (1974), 1-33.
8. W.M. Scruggs, *Unbounded representations of *-algebras*, Dissertation, University of Denver (1976).
9. B. Simon, *The $P(\phi)_2$ Euclidean (Quantum) Field Theory*, Princeton University Press, Princeton, New Jersey (1974).
10. R.F. Streater and A.S. Wightman, *PCT, Spin and Statistics and All That*, W.A. Benjamin, Inc., New York (1964).

Received January 13, 1977 and in revised form May 19, 1977.

UNIVERSITY OF DENVER
DENVER, CO 80210

PACIFIC JOURNAL OF MATHEMATICS
EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. A. BEAUMONT
University of Washington
Seattle, WA 98105

R. FINN AND J. MILGRAM
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$ 72.00 a year (6 Vols., 12 issues). Special rate: \$ 36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1977 Pacific Journal of Mathematics
All Rights Reserved

Pacific Journal of Mathematics

Vol. 70, No. 2

October, 1977

B. Arazi, <i>A generalization of the Chinese remainder theorem</i>	289
Thomas E. Armstrong, <i>Polyhedrality of infinite dimensional cubes</i>	297
Yoav Benyamini, Mary Ellen Rudin and Michael L. Wage, <i>Continuous images of weakly compact subsets of Banach spaces</i>	309
John Thomas Burns, <i>Curvature functions on Lorentz 2-manifolds</i>	325
Dennis F. De Raggi and Nelson Groh Markley, <i>Shear distality and equicontinuity</i>	337
Claes Fernström, <i>Rational approximation and the growth of analytic capacity</i>	347
Pál Fischer, <i>On some new generalizations of Shannon's inequality</i>	351
Che-Kao Fong, <i>Quasi-affine transforms of subnormal operators</i>	361
Stanley P. Gudder and W. Scruggs, <i>Unbounded representations of $*$-algebras</i>	369
Chen F. King, <i>A note on Drazin inverses</i>	383
Ronald Fred Levy, <i>Countable spaces without points of first countability</i>	391
Eva Lowen-Colebunders, <i>Completeness properties for convergence spaces</i>	401
Calvin Cooper Moore, <i>Square integrable primary representations</i>	413
Stanisław G. Mrówka and Jung-Hsien Tsai, <i>On preservation of E-compactness</i>	429
Yoshiomi Nakagami, <i>Essential spectrum $\Gamma(\beta)$ of a dual action on a von Neumann algebra</i>	437
L. Alayne Parson, <i>Normal congruence subgroups of the Hecke groups $G(2^{(1/2)})$ and $G(3^{(1/2)})$</i>	481
Louis Jackson Ratliff, Jr., <i>On the prime divisors of zero in form rings</i>	489
Caroline Series, <i>Ergodic actions of product groups</i>	519
Robert O. Stanton, <i>Infinite decomposition bases</i>	549
David A. Stegenga, <i>Sums of invariant subspaces</i>	567