

Uniform Density in Lindenbaum Algebras

V. Yu. Shavrukov and Albert Visser

Abstract In this paper we prove that the preordering \lesssim of provable implication over any recursively enumerable theory T containing a modicum of arithmetic is uniformly dense. This means that we can find a recursive extensional density function F for \lesssim . A recursive function F is a density function if it computes, for A and B with $A \lesssim B$, an element C such that $A \lesssim C \lesssim B$. The function is extensional if it preserves T -provable equivalence. Secondly, we prove a general result that implies that, for extensions of elementary arithmetic, the ordering \lesssim restricted to Σ_n -sentences is uniformly dense. In the last section we provide historical notes and background material.

1 Introduction

It is well known that the Lindenbaum algebras of theories that contain a modicum of arithmetic are dense with respect to the implication ordering. In this paper we will study a property that is stronger than *density*, to wit, *uniform density*. We prove that the Lindenbaum algebras of these theories are uniformly dense with respect to the implication ordering. We first provide the necessary definitions to formulate the result.

Consider any recursively enumerable theory T that interprets the theory R introduced by Tarski, Mostowski, and Robinson in [21]. We define

- $A \lesssim_T B$ if and only if $T + A \vdash B$,
- $A \lesssim_T B$ if and only if $A \lesssim_T B$ and not $B \lesssim_T A$,
- $A \sim_T B$ if and only if $A \lesssim_T B$ and $B \lesssim_T A$.

Here \lesssim_T is the “provable implication” ordering on \mathcal{L}_T , the Lindenbaum sentence algebra of T . It is well known that \lesssim_T is dense. We say that \mathcal{L}_T (or \lesssim_T) is *uniformly dense* if there is a recursive function F such that

1. F is a *density function*: that is, we have $A \lesssim_T F(A, B) \lesssim_T B$, whenever $A \lesssim_T B$, and if $A \sim_T B$, then $A \sim_T F(A, B) \sim_T B$;

Received January 28, 2012; accepted January 15, 2013

2010 Mathematics Subject Classification: Primary 03F40; Secondary 03D45, 03F30

Keywords: first-order theories, arithmetic, uniform density, Lindenbaum algebras

© 2014 by University of Notre Dame 10.1215/00294527-2798754

2. F is *extensional*: if $A \sim_T A'$ and $B \sim_T B'$, then $F(A, B) \sim_T F(A', B')$.

We show that \mathfrak{L}_T is uniformly dense for recursively enumerable theories T that interpret R . Moreover, we can take the function F to be elementary and, in some specific cases, even p-time computable.

We present our proof in Section 3. It consists of two stages. First we prove the desired result for Peano arithmetic PA or, more generally, for essentially reflexive theories (see Remark 1.1 for a discussion of essential reflexivity). Our construction delivers a p-time computable density function. This result is then generalized to all r.e. consistent theories that interpret R although the new density functions fall outside polynomial time—we don't know if this can be rectified.

A variant of the density question is obtained by imposing a restriction to a prescribed formula class. We explore this variant in Section 4. We prove a general result which implies, for example, that, for extensions T of elementary arithmetic, the ordering \lesssim_T restricted to Σ_n -sentences is uniformly dense.

The basic idea and the ingredients of our construction for PA come with a history. The sentences we produce are certain unique Rosser sentences of a kind studied by Smoryński [19], and they are the unique Gödel sentences of a certain Feferman predicate studied in Shavrukov [18]. Finally, they are Orey sentences. We will explain this background in Section 5. The reader who wants to just see the solution may, of course, skip Section 5.

Remark 1.1 A theory is *essentially reflexive* (*uniformly essentially reflexive*) if it proves reflection (resp., uniform reflection) for each of its finitely axiomatized subtheories. Here (uniform) reflection concerns a proof predicate that is formalized with respect to an interpretation N of a weak arithmetic, like S_2^1 , in the given theory. Thus, a theory U is essentially reflexive with respect to N if it proves all principles of the form $\vdash \Box_{U_0}^N A \rightarrow A$, where A is a *sentence* of the language of U and U_0 is any finitely axiomatized subtheory of U . A theory U is uniformly essentially reflexive with respect to N if it proves all principles of the form $\vdash \forall \vec{x} (\Box_{U_0}^N A\vec{x} \rightarrow A\vec{x})$, where $A\vec{x}$ is a *formula* of the language of U with all variables among those shown and U_0 is any finitely axiomatized subtheory of U . We refer the reader to Beklemishev [1, Section 2] for a more extensive discussion.

Uniform essential reflexivity implies full induction with respect to the designated interpretation of the numbers. Conversely, a theory that satisfies full induction and is sequential is uniformly essentially reflexive (for the definition of *sequential* see Hájek and Pudlák [5] or Visser [26]). If we drop uniformity, essentially reflexive theories can be much weaker. For example, the minimal essentially reflexive extension of elementary arithmetic EA (here $EA = I\Delta_0 + \text{Exp}$) is both a subtheory of PA and of EA plus all true Π_1 -sentences. See also Visser [28].

Essentially reflexive theories form a natural class in the study of interpretability. For example, they satisfy the same propositional schemes for interpretability. See, for example, Visser [27] for a discussion.

2 The Usual Proof of Density

Our proof of uniform density for \mathfrak{L}_{PA} is a specific instance of the usual proof of the density of \mathfrak{L}_T , where T is a recursively enumerable theory that interprets R . We first present this usual proof. This proof will form a frame of reference for the rest of the paper.

Suppose $A \not\lesssim_T B$. It follows that $T + \neg A + B$ is consistent.

Let C be any arithmetical sentence that is independent of $T + \neg A + B$; that is, $T + \neg A + B \not\vdash C$ and $T + \neg A + B \not\vdash \neg C$.

The essential ingredients of the proof that, for every consistent recursively enumerable theory U that interprets R , there exists a sentence R that is independent of U were provided by J. Barkley Rosser in his classical paper [16]. Here R is a very weak arithmetic introduced in [21].

We consider $D := A \vee (C \wedge B)$. We claim that $A \not\lesssim_T D \not\lesssim_T B$.

First, we clearly have $A \lesssim_T D \lesssim_T B$.

Suppose $T + B \vdash A \vee (C \wedge B)$. By propositional logic, we find $T + \neg A + B \vdash C$. Quod non.

Suppose $T + A \vee (C \wedge B) \vdash A$. By propositional logic, we have $T + \neg A + B \vdash \neg C$. Quod non.

We can squeeze a little bit more information out of our construction. If C is independent, then $\neg C$ is also independent. So we can construct $E := A \vee (\neg C \wedge B)$. We find that $A \not\lesssim_T E \not\lesssim_T B$. Moreover, we have $T \vdash (D \wedge E) \leftrightarrow A$ and $T \vdash (D \vee E) \leftrightarrow B$. So we have two sentences D and E strictly between A and B such that B is the supremum with respect to \lesssim_T of D and E and A is the infimum with respect to \lesssim_T of D and E .

Rosser's construction delivers a p-time mapping $A \mapsto C_A$, where C_A is independent over $T + A$, provided that $T + A$ is consistent, for a given r.e. theory T that interprets R . It is very unlikely, however, that the usual Rosser construction produces a uniform density function. For a brief discussion of this issue see Section 5.

In light of the proof given above, to prove p-time uniform density for PA, it is, modulo some simple details, sufficient to give a p-time construction $A \mapsto C_A$, where C_A is independent over $PA + A$, provided that $PA + A$ is consistent and, if $PA \vdash A_0 \leftrightarrow A_1$, then $PA \vdash C_{A_0} \leftrightarrow C_{A_1}$.

In the next section, we will provide a mapping $A \mapsto C_A$ that satisfies the desiderata.

3 Uniform Density

The first order of business for this section is the following.

Theorem 3.1 *\mathfrak{L}_{PA} is uniformly dense via a p-time computable density function. More generally, this result holds for all essentially reflexive sequential r.e. theories.*

This theorem will later be extended to more theories.

We are interested in getting our density function as efficient as possible. In this case, we will construct a p-time computable function. To obtain an algorithm of the desired complexity, we will use efficiently coded syntax and base 2 numerals. See, for example, Buss [2, Chapter 7.3] or Hájek and Pudlák [5, Chapter V, Section 3].

We consider the sequence of theories Ar_n , where Ar_0 is EA, also known as $I\Delta_0 + \text{Exp}$, and $Ar_{n+1} := I\Sigma_{n+1}$. These theories have the following important property.

Theorem 3.2 *(EA proves that) for all n , Ar_{n+1} proves uniform Π_{n+3} -reflection for Ar_n .*

The theorem claims formalizability in EA of a result that was presumably first stated in Ono [12, Theorem 4.4]. That result is readily seen to be equivalent to $I\Sigma_{n+1}$

proving uniform Π_{n+3} -reflection for EA or even for pure predicate calculus, for $\text{IS}_n + \text{Exp}$ can be axiomatized by a single Π_{n+2} -sentence (see [5, Theorems I.2.52 and V.5.6]). The latter equivalent coincides with the lemma in Leivant [7] modulo some ambiguity as to what Leivant's base theory Z_0 exactly is. The verification of EA-provability is best carried out along the proof of Теорема 7 in [1]. We note that we have the following as an immediate consequence.

Corollary 3.3 (EA proves that) for any sentence A in Σ_{n+3} , $\text{Ar}_{n+1} + A$ proves uniform Π_{n+3} -reflection for $\text{Ar}_n + A$.

We have the following definitions:

$$\begin{aligned} \Box_{A,x} B &\text{ stands for } \text{prov}_{\text{Ar}_x + A}(\ulcorner B \urcorner), \\ \Diamond_{A,x} B &\text{ stands for } \neg \text{prov}_{\text{Ar}_x + A}(\ulcorner \neg B \urcorner); \text{ that is, } \neg \Box_{A,x} \neg B. \end{aligned}$$

As a first step towards the proof of Theorem 3.1 we need the formula C_A :

$$C_A := A \wedge \forall x (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp).$$

We have the following useful lemma.

Lemma 3.4 Given any n , suppose $A \in \Sigma_{n+3}$. Then,

$$\text{Ar}_n \vdash C_A \leftrightarrow (A \wedge \forall x \geq n (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp)).$$

Proof By Corollary 3.3, $\text{Ar}_n + A$ proves Σ_1 -reflection for $\Box_{A,k}$ with $k < n$. Hence,

$$\text{Ar}_n + A \vdash \forall x < n (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp).$$

The desired result is immediate. \square

In the next lemma, we show that $A \mapsto C_A$ is extensional.

Lemma 3.5 Suppose $\text{PA} \vdash A_0 \leftrightarrow A_1$; then $\text{PA} \vdash C_{A_0} \leftrightarrow C_{A_1}$.

Proof Suppose $\text{PA} \vdash A_0 \leftrightarrow A_1$. Then, for some n , $\text{Ar}_n \vdash A_0 \leftrightarrow A_1$. We can pick n so large that $A_0, A_1 \in \Sigma_{n+3}$. We have

$$\begin{aligned} \text{Ar}_n \vdash C_{A_0} &\leftrightarrow A_0 \wedge \forall x \geq n (\Box_{A_0,x} \Box_{A_0,x} \perp \rightarrow \Box_{A_0,x} \perp) \\ &\leftrightarrow A_1 \wedge \forall x \geq n (\Box_{A_1,x} \Box_{A_1,x} \perp \rightarrow \Box_{A_1,x} \perp) \\ &\leftrightarrow C_{A_1}. \end{aligned}$$

Hence, $\text{PA} \vdash C_{A_0} \leftrightarrow C_{A_1}$. \square

Lemma 3.6 Suppose that $\text{PA} + A$ is consistent. Then C_A is independent over $\text{PA} + A$.

Proof Suppose that $\text{PA} + A \vdash C_A$. Then, for some n , $\text{Ar}_n + A \vdash C_A$. We may assume that $A \in \Sigma_{n+3}$. It follows that $\text{Ar}_n + A \vdash \Box_{A,n} \Box_{A,n} \perp \rightarrow \Box_{A,n} \perp$. Hence, by Löb's theorem, $\text{Ar}_n + A \vdash \Box_{A,n} \perp$. We may conclude that $\text{PA} + A \vdash \perp$. Quod non.

Suppose that $\text{PA} + A \vdash \neg C_A$. Then, for some n , $\text{Ar}_n + A \vdash \neg C_A$. We may assume that $A \in \Sigma_{n+3}$. We find, using Lemma 3.4,

$$\text{Ar}_n + A \vdash \exists x \geq n (\Box_{A,x} \Box_{A,x} \perp \wedge \Diamond_{A,x} \top).$$

But then $\text{Ar}_n + A \vdash \Diamond_{A,n} \top$, contradicting the second incompleteness theorem. \square

We put $F(A, B) := A \vee (C_{\neg A \wedge B} \wedge B)$. We note that this is defined for any arithmetical sentences A and B . Moreover, we always have $A \lesssim_{\text{PA}} F(A, B) \lesssim_{\text{PA}} A \vee B$. By Lemma 3.5, F is extensional. By Lemma 3.6, we have that, if $A \lesssim_{\text{PA}} B$, then $A \lesssim_{\text{PA}} F(A, B) \lesssim_{\text{PA}} B$.

This concludes the proof of Theorem 3.1. We note that F is p-time in A and B .

Remark 3.7 We can put

$$F_0(A, B) := A \vee (C_{\neg A \wedge B} \wedge B) \quad \text{and} \quad F_1(A, B) := A \vee (\neg C_{\neg A \wedge B} \wedge B),$$

and then construct an infinite \lesssim_{PA} -antichain between A and B by considering

$$\begin{aligned} D_{A,B,0} &:= F_0(A, B), & D_{A,B,1} &:= F_0(A, F_1(A, B)), \\ D_{A,B,2} &:= F_0(A, F_1(A, F_1(A, B))), & \dots & \end{aligned}$$

The mapping $H : A, B, n \mapsto D_{A,B,n}$ need not be p-time, but since this mapping is elementary we can represent it in PA. Since the $D_{A,B,n}$ have complexity (in the sense of the arithmetical hierarchy) bounded by the maximum of 2, the complexity of A and that of B —say the complexity is $k(A, B)$ —we can, using efficient numerals, replace $D_{A,B,n}$ by $E_{A,B,n} := \text{True}_{k(A,B)}(H(A, B, n))$. If we use a reasonable version of the definition of True_k , the mapping $A, B, n \mapsto E_{A,B,n}$ becomes p-time. (Note that we do not need to worry about the length of the verifications of the usual properties of the True_k . We are only interested in the size of the formulas.)

Our proof can be immediately adapted to any essentially reflexive theory—like ZF: all the ingredients of the construction of C are also present in such a theory.

Let $C_A^\circ := \forall x (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp)$. Note that C_A° is Δ_2 over IS_1 because it is IS_1 -provably equivalent to $\Diamond_A \top \vee \exists x (\Box_{A,x+1} \perp \wedge \Diamond_{A,x} \Diamond_{A,x} \top)$.

Let us consider the relationship between C_A° and $\text{con}(\text{PA} + A)$.

Proposition 3.8 We have $\text{PA} + \text{con}(\text{PA} + A) \vdash C_A^\circ$.

Proof Suppose that A is Σ_{n+3} . We reason in $\text{PA} + \text{con}(\text{PA} + A)$. Suppose that $\Box_{A,x} \Box_{A,x} \perp$. Then $\Box_{A, \max\{x+1, n\}} \perp$ and, hence, $\Box_A \perp$. Quod non. We may conclude $\neg \Box_{A,x} \Box_{A,x} \perp$, and, a fortiori, C_A° . \square

Since, as we will show in Section 5, C_A° is an Orey sentence of $\text{PA} + A$ and, provided that $\text{PA} + A$ is consistent, $\text{con}(\text{PA} + A)$ is not an Orey sentence, C_A° is strictly between $A + \text{con}(\text{PA} + A)$ and A over PA. In other words, C_A° is a reflection principle that is strictly between $\text{con}(\text{PA} + A)$ and \top over $\text{PA} + A$.

Theorem 3.1 generalizes to theories containing R thanks to the following theorem.

Theorem 3.9 (Pour-El and Kripke [14, Theorem 2]) *The Lindenbaum sentence algebras of all recursively enumerable, consistent theories that interpret R are effectively isomorphic.*

For us, “effective isomorphism” means a recursive function from sentences of one theory to those of the other theory that, through provable equivalence, quotients down to an isomorphism between the two Lindenbaum algebras. The functions constructed in [14], however, possess further nice properties.

Pulling the density function of Theorem 3.1 off \mathcal{L}_{PA} back to \mathcal{L}_T along an effective isomorphism $\mathcal{L}_T \rightarrow \mathcal{L}_{\text{PA}}$, we obtain the following.

Corollary 3.10 *The Lindenbaum sentence algebras of all recursively enumerable consistent theories that interpret R enjoy uniform density.*

It is seen from the proof of [14, Theorem 2] that the isomorphisms of Theorem 3.9 together with their inverses can be given by elementary functions (i.e., ones from Grzegorzczak class \mathcal{E}^3).

Accordingly, we are only able to claim elementarity rather than polynomial time for the second-hand density functions obtained via (the intended proof of) Corollary 3.10. Our proof also forfeits the ability to have uniform density achieved by just mixing in an appropriate Δ_2 -sentence.

4 Orderings of Σ_n -Sentences and Precomplete Lattices

We address the question of uniform density for restricted classes of formulas in a somewhat more general setting.

An *r.e. lattice* L is a pair of recursive functions \vee and \wedge defined on an r.e. subset field L of ω together with an r.e. equivalence relation \sim on field L which is a congruence for \vee and \wedge and such that the quotient is a lattice. If that lattice is Boolean, then L is called an *r.e. Boolean algebra* and, as is easily seen, has a recursive negation function. Lindenbaum sentence algebras of r.e. theories provide typical examples.

A *density function* for L is a function $D : (\text{field } L)^2 \rightarrow \text{field } L$ such that, if $a \not\approx b$, then $a \not\approx D(a, b) \not\approx b$, and $D(a, b) \sim a \sim b$ whenever $a \sim b$. The function D is *extensional* (with respect to \sim or L) if \sim is a congruence for D . The lattice L is *uniformly dense* if it admits an effective extensional density function—note that for \mathcal{L}_T this agrees with our earlier definition.

Montagna and Sorbi [11, Proposition 3.1(b)] extend Theorem 3.9 to all *effectively inseparable* (e.i.) r.e. Boolean algebras, that is, algebras where the \sim -equivalence classes of (Boolean) 0 and 1 are effectively inseparable within field L . Hence Corollary 3.10 also holds for all e.i. r.e. Boolean algebras.

When the proof of Theorem 3.1 works for a theory T , it works equally well for the sublattice Σ_n/T of \mathcal{L}_T determined by Σ_n -sentences provided that $n > 1$ because $D(a, b)$ is a lattice polynomial in a, b , and a Δ_2 -sentence. In this section we handle Σ_1/T using a different approach which starts with the definition of a precomplete numeration/equivalence.

A nontrivial equivalence relation \sim on an r.e. subset field \sim of ω is *precomplete* if to every partial recursive $f : \omega \rightarrow \text{field } \sim$ there is a total recursive $F : \omega \rightarrow \text{field } \sim$ that *makes f total modulo \sim* , that is, $F(n) \sim f(n)$ whenever $f(n)$ converges. Reducing f to a universal (field \sim)-valued partial recursive function, we see that an index for F can be found effectively in one for f .

An r.e. lattice L is *precomplete* if its associated (r.e.) equivalence relation \sim is (see [11, Section 2] or Selivanov [17, 4.4]). By \lesssim we denote the corresponding (r.e.) preorder on field L .

Example 4.1 (Visser [23, 1.6.6]) Σ_n/T is r.e. and precomplete whenever T is a consistent r.e. extension of EA.

Hint The mapping that assigns to k the Σ_n -sentence $\exists y (y = f(\vec{k}) \wedge \text{True}_n(y))$ makes f total modulo T -provable equivalence. \square

It is an open question whether Σ_1/S_2^1 or $\exists\Sigma_1^b/S_2^1$ is precomplete (see [2] for definitions of $\exists\Sigma_1^b$ and S_2^1).

Sentences of the form $\exists x T_0(\underline{n}, x)$, where T_0 is Kleene's T-predicate for the 0-ary case, form an example of a class Γ , such that Γ/S_2^1 is r.e., precomplete, and is, modulo S_2^1 -provability, a sublattice of $\mathfrak{L}_{S_2^1}$.

Mutual interpretability for finitely axiomatized sequential theories is also r.e. precomplete, since the interpretability ordering on finitely axiomatized sequential theories (modulo mutual interpretability) is p-time anti-isomorphic to Π_1/EA . This uses the Friedman characterization of interpretability between finitely axiomatized theories (see, e.g., [25, Theorem 3.6]). Thus, the lattice of finitely axiomatized sequential interpretability degrees is (p-time) isomorphic to Σ_1/EA .

The r.e. extensions of PA in the language of PA modulo interpretability give us under an appropriate indexing an example of a precomplete numeration that is *not* recursively enumerable.

Theorem 4.2 *Any r.e. precomplete lattice is uniformly dense.*

Note that the theorem needs neither distributivity nor boundedness.

Here is the plan: given a recursive F , we are going to craft a partial recursive f . In other words, the construction below will effectively associate to an index e for F an index $c(e)$ for f . An index $t(c(e))$ for some F' making f total modulo \sim is then effective in $c(e)$. By the second recursion theorem there is an e_0 indexing the same function as $t(c(e_0))$. For that e_0 we have $F' \simeq F$. We may therefore assume from the outset that F makes f total modulo \sim .

Lastly, we put $D(a, b) = a \vee (F(a, b) \wedge b)$, which will be the desired extensional density function for L .

We fix effective enumerations $(\sim_n)_{n \in \omega}$ and $(\lesssim_n)_{n \in \omega}$ of \sim and \lesssim , respectively, that satisfy the following:

- for each $n \in \omega$, field $\sim_n = \text{field } \lesssim_n$ is a finite nonempty subset of field L ;
- \sim_n is an equivalence relation;
- $\sim_n \subseteq \sim_{n+1}$ and $\lesssim_n \subseteq \lesssim_{n+1}$;
- $\sim = \bigcup_{n \in \omega} \sim_n$ and $\lesssim = \bigcup_{n \in \omega} \lesssim_n$.

Construction The construction of f proceeds in stages. The following happens at Stage n .

- (C1) Suppose that $a, b \in \text{field } \sim_n$ and $f(a, b)$ has not yet been defined. Let $a_0, b_0 \in \text{field } \sim_n$ be the minimal such that $a_0 \sim_n a$ and $b_0 \sim_n b$. Put $f(a, b) = F(a_0, b_0)$ unless $(a, b) = (a_0, b_0)$.
- (C2) Suppose that $a \lesssim_n b$, $f(a, b)$ has not yet been defined and $a \vee (F(a, b) \wedge b) \lesssim_n a$. Put $f(a, b) = b$.
- (C3) Suppose that $a \lesssim_n b$, $f(a, b)$ has not yet been defined and $b \lesssim_n a \vee (F(a, b) \wedge b)$. Put $f(a, b) = a$.

Claim 1 *If $a \lesssim b$, then $a \lesssim D(a, b) \lesssim b$. In particular, $a \sim b$ implies $a \sim D(a, b) \sim b$.*

Proof This holds by virtue of the definition $D(a, b) = a \vee (F(a, b) \wedge b)$ regardless of the value of $F(a, b)$. ⊥

Claim 2 *$f(a, b)$ is defined unless both a and b are minima of their respective \sim -equivalence classes.*

Proof Clause (C1) takes care of this. \dashv

Claim 3 If $f(a, b)$ is defined via clause (C2) or (C3), then $a \sim b$.

Proof Suppose that $f(a, b)$ is defined via clause (C2). We must then have $a \lesssim b$, $a \vee (F(a, b) \wedge b) \lesssim a$, and $F(a, b) \sim f(a, b) = b$, so $a \vee b \lesssim a$; hence $a \sim b$.

Clause (C3) is treated similarly. \dashv

Claim 4 D is extensional with respect to \sim .

Proof That the \sim -equivalence class of $D(a, b)$ only depends on those of a and b follows from Claim 1 for the case $a \sim b$. We may therefore assume $a \not\sim b$. This implies, by Claim 3, that the only way to define $f(a', b')$ for $a' \sim a$ and $b' \sim b$ is via clause (C1).

Assume that a_0 and b_0 are the minima of the \sim -equivalence classes of a and b , respectively. We show by induction on $a + b$ that $F(a, b) \sim F(a_0, b_0)$ for all $a \sim a_0$ and $b \sim b_0$. Suppose $(a, b) \neq (a_0, b_0)$. By Claim 2, $f(a, b)$ is defined—via clause (C1). So $f(a, b) = F(a', b')$ where $a' \sim a$, $b' \sim b$, and $a' + b' < a + b$. Accordingly,

$$F(a, b) \sim f(a, b) = F(a', b') \sim F(a_0, b_0)$$

with the last equivalence holding by the induction hypothesis. Thus, we may conclude that $D(a, b) \sim D(a_0, b_0)$. \dashv

Claim 5 If $a \not\lesssim b$, then $a \not\lesssim D(a, b) \not\lesssim b$.

Proof In view of Claim 1 it will suffice to exclude the situations $a \sim D(a, b)$ and $D(a, b) \sim b$.

Suppose $a \sim D(a, b) = a \vee (F(a, b) \wedge b)$. Let a_0, b_0 be the minima of a 's and b 's \sim -equivalence classes. Then a_0, b_0 also are minima of any \sim_n -equivalence classes they belong to. Thus clause (C1) cannot define $f(a_0, b_0)$. By Claim 3 neither can (C2) nor (C3). Yet clause (C2) will sooner or later define $f(a_0, b_0)$ if nothing else does, a contradiction.

$D(a, b) \sim b$ is outruled in a similar fashion. \dashv

Claims 1, 4, and 5 amount to a proof of Theorem 4.2.

Corollary 4.3 For r.e. consistent T extending EA the lattice Σ_n/T is uniformly dense.

Remark 4.4 Using $\text{True}_n(\dots)$ as in Remark 3.7, one can bring down to p-time the complexity of any recursive function with values in Σ_n/T . The density functions for Σ_n/T obtained through Theorem 4.2 however are already polynomial time because in Σ_n/T totalization works by substitution (see the hint to Example 4.1) as does, for that matter, the second recursion theorem.

Corollary 4.5 The finitely axiomatized sequential theories are uniformly dense with respect to the interpretability preordering \triangleleft . The density function can be taken to be p-time.

Open Question 4.6 Are Σ_1/S_2^1 and/or $\exists\Sigma_1^1/S_2^1$ uniformly dense?

Harvey Friedman shows in his Tarski lectures that the interpretability preordering on arbitrary finitely axiomatized theories of predicate logic has an effective density function. Is this ordering uniformly dense?

Open Question 4.7 Do Lindenbaum sentence algebras admit uniform density functions that are monotone with respect to provable implication in (ideally) both arguments? What about precomplete r.e. lattices?

5 Archaeology

In this section we provide assorted background material that makes our construction of C_A meaningful. We will sketch how the main ingredient of our formula C_A , to wit, the formula $C_A^\circ := \forall x (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp)$, can be viewed as either a unique Rosser or Gödel fixed point. We first discuss the Rosser construction.

A standard way to produce independent sentences is the Rosser construction, invented by J. Barkley Rosser. The original paper is Rosser [16]. Rosser’s construction has some extra good properties. The construction is verifiable in PA, and, after some careful inspection, even in EA.¹ A second point is that Rosser’s argument works for a very wide class of theories including the recursively enumerable extensions of the Tarski–Mostowski–Robinson theory R. Finally, the sentence produced by his construction, the Rosser sentence, is Σ_1 or Π_1 , more specifically, $\exists \Pi_1^b$ or $\forall \Sigma_1^b$.

Can we use the original Rosser construction to obtain independent sentences in a uniform way? This does not look very promising: the sentences delivered by that construction are quite sensitive to implementation details. For example, suppose that we use a standard fixed point construction to obtain a Rosser sentence R_A for $PA + A$ and a Rosser sentence $R_{A'}$ for $PA + A'$. Suppose further that A and A' are PA-provably equivalent. Then, R_A and $R_{A'}$ need not be PA-provably equivalent. The *intensionality* of the Rosser construction has, for example, been studied in Guaspari and Solovay [4] and Voorbraak [29]. However, several variants of the Rosser construction have been considered in the literature, and among these we find one that is sufficiently uniform. This Rosser construction was introduced by Craig Smoryński. As we will see, this Rosser construction can also be viewed as a Gödel construction.

Consider an r.e. extension T of PA in the same language. Let $\tau := (T_n)_{n \in \omega}$ be a recursive sequence of theories so that $I\Sigma_1$ proves that

1. for all n and k , if $n < k$, then T_n is a subtheory of T_k ;
2. the union of the T_n is T ;
3. for each n , $T_{n+1} \vdash \text{con}(T_n)$.

We need the following definitions:

$\Box_\tau^* B$ stands for $\exists x \Box_{T_x} B$; note that $\Box_\tau^* B$ is provably equivalent to $\Box_T B$.

Suppose that C and D are of the respective forms $\exists x C_0(x)$ and $\exists y D_0(y)$; then

$$C < D := \exists x (C_0(x) \wedge \forall y \leq x \neg D_0(y))$$

and

$$C \leq D := \exists x (C_0(x) \wedge \forall y < x \neg D_0(y)).$$

Note that $\Box_\tau^* B < \Box_\tau^* C$ is $I\Sigma_1$ -provably equivalent to $\Box_T B \wedge \neg(\Box_\tau^* C \leq \Box_\tau^* B)$. Thus, the formula $\Box_\tau^* B < \Box_\tau^* C$ is Δ_2 over $I\Sigma_1$. The case is similar for $\Box_\tau^* B \leq \Box_\tau^* C$.

The formula $\Box_\tau^* B < \Box_\tau^* C$ is equivalent over PA to $\exists x (\Box_{T_x} B \wedge \neg \Box_{T_x} C)$. It follows that the formula $\Box_\tau^* B < \Box_\tau^* \neg B$ is equivalent to $\Box_\tau^* B < \Box_\tau^* \perp$ which coincides

with the *Feferman predicate* for τ defined as

$$\Delta_\tau A := \Box_\tau^* A < \Box_\tau^* \perp.$$

We note that, over EA, $\Delta_\tau A$ is equivalent to $\exists x (\Box_{T_x} A \wedge \Diamond_{T_x} \top)$.

The Feferman predicate was introduced by Solomon Feferman in his classical paper [3]. The Feferman predicate is a sort of self-correcting provability predicate. It is related to trial and error predicates as studied in Putnam [15] and Jeroslow [6]. Feferman’s aim in introducing it was not just the study of ways to escape the second incompleteness theorem but also applications to the study of relative interpretability.

Here is the central insight. We write

- $\nabla_\tau A$ for $\neg \Delta_\tau \neg A$;
- $U \triangleright V$ for: there is a relative interpretation of V in U .

See, for example, [21], [3], Lindström [8] and Visser [25] for basic definitions concerning interpretations.

Theorem 5.1 *We have $(T + \nabla_\tau A) \triangleright (T + A)$.*

See [3] for the main ingredients of the proof. The basic idea of the result is that $\nabla_\tau A$ is a consistency statement of $T + A$. We can use the Henkin construction to build the desired interpretation.

We now consider the specialized sequences $v_A := (Ar_n + A)_{n \in \omega}$ for the theories $PA + A$. We simply write \Box_A^* for $\Box_{v_A}^*$, and so on.

By the Gödel fixed point lemma, we can find a sentence R_A , such that

$$PA \vdash R_A \leftrightarrow \neg (\Box_A^* R_A < \Box_A^* \neg R_A).$$

Thus R_A is a Rosser sentence for the \Box_A^* . By our previous remarks, the sentence R_A is also a Gödel sentence for the Feferman predicate Δ_A , that is:

$$PA \vdash R_A \leftrightarrow \neg \Delta_A R_A.$$

Smoryński investigates R_A in his paper [19]. That paper was inspired by a study of a variant of the Rosser construction in the context of set theory by Kenneth McAloon [9].

Theorem 2.1 in [19] implies that R_A is, up to provable equivalence, unique over $PA + A$. By a minor addition to Smoryński’s argument, one can show that the mapping $A \mapsto (A \wedge R_A)$ preserves PA-provable equivalence. Shavrukov [18] shows that uniqueness can fail under a choice of stratification sequence different from $(Ar_n)_{n \in \omega}$.

Smoryński also shows that R_A is independent over $PA + A$, provided that $PA + A$ is consistent. As we will see, Smoryński’s Rosser sentence R_A is $(PA + A)$ -provably equivalent to the sentence $\forall x (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp)$. So the independence of R_A also follows from our Lemma 3.6.

Since the Feferman–Smoryński predicate explicates a notion of provability, it can be studied modally. This study was taken up in Montagna [10], Visser [24], and Shavrukov [18]. The latter paper studies the Feferman predicate over PA based on the sequence Ar_n with conclusions translatable to the hierarchy $Ar_n + A$. Thus [18] contains an alternative, modal, proof of the uniqueness of R_A .

Recall that $C_A^\circ := \forall x (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp)$. We show that C_A° is a Gödel sentence for Δ_A .

Theorem 5.2 ([18, Exercise 2.7]) C_A° is a Gödel sentence of Δ_A over $PA + A$.

Proof We have, using Löb’s theorem in the third step,

$$\begin{aligned} \text{PA} + A \vdash \Delta_A C_A^\circ &\rightarrow \exists x (\Box_{A,x} \forall y (\Box_{A,y} \Box_{A,y} \perp \rightarrow \Box_{A,y} \perp) \wedge \Diamond_{A,x} \top) \\ &\rightarrow \exists x (\Box_{A,x} (\Box_{A,x} \Box_{A,x} \perp \rightarrow \Box_{A,x} \perp) \wedge \Diamond_{A,x} \top) \\ &\rightarrow \exists x (\Box_{A,x} \Box_{A,x} \perp \wedge \Diamond_{A,x} \top) \\ &\rightarrow \neg C_A^\circ. \end{aligned}$$

We treat the other direction. Suppose $A \in \Sigma_{n+3}$. We work in $\text{PA} + A$. Suppose $\neg C_A^\circ$, that is,

$$\exists x (\Box_{A,x} \Box_{A,x} \perp \wedge \Diamond_{A,x} \top).$$

Clearly, it follows that $\exists x (\Box_{A,x} \forall y \geq x (\Box_{A,y} \perp \wedge \Diamond_{A,x} \top))$, and hence:

$$\exists x (\Box_{A,x} \forall y \geq x (\Box_{A,y} \Box_{A,y} \perp \rightarrow \Box_{A,y} \perp) \wedge \Diamond_{A,x} \top). \tag{i}$$

We note that we may assume that $x \geq n$, since for any standardly finite k , $\Box_{A,k} \Box_{A,k} \perp$ implies \perp . Hence, by the fact that $\text{Ar}_x + A$ proves Σ_1 -reflection for $\text{Ar}_y + A$, for $y < x$, we find

$$\exists x (\Box_{A,x} \forall y < x (\Box_{A,y} \Box_{A,y} \perp \rightarrow \Box_{A,y} \perp) \wedge \Diamond_{A,x} \top). \tag{ii}$$

Combining (i) and (ii), we find

$$\exists x (\Box_{A,x} \forall y (\Box_{A,y} \Box_{A,y} \perp \rightarrow \Box_{A,y} \perp) \wedge \Diamond_{A,x} \top). \tag{iii}$$

Of course (iii) is $\Delta_A C_A^\circ$.

Thus, we have shown that $\text{PA} + A \vdash C_A^\circ \leftrightarrow \neg \Delta_A C_A^\circ$. □

So, C_A° is modulo PA-provable equivalence Smoryński’s Rosser sentence for $\text{PA} + A$.

Open Question 5.3 Our proof of the extensionality of $A \mapsto C_A$ as well as that of unprovability of C_A go through, with minor modifications, for any stratification sequence τ for PA satisfying our conditions. The consistency of C_A° with $\text{PA} + A$ is the only element of Theorem 3.1 that ostensibly depends on $\tau = (\text{Ar}_n)_{n \in \omega}$ (or, more generally, on the “fast-growing” property of τ that each level proves enough reflection for the previous ones).

This makes us wonder if there exists a consistent theory of the form $\text{PA} + A$ together with a stratification sequence $\tau = (T_n)_{n \in \omega}$ such that $\text{PA} + A$ refutes $\forall x (\Box_{A,T_x} \Box_{A,T_x} \perp \rightarrow \Box_{A,T_x} \perp)$.

A similar question can be asked of Theorem 5.2.

We end this section by showing that C_A° is an Orey sentence of $\text{PA} + A$.

Consider any theory T . A sentence O in the language of T is an *Orey sentence* of T if $T \triangleright (T + O)$ and $T \triangleright (T + \neg O)$. Note that the negation of an Orey sentence is an Orey sentence. An Orey sentence O of T is clearly independent of T . Neither an Orey sentence nor its negation add interpretability strength to the given theory.

The idea of Orey sentences was introduced by Orey [13], who also provided the first known Orey sentence for PA. There are many salient natural Orey sentences. Two well-known examples are the parallel axiom over a suitable version of neutral geometry and the continuum hypothesis over ZFC. For essentially reflexive sequential theories, the Gödel sentence of a Feferman predicate for the theory is an Orey sentence—see below.

We will show that C_A° is an Orey sentence of $\text{PA} + A$. This sentence is still meta-mathematical and does involve coding, but it is, at least, self-reference-free. Since

C_A° is a Gödel sentence of Δ_A , the desired insight is immediate by the following theorem. (This theorem was also proved in [24].)

Theorem 5.4 *Consider a consistent theory T given by a sequence τ satisfying the conditions given above. Then any Gödel sentence of Δ_τ is an Orey sentence for T .*

Proof Let G be a Gödel sentence of Δ_τ . We have

$$\begin{aligned} T + G &\vdash T + \nabla_\tau \neg G \\ &\triangleright T + \neg G, \\ T + \neg G &\vdash T + \Delta_\tau G \\ &\vdash T + \nabla_\tau G \\ &\triangleright T + G. \end{aligned}$$

In the second step of the second proof, we use $T \vdash \Delta_\tau \neg A \rightarrow \neg \Delta_\tau A$. Thus, we have both $(T + G) \triangleright (T + G)$, by the identity interpretation, and $(T + \neg G) \triangleright (T + G)$. So, using a disjunctive interpretation, we find $T \triangleright (T + G)$. Similarly, $T \triangleright (T + \neg G)$. \square

Note

1. A modified argument even works in $I\Delta_0 + \Omega_1$. The basic idea of this argument is due to Švejdar [20]. For the verification that Švejdar's assumptions are fulfilled, see Verbrugge and Visser [22].

References

- [1] Beklemishev, L. D., "Reflection schemes and provability algebras in formal arithmetic" (in Russian), *Uspekhi Matematicheskikh Nauk*, vol. 60, no. 2 (2005), pp. 3–78; English translation in *Russian Mathematical Surveys*, vol. 60 (2005), pp. 197–268. [Zbl 1097.03054](#). [MR 2152943](#). [DOI 10.1070/RM2005v060n02ABEH000823](#). [570](#), [572](#)
- [2] Buss, S. R., *Bounded Arithmetic*, vol. 3 of *Studies in Proof Theory: Lecture Notes*, Bibliopolis, Naples, Italy, 1986. [Zbl 0649.03042](#). [MR 0880863](#). [571](#), [574](#)
- [3] Feferman, S., "Arithmetization of metamathematics in a general setting," *Fundamenta Mathematicae*, vol. 49 (1960/61), pp. 35–92. [Zbl 0095.24301](#). [MR 0147397](#). [578](#)
- [4] Guaspari, D., and R. M. Solovay, "Rosser sentences," *Annals of Mathematical Logic*, vol. 16 (1979), pp. 81–99. [Zbl 0426.03062](#). [MR 0530432](#). [DOI 10.1016/0003-4843\(79\)90017-2](#). [577](#)
- [5] Hájek, P., and P. Pudlák, *Metamathematics of First-Order Arithmetic*, Perspectives in Mathematical Logic, Springer, Berlin, 1993. [Zbl 0889.03053](#). [MR 1219738](#). [DOI 10.1007/978-3-662-22156-3](#). [570](#), [571](#), [572](#)
- [6] Jeroslow, R. G., "Experimental logics and Δ_2^0 -theories," *Journal of Philosophical Logic*, vol. 4 (1975), pp. 253–67. [Zbl 0319.02024](#). [MR 0505406](#). [578](#)
- [7] Leivant, D., "The optimality of induction as an axiomatization of arithmetic," *Journal of Symbolic Logic*, vol. 48 (1983), pp. 182–84. [Zbl 0515.03018](#). [MR 0693260](#). [DOI 10.2307/2273332](#). [572](#)
- [8] Lindström, P., *Aspects of Incompleteness*, 2nd ed., vol. 10 of *Lecture Notes in Logic*, Association for Symbolic Logic, Urbana, Ill., A K Peters, Natick, Mass., 2003. [Zbl 1036.03002](#). [MR 2014250](#). [578](#)
- [9] McAloon, K., "Formules de Rosser pour ZF," *Comptes Rendus hebdomadaires des Séances de l'Académie des Sciences (Série A)*, vol. 281 (1975), pp. A669–72. [Zbl 0333.02046](#). [MR 0384543](#). [578](#)

- [10] Montagna, F., “On the algebraization of a Feferman’s predicate (the algebraization of theories which express Theor; X),” *Studia Logica*, vol. 37 (1978), pp. 221–36. [Zbl 0398.03054](#). [MR 0515168](#). [DOI 10.1007/BF02124724](#). 578
- [11] Montagna, F., and A. Sorbi, “Universal recursion-theoretic properties of r.e. preordered structures,” *Journal of Symbolic Logic*, vol. 50 (1985), pp. 395–406. [Zbl 0578.03026](#). [MR 0793120](#). [DOI 10.2307/2274228](#). 574
- [12] Ono, H., “Reflection principles in fragments of Peano arithmetic,” *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, vol. 33 (1987), pp. 317–33. [Zbl 0607.03017](#). [MR 0913295](#). [DOI 10.1002/malq.19870330407](#). 571
- [13] Orey, S., “Relative interpretations,” *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, vol. 7 (1961), pp. 146–53. [Zbl 0121.25501](#). [MR 0146082](#). 579
- [14] Pour-El, M. B., and S. Kripke, “Deduction-preserving ‘recursive isomorphisms’ between theories,” *Fundamenta Mathematicae*, vol. 61 (1967), pp. 141–63. [Zbl 0174.02003](#). [MR 0252226](#). 573, 574
- [15] Putnam, H., “Trial and error predicates and the solution to a problem of Mostowski,” *Journal of Symbolic Logic*, vol. 30 (1965), pp. 49–57. [Zbl 0193.30102](#). [MR 0195725](#). 578
- [16] Rosser, J. B., “Extensions of some theorems of Gödel and Church,” *Journal of Symbolic Logic*, vol. 1 (1936), pp. 87–91. [Zbl 0015.33802](#). 571, 577
- [17] Selivanov, V., “Positive structures,” pp. 321–50 in *Computability and Models: Perspectives East and West*, edited by S. B. Cooper and S. S. Goncharov, Kluwer Academic/Plenum, New York, 2003. [MR 2043604](#). 574
- [18] Shavrukov, V. Yu., “A smart child of Peano’s,” *Notre Dame Journal of Formal Logic*, vol. 35 (1994), pp. 161–85. [Zbl 0815.03011](#). [MR 1295557](#). [DOI 10.1305/ndjfl/1094061859](#). 570, 578
- [19] Smoryński, C., “Arithmetic analogues of McAloon’s unique Rosser sentences,” *Archive for Mathematical Logic*, vol. 28 (1989), pp. 1–21. [Zbl 0663.03049](#). [MR 0987763](#). [DOI 10.1007/BF01624079](#). 570, 578
- [20] Švejdar, V., “Modal analysis of generalized Rosser sentences,” *Journal of Symbolic Logic*, vol. 48 (1983), pp. 986–99. [Zbl 0543.03010](#). [MR 0727788](#). [DOI 10.2307/2273663](#). 580
- [21] Tarski, A. A., *Undecidable Theories*, in collaboration with A. Mostowski and R. M. Robinson, Studies in Logic and the Foundations of Mathematics, North-Holland, Amsterdam, 1953. [Zbl 0053.00401](#). [MR 0058532](#). 569, 571, 578
- [22] Verbrugge, R., and A. Visser, “A small reflection principle for bounded arithmetic,” *Journal of Symbolic Logic*, vol. 59 (1994), pp. 785–812. [Zbl 0814.03039](#). [MR 1295969](#). [DOI 10.2307/2275908](#). 580
- [23] Visser, A., “Numerations, λ -calculus & arithmetic,” pp. 259–84 in *To H. B. Curry: Essays on Combinatory Logic, Lambda Calculus and Formalism*, edited by J. P. Seldin and J. R. Hindley, Academic Press, London, 1980. [MR 0592807](#). 574
- [24] Visser, A., “Peano’s smart children: A provability logical study of systems with built-in consistency,” *Notre Dame Journal of Formal Logic*, vol. 30 (1989), pp. 161–96. [Zbl 0686.03033](#). [MR 1050516](#). [DOI 10.1305/ndjfl/1093635077](#). 578, 580
- [25] Visser, A., “Can we make the second incompleteness theorem coordinate free?” *Journal of Logic and Computation*, vol. 21 (2011), pp. 543–60. [Zbl 1262.03123](#). [MR 2823429](#). [DOI 10.1093/logcom/exp048](#). 575, 578
- [26] Visser, A., “What is sequentiality?” pp. 229–69 in *New Studies in Weak Arithmetics*, edited by P. Cégielski, Ch. Cornaros and C. Dimitracopoulos, vol. 211 of *CSLI Lecture Notes*, CSLI Publications/Presses universitaires du Pôle de Recherche et d’Enseignement Supérieur de Paris-Est, Stanford/Paris, 2013. 570
- [27] Visser, A., “The arithmetics of a theory,” to appear in *Notre Dame Journal of Formal Logic*, preprint, 2012, <http://dspace.library.uu.nl/handle/1874/234711>. 570

- [28] Visser, A., “Peano Corto and Peano Basso: A study of local induction in the context of weak theories,” *Mathematical Logic Quarterly*, vol. 60 (2014), pp. 92–117. DOI [10.1002/malq.201200102](https://doi.org/10.1002/malq.201200102). 570
- [29] Voorbraak, F., “A simplification of the completeness proofs for Guaspari and Solovay’s R,” *Notre Dame Journal of Formal Logic*, vol. 31 (1989), pp. 44–63. Zbl [0712.03010](https://zbmath.org/?q=ri:0712.03010). MR [1043791](https://www.ams.org/mathscinet-getitem?mr=1043791). DOI [10.1305/ndjfl/1093635332](https://doi.org/10.1305/ndjfl/1093635332). 577

Acknowledgments

The question concerning the uniform density of Lindenbaum algebras was raised by Sy Friedman in conversation. Sy’s question was inspired by the idea that perhaps the consistency operator is, in some sense, the least way to strengthen theories. This is not so, as is corroborated by the results of this paper. Both authors thank Lev Beklemishev for his consulting services. Visser’s work was supported by the Templeton Foundation through his participation in the CRM Infinity Project (Templeton Foundation project #13152).

Shavrukov
Nijenburg 24
1081GG Amsterdam
The Netherlands
v.yu.shavrukov@gmail.com

Visser
Department of Philosophy
Utrecht University
Janskerkhof 13A
3512BL Utrecht
The Netherlands
a.visser@uu.nl