

Universal metric properties of bifurcations of endomorphisms

B Derrida†, A Gervois‡ and Y Pomeau‡

† Institut Laue-Langevin, BP156, 38042 Grenoble Cédex, France

‡ C.E.A. Centre d'Etudes Nucléaires de Saclay, Division de la Physique, BP2, 91190 Gif-sur-Yvette, France

Received 6 March 1978

Abstract. Endomorphisms of the real axis with one extremum have some universal metric properties which depend only on their analytic dependence near the extremum (bifurcation velocity, reduction parameter). It is shown how this problem is similar to the renormalisation problem, and how the bifurcation velocity may be derived from a fixed-point theory.

Several renormalisation methods are discussed for the reduced endomorphism

$$\mathcal{T}_a: x \rightarrow \mathcal{T}_a(x) = 1 - a|x|^{1+\epsilon}.$$

They all give the same (exact) result for small ϵ .

1. Introduction

Because of their relative simplicity, endomorphisms of the real axis have been extensively studied, and a number of results have been obtained for some classes of functions (May 1976, Li and Yorke 1975).

In this paper, we intend to give some results for endomorphisms (or mappings) T_a of an interval $I \subset \mathbb{R}$ depending on one parameter a ; the function T_a is continuous, convex with a unique maximum (or critical point) on I . It is convenient to assume that the derivative $(d/dx)T_a(x)$ exists except perhaps at the critical point.

In most theoretical works which have been devoted to this class of mappings, supplementary conditions have been introduced, but it appears from numerical calculations that they are not essential. Some universal features appear which are independent of the shape of the function T_a . Roughly, they are of two kinds. Some of them depend only on the existence of a unique maximum and can be called structural, e.g. the order of occurrence of periods (Metropolis *et al* (1973) and the internal similarity (Derrida *et al* 1978, referred to hereafter as I). The other universal results may be called metric (Feigenbaum 1977 a,b) and depend on the shape of T_a near the extremum c .

For example, if $|T_a(x) - T_a(c)| \sim |x - c|^{1+\epsilon}$, the bifurcation velocity and reduction parameter (to be defined in § 2.3) depend on the coefficient ϵ only. Feigenbaum has already pointed out the connection between this property of universality and the idea of renormalisation group theory in critical phenomena.

We shall focus here on the metric properties. Some theoretical results may be obtained, and an extension of the renormalisation group approach is possible.

In § 2 we recall some definitions for periods and results obtained by Metropolis *et al* (1973) and by the authors. In § 3 we enlarge Feigenbaum's considerations and show how the bifurcation problem can be considered as being similar to the phase transition problem. Critical exponents may be defined, but there are an infinite number of critical points (and exponents). In § 4 we try to get these exponents by approximate renormalisation methods and compare their rapidity of convergence with the exact (numerical) results. Because of universality, we restrict ourselves to the reduced endomorphism \mathcal{T}_a of the interval $I = (-1, +1)$, where a is a parameter:

$$\mathcal{T}_a: x \rightarrow \mathcal{T}_a(x) = 1 - a|x|^{1+\epsilon}. \tag{1.1}$$

The piecewise linear mapping ($\epsilon = 0$) is already renormalised, and for small ϵ all methods give the same first-order expression, which can be expected to be in good agreement with the exact result. For the quadratic case ($\epsilon = 1$) the simplest method gives the exact result within an error of 20%, and more sophisticated ways reach an accuracy better than 1% with not too much calculation. In § 5 we derive directly for small ϵ the first-order correction to the piecewise linear case $\epsilon = 0$, and obtain the same results as by renormalisation theory in § 4. Even when we put $\epsilon = 1$ (quadratic case) into the first-order formulae, we get the exact result within an error of 2%!

2. Generalities on periodic points

2.1. Definitions

A point x belongs to a *period* k if

$$T_a^{(k)}(x) = x \tag{2.1a}$$

with

$$T_a^{(l)}(x) \neq x \quad \text{if } l < k \tag{2.1b}$$

where by definition

$$T_a^{(1)}(x) = T_a(x) \tag{2.2a}$$

and

$$T_a^{(l)}(x) = T_a(T_a^{(l-1)}(x)) = x_l \tag{2.2b}$$

is the l th iterate of x . The iterates $x (= x_0 = x_k), x_1, \dots, x_{k-1}$ form a *sequence*.

The period is *stable* if for a neighbourhood U of x_0

$$y \in U \Rightarrow |T_a^{(l)}(x_0) - T_a^{(l)}(y)| \rightarrow 0$$

for l sufficiently large.

When the derivative $(d/dx)T_a(x) = T'_a(x)$ exists, the above condition is equivalent to

$$-1 < s < 1 \tag{2.3}$$

with

$$s = T'_a(x_0)T'_a(x_1) \dots T'_a(x_{k-1}). \tag{2.4}$$

When a increases, a stable period appears for the value a_s of parameter a for which

$s = +1$ and disappears for the value a_u of a for which $s = -1$. Then a period $2k$ becomes stable; a_u is by definition a *bifurcation point* on the parameter axis. Interval (a_s, a_u) is the *stability zone*.

For a given value a of the parameter, three situations may occur: (i) no stable period exists; (ii) one stable period exists; (iii) two or more stable period exist. In the latter case it may happen that two points attracted by two different stable periods may be as near to each other as we like, and many complications occur. Presumably these difficulties may be avoided in general when one restricts oneself to the class of functions defined in the introduction, though Metropolis *et al* (1973) give an example where two stable periods coexist. At least it is believed that the number of these 'spurious' periods is finite, and we may exhibit endomorphisms where only one stable period exists. By definition a 'spurious' period is not found in the construction of Metropolis *et al*. This is due to the fact that, when the mapping is continuously changed, the critical point never belongs to this spurious period. For example, the piecewise linear mapping (Derrida and Pomeau 1979)

$$T_\lambda(x) = \begin{cases} \lambda x & x < 1 \\ \lambda & 1 < x < r \\ \lambda(1+r) - \lambda x & r < x \end{cases}$$

and its peculiar case $r = 1$ that we studied in I have no 'spurious' periods.

In the following, we shall assume that situation (iii) above does not occur.

2.2. Superstability and MSS sequences

The period k is *superstable* if $s = 0$. It corresponds to a value of the parameter a for which the critical point belongs to the period. Choosing then $c = x_1$, the $(k - 1)$ iterates of c may be represented by a formal sequence Q of $(k - 1)$ characters R (right) or L (left)

$$Q = \sigma_1 \dots \sigma_{k-1} \tag{2.5a}$$

where

$$\sigma_i = \begin{cases} R & \text{if } T_a^{(i)}(c) > c \\ L & \text{if } T_a^{(i)}(c) < c \end{cases} \tag{2.5b}$$

For example, the fixed point is represented by a sequence with 0 character (the 'blank' sequence b), the period 2 by the only character R , and so on.

All sequences of R and L characters do not represent a period. For example, the first two symbols are always RL . When a sequence Q of $(k - 1)$ characters represents an actual superstable period k , we say that the sequence Q is *allowed*. Its *length* is $k - 1$. For $k = 5$, we have only three allowed sequences: RL^3 , RL^2R and RLR^2 .

These sequences may be visualised by setting on an axis the iterates x_1, x_2, \dots of the extremum (see figure 1). When a goes from a_s (beginning of stability) to a_u (end of stability) and to larger values passing through the superstable value, *the ordering of the iterates is unchanged*. Superstable periods give then a good picture of the whole stability zone, hence their importance. We shall use them indifferently to denote either the whole corresponding stability zone or the point of superstability in the parameter space.

Except for periods 1, 2 and 3, many sequences exist corresponding to periods with the same number of points. The ordering of the point of the period and the location of the stability zone in the parameter space depend on the sequence. This is shown for periods with five points on figure 1.




Figure 1. Graphic representation of the iterates of the critical point 1 on an axis for the three allowed sequences with period 5. The last line shows for these sequences the non-intersecting stability zones on the parameter axis.

These sequences Q have been studied by Metropolis *et al* (1973) and by the present authors in I. We shall call them MSS sequences. We recall here the results which will be useful later on. From now on, we shall denote as a_P, a_Q, \dots the value of the parameter for which the periods P, Q, \dots are superstable.

(i) If P and Q are allowed sequences which are superstable for the values a_P and a_Q respectively of the parameter, we say that

$$P < Q \quad \text{whenever } a_P < a_Q. \tag{2.6}$$

When a increases, the MSS sequences always appear in the same order, independently of T_a (*universal sequences*). For example,

$$b < R < RLR < RLR^3LR < \dots < RL < RL^2RL < \dots$$

and

$$RL^{n_1} < RL^{n_2} \quad \text{if } n_1 < n_2.$$

Notice that, as shown in the above example, the first sequence after a given sequence is its harmonic: for instance RLR (period 4) follows R (period 2), RL^2RL (period 6) follows RL (period 3), and so on \dots

Between two given sequences P_1 and P_2 , we may always construct the sequence P_3 of minimal length. Thus the construction and ordering of all MSS sequences is possible (Metropolis *et al* 1973).

(ii) One may define on the set of allowed sequences an inner composition law denoted by $*$. Q and P being two allowed sequences corresponding to periods k and l , $Q * P$ is an allowed sequence of period kl . More precisely, if $P = \sigma_1\sigma_2 \dots \sigma_{l-1}(\sigma_i = R, L)$,

$$Q * P = Q\tau_1 Q\tau_2 \dots Q\tau_{l-1} Q \tag{2.7}$$

where

$$\tau_i \begin{cases} = \sigma_i & \text{if } Q \text{ has an even number of } R \text{ characters} \\ \neq \sigma_i & \text{otherwise} \end{cases} \quad (2.8)$$

For example, the harmonic of Q is $Q * R$.

By construction, the law $*$ is *not commutative*, but is associative with neutral element b (corresponding to period 1). The sequence $Q * P (Q \neq b)$ is said to be *factorisable* or *non-primary*. The number of non-primary sequences is small compared with the number of primary sequences (Derrida *et al* 1977, 1978).

One important consequence is the law of *internal similarity*. For given Q , the mapping $Q * : P \rightarrow Q * P$ maps the whole set of MSS sequences into one of its parts and preserves the ordering relation. The set of all sequences between b and RL^N (for every N) has the same structure as the set of sequences between Q and $Q * RL^N$, and conversely every $Q' (Q < Q' < Q * RL^N)$ can be written in the form $Q' = Q * P$. This result is to be related to the Sarkovskii theorem (Sarkovskii 1964, Stefan 1977, Cosnard and Eberhard 1977) and is of great importance in showing the connection with renormalisation. A similar property does not exist for the mapping $*P, *P : Q \rightarrow Q * P$, so we shall not deal with it any further.

With the graphic representation, it is easy to see what the non-primary sequences correspond to. Let us show it in an example. For $k = 6$, there exist two non-primary sequences $P_1 = R * RL = RLR^3$ and $P_2 = RL * R = RL^2RL$ which are represented in figure 2. For P_1 (figure 2a) we may draw a limit between points of the period numbered 5 and 4; by mapping T_a at every iteration, this limit is crossed. On a large scale—i.e. if one assimilates each of the clusters 1–3–5 and 4–6–2 to a single point—the dynamics is that of period 2; on a small scale—i.e. in each of the clusters 1–3–5 and 4–6–2—the dynamics is that of period 3. For sequence P_2 , there are two separations (figure 2b); on a large scale the dynamics is that of period 3, and on a small scale that of period 2. More generally, if $P = P_1 * \dots * P_n$, there are n successive scales. For instance, if $P = R * RL * R$ (figure 2c), the dynamics at large, medium and small scale is that of period 2, 3 and 2 respectively.




Figure 2. Different scales in non-primary sequences: (a) $P_1 = R * RL$; (b) $P_2 = RL * R$; (c) $P = R * RL * R$.

(iii) It is possible to recognise whether a given sequence is allowed or not and to compare it—in the sense of (2.6)—with other sequences without constructing all sequences with a smaller period. The classification has been found—because of the universality property (i)—by studying the simplest function T_a , i.e. the piecewise linear mapping \mathcal{L}_a

$$\mathcal{L}_a: x \rightarrow \mathcal{L}_a(x) = 1 - a|x| \tag{2.9}$$

(with $a \geq 1$, as no superstable period exists if $a < 1$).

The derivative of \mathcal{L}_a does not exist at $x = 1$, but the only difference with everywhere derivable endomorphisms T_a is that some non-primary periods cancel out.

We shall use in § 5 some results concerning the piecewise linear mapping \mathcal{L}_a which have been derived in I. We recall them here:

(a) The value \bar{a}_Q of the parameter a for which the sequence Q is superstable is the largest real root of the polynomial

$$\mathcal{P}_Q(x) = 1 + \alpha_{k-2}x + \dots + \alpha_{k-2} \dots \alpha_i x^{k-1-i} + \dots + \alpha_{k-2} \dots \alpha_0 x^{k-1}$$

where

$$\alpha_i = -\text{sgn } \mathcal{L}_a^{(i+1)}(0) \tag{2.10}$$

and

$$\text{sgn } x = \begin{cases} x/|x| & \text{if } x \neq 0 \\ 0 & \text{otherwise.} \end{cases} \tag{2.11}$$

The $\{\alpha_i\}$ are related in an obvious way to the R and L symbols.

(b) Because of the identity proved in I

$$\mathcal{P}_{Q*P}(x) = \mathcal{P}_Q(x)\mathcal{P}_P(x^k) \tag{2.12}$$

all sequences $Q * P$, Q being primary, define the same parameter \bar{a}_Q except when $Q = R$. The parameter \bar{a}_{R*P} (when $Q = R$) is $\bar{a}_P^{1/2}$.

Notice that, as $|(d/dx)\mathcal{L}_a(x)| = a$ except at the critical point, no period is stable except the fixed point at $a < 1$.

3. Metric universal properties†

3.1. Feigenbaum results

Feigenbaum (1977a, b) succeeded in calculating numerically the value a_n of the parameter a of the superstable harmonics R^{*n} (of period 2^n) for several functions T_a . He showed that:

(i) When n increases, the $\{a_n\}$ are an increasing sequence with limit a_* and

$$a_* - a_n \sim \delta_R^{-n}. \tag{3.1}$$

† In the bulk of the paper, the word ‘critical’ appears with two different meanings. In §§ 1, 2 and 6, a critical point for a mapping is a point where the Jacobian of the transformation vanishes (in one-dimensional mappings the point where the derivative $(d/dx)T_a(x)$ is zero). In §§ 3–5, critical points and critical exponents refer to the notion of critical points and critical exponents in phase transitions. These two definitions are traditional, and we maintain them since they do not appear at the same time in the paper. We hope that no ambiguity is possible.

More precisely, the ratio

$$\delta_n = (a_n - a_{n-1}) / (a_{n+1} - a_n) \tag{3.2}$$

has a limit δ_R (defined by equation (3.1)) when n goes to infinity, and it depends only on the shape near the extremum c (and not on the function). If

$$|T_a(x) - T_a(c)| \sim |x - c|^{1+\epsilon} \tag{3.3}$$

δ_R depends on ϵ only. We shall call δ_R the *bifurcation velocity*.

(ii) For large n , we have the asymptotic conjugation law

$$T_a^{2^n}(x) \underset{n \rightarrow \infty}{\sim} \gamma_R^{-n} \bar{\varphi}(\gamma_R^n x) \tag{3.4}$$

where $\bar{\varphi}$ is some (complicated) universal function, and γ_R is the *reduction parameter*. Both $\bar{\varphi}$ (up to a linear homomorphism) and γ_R do not depend on the details of T_a but only on the exponent ϵ . The renormalisation methods start from equation (3.4).

For $\epsilon = 1$ and different mappings T , Feigenbaum found $\delta_R = 4.669\ 201\ 609\ 03 \dots$, $\gamma_R = 2.502\ 907\ 875\ 0957 \dots$

3.2. Other numerical results

We repeated the calculations for more general sequences $P_1 * Q^{*n} * P_2$ and corresponding periods $p_1 q^n p_2$. We focused our attention on the parameter $a_{P_1 * Q^{*n} * P_2}$ in the particular transformation \mathcal{T}_a for several possible sequences Q and P_1 . Actually, for simplicity, we considered only sequences corresponding to small periods, i.e.

$$\begin{aligned} (a) \quad Q = R, \quad P_1 = b, RL, RL^2 \\ (b) \quad Q = RL, \quad P_1 = b, R, RL^2 \end{aligned}$$

and the case $P_2 = RL^\infty$, where RL^∞ is the limit for sequences RL^n when n is sufficiently large.

Though sequences R^{*n} behave in a slightly different manner than sequences Q^{*n} , from our results (see tables 1 and 2) it seems that properties (i) and (ii) of Feigenbaum can be extended in the following way:

(i) The parameter

$$a_n = a_{P_1 * Q^n * P_2} \tag{3.5a}$$

and the ratio

$$\delta_n = (a_n - a_{n-1}) / (a_{n+1} - a_n) \tag{3.5b}$$

tend to limits a_* and δ_Q respectively as $n \rightarrow \infty$. The limit value δ_Q of δ_n is a universal number which depends on Q and ϵ only.

(ii) When n becomes large, the asymptotic conjugation law (3.4) is true, γ and $\bar{\varphi}$ depending only on ϵ and Q . Sequence P_2 plays no role. This seems reasonable as, when n becomes large, the $p_1 q^n$ first characters of $P_1 * Q^{*n} * P_2$ depend on Q and P_1 only. The case $\epsilon = 1$ and $Q = R$ ($P = b$) is the one studied by Feigenbaum; for $Q = RL$ we find $\delta_{RL} = 55.247 \dots$, $\gamma_{RL} = 9.277\ 38 \dots$. The parameters γ_{RL} and δ_{RL} are much greater than γ_R and δ_R . We shall see in § 5 that this can be explained by computing δ_{RL} (and more generally δ_Q) by expansion near $\epsilon = 0$.

Table 1. Fixed-point and bifurcation velocity for the set of sequences $P * Q^{*n}$ ($n \rightarrow \infty$) for several MSS sequences Q and P in the quadratic case ($\epsilon = 1$) obtained numerically: (a) $Q = R$; (b) $Q = RL$.

Sequence P	(a) $Q = R, \epsilon = 1$		(b) $Q = RL, \epsilon = 1$	
	a_*	δ_R^{-1}	a_*	δ_{RL}^{-1}
b	1.401 155 189 09 ...	0.214 169 37 ...	1.786 440 255	0.018 1005
(a) RL , (b) R	1.779 818 07	0.214 1693	1.484 583 30	0.018 100
RL^2	1.942 173 796	0.214 169	1.942 550 324	0.018 100

Table 2. Fixed-point and bifurcation velocity for the set of sequences R^{*n} for different values of ϵ . For $\epsilon \leq 1$, we have plotted the values we get by using the 'first-order' approximation in ϵ (see § 5).

ϵ	Exact calculation		Methods $\epsilon \approx 0$	
	a_*	δ_R^{-1}	a_*	$\left(2 + \frac{\epsilon}{\varphi_*}\right)^{-1}$ for $\epsilon \leq 1$ $\varphi_* = a_* - 1$
0	1	0.5	1	0.5
0.1	1.1236	0.350	1.116	0.349
1	1.401 15	0.214 17	1.28	0.179
2	1.519 2995	0.167		
5	1.683 26	0.107 53		
9	1.772 6435	0.0808		
99	1.959 263 548	0.035		

Remarks

1. If $\epsilon = 0$ and $Q = R$, $\delta_R = 2$ (as $a_{Q^{*n} * P_2} = a_P^{1/2^n}$).
2. From definition (3.4), $\forall \epsilon$, if $Q = P * P$ then $\delta_Q = \delta_P^2$, and more generally $\delta_{P^{*n}} = (\delta_P)^n$, but if $Q = P_1 * P_2$, in general $\delta_{P_1 * P_2} \neq \delta_{P_1} \delta_{P_2}$. Equality is true at the lowest order when ϵ is small (and $P_i \neq R$). When ϵ is not small, we have the weaker property $\delta_{P_1 * P_2} = \delta_{P_2 * P_1}$. We shall prove it in § 3.4.

3.3. 'Critical' exponent ν_Q

One problem in bifurcation theory is to determine the stability zone for all periods, or equivalently to plot the length of the stable period k as a function of the parameter a : $k = k(a)$. From § 2.1 it is assumed to be unique for a given a . When no stable finite period exists, one may set $k(a) = \infty$, so that $k(a)$ is a step function with an infinite number of jumps. For example, the first periods when a increases are the harmonics R^{*n} ; the stability zones are adjacent and become smaller when n becomes large (see figure 3a). Setting $a_n = a_{R^{*n}}$ for the superstable value of the parameter as $a_* - a_n \sim \delta_R^{-n}$, we have on the n th step

$$k(a) = k(a_n) \sim (a_* - a_n)^{\nu_R} \tag{3.6a}$$

which diverges as a power at the limit value a_* , or more precisely

$$k(a) = (a_* - a)^{\nu_R} g_R(\ln(a_* - a)/\ln \delta_R) \tag{3.6b}$$




Figure 3. Study of the bifurcation points for the harmonics (a) R^{*n} and (b) $P * R^{*n}$ of period q . (For convenience, we took $P = RL$, $q = 3$.) Stability zones are adjacent. The upper figures plot the stable periods as a function of a . The superstable values of the parameter $a_i = a_{R^{*i}}$ and $b_j = a_{P * R^{*j}}$ are shown on the a axis. The lower curves show the length of the period as a function of a ; a_* and b_* are the limit parameters, and the period is multiplied by 2 at each harmonic.

where $g_R(x)$ is some discontinuous but periodic function: $g_R(x + 1) = g_R(x)$. The exponent ν_R depends on R and ϵ only. As $k(a_{n+1}) = 2k(a_n)$, we get for the critical exponent ν_R

$$\nu_R = -\ln 2 / \ln \delta_R. \tag{3.7}$$

For example, if $\epsilon = 1$, $\nu_R = -0.449\ 807$.

The same occurs if we consider the harmonics $P * R^{*n}$ of the sequence P of period p (figure 3b, with $P = RL$); setting $b_n = a_{P * R^{*n}}$ as $b_* - b_n \sim \delta_R^{-n}$ and $k(b_{n+1}) = 2k(b_n)$, we get the same power law $k(b) \sim (b_* - b_n)^{\nu_R}$, where ν_R is again given by (3.7).

The situation is more complicated if we consider the sequences $R^{*n} * P$, though the limit point is a_* , as for R^{*n} , and periods are multiplied by a factor of 2. The stability zones are no longer adjacent, and between each zone there is a complicated structure with an infinite number of discontinuities. Because of internal similarity, the fine structure between $c_n = a_{R^{*n} * P}$ and c_{n-1} is reproduced between c_{n+1} and c_n once the periods are multiplied by 2. A schematic plot of this situation can be found in figure 4. If again for the *superstable* values c_n of the parameter we have the power law $k(c_n) \sim (c_n - a_*)^{\nu_R}$, the true function $k(a)$ is of the form

$$k(a) \sim (a - a_*)^{\nu_R} g_R(\ln(a - a_*) / \ln \delta_R)$$

where g_R is periodic ($g_R(x + 1) = g_R(x)$) and has, in general, an infinite number of jumps. The above expression describes the behaviour of $k(a)$ near a_* , even for values of a which do not correspond to a superstable period.



Figure 4. Plot of $R^{*n} * RL$, $R^{*n} * RLR^\infty$ and $R^{*n} * RLR^8$ when $a \rightarrow a_*$ ($a > a_*$). The same patterns (multiplied by a factor of 2) may be found for each n , especially the infinite value of k for $R^{*n} * RLR^\infty$. The power law (3.6) is true for $P = RL$ (---) and $P = RLR^8$ (—) separately. For convenience, the scale is not respected.

More generally, we may consider all sequences $P_1 * Q^{*n} * P_2$, where $Q \neq R$ is some sequence of period q . All the superstable parameters $a_n = a_{P_1 * Q^{*n} * P_2}$ converge to a limit $a_* = a_{P_1 * Q^{*\infty}}$ (a_* depending on P_1 but not on P_2), with velocity δ_Q , and periods are multiplied by q . Again the stability zones are not adjacent; infinitely many discontinuities exist between them. But for superstable periods

$$k(a_n) \sim |a_* - a_n|^{\nu_Q} \tag{3.8}$$

where ν_Q is given by

$$\nu_Q = -\ln q / \ln \delta_Q \tag{3.9}$$

(if $\epsilon = 1$ and $Q = RL$, we get $\nu_Q = -0.27384$), and near a_* , we take into account all jumps and infinite discontinuities by writing

$$k(a) \sim |a_* - a|^{\nu_Q} g_Q(\ln|a_* - a| / \ln \delta_Q) \tag{3.10}$$

where g_Q is some complicated function which is periodic ($g_Q(x + 1) = g_Q(x)$) and may be infinite for certain values of its argument.

Remarks

1. The function g_Q is not the same for $a < a_*$ and $a > a_*$. This can be seen in figures 3(a) and 4 for $Q = R$.

2. From (3.9) and (3.5), $\nu_{Q^*1} = \nu_Q$, but there is no simple relation between $\nu_{Q_1^*Q_2}$, ν_{Q_1} and ν_{Q_2} . Nevertheless, as $\delta_{Q_1^*Q_2} = \delta_{Q_2^*Q_1}$, we have the weaker result $\nu_{Q_1^*Q_2} = \nu_{Q_2^*Q_1}$.

3.4. Theoretical background

If P and Q are MSS sequences, the parameter a_{Q^*P} corresponding to the superstability of the non-primary MSS sequence Q^*P is related to a_P by a function of the form

$$a_{Q^*P} = f_Q(a_P) \tag{3.11}$$

where f_Q is monotonic with respect to P (and thus to a_P). We may calculate f_Q by expansion near $\epsilon = 0$ around the broken linear case (see § 5.3), but when ϵ is not small it is possible to find this function f_Q by numerical approximations only.

In order to extend the definition of f_Q to the whole stability zone of the periods P and Q^*P , one may write

$$a' = f_Q(a) \tag{3.12}$$

whenever a and a' belong to the stability zone of P and Q^*P respectively, and $s_{Q^*P}(a') = s_P(a)$, the parameter of stability s being defined as in equation (2.4). The previous definition (3.11) corresponds to $s_P(a) = s_{Q^*P}(a') = 0$.

This pointwise construction of f_Q is possible owing to the property of internal similarity and to the fact that the zones of stability of the various periods do not overlap. It is apparently true from numerical calculations that these stability zones are dense. Thus one may assume (and we shall do it) that f_Q may be defined by continuity for any value of a . (As we already pointed out in § 2, a similar construction for mapping $*P$ is not possible; we may write $a_{Q^*P} = \tilde{f}_P(a_Q)$ instead of (3.11), but we cannot extend it to the whole stability zone.)

If q is the period of the sequence Q , obviously $k(a') = qk(a)$. The solution of

$$a_* = f_Q(a_*) \tag{3.13}$$

corresponds to a sequence P such that $Q^*P = P$; whence $P = Q^{*\infty}$, where the symbol $Q^{*\infty}$ represents the infinite sequence obtained as the limit of the sequences Q^{*N} when N goes to infinity[†]. If $a_n = a_{Q^{*n}}$, then

$$a_{n+1} = f_Q(a_n) \approx a_* - (a_* - a_n)df/da|_{a=a_*}$$

when n is large. This yields at once the bifurcation velocity

$$\delta_Q = (df_Q/da|_{a=a_*})^{-1}. \tag{3.14}$$

More generally, if we consider sequences $P_1^*Q^{*n}P_2$ corresponding to parameters $b_n = a_{P_1^*Q^{*n}P_2}$, we have $b_{n+1} = (f_{P_1} \circ f_Q \circ f_{P_1}^{-1})(b_n)$ as $b_n = f_{P_1} \circ f_Q^n \circ f_{P_2}$ (a_{blank}). The symbol \circ denotes the composition law for applications: $(f \circ g)(x) = f(g(x))$. The fixed point is $b_* \neq a_*$ corresponding to the infinite sequence $P_1^*Q^{*\infty}$, but because of the chain rule the bifurcation velocity is again δ_Q . As already pointed out, P_2 plays no role.

Notice that δ_Q and ν_Q are not only functions of period q but of the sequence Q itself. Two sequences of the same period q have distinct bifurcation velocities. For example $\delta_{RL^2} \neq \delta_{RLR}$. This will be shown for small ϵ in § 5.

We are now able to derive the commutation law

$$\delta_{Q_1^*Q_2} = \delta_{Q_2^*Q_1} \tag{3.15}$$

[†] We shall not enter here into the rigorous definition of infinite sequences. This has already been done in I.

and the corresponding law for critical exponents

$$\nu_{Q_1 \circ Q_2} = \nu_{Q_2 \circ Q_1}. \tag{3.16}$$

We have

$$f_{Q_1 \circ Q_2} = f_{Q_1} \circ f_{Q_2}, \quad f_{Q_2 \circ Q_1} = f_{Q_2} \circ f_{Q_1}$$

where \circ denotes again the usual composition law.

Let a_* and b_* be the fixed points for $f_{Q_1 \circ Q_2}$ and $f_{Q_2 \circ Q_1}$ respectively. Obviously

$$b_* = f_{Q_2}(a_*), \quad a_* = f_{Q_1}(b_*).$$

As

$$\begin{aligned} (\delta_{Q_1 \circ Q_2})^{-1} &= df_{Q_1 \circ Q_2}/da|_{a=a_*} = df_{Q_1}/da|_{a=f_{Q_2}(a_*)} \times df_{Q_2}/da|_{a=a_*} \\ &= df_{Q_1}/da|_{a=b_*} \times df_{Q_2}/da|_{a=a_*} \end{aligned}$$

is now a symmetric function of a_* and b_* , we easily obtain results (3.15) and (3.16).

The fixed points a_* and the universal metric constants ν_Q and δ_Q are known once the function f_Q is known. Actually, even in the simplest cases it is not possible to get this function f_Q exactly. The first-order correction when ϵ is small is given in § 5. When ϵ is not small, we suggest different kinds of approximation schemes which are based on conjugation law (3.4) and are to be compared with the usual renormalisation methods. This will be done in § 4.

3.5. Analogies and differences with critical phenomena

We end this section with some remarks. Parameter a plays in bifurcation problems the role of temperature—or coupling constants—in the theory of phase transitions, and operation f_Q is analogous to the recurrence relation which renormalises coupling constants. Similarly the critical temperature corresponds to the fixed point of f_Q , and exponent ν_Q is related to the linearised expression for f_Q near a_* .

Nevertheless, there is a big difference, as there are an infinite number of operations f_Q . Two functions f_{Q_1} and f_{Q_2} define two renormalisations which have different critical points and critical exponents.

If $a_{Q_1^\infty}$ and $a_{Q_2^\infty}$ are the fixed points of f_{Q_1} and f_{Q_2} , then function $k(a)$ diverges near $a_{Q_1^\infty}$ and $a_{Q_2^\infty}$, then near $f_{Q_2}(a_{Q_1^\infty})$, $f_{Q_2^2}(a_{Q_1^\infty})$, \dots , $f_{Q_1}(a_{Q_2^\infty})$, \dots and more generally for an infinite number of values of the parameter. The set of points for which $k(a)$ is infinite is stable by all operations f_Q and has the power of the continuum (even if these points are nowhere dense; see I).

In the theory of phase transitions, the renormalisation is a transformation of the coupling constants of some functional (in our case the coefficients of the Taylor expansion of the transform near its maximum could be seen as more or less similar to these coupling constants). This transformation can be made infinitesimal, so that it defines a semi-flow in the space of the coupling constants, or it can be made by finite steps in the so-called ‘real space’ renormalisation. In the problem we consider here, the discreteness is fundamental— f_R multiplies any period by 2, f_{RL} by 3, and so on—and the behaviour of function $k(a)$ (length of the period as a function of a) near a critical point (that is near $a = a_*$) is not

$$k(a) \sim |a - a_{Q^\infty}|^{\nu_Q}$$

but rather of the form (3.10)

$$k(a) \sim |a - a_{Q^{\infty}}|^{\nu_Q} g_Q(\ln|a_{Q^{\infty}} - a| / \ln \delta_Q)$$

where g_Q is periodic ($g_Q(x + 1) = g_Q(x)$) and has a very complicated structure. (For a discussion of phase transitions, see, for example, Domb and Green (1976)).

4. Approximate calculations of the function f_Q

In this section we investigate various approximations for computing the function f_Q that allow one to derive the quantities δ_Q and ν_Q (equations (3.9) and (3.13), (3.14)). Using rather simple arguments, we may approximate the raw numerical results with a relative error that may be of the order of one or two per cent. All our approximation schemes may be refined indefinitely, at least in principle. The numerical evidence shows that the next-order calculations yield improved numerical values, so that we may believe that these calculations do really converge towards well defined results which are actually independent of the approximation scheme.

We describe first our calculations of $a_{Q^{\infty}}$ and δ_Q for $Q = R$ and the C^∞ endomorphism $\mathcal{T}_a(x)$ with a single maximum

$$\mathcal{T}_a(x) = 1 - ax^2. \tag{4.1}$$

We then explain how these calculations must be modified in the case $Q = RL$. Although the extension of these computations to other sequences does not involve any difficulty, at least in principle, we shall not do it, as we have no numerical results for comparison.

At the end of the section, we consider the nearly broken linear case ($\epsilon \sim 0$) defined in equation (1.1); all the methods we propose give the same result at the first order in ϵ , and we shall see in § 5 that it is the first term of a systematic expansion near $\epsilon = 0$. We shall not restrict this calculation to the sequences $Q = R$ and $Q = RL$ as its extension to any Q is straightforward.

The approximation schemes we developed for $\epsilon = 1$ may be also used for $\epsilon \neq 1$ or ϵ not too small. However, when this exponent is no longer an integer, many expressions become very complicated and are difficult to handle from a practical point of view, but the principle of the method remains unchanged. We expect a good convergence to the exact value for small ϵ . The inaccuracy worsens as ϵ increases.

Finally, we give the *exact* expression for functions f_Q for the piecewise linear mappings with a flat top. Their derivation will be done in a forthcoming publication.

4.1. Quadratic case and $Q = R$

Let us consider the endomorphism $\mathcal{T}_a(x) = 1 - ax^2$ in the interval $[1, 2]$ of values of the parameter a . If x belongs to the interval $I(a) \equiv [1 - a, 1]$, then $\mathcal{T}_a(x) \in I(a)$ so that the interval $I(a)$ is stable under the action of \mathcal{T}_a , and \mathcal{T}_a has a single extremum in this stable interval.

Let us now consider the range of values of the parameter defined by $1 < a < a_0$, where a_0 is the real root of the polynomial $X^3 - 2X^2 + 2X - 2$. In this range of values of a , two disconnected intervals exist, $I_1(a) \equiv [1 - a, a - 1]$ and $I_2(a) \equiv [1 - a + 2a^2 - a^3, 1]$, which are both stable under the action of \mathcal{T}_a^2 , and \mathcal{T}_a^2 has a single extremum in each of these intervals; of course $\mathcal{T}_a(I_1(a)) \subset I_2(a)$ and $\mathcal{T}_a(I_2(a)) \subset I_1(a)$. This helps one to understand the way in which the law of internal similarity holds: for a

given value of a , there exists a value A of the parameter corresponding to a mapping \mathcal{T}_A and a homeomorphism h_1 between $I_1(a)$ and $I(A)$ such that

$$\mathcal{T}_a^2(x) = h_1 \circ \mathcal{T}_A(x) \circ h_1^{-1} \tag{4.2}$$

(see also figure 5).




Figure 5. Segments $I_1(a)$ and $I_2(a)$ stable by transformation \mathcal{T}_a^2 for a given a . They are represented as full lines with the corresponding parts of $\mathcal{T}_a(x)$ and $\mathcal{T}_a^2(x)$.

To our knowledge, there is no way of proving the existence of such a pointwise homeomorphism. This proof would require, in particular, a knowledge of all the invariant sets of \mathcal{T}_a^2 and \mathcal{T}_A in $I_1(a)$ and $I(A)$. In what follows we shall adopt a ‘practical’ point of view and try to find in an approximate way the homeomorphism h_1 and the function $f_R(a)$ that gives the value of A by imposing a limited number of conditions to h_1 and $f_R(A)$. As we shall choose very simple functions for h_1 , we shall assume implicitly that, at least for some values of a , h_1 is not a ‘wild’ function and can be actually fairly well approximated by this simple function.

Remarks

1. The same holds for the interval $I_2(a)$. We look for a homeomorphism h_2 of $I(A)$ onto $I_2(a)$ such that

$$h_2(I(A)) = I_2(a), \quad h_2^{-1} \circ \mathcal{T}_a^2 \circ h_2 = \mathcal{T}_A. \tag{4.3}$$

2. We are not able to say whether conjugation implies for h_1 and h_2 more than simple continuity. We think that, actually, h_1 and h_2 need to be differentiable at least.

3. To find the function f_R is equivalent to looking for homeomorphisms h_1 and h_2 (equations (4.2) and (4.3)). Practically, there is no hope of getting them from a series expansion. Even for the reduced functions $\mathcal{T}_a(x) = 1 - ax^2$ the homeomorphisms cannot be polynomials or rational functions, as can be seen by simple consideration of the degrees and factorisations for the polynomials that conjugation (4.2) or (4.3) would require.

We are led to do simple approximations. In the following, we indicate several possibilities.

4.1.1. Centred renormalisation with one parameter. We assume that conjugation law (4.2) holds and that in a first approximation h_1 is linear. Keeping in $\mathcal{T}_a^2(x)$ terms up to

second order, as (4.2) must hold for small x , we get

$$1 - a + 2a^2 h_1^2(x) = h_1(1 - Ax^2)$$

the solution of which is $h_1: x \rightarrow (1 - a)x$.

Parameters A and a are related through

$$A = 2a^2(a - 1) \tag{4.4}$$

which yields an implicit and approximate expression for f_R . From (3.13) and (3.14)

$$a_* = (1 + \sqrt{3})/2 \quad (\text{fixed point } a_* = 2a_*^2(a_* - 1))$$

and

$$\delta_R^{-1} = \partial A / \partial a |_{a=a_*} = (4 - \sqrt{3})/13.$$

We get a_* and δ_R^{-1} with an error of 2.5% and 19% respectively. The results are shown in table 3, row 1.

Table 3. Comparison of the different methods of renormalisation for the set of sequences R^{*n} .

Method of calculation	δ_R^{-1}	Relative error in δ_R^{-1} (%)	Critical point $a_{R^{*\infty}}$	Relative error in $a_{R^{*\infty}}$ (%)	γ_R^{-1}	Relative error in γ_R (%)
Centred renormalisation						
1 parameter	0.1745	19	1.366 025	2.5	0.366 025	8
2 parameters	0.2064	4	1.400 030	0.08	0.394 628	1.2
3 parameters	0.2214	3	1.401 393	0.02	0.403 403	1
Non-centred renormalisation						
1 parameter	0.2265	6	1.457 107	4		
2 parameters	0.2117	1.5	1.402 414	0.09		
3 parameters	0.2133	0.4	1.401 003	0.01		
Non-linear homeomorphism						
1 parameters	0.1874	12	1.384 574	1		
1st equality of slopes						
1 parameters	0.1952	9	1.390 388	0.7		
2nd equality of slopes						
1 parameter	0.2167	1	1.401 420	0.02		
Experimental (measured) value						
	0.214 169 37		1.401 155 189 09		0.399 535	

4.1.2. Non-centred renormalisation with one parameter. We proceed in the same way as above, except that we have (4.3) as a conjugation law. In the interval $I_2(a)$, \mathcal{T}_a^2 reaches its maximum for the positive solution x_0 of $\mathcal{T}_a(x_0) = 0$. Homeomorphism h_2 is assumed

to be linear:

$$h_2(x) = x_0 + \lambda x. \tag{4.5}$$

The choice of the constant term in this expression ensures that the extremum of \mathcal{T}_a^2 is mapped on the critical point of \mathcal{T}_A . For the reduced mapping (4.1) we get $x_0 = a^{-1/2}$, $x_0 + \lambda = 1$ (this last condition ensures that the common right boundaries of $I_2(a)$ and $I(A)$ are mapped onto each other).

Then, by imposing again that the quadratic terms in \mathcal{T}_a^2 and $h_2 \circ \mathcal{T}_A \circ h_2^{-1}$ are the same, one obtains

$$A = 4a^2(1 - a^{-1/2}). \tag{4.6}$$

This yields, by using (3.13) and (3.14), the following approximate expressions for a_* and δ_R^{-1} : $a_* = (3 + 2\sqrt{2})$ (relative error 4%), $\delta_R^{-1} = (3 - \sqrt{2})/7$ (relative error 6%) (see table 3, row 4).

4.1.3. Centred renormalisation with several parameters. To get a better approximation, we may consider a larger set of transformations depending on several parameters a, b, c, \dots :

$$T_{a,b,c,\dots}(x) = 1 - ax^2 + bx^4 + cx^6 + \dots$$

The coefficients of $T_{a,b,c,\dots}^2(x)$ are polynomials in a, b, c, \dots , and the stable segments I, I_1, I_2 depend now on a, b, c, \dots .

If we look for a centred approximation and assume that h_1 is linear

$$T_{a,b,c,\dots}^2(h_1(x)) = h_1(T_{A,B,C,\dots}(x)), \quad x \in I(a, b, c, \dots)$$

we get

$$h_1(x) = u_0x, \quad A = u_2u_0, \quad B = u_4u_0^3, \quad C = u_6u_0^5$$

where

$$T_{a,b,c,\dots}^2(x) = u_0 - u_2x^2 + u_4x^4 + u_6x^6 + \dots$$

and the u_i 's are polynomials in a, b, c, \dots .

Then the fixed point is the solution of $A = a, B = b, C = c, \dots$, and the unstable eigenvalues of the matrix of the derivatives give δ_R .

For example, with two parameters a, b we get

$$u_0 = 1 - a + b, \quad u_2 = 4ab - 2a^2, \quad u_4 = -a^3 + 6a^2b - 2ab + 4b^2.$$

Then

$$A = (1 - a + b)(4ab - 2a^2), \quad B = (1 - a + b)^3(-a^3 + 6a^2b - 2ab + 4b^2). \tag{4.7}$$

The calculations give very good results (see table 3, row 2). The accuracy is improved further with three parameters (table 1, row 3), but approximations, though more accurate, become very complicated with more parameters.

We have plotted on figure 6 the critical line and fixed point in the plane (a, b) . The value a_* for $\mathcal{T}_a(x) = 1 - ax^2$ corresponds to $b = 0$ on the critical line. The critical line stops at another fixed point $a = 0, b = -1.545$ corresponding to the sequence $R^{*\infty}$ and the transformation $T_b(x) = 1 + bx^4$.




Figure 6. Two-parameter centred renormalisation. The recurrence equations (4.7) have interesting fixed points. The origin is attractive; the fixed point of $T_b = 1 + bx^4$ is repulsive. The critical fixed point P is unstable except for the critical line which goes through the fixed points of the one-parameter transformation $T_b = 1 + bx^4$ (point M) and $\mathcal{T}_a = 1 - ax^2$ (point N).

Important remarks

1. This procedure is similar to that used in phase transition theory, where one assumes that the renormalised Hamiltonian is accurately described by a small number of ‘relevant’ coupling constants. As we pointed out in the introduction, if we were looking to $a_{R^{*}}$ and δ_R for another C^∞ transformation, for example

$$T_a(x) = 1 - a \sinh^2 x \sim 1 - a(x^2 + \frac{1}{3}x^4)$$

we should recover the same δ_R . The fixed point $a_{R^{*}}$ is approximately at the intersection of the critical line with $b = -a/3$.

2. *Non-centred renormalisation with several parameters.* The same method holds for I_2 and h_2 , but, as $T_{a,b,c,\dots}^2$ is no longer symmetric around x_0 , we must include odd powers of x . For example, with three parameters a, b, c

$$T_{a,b,c}(x) = 1 - ax^2 + bx^3 + cx^4$$

and $x_0 (\neq 0)$ is the solution of $T_{a,b,c}(x_0) = 0$. Expanding $T_{a,b,c}^2$ around x_0 we get

$$T_{a,b,c}^2(x_0 + y) = 1 - v_2y^2 + v_3y^3 + v_4y^4 + \dots$$

where again v_2, v_3, v_4 are functions of a, b, c and

$$A = v_2(1 - x_0), \quad B = v_3(1 - x_0)^2, \quad C = v_4(1 - x_0)^3.$$

We did calculations with two and three parameters (table 3, rows 5 and 6). The latter gives δ_R^{-1} with an error of 0.4% and $a_{R^{*}}$ with an error of 0.01%. This method is the most accurate among all we tried, and results should be still better with more parameters. Unfortunately, this method yields the longest renormalisation formulae!

4.1.4. *Nonlinear homeomorphism.* Up to now, we have approximated homeomorphisms h_1 and h_2 by linear functions. This is reasonable, as this is true asymptotically. Nevertheless we may ask whether a more elaborate choice for h_1 and h_2 gives better accuracy. As an example, we look here at a_* and δ_R^{-1} when the renormalisation is

centred and with the homeomorphism $h_1^{-1}(x) = \lambda x + \mu x^3$. From $h_1^{-1}(T_a^2(x)) = T_A(h_1^{-1}(x))$ and neglecting all terms $x^n, n \geq 5$ we get the equation for A

$$\alpha(\hat{a})A^2 + \beta(a)A + \gamma(a) = 0 \tag{4.8a}$$

where

$$\alpha(a) = 8$$

$$\beta(a) = 16a^2 - 32a^3 + 83a^4 - 151a^5 + 222a^6 - 318a^7 + 255a^8 - 75a^9 \tag{4.8b}$$

$$\gamma(a) = 64a^6 - 208a^7 + 368a^8 - 608a^9 + 736a^{10} - 464a^{11} + 112a^{12}$$

whence a_* and $\hat{\delta}_R^{-1}$ (table 3, row 7). The accuracy is better than with centred (linear) renormalisation with one parameter, but not as good as the other approximations we described in §§ 4.1.2 and 4.1.3.

4.1.5. Equality of the slopes. We deal only with sequences R^{*n} . The parameter s_{2^n} (equations (2.3) and (2.4)) which defines the stability zone for R^{*n} is the slope of $\mathcal{F}_a^{2^n}(x)$ at one point of the period.

Sequence b (equal to the fixed point) is stable if $|s_1| < 1$, i.e. $-\frac{1}{4} < a < \frac{3}{4}$, where $s_1(a) = 1 - (1 + 4a)^{1/2}$.

Sequence R is stable if $|s_2| < 1$, i.e. $\frac{3}{4} < a < \frac{5}{4}$, where $s_2(a) = 4(1 - a)$, and $s_4(a)$ is given by the implicit expression

$$1 + 2a^2 - 3a^3 + 3a^4 + a^6 + \frac{1}{16}s_4(a)(a^4 - a^3 - a^2 - 3) + \frac{s_4^2(a)}{16^2}(3 - a^2) - \frac{s_4^3(a)}{16^3} = 0$$

which holds for $\frac{3}{4} < a < \bar{a}$ (destabilisation value).

When $-\frac{1}{4} < A < \frac{3}{4}$, the function f_R is given implicitly by $s_1(A) = s_2(a)$, i.e.

$$a = [3 + (1 + 4A)^{1/2}]/4 \tag{4.9a}$$

and when $\frac{3}{4} < A < \frac{5}{4}$, $s_2(A) = s_4(a)$ gives for $a = f_R(A)$ the implicit equation

$$A^3 + A^2(9 - 4a^2) + A(27 + 24a^2 + 16a^3 - 16a^4) + 27 + 108a^2 - 208a^3 + 208a^4 - 192a^5 + 64a^6 = 0. \tag{4.9b}$$

The analytic form for $f_R(A)$ changes at $A = \frac{3}{4}$, though both f_R and df_R/dA are continuous. Using (4.9a) as an approximation for f_R , we get $a_* = (7 + \sqrt{17})/8$ and $\hat{\delta}_R^{-1} = (\sqrt{17} - 1)/16$. When (4.9b) is used, the accuracy is much better (actually, it is one of the best approximations; see table 3, rows 8-9), as the expression (4.9b) for f_R corresponds to an interval which is nearer the limit point $a_{R^{*\infty}}$.

Remark

One may notice that even the simplest methods sketched in this subsection give a good order of magnitude for $a_{R^{*\infty}}$ and $\hat{\delta}_R$; more elaborate methods (mainly non-centred linear renormalisation with three parameters, and the slope method of equation (4.9b)) give very accurate results. Nevertheless we are not able *a priori* to tell which method will give the best result. This is mainly due to our lack of knowledge of the manner in which these approximations converge to the exact result.

One could improve these approximation schemes by increasing the number of parameters, but again we do not know if this will actually increase the precision. We have no knowledge of a proof of the convergence of all these methods to a single result.

4.2. Quadratic case ($\epsilon = 1$ and $Q = RL$)

We indicate briefly how the above prescriptions are modified when we replace R by RL .

The main point is that there are three disconnected intervals J_1, J_2, J_3 which are stable by $\mathcal{T}_a^3(x), \mathcal{T}_a(J_1) \subset J_2, \mathcal{T}_a(J_2) \subset J_3, \mathcal{T}_a(J_3) \subset J_1$ for a range of the parameter $a_1 < a < a'_1; a_1$ and a'_1 being the values of the parameter a corresponding to sequences RL and $RL * RL^\infty$ respectively. On each interval \mathcal{T}_a^3 has one extremum only and only one of these (say J_1) is centred at $x=0$. Choosing any one of these intervals J_i ($i = 1, 2, 3$), we may write $\mathcal{T}_a^3 \sim \mathcal{T}_A$, which implies the existence of a homeomorphism j_i between $I(A)$ and $J_i(a)$ such as

$$j_i(I(A)) = J_i(a), \quad \mathcal{T}_a^3(j_i(x)) = j_i(\mathcal{T}_A(x)), \quad \forall x \in I(A).$$

We list here the analytic expressions for the simplest approximations for f_{RL} :

$$A = 4a^3(a-1)(a^3-2a^2+a-1) \quad \text{centred renormalisation with one parameter}$$

$$A = 8a^3\left(1-a+\frac{1}{\sqrt{a}}\right) \quad \text{first non-centred renormalisation with one parameter}$$

$$A = 16a^3\left(1+\frac{1}{\sqrt{a}}\right)\left[1-\frac{1}{\sqrt{a}}\left(1+\frac{1}{\sqrt{a}}\right)^{1/2}\right] \quad \text{second non-centred renormalisation with one parameter}$$

$$A = \frac{[4a-7+4a(4a-7)^{1/2}]^2-1}{4} \quad \text{equality of slopes } (s_1(A) = s_3(a)).$$

Again $a_{RL}^{*\infty}$ is known with a good accuracy and we have a good order of magnitude for δ_{RL}^{-1} . We get better results with a centred renormalisation with two parameters (table 4).

Table 4. Comparison of the different methods of renormalisation for the set of sequences $(RL)^{*\infty}$.

Method of calculation	δ_{RL}^{-1}	Relative error in δ_{RL}^{-1} (%)	Critical point $a_{RL}^{*\infty}$	Relative error in $a_{RL}^{*\infty}$ (%)	γ_{RL}^{-1}	Relative error in γ_{RL} (%)
Centred renormalisation						
1 parameter	0.015 077	17	1.784 998	0.08	0.099 953	7
2 parameters	0.018 118	0.1	1.786 466	0.0015	0.107 834	0.04
1st non-centred renormalisation						
1 parameter	0.017 176	5	1.787 164	0.04		
2nd non-centred renormalisation						
1 parameter	0.017 721	2	1.787 696	0.07		
Equality of slopes						
1 parameter	0.016 229	10	1.785 972	0.03		
Experimental (measured) value						
	0.018 1005		1.786 440 255		0.107 789	

These methods might be extended to all sequences Q of period q . In particular, it has already been shown that in some range of values of the parameter a , q disconnected intervals exist; each of them is stable under the action of $T_a^{(q)}(x)$, and this function has a single critical point in each interval (Guckenheimer 1977).

4.3. *Study of small ϵ*

The same methods are developed for the mapping $\mathcal{T}_a(x) = 1 - a|x|^{1+\epsilon}$ when ϵ is small.

By the centred and non-centred renormalisation with one parameter we get for f_R
 $A = a^2(1 + \epsilon)(a - 1)^\epsilon$ and $A = a^2(1 + \epsilon)^{1+\epsilon}(1 - a^{-1/(1+\epsilon)})^\epsilon$

respectively, which gives to first order the same results:

$$\begin{aligned} a_{R^*} &\approx 1 + \varphi, & \delta_R &\approx 2 + \epsilon/\varphi \text{ hence} \\ \nu_R &\approx -1 + \epsilon/2(\ln 2)\varphi \end{aligned} \tag{4.10}$$

where φ is the solution of $\varphi + \epsilon \ln \varphi + \epsilon = 0$. We shall see in § 5 that this is the *exact* first-order term.

Similarly for $Q = RL$ we get

$$\begin{aligned} A &= a^3|a - 1|^\epsilon(1 + \epsilon)^2(a|a - 1|^{1+\epsilon} - 1)^\epsilon \text{ and} \\ A &= a^3(1 + \epsilon)^{2+\epsilon}|a - a^{-1/(1+\epsilon)} - 1|^\epsilon \end{aligned}$$

which gives for $a_{(RL)^*}$ and δ_{RL} the same first-order terms

$$a_{(RL)^*} \approx a_{RL} + \varphi, \quad \delta_{RL} \approx \epsilon_{a_{RL}}/\varphi \tag{4.11}$$

where a_{RL} is the root of $1 - a_{RL}|a_{RL} - 1|^{1+\epsilon} = 0$, corresponding to the superstability point of sequence RL , and

$$\varphi \approx e^{-2}[(2a_{RL} - 1)(a_{RL} - 1)]^{-1} a_{RL}^{-2/\epsilon}. \tag{4.12}$$

Again this is the precise first-order result as we shall see in § 5. Notice that, in contrast to δ_R , δ_{RL} increases infinitely for $\epsilon \rightarrow 0$; as a consequence the critical exponent ν_{RL} goes to zero.

As even the simplest methods seem to give very good results for small ϵ , we may continue and do the same thing with any sequence Q of period q . With the centred renormalisation method, as

$$\begin{aligned} \mathcal{T}_a^{(k)}(x) &= \mathcal{T}_a^k(0) + (-a)^k(1 + \epsilon)^{k-1} \\ &\quad \times \text{sgn}(\mathcal{T}_a(0) \dots \mathcal{T}_a^{(k-1)}(0))|\mathcal{T}_a(0) \dots \mathcal{T}_a^{(k-1)}(0)|^\epsilon |x|^{1+\epsilon} + \dots \end{aligned}$$

we get

$$A = |\mathcal{T}_a(0) \dots \mathcal{T}_a^{(k)}(0)|^\epsilon \text{sgn}(\mathcal{T}_a(0) \dots \mathcal{T}_a^{(k-1)}(0))(1 + \epsilon)^{k-1} a^k (-1)^{k-1}.$$

$\mathcal{T}_a(0), \dots, \mathcal{T}_a^{(k-1)}(0)$ are all far from zero, but $\mathcal{T}_a^{(k)}(0)$ is small and can be written as

$$\mathcal{T}_a^{(k)}(0) = (a - a_Q)(d/da)\mathcal{T}_a^{(k)}(0) + \dots$$

where a_Q is the solution of $\mathcal{T}_{a_Q}^{(k)}(0) = 0$ and corresponds to the superstability point for sequence Q , whence the first-order approximation for a_*

$$a_* - a_Q \sim \frac{e^{-(k-1)} a_*^{-(k-1)/\epsilon}}{|\mathcal{T}_a(0) \dots \mathcal{T}_a^{(k-1)}(0)| |(d/da)\mathcal{T}_a^k(0)|}$$

which generalises result (4.12) and is exactly expression (5.12). We recover that, when $Q \neq R$, $\delta_Q \rightarrow \infty$ (and $\nu_Q \rightarrow 0$) in the limit $\epsilon \rightarrow 0$.

4.4. Piecewise linear mapping

We consider the transformation $T_{r,\lambda}(x)$ depending on two parameters r and λ and defined by (figure 7)

$$T_{r,\lambda}(x) = \begin{cases} \lambda x & x < 1 \\ \lambda & 1 < x < r \\ \lambda(1+r) - \lambda x & x < r. \end{cases}$$

For $r = 1$ we recover the so-called broken linear case $\epsilon = 0$ that we studied in I. Perhaps the broken linear transform in the limit $r \neq 0, \lambda \rightarrow \infty$ could have something to do with the behaviour of $\mathcal{T}_a(x)$ when $\epsilon \rightarrow \infty$.




Figure 7. Plot of $T_{r,\lambda}(x)$ as a function of x .

All calculations may be carried out. The iterates of $T_{r,\lambda}$ are piecewise linear functions with flat extrema. With two parameters, the linear renormalisation is then exact. For $Q = R$, $T_{r,\lambda}^2 \sim T_{R,\Lambda}$, where

$$\Lambda = \lambda^2, \quad (R - 1)/(R + 1) = (\lambda + 1)(r - 1)/(\lambda - 1)(r + 1).$$

Similarly, for $Q = RL$, $T_{r,\lambda}^3 \sim T_{R',\Lambda'}$, with

$$\Lambda' = \lambda^3, \quad (R' - 1)/(R' + 1) = (\lambda^2 + \lambda + 1)(r - 1)/(\lambda^2 - \lambda - 1)(r + 1).$$

Remark

The fixed points of these recursion relations go to infinity, and it can be shown that in this case the length of the periods diverges logarithmically near the critical values of the parameter. More generally, for Q with period k and characteristic polynomial $P_Q(x) = \alpha_0 \dots \alpha_{k-2} \mathcal{P}_Q(x)$ (see § 2.2), $T_{r,\lambda}^k \sim T_{R'',\Lambda''}$, with

$$\Lambda'' = \lambda^k, \quad (R'' - 1)/(R'' + 1) = (\lambda^k - 1)(r - 1)/(\lambda - 1)P_Q(\lambda)(r + 1).$$

5. Study of the superstable periods for $\epsilon \sim 0$

As we saw in the preceding sections, we need only the behaviour of T_a near its maximum. Then, in § 4, we considered the reduced transformation

$$\mathcal{T}_a(x) = 1 - a|x|^{1+\epsilon} \tag{5.1}$$

depending continuously on the parameter ϵ . For $\epsilon = 0$ we recover the piecewise linear mapping

$$\mathcal{L}_a(x) = 1 - a|x| \tag{5.2}$$

which we studied in some detail in I. The superstable (or MSS) parameters a_Q are then the so-called λ numbers. For small ϵ the parameter a_Q for sequence Q may be obtained by expansion near the λ number of this period. The knowledge of a_Q itself is not very important, but from equations (3.14) and (3.9) we may get the universal parameters δ_Q and ν_Q respectively. This is essentially what is explained in that section. The exponent ϵ is given and assumed to be small although a is varying, and we shall call a *the* parameter.

We use in a relaxed way the terminology first order, second order, . . . Rigorously it is not correct, as the dependence in ϵ will not be polynomial. When a quantity $\phi(\epsilon)$ may be written as a sum

$$\phi(\epsilon) = \phi_1(\epsilon) + \phi_2(\epsilon) + \phi_3(\epsilon) + \dots$$

with

$$\phi_2(\epsilon)/\phi_1(\epsilon) \rightarrow 0, \quad \phi_3(\epsilon)/\phi_2(\epsilon) \rightarrow 0, \dots$$

then $\phi_1(\epsilon)$, $\phi_2(\epsilon)$, $\phi_3(\epsilon)$ will be said to be of first, second and third-order respectively.

As the case $Q = R$ differs from the other sequences, it is studied separately in § 5.4.

In the following, \bar{a}_Q and a_Q are the values of the superstable sequence when $\epsilon = 0$ and $\epsilon \neq 0$ respectively; $x_n(\bar{a}_Q) = \mathcal{L}_{\bar{a}_Q}^{(n)}(0)$ and $y_n(a_Q) = \mathcal{T}_{a_Q}^{(n)}(0)$ are the iterates ($n = 1, \dots, k$) of the maximum when $\epsilon = 0$ and $\epsilon \neq 0$ respectively.

5.1. Shift of the parameters of superstable periods

When ϵ is small, we rewrite $\{y_n\}$ and a_Q as

$$y_n = x_n + \epsilon z_n + O(\epsilon^2), \quad a_Q = \bar{a}_Q + \epsilon \Delta a + O(\epsilon^2) \tag{5.3}$$

and to first order, the iteration formula (5.1) gives

$$z_n = -\bar{a}_Q(\text{sgn } x_{n-1})z_{n-1} - \Delta a|x_{n-1}| - \bar{a}_Q|x_{n-1}| \ln|x_{n-1}| \tag{5.4}$$

with ‘sgn’ being the sign function (equation (2.10)).

The solution of (5.4) with initial conditions $z_0 = z_1 = 0$ is

$$z_n = \sum_{i=2}^{n-1} \bar{a}_Q^{n-i} \alpha_{n-2} \dots \alpha_{i-1} x_i \ln|x_i| + (\Delta a/\bar{a}_Q) \sum_{i=0}^{n-1} \bar{a}_Q^{n-1-i} (n-1-i) \alpha_{n-2} \dots \alpha_i$$

the first sum being zero when $n < 3$.

As Q is a k period, $z_k = 0$, whence the shift of the parameter is of order ϵ and

$$\Delta a = -\frac{\sum_{i=2}^{k-1} \bar{a}_Q^{k-i} \alpha_{k-2} \dots \alpha_{i-1} x_i(\bar{a}_Q) \ln|x_i(\bar{a}_Q)|}{(d/d\bar{a}_Q)x_k(\bar{a}_Q)} \tag{5.5}$$

In (5.5) we have made explicit the dependence on the parameter \bar{a}_Q and $x_k(\bar{a}_Q) = \mathcal{P}_Q(\bar{a}_Q) = 0$, where $\mathcal{P}(x)$ is defined in (2.10).

Notice that, if $Q (= P_1 * P_2)$ is non-primary, as $\mathcal{P}_Q(\bar{a}_{P_1}) = \mathcal{P}_{P_1}(\bar{a}_{P_1})\mathcal{P}_{P_2}(\bar{a}_{P_1}^k)$, both numerator and denominator factorise in (5.5), $\bar{a}_Q = \bar{a}_{P_1}$ and $\Delta\bar{a}_Q = \Delta\bar{a}_{P_1}$. Again it is not possible to separate non-primary sequences by a first-order expansion. We shall see in § 5.3 that the splitting $a_{P_1+P_2} - a_{P_1}$ is very small.

5.2. Zone of stability

Now we assume that a_Q is known, $\{y_n\}$ is the sequence of the iterates of the maximum, and we look for a value b of the parameter and for k numbers $t_0(b), t_1(b), \dots, t_{k-1}(b)$ ($t_k(b) = t_0(b)$) such that

$$\prod_{i=1}^k (d/dt_i)\mathcal{T}_b(t_i) = \eta \tag{5.6}$$

where $t_i = \mathcal{T}_b(t_{i-1})$ is the i th iterate of t_0 , and $\eta = \pm 1$.

When $\eta = 0$, $b = a_Q$, although b_s and b_u mark the beginning ($\eta = +1$) and end ($\eta = -1$) of the stability zone of the period Q in the parameter space. Equation (5.6) can be rewritten

$$(-b)^k (1 + \epsilon)^k |t_1 \dots t_k|^\epsilon (\text{sgn } t_1) \dots (\text{sgn } t_k) = \eta. \tag{5.7}$$

As $b \approx a_Q$, then $t_i \approx y_i$ and $\text{sgn } t_i = \text{sgn } y_i = \text{sgn } x_i$ if $1 \leq i \leq k-1$. We have $t_k = t_0 \approx 0$, and from (5.7)

$$|t_0| = b^{-k/\epsilon} (1 + \epsilon)^{-k/\epsilon} / |t_1 \dots t_{k-1}|$$

with $\text{sgn } t_0 = (-1)^k \eta \text{sgn}(y_1 \dots y_{k-1})$. Then the stability zone is very small, and its width is proportional to $b^{-k/\epsilon}$. Setting

$$b - a_Q = a_Q^{-k/\epsilon} u, \quad \text{and} \quad t_n - y_n = a_Q^{-k/\epsilon} v_n \tag{5.8}$$

the recursion formula (5.1) gives to first order

$$v_n = -a_Q v_{n-1} \text{sgn } x_{n-1} - u |x_{n-1}|$$

with general solution (compare with (5.4))

$$v_n = A \alpha_{n-2} \dots \alpha_0 (a_Q)^n + u \sum_{i=0}^{n-1} (a_Q)^{n-2-i} (n-1-i) \alpha_{n-2} \dots \alpha_i$$

A being a parameter. With the initial conditions for t_0 above, we get

$$v_n = u (d/da_Q) x_n(a_Q) + a_Q^{n-k} e^{-k} \alpha_{n-2} \dots \alpha_0 / |y_1 \dots y_{k-1}|.$$

The condition $v_k = v_0 = t_0 a_Q^{k/\epsilon}$ reads to first order

$$u (d/da_Q) x_k(a_Q) - e^{-k} \alpha_{k-2} \dots \alpha_0 / |y_1 \dots y_{k-1}| = -\eta (\alpha_{k-2} \dots \alpha_0 e^{-k}) / |y_1 \dots y_{k-1}|.$$

(1) When $\eta = -1$ (end of stability zone) we get

$$b_u = a_Q + 2(a_Q^{-k/\epsilon} e^{-k} \alpha_{k-2} \dots \alpha_0) / |y_1 \dots y_{k-1}| (d/da_Q) x_k(a_Q). \tag{5.9}$$

For example, if $k = 3$, $Q = RL$ and

$$b_u \sim a_Q + 2a_Q^{-3/\epsilon} e^{-3} / (\bar{a}_Q - 1)(2\bar{a}_Q - 1)$$

where $\bar{a}_Q = (1 + \sqrt{5})/2$ is the golden number. Notice that, as $a_Q - \bar{a}_Q \sim O(\epsilon)$, $a_Q^{-k/\epsilon}$ and $\bar{a}_Q^{-k/\epsilon}$ are not equivalent in the limit $\epsilon \rightarrow 0$, although we may replace $y_i(a_Q)$ by $x_i(\bar{a}_Q)$.
 (2) When $\eta = +1$ (beginning of stability zone) we get $u = 0$. The correction to the superstable parameter is at least of the order $\epsilon a_Q^{-k/\epsilon}$, and to a first approximation the beginning of the stability zone and the point of superstability coincide. To get the second-order term, we replace (5.8) by

$$b - a_Q = \epsilon v a_Q^{-k/\epsilon}, \quad t_n - y_n = a_Q^{-k/\epsilon} (v_n + \epsilon v'_n)$$

and keep in the iteration formula (5.1) all terms up to second order, getting

$$b_s = a_Q - \epsilon a_Q^{-k/\epsilon} e^{-k} \alpha_{k-2} \dots \alpha_0 / |y_1 \dots y_{k-1}| (d/da_Q) x_k(a_Q). \tag{5.10}$$

As $\alpha_{k-2} \dots \alpha_0 (d/da_Q) x_k(a_Q) > 0$, we have $b_s < a_Q < b_u$. We thus verify that the width of the stability zone (that is $b_s - b_u$) is very small and reduces to a point when $\epsilon = 0$. Moreover, when $\epsilon = 0$ the measure of the total stability zone is zero as no period is stable. We expected this result as the set $\{\bar{a}_Q\}$ for $\epsilon = 0$ is countable.

5.3. Splitting of non-primary sequences

Sequence Q is now assumed to be primary with period $k > 2$, P is any other sequence, primary or not, of period l . We look for the value a_{Q*P} of the parameter for which the period $Q * P$ is superstable. When $\epsilon = 0$, $\overline{a_{Q*P}} = \bar{a}_Q$; near $\epsilon \rightarrow 0$ (§§ 5.2 and 4.3), $a_{Q*P} - a_Q \sim a_Q^{-k/\epsilon} a_P^{1/\epsilon}$. This implies that a_{Q*P_1} and $a_{Q*P_1*P_2}$ cannot be separated in a first-order approximation unless $a_{P_2} = 1$ (which corresponds to $P_2 = R$). This is of course no longer true when we take into account higher-order terms.

Proof. $\{t_n\}$ ($n = 1, 2, \dots, kl$) is the sequence of iterates of the maximum for the value a_{Q*P} of the parameter; we have $t_{kl} = 0$ and $t_{km} \neq 0$ (but small) if $1 < m < l$; $\{y_n\}$ denotes again the sequence of iterates of the maximum for Q with parameter a_Q . We have $y_k = y_{2k} = \dots = y_{lk} = 0$ and $y_{km+r} = y_r$ ($0 \leq r < k, 0 \leq m < l$).

Let us write

$$t_n = y_n + v_n, \quad a_{Q*P} = a_Q + u \tag{5.11}$$

and assume that terms of order uv_n, v_n^2, u^2 may be neglected compared with v_n and u .

When $y_{n-1} \neq 0$ (or $r \neq 1$) the iteration formula (5.1) is rewritten

$$v_{mk+r} = a_P \alpha_{r-2} |y_{r-1}|^\epsilon (1 + \epsilon) v_{mk+r-1} - u |y_{r-1}|^{1+\epsilon}$$

although, if $y_{n-1} = 0$ (i.e. $r = 1$), $v_{mk+1} = -a_Q |v_{mk}|^{1+\epsilon}$.

The ‘first-order’ solution of the above recurrence equations is not obvious. After tedious manipulations and using results of § 5.1 we derive the expression for v_{kl} ($l' \leq l$). As $v_{kl} = 0$ (periodicity condition) we get

$$\ln[u^\epsilon |y_1 \dots y_{k-1}|^\epsilon (a_Q^k/a_P) (1 + \epsilon)^{k-1} |(d/da_Q) x_k(a_Q)|^\epsilon] = 0$$

whence u and the formula

$$a_{Q*P} \sim a_Q + a_Q^{-k/\epsilon} a_P^{1/\epsilon} e^{-(k-1)/\epsilon} / |x_1 \dots x_{k-1}| |(d/da_Q) x_k(\bar{a}_Q)|. \tag{5.12}$$

The correction is small and the displacement is negligible for large periods.

More generally, to first order

$$a_{Q^{*n+1}*P} = a_Q + a_Q^{-k/\epsilon} a_Q^{1/\epsilon} a_{Q^{*n}*P} e^{-(k-1)/|x_1 \dots x_{k-1}|} (d/da_Q)x_k(\bar{a}_Q) + \dots$$

The limit point a_* is the solution of

$$a_* = a_Q + a_Q^{-k/\epsilon} e^{-(k-1)} a_*^{1/\epsilon} / |x_1 \dots x_{k-1}| (d/da_Q)x_k(\bar{a}_Q)$$

and, as $a_{Q^{*n}*P} = a_Q + O(a_Q^{-(k-1)/\epsilon})$,

$$\delta_Q^{-1} \sim \left. \frac{\partial a_{Q^{*n+1}*P}}{\partial a_{Q^{*n}*P}} \right|_{a_{Q^{*n}*P} = a_Q}$$

(equation (3.14)), whence

$$\delta_Q^{-1} \sim [e^{-(k-1)} / |x_1 \dots x_{k-1}| (d/da_Q)x_k(\bar{a}_Q)] \epsilon^{-1} a_Q^{-k/\epsilon} (a_*^{1/\epsilon}) / a_* \tag{5.13}$$

and both a_* and δ_Q do not depend on P , as expected. The bifurcation velocity for $Q \neq R$ goes to infinity when $\epsilon \rightarrow 0$, although $\delta_R \sim 2$ (see § 3.2). Notice that we get the same first-order formula as in § 4.

Expression (5.13) for the bifurcation velocity is a good approximation even when ϵ is not too small. If we set $\epsilon = 1$ and $Q = RL$, then $a_Q = 1.7549$, $\bar{a}_Q = 1.6180$, and we get from (5.13) $\delta_{RL}^{-1} = 0.01812$, which is very close to the exact result.

Using now (5.12) for sequences $(Q_1 * Q_2)^{*n} * P$ and $Q_2 * (Q_1 * Q_2)^{*n-1} * P$, Q_1 and Q_2 being primary and $Q_1, Q_2 \neq R$, we get coupled equations for $a_{(Q_1*Q_2)^{*n}*P}$ and $a_{Q_1*(Q_2*Q_1)^{*n-1}*P}$, whence the multiplication formula

$$\delta_{Q_1*Q_2} \sim \delta_{Q_1} \delta_{Q_2} \tag{5.14}$$

5.4. The case $Q = R$

The derivation of the above formulae does not work in the case $Q = R$, as $a_R = 1$ and the quantity $a_Q^{-k/\epsilon}$ which was considered as small in the previous case is now equal to 1. One consequence is that the values of superstability for the parameter of the harmonics R^{*n} behave roughly as $1 + c_n \epsilon \ln \epsilon$ (c_n all distinct). Another big difference is that the bifurcation velocity is no longer infinite at $\epsilon = 0$, since $\delta_R = 2$ at $\epsilon = 0$. We shall see that the corrective term is a very slowly varying function of ϵ as $\delta_R \sim 2 + \epsilon / (a_* - 1)$, with $a_* - 1 \sim -\epsilon \ln \epsilon$. For example, when $\epsilon = 10^{-3}$, δ_R is still noticeably larger than its limit value. We find $\delta_R \sim 2.25!$

Proof. For a given parameter $a = 1 + \varphi$, φ being small, the set $\{t_j(\varphi)\}_{j=1, \dots, 2^{n+1}}$ of the iterates of the maximum satisfy the recurrence relation

$$t_{2^n+p}(\varphi) = t_p(\varphi) + |\psi_n(\varphi)| (A_p + B_p \epsilon \ln |\psi_n(\varphi)| + C_p \epsilon + D_p \varphi)$$

where $1 \leq p \leq 2^n$, $\psi_n(\varphi) = t_{2^n}(\varphi)$, and the coefficients A_p, B_p, C_p and D_p are to be found by recursion and depend on the $(2^n - 1)$ numbers α_i defined from the sequence R^{*n} following the rules given in I or in equation (2.10). One has

$$\begin{aligned} A_p &= B_p = -\alpha_{p-2} \dots \alpha_0, & D_p &= p A_p \\ C_p &= -(p-1) \alpha_{p-2} \dots \alpha_0 - \alpha_{p-2} \dots \alpha_0 \ln |t_1(\varphi) \dots t_{p-1}(\varphi)| \end{aligned}$$

whence, as $\text{sgn } \psi_n = (-1)^n$,

$$\psi_{n+1}(\varphi) = \psi_n(\varphi)[\epsilon \ln|\psi_n(\varphi)| + (2^n - 1)\epsilon + \epsilon \ln|t_1(\varphi) \dots t_{2^n-1}(\varphi)| + 2^n\varphi] \tag{5.15}$$

for every small φ .

The value of φ for which $R^{*(n+1)}$ is superstable is the solution $\varphi (> 0)$ of

$$\epsilon \ln|\psi_n(\varphi)| + (2^n - 1)\epsilon + \epsilon \ln|t_1(\varphi) \dots t_{2^n-1}(\varphi)| + 2^n\varphi = 0$$

for example, for sequence R , $\psi_1(\varphi) = \varphi$ and $\varphi_1 = 0$, and for sequence $R * R$, $\psi_2(\varphi) = \varphi(\epsilon \ln \varphi + \epsilon + 2\varphi)$; and φ_2 is the solution of

$$\epsilon \ln \varphi + \epsilon + 2\varphi = 0, \quad \varphi_2 \sim -\frac{1}{2}\epsilon(\ln \epsilon + \ln|\ln \epsilon| + 1 - \ln 2).$$

The calculations quickly become very tedious in this way, and we substitute for sequence $\psi_n(\varphi)$ the sequence $\chi_{n+1}(\varphi) = \psi_{n+1}(\varphi)/\psi_n(\varphi)$ [$\chi_1 = \varphi$]. It is easy to prove that

$$\chi_{n+1}(\varphi) = 2\chi_n(\varphi) + \epsilon + \epsilon \ln \chi_n(\varphi) \tag{5.16}$$

(we use $\ln|t_1(\varphi) \dots t_{2^{n+1}-1}(\varphi)| \sim 2 \ln|t_1(\varphi) \dots t_{2^n-1}(\varphi)| + \ln|\psi_n(\varphi)|$) and φ_n is the root of $\chi_n(\varphi_n) = 0$.

As $\chi_n(\varphi_n) = \chi_{n+1}(\varphi_{n+1}) = 0$, we get from (5.16)

$$\chi_{n-1}(\varphi_n) = \chi_n(\varphi_{n+1}), \quad \chi_{n-2}(\varphi_n) = \chi_{n-1}(\varphi_{n+1}) \dots \chi_1(\varphi_n) = \varphi_n = \chi_2(\varphi_{n+1})$$

whence

$$2\varphi_{n+1} + \epsilon + \epsilon \ln \varphi_{n+1} = \varphi_n. \tag{5.17}$$

The limit value $\varphi_* = a_* - 1$ is then the solution of

$$\varphi_* + \epsilon + \epsilon \ln \varphi_* = 0 \tag{5.18}$$

and

$$\begin{aligned} \delta_R &= \lim_{n \rightarrow \infty} (\partial\varphi_n / \partial\varphi_{n+1}) = \lim_{n \rightarrow \infty} (2 + \epsilon / \varphi_{n+1}) \\ \delta_R &= 2 + \epsilon / \varphi_* \end{aligned} \tag{5.19}$$

which is equation (4.10).

Notice that for $\epsilon = 1$, $\varphi_* \sim 0.28$, and using formula (5.19) we get $\delta_R \sim 5.57$ which is not so bad as the relative error is less than 20%.

6. Conclusion

In I, the authors showed that the whole set of sequences of Metropolis, Stein and Stein has a property of internal similarity; in the present paper, it is shown that this internal similarity is not only of a purely algebraic nature. After a large number of applications of one of these similarity transformations, one gets a transform which is actually a contraction in the parameter space, with a constant ratio that is independent of the general shape of the transform, but depends only on its analytic structure near the maximum.

It is also rather striking to observe that this kind of property exists also in the case of invertible mapping, as encountered in dynamical systems. In particular we have studied

from this point of view the Hénon two-dimensional mapping $T_{a,b}(x,y) \rightarrow (1 - ax^2 + y, bx)$, a and b being parameters, and x and y real quantities (Hénon 1976).

There is no longer a critical point, as this mapping is invertible and multiplies the area by a constant factor b . Choosing for b the same value as Hénon did, $b = 0.3$, it is possible to follow on a computer the bifurcation from a stable period 1 to a stable period 2, to a stable period 4 . . . to a stable period 2^n . The accumulation point for this set of bifurcations is reached at $a_* \sim 1.058\ 048\ .\ .\ .$ when $b = 0.3$. In table 5 we give the values of a corresponding to the occurrence of a stable period 2^n with $0 \leq n \leq 8$ (and to the destabilisation of the period 2^{n-1}). The computations are noticeably more lengthy than in the case of endomorphisms of the line, as it is not possible to use the Newton method for finding the point of superstability which does not exist in this case, as the Hénon mapping has a constant Jacobian. It is rather striking to observe that all these values of a approach the limit a_* in a geometric fashion, and that the bifurcation velocity is the same (up to three digits) as in the case of C^∞ endomorphisms of the line. In order to justify this fact (in a very sloppy way of course!), one might just observe that after a large number of iterations, say N , the elementary area is multiplied by b^N and thus shrinks to zero, as $N \rightarrow \infty$ when $b < 1$, so that the mapping is, in some sense, 'one-dimensional'. We hope to come to this point in a future work.

Table 5. List of the zones of stability and approximations for the sequences R^{**n} in the Hénon mapping. The calculations correspond to $b = 0.3$. The accuracy of δ increases with n . We use formula (3.5) for $\delta(n)$.

n	Period, 2^n	a_s (beginning of stability)	Approximation for δ
0	1	$-(1-b)^2/4 = -0.1225$	
1	2	$3(1-b)^2/4 = 0.3675$	
2	4	$\frac{5+5b^2-6b}{4} = 0.9125$	4.844
3	8	1.026	4.3269
4	16	1.051	4.696
5	32	1.056 536	4.636
6	64	1.057 730 83	4.7748
7	128	1.057 980 8931	4.6696
8	256	1.058 034 452 15	4.6691
9	512	1.058 045 923 04	4.6691
10	1024	1.058 048 379 80	4.6694
11	2048	1.058 048 905 931	

Acknowledgment

The authors are grateful to Dr T W Burkhardt for many useful discussions on the comparison with phase transition problems.

References

Cosnard M Y and Eberhard A 1977 *Séminaire Analyse Numérique No. 274* Lab. Math. Appl., USMG, Grenoble

- Derrida B, Gervois A and Pomeau Y 1977 *C.R. Acad. Sci., Paris A* **285** 43
— 1978 *Ann. Inst. Henri Poincaré* **29** 305
Derrida B and Pomeau Y 1979 *Preprint*
Domb C and Green M S 1976 *Phase Transitions and Critical Phenomena* Vol 6 (New York: Academic)
Feigenbaum M J 1977a *Proc. Como Conf. on Stochasticity of Hamiltonian systems* ed. J Ford
— 1977b *SIAM J.*
Guckenheimer J 1977 *Invent. Math.* **39** 165
Hénon M 1976 *Commun. Math. Phys.* **50** 69
Li T Y and Yorke J A 1975 *Am. Math. Mon.* **82** 985
— 1979 *Trans. Am. Math. Soc.* in the press
May R M 1976 *Nature* **261** 459
Metropolis N, Stein M L and Stein P R 1973 *J. Comb. Theory A* **15** 25
Sarkovskii A N 1964 *Ukr. Math. J.* **16** 61
Stefan P 1977 *Commun. Math. Phys.* **54** 237