

Universal primers for amplification and sequencing a noncoding spacer between the *atpB* and *rbcL* genes of chloroplast DNA

Tzen-Yuh Chiang^{1,4}, Barbara A. Schaal² and Ching-I Peng³

¹Department of Biology, National Cheng-Kung University, Tainan 700, Taiwan

²Department of Biology, Washington University, St. Louis, Missouri 63130, USA

³Institute of Botany, Academia Sinica, Nankang, Taipei 115, Taiwan

(Received January 14, 1998; Accepted April 23, 1998)

Abstract. Universal primers for amplifying and sequencing a noncoding spacer between the *atpB* and *rbcL* genes of the chloroplast DNA were constructed from the published sequences of *Marchantia* (a liverwort), tobacco, and rice. Our results indicate an evolutionary trend of increasing spacer size from liverworts, through mosses, to vascular plants. This *atpB-rbcL* spacer is AT-rich, consistent with other chloroplast noncoding spacers. Due to weak functional constraints, the spacer is evolving rapidly. A sequence identity of 92.2% was observed between spacers of two closely related moss species, *Rhytidiadelphus loreus* and *R. triquetrus*. Insertion/deletion events are common in the evolution of this spacer. A 23 bp deletion occurs in *R. loreus*. Variation is found between two populations of *Amorphophallus henryi* (Araceae) and between individuals in a populations of *Pasania formosana* (Fagaceae). We suggest that this spacer will be useful for molecular systematics at the subspecific, specific, and generic levels and, in some species, for population genetics studies.

Keywords: *Amorphophallus henryi*; *atpB-rbcL* chloroplast spacer; PCR; *Rhytidiadelphus* sp.; Universal primers.

Introduction

The chloroplast genome evolves slowly in general (Clegg et al., 1991), making it an ideal system to assess phylogenetic relationships among genera (e.g., Onagraceae, Conti et al., 1993; Apiaceae, Downie et al., 1996), families (e.g., Asteridae, Olmstead et al., 1992) or higher levels (e.g., seed plants, Chase et al., 1993; monocots, Davis, 1995; angiosperms, Nickrent and Soltis, 1995). Nevertheless, some recent studies have revealed genetic variation between populations based on RFLP's of the total chloroplast genome (see Schaal and Leverich, 1996) or on specific sequences, such as noncoding spacers between several tRNAs of chloroplast DNA (Demesure et al., 1995, 1996).

Several universal primers for amplifying noncoding spacers of the chloroplast genome have been reported (Taberlet et al., 1991; Demesure et al., 1995; Dumolin-Lapegue et al., 1997). Most of the primers were designed for amplifying spacers between tRNA genes, which have been proved variable among species or populations (Demesure et al., 1996). In this study we describe one pair of universal primers for another noncoding spacer region that lies between the large subunit of ribulose-1,5-bisphosphate-carboxylase (*rbcL*) and the beta-subunit of the chloroplast ATP-synthase (*atpB*) genes. Compared to the primers designed for specific taxa (e.g., Rubiaceae,

Manen et al., 1994), the primers proposed here work for plants of a wider phylogenetic array from bryophytes to flowering plants. The spacer is useful for both phylogenetic assessment (Golenberg et al., 1993; Manen et al., 1994; Natali et al., 1995; Savolainen et al., 1994) and species level or population studies.

Materials and Methods

Material Collection

Species of mosses and vascular plants representing a phylogenetic array were collected for testing the primers designed in this study. Two mosses, *Rhytidiadelphus triquetrus* and *R. loreus* (Hylocomiaceae), one fern (*Angiopteris*), one gymnosperm [*Cunninghamia lanceolata* (Taxodiaceae)], three monocots [i.e., *Imperata cylindrica*, *Miscanthus transmorrisonensis* (Poaceae) and *Amorphophallus henryi* (Araceae)], and three dicots [i.e., *Begonia aptera* (Begoniaceae), *Glycine soja* (Fabaceae), and *Pasania formosana* (Fagaceae)] were collected in the field in Taiwan, China, Canada or USA (Table 1). Vouchers are deposited at HAST herbarium. Leaf tissue, fast dried in silica gel after collecting, was powdered in liquid nitrogen and kept in a -70°C freezer.

DNA Isolation

Genomic DNA was extracted by a CTAB method (Doyle and Doyle, 1987) from frozen tissue. Isolated DNA of mosses was purified on low melting agarose gels to remove secondary compounds and RNA (Chiang, 1994).

⁴Corresponding author. Tel: +886-6-2757575 ext. 65525; FAX: +886-6-2742583; E-mail: tychiang@mail.ncku.edu.tw

Table 1. Materials collected for amplification of *atpB-rbcL* spacer of cpDNA with size variation.

Taxon	Sources	Size (bp)
MOSSES		
<i>Rhytidiadelphus loreus</i> (Hypnaceae)	Canada: British Columbia	524
<i>R. triquetrus</i> (Hypnaceae)	USA: Smoky Mt.	549
FERN		
<i>Angiopteris lygodifolia</i> (Marattiaceae)	Taiwan: Yangmingshan	629
GYMNOSPERM		
<i>Cunninghamia lanceolata</i> (Taxodiaceae)	China: Chekiang Prov.	1,000
ANGIOSPERMS		
Monocots:		
<i>Miscanthus transmorrisonensis</i> (Poaceae)	Taiwan: Hohuanshan	800
<i>Imperata cylindrica</i> (Poaceae)	Taiwan: Tainan Co.	800
<i>Amorphophallus henryi</i> (clone 1) (Araceae)	Taiwan: Tainan Co.	800
<i>A. henryi</i> (clone 2) (Araceae)	Taiwan: Kaohsiung City	800
Dicots:		
<i>Glycine soja</i> (Fabaceae)	Taiwan: Taipei Hsien	900
<i>Begonia aptera</i> (Begoniaceae)	Taiwan: Taipei Hsien	841
<i>Pasania formosana</i> (Fagaceae)	Taiwan: Pingtung Hsien	900

Primer Design and PCR

One pair of universal primers were designed by comparing the sequences of *Marchantia* (a liverwort, Umesono et al., 1988), *Angiopteris* (a fern, EMBL X58429), tobacco (Shinozawa et al., 1986), and rice (Nishizawa and Hirai, 1987). Primers, *atpB*-1: 5'-ACATCKARTACKGGACC AATAA-3' and *rbcL*-1: 5'-AACACCAGCTTTRAATCCAA-3', were used for PCR and sequencing. PCR amplification conditions were 30 cycles of 94°C denaturing for 45 s, 49°C annealing for 1 min 15 s, and 72°C extension for 1 min 15 s, followed by 72°C extension for 10 min and 4°C for storing. PCR products were electrophoresed in a 1% agarose gel.

Nucleotide Sequencing

PCR products were separated and eluted using agarose gel purification (BM) and ligated to a pT7 Blue T-vector (Novagen). Plasmid DNA was purified using Wizard Plus SV kit (Promega) and quantified for further sequencing. Cycle sequencing based on dideoxy-mediated chain-termination methodology (Sanger et al., 1977) with *Taq* polymerase was performed using the fmol™ Sequencing System (Promega). Double strands were sequenced from both ends, using primers for PCR amplification and additional primers in between as well, with overlapping of about 100 bp. Clones of mosses and *Amorphophallus henryi* were sequenced completely. DNA fragments from the other species were sequenced only with one of the primers, *atpB*-1, to confirm the nature of the amplified segments.

Results and Discussion

PCR products show size variation of the *atpB-rbcL* spacer among groups. Our results indicate an evolutionary trend of increasing *atpB-rbcL* spacer size, from liv-

erworts (507 bp in *Marchantia*), through mosses (ca. 550 bp in *Rhytidiadelphus*), to vascular plants including ferns (629 bp in *Angiopteris*), conifers (ca. 1,000 bp in *Cunninghamia lanceolata*), monocots (ca. 800 bp) and dicots (ca. 850–950 bp) (Table 1).

This is also the first report of variation of cpDNA sequences between closely related species of bryophytes. 553 nucleotide bases of the *atpB-rbcL* noncoding spacer were aligned (Figure 1). Clones of *Rhytidiadelphus loreus* (EMBL no. L21722) and *R. triquetrus* (EMBL no. L23410) shared 92.2% identity. Substitutions occurred with no bias in transitions (5 times) versus transversions (6 times). Insertion/deletion is a common phenomenon in this spacer. Several length mutations were observed between the two taxa. *Rhytidiadelphus loreus* lacked a three bases (TAG) and a 13 bases (positions 381–403) sequences in contrast to *R. triquetrus*. Additionally, *R. triquetrus* lacked a three bases sequence (TTC) in a different portion of the sequence. The variability of this spacer among species is close to that of the internal transcribed spacer (ITS region) of nuclear ribosomal DNA, in which 87.6–93.7% identity between *Raillardella* species (family Asteraceae, Baldwin, 1992) has been reported.

Sequences (Figure 2) of the spacer in *Amorphophallus henryi* showed variation between two populations. In 810 aligned bases 11 sites (1.4%) were variable. Two indels occurred at positions 169 and 718. Nine substitutions were observed with more transitions (6 sites) than transversions (3 sites), which is consistent with reports for other genes (cf. Li, 1997). The *atpB-rbcL* spacer is AT rich. In the sequence of *A. henryi*, 29.9% of the bases (242 out of 810) were either C or G. Most noncoding spacers and pseudogenes are AT rich due to low functional constraints (Li, 1997). Lack of functional constraints also results in a faster rate of evolution for most spacer regions. Our preliminary data

R-1	1	TACAAC TCAA	TTGATAACTC	TTGCAAGGAT	TAGGTCTGCT	CGACAAATAA
R-t	
R-1	51	GTTTTTTTAA	TCCTATAAAA	AAACATCTTT	GGTATTTAAT	TTTGTCTATT
R-t	
R-1	101	TGTAGACAAA	GTTATTTTAC	TATAACAAAA	CAGTATCAAT	GTATAGTATT
R-t		.T.....
R-1	151	TTTTACATTT	GATGCAACTC	AGATTATTTT	TAGTAAATGA	--T---TTTT
R-t		GC.TAG....
R-1	201	ATTTGTATAA	GCATTGACCT	AATAATCTTT	TGAATGTTAA	ACTAATTAAT
R-t		...C.....
R-1	251	AAAAATAAGA	TTTAATTAAA	AACATAGTAA	GAAATAAAAA	AATAAACTAT
R-t	A.G..	.TA...-....
R-1	301	TACCTAATTC	AATTTATTTT	TTGAAAATAA	AATAAAATTT	TTTTATTTAA
R-t	T
R-1	351	ATATATTTAT	TATTTACTAA	TATTTT TAGA	-----	-----
R-t	G...	TTGTAAAAAA	TATGTATGTC
R-1	401	---CAGTTTT	TAATTTTTTTT	TATTTTAGAT	ACTTATATCT	ATATATAAGT
R-t		TTA..C....T..
R-1	451	AATTAAT TGA	AAGTAATTTT	TTTTCTTCAA	TAAT TAAATG	ATCGAGTTGA
R-t	---
R-1	501	TACTAAATAT	TTTTT-CGAT	ATAATCAGCA	ACTAATTTAT	TACTTCTAAA
R-t	T.A..
R-1	551	TTT				
R-t		...				

Figure 1. Alignment of sequences of *atpB-rbcL* spacer of *Rhytidiadelphus loreus* (R-1) and *R. triquetrus* (R-t), with identical sequences (•) and deletions (-).

on *Pasania formosana* shows much higher diversity (data not shown) in the nucleotide sequence of the *atpB-rbcL* spacer than has been detected in any other plant groups reported previously, such as the taxa of Rubiaceae (Manen et al., 1994; Natali et al., 1995). Here we find variation among individuals within populations of *Pasania*. Therefore, we suggest that this spacer will be useful for molecular systematics at the subspecific, specific, and generic levels and, in some cases, for population genetic studies.

Acknowledgments. We are indebted to T. F. Chiang for lab assistance. This research was granted by National Science Council, ROC to CIP and TYC and by National Science Foundation, USA to BS.

Literature Cited

- Baldwin, B.G. 1992. Phylogenetic utility of the internal transcribed

spacers of nuclear ribosomal DNA in plants: an example from the Compositae. *Mol. Phylog. Evol.* **1**: 3–16.

- Chase, M.W., D.E. Soltis, R.G. Olmstead, D. Morgan, D.H. Les, B.D. Mishler, M.R. Duvall, R.A. Price, H.G. Hills, and Y.L. Qiu. 1993. Phylogenies of seed plants: an analysis of nucleotide sequences from the plastid gene *rbcL*. *Ann. Missouri Bot. Gard.* **80**: 528–580.
- Chiang, T.Y. 1994. Phylogenetics and evolution of the Hylocomiaceae (Mosses, Hypnales) based on sequences of nrDNA ITS and cp DNA *atpB-rbcL* spacer. Ph.D. Dissertation, Department of Biology, Washington University, St. Louis, USA.
- Clegg, M.T., G.H. Learn, and E.M. Golenberg. 1991. Molecular evolution of chloroplast DNA. In R. K. Selander, A. G. Clark and T. S. Whittam (eds.), *Evolution at the Molecular Level*. Sinauer Publishers, Sunderland, pp. 135–149.
- Conti, E., A. Fischbach, and K.J. Sytsma. 1993. Tribal relationships in Onagraceae: implications from *rbcL* sequence data. *Ann. Missouri Bot. Gard.* **80**: 672–685.
- Davis, J.I. 1995. A phylogenetic structure of the monocotyledons,




Figure 2. Nucleotide sequences of *atpB-rbcL* chloroplast DNA spacer of clones (Hen 1 and Hen 2) of *Amorphophalus henryi*, with identical sequences (•) and deletions (■).

- as inferred from chloroplast DNA restriction site variation, and a comparison of measures of clade support. *Syst. Bot.* **20**: 503–527.
- Demesure, B., B. Comps, and R.J. Petit. 1996. Chloroplast DNA phylogeography of the common beech (*Fagus sylvatica* L.) in Europe. *Evolution* **50**: 2515–2520.
- Demesure, B., N. Sodzi, and R.J. Petit. 1995. A set of universal primers for amplification of polymorphic non-coding regions of mitochondrial and chloroplast DNA in plants. *Mol. Ecol.* **4**: 129–131.
- Downie, S.R., D.S. Katz-Downie, and K.J. Cho. 1996. Phylogenetic analysis of Apiaceae subfamily Apioideae using nucleotide sequences from the chloroplast *rpoC1* intron. *Mol. Phylogen. Evol.* **6**: 1–18.
- Doyle, J.J. and J.L. Doyle. 1987. A rapid procedure for DNA purification from small quantities of fresh leaf tissue. *Phytochem. Bull.* **19**: 11–15.
- Dumolin-Lapegue, S., M.H. Pemonge, and R.J. Petit. 1997. An enlarged set of consensus primers for the study of organelle DNA in plants. *Mol. Ecol.* **6**: 393–397.
- Golenberg, E.M., M.T. Clegg, M.L. Durbin, J. Doebley, and D.P. Ma. 1993. Evolution of a noncoding region of the chloroplast genome. *Mol. Phylogen. Evol.* **2**: 52–64.
- Li, W.H. 1997. *Molecular Evolution*. Sinauer Associates, Sunderland, 487 pp.
- Manen, J., A. Natali, and F. Ehrendorfer. 1994. Phylogeny of Rubiaceae-Rubieae inferred from the sequence of a cpDNA intergenic region. *Pl. Syst. Evol.* **190**: 195–211.
- Natali, A., J.F. Manen, and F. Ehrendorfer. 1995. Phylogeny of the Rubiaceae-Rubioideae, in particular the tribe Rubieae: evidence from a noncoding chloroplast DNA sequence. *Ann. Missouri Bot. Gard.* **82**: 428–439.
- Nickrent, D.L. and D.E. Soltis. 1995. A comparison of angiosperm phylogenies from nuclear 18S rDNA and *rbcL* sequences. *Ann. Missouri Bot. Gard.* **82**: 208–234.
- Nishizawa, Y. and A. Hirai. 1987. Nucleotide sequence and expression of the gene for the large subunit of rice ribulose 1,5-bisphosphate carboxylase. *Jap. J. Genet.* **62**: 223–229.
- Olmstead, R.G., H.J. Michaels, K.M. Scott, and J.D. Palmer. 1992. Monophyly of the Asteridae and identification of their major lineages inferred from DNA sequences of *rbcL*. *Ann. Missouri Bot. Gard.* **79**: 249–265.
- Sanger, F., G. Nicklen, and A.R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**: 5463–5467.
- Savolainen, V., J.F. Manen, E. Douzery, and R. Spichiger. 1994. Molecular phylogeny of families related to Celastrales based on 5' flanking sequences. *Mol. Phylogen. Evol.* **3**: 27–37.
- Schaal, B.A. and W.J. Leverich. 1996. Molecular variation in isolated plant populations. *Pl. Sp. Biol.* **11**: 33–41.
- Shinozaki, K., J. Chunwongse, H. Deno, N. Hayashida, T. Kamogashira, A. Kato, J. Kusuda, T. Matsubayashi, B.Y. Meng, J. Obokata, M. Ohme, C. Ohto, H. Shimada, M. Sugita, M. Sugiura, F. Takaiwa, M. Tanaka, N. Tohdoh, K. Torazawa, T. Wakasugi, K. Yamada, K. Yamaguchishinozaki, and N. Zaita. 1986. The complete nucleotide sequence of tobacco chloroplast genome: its gene organization and expression. *EMBO J.* **5**: 2043–2049.
- Taberlet, P., L. Gielly, G. Patou, and J. Bouvet. 1991. Universal primers for amplification of three noncoding regions of chloroplast DNA. *Pl. Mol. Biol.* **17**: 1105–1109.
- Umesono, K., H. Inokuchi, Y. Shiki, M. Takeuchi, Z. Chang, H. Fukuzawa, T. Kohchi, H. Shirai, K. Ohyama, and H. Ozeki. 1988. Structure and organization of *Marchantia polymorpha* chloroplast genome II. gene organization of the large copy region from *rps*'12 to *atpB*. *J. Mol. Biol.* **203**: 299–331.