

Foxcroft, D and Tsertsvadze, A

Universal school-based prevention programs for alcohol misuse in young people.

Foxcroft, D and Tsertsvadze, A (2011) Universal school-based prevention programs for alcohol misuse in young people. *Cochrane Database of Systematic Reviews*, 5 (CD009113).

Doi: 10.1002/14651858.CD009113

This version is available: <https://radar.brookes.ac.uk/radar/items/7f2037d4-c0f7-b1dc-0e03-a2833827fa78/1/>

Available on RADAR: July 2012

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the published version of the journal article. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

# Universal school-based prevention programs for alcohol misuse in young people (Review)

Foxcroft DR, Tsertsvadze A



**THE COCHRANE  
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2011, Issue 5

<http://www.thecochranelibrary.com>



---

Universal school-based prevention programs for alcohol misuse in young people (Review)  
Copyright © 2011 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
BACKGROUND . . . . .	2
OBJECTIVES . . . . .	3
METHODS . . . . .	3
Figure 1. . . . .	5
Figure 2. . . . .	6
Figure 3. . . . .	7
RESULTS . . . . .	8
DISCUSSION . . . . .	13
AUTHORS' CONCLUSIONS . . . . .	15
ACKNOWLEDGEMENTS . . . . .	15
REFERENCES . . . . .	16
CHARACTERISTICS OF STUDIES . . . . .	23
DATA AND ANALYSES . . . . .	120
APPENDICES . . . . .	120
HISTORY . . . . .	123
CONTRIBUTIONS OF AUTHORS . . . . .	123
DECLARATIONS OF INTEREST . . . . .	123
SOURCES OF SUPPORT . . . . .	123
NOTES . . . . .	123
INDEX TERMS . . . . .	123

[Intervention Review]

# Universal school-based prevention programs for alcohol misuse in young people

David R Foxcroft<sup>1</sup>, Alexander Tsertsvadze<sup>2</sup>

<sup>1</sup>School of Health and Social Care, Oxford Brookes University, Oxford, UK. <sup>2</sup>University of Ottawa Evidence-Based Practice Center, Clinical Epidemiology Methods Centre, Ottawa Hospital Research Institute, Ottawa, Canada

Contact address: David R Foxcroft, School of Health and Social Care, Oxford Brookes University, Marston Road, Jack Straws Lane, Marston, Oxford, England, OX3 0FL, UK. [david.foxcroft@brookes.ac.uk](mailto:david.foxcroft@brookes.ac.uk).

**Editorial group:** Cochrane Drugs and Alcohol Group.

**Publication status and date:** New, published in Issue 5, 2011.

**Review content assessed as up-to-date:** 3 March 2011.

**Citation:** Foxcroft DR, Tsertsvadze A. Universal school-based prevention programs for alcohol misuse in young people. *Cochrane Database of Systematic Reviews* 2011, Issue 5. Art. No.: CD009113. DOI: 10.1002/14651858.CD009113.

Copyright © 2011 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Alcohol misuse in young people is cause of concern for health services, policy makers, prevention workers, criminal justice system, youth workers, teachers, parents. This is one of three reviews examining the effectiveness of (1) school-based, (2) family-based, and (3) multi-component prevention programs.

### Objectives

To review evidence on the effectiveness of universal school-based prevention programs in preventing alcohol misuse in school-aged children up to 18 years of age.

### Search methods

Relevant evidence (up to 2002) was selected from the previous Cochrane review. Later studies, to July 2010, were identified from MEDLINE, Cochrane Central Register of Controlled Trials, EMBASE, Project CORK, and PsycINFO.

### Selection criteria

Randomized trials evaluating universal school-based prevention programs and reporting outcomes for alcohol use in students 18 years of age or younger were included. Two reviewers screened titles/abstracts and full text of identified records.

### Data collection and analysis

Two reviewers extracted relevant data independently using an *a priori* defined extraction form. Risk of bias was assessed.

### Main results

53 trials were included, most of which were cluster-randomised. The reporting quality of trials was poor, only 3.8% of them reporting adequate method of randomisation and program allocation concealment. Incomplete data was adequately addressed in 23% of the trials. Due to extensive heterogeneity across interventions, populations, and outcomes, the results were summarized only qualitatively.

Six of the 11 trials evaluating alcohol-specific interventions showed some evidence of effectiveness compared to a standard curriculum. In 14 of the 39 trials evaluating generic interventions, the program interventions demonstrated significantly greater reductions in alcohol use either through a main or subgroup effect. Gender, baseline alcohol use, and ethnicity modified the effects of interventions. Results from the remaining 3 trials with interventions targeting cannabis, alcohol, and/or tobacco were inconsistent.

## Authors' conclusions

This review identified studies that showed no effects of preventive interventions, as well as studies that demonstrated statistically significant effects. There was no easily discernible pattern in characteristics that would distinguish trials with positive results from those with no effects. Most commonly observed positive effects across programs were for drunkenness and binge drinking. Current evidence suggests that certain generic psychosocial and developmental prevention programs can be effective and could be considered as policy and practice options. These include the Life Skills Training Program, the Unplugged program, and the Good Behaviour Game. A stronger focus of future research on intervention program content and delivery context is warranted.

## PLAIN LANGUAGE SUMMARY

### Psychosocial and Developmental Alcohol Misuse Prevention in Schools can be effective

We conducted a Cochrane systematic review of 53 well-designed experimental studies that examined the effectiveness of school-based universal programs for the prevention of alcohol misuse in young people. The studies were divided into two major groups based on the nature of the prevention program: 1) programs targeting specifically prevention or reduction of alcohol misuse and 2) generic programs with wider focus for prevention (e.g., other drug use/abuse, antisocial behavior). In the review we found studies that showed no effects of the preventive program, as well as studies that demonstrated statistically significant effects. There was no easily discernible pattern in program characteristics that would distinguish studies with positive results from those with no effects. Most commonly observed positive effects across programs were for drunkenness and binge drinking. In conclusion, current evidence suggests that certain generic psychosocial and developmental prevention programs can be effective and could be considered as policy and practice options. These include the Life Skills Training Program, the Unplugged program, and the Good Behaviour Game.

## BACKGROUND

### Description of the condition

Alcohol misuse is defined as drinking levels of alcohol that can cause physical, psychological and social problems - both in the short term and the long term. Worldwide, alcohol misuse causes 1.8 million deaths (3.2% of total deaths) and 58.3 million Disability-Adjusted Life Years (DALYs) (4% of total). Accidental injuries are responsible for about one third of the 1.8 million deaths, while neuro-psychiatric conditions are responsible for nearly 40% of the 58.3 million DALYs (WHO 2008).

In many countries heavy episodic or binge drinking is prevalent amongst young people and presents an increased risk for accidents, violence, criminal activity, poorer health and social outcomes. Alcohol consumption is also limited by legislated age-related restrictions, and much alcohol use by young people under the age of 21 (e.g. United States) or 18 (e.g. United Kingdom and some other European countries) is therefore illegal.

The European Union (EU) is the heaviest drinking region of the world, drinking 11 litres of pure alcohol per adult each year (Anderson 2006). More than 1 in 4 deaths among men (aged 15-

29 years) and 1 in every 10 deaths among young women in the EU is alcohol related (Rehm 2005). Young people (aged 15-24 years) are responsible for a high proportion of this burden, with over 25% of youth male mortality and approximately 10% of young female mortality being due to alcohol (Anderson 2006). Some information exists on the extent of social harm in young people, for example a third of a million (6%) 15-16 year old school students in the EU report engaging in fights, and 200,000 (4%) report unprotected sex, due to their own drinking (Anderson 2006).

Amongst young people, early initiation of alcohol use has been shown to be linked to later binge drinking, heavy drinking and alcohol-related problems, in both prospective longitudinal studies (Pitkanen 2005; Warner 2003; Zakrajsek 2006) and large scale cross-sectional epidemiological studies from the United States (Dawson 2008; Hingson 2006; Hingson 2003a; Hingson 2003b). There is some evidence that early consumption may lead to neurological development problems and impairment (AMA 2004), and the Chief Medical Officer for England has recently advised that young people below the age of 15 should not be allowed to drink alcohol (CMO 2009).

## Description of the intervention

The United States Institute of Medicine (Mrazek 1994) proposed a framework for classifying prevention into universal or selective prevention interventions, as a replacement for the previous concepts of primary or secondary prevention. Universal prevention strategies address the entire population within a particular setting (schools, colleges, families, community). The aim of universal prevention is to deter or to delay the onset of a disorder or problem by providing all individuals the information and skills necessary to prevent the problem. Universal prevention programs are delivered to large groups without any prior screening for risk factors, so all members of the population share the same general risk, although the risk may vary greatly among individuals and sub-groups (EMCDDA 2010). In school settings, universal prevention typically takes the form of alcohol awareness education, social and peer resistance skills, normative feedback, or development of behavioural norms and positive peer affiliations. Prevention programs can be either specific curricula delivered as school lessons, or classroom behaviour management programs, and can be educational, psychosocial, or a combination. Psychosocial interventions aim to develop psychological and social skills (e.g. peer resistance) through modelling, understanding, norm-setting and social skill practice, so that young people are less likely to misuse alcohol. Educational interventions aim to raise awareness of the potential dangers of alcohol misuse (e.g. increased knowledge) so that young people are less likely to misuse alcohol (Foxcroft 2002).

## How the intervention might work

In a robust cost-benefit model (Caulkins 2004) it was estimated that even small effect sizes in universal prevention interventions, in terms of delaying initiation into substance use for a few years, could lead to important savings to society over an individual's lifetime. Similarly, the United Kingdom National Institute for Health and Clinical Excellence (NICE 2010) has estimated that a national alcohol misuse prevention program in schools would be a cost effective use of public money if it cost £75 million and achieved at least a 1.4% absolute reduction in alcohol consumption amongst young people, a very small effect size.

Both economic models assumed that delaying onset of alcohol misuse and use would avert some of the long-term adverse health outcomes associated with alcohol consumption. Therefore, an important consideration when evaluating the effectiveness of alcohol misuse prevention programs is how long a program's effects persist, ie a program's duration of impact. Those interventions that show persistence of effects over several years are more useful than those interventions that show some immediate or short-term effects but no evidence of any longer-term duration of impact over several years.

## Why it is important to do this review

A previous Cochrane systematic review (Foxcroft 2002) covered the primary prevention of alcohol misuse amongst young people, with 55 studies included for the period to January 2002. This review was broad in scope, extending across different intervention settings (e.g. schools, families, community, health clinics), age groups (up to age 25 years-old), population focus (universal and selective primary prevention programmes), and study designs (randomised controlled trials, matched comparison studies, interrupted times series studies). In updating the search for this previous Cochrane review, we found a large number of records (n=153, to July 2010) reporting new randomised trials and new results from existing randomised trials.

We therefore decided to narrow the scope of the Cochrane review and produce an updated review of randomised trials evaluating the effectiveness of universal school-based prevention programs for alcohol misuse amongst youth 18 years or younger. This is consistent with other reviews produced by the Cochrane Drugs and Alcohol Group, for example universal school-based prevention of drug misuse (Faggiano 2005). Two other, more focused, reviews have also been produced: one covers universal family-based prevention (Foxcroft 2011a), and the other universal multi-component prevention (Foxcroft 2011b).

Other Cochrane reviews, begun or published since 2002, have also focused on the prevention of alcohol misuse in young people, though typically in young adults including college student populations (Moreira 2008; Coombes 2008). A recent review published by an influential group in an influential book (Babor 2010) has concluded that education and persuasion prevention approaches are ineffective, but this was not a systematic review of the available evidence.

## OBJECTIVES

To systematically review evidence on the effectiveness of universal school-based prevention programs in preventing alcohol misuse in school-aged children up to 18 years of age. The specific aim of this review was to determine if psychosocial and educational prevention programs prevent alcohol misuse compared to standard school curriculum or other types of interventions.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

Randomized controlled trials (individual or cluster design).

### Types of participants

Young people up to 18 years attending school. For this review, young people are defined as children and adolescents.

### Types of interventions

**Experimental** - any universal school-based psychosocial or educational prevention program; psychosocial intervention is defined as one that specifically aims to develop psychological and social skills in young people (e.g., peer resistance) so that they are less likely to misuse alcohol; educational intervention is defined as one that specifically aims to raise awareness of the potential dangers of alcohol misuse so that young people are less likely to misuse alcohol; studies that evaluated interventions aiming specifically at preventing and reducing alcohol misuse as well as generic interventions (e.g., drug education programs, healthy school or community initiatives), or other types of interventions (e.g., screening for alcohol consumption) were eligible for inclusion in the review. **Control** - any alternative prevention program (e.g., school-, family-, office-based, multi-component, other) or standard curriculum

### Types of outcome measures

#### Primary outcomes

1. Any direct self-reported or objective measures of alcohol consumption or problem drinking. Outcome measures related to psychological perception/attitudes or awareness were deemed as indirect and therefore were not considered in this review. As an example, the following outcomes were included and considered as relevant:

2. Alcohol use (yes/no)
3. Alcohol use (quantity, frequency)
4. Drinking 5+ drinks at any one occasion (yes/no)
5. Incidence of drunkenness
6. Note that this list is simply indicative rather than exhaustive. Many authors of potentially relevant studies develop and report their own measures for recording and quantifying the misuse of alcohol.

#### Secondary outcomes

1. Alcohol initiation (age)
2. Drunkenness initiation (age)

### Search methods for identification of studies

All relevant studies published up to 2002 inclusively, evaluating the effectiveness of universal school-based prevention programs in reducing/preventing alcohol use or misuse in students 18 years of age or younger were identified and selected from the previously published Cochrane review (Foxcroft 2002). The selection was not restricted by language or status of publication.

### Electronic searches

Update searches were conducted to identify new relevant evidence for the period of 2002 January to 2010 July. No language restrictions were applied. Details of search terms are given in [Appendix 1](#). The following electronic databases were searched: MEDLINE (2002 January - July Week 4 2010) Cochrane Central Register of Controlled Trials (*The Cochrane Library* 2009, issue 4) EMBASE (2002 January - July Week 4 2010) Project CORK (2002 January - 2009 December) PsycINFO (up to July Week 4 2010)

### Searching other resources


The references of topic-related systematic reviews and included studies were hand searched in order to identify potentially relevant citations. Unpublished reports, abstracts, dissertations, brief and preliminary reports were eligible for inclusion.

### Data collection and analysis

#### Selection of studies

Two independent reviewers (D.F. and A.T.) completed broad screening of titles and abstracts of all identified records (screening level 1). Afterwards, the same reviewers independently assessed full-text reports of all potentially relevant for inclusion records that passed the initial screening level. Differences in opinion arising at both screening levels were resolved through discussions. After bibliographic searches were completed, all the retrieved records were assembled in a database and were de-duplicated (i.e., duplicate records identified and removed). The amount of evidence was maximized by using all companion publications reporting relevant outcomes for any given study. The study flow diagram of records identified from update search conducted in electronic databases is presented in [Figure 1](#).

Figure 1.



### Data extraction and management

Two reviewers (D.F. and A.T.) extracted relevant data independently using *a priori* defined extraction form and entered data into RevMan 5.0.24 (RevMan 2010). Differences in opinion arising during data extraction were resolved through discussions.

### Assessment of risk of bias in included studies

For each study included in the review, two authors (D.F. and A.T.) independently assessed the risk of bias using the Cochrane Collaboration's tool (Higgins 2009, section 8.5.1.) The risk of bias assessment was based on the recommended 6 methodological domains of validity:

1. Adequate sequence generation (High, Low, Unclear risk)
2. Adequate allocation concealment (High, Low, Unclear risk)
3. Blinding of personnel/outcome assessors (High, Low, Unclear risk)
4. Addressing incomplete outcome data (High, Low, Unclear risk)

5. Free of selective outcome reporting (High, Low, Unclear risk)

6. Free of other bias (High, Low, Unclear risk)


Each item was rated with one of three possible responses: yes, no, or unclear. For each response, an explanation was provided. Blinding of participants and program deliverers is not achievable for these sort of interventions, so our assessment of blinding focused on whether outcome assessors were blinded to study condition.

For addressing incomplete outcome data (item #4), a cut-off value of 20% for attrition rate (Fewtrell 2008) and reporting of intention to treat (ITT) analysis were considered. For example, studies with higher attrition rates (> 20%) not reporting ITT analysis were classified as 'No'. Studies with lower attrition rates ( $\leq$  20%) reporting ITT analysis were classified as 'Yes'. If only either of the two criteria was met (e.g.,  $\leq$  20% attrition but no ITT analysis reported), the study was classified as 'Unclear'. For the purpose of this review, the item # 6 was assessed for possibility of confounding (i.e., baseline between study group imbalance in important covariate such as gender and alcohol use) and contamination of program effects (e.g., if clusters of students were randomised to



the experimental or control program within one school). The risk of bias data for included trials was summarized in [Figure 2](#) (risk of bias graph) and [Figure 3](#) (risk of bias summary).

**Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.**



**Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.**

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding (performance bias and detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Aitken 1999	?	?	?	?	?	?
Boutreau 1999	?	?	?	?	?	?
Reid 2004	?	?	?	?	?	?
Boslin 1981	?	?	?	?	?	?
Dubin 1995	?	?	?	?	?	?
Bosni 2001	?	?	?	?	?	?
Bosni 2003	?	?	?	?	?	?
Brewer 1991	?	?	?	?	?	?
Caplan 1992	?	?	?	?	?	?
Coydon 1991	?	?	?	?	?	?
Cook 1994	?	?	?	?	?	?
Eitelman 1986	?	?	?	?	?	?
Dunbar 1986	?	?	?	?	?	?
Durjee 1991a	?	?	?	?	?	?
D'Arino 2002	?	?	?	?	?	?
Eisen 2002	?	?	?	?	?	?
Ellicson 1999	?	?	?	?	?	?
Ellicson 2003	?	?	?	?	?	?
Faggiano 2007	?	?	?	?	?	?
Fur-Holden 2004	?	?	?	?	?	?
Goldberg 2009	?	?	?	?	?	?
Roldaan 2007	?	?	?	?	?	?
Goodstadt 1993	?	?	?	?	?	?
Griffin 2009	?	?	?	?	?	?
Hansen 1998	?	?	?	?	?	?
Hansen 1991	?	?	?	?	?	?
Harit 2003	?	?	?	?	?	?
Johnson 2009	?	?	?	?	?	?
Kellam 2009	?	?	?	?	?	?
Koning 2009	?	?	?	?	?	?
McBride 2000	?	?	?	?	?	?
Morgenstern 2006	?	?	?	?	?	?
Moskowitz 1991	?	?	?	?	?	?
Newman 1992	?	?	?	?	?	?
Newton 2009a	?	?	?	?	?	?
Petrie 1988	?	?	?	?	?	?
Perry 2003	?	?	?	?	?	?
Reddy 2002	?	?	?	?	?	?
Ringsdorf 1991	?	?	?	?	?	?
Ringsdorf 2009	?	?	?	?	?	?
Roeggs 1985	?	?	?	?	?	?
Schinke 2009	?	?	?	?	?	?
Sheehan 1998	?	?	?	?	?	?
Shobana 2009	?	?	?	?	?	?
Speth 2002	?	?	?	?	?	?
St Pierre 2005	?	?	?	?	?	?
Stein 2000	?	?	?	?	?	?
Ver Lier 2009	?	?	?	?	?	?
Viel 2009	?	?	?	?	?	?
Wersch 2008	?	?	?	?	?	?
Wersch 2010	?	?	?	?	?	?
Willemsen 1991	?	?	?	?	?	?
Williams 1998	?	?	?	?	?	?

### Unit of analysis issues

Additional validity threats were ascertained regarding appropriate unit of analysis depending whether the randomisation was implemented at individual- or cluster-level (see Tables of Characteristics of included studies).

### Dealing with missing data

If important data was missing, attempts were made to contact the authors of included studies.

### Assessment of reporting biases

The extent of publication bias was to be assessed through visual inspection of asymmetry and running the regression-based method for a funnel plot (Egger 1997; Peters 2008).

### Data synthesis

The results of the review were arranged according to two strata: 1) interventions targeting specifically prevention or reduction of alcohol misuse and 2) generic interventions with wider focus for prevention (e.g., other drug use/abuse, antisocial behavior). The statistical pooling of results of individual studies was planned conditional on the absence of heterogeneity with respect to study populations (e.g., baseline characteristics, gender), interventions (e.g., type, differences in target/focus), and outcome measures (e.g., different tools, instruments, scales) as well as the methodology of conduct (e.g., units of randomisation and analysis, cluster vs. individual trials).

### Subgroup analysis and investigation of heterogeneity

The extent of heterogeneity was planned through examination of forest plots (Chi square statistic and p-value;  $I^2$  statistic) and qualitative subgroup analysis. The subgroup analyses would explore whether or not the effects of universal school-based prevention programs differed in certain subgroups of study participants. The following *a priori* determined participant-level subgroups were based on age, race (Caucasians vs. Blacks, Hispanics), gender, and levels of alcohol use/consumption (at baseline).

### Sensitivity analysis

These analysis were planned, if data allowed, to investigate whether the effects of universal school-based prevention programs in reducing alcohol misuse were different in the following trial-level defined groups:

1. Cluster (ones appropriately analysed)- vs. individually randomised trials.
2. Cluster-randomized trial appropriately analysed (i.e., units of randomisation and analysis are matching) vs. cluster-randomised trial inappropriately analysed (i.e., units of randomisation and analysis not matching).
3. Trials with attrition > 20% (1<sup>st</sup> follow-up) vs. trials with attrition ≤ 20% (1<sup>st</sup> follow-up)

## RESULTS

### Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

See: [Characteristics of included studies](#) and [Characteristics of excluded studies](#).

### Results of the search

Initially, we examined the previously published Cochrane review (Foxcroft 2002) to identify trials published up to 2002 January inclusively by applying our eligibility criteria (see the Methods section). In total, 27 trials were deemed eligible for inclusion in the review (Allison 1990; Beaulieu 1998; Botvin 1984; Botvin 1995; Botvin 2001; Brewer 1991; Caplan 1992; Clayton 1991; Cook 1984; Dielman 1986; Durrant 1986; Duryea 1984a; Ellickson 1990; Goldberg 2000; Goodstadt 1983; Hansen 1988; Hansen 1991; McBride 2000; Moskowitz 1984; Newman 1992; Perry 1988; Ringwalt 1991; Scaggs 1985; Schinke 2000; Sheehan 1996; Wilhelmsen 1994; Williams 1968).

The updated electronic searches (2002 January - 2010 July) identified 1,874 bibliographic records (1,684 through MEDLINE, EMBASE, and PsycINFO; 138 through Project Cork; 52 through the Cochrane Trial Register). The process of de-duplication resulted in 1,801 unique bibliographic records. Through the screening of titles and abstracts, 1,648 records were excluded as obviously irrelevant. The full text reports of the remaining 153 records were examined of whom 113 were excluded due to ineligibility of intervention (n=79; selective, indicated, family-based, multi-component, other), design (n=4; non-randomised study), study participants age (n=16; age > 18 years), and outcomes (n=14; non-alcohol related). The screening process left 40 records representing 26 unique trials, which were included in this review (Bond 2004; Botvin 2003; D'Amico 2002; Eisen 2002; Ellickson 2003; Faggiano 2007; Furr-Holden 2004; Goldberg 2007; Griffin 2009; Johnson 2009; Kellam 2008; Koning 2009; Hecht 2003;

Morgenstern 2009; Newton 2009a; Perry 2003; Reddy 2002; Ringwalt 2009; Sloboda 2009; Spoth 2002; St. Pierre 2005; Sun 2008; van Lier 2009; Vogl 2009; Werch 2008; Werch 2010). The study flow diagram of the update search is depicted in Figure 1. In total, 53 trials evaluating universal school-based alcohol misuse prevention programs in young people were included in the review. Of the 53 trials, 50 were published in peer-reviewed journals and three were unpublished dissertations (Brewer 1991; Durrant 1986; Scaggs 1985). The following 12 trials were reported in multiple companion publications: Clayton 1991 (Clayton 1996; Lynam 1999), Dielman 1986 (Dielman 1989; Wynn 2000), Duryea 1984a (Duryea 1984b; Duryea 1988), Eisen 2002 (Eisen 2003), Ellickson 1990 (Ellickson 1993a; Ellickson 1993b), Faggiano 2007 (Faggiano 2008; Vigna-Taglianti 2009; Faggiano 2010; Caria 2010), Hecht 2003 (Kulis 2005; Kulis 2007a; Kulis 2007b), McBride 2000 (McBride 2003; McBride 2004), Newton 2009a (Newton 2009b), Perry 1988 (Perry 1989), Ringwalt 2009 (Ringwalt 2010), and Spoth 2002 (Spoth 2005; Spoth 2008).

### Included studies

All 53 included studies were parallel-group randomised controlled trials. The unit of randomisation in 46 trials was a cluster (e.g., school, class) and in 7 trials - an individual student (Brewer 1991; Cook 1984; D'Amico 2002; Duryea 1984a; Werch 2008; Werch 2010; Williams 1968).

The total number of students randomised in cluster-randomized trials ranged from 86 (Hansen 1988) to 19,529 (Sloboda 2009). The total number of students randomised in trials with individual unit of randomisation ranged from 54 (Brewer 1991) to 416 (Werch 2010).

Forty-one trials were conducted in North America (US and Canada), 6 trials in Europe (Austria, Belgium, Greece, Italy, Spain, Sweden, Netherlands, Norway, Germany) and 6 trials in Australia. One trial was conducted in India (Reddy 2002) and one trial in Swaziland (Perry 1988). Two trials were conducted in multiple countries (Faggiano 2007; Perry 1988).

The study participants' age at baseline in the included trials ranged from 5 years (Furr-Holden 2004) to 18 years (D'Amico 2002; Perry 1988; Sun 2008). Authors of 9 trials failed to report the age of study participants at baseline (Botvin 2003; Ellickson 2003; Goodstadt 1983; Griffin 2009; Johnson 2009; Perry 2003; Ringwalt 2009; Spoth 2002; St. Pierre 2005). The proportion of males amongst the included trials ranged from 36.5% (Werch 2010) to 62% (Griffin 2009). Two trials enrolled only male students (Goldberg 2000; Williams 1968). The gender-specific proportion was not reported for 10 trials (Allison 1990; Beaulieu 1998; Botvin 1984; Dielman 1986; Duryea 1984a; McBride 2000; Moskowitz 1984; Newman 1992; van Lier 2009; Wilhelmsen 1994). The distribution of ethnic background of study participants across trials varied. Many trials conducted in North America included mixtures of Caucasian, Black American,

Asian, and Hispanic students. The majority (> 70%) of students were Caucasians in 8 trials (Botvin 1984; Botvin 1995; Clayton 1991; Ellickson 2003; Goldberg 2007; Moskowitz 1984; Spoth 2002; St. Pierre 2005) and Black Americans in three trials (Caplan 1992; Furr-Holden 2004; Griffin 2009). In one trial (Beaulieu 1998), all study participants were Black Americans and in another (Schinke 2000) - Native Americans. The ethnic composition of the study sample was not reported for 24 trials (Allison 1990; Bond 2004; Brewer 1991; Dielman 1986; Durrant 1986; Duryea 1984a; Faggiano 2007; Goldberg 2000; Goodstadt 1983; Johnson 2009; Kellam 2008; Koning 2009; McBride 2000; Morgenstern 2009; Newman 1992; Newton 2009a; Perry 1988; Reddy 2002; Scaggs 1985; Sheehan 1996; van Lier 2009; Vogl 2009; Wilhelmsen 1994; Williams 1968).

In all 53 trials, universal school-based prevention programs were implemented. In 39 trials (71%), the target of the intervention programs was of generic nature, focusing on prevention of multiple factors (i.e., alcohol, tobacco, drugs, anti-social behavior). Programs focusing exclusively on the prevention of alcohol misuse were evaluated in 11 trials (Dielman 1986; Duryea 1984a; Goodstadt 1983; McBride 2000; Morgenstern 2009; Newman 1992; Perry 1988; Sheehan 1996; Vogl 2009; Wilhelmsen 1994; Williams 1968). In the three remaining trials, the focus of preventive intervention was misuse of alcohol-cannabis (Newton 2009a), drug-alcohol (Goldberg 2007) or tobacco only (Reddy 2002). In 11 trials at least two different intervention programs were compared (Brewer 1991; D'Amico 2002; Goodstadt 1983; Hansen 1988; Hansen 1991; Koning 2009; Perry 2003; Reddy 2002; Spoth 2002; Sun 2008; Werch 2008). For example, in one trial, three groups of randomised students received Behaviour Image Model (BIM)-based brief multiple behaviour interventions: Goal Survey, Goal Survey plus Contract, and Goal Survey plus Consult (Werch 2008). In another trial, three programs of alcohol education were compared: cognitive, decision-making, and values-clarification interventions (Goodstadt 1983).

The components of the evaluated intervention programs in the majority of trials were the promotion of awareness (e.g., benefits, consequences, risks), resilient behaviour, change in normative beliefs/attitudes, self-esteem, social networking, peer resistance, as well as the development of problem solving, refusal, and/or decision-making skills. The duration of intervention programs across the included trials ranged from a single 50-minute session (D'Amico 2002) to 3 years (Botvin 1995; Botvin 2003).

In the majority of trials (85%), the effectiveness of prevention programs was compared to that of a standard curriculum. In one trial that evaluated the effectiveness of random alcohol-level testing, the control group of students received only deferred testing (Goldberg 2007).

The outcome measures of alcohol use differed greatly across the trials. For example, the outcomes varied with respect to their definition (e.g., alcohol use, frequency of use, mean number of drinks, proportion of alcohol non-users, weekly drinking, hard liquor use,

frequency of drunkenness, drunkenness in the last month, incidence of drinking and driving, binge drinking), scales of measurement (means, percentages, odds ratios, risk ratios), and the period to which they pertained (e.g., past month, past 2 months, current, past year, ever). The last follow-up at which the outcomes were measured ranged from one month (Werch 2008) to 12 years (Kellam 2008) post-randomisation.

### Excluded studies

Many studies were excluded at screening because they clearly did not meet the inclusion criteria. Forty-eight studies required closer scrutiny before they were excluded on the basis that they did not meet the exclusion criteria. These 48 excluded studies are listed in the [Characteristics of excluded studies](#) table.

### Risk of bias in included studies

The assessment results of risk of bias for the included trials are presented in [Figure 2](#) and [Figure 3](#). All trials were randomised.

### Allocation

The adequate method of randomisation and program allocation concealment was reported only for about 3.8% of the trials. It was unclear whether the remaining 96% of the trials utilized adequate methods for randomisation or program allocation concealment.

### Blinding

No blinding of study personnel was carried out and it was unclear whether or not blinding of outcome assessors was carried out in the reviewed trials; this information was not explicitly reported. It is difficult to see how blinding of students or teachers or program deliverers could be achieved and this is a methodological limitation of such social and preventive intervention studies.

### Incomplete outcome data

Incomplete data was adequately addressed in about 23% of the trials and this information was unclear for about 31% of the trials. The attrition rates (at first follow-up) of 26 trials were acceptable ( $\leq 20\%$ ) and for 21 trials not acceptable ( $> 20\%$ ). One trial reported no loss to follow-up (Brewer 1991). The attrition rates were not reported for 6 trials (D'Amico 2002; Durrant 1986; Kellam 2008; McBride 2000; Morgenstern 2009; Newman 1992). see [Characteristics of included studies](#).

### Selective reporting

The majority of the trials (88.5%) were free of selective outcome reporting, whereas only 11.5% of them were not

### Other potential sources of bias

A quarter of all trials were found free of other bias (i.e., confounding, contamination), whereas results in 42% of the trials were deemed to be prone to confounding, contamination, or both. It was unclear for 32.7% of the trials whether or not their results may have been biased due to confounding and/or contamination. Of the 46 cluster-randomised trials, 37 reported at least some efforts of adjusting the intervention effect estimates for clustering effects, whereas for 9 trials, no such efforts were reported (Beaulieu 1998; Durrant 1986; Goodstadt 1983; Griffin 2009; Perry 1988; Scaggs 1985; Schinke 2000; Sheehan 1996; Wilhelmsen 1994). In 7 other trials, students were randomised individually (Brewer 1991; Cook 1984; D'Amico 2002; Duryea 1984a; Werch 2008; Werch 2010; Williams 1968).

The results based on ITT analysis were reported for only 12 out of 52 trials (Bond 2004; Brewer 1991; Eisen 2002; Faggiano 2007; Furr-Holden 2004; Goldberg 2007; Kellam 2008; Morgenstern 2009; Perry 2003; Ringwalt 2009; Spoth 2002; Vogl 2009). For three trials, it was unclear whether or not the reported results were ITT-based (Hecht 2003; Newman 1992; van Lier 2009).

The instruments or questionnaires used for measurement of alcohol misuse/consumption were reported to be validated for only 24 trials (Beaulieu 1998; Bond 2004; Cook 1984; D'Amico 2002; Dielman 1986; Furr-Holden 2004; Goldberg 2000; Goodstadt 1983; Griffin 2009; Hansen 1991; McBride 2000; Moskowitz 1984; Newman 1992; Newton 2009a; Perry 1988; Perry 2003; Scaggs 1985; Spoth 2002; Sun 2008; Vogl 2009; Werch 2008; Werch 2010; Wilhelmsen 1994; Williams 1968). It was unclear whether or not the outcome measures reported in the remaining 29 trials had been validated.

### Effects of interventions

see [Characteristics of included studies](#)

### Quantitative data synthesis

The reviewers could not pool the results from individual trials due to heterogeneity in study populations (baseline characteristics), interventions (differences in target/focus), and the outcome measures of alcohol misuse (different tools, instruments, scales, outcome definitions). For example, Bond 2004 reported a whole school intervention which included teaching resources and a school liaison officer, with 20 sessions; whereas Dielman 1986 reported a 4-session intervention that focused on awareness and refusal skills. Similarly, in the study by Faggiano 2007, outcomes were drinking frequency and drunkenness measures, but in the study by Furr-Holden 2004, outcomes were drinking without parents or without permission. More generally, there is a lack of clear information about the content of the interventions, and this is necessary for ensuring that appropriate comparisons and pooling of results is achieved. This is a general problem for the prevention

field, as identified by Abraham 2008. Therefore, the main results are presented in tabular form and compared in the style of a narrative systematic review.

### 1. Alcohol-Specific Programs

This section included 11 trials that evaluated the effectiveness of universal school-based intervention programs specifically focusing on the prevention of alcohol misuse in young students (Dielman 1986; Duryea 1984a; Goodstadt 1983; McBride 2000; Morgenstern 2009; Newman 1992; Perry 1988; Sheehan 1996; Vogl 2009; Wilhelmsen 1994; Williams 1968).

In 5 trials (Duryea 1984a; Goodstadt 1983; Newman 1992; Sheehan 1996; Williams 1968), the intervention effects on alcohol misuse (e.g., alcohol use in past year, frequency of drinking, mean number of drinks, proportion of weekly drinkers, drinking and driving in the past month) were not statistically significantly different from those of standard curriculum at the follow-up from 6 months to 3 years post-randomisation. For example, the mean numbers of drinking occasions in the intervention and control groups in one trial were 2.06 (1.11) vs. 2.05 (0.97), respectively (Duryea 1984a). In another trial (Newman 1992), the corresponding mean monthly frequencies of drinking occasions were 3.06 (intervention group) vs. 3.43 (control group). In one trial (Sheehan 1996), although the proportions of weekly drinkers increased in both the intervention (from 10% to 36%) and the control group (from 13% to 34%), the between-group difference was not statistically significant (36% vs. 34%,  $p=0.09$ ). In the same trial, there was no difference in drinking and driving frequency between the intervention and control groups regardless of drunk driving status at baseline. In one trial (Williams 1968), in the intervention and control groups, the proportion of students using alcohol in the past year was 85% ( $p>0.05$ ). Note that, three (Duryea 1984a; Goodstadt 1983; Williams 1968) of the 5 trials had a relatively small sample size of randomised students (range: 155-540).

In the remaining 6 trials, the intervention groups had statistically significant reductions in the outcomes of alcohol misuse compared with control groups (McBride 2000; Dielman 1986; Morgenstern 2009; Perry 1988; Vogl 2009; Wilhelmsen 1994). For example, in two trials, the post-intervention effects were modified by prior drinking experience or gender (Dielman 1986; Vogl 2009). Specifically, one-to-three years post-randomisation, significantly greater reduced rates of mean weekly alcohol use were found in subgroups of students with prior drinking experience (Dielman 1986) or female students (Vogl 2009). In the same trials, the between-group differences in the mean weekly alcohol use rates amongst the subgroups of students without prior drinking experience (Dielman 1986) or males (Vogl 2009) were not statistically significant. In another study (McBride 2000), the intervention compared to standard curriculum, significantly reduced risky alcohol consumption (at least once a month) in baseline non-drinkers (21.5% vs. 32.5%,  $p<0.05$ ) but not in baseline drinkers 20 months after the randomization. Note that in this trial, the intervention group at baseline

included significantly more non-drinkers than the control group. In the trial by Morgenstern and colleagues (Morgenstern 2009), students in the intervention group, compared to the standard curriculum group, experienced significantly reduced risk of lifetime binge drinking at both follow-ups: four months (OR=0.56, 95% CI: 0.41, 0.77) and 12 months (OR=0.74, 95% CI: 0.57, 0.97) after the randomisation. The intervention and standard curriculum groups were not significantly different in the post-treatment lifetime alcohol use and drunkenness. In the World Health Organization (WHO) Collaborative Study (Perry 1988), the peer-led program, compared to the teacher-led program or standard curriculum, significantly reduced the immediate post-intervention mean alcohol consumption (using a composite alcohol score) but only in baseline non-drinkers (3.15 vs. 3.46 vs. 3.52,  $p < 0.005$ ). In the same study, the post-intervention mean alcohol consumption in baseline drinkers did not significantly differ for the three groups (5.14 vs. 5.84 vs. 5.71, respectively). Finally, one trial (Wilhelmsen 1994) compared the effectiveness of two programs (Highly Role-Specified and Less Role-Specified) and standard curriculum, and found a significantly lower 4 months post-intervention mean frequency of alcohol use (6-point scale) in the Highly Role-Specified intervention group ( $0.53 \pm 1.4$ ) versus the Less Role-Specified intervention ( $0.90 \pm 1.0$ ) or the control/standard curriculum ( $0.69 \pm 1.3$ ).

The duration of post-intervention impact (i.e., difference between the duration of intervention and last follow-up to which positive result persisted) for the 6 trials (McBride 2000; Dielman 1986; Morgenstern 2009; Perry 1988; Vogl 2009; Wilhelmsen 1994) ranged from 0 days (Vogl 2009) to 2 years (Dielman 1986).

### 2. Generic Programs

This section included 39 trials that evaluated the effectiveness of universal school-based intervention programs with respect to the prevention of multiple factors such as misuse of alcohol, tobacco, drugs, and anti-social behaviour in young students (Allison 1990; Beaulieu 1998; Bond 2004; Botvin 1984; Botvin 1995; Botvin 2001; Botvin 2003; Brewer 1991; Caplan 1992; Clayton 1991; Cook 1984; D'Amico 2002; Durrant 1986; Eisen 2002; Ellickson 1990; Ellickson 2003; Faggiano 2007; Furr-Holden 2004; Goldberg 2000; Griffin 2009; Hansen 1988; Hansen 1991; Johnson 2009; Kellam 2008; Koning 2009; Hecht 2003; Moskowitz 1984; Perry 2003; Ringwalt 1991; Ringwalt 2009; Scaggs 1985; Schinke 2000; Sloboda 2009; Spoth 2002; St. Pierre 2005; Sun 2008; van Lier 2009; Werch 2008; Werch 2010).

In 24 trials, there was no statistically significant difference in the effectiveness between the intervention programs and the control/standard curriculum groups (Allison 1990; Beaulieu 1998; Bond 2004; Botvin 2003; Brewer 1991; Clayton 1991; D'Amico 2002; Durrant 1986; Ellickson 1990; Furr-Holden 2004; Goldberg 2000; Hansen 1988; Hansen 1991; Johnson 2009; Koning 2009; Moskowitz 1984; Perry 2003; Ringwalt 1991; Ringwalt 2009; Spoth 2002; St. Pierre 2005; Sun 2008; Werch 2008; Werch



2010). The follow-up period when the alcohol use outcome measures were ascertained in these trials ranged from one month to 10 years post-randomisation (Werch 2008; Clayton 1991). In 7 trials (Hansen 1988; Hansen 1991; Koning 2009; Perry 2003; Spoth 2002; Sun 2008; Werch 2008), two or more different intervention programs or their combinations were also compared and no significant differences between the effectiveness of these programs (e.g., social influences curriculum, affective education, drug abuse resistance education, life skills training, strengthening families program, cognitive perception information, behavioural skills curriculum, behaviour image model-based interventions) on measures of alcohol use were observed. The number of randomized students across 19 of the 24 trials ranged from 54 (Brewer 1991) to 8,338 (Ringwalt 2009). The randomised number of students for the remaining five trials was not reported (Botvin 2003; Clayton 1991; Durrant 1986; Goldberg 2000; St. Pierre 2005). In one large trial (83 randomised clusters and 17,320 participating students) conducted by Sloboda and colleagues (Sloboda 2009), the intervention program (i.e., take charge of your life) had a significantly negative effect compared to the standard curriculum in preventing alcohol use within 30 days (45.7% vs. 41.9%, OR = 1.09, 95% CI: 1.01, 1.18). In the remaining 14 trials, the program interventions demonstrated significantly greater reductions in alcohol use with respect to main or subgroup effects (i.e., gender, ethnicity, baseline alcohol use status) compared to standard curriculum (Botvin 1984; Botvin 1995; Botvin 2001; Caplan 1992; Cook 1984; Eisen 2002; Ellickson 2003; Faggiano 2007; Griffin 2009; Kellam 2008; Hecht 2003; Scaggs 1985; Schinke 2000; van Lier 2009). The number of randomised students across 11 of the 14 trials ranged from 127 (Scaggs 1985) to 6,900 (Hecht 2003). The randomized number of students for the remaining three trials was not reported (Botvin 1995; Faggiano 2007; Kellam 2008). In two trials, the single intervention program - Life Skills Training (LST) was delivered through formal teacher, older students, or video training (Botvin 1984; Botvin 1995). In the first trial (Botvin 1984), the peer-led program significantly reduced frequency of drunkenness (ANCOVA-based  $F=4.54$ ,  $p < 0.01$ ) and the amount of consumption per occasion (ANCOVA-based  $F=5.10$ ,  $p < 0.006$ ) compared to the teacher-led program or standard curriculum at 4 months of follow-up. In the second trial (Botvin 1995), the LST program delivered either through teacher or video training was significantly more effective ( $0.34 \pm 0.02$  or  $0.33 \pm 0.03$ , respectively) in reducing the mean number of drunkenness episodes in the last month compared to standard curriculum ( $0.40 \pm 0.02$ ) at 6 months of follow-up. In the same trial, post-intervention alcohol use (monthly or weekly) and the frequency of three or more drinks per occasion did not significantly differ between the intervention programs and the control group. In another trial that evaluated the LST program (Botvin 2001), the program was significantly more effective in reducing binge drinking (> 5 drinks per occasion) at one year (OR = 0.41, 95% CI: 0.18, 0.93) and two

years (OR = 0.40, 95% CI: 0.22, 0.74) of follow-up. In one trial including ethnically diverse composition of students, three different versions of one program (Multicultural, Mexican American, and Black/White) were compared and Multicultural and Mexican versions of the program were significantly more effective in reducing the mean number of alcohol drinks compared to Black/White version of the program (Hecht 2003). Similarly, 8 other trials demonstrated statistically significant superiority in the main effects of intervention programs for reducing alcohol use (e.g., 3+ drinks per occasion, daily hard liquor use, monthly frequency of alcohol drinking, 4+ drinking weekly, lifetime alcohol abuse, alcohol use in the past week) compared to standard curricula at one to six years of follow-up (Caplan 1992; Cook 1984; Faggiano 2007; Griffin 2009; Kellam 2008; Scaggs 1985; Schinke 2000; van Lier 2009). For example, in the Faggiano et al. trial, the intervention program, compared to standard curriculum, significantly reduced any or frequent drunkenness in the past month at three months (any drunkenness; OR = 0.69, 95% CI: 0.48, 0.99) and 18 months (any drunkenness; OR = 0.80, 95% CI: 0.67, 0.97) of follow-up (Faggiano 2007). In another trial (van Lier 2009), a significantly reduced growth rate of alcohol use (in the past week) for the intervention group - Good Behaviour Game program (GBG) was shown after 3-6 years of follow-up (between level slope estimate for GBG:  $\beta = -0.43$ ,  $p < 0.05$ ). However, no such reductions were found for alcohol use during last month ( $\beta = -0.31$ ,  $p > 0.05$ ) or last year ( $\beta = 0.05$ ,  $p > 0.05$ ).

In three trials, intervention programs were shown to be significantly more effective than standard curriculum but only in certain subgroups, which were defined by gender (Faggiano 2007), ethnicity (Eisen 2002), and baseline alcohol use status (Ellickson 2003). For example, in one trial (Faggiano 2007), the intervention program was significantly more effective in reducing any drunkenness compared to standard curricula in male (OR = 0.64, 95% CI: 0.49, 0.85) but not in female students (OR = 0.86, 95% CI: 0.63, 1.18) at three months follow-up. The trial by Eisen et al., showed that the intervention program was significantly more effective in reducing alcohol use in the past 30 days and binge drinking (three or more drinks per occasion) in Hispanics but not in non-Hispanic students after two years of follow-up. Similarly, in one trial (Ellickson 2003), the significant effect of the intervention compared to the standard curriculum in reducing overall alcohol misuse was observed after 18 months of follow-up in baseline alcohol users (1.78 vs. 2.23,  $p < 0.05$ ) but not in baseline alcohol non-users (0.22 vs. 0.30,  $p > 0.05$ ).

The duration of post-intervention impact in the 14 trials (Botvin 1984; Botvin 1995; Botvin 2001; Caplan 1992; Cook 1984; Eisen 2002; Ellickson 2003; Faggiano 2007; Griffin 2009; Kellam 2008; Hecht 2003; Scaggs 1985; Schinke 2000; van Lier 2009) ranged from 0 days (Ellickson 2003; Griffin 2009; Caplan 1992; Botvin 2001) to 10 years (Kellam 2008).

### 3. Other Programs

This section included three trials that evaluated the effectiveness of universal school-based intervention programs whose focus was the prevention of alcohol-cannabis (Newton 2009a), drug-alcohol (Goldberg 2007), or tobacco only (Reddy 2002). In all three trials, students were individually randomised to intervention or control groups. In two trials the numbers of randomized students were 1,296 (Newton 2009a) and 4,776 (Reddy 2002). No sample size (i.e., number randomised) was reported for the third trial (Goldberg 2007).

In one trial (Goldberg 2007), the mean past month/year illicit drug and alcohol use index (ranging from 0 = no use to 3=heavy use) in students allocated to the intervention program –a random Drug and Alcohol Testing (DAT) did not significantly differ from that in students allocated to only deferred random DAT after two years of follow-up (past month: 0.165 vs. 0.261,  $p > 0.05$ ; past year: 0.917 vs. 1.033,  $p > 0.05$ ).

In the remaining two trials (Newton 2009a; Reddy 2002), the intervention programs were shown to be statistically significantly better in reducing certain alcohol use outcome measures compared to standard curricula. In the first trial (Newton 2009a), students in the intervention group ( $-0.63 \pm 1.14$ ) compared to the standard curriculum group ( $5.30 \pm 1.50$ ) had a significantly reduced ‘average weekly alcohol use’ from baseline to 18 months of follow-up ( $p < 0.02$ ). In the same trial, the between-group differences with respect to ‘frequency of drinking to excess’ and ‘harms related to own use of alcohol’ were not statistically significant ( $p > 0.05$ ). In the second trial (Reddy 2002), both the school-based program alone (i.e., health-related information and dissemination among youth) or combined with a family-based program were significantly more effective than the standard curriculum in reducing alcohol use (‘ever having a drink of alcohol’) after 17 months of follow-up (0.128 vs. 0.144 vs. 0.288, respectively;  $p < 0.001$ ).

The duration of post-intervention impact in the 3 trials (Goldberg 2007; Newton 2009a; Reddy 2002) ranged from 1 month (Reddy 2002) to 1 year (Goldberg 2007; Newton 2009a).

### Subgroup analysis

Given the fact that studies could not be pooled in this review, the extent of between-study statistical heterogeneity in the intervention effects (e.g., forest plots; Chi square statistic and  $p$ -value;  $I^2$  statistic) could not be assessed quantitatively.

The study-level subgroup analysis qualitatively exploring whether or not the effect of any given universal school-based prevention program differed across the subgroups defined by age, gender, ethnicity, or prior alcohol use of study participants could not be carried out due to unavailability or non-comparability of relevant data. For example, there were only four trials that included only males (Goldberg 2000; Williams 1968), Black Americans (Beaulieu 1998), or Native Americans (Schinke 2000) and the interventions evaluated in these trials differed.

Within-study subgroup effects of the interventions (if reported

that such effects existed) are qualitatively summarized in the Results section (see ‘the effects of interventions’ sub-section) and presented in [Characteristics of included studies](#).

### Sensitivity analysis

The study-level sensitivity analysis qualitatively exploring whether or not the effect of any given universal school-based prevention program differed across methodological aspect defined by unit of randomisation (individual vs. cluster), the appropriateness of analysis (matching units of randomisation and analysis), and attrition rates ( $> 20\%$  vs.  $\leq 20\%$ ) was performed but could not reveal any specific methodological aspect(s) that would potentially account for differences in the study results; the reason being that the studies evaluating the same or similar intervention program(s) reported relatively uniform results. For example, all trials that evaluated the Life Skills Training (LST) program yielded positive results in favour of the intervention (Botvin 1984; Borvin 1995; Botvin 2001; Botvin 2003; Schinke 2000; Spoth 2002). Similarly, two of the three trials that evaluated the GBG program (van Lier 2009, Furr-Holden 2004, Kellam 2008) demonstrated positive results in favour of the intervention. Trials that evaluated the ALERT (Ellickson 1990; Ellickson 2003; Ringwalt 2009; St. Pierre 2005) or drug abuse resistance education program (DARE) (Clayton 1991; Perry 2003; Ringwalt 1991) showed no effects (i.e., statistically non-significant).

### Publication bias

Many studies did not report sufficient information for calculation of effect sizes for use in a funnel plot (Allison 1990; Botvin 1984; Caplan 1992; Duryea 1984a; Eisen 2002; Ellickson 1990; Ellickson 2003; Furr-Holden 2004; Goldberg 2000; Goldberg 2007; Goodstadt 1983; Hansen 1988; Hansen 1991; Hecht 2003; Johnson 2009; Moskowitz 1984; Newman 1992; Newton 2009a; Perry 2003; Sheehan 1996; St. Pierre 2005; van Lier 2009) and therefore we were not able to construct a funnel plot and assess risk of publication bias for this review.

## DISCUSSION

### Summary of main results

In this systematic review we have found studies that observed no effects of preventive interventions, as well as studies that demonstrated some statistically significant effects, for both alcohol-specific and generic prevention interventions. In this review, the number of studies that evaluated generic interventions was greater compared to that of studies that evaluated alcohol-specific interventions (39 vs. 11). Five of the 11 studies that evaluated alcohol-specific interventions did not find any statistically significant effects,



whereas 6 studies found significant beneficial effects of the intervention programs (McBride 2000; Dielman 1986; Morgenstern 2009; Perry 1988; Vogl 2009; Wilhelmsen 1994). Similarly, 24 of the 39 trials that evaluated generic programs did not find statistically significant effects, whereas the remaining 15 studies reported significantly beneficial effects of the programs with regards to some of the measures of alcohol use.

For both alcohol-specific and generic intervention programs, there was no clearly discernible pattern in characteristics (e.g., sample size, appropriate analysis, attrition rates, subgroups, intervention duration, unit of randomisation, or baseline use) that would distinguish trials with positive results from those with negative results. Most commonly observed positive effects across programs were for drunkenness and binge drinking.

Duration of intervention impact tended to be longer for generic vs. alcohol-specific or other programs. In general, studies that evaluated generic programs reported longer-term follow-up evaluations than those that evaluated alcohol-specific interventions, providing evidence of persistent effects over time (i.e., duration of impact). Positive effects of programs observed in cluster-randomised trials, which did not account for clustering effects in the analyses, may have been spurious (Scaggs 1985; Schinke 2000; Griffin 2009; Perry 1988; Wilhelmsen 1994).

Amongst the generic prevention programs, those based on psychosocial or developmental approaches (e.g., life skills through the LST program in the United States; social skills and norms through the Unplugged program in Europe; development of behaviour norms and peer affiliation through the GBG in the United States and in Europe) were more likely to report statistically significant effects over several years (up to 12 years with the GBG) when compared to standard school curriculum or other types of interventions, with effect sizes that are often small but potentially important based on economic models (Caulkins 2004; NICE 2010). Generic programs offer the additional advantage of potentially impacting on a broader set of problem behaviours, for example cannabis, tobacco, harder drugs, antisocial behaviour. Overall, we conclude that the evidence supports certain generic prevention programs over alcohol-specific prevention programs.

Some trials observed subgroup effects (e.g., by gender, baseline alcohol use, or levels of disruptive behaviour) or after adjusting or stratifying the main effects, some studies found significant effects only in certain subgroups. Few studies reported planned subgroup analyses, and on the one hand, because examining smaller samples reduces statistical power, potentially important effects in some subgroups may not have reached statistical significance (e.g. weaker effects in females). Such analyses should be regarded as hypothesis generating. On the other hand, subgroup analyses through multiple testing may result in type I error and spuriously significant associations (e.g., positive effects shown in males but not in females). It is also possible that some studies that looked only at main effects, without adjusting for potential confounders or effect modifiers, may have concealed possible subgroup effects (e.g. stronger effects

in males). Characteristics such as gender and baseline alcohol use are potential effect moderators, so by not accounting for them in the analysis, subgroup effects may be missed.

One study reported unexpected effects, in that the intervention seemed to increase the risk of alcohol misuse (i.e. a statistically significant increase in drinking in the intervention group). However, before any attribution of iatrogenic effects of particular interventions can be made, it is important to rule out the possibility that occasional unexpected results did not arise by chance, differential attrition or confounding.

One interpretation of the overall picture - some studies showing some effect and other studies showing no effect - is that this is a reflection of the reality that school-based alcohol prevention programs do not work, i.e., they are ineffective, and that there is simply a variation of individual study (and sub-group analysis) effect size estimates around an actual zero effect, with some achieving statistical significance by chance (Ioannidis 2005). However, we regard this as unlikely given the proportion and sample size of studies that found statistically significant effects coupled with the likelihood that many studies were underpowered to find small effects. A more likely interpretation of the overall picture is that some school-based psychosocial and developmental prevention interventions are effective in particular settings for reducing alcohol misuse amongst young people. However, we have also found in this systematic review that some social or life-skills based prevention interventions are not effective (e.g. Sloboda 2009). It is not clear why some prevention interventions seem to work in some studies but not in others, so further investigation of the specific content of prevention programs, and the context of their delivery, is warranted, so that clear recommendations regarding the transfer of particular prevention interventions to new settings can be made.

## Quality of the evidence

In previous systematic reviews of alcohol misuse prevention for young people (Foxcroft 1997; Foxcroft 2002) we have pointed to methodological limitations in included studies. Over this period, consensus statements have been published providing guidance on reporting of randomised controlled trials generally (CONSORT 2010) or more specifically for prevention trials (Flay 2005). Cochrane reviews have also become better at systematically identifying methodological limitations through the risk of bias analysis. Our assessment is that the methodological quality of trials of alcohol misuse prevention for young people has improved over time, between 1997, 2002 and 2010. However, despite these improvements, there remain important methodological limitations and reporting problems. The failure of some studies to account for clustering effects in design or analysis is a significant limitation in studies of universal school-based alcohol misuse prevention programs, given the need for large studies that have sufficient statistical power to detect small effect sizes.

High attrition rates remain a challenge, with few studies in this review achieving the standard 80% follow-up rates expected of good trials beyond the first follow-up. High attrition rates may limit the study power to detect pre-specified between-group differences and/or extent of applicability of study results (Fewtrell 2008). More importantly, in case of differential attrition, study results may be seriously biased due to selection bias/confounding. Alongside this, few studies reported using more advanced techniques for missing data imputation and analysis within an intention to treat approach (Brown 2008). Moreover, in this review over 40% of the studies included were deemed to be susceptible to other bias in the form of confounding or contamination. . Reporting of salient features of RCTs (CONSORT 2010) is also poor in some aspects, notably allocation concealment, randomisation technique, and blinding specifically of outcome assessors. Moreover, reporting of results varied markedly across studies, with many studies not reporting important statistical information such as sample sizes in each group, or standard deviations or standard errors associated with mean scores. This inconsistent reporting meant that calculation of effect sizes for a funnel plot to assess any potential publication bias was not possible.

### **Content and Context: further considerations**

The content or ingredients of effective prevention programs, as distinct from the content of ineffective prevention programs, needs to be more clearly understood. For example, social or life skills curricula may, or may not, vary importantly across different programs. Unfortunately, standard scientific reporting of prevention trials does not include sufficient information about the content detail of prevention interventions to make an analysis of effective ingredients straightforward. Importantly, this lack of information is also one factor that limits a pooling of results across different studies in a meta-analysis, because it is not clear whether interventions have similar or different components. Rather, program manuals and unpublished reports have to be scrutinised, coded for different ingredients, and then analysed, which is a labour intensive and costly approach. Some early review work that has taken this approach has analysed the contribution of different ingredients of prevention programs and these studies have highlighted a number of methodological and analytical challenges (Hansen 2007; Abraham 2008).

Alternatively, it may be that program content is less important than context in discriminating effective from ineffective interventions. It may be that characteristics of program delivery, including program setting, key personnel, or target age are important moderators of program effects. For example, a prevention program which has been shown to be effective in a low prevalence adoles-

cent alcohol misuse setting or country may be ineffective where adolescent drinking is the norm and social and cultural pressures to drink are more powerful.

In order to better understand the importance of content and context for effective prevention, replication studies and more systematic reporting of program content details and delivery contexts are needed. Meta-analysis, via sub-group analysis or using meta-regression techniques, could then be used to illuminate the important aspects of content and context for effective prevention interventions.

## **AUTHORS' CONCLUSIONS**

### **Implications for practice**

Current evidence suggests that certain generic psychosocial and developmental prevention programs can be effective and could be considered as policy and practice options. These include the Life Skills Training Program in the United States, the Unplugged program in Europe, and the Good Behaviour Game in both the United States and Europe. However, given variability in effects between studies and between subgroups within studies, it is recommended that particular attention is paid to program content and delivery context, ideally through conducting further evaluation studies alongside any further implementation in different settings.

### **Implications for research**

As small effects could provide important cost-benefits for prevention programs, it is important to undertake studies with sufficient statistical power to detect small effects. Such small effects may vary in size and importance between subgroups, so further research should also be powered to detect hypothesized subgroup effects. The relevance of content and context of prevention program delivery for program effects is poorly understood, so studies should undertake more rigorous process evaluations alongside outcome evaluations. Reporting of program content and context should be more detailed and systematic to enable comparison of these aspects across studies. Further improvement to study design, analysis and reporting, in line with accepted guidance is required (Flay 2005; CONSORT 2010).

## **ACKNOWLEDGEMENTS**

We are grateful to the UK Cochrane Collaboration and the NIHR for providing support to update the 2002 systematic review.

## REFERENCES

### References to studies included in this review

**Allison 1990** *{published data only}*

Allison KR, Silverman G, Dignam C. Effects on students of teacher training in use of a drug education curriculum. *Journal of Drug Education* 1990;**20**:31–46.

**Beaulieu 1998** *{published data only}*

Beaulieu MA, Jason LA. A drug abuse prevention program aimed at teaching seventh grade students problem solving strategies. *Children and Youth Services Review* 1988;**10**: 131–49.

**Bond 2004** *{published data only}*

Bond L, Patton G, Glover S, Carlin JB, Butler H, Thomas L, Bowes G. The Gatehouse Project: can a multilevel school intervention affect emotional wellbeing and health risk behaviours?. *Journal of Epidemiology & Community Health* 2004;**58**(12):997–1003.

**Botvin 1984** *{published data only}*

Botvin GJ, Baker E, Renick NL, Filazzola AD, Botvin EM. A cognitive behavioral approach to substance abuse prevention. *Addictive Behaviors* 1984;**9**:137–47.

**Botvin 1995** *{published data only}*

Botvin GJ, Baker E, Dusenbury L, Botvin EM, Diaz T. Long term follow-up results of a randomised drug abuse prevention trial. *Journal of the American Medical Association* 1995;**273**:1106–12.

**Botvin 2001** *{published data only}*

Botvin GJ, Griffin KW, Diaz T, Ifill-Williams M. Preventing binge drinking during early adolescence: one- and two-year follow-up of a school-based preventive intervention. *Psychology of Addictive Behaviors* 2001;**15**(4):360–5.

**Botvin 2003** *{published data only}*

Botvin GJ, Griffin KW, Paul E, Macaulay AP. Preventing tobacco and alcohol use among elementary school students through life skills training. *Journal of Child and Adolescent Substance Abuse* 2003;**12**(4):1–17.

**Brewer 1991** *{unpublished data only}*

Brewer LC. Social skills training as a deterrent to entry level drug experimentation among 15-year-old adolescents. Unpublished Ph.D. Dissertation, University of Pennsylvania 1991.

**Caplan 1992** *{published data only}*

Caplan M, Weissberg RP, Grober JS, Sivo PJ, Grady K, Jacoby C. Social competence promotion with inner city and suburban young adolescents: Effects on social adjustment and alcohol use. *Journal of Consulting and Clinical Psychology* 1992;**60**:56–63.

**Clayton 1991** *{published data only}*

Clayton RR, Cattarello A, Johnstone BM. The Effectiveness of Drug Abuse Resistance Education (Project DARE): 5-Year Follow-Up Results. *Preventive Medicine* 1996;**25**: 307–18.

\* Clayton RR, Cattarello A, Walden KP. Sensation seeking as a potential mediating variable for school based prevention

intervention: A two year follow-up of DARE. *Health Communication* 1991;**3**(4):229–39.

Lynam DR, Milich R, Zimmerman R, Novak SP, Logan TK, Martin C, et al. Project DARE: No Effects at 10-Year Follow-Up. *Journal of Consulting and Clinical Psychology* 1999;**67**(4):590–3.

**Cook 1984** *{published data only}*

Cook R, Lawrence H, Morse C, Roehl J. An evaluation of the alternatives approach to drug abuse prevention. *International Journal of the Addictions* 1984;**19**(7):767–87.

**Dielman 1986** *{published data only}*

\* Dielman TE, Shope JT, Butchart AT, Campanelli PC. Prevention of adolescent alcohol misuse: An elementary school program. *Journal of Pediatric Psychology* 1986;**11**(2): 259–82.

Dielman TE, Shope JT, Leech SL, Butchart AT. Differential Effectiveness of an Elementary School-based Alcohol Misuse Prevention Programme. *Journal of School Health* 1989;**59**: 255–63.

Wynn SR, Schulenberg J, Maggs JL, Zucker RA. Preventing Alcohol Misuse: The Impact of Refusal Skills and Norms. *Psychology of Addictive Behaviours* 2000;**14**(1):36–47.

**Durrant 1986** *{unpublished data only}*

Durrant LH. A multi component approach to prevention of adolescent substance abuse. Unpublished Ph.D. Dissertation, University of Utah 1986.

**Duryea 1984a** *{published data only}*

Duryea E, Mohr P, Newman IM, Martin GL, Egwaoje E. Six month follow up results of a preventive alcohol education intervention. *Journal of Drug Education* 1984;**14** (2):97–104.

Duryea E, Okwumabua J. Effects of a preventive alcohol education programme after three years. *Journal of Drug Education* 1988;**18**(1):23–31.

\* Duryea E, Okwumabua J, Rouse J. Preliminary 6 month follow-up results of a preventive alcohol intervention. *Australian Council for Health, Physical Education and recreation national Journal* 1984;**104**:15–8.

**D'Amico 2002** *{published data only}*

D'Amico EJ, Fromme K. Brief prevention for adolescent risk-taking behavior. *Addiction* 2002;**97**(5):563–74.

**Eisen 2002** *{published data only}*

\* Eisen M, Zellman GL, Massett HA, Murray DM. Evaluating the Lions-Quest "Skills for Adolescence" drug education program: first-year behavior outcomes. *Addictive Behaviors* 2002;**27**(4):619–32.

Eisen M, Zellman GL, Murray DM. Evaluating the Lions-Quest "Skills for Adolescence" drug education program: second-year behavior outcomes. *Addictive Behaviors* 2003; **28**:883–97.

**Ellickson 1990** *{published data only}*

\* Ellickson PL, Bell RM. Drug prevention in junior high: A multi site longitudinal test. *Science* 1990;**247**:1299–305.  
Ellickson PL, Bell RM, Harrison ER. Changing adolescent propensities to use drugs: Results from Project ALERT. *Health Education Quarterly* 1993;**20**:227–42.  
Ellickson PL, Bell RM, McGuigan K. Preventing adolescent drug use: long-term results of a junior high programme. *American Journal of Public Health* 1993;**83**(6):856–61.

**Ellickson 2003** *{published data only}*

Ellickson PL, McCaffrey DF, Ghosh-Dastidar B, Longshore DL. New Inroads in Preventing Adolescent Drug Use: Results from a Large-Scale Trial of Project ALERT in Middle Schools. *American Journal of Public Health* 2003;**93** (11):1830–6.

**Faggiano 2007** *{published and unpublished data}*

Caria MP, Faggiano F, Bellocco R, Galanti MR, and the EU-Dap Study group. Effects of a School-based Prevention Program on European Adolescents Patterns of Alcohol Use. *Journal of Adolescent Health* 2010; Vol. (in press):DOI: 10.1016/j.jadohealth.2010.06.003 .

Faggiano F, Galanti MR, Bohrn K, Burkhart G, Vigna-Taglianti F, Cuomo L, et al. The effectiveness of a school-based substance abuse prevention program: EU-Dap cluster randomised controlled trial. *Preventive Medicine* 2008;**47**: 537–43.

\* Faggiano F, Richardson C, Bohrn K, Galanti MR, EU-Dap Study Group. A cluster randomised controlled trial of school-based prevention of tobacco, alcohol and drug use: The EU-Dap design and study population. *Preventive Medicine* 2007;**44**(2):170–3.

Faggiano F, Vigna-Taglianti F, Burkhart G, Bohrn K, Cuomo L, Gregori D, Panella M, et al. The effectiveness of a school-based substance abuse prevention program: 18-month follow-up of the Eu-Dap cluster randomised controlled trial. *Drug and Alcohol Dependence* 2010;**108**: 56–64.

Vigna-Taglianti F, Vadrucchi S, Faggiano F, Burkhart G, Siliquini R, Galanti MR. Is universal prevention against youths' substance misuse really universal? Gender-specific effects in the EU-Dap school-based prevention trial. *Journal of Epidemiology and Community Health* 2009;**63**:722–8.

**Furr-Holden 2004** *{published data only}*

Furr-Holden CD, Ialongo NS, Anthony JC, Petras H, Kellam SG. Developmentally inspired drug prevention: Middle school outcomes in a school-based randomised prevention trial. *Drug and Alcohol Dependence* 2004;**73**: 149–58.

**Goldberg 2000** *{published data only}*

Goldberg L, MacKinnon DP, Elliot DL, Moe EL, Clarke G, Cheong JW. The adolescents training and learning to avoid steroids program. *Archives of Pediatric Adolescence Medicine* 2000;**154**:332–8.

**Goldberg 2007** *{published data only}*

Goldberg L, Elliot DL, MacKinnon DP, Moe EL, Kuehl KS, Yoon M, et al. Outcomes of a prospective trial of student-athlete drug testing: The Student Athlete Testing

Using Random Notification (SATURN) Study. *Journal of Adolescent Health* 2007;**41**(5):421–9.

**Goodstadt 1983** *{published data only}*

Goodstadt MS, Sheppard MA. Three approaches to alcohol education. *Journal of Studies on Alcohol* 1983;**44**(2):362–80.

**Griffin 2009** *{published data only}*

Griffin JP, Holliday RC, Frazier E, Braithwaite RL. The BRAVE (Building Resiliency and Vocational Excellence) Program: evaluation findings for a career-oriented substance abuse and violence preventive intervention. *Journal of Health Care for the Poor & Underserved* 2009;**20**(3): 798–816.

**Hansen 1988** *{published data only}*

Hansen WB, Johnson CA, Flay BR, Graham JW, Sobel J. Affective and social influences approaches to the prevention of multiple substance abuse among seventh grade students: Results from project SMART. *Preventive Medicine* 1988;**17**: 135–54.

**Hansen 1991** *{published data only}*

Hansen WB, Graham JW. Preventing alcohol, marijuana, and cigarette use among adolescents: Peer pressure resistance training versus establishing conservative norms. *Preventive Medicine* 1991;**20**:414–30.

**Hecht 2003** *{published data only}*

\* Hecht ML, Marsiglia FF, Elek E, Wagstaff DA, Kulis S, Dustman P, et al. Culturally grounded substance use prevention: an evaluation of the 'keeping' it R.E.A.L. curriculum. *Prevention Science* 2003;**4**(4):233–48.

Kulis S, Marsiglia FF, Elek E, Dustman P, Wagstaff DA, Hecht ML. Mexican/Mexican American adolescents and 'keeping' it REAL: An evidence-based substance use prevention program. *Children & Schools* 2005;**27**(3): 133–45.

Kulis S, Nieri T, Yabiku S, Stromwall LK, Marsiglia FF. Promoting reduced and discontinued substance use among adolescent substance users: effectiveness of a universal prevention program. *Prevention Science* 2007;**8**(1):35–49.  
Kulis S, Yabiku S, Marsiglia FF, Nieri T, Crossman A. Differences by gender, ethnicity, and acculturation in the efficacy of the 'keeping' it REAL model prevention program. *Journal of Drug Education* 2007;**37**(2):123–44.

**Johnson 2009** *{published data only}*

Johnson KW, Shamblen SR, Ogilvie KA, Collins D, Saylor B. Preventing youths' use of inhalants and other harmful legal products in frontier Alaskan communities: a randomised trial. *Prevention Science* 2009;**10**(4):298–12.

**Kellam 2008** *{published data only}*

Kellam SG, Brown CH, Poduska JM, Ialongo NS, Wang W, Toyinbo P, et al. Effects of a universal classroom behavior management program in first and second grades on young adult behavioral, psychiatric, and social outcomes. *Drug and Alcohol Dependence* 2008;**95S**:S5–S28.

**Koning 2009** *{published data only}*

Koning IM, Vollebergh W, Smit F, Verdurmen J, van den Eijnden R, ter Bogt T, et al. Preventing heavy alcohol use in adolescents (PAS): Cluster randomised trial of a parent and

- student intervention offered separately and simultaneously. *Addiction* 2009;**104**(10):1669–78.
- McBride 2000** *{published data only}*  
 McBride N, Farrington F, Midford R, Meuleners L, Phillips M. Early unsupervised drinking - reducing the risks. The School health and Alcohol Harm Reduction Project. *Drug and Alcohol Review* 2003;**22**:263–76.  
 McBride N, Farrington F, Midford R, Meuleners L, Phillips M. Harm minimization in school drug education: Final results of the School Health and Alcohol Harm Reduction Project (SHAHRP). *Addiction* 2004;**99**(3):278–91.  
 \* McBride N, Midford R, Farrington F, Phillips M. Early results from a school alcohol harm minimization study: the School Health and Alcohol Harm Reduction Project. *Addiction* 2000;**95**(7):1021–42.
- Morgenstern 2009** *{published data only}*  
 Morgenstern M, Wiborg G, Isensee B, Hanewinkel R. School-based alcohol education: results of a cluster-randomised controlled trial. *Addiction* 2009;**104**(3):402–12.
- Moskowitz 1984** *{published data only}*  
 Moskowitz JM, Malvin JH, Schaeffer GA, Schaps E. An experimental evaluation of a drug education course. *Journal of Drug Education* 1984;**14**(1):9–22.
- Newman 1992** *{published data only}*  
 Newman IM, Anderson CS, Farrell KA. Role rehearsal and efficacy: Two 15 month evaluations of a ninth grade alcohol education programme. *Journal of Drug Education* 1992;**22**(1):55–67.
- Newton 2009a** *{published data only}*  
 \* Newton NC, Andrews G, Teesson M, Vogl LE. Delivering prevention for alcohol and cannabis using the Internet: a cluster randomised controlled trial. *Preventive Medicine* 2009;**48**(6):579–84.  
 Newton NC, Teesson M, Vogl LE, Andrews G. Internet-based prevention for alcohol and cannabis use: final results of the Climate Schools course. *Addiction* 2009;**105**:749–59.
- Perry 1988** *{published data only}*  
 \* Perry CL, Grant M. Comparing peer-led to teacher-led youth alcohol education in four countries (Australia, Chile, Norway and Swaziland). *Alcohol Health and Research World* 1988;**12**:322–6.  
 Perry CL, Grant M, Ernberg G, Florenzano RU, Langdon MC, Myeni AD, et al. WHO collaborative study of alcohol education and young people: Outcomes of a four country pilot study. *International Journal of the Addictions* 1989;**24**(12):1145–71.
- Perry 2003** *{published data only}*  
 Perry CL, Komro KA, Mortenson-Veblen S, Bosma LM, Farbaksh K, Munson KA, et al. A randomised controlled trial of the middle and junior high school D.A.R.E. and D.A.R.E. Plus programs. *Archives of Pediatrics & Adolescent Medicine* 2003;**157**:178–184.
- Reddy 2002** *{published data only}*  
 Reddy KS, Arora M, Perry CL, Nair B, Kohli A, Lytle LA, et al. Tobacco and alcohol use outcomes of a school-based intervention in New Dehli. *American Journal of Health Behavior* 2002;**26**(3):173–81.
- Ringwalt 1991** *{published data only}*  
 Ringwalt C, Ennett S, Holt KD. An outcome evaluation of project DARE. *Health Education Research* 1991;**6**(3):327–37.
- Ringwalt 2009** *{published data only}*  
 \* Ringwalt CL, Clark HK, Hanley S, Shamblen SR, Flewelling RL. Project ALERT: a cluster randomised trial. *Archives of Pediatrics and Adolescent Medicine* 2009;**163**(7):625–32.  
 Ringwalt CL, Clark HK, Hanley S, Shamblen SR, Flewelling RL. The effects of Project ALERT one year past curriculum completion. *Prevention Science* 2010;**11**:172–84.
- Scaggs 1985** *{unpublished data only}*  
 Scaggs LS. A substance abuse awareness prevention program: knowledge, attitudes and behaviours. Unpublished Ph.D. Dissertation, Ohio State University 1985.
- Schinke 2000** *{published data only}*  
 Schinke SP, Tepavac L, Cole KC. Preventing substance use among Native American youth: three-year results. *Addictive Behaviours* 2000;**25**(3):387–97.
- Sheehan 1996** *{published data only}*  
 Sheehan M, Schonfeld C, Ballard R, Schofield F, Najman J, Siskind V. Three Year Outcome Evaluation of a Theory Based Drink Driving Education Program. *Journal of Drug Education* 1996;**26**(3):295–312.
- Sloboda 2009** *{published data only}*  
 Sloboda Z, Stephens RC, Stephens PC, Grey SF, Teasdale B, Hawthorne RD, et al. The Adolescent Substance Abuse Prevention Study: A randomised field trial of a universal substance abuse prevention program. *Drug and Alcohol Dependence* 2009;**102**:1–10.
- Spoth 2002** *{published data only}*  
 Spoth RL, Randall GK, Trudeau L, Shin C, Redmond C. Substance use outcomes 5 1/2 years past baseline for partnership-based, family-school preventive interventions. *Drug and Alcohol Dependence* 2008;**96**:57–68.  
 Spoth RL, Randall K, Shin C, Redmond C. Randomized study of combined universal family and school preventive interventions: Patterns of long-term effects on initiation, regular use, and weekly drunkenness. *Psychology of Addictive Behaviors* 2005;**19**(4):372–81.  
 \* Spoth RL, Redmond C, Trudeau L, Shin C. Longitudinal substance initiation outcomes for a universal preventive intervention combining family and school programs. *Psychology of Addictive Behaviors* 2002;**16**(2):129–34.
- St. Pierre 2005** *{published data only}*  
 St. Pierre TL, Osgood DW, Mincemoyer CC, Kaltreider DL, Kauh TJ. Results of an independent evaluation of Project ALERT delivered in schools by Cooperative Extension. *Prevention Science* 2005;**6**(4):305–17.

**Sun 2008** {published data only}

Sun P, Sussman S, Dent CW, Rohrbach A. One-year follow-up evaluation of Project Towards No Drug Abuse (TND-4). *Preventive Medicine* 2008;**47**(4):438–42.

**van Lier 2009** {published data only}

van Lier PAC, Huizink A, Crijnen A. Impact of a preventive intervention targeting childhood disruptive behavior problems on tobacco and alcohol initiation from age 10 to 13 years. *Drug and Alcohol Dependence* 2009;**100**(3):228–33.

**Vogl 2009** {published data only}

Vogl L, Teesson M, Andrews G, Bird K, Steadman B, Dillon P. A computerized harm minimization prevention program for alcohol misuse and related harms: Randomized controlled trial. *Addiction* 2009;**104**(4):564–75.

**Werch 2008** {published data only}

Werch CE, Bian H, Moore MJ, Ames SC, DiClemente CC, Thombs D, et al. Brief multiple behavior health interventions for older adolescents. *American Journal of Health Promotion* 2008;**23**(2):92–6.

**Werch 2010** {published data only}

Werch CE, Bian H, DiClemente CC, Moore MJ, Thombs D, Ames SC, et al. A brief image-based prevention intervention for adolescents. *Psychology of Addictive Behaviors* 2010;**24**(1):170–5.

**Wilhelmsen 1994** {published data only}

Wilhelmsen BU, Laberg JC, Klepp KI. Evaluation of two student and teacher involved alcohol prevention programmes. *Addiction* 1994;**89**(9):1157–65.

**Williams 1968** {published data only}

Williams AF, DiCicco LM, Unterberger H. Philosophy and evaluation of an alcohol education programme. *Quarterly Journal of Studies on Alcohol* 1968;**29**:685–702.

## References to studies excluded from this review

**Ackermann 2008** {published data only}

Ackermann ME. The importance of gender and readiness to change in the prediction of drinking and negative consequences of first-year student drinkers. Ackermann, Margot E: Old Dominion U, US. 2008.

**Amaro 2009** {published data only}

Amaro H, Ahl M, Matsumoto A, Prado G, Mulé C, Kemmemer A, et al. Trial of the university assistance program for alcohol use among mandated students. *Journal of Studies on Alcohol & Drugs Supplement* 2009;**16**:45–56.

**Anderson 2004** {published data only}

Anderson KL. Child and adolescent predictors of youth alcohol use to intoxication. The Johns Hopkins University, US 2004.

**Bailey 2004** {published data only}

Bailey K A, Baker AL, Webster RA, Lewin TJ. Pilot randomised controlled trial of a brief alcohol intervention group for adolescents. *Drug and Alcohol Review* 2004;**23**(2):157–66.

**Bauman 2002** {published data only}

Bauman KE, Ennett ST, Foshee VA, Pemberton M, King TS, Koch GG. Influence of a family program on adolescent smoking and drinking prevalence. *Prevention Science* 2002;**3**(1):35–42.

**Bell 2005** {published data only}

Bell ML, Baker TK, Falb T, Roberts-Gray C. Protecting You/Protecting Me: Evaluation of a student-led alcohol prevention and traffic safety program for elementary students. *Journal of Alcohol and Drug Education* 2005;**49**(1):33–53.

**Benner 2008** {published data only}

Benner TA. *Aban Aya Youth Project: Preventing high-risk behaviours among African American youth in grades 5-8. Model programs for adolescent sexual health: Evidence-based HIV, STI, and pregnancy prevention interventions*. New York: J. J. Card and T. A. Benner. New York, NY, Springer Publishing Co, 2008:157–164.

**Bersamin 2007** {published data only}

Bersamin M, Paschall MJ, Fearnow-Kenney M, Wyrick D. Effectiveness of a Web-based alcohol-misuse and harm-prevention course among high- and low-risk students. *Journal of American College Health* 2007;**55**(4):247–54.

**Boekeloo 2004** {published data only}

Boekeloo BO, Jerry J, Lee-Ougo WI, Worrell KD, Hamburger EK, Russek-Cohen E, et al. Randomized trial of brief office-based interventions to reduce adolescent alcohol use. *Archives of Pediatrics and Adolescent Medicine* 2004;**158**(7):635–42.

**Botvin 2006** {published data only}

Botvin GJ, Griffin KW, Nichols TD. Preventing youth violence and delinquency through a universal school-based prevention approach. *Prevention Science* 2006;**7**(4):403–8.

**Brody 2004** {published data only}

Brody GH, Murry VM, Gerrard M, Gibbons FX, Molgaard V, McNair L, et al. The Strong African American Families Program: translating research into prevention programming. *Child Development* 2004;**75**(3):900–17.

**Brody 2005** {published data only}

Brody GH, Murry VM, McNair L, Chen YF, Gibbons FX, Gerrard M, et al. Linking Changes in Parenting to Parent-Child Relationship Quality and Youth Self-Control: The Strong African American Families Program. *Journal of Research on Adolescence* 2005;**15**(1):47–69.

**Brody 2006** {published data only}

Brody GH, Murry VM, Kogan SM, Gerrard M, Gibbons FX, Molgaard V, et al. The Strong African American Families Program: A cluster-randomised prevention trial of long-term effects and a mediational model. *Journal of Consulting and Clinical Psychology* 2006;**74**(2):356–66.

**Brody 2008** {published data only}

Brody GH, Kogan SM, Chen YF, McBride, Murry V. Long-term effects of the strong African American families program on youths' conduct problems. *Journal of Adolescent Health* 2008;**43**(5):474–81.

- Brown 2005** *{published data only}*  
Brown EC, Catalano RF, Fleming CB, Haggerty KP, Abbott RD. Adolescent substance use outcomes in the Raising Healthy Children project: a two-part latent growth curve analysis. *Journal of Consulting and Clinical Psychology* 2005;**73**(4):699–710.
- Brown 2007** *{published data only}*  
Brown CH, Guo J, Singer LT, Downes K, Brinales JM. Examining the effects of school-based drug prevention programs on drug use in rural settings: methodology and initial findings. *Journal of Rural Health* 2007;**23** Suppl: 29–36.
- Bryan 2009** *{published data only}*  
Bryan AD, Schmiege SJ, Broaddus MR. HIV risk reduction among detained adolescents: a randomised, controlled trial. *Pediatrics* 2009;**124**(6):e1180–1188.
- Castellanos 2006** *{published data only}*  
Castellanos N, Conrod P. Brief interventions targeting personality risk factors for adolescent substance misuse reduce depression, panic and risk-taking behaviours. *Journal of Mental Health* 2006;**15**(6):645–58.
- Caudill 2007** *{published data only}*  
Caudill BD, Luckey B, Crosse SB, Blane HT, Ginexi EM, Campbell B. Alcohol risk-reduction skills training in a national fraternity: a randomised intervention trial with longitudinal intent-to-treat analysis. *Journal of Studies on Alcohol & Drugs* 2007;**68**(3):399–409.
- Connell 2007** *{published data only}*  
Connell AM, Dishion TJ, Yasui M, Kavanagh K. An adaptive approach to family intervention: linking engagement in family-centred intervention to reductions in adolescent problem behavior. *Journal of Consulting and Clinical Psychology* 2007;**75**(4):568–79.
- Conrod 2006** *{published data only}*  
Conrod PJ, Stewart SH, Comeau N, Maclean AM. Efficacy of Cognitive-Behavioral Interventions Targeting Personality Risk Factors for Youth Alcohol Misuse. *Journal of Clinical Child and Adolescent Psychology* 2006;**35**(4):550–63.
- Conrod 2008** *{published data only}*  
Conrod PJ, Castellanos N, Mackie C. Personality-targeted interventions delay the growth of adolescent drinking and binge drinking. *Journal of Child Psychology and Psychiatry and Allied Disciplines* 2008;**49**(2):181–90.
- Croom 2009** *{published data only}*  
Croom KD, Lewis D, Marchell T, Lesser ML, Reyna VF, Kubicki-Bedford L, et al. Impact of an online alcohol education course on behavior and harm for incoming first-year college students: short-term evaluation of a randomised trial. *Journal of American College Health* 2009;**57**(4): 445–54.
- D'Amico 2008** *{published data only}*  
D'Amico EJ, Miles JN, Stern SA, Meredith LS. Brief motivational interviewing for teens at risk of substance use consequences: a randomised pilot study in a primary care clinic. *Journal of Substance Abuse Treatment* 2008;**35**(1): 53–61.
- DeGarmo 2009** *{published data only}*  
DeGarmo DS, Eddy JM, Reid JB, Fetrow RA. Evaluating mediators of the impact of the Linking the Interests of Families and Teachers (LIFT) multimodal preventive intervention on substance use initiation and growth across adolescence. *Prevention Science* 2009;**10**(3):208–20.
- Dembo 2002** *{published data only}*  
Dembo R, Wothke W, Livingston S, Schmeidler J. The impact of a family empowerment intervention on juvenile offender heavy drinking: a latent growth model analysis. *Substance Use and Misuse* 2002;**37**(11):1359–90.
- Donohue 2004** *{published data only}*  
Donohue BD, Allen N, et al. A Controlled Evaluation of Two Prevention Programs in Reducing Alcohol Use Among College Students at Low and High Risk for Alcohol Related Problems. *Journal of Alcohol and Drug Education* 2004;**48** (1):13–33.
- Eddy 2003** *{published data only}*  
Eddy JM, Reid JB, Stoolmiller M, Fetrow RA. Outcomes During Middle School for an Elementary School-Based Preventive Intervention for Conduct Problems: Follow-Up Results From a Randomized Trial. *Behavior Therapy* 2003; **34**(4):535–52.
- Elder 2002** *{published data only}*  
Elder JP, Litrownik AJ, Slymen DJ, Campbell NR, Parra-Medina D, Choe S, et al. Tobacco and alcohol use-prevention program for Hispanic migrant adolescents. *American Journal of Preventive Medicine* 2002;**23**(4): 269–75.
- Elliot 2004** *{published data only}*  
Elliot DL, Goldberg L, Moe EL, Defrancesco CA, Durham MB, Hix-Small H. Preventing substance use and disordered eating: Initial outcomes of the ATHENA (Athletes Targeting Healthy Exercise and Nutrition Alternatives) program. *Archives of Pediatrics and Adolescent Medicine* 2004;**158**(11):1043–9.
- Friedman 2002** *{published data only}*  
Friedman AS, Terras A, Glassman K. Multimodal substance use intervention program for male delinquents. *Journal of Child and Adolescent Substance Abuse* 2002;**11**(4):43–65.
- Fromme 2004** *{published data only}*  
Fromme K, Corbin W. Prevention of heavy drinking and associated negative consequences among mandated and voluntary college students. *Journal of Consulting and Clinical Psychology* 2004;**72**(6):1038–49.
- Griffin 2003** *{published data only}*  
Griffin KW, Botvin GJ, Nichols TR, Doyle MM. Effectiveness of a universal drug abuse prevention approach for youth at high risk for substance use initiation. *Preventive Medicine* 2003;**36**(1):1–7.
- Griffin 2004** *{published data only}*  
Griffin KW, Botvin GJ, Nichols TR. Long-term follow-up effects of a school-based drug abuse prevention program on adolescent risky driving. *Prevention Science* 2004;**5**(3): 207–12.

**Griffin 2006** {published data only}

Griffin KW, Botvin GJ, Nichols TR. Effects of a school-based drug abuse prevention program for adolescents on HIV risk behavior in young adulthood. *Prevention Science* 2006;7(1):103–12.

**Haggerty 2006** {published data only}

Haggerty KP, Fleming CB, Catalano RF, Harachi TW, Abbott RD. Raising healthy children: examining the impact of promoting healthy driving behavior within a social development intervention. *Prevention Science* 2006;7(3):257–67.

**Haggerty 2007** {published data only}

Haggerty KP, Skinner ML, MacKenzie EP, Catalano RF. A randomised trial of Parents Who Care: effects on key outcomes at 24-month follow-up. *Prevention Science* 2007;8(4):249–60.

**Haggerty 2008** {published data only}

Haggerty KP, Skinner M, Fleming CB, Gaaney RR, Catalano RF. Long-term effects of the Focus on Families project on substance use disorders among children of parents in methadone treatment. *Addiction* 2008;103(12):2008–16.

**Hembroff 2007** {published data only}

Hembroff L, Atkin C, Martell D, McCue C, Greenamyre JT. Evaluation results of a 21st birthday card program targeting high-risk drinking. *Journal of American College Health* 2007;56(3):325–32.

**Jemmott 2005** {published data only}

Jemmott JB 3rd, Jemmott LS, Braverman PK, Fong GT. HIV/STD risk reduction interventions for African American and Latino adolescent girls at an adolescent medicine clinic: A randomized controlled trial. *Archives of Pediatrics and Adolescent Medicine* 2005;159(5):440–9.

**Jones 2005** {published data only}

Jones DJ, Olson AL, Forehand R, Gaffney CA, Zens MS, Bau JJ. A Family-Focused Randomized Controlled Trial to Prevent Adolescent Alcohol and Tobacco Use: The Moderating Roles of Positive Parenting and Adolescent Gender. *Behavior Therapy* 2005;36(4):347–55.

**Martinez 2005** {published data only}

Martinez CR, Eddy JM. Effects of culturally adapted parent management training on Latino youth behavioral health outcomes. *Journal of Consulting and Clinical Psychology* 2005;73(5):841–51.

**Poduska 2008** {published data only}

Poduska JM, Kellam SG, Wang W, Brown CH, Ialongo NS, Toyinbo P. Impact of the Good Behavior Game, a universal classroom-based behavior intervention, on young adult service use for problems with emotions, behavior, or drugs or alcohol. *Drug and Alcohol Dependence* 2008;95S:S29–S44.

**Simons-Morton 2005** {published data only}

Simons-Morton B, Haynie D, Saylor K, Crump AD, Chen R. Impact analysis and mediation of outcomes: the Going Places program. *Health Education and Behavior* 2005;32(2):227–41.

**Sussman 2002** {published data only}

Sussman S, Dent CW, Stacy AW. Project towards no drug abuse: a review of the findings and future directions. *American Journal of Health Behavior* 2002;26(5):354–65.

**Wagenaar 2005** {published data only}

Wagenaar AC, Toomey TL, Erickson DJ. Preventing youth access to alcohol: outcomes from a multi-community time-series trial. *Addiction* 2005;100(3):335–45.

**Wolchik 2002** {published data only}

Wolchik SA, Sandler IN, Millsap RE, Plummer BA, Greene SM, Anderson ER, et al. Six-year follow-up of preventive interventions for children of divorce: A randomised controlled trial. *Journal of the American Medical Association* 2002;288(15):1874–81.

**Wu 2003** {published data only}

Wu Y, Stanton BF, Galbraith J, Kaljee L, Cottrell L, Li X, et al. Sustaining and broadening intervention impact: a longitudinal randomised trial of 3 adolescent risk reduction approaches. *Pediatrics* 2003;111(1):32–8.

## Additional references

**Abraham 2008**

Abraham C, Michie S. A taxonomy of behavior change techniques used in interventions. *Health Psychology* 2008;27:379–87.

**AMA 2004**

American Medical Association. Harmful consequences of alcohol use on the brains of children, adolescents, and college students. American Medical Association 2004. <http://www.ama-assn.org/ama/pub/category/9416.html> (accessed July 16, 2010).

**Anderson 2006**

Anderson P, Baumberg B. Alcohol in Europe. Institute of Alcohol Studies 2006. <http://ec.europa.eu/health-eu/news/alcoholineurope/en.htm> (accessed July 16, 2010).

**Babor 2010**

Babor TF, Caetano RS, Casswell G, Edwards N, Giesbrecht K, Graham J, et al. *Alcohol: No Ordinary Commodity. Research and Public Policy*. Oxford: Oxford University Press, 2010.

**Brown 2008**

Brown CH, Wang W, Kellam SG, Muthén BO, Petras H, Toyinbo P, et al. Methods for testing theory and evaluating impact in randomised field trials: intent-to-treat analyses for integrating the perspectives of person, place, and time. *Drug and Alcohol Dependence* 2008;95:S74–S104.

**Caulkins 2004**

Caulkins JP, Pacula RL, Paddock S, Chiesa J. What we can-and cannot-expect from school-based drug prevention. *Drug and Alcohol Review* 2004;23:79–87.

**CMO 2009**

Chief Medical Officer. Guidance on the consumption of alcohol by children and young people. Chief Medical Officer 2009. <http://www.dh.gov.uk/en/Publicationsandstatistics/>



- Publications/PublicationsPolicyAndGuidance/DH:110258 (accessed 16 July 2010).
- CONSORT 2010**  
Consolidated Standards of Reporting Trials. Library for health research reporting. <http://www.equator-network.org/resource-centre/library-of-health-research-reporting/> (accessed 16 July 2010).
- Coombes 2008**  
Coombes L, Allen D, Foxcroft D, Guydish J. Motivational interviewing for the prevention of alcohol misuse in young people. *Cochrane Database of Systematic Reviews* 2008, Issue 2. [DOI: 10.1002/14651858.CD007025]
- Dawson 2008**  
Dawson DA, Goldstein RB, Chou SP, Ruan WJ, Grant BF. Age at first drink and the first incidence of adult-onset DSM-IV alcohol use disorders. *Alcoholism, Clinical and Experimental Research* 2008;**32**:2149–60.
- Egger 1997**  
Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629–34.
- EMCDDA 2010**  
European Monitoring Centre for Drugs and Drug Addiction. Prevention responses to drug use in the EU. <http://www.emcdda.europa.eu/themes/prevention/responses-in-eu> (last updated 15 June 2010; accessed 15 July 2010).
- Faggiano 2005**  
Faggiano F, Vigna-Taglianti F, Versino E, Zambon A, Borraccino A, Lemma P. School-based prevention for illicit drugs' use. *Cochrane Database of Systematic Reviews* 2005, Issue 2. [DOI: 10.1002/14651858.CD003020.pub2]
- Fewtrell 2008**  
Fewtrell MS, Kennedy K, Singhal A, Martin RM, Ness A, Hadders-Algra M, et al. How much loss to follow-up is acceptable in long-term randomised trials and prospective studies?. *Archives of Disease in Childhood* 2008;**93**(6): 458–61.
- Flay 2005**  
Flay BR, Biglan A, Boruch RF, Castro FG, Gottfredson D, Kellam S, et al. Standards of evidence: criteria for efficacy, effectiveness and dissemination. *Prevention Science* 2005;**6**: 151–75.
- Foxcroft 1997**  
Foxcroft DR, Lister-Sharp DJ, Lowe G. Alcohol misuse prevention for young people: a systematic review reveals methodological concerns and lack of reliable evidence of effectiveness. *Addiction* 1997;**92**:531–7.
- Foxcroft 2002**  
Foxcroft D, Ireland D, Lowe G, Breen R. Primary prevention for alcohol misuse in young people. *Cochrane Database of Systematic Reviews* 2002, Issue 3. [DOI: 10.1002/14651858.CD003024]
- Hansen 2007**  
Hansen WB, Dusenbury L, Bishop D, Derzon JH. Substance abuse prevention program content: systematizing the classification of what programs target for change. *Health Education Research* 2007;**22**:351–60.
- Higgins 2009**  
Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.0.2 [updated September 2009]*. The Cochrane Collaboration, 2009. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org). 5.0.2. Cochrane Collaboration, 2009.
- Hingson 2003a**  
Hingson R, Heeren T, Winter MR, Wechsler H. Early age of first drunkenness as a factor in college students' unplanned and unprotected sex attributable to drinking. *Pediatrics* 2003;**111**:34–41.
- Hingson 2003b**  
Hingson R, Heeren T, Zackos R, Winter M, Wechsler H. Age of first intoxication, heavy drinking, driving after drinking, and risk of unintentional injury among U.S. college students. *Journal of Studies on Alcohol* 2003;**64**: 23–31.
- Hingson 2006**  
Hingson RW, Heeren T, Winter MR. Age at drinking onset and alcohol dependence: age at onset, duration, and severity. *Archives of Pediatrics Adolescent Medicine* 2006; **160**:739–46.
- Ioannidis 2005**  
Ioannidis JPA. Why Most Published Research Findings Are False. *PLoS Med* 2005;**2**(8):e124 doi:10.1371/journal.pmed.0020124.
- Moreira 2008**  
Moreira MT, Smith LA, Foxcroft D. Social norms interventions to reduce alcohol misuse in University or College students. *Cochrane Database of Systematic Reviews* 2009, Issue 3. [DOI: 10.1002/14651858.CD006748.pub2]
- Mrazek 1994**  
Mrazek PJ, Haggerty RJ (eds). *Reducing risks for mental disorders: Frontiers for preventive intervention research*. Institute of Medicine. Washington DC: National Academy Press, 1994.
- NICE 2010**  
National Institute for Health and Clinical Excellence. PHSE Evidence Review - Secondary Education and FE. <http://www.nice.org.uk/nicemedia/live/11673/49240/49240.pdf> (accessed 15 July 2010).
- Peters 2008**  
Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Contour-enhanced meta-analysis funnel plots help distinguish publication bias from other causes of asymmetry. *Journal of Clinical Epidemiology* 2008;**61**(10):991–6.
- Pitkanen 2005**  
Pitkanen T, Lyyra AL, Pulkkinen L. Age of onset of drinking and the use of alcohol in adulthood: a follow-up study

from age 8-42 for females and males. *Addiction* 2005;**100**: 652-61.

**Rehm 2005**

Rehm J, Room R, van den Brink W, Jacobi F. Alcohol use disorders in EU countries and Norway: An overview of the epidemiology. *European Neuropsychopharmacology* 2005;**15**: 377-88.

**Warner 2003**

Warner LA, White HR. Longitudinal effects of age at onset and first drinking situations on problem drinking. *Substance Use and Misuse* 2003;**38**:1983-2016.

**WHO 2008**

World Health Organization. Management of substance abuse - Alcohol. World Health Organization 2008. [http://www.who.int/substance\\_abuse/facts/alcohol/en/index.html](http://www.who.int/substance_abuse/facts/alcohol/en/index.html) (accessed July 16, 2010).

**Zakrajsek 2006**

Zakrajsek JS, Shope JT. Longitudinal examination of underage drinking and subsequent drinking and risky driving. *Journal of Safety Research* 2006;**37**:443-51.

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Allison 1990

Methods	<p>Design: RCT  FU: 1 yr (post-randomisation)  Attrition: 18%  ITT: no  Unit of randomisation: class  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int-1: 4 (92)  Int-2: 5 (107)  Ctrl: 3 (67)  Total N: 12 (266)  <u>N of Clusters (subjects) participated</u>  Int-1: 4 (92)  Int-2: 5 (107)  Ctrl: 3 (67)  Total N: 12 (266)</p> <p><u>Analysed sample</u>  Int-1: 4 (70)  Int-2: 5 (80)  Ctrl: 3 (59)  Total N: 12 (209)  Age: 10-11 yrs  Sex (male): NR  Ethnicity: NR  Alcohol users at baseline: 25%-33% had drunk wine with parents  Country: Canada</p>
Interventions	<p><u>Intervention-1</u>: Intensive staff training  <u>Intervention-2</u>: less intensive staff training: 1-2 hours only  <u>Intervention description</u>: a set of resource documents to assist teachers implement the Ontario guidelines for alcohol, tobacco and other drug education  <u>Focus/target</u>: closely related to Life Skills curriculum, the aim was to provide specific substance use information  <u>Components</u>: promote awareness, problem-solving, decision-making, refusal skills  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 5 x 3hr sessions  <u>Control</u>: no staff training for curriculum</p>
Outcomes	<p>FU: 1 yr  <u>Ever had a sip of alcohol (pre-post diff.)</u>  Int-1: 0% vs. Int-2: -2% vs. Ctrl: 3%, p=0.95</p>

Allison 1990 (Continued)

	<p>Ever drank wine with parents (pre-post diff.) Int-1: 8% vs. Int-2: 10% vs. Ctrl: 11%, p=0.96</p> <p>Ever drank wine or beer (pre-post diff.) Int-1: -1%; Int-2: 0%; Ctrl: 0% (p=0.68)</p>
--	--

Notes	<p><b>Drug Abuse Prevention Program - An Education Resource (DAPPER)</b> Teacher training had no clear effect on outcome measures. Small sample size. Possibility of contamination between Intervention groups in some schools</p> <p><u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); diff=difference; ITT=intention to treat (analysis); NA=not applicable; mo=month(s)</p>
-------	---

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No intention to treat analysis
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Possible contamination

**Beaulieu 1998**

Methods	<p>Design: RCT FU: immediate post-intervention Attrition: 47% ITT: no Unit of randomisation: class Clustering effect adjusted: no</p>
Participants	<p><u>N of Clusters (subjects) randomised</u> Int: 2 (40) Ctrl: 3 (72)</p>

**Beaulieu 1998** (Continued)

	<p>Total N: 5 (112)  <u>N of Clusters (subjects) participated</u>            Int: 2 (40)            Ctrl: 3 (72)            Total N: 5 (112)</p> <p><u>Analysed sample</u>            Int: 2 (16)            Ctrl: 3 (44)            Total N: 5 (60)            Age: 12-13 yrs            Sex (male): NR            Ethnicity: 100% Black            Alcohol users at baseline: 31%            Alcohol use mean (SD) frequency (Int vs. Ctrl) at baseline: 1.06 (0.24) vs. 1.11 (0.58)            Country: US</p>	
Interventions	<p><u>Intervention description</u>: primary preventive intervention to reduce tobacco, alcohol and marijuana use  <u>Focus/target</u>: strategies addressing the social and personal conditions underlying abusive drug behavior, delivered by program staff with eighth-grade peer helpers  <u>Components</u>: information giving, decision making, problem solving techniques, social competency-building skills, peer helpers and role models  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: one 45 min session per week for eight weeks  <u>Control</u>: standard curriculum</p>	
Outcomes	<p>FU: 2 mo  <u>Current alcohol use - mean (SD) frequency</u>            Int: 1.12 (0.33) vs. Ctrl: 1.07 (0.33), p&gt;0.05</p>	
Notes	<p><b>Drug Abuse Prevention Program</b>            High attrition; Intervention and control groups may not have been comparable; baseline differences in drug knowledge not accounted.  <u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); SD=standard deviation; NA=not applicable; mo=month(s)</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described

**Beaulieu 1998** (Continued)

Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition > 40%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Differences in some baseline factors between study arms

**Bond 2004**

Methods	Design: RCT FU: 3 yrs (post-randomisation) Attrition: 3% (year 1), 8% (year 2), and 10% (year 3) ITT: yes Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: 16 (NR) Ctrl: 16 (NR) Total N: 32 (3623) <u>N of Clusters (subjects) participated</u> Int: 12 (1335) Ctrl: 14 (1343) Total N: 26 (2678) Age: 14 yrs Sex (male): 47% Ethnicity: NR Alcohol users at baseline: 31% Country: Australia
Interventions	<u>Intervention description</u> : a multilevel program, whole school, teaching resources, and school liaison <u>Focus/target</u> : promote emotional and behavioral well-being to reduce rates of substance use <u>Components</u> : Institutional and individual <u>Fidelity</u> : one school did not deliver the intervention for the 1 <sup>st</sup> year <u>Duration/frequency</u> : 20/yr for 10 wks <u>Control</u> : standard curriculum

**Bond 2004** (Continued)

Outcomes	<p>FU: 1, 2, 3, yrs</p> <p><u>Any drinking (Int vs. Ctrl)</u></p> <p>OR=0.93, 95% CI: 0.71, 1.21 (year 1)</p> <p>OR=1.00, 95% CI: 0.78, 1.28 (year 2)</p> <p>OR=0.96, 95% CI: 0.69, 1.33 (year 3)</p> <p><u>Regular drinking (Int vs. Ctrl)</u></p> <p>OR=1.09, 95% CI: 0.77, 1.57 (year 1)</p> <p>OR=1.05, 95% CI: 0.70, 1.57 (year 2)</p> <p>OR=1.13, 95% CI: 0.77, 1.66 (year 3)</p> <p><u>Binge drinking</u></p> <p>OR=0.95, 95% CI: 0.69, 1.32 (year 1)</p> <p>OR=0.99, 95% CI: 0.70, 1.38 (year 2)</p> <p>OR=1.02, 95% CI: 0.71, 1.46 (year 3)</p>	
Notes	<p><b>Gatehouse Project</b></p> <p>Response rate was &gt; 89% and non-differential</p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); OR=odds ratio; 95% CI: ninety-five percent confidence interval; NA=not applicable; mo=month(s)</p>	
<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT and attrition < 10%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	No obvious differences between two study arms

**Botvin 1984**

Methods	Design: RCT FU: 4 months (post-randomisation) Attrition < 10% ITT: no Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 4 (NR) Int-2: 4 (NR) Ctrl: 2 (NR) Total N: 10 (1311) <u>N of Clusters (subjects) participated</u> Int-1: 4 (NR) Int-2: 4 (NR) Ctrl: 2 (NR) Total N: 10 (1311)  <u>Analysed sample</u> Int-1: 4 (NR) Int-2: 4 (NR) Ctrl: 2 (NR) Total N: 10 (1185) Age: 12-13 yrs (7th grade) Sex (male): NR Ethnicity: mostly White Alcohol users at baseline: NR Country: US
Interventions	<u>Intervention-1</u> : multi-component substance abuse prevention curriculum (LST) delivered by older students <u>Intervention-2</u> : a multi-component substance abuse prevention curriculum (LST) delivered by classroom teachers <u>Focus/target</u> : the major social, psychological, cognitive, and attitudinal factors that promote the use of tobacco, alcohol and marijuana <u>Components</u> : personal self-management, general social skills <u>Fidelity</u> : NR <u>Duration/frequency</u> : 20 sessions over 3-4 months Control: Standard curriculum
Outcomes	FU: 4 months Frequency of drunkenness (ANCOVA): $F(2,929) = 4.54, p < 0.01$ Consumption per occasion (ANCOVA): $F(2,876) = 5.10, p < 0.006$ No specific details given but authors report that alcohol consumption was less in peer-led group. No differences were found in levels of drunkenness between any groups.



**Botvin 1984** (Continued)

Notes	<b>Life Skills Training (LST)</b> <u>Abbreviations</u> FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); ANCOVA=analysis of covariance; LST=life skills training; NA=not applicable; mo=month(s)	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT; attrition < 10%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Not described

**Botvin 1995**

Methods	Design: RCT FU: 6 yrs (post-randomisation) Attrition: 25% (year 3) and 40% (year 6) ITT: no Unit of randomisation: school (stratified by smoking behavior) Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: NR (NR) Int-2: NR (NR) Ctrl: NR (NR) Total N: 57 (NR) <u>N of Clusters (subjects) participated</u> Int-1: 18 (NR) Int-2: 16 (NR) Ctrl: 22 (NR) Total N: 56 (5954)

**Botvin 1995** (Continued)

	<p><u>Sample analysed: 3 yrs</u>            Int-1: 18 (NR)            Int-2: 16 (NR)            Ctrl: 22 (NR)            Total N: 56 (4466)</p> <p><u>Sample analysed: 6 yrs</u>            Int-1: 18 (1128)            Int-2: 16 (1327)            Ctrl: 22 (1142)            Total N: 56 (3597)            Age: 12-13 yrs (7th grade)            Sex (male): 52%            Ethnicity: 91% White            Alcohol users at baseline: 18% monthly drinkers            Country: US</p>
Interventions	<p><u>Intervention-1</u>: a multi-component substance abuse prevention curriculum through formal teacher training and delivery with implementation feedback  <u>Intervention-2</u>: a multi-component substance abuse prevention curriculum through video training for teacher with no implementation feedback  <u>Focus/target</u>: the major social, psychological, cognitive, and attitudinal factors that promote the use of tobacco, alcohol and marijuana  <u>Components</u>: cognitive behavioral skills to raise self-esteem, resistance, assertiveness, relationship, anxiety management &amp; communication skills  <u>Fidelity</u>: one school post-randomisation was lost. Randomly selected classes monitored and average implementation fidelity scored at 68%  <u>Duration/frequency</u>: 15 sessions (1 year) and 10 + 5 booster sessions in the following 2 years  <u>Control</u>: standard curriculum</p>
Outcomes	<p><u>FU: 3 yrs</u>            Results for FU of 3 yrs were reported for “high fidelity” sample only and as this breaks the randomisation and increases the risk of bias then these results are not reported here  <u>FU: 6 yrs</u> (Int-1 vs. Int-2 vs. Ctrl)            Monthly alcohol use (0=no, 1=yes): 0.61 (0.02) vs. 0.57 (0.03) vs. 0.60 (0.02), p &gt; 0.05            Weekly alcohol use (0=no, 1=yes): 0.29 (0.02) vs. 0.24 (0.023) vs. 0.29 (0.02), p &gt; 0.05            3+ drinks per occasion (0=no, 1=yes): 0.57 (0.02) vs. 0.55 (0.03) vs. 0.59 (0.02), p &gt; 0.05            Drunkenness in last month (0=no, 1=yes): 0.34 (0.02)* vs. 0.33 (0.03)* vs. 0.40 (0.02), *p&lt;0.05 (vs. Ctrl; 1-tailed test)</p>
Notes	<p><b>Life Skills Training (LST)</b>            Response rate was non-differential  <u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s);            ITT=intention to treat (analysis); LST=life skills training; NA=not applicable; mo=</p>

**Botvin 1995** (Continued)

month(s)		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	no ITT; attrition: 25%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	No baseline imbalance

**Botvin 2001**

Methods	Design: RCT FU: 2 yrs (post-randomisation) Attrition: 20% (year 1), 42% (year 2) ITT: no Unit of randomisation: school (stratified by smoking prevalence) Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: 16 (NR) Ctrl: 13 (NR) Total N: 29 (5233) <u>N of Clusters (subjects) participated</u> Int: 16 (NR) Ctrl: 13 (NR) Total N: 29 (5233)  <u>Sample analysed (year 2)</u> Int: 16 (NR) Ctrl: 13 (NR) Total N: 29 (3041) Age: 12-13 yrs (7th grade) Sex (male): 44% Ethnicity: 57% African American, 24% Hispanic, 8% Asian, 3% White

**Botvin 2001** (Continued)

	Alcohol users: 1-2% already binge drinkers Country: US
Interventions	<u>Intervention</u> : a multi-component substance abuse prevention curriculum <u>Focus/target</u> : the major social, psychological, cognitive, and attitudinal factors that promote the use of tobacco, alcohol and marijuana <u>Components</u> : cognitive behavioral skills to raise self-esteem, resistance, assertiveness, relationship, anxiety management & communication skills <u>Fidelity</u> : randomly selected classes monitored, with average of 48% coverage achieved <u>Duration/frequency</u> : 15 sessions in 7 <sup>th</sup> Grade and 10 booster sessions in 8 <sup>th</sup> Grade <u>Control</u> : standard curriculum
Outcomes	FU: 1 yr Binge drinking (> 5 drinks per occasion): Int: 1.8% vs. Ctrl: 4.3% OR (adjusted)=0.41, 95% CI: 0.18, 0.93, N=2982  FU: 2 yrs Binge drinking (> 5 drinks per occasion): Int: 2.2% vs. Ctrl: 5.2% OR (adjusted)=0.40, 95% CI: 0.22,0.74, N=2982
Notes	<b>Life Skills Training (LST)</b> Notable baseline differences in ethnicity and free lunch status. Response rate was < 60% and non-differential <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); LST=life skills training; NA=not applicable; mo=month(s); OR=odds ratio; CI=confidence interval

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	no ITT; attrition: 20%
Selective reporting (reporting bias)	High risk	some drinking outcomes not reported

**Botvin 2001** (Continued)

Other bias	High risk	baseline differences in ethnicity and free lunch status
------------	-----------	---

**Botvin 2003**

Methods	<p>Design: RCT  FU: 3 yrs (post-randomisation)  Attrition: 42% unmatched students; matched students, relative to unmatched, had lower smoking prevalence (8.6% vs. 11%) and higher proportion of White students (48% vs. 38%)  ITT: No (778 unmatched students were not included in the analyses)  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p>N of Clusters (subjects) randomized: NR  N of Clusters (subjects) participated: Int: 9 (426) vs. Ctrl: 11 (664)  Total N Clusters (subjects): 20 (1090)  Age: NR (3-6 graders)  Sex (male): 52%  Ethnicity: White (48%), Hispanic (26%), African-American (13%), Asian (3%)  Alcohol users: 35%  Country: US</p>
Interventions	<p><u>Intervention</u>: LST  <u>Focus/target</u>: provide knowledge and skills for resisting social influence to use alcohol/tobacco leading to reduction of alcohol/tobacco use  <u>Components</u>: personal self-management, general social skills  <u>Fidelity</u>: random monitoring of teachers  <u>Duration/frequency</u>: 8 classes/yr for 3 yrs  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 3 yrs  <u>Individual level analysis</u>  Drinking frequency (Int vs. Ctrl): 0.44 (±0.04) vs. 0.41 (±0.03), p=0.28  Drink in past yr (Int vs. Ctrl): 0.15 (±0.01) vs. 0.16 (±0.01), p=0.17    <u>School level analysis</u>  Drinking frequency (Int vs. Ctrl): 0.43 (±0.06) vs. 0.52 (±0.05), p=0.14  Proportion drink in past yr (Int vs. Ctrl)  0.13 (±0.02) vs. 0.17 (±0.02), p=0.054</p>
Notes	<p><b>Life Skills Training (LST)</b>  Not clear if attrition was differential; int arm had lower proportion of Whites than Ctrl (44.1% vs. 50.9%, p=0.017)  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); ITT=intention to treat (analysis); LST=life skills training; NA=not ap-</p>

**Botvin 2003** (Continued)

	plicable; mo=month(s)	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	no ITT; attrition: 42%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	differences across study arms

**Brewer 1991**

Methods	Design: RCT FU: 6 mo (post-randomisation) Attrition: 0% (year 1) ITT: yes Unit of randomisation: student Clustering effect adjusted: NA
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: NR (18) vs. Int-2: NR (18) vs. Ctrl: NR (18) Total N: NR (54) <u>N of Clusters (subjects) participated</u> Int-1: NR (18) Int-2: NR (18) Ctrl: NR (18) Total N: NR (54)  Age: 15-16 yrs Sex (male): 50% Ethnicity: NR Alcohol users: 9% used monthly Country: US

**Brewer 1991** (Continued)

Interventions	<p><u>Intervention-1</u>: social skills development curriculum  <u>Intervention-2</u>: placebo - video tapes of drug and alcohol material  <u>Focus/target</u>: decrease onset and use of tobacco and alcohol  <u>Components</u>: development of social skills through 5 steps; recognizing need for skills; modelling 5 steps; role playing; rehearsal; feedback  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 9 x 40 minute sessions over 9 weeks  <u>Control</u>: standard curriculum</p>	
Outcomes	<p>FU: 6 mo  Alcohol use (0-19 scale; 0=non-user): Int-1: 10.7 (+3.1) vs. Int-2: 13.0 (no change) vs. Ctrl: 13.1 (-0.7)  (F = 4,863, df = 2, p = 0.014; higher alcohol use in the experimental group due to some methodological errors)  No significant effect (level of significance adjusted for family wise error rate) for initiation, experimental and regular alcohol use</p>	
Notes	<p><b>HLAY2</b>  Response rate was 100%. Possible contamination as study in one school only. Some concerns over methodological errors in recording alcohol use and validity/reliability of questionnaire measures  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); df=degrees of freedom; HLAY2= here's looking at you</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT done; Attrition: < 10%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported

**Brewer 1991** (Continued)

Other bias	High risk	Possible contamination as study in one school only. Some concerns over methodological errors in recording alcohol use and validity/reliability of questionnaire measures
------------	-----------	--

**Caplan 1992**

Methods	Design: RCT FU: 15 wks (post-randomisation) Attrition: 17% ITT: no Unit of randomisation: class, stratified by ability Clustering effect adjusted: yes
Participants	Total N of Clusters (subjects) randomised: NR (298) N of Clusters (subjects) participated: Int: NR (NR) vs. Ctrl: NR (NR) Total N of Clusters (subjects) participated: NR (282) Age: 11-13 yrs Sex (male): 55% Ethnicity: 90% Black Alcohol users: NR Country: US
Interventions	<u>Intervention</u> : a school-based social competence / skills curriculum <u>Focus/target</u> : promoting personal and social competence to reduce self-reported substance use intentions and excessive alcohol use <u>Components</u> : social competence training; stress management; self-esteem; problem solving; substances and health; assertiveness; social networks <u>Fidelity</u> : NR <u>Duration/frequency</u> : 20 sessions over 15 weeks <u>Control</u> : standard curriculum
Outcomes	FU: 15 wks <u>3+ drinks per occasion (ANOVA)</u> F (1,213)=3.65, p<0.05; Ctrl higher <u>Drinking too much (ANOVA)</u> F (1,213)=3.68, p<0.05; Ctrl higher <u>Amount usually consumed per occasion (ANOVA)</u> F (1,213)=5.65, p<0.02; Ctrl higher Other measures of drinking behaviour were listed in method but not reported in results
Notes	<b>Positive Youth Development Program (PDYP)</b> Risk of contamination within schools. One class reassigned to Ctrl group because of "scheduling difficulties" <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;



**Caplan 1992** (Continued)

	wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); ANOVA=analysis of variance; PDYP= positive youth development program	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT
Selective reporting (reporting bias)	High risk	Some measures of alcohol drinking listed but not reported
Other bias	High risk	Possible contamination since classes were randomized within schools

**Clayton 1991**

Methods	Design: RCT FU: 2, 5, 10 yrs post-randomisation Attrition: 21% (year 2 post-randomisation), 45% (year 5 post-randomisation), 52% (year 10 post-randomisation) ITT: no Unit of randomisation: school Clustering effect adjusted: yes (at 5 and 10 yrs)
Participants	N of Clusters (subjects) randomised: Int: 23 (NR) vs. Ctrl: 8 (NR) Total N of Clusters (subjects) randomised: 31 (NR) Analysed sample (10 years): Int: 23 (NR) vs. Ctrl: 8 (NR) Total N analysed clusters (subjects) at year 10: 31 (1002) Age: 11-12 yrs Sex (male): 51% Ethnicity: 75% White Past yr of alcohol use: Int (19.8%) vs. Ctrl (16.2%), p<0.05 Country: US
Interventions	<u>Intervention</u> : DARE - social and life skills curriculum <u>Focus/target</u> : teaching students skills for recognizing and resisting social pressure to use

Clayton 1991 (Continued)

	<p>drugs  <u>Components:</u> resistance training, self-esteem, social skills, information, role-play. Delivered by trained uniformed police officer  <u>Fidelity:</u> NR  <u>Duration/frequency:</u> 17 1-hr sessions / 17 weeks  <u>Control:</u> standard drug education</p>	
Outcomes	<p>FU: 2 yrs  <u>Past yr of alcohol use (Int vs. Ctrl):</u> 38.3% vs. 38.3%, <math>p \geq 0.05</math>  FU: 5 yrs  <u>Frequency of past yr alcohol use:</u> standardized beta coefficient for DARE status in HLM = -0.12 (<math>p \geq 0.05</math>)  FU: 10 yrs  <u>Frequency of past mo alcohol use:</u> standardized beta coefficient for DARE status in HLM = -0.18 (<math>p \geq 0.05</math>)  DARE status was not related to alcohol use at age 20</p>	
Notes	<p><b>Project DARE</b>  Some evidence of differential attrition. Baseline imbalance in alcohol use  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s);  DARE=drug abuse resistance education;  HLM= hierarchical linear model</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition > 20% (possibly differential)
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Baseline imbalance in alcohol use

**Cook 1984**

Methods	<p>Design: RCT  FU: 2 yrs (post-randomisation)  Attrition: 16%  ITT: no  Unit of randomisation: individual  Clustering effect adjusted: no</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: NA (90)  Ctrl: NA (64)  Total N: NA (154)  <u>N of Clusters (subjects) participated</u>  Int: NA (90)  Ctrl: NA (64)  Total N: NA (154)</p> <p><u>Analysed sample (year 1)</u>  Int: NA (77)  Ctrl: NA (57)  Total N: NA (134)</p> <p><u>Analysed sample (year 2)</u>  Int: NA (58)  Ctrl: NA (50)  Total N: NA (108)  Age: 15-16 yrs  Sex (male): 53%  Ethnicity: 66.4% (White), 32% (Black), and 1.5% (Oriental)  Alcohol users: 31%  Country:US</p>
Interventions	<p><u>Intervention</u>: additional classes meeting during school-time to present healthy alternatives to drug abuse  <u>Focus/target</u>: promote consideration of alternative activities to reduce substance use  <u>Components</u>: PAY orientation, communication, self-concept, self-care, activities, physical, creative self-expression, consciousness alteration  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 2-3 sessions/week over 1 semester  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 1 yr  <u>Hard liquor use in the last 2 mo (not at all - daily)</u>  Int (pre: 1.53 vs. post: 1.40)  Change: -0.13  Ctrl (pre: 1.49 vs. post: 1.55)  Change: +0.06  F=6.72, p=0.01 (in favour of Int)  <u>Beer/wine use in the last 2 mo (not at all - daily)</u></p>

**Cook 1984** (Continued)

	(F=0.298, p=0.58)	
	FU: 2 yrs No results were presented as all analyses reportedly were non-significant	
Notes	<p><b>Positive Alternatives for Youth (PAY)</b> Marked differences at baseline between Int. and Ctrl. Differences in baseline characteristics only taken into account in Study 2 analyses</p> <p><u>Abbreviations</u> FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); PAY=positive alternatives for youth; pre=pre-treatment; post=post-treatment</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT
Selective reporting (reporting bias)	High risk	Second year results of alcohol use not reported
Other bias	High risk	Differences in baseline characteristics for Study 1 not taken into account

**Dielman 1986**

Methods	<p>Design: RCT FU: 3-4 yrs (post-randomisation) Attrition: 12% (post-test), 33% (year 2), 36% (year 4) ITT: no Unit of randomisation: school building Clustering effect adjusted: yes</p>
---------	---

**Dielman 1986** (Continued)

Participants	<p><u>N of Clusters (subjects) randomised</u>            Int: NR (NR)            Ctrl: NR (NR)            Total N: 49 (5635)</p> <p><u>N of Clusters (subjects) participated</u>            Int: NR (NR)            Ctrl: NR (NR)            Total N: NR (NR)</p> <p>Only half the randomised sample were pre-tested and analysed at follow-up:</p> <p><u>Analysed sample</u>            Int: NR (1499)            Ctrl: NR (706)            Total N: NR (2280)            Age: 10-12 yrs            Sex (male): NR            Ethnicity: NR            Alcohol users: 7%-13%            Country: US</p>
Interventions	<p><u>Intervention</u>: social learning approach to alcohol misuse prevention  <u>Focus/target</u>: the acquisition of behavioral skills to resist pressures to misuse alcohol  <u>Components</u>: providing information and social skills; awareness of risks and pressures to drink; peer pressure resistance &amp; refusal skills  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 4 sessions (each 45 min) + booster sessions  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU - Immediate post-intervention            [Unadjusted for prior drinking]  <u>Mean (SD) alcohol weekly use in past 12 mo</u> [Never=0 to every day=5]            Int: pre 0.26 (0.70) vs. post 0.41 (0.77)            Ctrl: pre 0.29 (0.69) vs. post 0.44 (0.79)            Between-group: NS</p> <p><u>Mean (SD) alcohol weekly misuse in past 12 mo (overindulgence)</u> [Never=0 to three or more times =3]            Int: pre 0.31 (0.70) vs. post 0.39 (0.78)            Ctrl: pre 0.36 (0.75) vs. post 0.43 (0.82)            Between-group: NS</p> <p>FU: 3 yrs            [Adjusted for prior drinking]            In those with prior alcohol drinking experience, the treatment group had greater post-intervention reduction in the rate of increase of alcohol use; in those with no prior alcohol drinking experience, there was no difference between the treatment</p>

**Dielman 1986** (Continued)

and control groups (total alcohol misuse -  $p < 0.01$ ; alcohol frequency-quantity index -  $p < 0.05$ )

FU: 4 yrs  
Follow-up on a sample subset found no significant effect of the AMPS curriculum on tenth grade alcohol misuse

**Notes**

**Alcohol Misuse Prevention Study (AMPS)**  
At FU immediately after treatment, prior alcohol drinking was not controlled and crude analysis did not reveal treatment effects on alcohol misuse; at FU 3yrs, controlling for prior alcohol drinking revealed treatment effects in reducing rates of alcohol misuse/use. Pre-test and analysis only undertaken in half of participating schools, reducing statistical power; attrition rates not clearly reported across publications; students who missed all booster or intervention sessions were coded as not having received intervention.

Abbreviations  
FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); AMPS=alcohol misuse prevention study; NS=statistically non-significant; SD=standard deviation; pre=pre-treatment; post=post-treatment

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition > 33%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Prior alcohol drinking not controlled

**Durrant 1986**

Methods	Design: RCT FU: 22 wks (post-randomisation) Attrition: NR ITT: NR Unit of randomisation: school Clustering effect adjusted: no
Participants	<u>N of Clusters (subjects) randomised</u> Int: 4 (NR) Ctrl: 4 (NR) Total N: 8 (NR) <u>N of Clusters (subjects) participated</u> Int: 4 (NR) Ctrl: 4 (NR) Total N: 8 (NR) <u>Analysed sample -N of Clusters (subjects)</u> Int: 4 (102) Ctrl: 4 (89) Total N: 8 (191) Age: 11-12 yrs Sex (male): 48% Ethnicity: NR Alcohol users: NR Country: US
Interventions	<u>Intervention: multi-component "inoculation programme"</u> <u>Focus/target:</u> health education through lectures, class discussions, and student exercises <u>Components:</u> anti-smoking strategy applied to alcohol & other drugs: emphasizing healthy lifestyles; negative effects of drug use; decision making skills; skills to resist social pressures <u>Fidelity:</u> NR <u>Duration/frequency:</u> 9 x 45 min sessions over 22 weeks <u>Control:</u> NR
Outcomes	FU: 22 wks <u>Mean (SD) liquor use (6-point scale)</u> Int: 1.54 (0.35) vs. Ctrl: 1.15 (0.64), $p > 0.05$ , NS F (1,184) < 1
Notes	<b>Multi-component "inoculation programme"</b> Randomization was stratified by school size, minority population and the number of free lunches <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; SD=standard deviation

***Risk of bias***

**Durrant 1986** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not described
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Not described

**Duryea 1984a**

Methods	Design: RCT FU: 3 yrs (post-randomisation) Attrition: 16% (year 3) ITT: no Unit of randomisation: individual Clustering effect adjusted: NA
Participants	<u>N of Clusters (subjects) randomised</u> Int: NA (NR) Ctrl: NA (NR) Total N: NA (155) <u>N of Clusters (subjects) participated</u> Int: NA (NR) Ctrl: NA (NR) Total N: NA (155) <u>Analysed sample</u> Int: NA (91) Ctrl: NA (39) Total N: NA (130) Age: 14-15 yrs Sex (male): NR Ethnicity: NR Alcohol users: NR Country: US



Duryea 1984a (Continued)

Interventions	<p><u>Intervention</u>: a psychological “immunization” program  <u>Focus/target</u>: forewarn students about the dangers of drugs to “immunize” them, especially with regard to peer pressure  <u>Components</u>: film, question &amp; answer session; highlighting specific alcohol related concepts; role playing and ability to refuse  Some booster activities  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 6 sessions/2 weeks  <u>Control</u>: NR</p>	
Outcomes	<p><u>FU</u>: 6 mo  Frequency of self-reported alcohol drinking did not significantly differ in Int vs. Ctrl, <math>p &gt; 0.05</math> (NS)  <u>FU</u>: 3 yrs  Mean (SD) drinking occasions in past 2 weeks - Int: 2.06 (1.11) vs. Ctrl: 2.05 (0.97), T (128)=-0.07, <math>p = 0.94</math> (NS)  Mean (SD) N times drank too much in past 2 weeks - Int: 1.48 (0.82) vs. Ctrl: 1.20 (0.47), T (128)=-1.98, <math>p = 0.05</math> (NS)</p>	
Notes	<p><b>Preventive Alcohol Education Program</b>  Compliance Increased in Int group (6 mo post-test vs. baseline, <math>p &lt; 0.05</math>) and decreased in Ctrl (6 mo post-test vs. baseline, <math>p &lt; 0.0005</math>). Response rate was c.84% but differential attrition rates between groups  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant;  SD=standard deviation; pre=pre-treatment; post=post-treatment</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors’ judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	High risk	Assessors were not blinded
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; differential attrition
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported

Duryea 1984a (Continued)

Other bias	Unclear risk	Not described
------------	--------------	---------------

D'Amico 2002

Methods	<p>Design: RCT  FU: 6 mo (post-randomisation)  Attrition: More students dropped from Ctrl vs. Int 1-2 arms (p&lt;0.01); amongst control students, male sex, low SES, and endorsement of strong positive expectancy were associated with higher dropout rates  ITT: No  Unit of randomisation: student  Clustering effect adjusted: NA</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int 1: NA (75)  Int 2: NA (75)  Ctrl: NA (150)  Total N: NA (300)  <u>N of Clusters (subjects) participated</u>  Int 1: NA (NR)  Int 2: NA (NR)  Ctrl: NA (NR)  Total N: NA (NR)</p> <p>Age: 14-19 yrs  Sex (male): 42%  Ethnicity: White (63%), Hispanic (17%), African-American (10%), Asian (2%), Other (8%)  Alcohol users: heavy drinkers (8%), moderate drinkers (13%), light drinkers (18%), abstainers (46%)  Country: US</p>
Interventions	<p><u>Intervention -1: RSTP</u> - a brief interactive and motivational program  <u>Focus/target:</u> multiple risk behaviours, personal beliefs, and experienced consequences to reduce participation in risk behaviors  <u>Components:</u> Interactive group session, exchange of confidential and personalized graphic feedback regarding risk taking behavior, discussion on peer influence  <u>Fidelity:</u> 5.3 (±0.58)  <u>Duration/frequency:</u> single 50 min session  <u>Intervention-2: DARE-A</u> is a brief didactic, education based program  <u>Focus/target:</u> Increasing knowledge of deleterious effects of substance use  <u>Components:</u> how substances change mind/body, drugs and the law, consequences, assertive resistance  <u>Fidelity:</u> 2.0 (±0.00)  <u>Duration/frequency:</u> single 50 min session  <u>Control:</u> standard curriculum</p>

D'Amico 2002 (Continued)

Outcomes	<p>FU-2 mo (Int-1 vs. Int-2 vs. Ctrl)</p> <p><u>Weekly drinking (0-21 drinks)</u></p> <p>2.82 (±4.19) vs. 2.64 (±4.42) vs. 2.48 (±4.14), p&gt;0.05 (NS)</p> <p><u>Risky drinking (0-41 drinks in 3 mo)</u></p> <p>1.42 (±2.98) vs. 0.84 (±2.85) vs. 1.59 (±4.92), p&gt;0.05 (NS)</p> <p>FU-6 mo (Int-1 vs. Int-2 vs. Ctrl)</p> <p><u>Weekly drinking (0-21 drinks)</u></p> <p>2.76 (±4.05) vs. 1.78 (±3.23) vs. 3.44 (±4.74), p&gt;0.05 (NS)</p> <p><u>Risky drinking (0-41 drinks in 3 mo)</u></p> <p>1.90 (±3.68) vs. 1.06 (±2.76) vs. 2.36 (±4.70), p&gt;0.05 (NS)</p>	
Notes	<p>No significant baseline differences between the trial arms.</p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; SD=standard deviation; RSTP=risk skills training program; DARE-A=drug abuse and resistance education</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; differential attrition rates
Selective reporting (reporting bias)	High risk	One relevant outcome not reported
Other bias	Low risk	No significant baseline differences

Eisen 2002

Methods	<p>Design: RCT  FU: 2 yrs (post-randomisation)  Attrition: 16% (1<sup>st</sup> yr) dropout rates were associated with marijuana use, Hispanic American race, not from two-parent household  ITT: Yes  Unit of randomisation: school  Clustering effect adjusted: Yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 17 (NR)  Ctrl: 17 (NR)  Total N: 34 (7426)  <u>N of Clusters (subjects) participated</u>  Int: NR (NR)  Ctrl: NR (NR)  Total N: NR (6239) 1<sup>st</sup> yr FU  Total N: NR (5691) 2<sup>nd</sup> yr FU</p> <p>Age: ≤14 yrs (6<sup>th</sup> graders)  Sex (male): 48.3%  Ethnicity: White (25.7%), Hispanic (34%), African-American (17.6%), Asian (7.1%), Other (6.3%)  Recent (30 days) use of alcohol: 9.5%  Country: US</p>
Interventions	<p><u>Intervention</u>: SFA - a multi-strategy program to teach social competency and refusal skills  <u>Focus/target</u>: teaches cognitive behavioral skills for building self-esteem and personal responsibility, communicating effectively, making better decisions, and resisting social influences  <u>Components</u>: 3 sessions on challenges of entering the teen yrs, 4 sessions on building self-confidence and communication skills, 5 sessions on managing emotions in positive way, 8 sessions on improving peer relationships, resisting peer pressure, and 20 sessions on living healthy and drug free  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 40 sessions over 1 yr (35-45 min each)  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU-1 yr (Int vs. Ctrl)  <i>Substance non-users at baseline</i>  Alcohol use-lifetime: 29.61% vs. 30.19% (RD=-0.58, 95% CI: -3.11, 4.27)  Alcohol use-recent: 7.17% vs. 7.25% (RD=-0.08, 95% CI: -2.33, 1.57)  Binge drinking: 3.15% vs. 3.58% (RD=-0.43, 95% CI: -1.91, 0.66)</p> <p><i>Substance users at baseline</i>  Binge drinking: 16.98% vs. 20.45% (RD=-3.47, 95% CI: -15.07, 8.14)</p> <p>FU-2 yr (Int vs. Ctrl)  Alcohol use-lifetime: 66.97% vs. 66.33% (RD=0.64, 95% CI: -2.25, 3.53)  Alcohol use-recent: 22.85% vs. 23.18% (RD=-0.33, 95% CI: -3.01, 2.35)</p>

**Eisen 2002** (Continued)

	Binge drinking: 12.67% vs. 13.11% (RD=-0.44, 95% CI: -2.78, 1.91)
Notes	<p><b>Lions-Quest “Skills for Adolescence”</b>            The effect of SFA was greater in Hispanic compared with non-Hispanic Americans</p> <p><u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;            wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s);            NS=statistically non-significant;            SD=standard deviation; RSTP=risk skills training program; DARE-A=drug abuse and resistance education;            SFA=skills for adolescence; min=minute(s); RD=risk difference; 95% CI: ninety-five percent confidence interval</p>

**Risk of bias**

Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT and attrition < 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Not described

**Ellickson 1990**

Methods	Design: RCT FU: 3 mo, 12 mo, 5 yrs (post-randomisation) Attrition: 40% (15 mo), 25% (year 2), 45% (year 5) ITT: no Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 10 (NR) Int-2: 10 (NR) Ctrl: 10 (NR)

Ellickson 1990 (Continued)

	<p>Total N: 30 (6527)  <u>N of Clusters (subjects) participated</u>            Int-1: 10 (NR)            Int-2: 10 (NR)            Ctrl: 10 (NR)            Total N: 30 (6527)  <u>Analysed sample</u>            Int-1: 10 (NR)            Int-2: 10 (NR)            Ctrl: 10 (NR)            Total N: 30 (3852)            Age: 12-14 yrs (Grades 7 &amp; 8)            Sex (male): 52%            Ethnicity: White (67%), Hispanic (10%), Black (10%), Asian (8%), and Indian/mixed (5%)            Alcohol users (ever): 77%            Country: US</p>
Interventions	<p>Intervention-1: teen leader (curriculum synthesized aspects of health belief model, social learning theory and self-efficacy theory)            Intervention-2: adult health educator (curriculum synthesized aspects of health belief model, social learning theory and self-efficacy theory)  <u>Focus/target</u>: change norms and beliefs about drug use; identify and resist peer pressure  <u>Components</u>: develop reasons not to use drugs, discuss pressures to use drugs, resistance skills, prevalence of drug use  <u>Fidelity</u>: 92% observed classes delivered curriculum as intended  <u>Duration/frequency</u>: 2 yrs; 7th grade - 8 sessions; 8th grade - 3 sessions  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 3 mo  <i>Non-users of alcohol at baseline (N=953)</i>  <u>Alcohol use - ever (%)</u>            Int-1: 16.3* vs. Int-2: 18.0 vs. Ctrl: 22.8  <u>Alcohol use past mo (%)</u>            Int-1: 5.9* vs. Int-2: 8.0 vs. Ctrl: 10.8    <i>Users of alcohol at baseline (N=1130)</i>  <u>Alcohol use past mo (%)</u>            Int-1: 69.6 vs. Int-2: 62.7 vs. Ctrl: 69.5              FU: 12 mo  <i>Non-users of alcohol at baseline (N=953)</i>  <u>Alcohol use - ever (%)</u>            Int-1: 47.4 vs. Int-2: 45.5 vs. Ctrl: 50.0  <u>Alcohol use past mo (%)</u>            Int-1: 14.4 vs. Int-2: 10.5 vs. Ctrl: 14.6    <i>Users of alcohol at baseline (N=1130)</i>  <u>Alcohol use past mo (%)</u></p>

	Int-1: 73.0 Int-2: 70.7 Ctrl: 71.6  *p<0.05 (vs. Ctrl)  FU: 5 yrs Alcohol use [absolute amount in % greater or less than Ctrl group prevalence] stratified by baseline risk (1 = nonuser, 2 = experimenter, 3 = user)			
	<b>Baseline risk</b>	<b>1</b>	<b>2</b>	<b>3</b>
	<b>N</b>	<b>(855)</b>	<b>(1569)</b>	<b>(1042)</b>
	<u>Lifetime (%)</u>			
	Ctrl	82.5	100	100
	Int-1	0	na	na
	Int-2	2.7	na	na
	<u>Past yr (%)</u>			
	Ctrl	54.0	80.5	91.3
	Int-1	3.4	-0.6	2.2
	Int-2	0.9	2.4	-0.4
	<u>Past mo (%)</u>			
	Ctrl	34.1	56.2	70.2
	Int-1	1.3	-3.3	4.6
	Int-2	0.2	0.4	-4.1
	<u>Monthly (%)</u>			
	Ctrl	24.2	41.3	59.9
	Int-1	-1.7	1.3	-1.0
	Int-2	-4.2	-1.9	-6.8
	<u>Weekly (%)</u>			
	Ctrl	8.3	10.2	13.9
	Int-1	-1.1	-0.3	1.3
	Int-2	-4.4*	-0.9	0
	<u>Daily (%)</u>			
	Ctrl	0.4	0.2	1.0
	Int-1	na	na	na
	Int-2	na	na	na
	*p<0.05 (vs. Ctrl)			
Notes	<p><b>Project ALERT</b>                  Attrition 25% to 45% but no evidence of differential attrition. No adjustment for multiple testing</p> <p><u>Abbreviations</u>                  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;                  wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant;                  SD=standard deviation; na=not analysed (because frequency was either 100% or &lt;2%)</p>			

Ellickson 1990 (Continued)

<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	Attrition > 20% but not differential
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	No evidence of baseline difference or Columbia

Ellickson 2003

Methods	Design: RCT FU: 18 mo (post-randomisation) Attrition: 9% ITT: No Unit of randomisation: school Clustering effect adjusted: Yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: NR (2810) Ctrl: NR (1879) Total N: 55 (4689)  <u>N of Clusters (subjects) participated</u> Int: 34 (2553) Ctrl: 21 (1723) Total N: 55 (4276) Age: NR (7 <sup>th</sup> and 8 <sup>th</sup> graders) Sex (male): 50% Ethnicity: Non-White (12.5%) Ever tried alcohol: 60% Country: US



Interventions	<p><u>Intervention:</u> ALERT - program</p> <p><u>Focus/target:</u> the program includes 3 theories in behavioral change:</p> <ol style="list-style-type: none"> <li>1) health belief model which focuses on cognitive factors that motivate healthy behavior</li> <li>2) social learning model which emphasizes social norms and significant others as key determinants of behavior</li> <li>3) self-efficacy theory of behavior change to accomplish a task as essential to effective action</li> </ol> <p><u>Components:</u> interactive teaching methods such as question and answer technique and small group activities done in 11 sessions in 7<sup>th</sup> grade and 3 sessions in 8<sup>th</sup> grade</p> <p><u>Fidelity:</u> 88% activities accomplished for 7<sup>th</sup> grade and 93% for 8<sup>th</sup> grade</p> <p><u>Duration/frequency:</u> 14 lessons in 18 mo</p> <p><u>Control:</u> standard curriculum</p>
Outcomes	<p>FU-18 mo</p> <p>The mean scores (using specific alcohol scales) for overall misuse and high risk drinking were significantly lower (i.e., improved) in Int vs. Ctrl (p&lt;0.05)</p> <p><i>Baseline non-users</i> (Int vs. Ctrl)</p> <p>Overall misuse: 0.22 vs. 0.30, NS</p> <p>Alcohol-related consequence: 0.13 vs. 0.18, NS</p> <p>High risk use: 0.10 vs. 0.11, NS</p> <p><i>Baseline users</i> (Int vs. Ctrl)</p> <p>Overall misuse: 1.78 vs. 2.23, p&lt;0.05</p> <p>Alcohol-related consequence: 1.04 vs. 1.29, p&lt;0.05</p> <p>High risk use: 0.74 vs. 0.92, p&lt;0.01</p>
Notes	<p><b>Project ALERT</b></p> <p>Although dropouts were more likely to be male, non-White, having low grades, single-parent family, users of alcohol or marijuana, the attrition rates did not significantly differ across the arms</p> <p>The effect of ALERT was observed only in baseline user (high risk) students but not in non-user students</p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant</p>

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described

**Ellickson 2003** (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Dropouts and analyzed sample compared; attrition < 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	No obvious differences across study arms

**Faggiano 2007**

Methods	<p>Design: RCT  FU: 3, 18 mo (post-randomisation)  Attrition: 10.0% (3 mo post-baseline), 21.7% (18 mo post-baseline)  ITT: yes  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 102 (NR)  Ctrl: 68 (NR)  Total N: 170 (NR)  <u>N of Clusters (subjects) participated at 18 mo FU</u>  Int: NR (2811)  Ctrl: NR (2730)  Total N: NR (5541)  Analyzed sample: N = 5541  Age: 12-14 yrs (7th-9<sup>th</sup> grades)  Sex (male): 52%  Ethnicity: NR  Alcohol use: 24.1%  Any drunkenness: 6%  Frequent drunkenness: 1.8%  Multi-center study - EU-DAP (Austria, Belgium, Germany, Greece, Italy, Spain, and Sweden)</p>
Interventions	<p><u>Intervention</u>: EU-Dap school program in 3 formats:  1) class curriculum alone, complemented with activities involving either 2) family or 3) peer  <u>Focus/target</u>: experimental and regular use of alcohol, tobacco, and illicit drugs  <u>Components</u>: Social skills, personal skills, knowledge and normative education  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 12 sessions (1 hr each) weekly  <u>Control</u>: standard curriculum</p>

<p>Outcomes</p>	<p><b>FU: 3 mo (Int vs. Ctrl)</b></p> <p><u>Any drunkenness in the past mo (%)</u>  OR [3-level adjusted] = 0.69 (95% CI: 0.48, 0.99); ARR=1.2%; NNT=82 (95% CI: 47, 305)</p> <p><u>Frequent drunkenness in the past mo (%)</u>  OR [3-level adjusted] = 0.77, 95% CI: 0.60, 1.00; ARR=1.7%; NNT=60, 95% CI: 34, 223</p> <p><u>Any drunkenness in the past mo (%)</u>  All Boys: OR [3-level adjusted] = 0.64 (95% CI: 0.49, 0.85)  Boys with high self-esteem: OR [3-level adjusted] = 0.69 (95% CI: 0.50, 0.94)  Boys with low self-esteem: OR [3-level adjusted] = 0.58 (95% CI: 0.29, 1.17)  All Girls: OR [3-level adjusted] = 0.86 (95% CI: 0.63, 1.18)  Girls with high self-esteem: OR [3-level adjusted] = 0.83 (95% CI: 0.58, 1.19)  Girls with low self-esteem: OR [3-level adjusted] = 1.23 (95% CI: 0.66, 2.29)</p> <p><u>Frequent drunkenness in the past mo (%)</u>  All Boys: OR [3-level adjusted] = 0.68 (95% CI: 0.45, 1.04)  Boys with high self-esteem: OR [3-level adjusted] = 0.71 (95% CI: 0.43, 1.14)  Boys with low self-esteem: OR [3-level adjusted] = 0.75 (95% CI: 0.25, 2.19)  All Girls: OR [3-level adjusted] = 0.66 (95% CI: 0.37, 1.18)  Girls with high self-esteem: OR [3-level adjusted] = 0.59 (95% CI: 0.31, 1.12)  Girls with low self-esteem: OR [3-level adjusted] = 1.71 (95% CI: 0.49, 5.92)</p> <p><b>FU: 18 mo (Int vs. Ctrl)</b></p> <p><u>Weekly drinking (%)</u>  Total sample: OR [3-level adjusted] =0.93, (95% CI: 0.79, 1.09)  Baseline drinkers: OR [3-level adjusted] =0.95, (95% CI: 0.72, 1.27)  Baseline non-drinkers: OR [3-level adjusted] =0.88, (95% CI: 0.73, 1.07)</p> <p><u>Alcohol drinking (current)</u></p> <p><i>Baseline non-drinkers</i>  None: 73.9% vs. 71.9%, p&gt;0.05  Occasional*: 12.5% vs. 12.1%, p&gt;0.05  Frequent**: 13.6% vs. 16.0%, p&gt;0.05</p> <p><i>Baseline frequent drinkers</i>  None: 17.7% vs. 20.2%, p&gt;0.05  Occasional: 16.1% vs. 13.1%, p&gt;0.05  Frequent: 66.1% vs. 66.7%, p&gt;0.05  * Monthly but not weekly drinking  ** At least weekly drinking</p> <p><u>Any drunkenness in past 30 d</u>  Total sample: OR [3-level adjusted] =0.80 (95% CI: 0.67, 0.97); NNT=26</p> <p><u>Frequent drunkenness in past 30 d</u>  Total sample: OR [3-level adjusted] =0.62 (95% CI: 0.47, 0.81); NNT=40</p>
<p>Notes</p>	<p><b>EU-DAP - European drug abuse prevention trial</b></p> <p>Results of three intervention programs were not presented separately but instead were pooled and compared with control group</p> <p>At 3 mo post-randomisation, the program was shown to be protective in terms of 'any</p>

**Faggiano 2007** (Continued)

drunkenness' in boys aged 13-18 yrs (OR=0.63, 95% CI: 0.48, 0.83), but not in boys aged 11-12 yrs old (OR=0.56, 95% CI: 0.25, 1.28) or girls of any similar age group. There was similar but non-significant trend for 'frequent drunkenness' in boys aged 13-18 yrs (OR=0.69, 95% CI: 0.46, 1.04). At 18 mo post-randomization, neither baseline drinking nor gender seemed to modify the non-significant effect of program in reducing the rate of weekly drinking. At 18 mo post-randomization, the proportion of non-drinking students did not differ in the program and control groups irrespective of the baseline drinking intensity

Abbreviations  
 FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; EU-DAP= European drug abuse prevention trial; ARR=absolute risk difference; NNT=number needed to treat; OR=odds ratio; CI=confidence interval

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	Central randomization
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT analysis; attrition: 10% (3 months post-randomisation)
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	Regression adjusted for covariate

**Furr-Holden 2004**

Methods	Design: RCT FU: 7 yrs (post-randomisation) Attrition: 16% ITT: Yes Unit of randomisation: classroom Clustering effect adjusted: Yes
---------	--

<p>Participants</p>	<p><u>N of Clusters (subjects) randomised</u>            Int 1: NR (NR)            Int 2: NR (NR)            Ctrl: NR (NR)            Total N: 27 (678)</p> <p><u>N of Clusters (subjects) participated</u>            Int 1: NR (192)            Int 2: NR (196)            Ctrl: NR (178)            Total N: 27 (566)            Age: 5.3 - 7.7 yrs (1<sup>st</sup> graders)</p> <p>Sex (male): 50%            Ethnicity: African American (85%-90%), Euro-American (10%-15%)            Ever tried alcohol: NR            Country: US</p>
<p>Interventions</p>	<p><u>Intervention 1: CC intervention</u>  <u>Focus/target:</u> to modify youth's characteristics, conditions and processes in life to achieve more healthy behavior later in life;            reduction of risk of use of alcohol, tobacco, marijuana, and other drugs  <u>Components:</u> 1) curricular enhancements, 2) improved classroom behavior management practice, and            3) supplementary strategies for children not performing adequately</p> <p><u>Intervention 2: FSP intervention</u>  <u>Focus/target:</u> to improve achievement and reduce early aggression and shy behavior by enhancing parent-school communication  <u>Components:</u> 1) training for teachers and other staff, 2) weekly home-school learning and communication activities,            3) 9 workshops for parents led by the 1<sup>st</sup> grade teacher, psychologist or social worker  <u>Fidelity:</u> NR  <u>Duration/frequency:</u> 1 yr  <u>Control:</u> standard curriculum</p>
<p>Outcomes</p>	<p>FU-7 yrs  <u>Alcohol use without parents</u>            Int 1 (34%) vs. Int 2 (37%) vs. Ctrl (33%), NS  <u>Alcohol use without permission</u>            RR (adjusted; Int 1 vs. Ctrl) = 0.95, 95% CI: 0.58, 1.54</p>
<p>Notes</p>	<p><b>JHU-PIRC trial</b>  <b>Good Behavior Game (GBG)</b>            Attrition rates did not significantly differ across the arms and the dropouts were similar to completers with respect to race, academic achievement, or gender.  <u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=</p>

**Furr-Holden 2004** (Continued)

	control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; RR=relative risk; FSP=family school partnership; CC=classroom centred	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Dropouts similar between study arms and similar to completers in race, grades, and gender
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Not described

**Goldberg 2000**

Methods	Design: RCT FU: 1 yr (post-randomisation) Attrition: 20% ITT: yes Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomized</u> Int: 19 (NR) Ctrl: 15 (NR) Total N: 34 (NR)  <u>N of Clusters (subjects) participated</u> Int: 15 (NR) Ctrl: 16 (NR) Total N: 31 (3207) Age: 15-16 yrs (grades 9-12) Sex (male): 100% Ethnicity: NR (in Int group there were more African Americans than in Ctrl group)

Goldberg 2000 (Continued)

	Alcohol users: NR Country: US
Interventions	<u>Intervention</u> : an informational program to raise awareness about anabolic steroid (AS) and other drug dangers <u>Focus/target</u> : Primarily prevent anabolic steroid use but also other substances <u>Components</u> : classroom curriculum addressing physiology and effects of AS - Exercise and weight training; pocket sized guides on diet <u>Fidelity</u> : NR <u>Duration/frequency</u> : 8-14 sessions <u>Control</u> : anti-AS leaflet
Outcomes	<u>Cumulative occurrence (incidents) of drinking and driving (mean score)</u> Baseline: Int: 5.0 (NR) vs. Ctrl: 4.6 (NR) FU-1 yr: Int: 10.7 vs. Ctrl: 12.1 (p<0.08 by school and p=0.004 by individual)
Notes	<b>Adolescents Training and Learning to Avoid Steroids Program</b> There were some baseline differences between the study groups. Three schools in experimental arm withdrew after randomization, so one Ctrl school was randomly reassigned to Int group. <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; AS=anabolic steroids

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT analysis; attrition 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	There were some baseline differences between the study groups

**Goldberg 2007**

Methods	<p>Design: RCT  FU: 2 yrs (post-randomisation)  Attrition: similar across the arms (66% vs. 70%) and the dropouts were similar to completers with respect to baseline variables  ITT: No  Unit of randomisation: school  Clustering effect adjusted: Yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 9 (NR)  Ctrl: 9 (NR)  Total N: 18 (NR)</p> <p><u>N of Clusters (subjects) participated</u>  Int: 5 (653)  Ctrl: 6 (743)  Total N: 11 (1396)  Age: mean range 15.4 - 15.6 yrs (high school athletes)  Sex (male): 54%  Ethnicity: White (91%)  Ever tried alcohol: 21%  Any use of alcohol in prior year: 49%  Country: US</p>
Interventions	<p><u>Intervention</u>: random DAT (100 mL urine analyzed using positive enzymatic immunoassay and gas chromatography-mass spectrometry testing; breath test for alcohol was also administered using approved analyzer)  <u>Focus/target</u>: prevent, identify, and treat substance/drug use  <u>Components</u>: Urine and breath test  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 15 visits biweekly per school year  <u>Control</u>: deferred DAT  <u>Focus/target</u>: NA  <u>Components</u>: NA</p>
Outcomes	<p>FU: 2 yrs  <u>Illicit drug and alcohol use (Index score range: 0 = no use, 1 = light use, 2 = moderate use, and 3 = heavy use)</u>  Past month: Int (0.165) vs. Ctrl (0.261), NS  Past year: Int (0.917) vs. Ctrl (1.033), NS</p>
Notes	<p><b>SATURN</b>  There was some imbalance at baseline between the arms in the mean age, grade, and gender (students in DAT arm being older, having higher grades, and more male)  <u>Abbreviations</u>  FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;</p>



**Goldberg 2007** (Continued)

	wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; AS=anabolic steroids; DAT=drug and alcohol testing	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition 66%-70%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Potential for confounding due to imbalance in covariate

**Goodstadt 1983**

Methods	Design: RCT FU: 6 months (post-randomisation) Attrition: 21.3% ITT: no Unit of randomisation: class Clustering effect adjusted: no
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 4 (111) Int-2: 4 (116) Int-3: 4 (105) Ctrl: 8 (208) Total N: 20 (540)  <u>N of Clusters (subjects) participated</u> Int-1: 4 (111) Int-2: 4 (116) Int-3: 4 (105) Ctrl: 8 (208) Total N: 20 (540)

**Goodstadt 1983** (Continued)

	Age: no details (High School) Sex (male): 41% Ethnicity: NR Alcohol users: NR Country: Canada	
Interventions	<u>Intervention-1</u> : cognitive - detailed examination of alcohol and its role in society <u>Focus/target</u> : reduce alcohol use <u>Intervention-2</u> : Decision- making skills; facts, social influence, behavioural options; <u>Focus/target</u> : reduce alcohol use <u>Intervention-3</u> : reinforcement of values in life <u>Focus/target</u> : reduce alcohol use  Fidelity: NR <u>Duration/frequency</u> : 10 sessions / 10 days <u>Control</u> : standard curriculum with no alcohol education	
Outcomes	FU: 6 months No significant differences between intervention groups and Control for frequency of drinking or usual quantity of drinking The "Values" Group reported significantly more drinking in the previous 6 months than the other programme groups, though no specific scores/details are given	
Notes	<b>"Three Approaches" study</b> Many significant differences found in pre-test measures raising possibility of confounding <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s)	
<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT analysis; 21% attrition
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported

Goodstadt 1983 (Continued)

Other bias	High risk	Potential for confounding due to some baseline differences across study arms
------------	-----------	--

Griffin 2009

Methods	<p>Design: RCT          FU: 1 yr (post-randomisation)          Attrition: 10%          ITT: No          Unit of randomisation: classroom          Clustering effect adjusted: No</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>          Int: 6 (NR)          Ctrl: 6 (NR)          Total N: 12 (199)</p> <p><u>N of Clusters (subjects) participated</u>          Int: 6 (92)          Ctrl: 6 (86)          Total N: 12 (178)          Age: NR (8<sup>th</sup> grade African American students)          Sex (male): 62%          Ethnicity: African American (99%)          Ever tried alcohol: 21.1%-25.0%          Country: US</p>
Interventions	<p><u>Intervention</u>: BRAVE - school-based ATOD and violence prevention program based on social learning theory and resiliency research findings  <u>Focus/target</u>: encouraged the development of resilient behaviours including a sense of purpose and future, autonomy, social competence, and willingness to embrace new experiences  <u>Components</u>: development and monitoring of career goals, mentoring, peer-to-peer goal monitoring and reinforcement, vocational field trips, vocational speakers' bureau, and case referral  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 90 min sessions for 12 mo  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU-12 mo (Int vs. Ctrl)          Alcohol drinking frequency in the past 30 days (0=not at all, 2=once or twice, 3= 1-2 per wk, 4=3-5 x per wk, 5=about once a day, 6=more than once a day)          1.14 (± 0.15) vs. 1.66 (± 0.19), p=0.04 (absolute mean score change between pre- and post-treatment)          7.1 (± 4.8) vs. 37.5 (± 9.9), p=NR (% mean score change between pre- and post-treatment)</p>

**Griffin 2009** (Continued)

	<p>Drunk from alcohol (frequency in the past 30 days: 0 occasions, 1-2 occasions, 3-5 occasions, 6-9 occasions, 10-19 occasions, 20-39 occasions, &gt; 40 occasions)            1.09 (± 0.11) vs. 1.16 (± 0.14), p=0.16 (absolute mean score change between pre- and post-treatment)            3.3 (± 3.3) vs. 8.7 (± 5.9), p=NR (% mean score change between pre- and post-treatment)</p>	
Notes	<p><b>Project BRAVE</b>            non-responders excluded from analyses  <u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;            wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s);            BRAVE=building resiliency and vocational excellence; ATOD=alcohol, tobacco, and other drugs</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT; 10% attrition; the attrition rates did not differ between study arms
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Some baseline differences between study groups

**Hansen 1988**

Methods	Design: RCT FU: 1 and 2 yrs (post-randomisation) Attrition: 37% (year 1), 52% (year 2) ITT: no Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 25 (NR) Int-2: 25 (NR) Ctrl: 36 (NR) Total N: 86 (NR)  <u>N of Clusters (subjects) participated</u> Int-1: 25 (NR) Int-2: 25 (NR) Ctrl: 36 (NR) Total N: 86 (NR)  <u>Analysed sample</u> Int-1: 25 (NR) Int-2: 25 (NR) Ctrl: 36 (NR) Total N: 86 (2863) Age: 12-13 yrs (7 <sup>th</sup> grade) Sex (male): 51% (missing on 95 students) Ethnicity: Hispanic (38.4%), Black (30.5%), White (22.7%), Asians (5.8%), Others (3.5%) Alcohol users: NR Country: US
Interventions	<u>Intervention-1: a social influences curriculum</u> <u>Focus/target:</u> "Gateway drugs" (tobacco, alcohol, marijuana) <u>Components:</u> social motivations, norms, consequences, resistance, role play, adult and media influences  <u>Intervention-2: affective education</u> <u>Focus/target:</u> "Gateway drugs" (tobacco, alcohol, marijuana) <u>Components:</u> motivations, alternatives, goal setting, consequences, self-esteem, decision making, assertiveness <u>Fidelity:</u> NR <u>Duration/frequency:</u> 1 d per wk in 12 sessions <u>Control:</u> standard curriculum
Outcomes	FU: 2 yrs <u>Alcohol use incidence</u> Int-1 vs. Ctrl: F=0.93, p=0.3 Int-2 vs. Ctrl: F=2.14, p=0.2  <u>Alcohol use prevalence</u>

Hansen 1988 (Continued)

	<p>Int-1 vs. Ctrl: F=3.57, p=0.06          Int-2 vs. Ctrl: F=13.18, p=0.0003*          *Int-2 drinking alcohol more vs. Ctrl</p> <p><u>Increased use (in baseline users) %</u>          Int-1 (22.6%) vs. Int-2 (48.0%) vs. Ctrl (37.7%), p &lt; 0.05 (Int-1 vs. Ctrl or Int-2)*          *Int-2 more increase in alcohol use vs. Ctrl or Int-1</p> <p><u>Reduced use (in baseline users) %</u>          Int-1 (43.8%) vs. Int-2 (69.8%) vs. Ctrl (63.3%), p &lt; 0.05 (Int-1 vs. Ctrl or Int-2)*          *Int-1 less reduced alcohol use vs. Ctrl or Int-2</p>	
Notes	<p><b>Project SMART</b>          Attrition rate was differential by race, Whites less likely to drop than Blacks. Attrition rate was differential by the study group at follow-ups with higher attrition in Ctrl than Int-2. There were pre-test differences between groups in drinking measures, the Ctrl group reporting more drinking (Int-1 vs. Ctrl).</p> <p><u>Abbreviations</u>          FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s)</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition > 30% (differential)
Selective reporting (reporting bias)	High risk	30-day, 7-day customary and lifetime use not reported in results section
Other bias	High risk	pre-test differences between groups in drinking measures,

## Hansen 1991

Methods	<p>Design: RCT  FU: 2 yrs (post-randomisation)  Attrition: 22% (year 1), 46% (year 2)  ITT: no  Unit of randomisation: school (stratified by size, ability and ethnicity)  Clustering effect adjusted: only at 2-year FU</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int-1: NR (NR)  Int-2 NR (NR)  Int-3 NR (NR)  Ctrl: NR (NR)  Total N: 12 (3011)</p> <p><u>N of Clusters (subjects) participated</u>  Int-1: NR (NR)  Int-2 NR (NR)  Int-3 NR (NR)  Ctrl: NR (NR)</p> <p><u>Analysed sample (year 1)</u>  Int-1: NR (NR)  Int-2 NR (NR)  Int-3 NR (NR)  Ctrl: NR (NR)  Total N: 12 (2416)  Age: 12-13 yrs  Sex (male): 45-51%  Ethnicity: White (33.2%-52.2%), Hispanic (11.3%-42.6%), Black (0.8%-3.0%), and Asian (9.8%-26.0%)  Alcohol users: 29%  Country: US</p>
Interventions	<p><u>Intervention-1: RT</u>  <u>Components:</u> negative/positive consequences of drinking using Socratic method, definition of drugs/alcohol, techniques/practice to refuse, assertiveness, role played refusals, film discussion, parent interview homework, types of social pressure</p> <p><u>Intervention-2: NE</u>  <u>Components:</u> negative/positive consequences of drinking using Socratic method, definition of drugs/alcohol, parent interview homework, review of consequences of drinking, survey about prevalence of alcohol use in students, survey results, agree or disagree opinion, class discussions, non-drinker interview homework</p> <p><u>Intervention-3: RT + NE</u>  <u>Components:</u> see above</p>

Hansen 1991 (Continued)

	<p><u>Focus/target</u>: develop social and life skills, including social norms  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 4-10 sessions x 45 min each  <u>Control</u>: ICU</p>	
Outcomes	<p>FU: 1 yr  <u>Ever used</u>: RT: F=3.22; NE: F=8.17**  NE: 11.0% increase (pre-post)  Non-NE: 14.2% increase (pre-post)  <u>30-day use</u>: RT: F=0.58; NE: F=5.10*  NE: 6.2% increase (pre-post)  Non-NE: 9.1% increase (pre-post)  <u>7-day use</u>: RT: F=0.88; NE: F=5.93*  NE: 2.8% increase (pre-post)  Non-NE: 5.1% increase (pre-post)  <u>Drunkenness</u>: RT: F=0.03; NE: F=25.19***  NE: 4.2% increase (pre-post)  Non-NE: 11.1% increase (pre-post)  <u>Problem use</u>: RT: F=0.11; NE: F=4.26*  NE: 0.3% increase (pre-post)  Non-NE: 2.4% increase (pre-post)</p> <p>* p&lt;0.05; ** p&lt;0.01; *** p&lt;0.0001</p> <p>FU: 2 yrs  Palmer et al (2000) re-analysed data taking into account unit (individual, class, school) effects and found no significant effects at any of these levels using the ICU group as the Ctrl</p>	
Notes	<p><b>Adolescent Alcohol Prevention Trial (AAPT)</b>  Some differences between groups at baseline in ethnic mix. Year 1 analysis mixed Ctrl and non-NE classes from other groups therefore muddying the comparison.  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); RT=resistance training; NE=normative education; ICU=information on consequences of drug use</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described



**Hansen 1991** (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition > 21%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Some differences between groups at baseline in ethnic mix. Year 1 analysis mixed Ctrl and non-NE classes from other groups therefore muddying the comparison.

**Hecht 2003**

Methods	Design: RCT FU: 3 yrs (post-randomisation) Attrition: 12.5% (post-randomisation), 16% (of 6035) at last follow-up ITT: Unclear Unit of randomisation: school Clustering effect adjusted: Yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: 25 (NR) Ctrl: 10 (NR) Total N: 35 (6900)  <u>N of Clusters (subjects) participated</u> Int: 25 (NR) Ctrl: 10 (NR) Total N: 35 (6035) Age: 12.5 [11-18] yrs (7th grade students) Sex (male): 53% Ethnicity: Hispanic (74%), White (17.4%), African American (8.6%) Alcohol ever use (30 days prior to survey): 22.6% (1364/6035) Country: US
Interventions	<u>Intervention</u> : REAL curriculum given in 3 versions (Mexican American, Black/White, Multicultural) <u>Focus/target</u> : to promote anti drug norms and teaching resistance/social skills reinforced by booster activities and media campaign <u>Components</u> : 10 sessions and booster activities <u>Fidelity</u> : mean score of 5.8 (appropriateness in conveying the curriculum ranging from 1 = inappropriately to 7 = appropriately) <u>Duration/frequency</u> : 2 yrs 1 booster activity per mo

Hecht 2003 (Continued)

	Control: standard curriculum
Outcomes	<p><u>FU- 1 yr (post-baseline)</u>  Mean # of alcohol drinks (range 1= none to 9 = more than 30): mean difference -0.148 (Int vs. Ctrl), p&lt;0.05  Multicultural and Mexican American version intervention groups had smaller mean increase in alcohol use than Black/White version group</p> <p><u>FU- 2 yr (post-baseline)</u>  Mean # of alcohol drinks (range 1= none to 9 = more than 30): mean difference -0.144 (Int vs. Ctrl), p&lt;0.01  No differences between the 3 cultural versions of curriculum in alcohol use change</p> <p><u>FU- 3 yr (post-baseline)</u>  Mean # of alcohol drinks (range 1= none to 9 = more than 30): mean difference -0.232 (Int vs. Ctrl), p&lt;0.001  Multicultural and Mexican American version intervention groups had smaller mean increase in alcohol use than Black/White version group</p>

Notes	<p><b><i>Keepin' it R.E.A.L.</i></b>  Imputation of missing data was performed; there were no gender differences in program effects on alcohol use (Kulis 2007a); amongst alcohol users at baseline, the discontinuation rate in the intervention group was 61% greater than in the control group (Kulis 2007b); students receiving multicultural version intervention had less overall alcohol use compared to the controls</p> <p><u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); REAL=refuse, explain, avoid, and leave</p>
-------	---

<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear if ITT was done;

**Hecht 2003** (Continued)

Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	baseline covariate distributions not reported; analysis adjusted for covariate

**Johnson 2009**

Methods	<p>Design: RCT  FU: 6 mo (post-randomisation)  Attrition: 29%; dropouts and participants did not differ in alcohol use and other characteristics  ITT: No  Unit of randomisation: community  Clustering effect adjusted: Yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: NR (317)  Ctrl: NR (289)  Total N: 14 (606)</p> <p><u>N of Clusters (subjects) participated</u>  Int: NR (238)  Ctrl: NR (222)  Total N: 14 (460)  Age: NR (5<sup>th</sup> and 6<sup>th</sup> grades Alaskan students)  Sex (male): 50%  Ethnicity: Caucasians (NR) and Native Americans (NR)  Alcohol use: NR  Country: US</p>
Interventions	<p><u>Intervention:</u> <i>Think Smart</i> is a modified form of Personal Intervention Curriculum by Schinke  <u>Focus/target:</u> designed to reduce use of inhalants, over the counter drugs, alcohol, tobacco, other drugs through teaching refusal skills, peer normative beliefs, drug-related consequences, assertiveness skills, and cultural identity  <u>Components:</u> 12 sessions (stereotypes-peer norms, cultural identity, drug facts, refusal and self-assertiveness skills) and 3 booster sessions (practicing problem-solving model)  <u>Fidelity:</u> NR  <u>Duration/frequency:</u> 12 core sessions in 5 mo and booster sessions 2-3 mo later  <u>Control:</u> standard curriculum</p>
Outcomes	<p>FU-6 mo  Use of alcohol in the past 30 days: OR [Int vs. Ctrl] = 0.39, p=NR (NS)</p>

**Johnson 2009** (Continued)

Notes	<p><b>Think Smart</b>            Baseline differences were adjusted for covariate</p> <p><u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; OR=odds ratio</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT; attrition > 20%; dropouts similar across study arms and to completers; missing data imputation
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	Baseline differences were adjusted for covariate

**Kellam 2008**

Methods	Design: RCT FU: 12 yrs (post-randomisation) Attrition: NR ITT: Yes Unit of randomisation: classrooms and teachers Clustering effect adjusted: Yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: NR (NR) Ctrl: NR (NR) Total N: NR (NR)  <u>N of Clusters (subjects) participated</u> Int: 8 (238) Ctrl: 6 (169)

**Kellam 2008** (Continued)

	<p>Total N: 14 (407)</p> <p>Additional set of 515 external controls (18 classes) were employed for certain analyses            Age: 6-7 yrs (1<sup>st</sup> and 2<sup>nd</sup> grade students)            Sex (male): 50%            Ethnicity: NR            Alcohol use: NR            Country: US</p>	
Interventions	<p><u>Intervention</u>: GBG a method of classroom behavior management  <u>Focus/target</u>: to create a classroom environment that is conducive to learning by focusing on the social context of the classroom;            to socialize children into the role of student and to teach them how to regulate their own and classmates behavior through team behavior contingent reinforcement; to reduce early aggressive, disruptive behavior (at classroom and individual level) - antecedents of later substance use, and dependence disorders  <u>Components</u>: 1) students were assigned to one of 3 heterogeneous teams that contained equal numbers of boys and girls, equal numbers of aggressive, shy, and socially isolated children given the baseline measurements; 2) teacher posted basic classroom behavior rules and teams were rewarded if they committed 4 or fewer infractions of these rules  <u>Fidelity</u>: not measured  <u>Duration/frequency</u>: 10 min - 3 hrs long sessions 3x per wk for 2 yrs  <u>Control</u>: standard curriculum</p>	
Outcomes	<p>FU-12 yrs  <u>Lifetime alcohol abuse and dependence disorders</u>:            Int (13%) vs. Ctrl (20%), p=0.08 (Unadjusted for baseline differences and based on 169 internal controls)            Int (13%) vs. Ctrl (29%), p=0.03 (Unadjusted for baseline differences and based on all 922 controls, including 515 external controls)            Adjusted OR [Int vs. Ctrl] = 0.5, p=0.05 (Individual-level risk factors and classroom variation considered)</p>	
Notes	<p><b>JHU-PIRC trial</b>  <b>Good Behavior Game (GBG)</b>            Greater effect of GBG on lifetime alcohol abuse and dependence disorder was found in males compared to females  <u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;            wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); OR=odds ratio; GBG=good behavior game</p>	
<i>Risk of bias</i>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>

**Kellam 2008** (Continued)

Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	ITT done but attrition not reported
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	baseline differences between study arms not significant; results adjusted for residual confounding

**Koning 2009**

Methods	<p>Design: RCT  FU: 22 mo (post-randomisation)  Attrition: 12.5% (2570/2937); dropouts were older, drank more, and had parents with lower education levels than completers  ITT: No  Unit of randomisation: school  Clustering effect adjusted: Yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int 1: NR (801)  Int 2: NR (942)  Int 3: NR (812)  Ctrl: NR (935)  Total N: 20 (3490)</p> <p><u>N of Clusters (subjects) participated</u>  Int 1: NR (608)  Int 2: NR (675)  Int 3: NR (588)  Ctrl: NR (699)  Total N: 19 (2570)  Age: 12.7 yrs (1<sup>st</sup> and 2<sup>nd</sup> yr high school students)  Sex (male): 51%  Ethnicity: NR  Alcohol use: not heavy drinkers  Country: The Netherlands</p>

Interventions	<p><u>Intervention 1: PI</u>  <u>Focus/target:</u> parental rules for their children's alcohol use  <u>Components:</u> 1) 20 min presentation about adverse effects of alcohol use at young age; 2) parents meet with the class mentor to discuss rules and reach a consensus; 3) Information leaflet with summary information sent to parents' home addresses as reminder of rules and consensus reached</p> <p><u>Intervention 2: SI</u>  <u>Focus/target:</u> based on HSD prevention program  <u>Components:</u> 1) coordinating committee; 2) 3 series of educational lessons about tobacco, alcohol, cannabis, ecstasy, games; 3) school regulations on drug use; 4) system of detection of drug problems; and 5) parental involvement</p> <p><u>Intervention 3: [PI + SI] combined</u>  <u>Fidelity:</u> NR  <u>Duration/frequency:</u> 2 mo</p> <p><u>Control:</u> standard curriculum</p>
Outcomes	<p>FU-10 mo  <u>Heavy weekly drinking</u>  Int 1 (3.5%) vs. Int 2 (3.4%) vs. Int 3 (1.2%) vs. Ctrl (3.2%), P&lt;0.05 (Int 3 vs. Ctrl)  <u>Weekly drinking</u>  Int 1 (12.6%) vs. Int 2 (16.1%) vs. Int 3 (11.8%) vs. Ctrl (16.6%), P&lt;0.05 (Int 3 vs. Ctrl)  <u>Frequency of drinking</u>  Students in Int 3 (combined intervention: PI + SI) drank significantly less frequently than students in the Ctrl arm</p> <p>FU-22 mo  <u>Heavy weekly drinking</u>  Int 1 (10.5%) vs. Int 2 (8.2%) vs. Int 3 (7.6%) vs. Ctrl (9.9%), NS  - OR=1.13, 95% CI: 0.73, 1.73 (adjusted; Int 1 vs. Ctrl), NNT = 48.9  - OR=0.85, 95% CI: 0.56, 1.29 (adjusted; Int 2 vs. Ctrl), NNT = 84.4  - OR=0.80, 95% CI: 0.48, 1.32 (adjusted; Int 3 vs. Ctrl), NNT = 58.7  <u>Weekly drinking</u>  Int 1 (33.2%) vs. Int 2 (36.1%) vs. Int 3 (31.5%) vs. Ctrl (41.5%), NS  - OR=0.86, 95% CI: 0.63, 1.16 (adjusted; Int 1 vs. Ctrl), NNT = 181.8  - OR=0.92, 95% CI: 0.71, 1.19 (adjusted; Int 2 vs. Ctrl), NNT = 67.9  - OR=0.71, 95% CI: 0.53, 0.94 (adjusted; Int 3 vs. Ctrl), NNT = 17.2  <u>Frequency of drinking</u>  Students in Int 3 (combined intervention: PI + SI) drank significantly less frequently than students in the Ctrl arm</p>
Notes	<p>Even though the authors reported to have used ITT analysis (based on 2937 students) with imputations, they had already excluded 431 (baseline drinkers) + 122 (refusals/not present) students from the initial</p>

**Koning 2009** (Continued)

sample of 3490 students  
Abbreviations  
 FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  
 wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s);  
 NS=statistically non-significant;  
 PI=parent intervention; SI=student intervention; HSD=healthy school and drugs;  
 NNT=number needed to treat; OR=odds ratio;  
 95% CI=ninety-five percent confidence interval

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	Central randomisation
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	ITT was done but not on the original sample
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	Baseline differences adjusted

**McBride 2000**

Methods	Design: RCT FU: 32 mo (post-randomisation) Attrition: 24.1% (at 32 mo) ITT: No Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: NR (NR) Ctrl: NR (NR) Total N: NR (NR)  <u>N of Clusters (subjects) participated</u> Int: NR (1111) Ctrl: NR (1232)



	<p>Total N: 14 (2343)</p> <p><u>N of Clusters (subjects) analysed (yr 1)</u>            Int: NR (855)            Ctrl: NR (872)            Total N: 14 (1727)</p> <p><u>N of Clusters (subjects) analysed (yr 2)</u>            Int: NR (970)            Ctrl: NR (1037)            Total N: 14 (2007)</p> <p><u>N of Clusters (subjects) analyzed (yr 3)</u>            Int: NR (863)            Ctrl: NR (915)            Total N: 14 (1778)            Age: 13 yrs (8<sup>th</sup> grade)            Sex (male): NR            Ethnicity: NR            Alcohol users: NR</p> <p><u>No. of standard drinks in last 12 mo (mean and 95% CI)</u>            Int: 30.2 (11, 49) vs. Ctrl: 19.7 (12, 27)</p> <p><u>Hazardous or harmful drinking once a month or more often (% and 95% CI)</u>            Int: 11.3 (8.8,14.3) vs.            Ctrl: 13.3 (10.5,16.8)</p> <p><u>Own harm index (mean and 95%CI)</u>            Int: 2.0 (1.5, 2.5) vs. Ctrl: 1.5 (1.1, 1.9)</p> <p><u>Country: Australia</u></p>
Interventions	<p><u>Intervention: SHAHRP</u>  <u>Focus/target:</u> to reduce alcohol-related harm  <u>Components:</u> 2 phases, phase 1 included 17 skill-based activities in 10 lessons; phase 2 included 12 activities over 5-7 wks and trigger video; these activities were skill rehearsal, individual or group decision making, discussions around scenarios suggested by students with identification of alcohol related harm and strategies to reduce the harm</p> <p><u>Fidelity:</u> 80.7%</p> <p><u>Duration/frequency:</u> 24 mo; see also above</p> <p><u>Control:</u> standard curriculum</p> <p><u>Duration:</u> 10 wks</p>
Outcomes	<p><u>FU: 8 mo (Int vs. Ctrl)</u>            Non-users of alcohol: 13.1% vs. 10.1%, p&gt;0.05            Users of alcohol (at least once per wk): 15.0% vs. 18.9%, p&gt;0.05            Drinking alone: 23.0% vs. 25.0%, p&gt;0.05            Supervised drinker: 33.0% vs. 30.0%, p&gt;0.05            Unsupervised drinker: 42.5% vs. 46.0%, p&gt;0.05</p> <p><u>FU: 20 mo (Int vs. Ctrl)</u></p>

	<p>Alcohol consumption 31.7% less in Int group  Risky levels of alcohol consumption 33.8% less in Int group  Harm related to alcohol own use 16.7% less in Int group  % of non-drinkers: 16.3 (95% CI: 13.9, 19.0) vs. 19.0 (95% CI: 16.5, 21.7)  Risky drinking at least once a month: 21.5 (95% CI: 18.0, 25.0) vs. 32.5 (95% CI: 28.2, 36.9)</p> <p><u>FU: 32 mo (Int vs. Ctrl)</u>  Alcohol consumption 9.2% less in Int group  Risky levels of alcohol consumption 4.2% less in Int group  Harm related to alcohol own use: 22.9% less in Int group  % of non-drinkers: 27.3 (95% CI: 24.4, 30.4) vs. 28.3 (95% CI: 25.4, 31.4)  Risky drinking at least once a mo: 32.2 (95% CI: 28.2, 36.2) vs. 33.9 (95% CI: 29.7, 39.8)  No. of standard drinks in last 12 mo (mean and 95% CI): Int: 273.8 (217, 330) vs. Ctrl: 362.7 (283, 443)  Hazardous or harmful drinking once a month or more often (% and 95% CI): Int: 32.2 (28.2, 36.2) vs. Ctrl: 33.9 (29.7, 39.8)  Own harm index (mean and 95% CI): Int: 9.8 (8.3, 11.3) vs. Ctrl: 12.5 (10.5, 14.4)</p>	
Notes	<p><b>SHAHRP</b>  Results were stratified by baseline use of alcohol and there was no intervention effect in baseline supervised drinkers compared to controls; but in baseline non-drinkers and unsupervised drinkers, risky alcohol consumption was significantly reduced in the Int group vs. Ctrl group. One school refused to be randomised so was assigned to control group; authors assumed randomization after sensitivity analysis. Baseline differences between the intervention and control groups were statistically significant for both context of use and harms associated with their own use of alcohol</p> <p><u>Abbreviations</u>  FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; NNT=number needed to treat; 95% CI=ninety-five percent confidence interval; SHAHRP=the school health and alcohol harm reduction project</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described

McBride 2000 (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Baseline distributions of covariate not reported

Morgenstern 2009

Methods	<p>Design: RCT  FU: 12 mo (post-randomisation)  Attrition: 23.6% (12 mo); dropout rates did not differ across the assigned conditions  ITT: yes  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 16 (911)  Ctrl: 14 (964)  Total N: 30 (1875)</p> <p><u>N of Clusters (subjects) participated</u>  Int: 16 (714)  Ctrl: 14 (719)  Total N: 30 (1433)  Age: 13 [12-15] yrs (7<sup>th</sup> grade)  Sex (male): 52%  Ethnicity: NR  Alcohol use (past mo; range 0 - 6 d): Int (0.62 ± 1.34) vs. Ctrl (0.63 ± 1.35), p = 0.79  LTA-WPK: Int (34.5%) vs. Ctrl (35.8%), p=0.56  Life-time drunkenness: Int (18.5%) vs. Ctrl (18.3), p=0.24  Life-time binge-drinking: Int (12%) vs. Ctrl (13.1%), p=0.50  Country: Germany</p>
Interventions	<p><u>Intervention</u>: alcohol education intervention  <u>Focus/target</u>: to reduce/prevent alcohol use with a main message 'no alcohol for minors'  <u>Components</u>:  1) 4 specified interactive class units (<i>what is permitted? advertisement, temptations, when is drinking alcohol OK?</i>)  2) booklet for students (knowledge about alcohol and consequences of its use)  3) booklet for parents (behavioral advices)  <u>Fidelity</u>: NR</p>

	<p><u>Duration/frequency</u>: 4 lessons in 3 mo</p> <p><u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 4 mo (Int vs. Ctrl)</p> <p>Alcohol use (past mo; range 0 - 6 d): 0.81 (<math>\pm</math> 0.046) vs. 0.89 (<math>\pm</math> 0.044), p=0.178</p> <p>Alcohol use (life-time): OR [adjusted]=0.81, 95% CI: 0.57, 1.16</p> <p>Life-time drunkenness: OR [adjusted]=0.70, 95% CI: 0.48, 1.02</p> <p>Life-time binge-drinking: OR [adjusted]=0.56, 95% CI: 0.41, 0.77</p> <p>FU: 12 mo (Int vs. Ctrl)</p> <p>Alcohol use (past mo; range 0 - 6 d): 0.89 (<math>\pm</math> 0.075) vs. 0.98 (<math>\pm</math> 0.081), p=0.419</p> <p>Alcohol use (life-time): OR [adjusted]=0.90, 95% CI: 0.67, 1.21</p> <p>Life-time drunkenness: OR [adjusted]=0.77, 95% CI: 0.52, 1.12</p> <p>Life-time binge-drinking: OR [adjusted]=0.74, 95% CI: 0.57, 0.97</p>
Notes	<p>Data imputation for missing values was performed. There were more current and lifetime smokers at baseline in Int (50.9% and 15.6%, respectively) vs. Ctrl (45.2% and 11.3%, respectively) groups (p=0.01)</p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); OR=odds ratio; 95% CI=ninety-five percent confidence interval</p>

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	Allocating person was blinded to intervention and research question
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT and data imputation done; attrition < 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Baseline imbalance by smokers

**Moskowitz 1984**

Methods	Design: RCT FU: 1 yr (post-randomisation) Attrition: 26% (year 1) ITT: no Unit of randomisation: class (stratified by attitudes and involvement) Clustering effect adjusted: yes	
Participants	<u>N of Clusters (subjects) randomised</u> Int: 8 (NR) Ctrl: 8 (NR) Total N: 16 (473)  <u>N of Clusters (subjects) participated</u> Int: 8 (NR) Ctrl: 8 (NR) Total N: 16 (473) Age: 12 yrs Sex (male): NR Ethnicity: Predominantly White Alcohol users: NR Country: US	
Interventions	<u>Intervention</u> : a drug education course incorporating three approaches to behavior change <u>Focus/target</u> : prevent, delay, reduce drug use including alcohol <u>Components</u> : models of motivation and decision making, advertising influence, as- sertiveness training/role play, knowledge of drugs <u>Fidelity</u> : NR <u>Duration/frequency</u> : 12 sessions (45 min each) per 12 weeks <u>Control</u> : standard curriculum	
Outcomes	FU: 1 yr No significant effects of intervention group over controls for alcohol involvement at 1 year follow-up ( $F < 1$ for males and females)	
Notes	<b>Drug Education Course</b> <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl= control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo= month(s)	
<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described

**Moskowitz 1984** (Continued)

Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	no ITT; attrition: 26%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Possible contamination

**Newman 1992**

Methods	Design: RCT FU: 1 yr (post-randomisation) Attrition: NR ITT: unclear Unit of randomisation: school Clustering effect adjusted: by class
Participants	<u>N of Clusters (subjects) randomised</u> Int: 5 (NR) Ctrl: 4 (NR) Total N: 9 (c.3500)  <u>N of Clusters (subjects) participated</u> Int: 5 (NR) Ctrl: 4 (NR) Total N: 9 (c.3500) Age: 15-16 yrs Sex (male): NR Ethnicity: NR Alcohol users: NR Country: US
Interventions	<u>Intervention</u> : study of alcohol behavior changes following a video-based alcohol education program <u>Focus/target</u> : enhancing refusal skills <u>Components</u> : video showed typical adolescent drinking situations, followed by role play; information about alcohol <u>Fidelity</u> : NR <u>Duration/frequency</u> : 10 lessons incorporating 5 videos (20 min each) <u>Control</u> : standard alcohol education curriculum
Outcomes	FU: 1 yr (Int vs. Ctrl) <u>Consumed 1+ drinks</u> Pretest: 64.9% vs. 68.5%

Newman 1992 (Continued)

	FU: 79.2% vs. 81.3% <u>Mean N drinks at last party</u> Pretest: 1.64 vs. 1.9 FU: 2.46 vs. 2.63 <u>Mean N of drinking events (last mo)</u> Pretest: 1.64 vs. 1.88 FU: 3.06 vs. 3.43 No significant effects of the intervention	
Notes	<b>Resisting Pressure to Drink and Drive (RPDD)</b> Random allocation by school but class was the unit of analysis <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); RPDD=resisting pressure to drink and drive	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	unclear if ITT was done; attrition rate: NR
Selective reporting (reporting bias)	Unclear risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Random allocation by school but class was the unit of analysis

**Newton 2009a**

Methods	<p>Design: RCT  FU: 1.5 yr (post-randomisation)  Attrition: 51.3% (666/1296) (1 yr); dropout rates did not differ across the assigned conditions  ITT: no  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 5 (728)  Ctrl: 5 (568)  Total N: 10 (1296)</p> <p><u>N of Clusters (subjects) participated</u>  Int: 5 (343)  Ctrl: 5 (287)  Total N: 10 (630)</p> <p>Analyzed sample: N = 764 (Int: 397 vs. Ctrl: 367)  Age: 13 yrs  Sex (male): 60%  Ethnicity: NR  Alcohol use: NR  Country: Australia</p>
Interventions	<p><u>Intervention</u>: alcohol and cannabis prevention course  <u>Focus/target</u>: to decrease alcohol and cannabis misuse  <u>Components</u>: 2 sets of six 40 min lessons including alcohol and cannabis modules; each lesson included 15-20 min internet based session completed individually; students also viewed a cartoon; 2<sup>nd</sup> part of each lesson was a predetermined activity to reinforce the information learnt in the cartoons  <u>Fidelity</u>: 91%  <u>Duration/frequency</u>: 6 mo  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 1 yr (Int vs. Ctrl)  <u>Average weekly alcohol use</u>  Mean difference (from baseline): <math>-0.88 \pm 0.91</math> vs. <math>2.67 \pm 1.09</math>, <math>p &lt; 0.05</math>  <u>Frequency of drinking to excess</u>  Mean difference (from baseline): <math>0.32 \pm 0.18</math> vs. <math>0.23 \pm 0.11</math>, <math>p = 0.69</math>  <u>Harms related to own use of alcohol</u>  Mean difference (from baseline): <math>0.34 \pm 1.63</math> vs. <math>2.73 \pm 1.47</math>, <math>p = \text{NR (NS)}</math></p> <p>FU: 1.5 yrs (Int vs. Ctrl)  <u>Average weekly alcohol use</u>  Mean difference (from baseline): <math>-0.63 \pm 1.14</math> vs. <math>5.30 \pm 1.50</math>, <math>p &lt; 0.02</math>  <u>Frequency of drinking to excess</u>  Mean difference (from baseline): <math>0.05 \pm 0.16</math> vs. <math>0.85 \pm 0.30</math>, <math>p &gt; 0.05</math>  <u>Harms related to own use of alcohol</u>  Mean difference (from baseline): <math>3.06 \pm 2.12</math> vs. <math>9.17 \pm 2.23</math>, <math>p &gt; 0.05</math></p>



**Newton 2009a** (Continued)

Notes	<p><b>CLIMATE Schools</b>            There were fewer males (54% vs. 65%, <math>p &lt; 0.05</math>), higher weekly alcohol use (<math>p &lt; 0.05</math>), higher frequency of bingeing (<math>p &lt; 0.01</math>) than in Ctrl group at baseline.</p> <p><u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant</p>
-------	---

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	using online randomisation system
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT; attrition > 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Imbalance in baseline covariate

**Perry 1988**

Methods	<p>Design: RCT            FU: 8 wks post-randomisation            Attrition: 7-8%            ITT: no            Unit of randomisation: school            Clustering effect adjusted: no</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>            Int-1: NR (NR)            Int-2: NR (NR)            Ctrl: NR (NR)            Total N: 25 (2536)</p> <p><u>N of Clusters (subjects) participated</u>            Int-1: NR (NR)</p>

**Perry 1988** (Continued)

	<p>Int-2: NR (NR)  Ctrl: NR (NR)  Total N: 25 (2536)  Age: 11-18 yrs  Sex (male): 50%  Ethnicity: NR  Alcohol users: 56% used in last 12 mo  Country: Australia, Chile, Norway and Swaziland</p>
Interventions	<p><u>Intervention-1</u>: Peer-led  <u>Focus/target</u>: social and normative influences to drink and resistance skills  <u>Intervention-2</u>: Teacher-led  <u>Focus/target</u>: social and normative influences to drink and resistance skills  <u>Components (Int1-2)</u>: School-based alcohol education developed from early LST - normative expectancies; peer influences; consequences of alcohol use; understanding mass media influences. Booster sessions to reinforce abstinence  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 5 sessions (4 weeks) + booster over 2 mo  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 8 wks  <u>Baseline non-drinkers (total) - Post-test composite alcohol score (Mean and SE)</u>  Int-1: 3.15 (±0.12) vs. Int-2: 3.46 (±0.12) vs. Ctrl: 3.52 (±0.16), p&lt;0.005 (Int-1 vs. Int-2 or Ctrl; ANCOVA)</p> <p><u>Baseline drinkers (total) - Post-test composite alcohol score (Mean and SE)</u>  Int-1: 5.14 (±0.35) vs. Int-2: 5.84 (±0.27) vs. Ctrl: 5.71(±0.37)</p> <p><u>Baseline non-drinkers (males) - Post-test composite alcohol score (Mean and SE)</u>  Int-1: 3.19 (±0.18) vs. Int-2: 3.48 (±0.17) vs. Ctrl: 3.53 (±0.25), p&lt;0.005 (Int-1 vs. Int-2 or Ctrl; ANCOVA)</p> <p><u>Baseline non-drinkers: (females) - Post-test composite alcohol score (Mean and SE)</u>  Int-1: 3.11 (±0.14) vs. Int-2: 3.45 (±0.15) vs. Ctrl: 3.44 (±0.20), p&lt;0.005 (Int-1 vs. Int-2 or Ctrl; ANCOVA)</p> <p><u>Baseline drinkers (males) - Post-test composite alcohol score (Mean and SE)</u>  Int-1: 5.28 (±0.53) vs. Int-2: 5.81 (±0.44) vs. Ctrl: 5.74 (±0.50)  <u>Baseline drinkers (females) - Post-test composite alcohol score (Mean and SE)</u>  Int-1: 5.01 (±0.45) vs. Int-2: 5.79 (±0.37) vs. Ctrl: 5.81 (±0.54), p&lt;0.05 (Int-1 vs. Int-2 or Ctrl; ANCOVA)</p>
Notes	<p><b>WHO Collaborative Study</b>  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant;</p>

**Perry 1988** (Continued)

LST=life skills training; ANCOVA=analysis of covariance; SE=standard error		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear because no ITT but low attrition of 8%
Selective reporting (reporting bias)	High risk	12-month follow-up outcomes not reported
Other bias	Unclear risk	baseline covariate distribution not reported

**Perry 2003**

Methods	Design: RCT FU: 20 mo (post-randomisation) Attrition: 16%; dropout rates did not differ across the study conditions ITT: Yes Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 8 (2518) Int-2: 8 (2635) Ctrl: 8 (2108) Total N: 24 (7261)  <u>N of Clusters (subjects) participated</u> Int-1: 8 (2226) Int-2: 8 (2221) Ctrl: 8 (1790) Total N: 24 (6237) Analyzed sample: N = 7261 Age: NR yrs (7 <sup>th</sup> or 8 <sup>th</sup> grade) Sex (male): 51.6% Ethnicity: White (67.3%), African American (7.5%), Asian American (12.7%), Hispanic (3.6%), American Indian (4%),

Perry 2003 (Continued)

	<p>other/mixed race (4.9%)          Alcohol use: NR          Country: US</p>	
Interventions	<p><u>Intervention 1: DARE</u>  <u>Focus/target:</u> to provide skills in resisting influences to use drugs and handling violent situations, character building and citizenship skills  <u>Components:</u> 10 sessions in 2 semesters  <u>Intervention 2: DARE +</u>  <u>Focus/target:</u> see above  <u>Components:</u> 'on the VERGE' - 4-session program implemented by trained teachers once a week for 4 weeks and extracurricular activities  <u>Fidelity:</u> NR  <u>Duration/frequency:</u> 18 mo          Control: no Intervention</p>	
Outcomes	<p>FU: 20 mo (Int-1 vs. Int-2 vs. Ctrl)  <u>Boys</u>          Alcohol use growth rate (past yr): 0.21 (<math>\pm 0.03</math>) vs. 0.19 (<math>\pm 0.03</math>) vs. 0.26 (<math>\pm 0.03</math>), <math>p=0.04</math> (Int-2 vs. Ctrl), other pair-wise comparisons NS          Alcohol use growth rate (past month): 0.11 (<math>\pm 0.02</math>) vs. 0.08 (<math>\pm 0.02</math>) vs. 0.14 (<math>\pm 0.02</math>), <math>p=0.01</math> (Int-2 vs. Ctrl), other pair-wise comparisons NS          Ever drunk growth rate: 0.11 (<math>\pm 0.02</math>) vs. 0.11 (<math>\pm 0.02</math>) vs. 0.15 (<math>\pm 0.02</math>), <math>p=0.07</math> (Int-1 or Int-2 vs. Ctrl), other pair-wise comparisons also NS    <u>Girls</u>          Alcohol use growth rate (past yr): 0.27 (<math>\pm 0.04</math>) vs. 0.23 (<math>\pm 0.04</math>) vs. 0.25 (<math>\pm 0.04</math>), all pair-wise comparisons NS (<math>p&gt;0.05</math>)          Alcohol use growth rate (past month): 0.13 (<math>\pm 0.02</math>) vs. 0.08 (<math>\pm 0.03</math>) vs. 0.12 (<math>\pm 0.03</math>), all pair-wise comparisons NS (<math>p&gt;0.05</math>)          Ever drunk growth rate: 0.13 (<math>\pm 0.02</math>) vs. 0.07 (<math>\pm 0.02</math>) vs. 0.12 (<math>\pm 0.02</math>), <math>p=0.04</math> (Int-1 vs. Int-2), other pair-wise comparisons NS</p>	
Notes	<p><b>DARE</b>          No significant differences across the conditions at baseline. Ctrl group received delayed DARE + program (started after final FU at 20 mo post-randomisation).  <u>Abbreviations</u>          FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); SE=standard error</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described

**Perry 2003** (Continued)

Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT done; attrition < 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	no baseline imbalance

**Reddy 2002**

Methods	Design: RCT FU: 17 mo (post-randomisation) Attrition: 7%; dropout rates did not differ across the study conditions ITT: No Unit of randomisation: school Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 10 (1439) Int-2: 10 (1863) Ctrl: 10 (1474) Total N: 30 (4776)  <u>N of Clusters (subjects) participated</u> Int-1: 10 (1293) Int-2: 10 (1769) Ctrl: 10 (1390) Total N: 30 (4452) Analyzed sample: N = 4452 Age: 12 yrs (7 <sup>th</sup> grade) Sex (male): 50.5% Ethnicity: NR Alcohol use: NR Country: India
Interventions	<u>Intervention-1: HRIDAY</u> <u>Focus/target:</u> to improve CV health (diet, physical activity, and smoking prevention); alcohol was not a focus of intervention <u>Components:</u> 1) set of 10 posters showing different aspects of CV health, 2) booklets of HRIDAY providing basic information on heart health circulated to students, 3) 30-60 min classroom activities selected by teachers, 4) debates on banning tobacco sponsorship, 5) discussions on nutrition and food policy; 3 of 20 classroom activities concerned ways

Reddy 2002 (Continued)

	<p>to refuse offers to smoke  <u>Intervention-2</u>: HRIDAY plus family-based program  <u>Focus/target</u>: see above  <u>Components</u>: students bringing home 6 booklets with information and activities to share with families  <u>Fidelity</u>: 80% of the schools reported booklet distribution; the degree of fidelity did not differ across the intervention arms  <u>Duration/frequency</u>: 10 mo (HRIDAY), 6 mo (booklet distribution)  <u>Control</u>: standard curriculum</p>	
Outcomes	<p>FU: 17 mo (Int-1 vs. Int-2 vs. Ctrl)  <u>Having a drink of alcohol (ever)</u>  0.1286 (95% CI: 0.0958, 0.1706) vs. 0.1447 (95% CI: 0.1096, 0.1886) vs. 0.2886 (95% CI: 0.2298, 0.3555),  p &lt; 0.001 (Int1-2 vs. Ctrl)</p>	
Notes	<p><b>HRIDAY</b>  <u>Abbreviations</u>  FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); SE=standard error  NS=statistically non-significant; HRIDAY = health-related information and dissemination among youth; 95% CI=ninety-five percent confidence interval; CV=cardiovascular</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	no ITT and attrition < 20%, attrition rates similar between study arms
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	no imbalance in baseline covariate

## Ringwalt 1991

Methods	Design: RCT FU: 17 wks (post-randomisation) Attrition: 9% ITT: no Unit of randomisation: school Clustering effect adjusted: yes	
Participants	<u>N of Clusters (subjects) randomised</u> Int: 10 (685) Ctrl: 10 (585) Total N: 20 (1270)  <u>N of Clusters (subjects) participated</u> Int: 10 (NR) Ctrl: 10 (NR) Total N: 20 (3.2% parents refused permission to participate) Age: 10-12 yrs Sex (male): 48% Ethnicity: 50% Black Alcohol users: 37% had tried beer and 20% had tried wine Country: US	
Interventions	<u>Intervention</u> : school-based drug education program delivered by specially trained police officers <u>Focus/target</u> : Promote knowledge, attitudes and social skills to reduce drug use <u>Components</u> : resistance training, self-esteem, social skills, information, role-play. Delivered by trained, uniformed, police officer <u>Fidelity</u> : NR <u>Duration/frequency</u> : 17 weekly sessions (45-60 min each) in 17 weeks <u>Control</u> : standard curriculum	
Outcomes	FU: 17 wks <u>Lifetime involvement with alcohol (pre vs. post)</u> Int: 50.2% vs. 54.8% Ctrl: 40.0% vs. 49.8% p>0.05	
Notes	<b>DARE</b> Baseline differences on seven different variables. <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); SE=standard error NS=statistically non-significant; DARE=drug abuse resistance education	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

**Ringwalt 1991** (Continued)

Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT but low attrition of 9%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Differences in baseline covariate

**Ringwalt 2009**

Methods	<p>Design: RCT  FU: 3 yrs (post-randomisation)  Attrition: 21%; dropout rates non-differential; dropouts were less likely to be Whites (52.5% vs. 47.6%, p=0.03)  ITT: Yes  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 20 (3990)  Ctrl: 20 (4348)  Total N: 40 (8338)</p> <p><u>N of Clusters (subjects) participated</u>  Int: 17 (2983)  Ctrl: 17 (3045)  Total N: 34 (6028)</p> <p>Analyzed sample: N = 5883 (FU-2 yrs)  Analyzed sample: N = 4607 (FU-3 yrs)  Age: NR yrs (6<sup>th</sup>-8<sup>th</sup> grades)  Sex (male): 49.0%  Ethnicity: White (50.0%), African American (16.0%), Hispanic (26.0%)  Alcohol use (lifetime): Int (39.4%) vs. Ctrl (34.4%), p=0.06  30-day use of alcohol: 7.5% vs. 5.5%, p=0.03  Country: US</p>



Ringwalt 2009 (Continued)

Interventions	<p><u>Intervention</u>: ALERT</p> <p><u>Focus/target</u>: cigarettes, alcohol, marijuana, and inhalant use; motivates not to use substances, and provides skills for resisting inducements</p> <p><u>Components</u>: 11 lessons of 45 min (1<sup>st</sup> yr), followed by 3 booster lessons the following year; activities were guided class discussions, role playing exercises and videos</p> <p><u>Fidelity</u>: 97.4% of all lessons were delivered based on logs/records</p> <p><u>Duration/frequency</u>: weekly lessons (1 per wk) for 2 yrs</p> <p><u>Control</u>: Standard curriculum</p>	
Outcomes	<p>FU: 2 yrs (Int vs. Ctrl)</p> <p><u>Lifetime use of alcohol</u> 55.4% vs. 52.4% (OR=0.99, p=NR)</p> <p><u>30-day use of alcohol</u> 14.3% vs. 14.0% (OR=1.32, p=NR)</p> <p>FU: 3 yrs (Int vs. Ctrl)</p> <p><i>Mean 30-day alcohol use from baseline to FU (none, 1-2 days, 3-5 days, 6-19 days, and 20 or more days in last mo)</i></p> <p>Int: from 7.1 (25.6) to 22.1 (41.5)</p> <p>Ctrl: from 5.2 (22.2) to 19.7 (39.8)</p> <p>Between-group p=0.91</p> <p><i>% Students with lifetime alcohol use from baseline to FU</i></p> <p>Int: from 39.6 to 63.5</p> <p>Ctrl: from 34.6 to 59.9</p> <p>Between-group p=0.97</p>	
Notes	<p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); SE=standard error</p> <p>NS=statistically non-significant</p>	
<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer generated random numbers
Allocation concealment (selection bias)	Unclear risk	students unaware of their assignment
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described

Ringwalt 2009 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT and imputation done; attrition 21% but dropout non differential
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Differences in baseline covariate

Scaggs 1985

Methods	Design: RCT FU: 2 yrs (post-randomisation) Attrition: 26% ITT: No Unit of randomisation: class Clustering effect adjusted: No
Participants	<u>N of Clusters (subjects) randomised</u> Int: 3 (NR) Ctrl: 2 (NR) Total N: 5 (127)  <u>N of Clusters (subjects) participated</u> Int: 3 (NR) Ctrl: 2 (NR) Total N: 5 (NR)  <u>Analyzed sample</u> Int: 3 (48) Ctrl: 2 (45) Total N: 5 (93) Age: 14-15 yrs Sex (male): 48% Ethnicity: NR Alcohol users: 31% Country: US
Interventions	<u>Intervention</u> : a group-guidance based prevention program focusing on the knowledge, attitudes and behaviors of alcohol and other drug use <u>Focus/target</u> : knowledge and skills around alcohol and drugs <u>Components</u> : focus on personal use, decision making skills, family drinking, drinking and driving, knowledge of substances <u>Fidelity</u> : NR <u>Duration/frequency</u> : 45 min/day for 15 days <u>Control</u> : normal science lessons

Scaggs 1985 (Continued)

Outcomes	<p>FU: 2 yrs</p> <p><u>Reported alcohol use (males):</u>            Int: N=12 (54.6%; -0.2% from baseline)            Ctrl: N = 12 (60.0%; +4.4% from baseline)</p> <p><u>Reported alcohol use (females):</u>            Int: N=9 (33.4%; -7.7% from baseline)            Ctrl: N=10 (58.9%; +29.4% from baseline)</p> <p>No results of formal statistical analyses were reported</p>
----------	---

Notes	<p><b>The Substance Abuse Awareness Program Prevention Model</b>            Described as a quasi-experimental study, but details indicate a RCT. Discrepancy between reported attrition rates and cell sizes in analyses. Possible contamination between groups</p> <p><u>Abbreviations</u>            FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s)</p>
-------	--

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	no ITT and high attrition of 264%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Possible contamination between study arms

Schinke 2000

Methods	<p>Design: RCT  FU: 3.5 yrs (post-randomisation)  Attrition: 14%  ITT: no  Unit of randomisation: school  Clustering effect adjusted: no</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: NR (NR)  Ctrl: NR (NR)  Total N: 27 (1396)</p> <p><u>N of Clusters (subjects) participated</u>  Int: NR (1335)  Ctrl: NR (1343)  Total N: 27 (1396)  Age (mean): 10.3 yrs (3<sup>rd</sup>-5<sup>th</sup> Grades)  Sex (male): 51%  Ethnicity: 100% Native Americans  Alcohol users: 9% weekly drinkers (4+ drinks)  Country: US</p>
Interventions	<p><u>Intervention</u>: program derived from LST tailored to cultural setting and everyday context for Native American youth  <u>Focus/target</u>: cognitive and behavioral skills for substance abuse prevention  <u>Components</u>: problem-solving, personal coping, interpersonal communication - all incorporating Native American myths, legends and stories  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 15 x 50-minute weekly sessions + bi-annual booster sessions  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 0.5, 1.5, 2.5 and 3.5 yrs  <u>Weekly drinking: 4+ (Int vs. Ctrl)</u>  9.13% vs. 8.72% (baseline)  7.65% vs. 8.31% (0.5 yrs)  12.57% vs. 15.55% (1.5 yrs)  15.89% vs. 19.06% (2.5 yrs) *  22.87% vs. 30.17% (3.5 yrs) *  * p&lt;0.01</p>
Notes	<p><b>Culturally tailored LST</b>  Baseline descriptive information not reported  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); LST=life skills training</p>

Schinke 2000 (Continued)

<i>Risk of bias</i>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	no ITT but low attrition of 14%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Baseline descriptive information not reported

Sheehan 1996

Methods	Design: RCT FU: 3 yrs (post-randomisation) Attrition: 38% ITT: no Unit of randomisation: school Clustering effect adjusted: no
Participants	<u>N of Clusters (subjects) randomised</u> Int: 21 (NR) Ctrl: 20 (NR) Total N: 41 (4545)  <u>N of Clusters (subjects) participated</u> Int: 21 (NR) Ctrl: 20 (NR) Total N: 41 (4545) Age: 17 yrs Sex (male): 41% Ethnicity: NR Alcohol users: 10-13% weekly drinkers Country: Australia

Interventions	<p><u>Intervention</u>: program based on Theory of Reasoned Action</p> <p><u>Focus/target</u>: prevent drinking and driving through weakening student intentions to drink and drive</p> <p><u>Components</u>: modification of students' attitudes and beliefs towards drink driving, subjective beliefs and norms, and perceived control over their own behaviour. Extensive use of role play and interactional activities</p> <p><u>Fidelity</u>: NR</p> <p><u>Duration/frequency</u>: 12 lessons over 2 yrs</p> <p><u>Control</u>: NR</p>	
Outcomes	<p>FU: 3 yrs (pre- vs. post-test)</p> <p><u>Weekly drinkers</u></p> <p>Int: 10% vs. 36%</p> <p>Ctrl: 13% vs. 34%</p> <p>Chi-squared = 6.54(3), p=0.09</p> <p><u>Drinking and driving in past mo</u></p> <p>Reported drink driving at baseline: OR=0.59 (95% CI: 0.15, 2.1)</p> <p>No reported drink driving at baseline: OR=0.90 (95% CI: 0.61, 1.31)</p>	
Notes	<p><b>Plan a Safe Strategy (PASS)</b></p> <p>No differential attrition. FU described as a random sample of 62% of baseline sample</p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;</p> <p>wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); LST=life skills training;</p> <p>PASS=plan safe strategy; OR=odds ratio; 95% CI: ninety-five percent confidence interval</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT and high attrition of 38%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported

Sheehan 1996 (Continued)

Other bias	Unclear risk	Baseline covariate distribution not described
------------	--------------	---

Sloboda 2009

Methods	<p>Design: RCT          FU: 5 yrs (post-randomisation)          Attrition: 45%          ITT: No          Unit of randomisation: school          Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomized</u>          Int: 41 (11314)          Ctrl: 42 (8215)          Total N: 83 (19529)</p> <p><u>N of Clusters (subjects) participated</u>          Int: 40 (10028)          Ctrl: 40 (7292)          Total N: 80 (17320)          Analyzed sample: N = 17320          Age: 12.5 yrs (7<sup>th</sup> grade)          Sex (male): 49.0%          Ethnicity: White (33% vs. 40%), Black (12.6% vs. 15.4%), Hispanic (27.8% vs. 17.9%)          , Asian (4.5%), American Indian (8.3%), and          Other (11.8% vs. 12.9%)          Alcohol use (past 30-d): 13.8% vs. 11.8%          Country: US</p>
Interventions	<p><u>Intervention</u>: TCYL  <u>Focus/target</u>: to prevent use of alcohol, tobacco, and other drugs by demonstrating consequences of the use of alcohol and drugs to students  <u>Components</u>: 10 lessons (for 7<sup>th</sup> grade) and booster of 7 lessons (for 9<sup>th</sup> grade); the lessons cover normative beliefs, consequences, decision making and resistance skills  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 1 yr (in 7<sup>th</sup> grade) and 1 yr (in 9<sup>th</sup> grade)  <u>Control</u>: standard curriculum</p>
Outcomes	<p>FU: 5 yrs (Int vs. Ctrl)  <u>Binge drinking (within 14 days)</u>          28.1% vs. 24.7% (OR= 1.14, 95% CI: 1.01, 1.27)  <u>Alcohol use (within 30 days)</u>          45.7% vs. 41.9% (OR= 1.09, 95% CI: 1.01, 1.18)  <u>Got drunk (within 30 days)</u>          30.0% vs. 27.3% (OR= 1.10, 95% CI: 0.98, 1.22)</p>

	<p><u>Alcohol use (within 12 mo)</u> 61.1% vs. 58.7% (OR= 1.04, 95% CI: 0.98, 1.10)</p> <p><u>Got drunk (within 12 mo)</u> 43.3% vs. 41.2% (OR= 1.05, 95% CI: 0.96, 1.14)</p> <p>The negative effect of intervention on 30 day and 12 mo alcohol use measures persisted in males but not in females; in White students, the intervention was associated with significantly higher risk for alcohol use compared with Ctrl; whereas no such associations were present in non-White students; non-users of alcohol in treatment group were at higher risk for alcohol use</p>
--	---

Notes	<p><b>TCYL - Take Charge of Your Life</b> Dropouts were more likely older, female, non-White, alcohol, marijuana or drug users; differential rates were for race with more of those classified in other race being in the Ctrl arm; missing data were imputed</p> <p><u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); PASS=plan safe strategy; OR=odds ratio; 95% CI: ninety-five percent confidence interval; TCYL= take charge of your Life</p>
-------	--

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT and high possibly differential attrition > 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Imbalance in baseline covariate may have confounded the association



## Spoth 2002

Methods	<p>Design: RCT  FU: 5.5 yrs (post-randomisation)  Attrition: 26%  ITT: yes  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int-1: NR (618)  Int-2: NR (541)  Ctrl: NR (491)  Total N: 36 (1650)</p> <p><u>N of Clusters (subjects) participated</u>  Int-1: NR (428)  Int-2: NR (450)  Ctrl: NR (347)  Total N: 36 (1225)  Analyzed sample: N = 1650  Age: NR (7<sup>th</sup> grade)  Sex (male): 52.0%  Ethnicity: White (96%)  Alcohol use (ever): Int-1/2 (56%) vs. Ctrl (46.9%)  Country: US</p>
Interventions	<p><u>Intervention-1: LST</u>  <u>Focus/target:</u> to promote skill development, social resistance, self-management  <u>Components:</u> 15 sessions and 5 booster sessions a year later and in 11<sup>th</sup> grade in 6 schools</p> <p><u>Intervention-2: SFP + LST</u>  <u>Focus/target:</u> to reduce youth substance use  <u>Components:</u> 7 sessions, 1 h parent and youth skill-building curricula, followed by conjoint family curricula to practice the skills,  4 booster sessions a yr later, and in 11<sup>th</sup> grade in 6 schools  <u>Fidelity:</u> 92%-98% (SFP), 77%-85% (LST)  <u>Duration/frequency:</u> 7 wks (SFP) 1 school yr (LST)  <u>Control:</u> leaflets on teen development sent to parents</p>
Outcomes	<p><u>FU: 1 yr (Int-1 vs. Int-2 vs. Ctrl)</u>  New users of alcohol: 35.2% vs. 25.7% vs. 36.7%, p&lt;0.05 (Int-2 vs. Int-1 or Ctrl)</p> <p><u>FU: 2.5 yr (Int-1 vs. Int-2 vs. Ctrl)</u>  RAU (mean/SE): 0.23±0.02 vs. 0.19±0.02 vs. 0.24±0.02, p&gt;0.05  RAU (growth trajectories): between-arm differences - NS  WD (mean/SE): 0.04±0.01 vs. 0.03±0.01 vs. 0.05±0.01, p&lt;0.05 (Int-2 vs. Ctrl)  WD (growth trajectories) between-arm differences: NS</p> <p><u>FU: 5.5 yrs (Int-1 vs. Int-2 vs. Ctrl)</u>  Alcohol initiation (mean/SE): 0.94±0.02 vs. 0.93±0.02 vs. 0.96±0.02, p&gt;0.05  Alcohol initiation (growth trajectories): p&lt;0.05 in favour of Int-2 (Int-2 vs. Ctrl)  Drunkness initiation (mean/SE): 0.60±0.03 vs. 0.64±0.03 vs. 0.68±0.03, p&gt;0.05</p>

**Spoth 2002** (Continued)

	Drunkness initiation (growth trajectories): $p < 0.05$ in favour of Int-1 and Int-2 (Int-2 or Int-1 vs. Ctrl)
Notes	<p><b>LST - Life Skills Training</b></p> <p>The effects of interventions were assessed in a subset of risky drinker students in whom alcohol and drunkness frequency was significantly reduced in Int-1 compared to Ctrl, but respective growth trajectories were similar. There was some imbalance in baseline factors, % students living with both parents was greater in Ctrl (78%) vs. Int-1 (69.6%) and alcohol users in Int-1 and Int-2 (56-57% vs. 47%)</p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; LST=life skills training; SFP=strengthening families program; RAU=regular alcohol use; WD=weekly drunkness</p>

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT done; imputation of missing values; attrition 25% but non-differential
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Baseline imbalance in covariate

Methods	<p>Design: RCT  FU: 2 yrs (post-randomisation)  Attrition: 18%  ITT: No  Unit of randomisation: class within school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int-1: NR (NR)  Int-2: NR (NR)  Ctrl: NR (NR)  Total N: NR (NR)</p> <p><u>N of Clusters (subjects) participated</u>  Int-1: NR (NR)  Int-2: NR (NR)  Ctrl: NR (NR)  Total N: 8 (1649)</p> <p><u>N of Clusters (subjects) analysed</u>  Int-1: NR (NR)  Int-2: NR (NR)  Ctrl: NR (NR)  Total N: NR (NR)  Age: NR (Grade 7)  Sex (male): 50.5%  Ethnicity: Caucasians (81.4%), African Americans (5.4%), Native Americans (2.2%), Hispanic (1.3%), Asians (1.1%), and Other (8.5%)  Alcohol users: NR  Country: US</p>
Interventions	<p><u>Intervention-1: adult-led ALERT</u>  <u>Focus/target:</u> drug and alcohol use  <u>Intervention-2: adult-led and teen-assisted ALERT</u>  <u>Focus/target:</u> drug and alcohol use  <u>Components (Int-1/2):</u> To develop reasons not to use drugs, discuss pressure leading to drug use, learn resistance skills, and prevalence of drug use  <u>Fidelity:</u> high; mean rating 5.8-6.8 (7<sup>th</sup> grade) and 4.8-6.7 (8<sup>th</sup> grade)  <u>Duration/frequency:</u> 7th grade - 11 sessions weekly; 8th grade - 3 sessions  <u>Control:</u> NR</p>
Outcomes	<p>FU: 2 yrs  The graph showed no difference between ALERT and Control Group for alcohol use in the past mo at final follow-up  <u>Multivariate statistical analysis</u>  Past month alcohol use: beta = -0.011 (SE=0.17), NS (Int-1 vs. Int-2 vs. Ctrl)  Past year alcohol use: beta = -0.071 (SE=0.15), NS (Int-1 vs. Int-2 vs. Ctrl)  Binge drinking: beta = -0.001 (SE=0.25), NS (Int-1 vs. Int-2 vs. Ctrl)</p>

St. Pierre 2005 (Continued)

Notes	<p><b>Project ALERT</b> Possible contamination because of allocation by class within school, although authors discount this by comparison with substance use rates in other schools. There was no program effect modification by gender</p> <p><u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; SE=standard error</p>
-------	--

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT but low attrition of 18%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Possible contamination

**Sun 2008**

Methods	<p>Design: RCT FU: 1 yr (post-randomisation) Attrition: 27% ITT: No Unit of randomisation: school district Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u> Int-1: NR (NR) Int-2: NR (NR) Ctrl: NR (NR) Total N: 9 (3908)</p> <p><u>N of Clusters (subjects) participated</u></p>

	<p>Int-1: 3 (NR)  Int-2: 3 (NR)  Ctrl: 3 (NR)  Total N: 9 (2734)</p> <p><u>Analyzed sample</u>  Int-1: 3 (767)  Int-2: 3 (688)  Ctrl: 3 (609)  Total N: 9 (2064)  There were 18 schools in the randomised groups, 6 schools per arm  Age: 13-19 yrs (9<sup>th</sup> grade)  Sex (male): 52.1%  Ethnicity: White (18.2%), Black (8.1%), Hispanic (62.1%), Asian (8.4%), and Other (3.2%)  Alcohol use (past 30-d): Int-1 (38.7%) vs. Int-2 (37.4%) vs. Int-3 (38.6%)  Country: US</p>
Interventions	<p><u>Intervention-1: CPI</u>  <u>Focus/target:</u> to change youths' attitudes/beliefs on their drug use  <u>Components:</u> classes with program curriculum</p> <p><u>Intervention-2: CPI + BSC</u>  <u>Focus/target:</u> BSC gives instruction in social skills and behavioral self management to allow youth flexible bond with peer groups, seek out social support when needed, and minimize stressful or conflict-type interactions  <u>Components:</u> classes with program curriculum</p> <p><u>Fidelity:</u> NR  <u>Duration/frequency:</u> 3 times/week for 4 wks  <u>Control:</u> standard curriculum</p>
Outcomes	<p><u>FU:</u> 1 yr  <u>Int-1 vs. Ctrl</u>  Alcohol use (past 30-d): OR=0.98, 95% CI: 0.63, 1.50  N of times of alcohol use (past 30-d): RR=0.92, 95% CI: 0.70, 1.21</p> <p><u>Int-2 vs. Ctrl</u>  Alcohol use (past 30-d): OR=1.03, 95% CI: 0.66, 1.58  N of times of alcohol use (past 30-d): RR =0.84, 95% CI: 0.64, 1.11</p> <p><u>Int-2 vs. Int-1</u>  Alcohol use (past 30-d): OR=1.05, 95% CI: 0.71, 1.55  N of times of alcohol use (past 30-d): RR =0.91, 95% CI: 0.69, 1.20</p>
Notes	<p><b>TND-4 - Project Towards no Drug Abuse</b>  Dropouts were more likely older, female, non-White, smoker; differential rates were not observed across the study arms  Imbalance in ethnicity and school type at baseline; statistical adjustment was done</p> <p><u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;  wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=</p>

Sun 2008 (Continued)

	month(s); NS=statistically non-significant; RR=relative risk; OR=odds ratio; 95% CI=ninety-five percent confidence interval; CPI=cognitive perception information; BSC=behavioral skills curriculum; TND=project towards no drug abuse	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	High risk	No ITT, attrition > 20%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Imbalance in baseline covariate

van Lier 2009

Methods	Design: RCT FU: 6 yrs (post-randomisation) Attrition: 28% ITT: unclear Unit of randomisation: class Clustering effect adjusted: yes
Participants	<u>N of Clusters (subjects) randomised</u> Int: NR (NR) Ctrl: NR (NR) Total N: NR (666)  <u>N of Clusters (subjects) participated</u> Int: NR (NR) Ctrl: NR (NR) Total N: NR (525) Analyzed sample: N = 477 Age: 7 yrs (1 <sup>st</sup> grade) Sex (male): NR% Ethnicity: NR

	Alcohol use (past 30-d): NR Country: Netherlands
Interventions	<u>Intervention</u> : GBG <u>Focus/target</u> : to prevent aggressiveness, opposition and ADHD symptoms <u>Components</u> : done in 3 stages; choosing class rules, accompanied by pictograms; teachers assigned students to one of 4 teams and members are encouraged to manage their own and team-mates behavior; each team receives a number of cards and one card is taken when a student violates a rule and are rewarded if at least one card remains <u>Fidelity</u> : in 9 out of 13 schools GBG was carried completely and in 4 - it was incomplete <u>Duration/frequency</u> : 3 times a wk (10 min sessions) for 2 yrs <u>Control</u> : standard curriculum
Outcomes	FU: 3 to 6 yrs (Int vs. Ctrl) <u>Alcohol use growth parameter</u> Slope: 0.05±0.19, NS (past yr) Slope: -0.31±0.15, NS (past mo) Slope: -0.43±0.17, p<0.05 (past wk)
Notes	<b>GBG - Good Behavior Game</b> <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; GBG=good behavior game; ADHD=attention deficit hyperactive disorder

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	ITT not done; dropouts differ from retainers in socioeconomic status
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Not described

Methods	<p>Design: RCT  FU: 12 mo (post-randomisation)  Attrition: 49%  ITT: yes  Unit of randomisation: school  Clustering effect adjusted: yes</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: 8 (833)  Ctrl: 8 (1159)  Total N: 16 (1992)</p> <p><u>N of Clusters (subjects) participated</u>  Int: 8 (611)  Ctrl: 8 (835)  Total N: 16 (1446)  Analyzed sample: N = 1434  Age: 13 yrs (8<sup>th</sup> grade)  Sex (male): 59%  Ethnicity: NR  Alcohol use in males (weekly frequency over past 3 mo): Int (1.56±6.94) vs. Ctrl (1.24±6.45)  Alcohol use in females (weekly frequency over past 3 mo): Int (0.66±3.16) vs. Ctrl (0.35±1.37)  Drinking to excess on single occasion (frequency) in males: Int (0.39±2.40) vs. Ctrl (0.39±2.30)  Drinking to excess on single occasion (frequency) in females: Int (0.21±1.01) vs. Ctrl (0.18±0.61)  Country: Australia</p>
Interventions	<p><u>Intervention</u>: CLIMATE curriculum integrated computerized harm minimization program  <u>Focus/target</u>: to reduce alcohol misuse  <u>Components</u>: 40-min lessons broken by two parts: 1) 15-20 min computer-based lesson and 2) activities for teachers and students (discussions, decision-making, problem-solving, and skills)  <u>Fidelity</u>: only 1 teacher failed to deliver complete curriculum  <u>Duration/frequency</u>: 1 yr  <u>Control</u>: CLIMATE curriculum integrated harm minimization program delivered by a classroom teacher</p>
Outcomes	<p>FU: 12 mo (Int vs. Ctrl)  Alcohol use in males (weekly frequency over past 3 mo): Int (3.86±14.54) vs. Ctrl (3.50±13.12), NS  Alcohol use in females (weekly frequency over past 3 mo): Int (0.99±4.07) vs. Ctrl (2.25±10.16), p &lt;0.05  Drinking frequency to excess on single occasion in males over past 3 mo: Int (1.07±3.69) vs. Ctrl (1.16±4.72), NS  Drinking frequency to excess on single occasion in females over past 3 mo: Int (0.38±1.16)</p>



Vogl 2009 (Continued)

	vs. Ctrl (0.93±3.45), p <0.05 Alcohol-related harms in males (own): Int (11.67±27.51) vs. Ctrl (10.79±29.5), NS Alcohol-related harms in females (own): Int (3.30±9.69) vs. Ctrl (7.15±22.93), p <0.05
Notes	<b>CLIMATE</b> - Dropouts were more likely to be male, higher alcohol use; attrition was not differential - High attrition possibly leading to confounding - More males in Ctrl (69%) vs. Int (45%); analysis adjusted and stratification by gender was also done <u>Abbreviations</u> FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	ITT done; high attrition of 49%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Unclear risk	Imbalance in baseline covariate

**Werch 2008**

Methods	Design: RCT FU: 1 mo (post-randomisation) Attrition: 19%; attrition rates did not significantly differ across study conditions ITT: no Unit of randomisation: individual level Clustering effect adjusted: NA
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: NA (NR) Int-2: NA (NR) Int-3: NA (NR)

	<p>Total N: NA (385)</p> <p><u>N of Clusters (subjects) participated</u></p> <p>Int-1: NA (NR)</p> <p>Int-2: NA (NR)</p> <p>Int-3: NA (NR)</p> <p>Total N: NA (375)</p> <p><u>Analyzed sample:</u></p> <p>Int-1: NA (113)</p> <p>Int-2: NA (113)</p> <p>Int-3: NA (109)</p> <p>Total N: NA (335)</p> <p>Age: 17 yrs (11<sup>th</sup> and 12<sup>th</sup> grades)</p> <p>Sex (male): 43%</p> <p>Ethnicity: White (49%), Black (23%), and Hispanic (6%)</p> <p>Alcohol use (over 30 d): 44%</p> <p>Country: US</p>
Interventions	<p><u>Intervention-1:</u> Goal Survey (based on BIM)</p> <p><u>Focus/target:</u> to influence multiple health habits by addressing images of college and career success</p> <p><u>Components:</u> printed text and scripted messages showing images of successful young adults while avoiding alcohol and tobacco smoking</p> <p><u>Intervention-2:</u> Goal Survey + Contract (based on BIM)</p> <p><u>Focus/target:</u> Contract was designed to help students in selecting goals leading to successful/happy life</p> <p><u>Components:</u> see above</p> <p><u>Intervention-3:</u> Goal Survey + Consult (based on BIM)</p> <p><u>Focus/target:</u> to provide image-based feedback tailored to personal development and health behaviors</p> <p><u>Components:</u> see above</p> <p><u>Fidelity:</u> NR</p> <p><u>Duration/frequency:</u> 1 mo</p>
Outcomes	<p>FU: 1 mo (Int-1 vs. Int-2 vs. Int-3)</p> <p><u>Intention to use alcohol (1='definitely not' to 4='definitely will')</u></p> <p>2.28±0.1 vs. 2.27±0.1 vs. 2.24±0.1, p&lt;0.001</p> <p><u>Length of alcohol use (1='do not use' to 5='1 year or more')</u></p> <p>2.69±0.2 vs. 2.47±0.2 vs. 2.61±0.2, p=0.05</p> <p><u>30-day alcohol frequency (1='0 days' to 7='all 30 days')</u></p> <p>1.91±0.1 vs. 1.74±0.1 vs. 1.81±0.1, p=0.13</p> <p><u>30-day alcohol quantity (1='do not drink' to 12='?11 drinks')</u></p> <p>3.27±0.3 vs. 3.32±0.3 vs. 3.42±0.3, p=0.82</p>
Notes	<p><b>Behavior-image model (BIM)</b></p> <p><u>Abbreviations</u></p> <p>FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=</p>

**Werch 2008** (Continued)

	control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); NS=statistically non-significant; BIM= behavior-image model	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	no ITT; attrition 19% and nondifferential
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	No significant imbalance in covariates between study arms

**Werch 2010**

Methods	Design: RCT FU: 3 mo (post-randomisation) Attrition: 13% ITT: no Unit of randomisation: individual Clustering effect adjusted: NA
Participants	<u>N of Clusters (subjects) randomised</u> Int: NA (NR) Ctrl: NA (NR) Total N: NA (416)  <u>N of Clusters (subjects) participated</u> Int: NA (179) Ctrl: NA (181) Total N: NA (360) Age: 15.8 yrs  Sex (male): 36.5%

Werch 2010 (Continued)

	<p>Ethnicity: Hispanics (15%), Asian (7%), Black (23%), White (46%), Other (24.4%)          Alcohol users (past 30-d): 24.5%          Any alcohol or drug problem: 30.5%          Country: US</p>	
Interventions	<p><u>Intervention</u>: BIM  <u>Focus/target</u>: alcohol and drugs; printed text and scripted messages designed to elicit a social image of successful young adult with life goals to increase positive behaviours while avoiding those that interfere with successful goals  <u>Components</u>: screening survey, consultation, and goal plan  <u>Fidelity</u>: research staff conducted independent rating of audio-taped sessions  <u>Duration/frequency</u>: 20 min consultation each for 3 mo  <u>Control</u>: standard curriculum</p>	
Outcomes	<p>FU: 3 mo  <u>30-day mean alcohol frequency from baseline to FU (1='0 days' to 11='all 30 days')</u>          Int: from 1.49 (0.08) to 1.35 (0.07)          Ctrl: from 1.31 (0.08) to 1.39 (0.07)          Between-group p=0.06</p> <p><u>30-day alcohol quantity (1='0 drinks per day' to 12=11 or more drinks per day )</u>          Int: from 2.05 (0.16) to 1.89 (0.16)          Ctrl: from 1.61 (0.16) to 1.83 (0.16)          Between-group p=0.05</p> <p><u>Heavy use of alcohol in past 30 days (4-5 more drinks in a row)</u>          Int: from 1.32 (0.08) to 1.20 (0.05)          Ctrl: from 1.17 (0.08) to 1.15 (0.05)          Between-group p=0.44</p> <p><u>Total number of alcohol/drug problems in past 30 days (17-item instrument)</u>          Int: from 1.35 (0.20) to 1.11 (0.20)          Ctrl: from 1.22 (0.20) to 1.11 (0.20)          Between-group p=0.65</p>	
Notes	<p><b>Behavior-image model (BIM)</b>          In baseline substance users subgroup, the intervention was considered as selective and the data was not abstracted</p> <p><u>Abbreviations</u>          FU=follow-up; RCT=randomized controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); BIM= behavior-image model</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>

**Werch 2010** (Continued)

Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	no ITT but low attrition of 13%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	Low risk	No significant imbalance in covariate between study arms

**Wilhelmsen 1994**

Methods	Design: RCT FU: 4 months (post-randomisation) Attrition < 5% ITT: no Unit of randomisation: school, stratified by location and SES Clustering effect adjusted: no
Participants	<u>N of Clusters (subjects) randomised</u> Int-1: 4 (NR) Int-2: 4 (NR) Ctrl: 4 (NR) Total N: 12 (1042)  <u>N of Clusters (subjects) participated</u> Int-1: 4 (NR) Int-2: 4 (NR) Ctrl: 4 (NR) Total N: 12 (955) Age: 12-13 yrs Sex (male): NR Ethnicity: NR Alcohol users: 31% Country: Norway
Interventions	<u>Intervention-1: HRS - highly structured intervention</u> <u>Focus/target:</u> Targeting social cognitive structures predictive of drug use <u>Components:</u> School-based prevention program providing knowledge of alcohol use and local traditions, norms, managing

Wilhelmsen 1994 (Continued)

	<p>pressures and attitudes  <u>Intervention-2</u>: LRS - loosely structured intervention  <u>Focus/target</u>: Targeting social cognitive structures predictive of drug use  <u>Components</u>: school-based prevention program providing knowledge of alcohol use and local traditions, norms, managing pressures and attitudes</p> <p><u>Fidelity</u>: NR  <u>Duration/frequency</u>: 10 sessions / 2 weeks  <u>Control</u>: NR</p>	
Outcomes	<p>FU: 4 months  <u>Mean (SD) frequency of use (6-point scale)</u>  Int-1: 0.53 (1.4)  Int-2: 0.90 (1.0)  Ctrl: 0.69 (1.3)  In ANCOVA a SS interaction between program and time <math>F(2,838) = 6.22, (p &lt; 0.01)</math> was due to an increase in Int-2 scores (<math>p &lt; 0.05</math>) and a decrease in Int-1 scores (<math>p &lt; 0.01</math>) and no change in the Ctrl group</p>	
Notes	<p><b>Alcohol Prevention Programmes</b>  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported; wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s); BIM= behavior-image model; SS=statistically significant; HRS=highly role-specified; LRS= less role-specified; ANCOVA=analysis of covariance</p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No ITT but low attrition 5%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported

Wilhelmsen 1994 (Continued)

Other bias	Low risk	No significant imbalance in covariate between study arms
------------	----------	--

Williams 1968

Methods	<p>Design: RCT  FU: 1 yr (post-randomisation)  Attrition: 14%  ITT: no  Unit of randomisation: individual  Clustering effect adjusted: NA</p>
Participants	<p><u>N of Clusters (subjects) randomised</u>  Int: NA (111)  Ctrl: NA (94)  Total N: NA (205)</p> <p><u>N of Clusters (subjects) participated</u>  Int: NA (111)  Ctrl: NA (94)  Total N: NA (205)  Age: 16-17 yrs  Sex (male): 100%  Ethnicity: NR  Alcohol users: NR  Country: US</p>
Interventions	<p><u>Intervention</u>: Massachusetts alcohol education program to reduce alcoholism in the population  <u>Focus/target</u>: increase knowledge about alcohol  <u>Components</u>: discussion group - examine own and peer attitudes to drinking. Factual information  <u>Fidelity</u>: NR  <u>Duration/frequency</u>: 5 sessions per yr  <u>Control</u>: alternative topics chosen for discussion</p>
Outcomes	<p>FU: 1 yr (Int vs. Ctrl)  Alcohol use in past yr: 85% (77% at pre-test) vs. 85% (74% at pre-test), <math>p &gt; 0.05</math>  Drank &gt; 5 times in past yr (drinkers): 72% (50% at pre-test) vs. 61% (43% at pre-test), <math>p &gt; 0.05</math>  Intoxicated in past yr (drinkers): 64% (25% at pre-test) vs. 57% (29% at pre-test), <math>p &gt; 0.05</math></p>
Notes	<p><b>Massachusetts Program</b>  Risk of contamination as same school provided both groups  <u>Abbreviations</u>  FU=follow-up; RCT=randomised controlled trial; N=number; Int=intervention; Ctrl=control; yr(s)=year(s); NR=not reported;</p>

**Williams 1968** (Continued)

wk(s)=week(s); d=day(s); ITT=intention to treat (analysis); NA=not applicable; mo=month(s)		
<i>Risk of bias</i>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding (performance bias and detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	no ITT low attrition 14%
Selective reporting (reporting bias)	Low risk	All outcomes of interest specified in study methods section were reported
Other bias	High risk	Risk of contamination as same school provided both groups

**Characteristics of excluded studies** [ordered by study ID]

Study	Reason for exclusion
Ackermann 2008	Participant age > 18 years old
Amaro 2009	Selective/indicated prevention
Anderson 2004	No relevant outcomes
Bailey 2004	Community-based intervention
Bauman 2002	Family-based intervention
Bell 2005	No relevant outcomes
Benner 2008	No relevant outcomes
Bersamin 2007	Participant age > 18 years old
Boekeloo 2004	Office-based intervention



(Continued)

Botvin 2006	No relevant outcomes
Brody 2004	No relevant outcomes
Brody 2005	No relevant outcomes
Brody 2006	Family-based intervention
Brody 2008	No relevant outcomes
Brown 2005	Multi-component intervention
Brown 2007	Meta-analysis
Bryan 2009	Selective/indicated prevention
Castellanos 2006	No relevant outcomes
Caudill 2007	Selective/indicated prevention
Connell 2007	Family/multi-component intervention
Conrod 2006	Selective/indicated prevention
Conrod 2008	Selective/indicated prevention
Croom 2009	Participant age > 18 years old
D'Amico 2008	Selective/indicated prevention
DeGarmo 2009	Multi-component intervention
Dembo 2002	Selective/indicated prevention
Donohue 2004	Participant age > 18 years old
Eddy 2003	Multi-component intervention
Elder 2002	Selective/indicated prevention
Elliot 2004	No relevant outcomes
Friedman 2002	Selective/indicated prevention
Fromme 2004	Participant age > 18 years old
Griffin 2003	Selective/indicated prevention

(Continued)

Griffin 2004	No relevant outcomes
Griffin 2006	Family-based intervention
Haggerty 2006	Family-based intervention
Haggerty 2007	Family-based intervention
Haggerty 2008	Selective/indicated prevention
Hembroff 2007	Participant age > 18 years old
Jemmott 2005	Participant age > 18 years old
Jones 2005	Family-based intervention
Martinez 2005	Selective/indicated prevention
Poduska 2008	No relevant outcomes
Simons-Morton 2005	Multi-component intervention
Sussman 2002	Not randomised trial
Wagenaar 2005	Not randomised trial
Wolchik 2002	Selective/indicated prevention
Wu 2003	Family-based intervention

## DATA AND ANALYSES

This review has no analyses.

## APPENDICES

### Appendix I. Medline search Strategy

1. exp ALCOHOLS/ad, ae
2. exp Alcohol Drinking/
3. exp Alcohol Abuse/
4. exp Alcohol, Ethyl/ae
5. exp Alcohol Abuse/mo, pc, rh, th
6. alcohol\$.ti,ab.
7. drink\$.ti,ab.
8. drunk\$.ti,ab.
9. intoxicat\$.ti,ab.
10. or/1-9
11. teenage\$.ti,ab. or youth.ti,ab or adolescent\$.ti,ab
12. (early adj2 adult\$).ti,ab.
13. (young adj2 adult\$).ti,ab.
14. exp adolescent/ or exp child/ or exp students/
15. or/11-14
16. intervention\$.ti,ab.
17. educat\$.ti,ab.
18. promot\$.ti,ab.
19. adverti\$.ti,ab.
20. campaign\$.ti,ab.
21. (mass adj2 media).ti,ab.
22. (primary adj5 prevention).ti,ab.
23. (secondary adj5 prevention).ti,ab.
24. (universal adj5 prevention).ti,ab.
25. (selective adj5 prevention).ti,ab.
26. (target\$ adj5 prevention).ti,ab.
27. exp HEALTH EDUCATION/
28. School\$.ti,ab or peer\$.ti,ab or Curriculum.ti,ab or skill\$.ti,ab or program\$.ti,ab
29. or/16-28
30. randomized controlled trial.pt
31. controlled clinical trial.pt
32. random\$.ti,ab
33. placebo.ti,ab
34. drug therapy.fs
35. trial.ab
36. groups.ab
37. OR/ 31-37
38. exp animals/ NOT humans.sh
39. 37 NOT 38
40. 10 and 15 and 29 and 39
41. limit 40 to yr="2002 -Current"

## Appendix 2. EMBASE Search strategy

- 1 random\$.ab,ti.
- 2 placebo.ab,ti.
- 3 ((singl\$ or doubl\$ or trebl\$ or tripl\$) and (blind\$ or mask\$)).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer name]
- 4 (cross-over\$ or crossover\$).tw.
- 5 randomized controlled trial/
- 6 phase-2-clinical-trial/
- 7 phase-3-clinical-trial/
- 8 double blind procedure/
- 9 single blind procedure/
- 10 crossover procedure/
- 11 Latin square design/
- 12 exp PLACEBOS/
- 13 multicenter study/
- 14 or/1-13
- 15 limit 14 to human
- 16 exp alcohol/
- 17 Drinking Behavior/
- 18 Alcoholism/
- 19 exp alcohol abuse/
- 20 exp Alcohol Drinking/
- 21 drink\$.ti,ab.
- 22 drunk\$.ti,ab.
- 23 intoxicat\$.ti,ab.
- 24 alcohol.ti,ab.
- 25 or/16-24
- 26 adolescen\$.ti,ab.
- 27 teenage\$.ti,ab.
- 28 (young adj2 people).ti,ab.
- 29 (early adj2 adult\$).ti,ab.
- 30 (young adj2 adult\$).ti,ab.
- 31 youth\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer name]
- 32 exp adolescent/ or exp child/ or exp students/
- 33 exp youth/
- 34 or/26-33
- 35 intervention\$.ti,ab.
- 36 educat\$.ti,ab.
- 37 promot\$.ti,ab.
- 38 adverti\$.ti,ab.
- 39 campaign\$.ti,ab.
- 40 (mass adj2 media).ti,ab.
- 41 (primary adj5 prevention).ti,ab.
- 42 (secondary adj5 prevention).ti,ab.
- 43 (universal adj5 prevention).ti,ab.
- 44 (selective adj5 prevention).ti,ab.
- 45 (target\$ adj5 prevention).ti,ab.
- 46 exp education/
- 47 or/35-46
- 48 15 and 25 and 34 and 47
- 49 limit 48 to yr="2002 -Current"

### Appendix 3. PsycINFO search strategy

- 1 clinical trials.sh.
- 2 placebo.sh.
- 3 (Single adj blind\*).ab,ti.
- 4 (Single adj dumm\*).ab,ti.
- 5 (Single adj mask\*).ab,ti.
- 6 (Double adj blind\*).ab,ti.
- 7 (Double adj dumm\*).ab,ti.
- 8 (Double adj mask\*).ab,ti.
- 9 (triple adj blind\*).ab,ti.
- 10 (triple adj dumm\*).ab,ti.
- 11 (triple adj mask\*).ab,ti.
- 12 (treble adj blind\*).ab,ti.
- 13 (treble adj dumm\*).ab,ti.
- 14 (treble adj mask\*).ab,ti.
- 15 (control\* adj study).ab,ti.
- 16 (control\* adj studies).ab,ti.
- 17 (control\* adj trial\*).ab,ti.
- 18 (Random\* or sham or shams or placebo\* or RCT\*).ab,ti.
- 19 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18
- 20 alcohol\$.ti,ab.
- 21 drink\$.ti,ab.
- 22 drunk\$.ti,ab.
- 23 intoxicat\$.ti,ab.
- 24 exp sobriety/ or exp alcohol withdrawal/ or exp alcohol intoxication/ or exp alcoholism/ or exp alcohols/ or exp blood alcohol concentration/ or exp binge drinking/ or exp driving under the influence/ or exp alcohol abuse/ or exp alcoholic psychosis/ or exp alcohol rehabilitation/ or exp alcohol drinking patterns/
- 25 or/20-24
- 26 adolescen\$.ti,ab.
- 27 teenage\$.ti,ab.
- 28 (young adj2 people).ti,ab.
- 29 (early adj2 adult\$).ti,ab.
- 30 (young adj2 adult\$).ti,ab.
- 31 youth\$.mp. [mp=title, abstract, heading word, table of contents, key concepts]
- 32 exp adolescent/ or exp child/ or exp infant/ or exp students/
- 33 exp youth/
- 34 or/26-33
- 35 intervention\$.ti,ab.
- 36 educat\$.ti,ab.
- 37 promot\$.ti,ab.
- 38 adverti\$.ti,ab.
- 39 campaign\$.ti,ab.
- 40 (mass adj2 media).ti,ab.
- 41 (primary adj5 prevention).ti,ab.
- 42 (secondary adj5 prevention).ti,ab.
- 43 (universal adj5 prevention).ti,ab.
- 44 (selective adj5 prevention).ti,ab.
- 45 (target\$ adj5 prevention).ti,ab.
- 46 exp education/
- 47 or/35-46
- 48 19 and 25 and 34 and 47
- 49 limit 48 to yr="2002 -Current"

## HISTORY

Review first published: Issue 5, 2011

Date	Event	Description
6 September 2010	New search has been performed	This review represents a substantial update of the review “Primary prevention for alcohol misuse in young people” that has been split into three reviews. This represents one of the three. The other two reviews focus on universal family based prevention and on universal multi-component prevention

## CONTRIBUTIONS OF AUTHORS

DF conceived and led on the scope and design of the review. DF and AW both undertook searches, screening and data extraction. AW led on data analysis. Both DF and AW contributed to drafting and writing the review.

## DECLARATIONS OF INTEREST

DFs Department has received funding from the alcohol industry for adapting and evaluating a family based prevention program.

## SOURCES OF SUPPORT

### Internal sources

- Oxford Brookes University, UK.

Funding to employ a co-reviewer

### External sources

- NIHR, UK.

Small grant for updating the previous review

## NOTES

This review represents a substantial update of the review “Primary prevention for alcohol misuse in young people” that has been split into three reviews. This represents one of the three. The other two reviews focus on universal family based prevention and on universal multi-component prevention.

## **INDEX TERMS**

### **Medical Subject Headings (MeSH)**

\*School Health Services; Adolescent; Alcohol Drinking [\*prevention & control]; Alcoholic Intoxication [\*prevention & control]; Central Nervous System Depressants [poisoning]; Ethanol [poisoning]; Program Evaluation; Randomized Controlled Trials as Topic; Substance-Related Disorders [prevention & control]

### **MeSH check words**

Child; Child, Preschool; Female; Humans; Male