

 Open access • Journal Article • DOI:10.1103/PHYSREVLETT.123.140403

## Universal Spectra of Random Lindblad Operators. — [Source link](#)

[Sergey Denisov](#), [Tetyana Lapyeva](#), [Wojciech Tarnowski](#), [Dariusz Chruściński](#) ...+2 more authors

**Institutions:** [Metropolitan University](#), [Jagiellonian University](#), [Nicolaus Copernicus University in Toruń](#), [Polish Academy of Sciences](#)

**Published on:** 01 Oct 2019 - [Physical Review Letters](#) (American Physical Society)

**Topics:** [Open quantum system](#), [Random matrix](#), [Quantum chaos](#), [Dissipative system](#) and [Quantum evolution](#)

Related papers:

- [Spectral Gaps and Midgap States in Random Quantum Master Equations.](#)
- [Complex Spacing Ratios: A Signature of Dissipative Quantum Chaos](#)
- [Random Lindblad dynamics](#)
- [Hierarchy of Relaxation Timescales in Local Random Liouvillians.](#)
- [On the Generators of Quantum Dynamical Semigroups](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/universal-spectra-of-random-lindblad-operators-4zpxrc01w>

## Universal Spectra of Random Lindblad Operators

Sergey Denisov,<sup>1,2</sup> Tetyana Lapyteva,<sup>2</sup> Wojciech Tarnowski,<sup>3</sup> Dariusz Chruściński,<sup>4</sup> and Karol Życzkowski<sup>3,5</sup>


<sup>1</sup>*Department of Computer Science, OsloMet–Oslo Metropolitan University, NO-0130 Oslo, Norway*

<sup>2</sup>*Department of Control Theory and Systems Dynamics, Lobachevsky University,  
Gagarina Avenue 23, Nizhny Novgorod, 603950, Russia*

<sup>3</sup>*Marian Smoluchowski Institute of Physics, Uniwersytet Jagielloński, Krakow, Poland*

<sup>4</sup>*Institute of Physics, Faculty of Physics, Astronomy and Informatics Nicolaus Copernicus University,  
Grudziadzka 5/7, 87–100 Torun, Poland*

<sup>5</sup>*Center for Theoretical Physics, Polish Academy of Sciences, 02-668 Warszawa, Poland*

 (Received 4 December 2018; published 1 October 2019; corrected 04 October 2019)

To understand the typical dynamics of an open quantum system in continuous time, we introduce an ensemble of random Lindblad operators, which generate completely positive Markovian evolution in the space of the density matrices. The spectral properties of these operators, including the shape of the eigenvalue distribution in the complex plane, are evaluated by using methods of free probabilities and explained with non-Hermitian random matrix models. We also demonstrate the universality of the spectral features. The notion of an ensemble of random generators of Markovian quantum evolution constitutes a step towards categorization of dissipative quantum chaos.

DOI: [10.1103/PhysRevLett.123.140403](https://doi.org/10.1103/PhysRevLett.123.140403)

*Introduction.*—Any real system is never completely isolated from its environment and the theory of open quantum systems [1–3] provides appropriate tools to deal with such phenomena as quantum dissipation and decoherence. In the Markovian regime (which assumes a weak interaction between the system and its environment and separation of system and environmental timescales), the evolution of an  $N$ -level open quantum system can be modeled by using the master equation  $\dot{\rho}_t = \mathcal{L}(\rho_t)$ . The corresponding Markovian generator  $\mathcal{L}$  (often called a Lindblad operator or simply “Lindbladian” [1,4]) has the well-known Gorini-Kossakowski-Sudarshan-Lindblad form (GKSL) [5,6]

$$\mathcal{L}(\rho) = -i[H, \rho] + \mathcal{L}_D(\rho) = \mathcal{L}_U(\rho) + \mathcal{L}_D(\rho), \quad (1)$$

with the dissipative part

$$\mathcal{L}_D(\rho) = \sum_{m,n=1}^{N^2-1} K_{mn} \left[ F_n \rho F_m^\dagger - \frac{1}{2} (F_m^\dagger F_n \rho + \rho F_m^\dagger F_n) \right], \quad (2)$$

where traceless matrices  $\{F_n\}$ ,  $n = 1, 2, 3, \dots, N^2 - 1$ , satisfy the orthonormality  $\text{Tr}(F_n F_m^\dagger) = \delta_{n,m}$ . Finally, the complex “Kossakowski matrix”  $K = \{K_{mn}\}$  is positive semidefinite. The solution of the master equation  $\dot{\rho}_t = \mathcal{L}(\rho_t)$  gives rise to the celebrated Markovian semigroup  $\Lambda_t = e^{t\mathcal{L}}$ , such that for any  $t \geq 0$  map  $\Lambda_t$  represents a quantum channel, a completely positive and trace-preserving linear map [7].

In this Letter, we analyze the spectral properties of random Lindblad operators. Spectral analysis lies in the

heart of quantum physics. In the static case, the spectrum of the Hamiltonian provides full information about the possible state of the system and system evolution. Such an analysis plays also a key role in the study of dissipative quantum evolution—eigenvalues and eigenvectors of the Lindblad operator provide the full information about the dynamical properties of the open system [9]. Spectra of dynamical maps were recently addressed in Ref. [10] in connection to quantum non-Markovian evolution. This connection was experimentally verified recently [11], which proves that spectral techniques can be used to characterize non-Markovian behavior as well. Here, instead of analyzing specific physical models (like in Refs. [10,11]), we look for universal spectral properties displayed by “typical” Lindblad operators. It should be stressed that the standard examples of generators of order  $N = 2$ , usually considered in the literature, do not display universal features. We address the problem in the limit  $N \gg 1$  by using the apparatus of the random matrix theory (RMT) [12].

The RMT has already found many applications in physics. It started from the Wigner statistical approach to nuclear physics [13] and a series of Dyson papers on statistical theory of spectra [14–17]. It was soon realized that quantum dynamics corresponding to classically chaotic dynamics can be described by suitable ensembles of random matrices [18–20]. Depending on the symmetry properties of the system investigated, one may use orthogonal, unitary, or symplectic ensembles [12]. Similar ideas found applications in disorder systems and single-particle [21] and many-body [22] ones. From a different perspective, a deep connection to RMT was observed in




FIG. 1. Eigenvalue localization areas (gray) of a (a) random quantum channel, Eq. (3), and (b) random purely dissipative Lindblad operator, Eq. (7), and (c) random generic Lindbladian with relative weight of the unitary component  $\alpha$ , Eq. (8). While for the random channel (a) the distribution approaches a Girko's disk upon the increase of the number of system levels  $N$  [27], eigenvalues of the random purely dissipative Lindblad operator (b) fill the interior of the universal lemonlike contour, Eq. (7). For the Lindblad operator with the unitary component (c), the spectral boundary approaches an ellipse upon increase of  $\alpha$ .

the models of 2D quantum gravity [23,24] and gauge theories [25].

In the case of discrete dynamics described by quantum operations, various ensembles of random channels are known [26,27]. Recently, the RMT found interesting applications in quantum information theory [28–30]. In the case of continuous quantum dynamics, a class of Lindblad equations with decay rates obtained by tracing out a random reservoir was studied in Refs. [31,32]. RMT was applied to open quantum systems in the context of scattering matrices and non-Hermitian effective Hamiltonians [33].

Our perspective in this paper is different: We introduce an ensemble of random Lindblad operators, which describe continuous time evolution of an  $N$ -level open quantum system, and evaluate universal properties of operator spectra. Namely, we analyze the distribution of eigenvalues of a randomly chosen operator  $\mathcal{L}$  and study the scaling of spectral characteristics with  $N$ . For generators of classical Markovian evolution, a similar program was initiated and realized by Timm in Ref. [34].

We start by briefly recalling main results concerning random quantum channels [27,35]. Next, we analyze the extreme case of purely dissipative evolution,  $H = 0$  and  $\mathcal{L} = \mathcal{L}_D$ . This part can be considered as a quantum extension of the program outlined in Ref. [34] [from the opposite perspective, classical Pauli rate equations can be obtained as a reduction of the quantum master equation (1)]. Finally, we address the general situation, when both unitary and dissipative components  $\mathcal{L}_U$  and  $\mathcal{L}_D$  are present.

*Random quantum channels.*—An ensemble of random channels (i.e., completely positive and trace-preserving transformations [1–3])  $\Phi: M_N(\mathbb{C}) \rightarrow M_N(\mathbb{C})$  can be defined by the flat Hilbert-Schmidt measure in the space of all quantum operations. The spectrum of  $\Phi$  includes the leading eigenvalue  $\lambda_1 = 1$ , corresponding to the invariant state, while all remaining eigenvalues fill a disk of radius  $R = 1/N$  centered at zero; see Fig. 1(a). The bulk of the spectrum can be obtained by sampling random matrices

$(1/N)G_R$  [27], with  $G_R$  being a member of a real Ginibre ensemble [36–39].

Thus, for a generic superoperator  $\Phi$  the size of its spectral gap,  $\Delta_N = \lambda_1 - |\lambda_2| = 1 - 1/N$ , increases with dimension  $N$ , so the convergence to an invariant state becomes exponentially fast. For a large  $N$ , a typical channel becomes close to a one-step contraction, which sends any initial state into the invariant state,  $\Phi(\rho) = \rho_{\text{inv}} = \Phi(\rho_{\text{inv}})$ . It is known [40] that a typical channel is close to a unital one and the correction term  $\Phi(1) - 1$  behaves like a random Hermitian matrix of the Gaussian unitary ensemble with an asymptotically vanishing norm.

*Purely dissipative random Lindblad operators.*—To generate a random operator  $\mathcal{L}_D$ , we fix an orthonormal Hilbert-Schmidt basis  $\{F_n\}$  [41] and first sample a random Kossakowski matrix  $K$ . There are many ways to do such sampling. However, as we show below, a particular way in which this non-negative order  $N^2 - 1$  matrix is sampled is not important: The spectral features of random purely dissipative Lindbladians are *universal*.

The most intuitive way is to sample  $K$  from the ensemble of square complex Wishart matrices [50]. A Wishart matrix [51] has the structure  $W = GG^\dagger \geq 0$ , where  $G$  is a complex square Ginibre matrix with independent complex Gaussian entries [52]. Such a choice is also physically motivated by the fact that these ensembles of random matrices correspond to nonunitary evolution of quantum dynamical systems under the assumption of classical chaos [18,35].

We use the following normalization condition  $\text{Tr}K = N$ , that is,  $K = NGG^\dagger/\text{Tr}GG^\dagger$ . Note that eigenvalues of  $K$ ,  $\gamma_m$ ,  $m = 1, \dots, N^2 - 1$ , which can be interpreted as decay rates [1], are distributed according to the universal Marchenko-Pastur law [51] with the mean value  $\langle \gamma \rangle \sim 1/N$ . By diagonalizing the Kossakowski matrix, one can reduce the form of  $\mathcal{L}_D$  to

$$\mathcal{L}_D(\rho) = \sum_{m=1}^{N^2-1} \gamma_m \left[ V_m \rho V_m^\dagger - \frac{1}{2} (V_m^\dagger V_m \rho + \rho V_m^\dagger V_m) \right], \quad (3)$$




FIG. 2. Spectral density  $P[\text{Re}(\ell'), \text{Im}(\ell')]$  of the rescaled eigenvalues,  $\ell' = N(\ell + 1)$ , from the spectrum of random purely dissipative Lindblad operators  $\mathcal{L}_D$  for  $N = 50$  and  $100$ . We use two different sampling procedures for  $N = 100$  (a),(b), sampling the Kossakowski matrix from the Wishart ensemble and (c) an alternative procedure (Supplemental Material [41]). Note a perfect agreement with the asymptotic boundary of the spectral bulk, Eq. (7) (thick black line), derived with the random matrix model (6). Observe also a concentration of eigenvalues along the real axis, accompanied by depletion nearby—compare to Fig. 3(a)—which decreases with  $N$ . Each distribution was sampled with  $10^3$  realizations.

where  $V_m$  are called “jump operators” [53].  $\Phi(\rho) = \sum_m \gamma_m V_m \rho V_m^\dagger$  defines a Kraus representation of a completely positive map. Moreover,  $\sum_m \gamma_m V_m^\dagger V_m = \Phi^\dagger(\mathbb{1})$ , where  $\Phi^\dagger$  is the dual map,  $\text{Tr}[A \cdot \Phi^\dagger(B)] = \text{Tr}[\Phi(A) \cdot B]$ , and  $\mathbb{1}$  is the identity matrix in  $M_N(\mathbb{C})$ . Therefore, Eq. (3) can be rewritten as

$$\mathcal{L}_D(\rho) = \Phi(\rho) - \frac{1}{2}[(\Phi^\dagger(\mathbb{1})\rho + \rho\Phi^\dagger(\mathbb{1}))], \quad (4)$$

which shows that the purely dissipative Lindblad generator is fully determined by a completely positive map  $\Phi$ . If, in addition,  $\Phi$  is trace preserving, i.e., it is a quantum channel, we have  $\mathcal{L}(\rho) = \Phi(\rho) - \rho$ . This is not the case in general, and the Hermitian translation matrix  $X = \Phi^\dagger(\mathbb{1}) - \mathbb{1}$  does not vanish. Making use of this notation, we rewrite the Lindblad operator as  $\mathcal{L}_D(\rho) = [\Phi(\rho) - \rho] - \frac{1}{2}(X\rho + \rho X)$ .

Due to the trace-preserving quantum Markovian dynamics, a Lindblad generator always has a zero eigenvalue. If  $\Phi$  is a quantum channel, the spectrum of  $\mathcal{L}_D$  is the spectrum of  $\Phi$  shifted by  $-1$ . Thus, the leading eigenvalue  $\lambda_1 = 1$  is translated into  $\ell_1 = 0$  and the Girko disk is now centered at  $z = -1$ .

To sample spectra of random Lindbladians, we generate  $10^3$  realizations for different values of  $N$ , ranging from 30 to 100. In order to reveal the universality of spectra of the operators, it is useful to apply an affine transformation,  $\mathcal{L}'_D = N(\mathcal{L}_D + 1)$  [41]. Then the bulk of the spectrum of  $\mathcal{L}'$  becomes scale invariant and independent of  $N$ , see Figs. 2 and 3(a).

Sampled eigenvalue distributions  $P[\text{Re}(\ell'), \text{Im}(\ell')]$  are significantly different from the Girko disk and display a universal lemonlike shape. Already for  $N = 50$ , a single realization is enough to reproduce the universal shape, see Fig. 3(b). From the scale invariance, it follows that the spectral gap of  $\mathcal{L}_D$  scales as  $\Delta_N \simeq 1 - (2/N)$ . It is clear that the very term  $(X\rho + \rho X)$  is responsible for the

“disk  $\rightarrow$  lemon” deformation. The density inside the lemon is manifestly nonuniform. Also notable is the eigenvalue concentration along the real axis and the corresponding depletion nearby, see Fig 3(a). Although  $\mathcal{L}_D$  is represented by a complex matrix, it can be made real by a similarity transformation, which explains the concentration [37,39,55].

Finally, we performed sampling by using alternative algorithms (see Supplemental Material [41]) and obtained near identical results (differences are within the sampling errors) for  $N \geq 50$ ; see Fig. 2(c). This confirms the universality of the spectral distribution. It is noteworthy that a similar-looking shape of the eigenvalue distribution was observed with classical random transition rate matrices [see Fig. 7(b) in Ref. [34]]. However, the singularities at the real poles are much stronger in the classical case [56].

*Random matrix model.*—Let us recall that the spectrum of  $\mathcal{L}_D$  represented in (4) coincides with the spectrum of the following  $N^2 \times N^2$  complex matrix  $\mathcal{L}_{mn} = \text{Tr}[F_m \mathcal{L}_D(F_n)]$ .




FIG. 3. (a) Marginal distribution,  $\text{Re}(\ell') = 0$ , of rescaled eigenvalues for three different values of  $N$ . (b) Rescaled eigenvalues  $\ell'$  (empty dots) of a single Lindblad operator realization for  $N = 50$ . Red outer contour is the boundary derived from the random matrix model, Eq. (6).



This matrix becomes real if basis matrices  $F_m$  are Hermitian, due to the fact that  $\mathcal{L}_D$  is Hermiticity preserving. Another well-known matrix representation of the Lindblad operator reads

$$\hat{\mathcal{L}}_D = \hat{\Phi} - \mathbb{1} \otimes \mathbb{1} - \frac{1}{2}(X \otimes \mathbb{1} + \mathbb{1} \otimes \bar{X}), \quad (5)$$

where  $\hat{\Phi} = \sum_{m=1}^{N^2-1} \gamma_m V_m \otimes \bar{V}_m$ , and  $\bar{V}_m$  stands for the complex conjugation. Note that  $\hat{\Phi}$  is neither Hermitian nor real; however, the term  $X \otimes \mathbb{1} + \mathbb{1} \otimes \bar{X}$  is perfectly Hermitian. To understand the spectrum of  $\mathcal{L}'_D$ , we use the matrix representation (5) and approximate its rescaled version with the following random matrix (RM) model:

$$\hat{\mathcal{L}}'_D \approx G_R - (C \otimes \mathbb{1} + \mathbb{1} \otimes C). \quad (6)$$

The  $N^2 \times N^2$  matrix  $G_R$  is sampled from the real Ginibre ensemble, while  $C$  approximates  $X$  by a symmetric  $N \times N$  Gaussian orthogonal ensemble (GOE) matrix [40].

Matrices are normalized as  $\text{Tr} G_R G_R^\dagger = N^2$ , so that its spectrum covers uniformly a disk of radius 1, while  $\text{Tr} C^2 = N/4$  assures that its density forms the Wigner semicircle of radius 1. The scaling and parameters of the model follow from the normalization of the Kossakowski matrix (Supplemental Material [41]).

We approach spectral properties of the RM model (6) with the quaternionic extension of free probability [57–61]. Within this framework, we determine the border of the spectrum of  $\mathcal{L}'_D$  as given by the solution of the following equation involving a complex variable  $z$  [41]:

$$\text{Im}[z + G(z)] = 0, \quad (7)$$

with

$$G(z) = 2z - \frac{2z}{3\pi} \left[ (4 + z^2)E\left(\frac{4}{z^2}\right) + (4 - z^2)K\left(\frac{4}{z^2}\right) \right],$$




FIG. 4. Probability distributions  $P[\text{Re}(\ell'), \text{Im}(\ell')]$  of the rescaled eigenvalues,  $\ell' = N(\ell + 1)$ , from the spectrum of random Lindblad operators  $\mathcal{L}$ , Eq. (8), for  $N = 100$  and different values of the unitary component weight  $\alpha$ . (a),(b),(d) We present here the results of the sampling of the Kossakowski matrix from the Wishart ensemble, emphasizing that alternative generation procedures (Supplemental Material [41]) yield the same results. (c) Spectral distribution obtained with the random matrix model, Eq. (10). Each distribution was sampled with  $10^3$  realizations. Additional normalization of the densities is performed in order to keep maximal values of all distributions equal.

where  $E(k)$  and  $K(k)$  are complete elliptic integrals of the first and second kind, respectively. The results of the sampling are in perfect agreement with this border, see Figs. 2 and 3. Evaluation of the spectral density inside the “lemon” is a much harder task; it could potentially be performed with diagrammatic techniques [58].

*General case of random Lindbladians.*—Finally, we include the unitary component  $\mathcal{L}_U$  into the Lindblad operator  $\mathcal{L}$ . For that we use random Hamiltonian  $H$ , which is sampled from the Gaussian unitary ensemble (GUE). To compare the spectra of the general and the purely dissipative Lindblad operators, we normalize the Hamiltonian,  $\text{Tr} H^2 = 1/N$ , and introduce a parameter  $\alpha \geq 0$ , which weights contribution of the unitary component. The corresponding Lindbladians can be written as [see Eq. (4)]

$$\mathcal{L}(\rho) = -\frac{i\alpha}{\hbar}(H\rho - \rho H) + \Phi(\rho) - \frac{1}{2}[\Phi^\dagger(\mathbb{1})\rho + \rho\Phi^\dagger(\mathbb{1})]. \quad (8)$$

The sampled spectra of the operator  $\mathcal{L}' = N(\mathcal{L} + 1)$ , for different values of  $\alpha$ , are shown in Figs. 4(a), 4(b), and 4(d) (see also Supplemental Material [41]). Similar to the case of purely dissipative evolution, we find a perfect scale invariance starting from  $N \geq 50$ .

To capture the shape of the spectra with the RMT, we transform expression (8) into

$$\hat{\mathcal{L}} = \hat{\Phi} - \mathbb{1} \otimes \mathbb{1} - \left(\frac{1}{2}X + i\alpha H\right) \otimes \mathbb{1} + \mathbb{1} \otimes \left(\frac{1}{2}\bar{X} - i\alpha \bar{H}\right). \quad (9)$$

The spectrum of  $\hat{\mathcal{L}}$  can be explained by updating the matrix model (6),

$$\hat{\mathcal{L}}' \approx G_R - (W \otimes \mathbb{1} + \mathbb{1} \otimes \bar{W}), \quad (10)$$

where  $G_R$  is again taken from the real Ginibre ensemble, while the extended correction term  $W = C + i\alpha H'$  contains

now a random GOE matrix  $C$  and an anti-Hermitian term proportional to a GUE matrix  $H'$  normalized as  $\text{Tr}H'^2 = N$ . Spectral density of the RMT model for  $\alpha = 1$  is shown in Fig. 4(c). It reproduces spectral density of the corresponding Lindbladian ensemble (except of eigenvalue concentration at the real axis [37,39,55]).

Eigenvalues of  $W$  uniformly cover an ellipse with semiaxes  $1/\sqrt{1+4\alpha^2}$  and  $4\alpha^2/\sqrt{1+4\alpha^2}$ . Spectral density of  $\hat{\mathcal{L}}'$  is therefore a (classical) convolution of two uniform densities supported on these ellipses followed by free convolution with the Girko disk of unit radius; see Fig. 1(c). Contrary to the case of purely dissipative Lindbladians, it is not possible to find an analytically spectral boundary in this case. However, when  $\alpha = \frac{1}{2}$ , it immediately follows (since a convolution of two disks is a disk) that the spectral boundary is a circle. This is in full agreement with the results of the sampling; see Fig. 4(a).

*Conclusions.*—Universal spectral features of different ensembles of unitary evolution generators—that are Hamiltonians—are the main pillar of the quantum chaos (QC) theory [18]. A notion of an ensemble of random operators of quantum Markovian evolution is therefore a first step in generalization of QC to open quantum systems. Two next steps would be (i) establishing links between the idea of “typical Lindbladian” and physical models [62] and (ii) evaluation of spectral properties of steady states of random Lindbladians.

Finally, our approach works equally well in the classical limit, where continuous dynamics in the space of probability distributions is determined by random transition rate matrices. By using a free probabilities apparatus, we were able to derive analytically [56] the boundary of the eigenvalue distributions reported earlier in Ref. [34].

S. D., D. C., and K. Ż. appreciate the hospitality of the Center for Theoretical Physics of Complex Systems (IBS, South Korea) where this project was started. The numerical simulations were supported by the Russian Science Foundation Grant No. 19-72-20086 (S. D. and T. L.) and were performed on the supercomputer “Lomonosov-2” of the Moscow State University. D. C. and K. Ż. are grateful to Ł. Paweła and Z. Puchała for numerous discussions on random operations; they also acknowledge support from Narodowe Centrum Nauki under Grants No. 2018/30/A/ST2/00837 and No. 2015/18/A/ST2/00274, respectively. T. L. acknowledges support by the Basis Foundation (Grant No. 18-1-3-66-1). W. T. appreciates the financial support from the Polish Ministry of Science and Higher Education through the “Diamond Grant” 0225/DIA/2015/44 and the doctoral scholarship ETIUDA 2018/28/T/ST1/00470 from the National Science Center.

*Note added.*—One of the authors (W. T.) attended a talk by Tankut Can given at the conference in Yad

Hashmona (Israel), in which a parallel project on random Lindblad operators was presented. Recently, three other papers on related subjects have appeared [65–67].

- [1] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
- [2] U. Weiss, *Quantum Dissipative Systems* (World Scientific, Singapore, 2000).
- [3] Á. Rivas and S. F. Huelga, *Open Quantum Systems. An Introduction* (Springer, Heidelberg, 2011).
- [4] R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications*, Lecture Notes in Physics Vol. 286 (Springer, Berlin, 1998).
- [5] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of  $N$ -level systems, *J. Math. Phys. (N.Y.)* **17**, 821 (1976).
- [6] G. Lindblad, On the generators of quantum dynamical semigroups, *Commun. Math. Phys.* **48**, 119 (1976).
- [7] For an account on the history and importance of the GKSL equation, consult a recent review [8].
- [8] D. Chruściński and S. Pascazio, A brief history of the GKLS equation, *Open Syst. Inf. Dyn.* **24**, 1740001 (2017).
- [9] J. Marshall, L. C. Venuti, and P. Zanardi, Noise suppression via generalized-Markovian processes, *Phys. Rev. A* **96**, 052113 (2017).
- [10] D. Chruściński, C. Macchiavello, and S. Maniscalco, Detecting Non-Markovianity of Quantum Evolution via Spectra of Dynamical Maps, *Phys. Rev. Lett.* **118**, 080404 (2017).
- [11] S. Yu *et al.*, Experimental Investigation of Spectra of Dynamical Maps and Their Relation to Non-Markovianity, *Phys. Rev. Lett.* **120**, 060406 (2018).
- [12] M. L. Mehta, *Random Matrices*, 3rd ed. (Elsevier, Amsterdam, 2004).
- [13] E. P. Wigner, in Conference on Neutron Physics by Time of Flight Oak Ridge National Laboratory Report No. 2309, 1957, p. 59.
- [14] F. Dyson, Statistical theory of the energy levels of complex systems I-III, *J. Math. Phys. (N.Y.)* **3**, 140 (1962); Statistical theory of the energy levels of complex systems. II, *J. Math. Phys. (N.Y.)* **3**, 157 (1962); Statistical theory of the energy levels of complex systems. III, *J. Math. Phys. (N.Y.)* **3**, 166 (1962).
- [15] F. Dyson and M. L. Mehta, Statistical theory of the energy levels of complex systems IV-V, *J. Math. Phys. (N.Y.)* **4**, 701 (1963).
- [16] F. Dyson, A Brownian-motion model for the eigenvalues of a random matrix, *J. Math. Phys. (N.Y.)* **3**, 1191 (1962).
- [17] F. Dyson, The threefold way. Algebraic structure of symmetry groups and ensembles in quantum mechanics, *J. Math. Phys. (N.Y.)* **3**, 1199 (1962).
- [18] F. Haake, S. Gnutzmann, and M. Kuś, *Quantum Signatures of Chaos*, 4th ed. (Springer, Berlin, 2018).
- [19] H.-J. Stoeckmann, *Quantum Chaos: An Introduction* (Cambridge University Press, Cambridge, England, 1999).
- [20] D. Braun, *Dissipative Quantum Chaos and Decoherence* (Springer Tracts in Modern Physics, Berlin, 2001).
- [21] F. Evers and A. D. Mirlin, Anderson transitions, *Rev. Mod. Phys.* **80**, 1355 (2008).

- [22] A. Pal and D.A. Huse, Many-body localization phase transition, *Phys. Rev. B* **82**, 174411 (2010).
- [23] É. Brézin and V. Kazakov, Exactly solvable field theories of closed strings, *Phys. Lett. B* **236**, 144 (1990).
- [24] D. Gross and A. Migdal, Nonperturbative Two-Dimensional Quantum Gravity, *Phys. Rev. Lett.* **64**, 127 (1990).
- [25] É. Brézin, I. Itzykson, G. Parisi, and J.-B. Zuber, Planar diagrams, *Commun. Math. Phys.* **59**, 35 (1978).
- [26] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States* (Cambridge University Press, Cambridge, England, 2017).
- [27] W. Bruzda, V. Cappellini, H.-J. Sommers, and K. Życzkowski, Random quantum operations, *Phys. Lett. A* **373**, 320 (2009).
- [28] B. Collins and I. Nechita, Random matrix techniques in quantum information theory, *J. Math. Phys. (N.Y.)* **57**, 015215 (2016).
- [29] G. Aubrun and S. J. Szarek, *Alice and Bob Meet Banach: The Interface of Asymptotic Geometric Analysis and Quantum Information Theory* (AMS, Providence, 2017).
- [30] M. B. Hastings, Superadditivity of communication capacity using entangled inputs, *Nat. Phys.* **5**, 255 (2009).
- [31] A. A. Budini, Random Lindblad equations from complex environments, *Phys. Rev. E* **72**, 056106 (2005).
- [32] A. A. Budini and P. Grigolini, Non-Markovian nonstationary completely positive open-quantum-system dynamics, *Phys. Rev. A* **80**, 022103 (2009).
- [33] H. Schomerus, Random matrix approaches to open quantum systems, [arXiv:1610.05816](https://arxiv.org/abs/1610.05816).
- [34] C. Timm, Random transition-rate matrices for the master equation, *Phys. Rev. E* **80**, 021140 (2009).
- [35] W. Bruzda, M. Smaczyński, V. Cappellini, H.-J. Sommers, and K. Życzkowski, Universality of spectra for interacting quantum chaotic systems, *Phys. Rev. E* **81**, 066209 (2010).
- [36] J. Ginibre, Statistical ensembles of complex, quaternion and real matrices, *J. Math. Phys. (N.Y.)* **6**, 440 (1965).
- [37] P. J. Forrester and T. Nagao, Eigenvalue Statistics of the Real Ginibre Ensemble, *Phys. Rev. Lett.* **99**, 050603 (2007).
- [38] V. L. Girko, Circular law, *Theory Probab. Appl.* **29**, 694 (1983).
- [39] T. Tao and V. Vu, Random matrices: The circular law, *Commun. Contemp. Math.* **10**, 261 (2008).
- [40] I. Nechita, Z. Puchała, Ł. Paweła, and K. Życzkowski, Almost all quantum channels are equidistant, *J. Math. Phys. (N.Y.)* **59**, 052201 (2018).
- [41] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.123.140403> for more information, which includes Refs. [42–49].
- [42] I. Meyerov, A. Linirov, E. Kozinov, V. Volokitin, M. Ivanchenko, and S. Denisov, Unfolding quantum master equation into a system of linear equations: Computationally effective expansion over the basis of  $SU(N)$  generators, [arXiv:1812.11626](https://arxiv.org/abs/1812.11626).
- [43] S. T. Belinschi, P. Śniady, and R. Speicher, Eigenvalues of non-Hermitian random matrices and Brown measure of non-normal operators: Hermitian reduction and linearization method, *Linear Algebra Appl.* **537**, 48 (2018).
- [44] M. A. Nowak and W. Tarnowski, Spectra of large time-lagged correlation matrices from random matrix theory, *J. Stat. Mech.* (2017) 063405.
- [45] H. J. Sommers, A. Crisanti, H. Sompolinsky, and Y. Stein, Spectrum of Large Random Asymmetric Matrices, *Phys. Rev. Lett.* **60**, 1895 (1988).
- [46] J. T. Chalker and B. Mehlige, Eigenvector Statistics in Non-Hermitian Random Matrix Ensembles, *Phys. Rev. Lett.* **81**, 3367 (1998).
- [47] B. Mehlige and J. T. Chalker, Statistical properties of eigenvectors in non-Hermitian Gaussian random matrix ensembles, *J. Math. Phys. (N.Y.)* **41**, 3233 (2000).
- [48] R. A. Janik, W. Nörenberg, M. A. Nowak, G. Papp, and I. Zahed, Correlations of eigenvectors for non-Hermitian random-matrix models, *Phys. Rev. E* **60**, 2699 (1999).
- [49] J. A. Mingo and R. Speicher, *Free Probability and Random Matrices* (Springer Science, New York, 2017).
- [50] K. Życzkowski, K. A. Penson, I. Nechita, and B. Collins, Generating random density matrices, *J. Math. Phys. (N.Y.)* **52**, 062201 (2011).
- [51] L. A. Pastur and M. Shcherbina, *Eigenvalue Distribution of Large Random Matrices* (AMS Press, Providence, 2011).
- [52] In case of a nonsquare (rectangular)  $N^2 - 1 \times M$  Ginibre matrix  $Z$ , used to construct the Kossakowski matrix  $K = ZZ^\dagger$ , the spectral distribution remains the same up to the scaling. That is, the spectrum remains invariant under the transformation  $\mathcal{L}'_D = \sqrt{M}(\mathcal{L}_D + 1)$ ; see Ref. [41] for a more detailed discussion.
- [53] Recently, Lindbladians with operators  $V_m$  sampled from the Gaussian unitary ensemble were considered in Ref. [54]. This choice leads to dephasing-governed dynamics with the uniform asymptotic state  $\rho_{\text{inv}} = \mathbb{1}/N$ .
- [54] Z. Xu, L. P. García-Pintos, A. Chenu, and A. del Campo, Extreme Decoherence and Quantum Chaos, *Phys. Rev. Lett.* **122**, 014103 (2019).
- [55] A. Edelman, E. Kostlan, and M. Shub, How many eigenvalues of a random matrix are real? *J. Am. Math. Soc.* **7**, 247 (1994).
- [56] W. Tarnowski, S. Denisov, D. Chruściński, and K. Życzkowski (to be published).
- [57] R. A. Janik, M. A. Nowak, G. Papp, J. Wambach, and I. Zahed, Non-Hermitian random matrix models: Free random variable approach, *Phys. Rev. E* **55**, 4100 (1997).
- [58] R. A. Janik, M. A. Nowak, G. Papp, and I. Zahed, Non-Hermitian random matrix models, *Nucl. Phys.* **B501**, 603 (1997).
- [59] J. Feinberg and A. Zee, Non-Gaussian non-Hermitian random matrix theory: Phase transition and addition formalism, *Nucl. Phys.* **B501**, 643 (1997).
- [60] J. Feinberg and A. Zee, Non-Hermitian random matrix theory: Method of Hermitian reduction, *Nucl. Phys.* **B504**, 579 (1997).
- [61] A. Jarosz and M. A. Nowak, Random Hermitian versus random non-Hermitian operators—unexpected links, *J. Phys. A* **39**, 10107 (2006).
- [62] Open systems that exhibit “many-body localization-ergodicity” transition at the Hamiltonian limit are interesting candidates [63,64].

- [63] M. V. Medvedyeva, T. Prosen, and M. Žnidarič, Influence of dephasing on many-body localization, *Phys. Rev. B* **93**, 094205 (2016).
- [64] I. Vakulchyk, I. Yusipov, M. Ivanchenko, S. Flach, and S. Denisov, Signatures of many-body localization in steady states of open quantum systems, *Phys. Rev. B* **98**, 020202(R) (2018).
- [65] T. Can, Random Lindblad dynamics, [arXiv:1902.01442](#).
- [66] T. Can, V. Oganesyan, D. Orgad, and S. Gopalakrishnan, Spectral gaps and mid-gap states in random quantum master equations, [arXiv:1902.01414](#).
- [67] L. Sá, P. Ribeiro, and T. Prosen, Spectral and steady-state properties of random Liouvillians, [arXiv:1905.02155](#).

*Correction:* An incorrect affiliation for the fourth author was processed and has been fixed.