

Original Paper

Upregulation of Na⁺,Cl⁻-Coupled Betaine/ γ -Amino-Butyric Acid Transporter BGT1 by Tau Tubulin Kinase 2

Ahmad Almilaji Carlos Munoz Zohreh Hosseinzadeh Florian Lang

Department of Physiology, University of Tübingen, Tübingen, Germany

Key Words

Osmolyte transporter • Neurodegeneration • Kinase • Electrophysiology

Abstract

Background/Aims: The serine/threonine kinase Tau-tubulin-kinase 2 (TTBK2) is expressed in various tissues including kidney, liver and brain. Loss of function mutations of TTBK2 lead to autosomal dominant spinocerebellar ataxia type 11 (SCA11). Cell survival is fostered by cellular accumulation of organic osmolytes. Carriers accomplishing cellular accumulation of organic osmolytes include the Na⁺,Cl⁻-coupled betaine/ γ -amino-butyric acid transporter BGT1. The present study explored whether TTBK2 participates in the regulation of BGT1 activity.

Methods: Electrogenic transport of GABA was determined in *Xenopus* oocytes expressing BGT1 with or without wild-type TTBK2, truncated TTBK2[1-450] or kinase inactive mutants TTBK2-KD and TTBK2[1-450]-KD. **Results:** Coexpression of wild-type TTBK2, but not of TTBK2[1-450], TTBK2-KD or TTBK2[1-450]-KD, increased electrogenic GABA transport. Wild-type TTBK2 increased the maximal transport rate without significantly modifying affinity of the carrier. Coexpression of wild-type TTBK2 significantly delayed the decline of transport following inhibition of carrier insertion with brefeldin A, indicating that wild-type TTBK2 increased carrier stability in the cell membrane. **Conclusion:** Tau-tubulin-kinase 2 TTBK2 is a powerful stimulator of the osmolyte and GABA transporter BGT1.

Copyright © 2013 S. Karger AG, Basel

Introduction

The Tau tubulin Kinase 2 (TTBK2), a serine/threonine kinase [1] expressed in various tissues including the brain [2-4], heart [5], kidney [5], intestine [5] and tumor cells [6], contributes to the maintenance of neuron survival [2]. Accordingly, loss of function mutations of TTBK2 lead to autosomal dominant spinocerebellar ataxia type 11 (SCA11) [7]. Moreover,

TTBK2 may contribute to the resistance of kidney tumor cells and melanoma cells to therapy [6]. Gene targeted mice lacking functional TTBK2 die at embryonic day 10 [7].

The maintenance of cell survival is supported by cellular accumulation of organic osmolytes, which have been shown to counteract apoptosis [8]. Mechanisms contributing to cellular accumulation of organic osmolytes include Na⁺ coupled osmolyte transporters, such as the betaine/ γ -amino-butyric acid (GABA) transporter BGT1 (SLC6A12) [9, 10]. BGT1 is a member of the Na⁺,Cl⁻ coupled transporter superfamily accomplishing the transport of neurotransmitters (e.g. dopamine, GABA, serotonin and norepinephrine), amino acids (e.g. glycine) [11], creatine [12], and the organic osmolytes betaine [13] and taurine [14]. BGT1 is expressed in a wide variety of tissues including brain [15, 16], liver [16-18], kidney [16] and airways [19]. BGT1 expression is stimulated by osmotic cell shrinkage [20-22]. Activity of the carrier is regulated by Ca²⁺ [23], protein kinase C [24], prostaglandin E₂ [25, 26], cytoskeleton [27, 28] and ATP [29]. The carrier is upregulated following epileptic seizures [30]. However, the BGT1 deficient mouse does not develop seizures [31].

At least in theory, the protective effect of TTBK2 on neurons could involve modification of transport systems. As a matter of fact, a previous study revealed that TTBK2 upregulates the Na⁺ coupled glucose transporter SGLT1. Carriers with potentially protective effect include the BGT1. The present study thus explored, whether TTBK2 modifies the activity of BGT1. To this end, BGT1 was expressed in *Xenopus* oocytes with or without additional coexpression of wild-type TTBK2, truncated TTBK2^[1-450], kinase inactive full-length TTBK2-KD or kinase inactive truncated TTBK2-KD^[1-450]. BGT1 activity was determined from γ -amino-butyric acid (GABA)-induced current utilizing two-electrode voltage clamp.

Materials and Methods


Constructs

For generation of cRNA, constructs were used encoding wild-type human BGT1 (SLC6A12) [32, 33], wild-type human full-length TTBK2 and truncated mutant TTBK2^[1-450] containing the first 450 residues of the kinase. TTBK2^[1-450] has been identified in SCA11 patients [34] and leads to decreased kinase activity [7]. Further constructs used were the kinase inactive mutants TTBK2-KD and TTBK2^[1-450]-KD in which an aspartic acid at position 163 was replaced by alanine [D163A]. Constructs encoding wild type and mutated TTBK2 have all been kindly provided by Dario Alessi, University of Dundee. The cRNA was generated as described previously [35, 36].

Voltage clamp

For voltage clamp analysis, *Xenopus* oocytes were prepared as previously described [37]. On the day following oocyte preparation, *Xenopus* oocytes were injected with water, 15 ng of cRNA encoding BGT1 and 10 ng of cRNA encoding TTBK2^{WT}, TTBK2-KD, TTBK2^[1-450] or TTBK2^[1-450]-KD. The oocytes were maintained at 17°C in ND96 solution containing 88.5 mM NaCl, 2mM KCl, 1mM MgCl₂, 1.8mM CaCl₂, 5mM HEPES. Tetracycline (Sigma, 0.11mM), Ciprofloxacin (Sigma, 4 μ M), Gentamycin (Refobacin © 0.2mM) and Theophyllin (Euphyllong ©, 0.5mM) as well as sodium pyruvate (Sigma, 5mM) were added to ND96, pH was adjusted to 7.5 by addition of NaOH. The two-electrode voltage clamp recordings were performed at room temperature 4 days after the injection [38] at a holding potential of -70 mV. The data were filtered at 10 Hz and recorded with a DigidataA/D-D/A converter and ClampexV 9.2 software for data acquisition and analysis (Axon Instruments, Union City, CA, USA) [39, 40]. The control superfusate was ND96 containing 96 mM NaCl, 2 mM KCl, 1.8 mM CaCl₂, 1 mM MgCl₂ and 5 mM HEPES, pH was adjusted to 7.4 by addition of NaOH. The substrate γ -amino-butyric acid (GABA) was added to the solutions at the indicated concentrations. The flow rate of the superfusion was approximately 20 ml/min, and a complete exchange of the bath solution was reached within about 10 sec [41]. Where indicated, brefeldin A (5 μ M) was added to the solutions in order to test for alterations of BGT1 protein stability in the cell membrane. Brefeldin disrupts the organization of the microtubule and the actin network [42] and interferes with the function of Golgi-specific coat proteins involved in the regulation of membrane transport in the secretory pathway [43]. Treatment of *Xenopus*

Fig. 1. GABA induced current in *Xenopus* oocytes expressing BGT1 with or without wild-type TTBK2. A. Original tracings of GABA (1 mM) induced currents in *Xenopus* oocytes injected with water (a), or with cRNA encoding wild-type TTBK2 alone (b), BGT1 alone (c) or BGT1 together with wild-type TTBK2 (d). B. Arithmetic means \pm SEM (n = 5-18) of GABA (1 mM) induced current in *Xenopus* oocytes injected without (left bars) and with (right bars) BGT1, without (white bars) or with (black bars) cRNA encoding wild-type TTBK2. ** (p<0.01) indicates statistically significant difference from *Xenopus* oocytes expressing BGT1 alone.



oocytes with brefeldin A prevents insertion of new channel protein into the cell membrane and the decay of transporter activity could be taken as a measure of channel protein clearance from the cell membrane [44].

Statistical analysis

Data are provided as arithmetic means \pm SEM; n represents the number of oocytes investigated. All oocyte experiments were repeated for at least three batches of oocytes and in all repetitions qualitatively similar data were obtained [45]. All data were tested for significance by using ANOVA or t-test, as appropriate. Results with p< 0.05 were considered statistically significant.

Results

In order to explore, whether the Tau tubulin Kinase 2 (TTBK2) influences the activity of the Na⁺,Cl⁻-coupled betaine/GABA transporter BGT1, cRNA encoding BGT1 was injected into *Xenopus* oocytes without or with additional cRNA encoding TTBK2 and two-electrode voltage clamp experiments were performed to determine BGT1-mediated electrogenic GABA transport. As illustrated in Fig. 1, GABA (1 mM) did not induce an appreciable inward current in water injected *Xenopus* oocytes, indicating that *Xenopus* oocytes do not express appreciable endogenous electrogenic GABA transporters. Similarly, no significant GABA-induced current was observed in *Xenopus* oocytes expressing wild-type TTBK2 alone (Fig. 1). In *Xenopus* oocytes injected with cRNA encoding BGT1, however, addition of GABA (1 mM) was followed by an inward current, reflecting electrogenic entry of Na⁺, Cl⁻ and GABA. The GABA induced current was significantly enhanced by additional coexpression of wild-type TTBK2 (Fig. 1).

In contrast to wild-type TTBK2, truncated TTBK2^[1-450] did not significantly increase the GABA induced current in BGT1 expressing *Xenopus* oocytes. Moreover, neither coexpression of the full-length kinase inactive mutant TTBK2-KD nor coexpression of the truncated kinase inactive mutant TTBK2-KD^[1-450] significantly modified the GABA induced current in BGT1 expressing *Xenopus* oocytes (Fig. 3). Thus, kinase activity was required for the effect of TTBK2 on BGT1-induced currents.

Kinetic analysis of the GABA-induced currents in BGT1-expressing *Xenopus* oocytes (Fig. 2) yielded a maximal current of 15.7 \pm 1.3 nA (n = 7). Coexpression of wild-type TTBK2 significantly enhanced the maximal current to 35.0 \pm 1.4 nA (n = 6). Calculation of the GABA concentration required for the half-maximal current (K_M) yielded a value of 416 \pm 116 μ M

Fig. 2. Kinetics of GABA induced current in *Xenopus* oocytes expressing BGT1 with or without wild-type TTBK2. Arithmetic means \pm SEM (n = 6 - 7) of GABA induced currents as a function of GABA concentration in *Xenopus* oocytes expressing BGT1 alone (open circles) or BGT1 together with wild-type TTBK2 (closed circles). * (p<0.05) indicates statistically significant difference from *Xenopus* oocytes expressing BGT1 alone.





Fig. 3. GABA induced current in *Xenopus* oocytes expressing BGT1 with or without wild type TTBK2, truncated TTBK2^[1-450] mutant, kinase inactive mutant TTBK2-KD or truncated kinase inactive mutant TTBK2^[1-450]-KD. A. Original tracings of GABA (1 mM) induced currents measured in *Xenopus* oocytes injected with water (a), or expressing BGT1 without (b), or with (c) additional co-expression of wild-type TTBK2^{WT}, truncated wild-type TTBK2^[1-450] (d), full-length kinase inactive mutant TTBK2-KD (e) or truncated kinase inactive mutant TTBK2^[1-450]-KD (f). B. Arithmetic means \pm SEM (n = 11-15) of GABA-induced current (1 mM) in *Xenopus* oocytes injected with water (dotted bar), expressing BGT1 without (white bar) or with additional coexpression of wild-type TTBK2 (black bar), truncated wild-type TTBK2^[1-450] (1st grey bar), full-length kinase inactive mutant TTBK2-KD (2nd grey bar) or truncated kinase inactive mutant TTBK2^[1-450]-KD (3rd grey bar).




BGT1 without (white bar) or with additional coexpression of wild-type TTBK2 (black bar), truncated wild-type TTBK2^[1-450] (1st grey bar), full-length kinase inactive mutant TTBK2-KD (2nd grey bar) or truncated kinase inactive mutant TTBK2^[1-450]-KD (3rd grey bar). *** (p<0.001) indicates statistically significant difference from *Xenopus* oocytes expressing BGT1 alone. ### (p<0.001) indicates statistically significant difference from *Xenopus* oocytes coexpressing BGT1 and wild-type TTBK2.

(n = 7) in the absence, and a value of $413 \pm 58 \mu\text{M}$ (n = 6) in the presence of wild-type TTBK2. The K_M was not significantly different between *Xenopus* oocytes expressing BGT1 together with wild-type TTBK2 and *Xenopus* oocytes expressing BGT1 alone. Thus, wild-type TTBK2 did not significantly modify K_M but significantly increased the maximal current.

The enhanced BGT1 activity could have resulted from stimulation of carrier activity, from accelerated insertion of new carriers into the cell membrane, or from delayed clearance of carriers from the cell membrane. The effect on maximal transport rate without influence on substrate affinity pointed to increased protein abundance in the cell membrane. To discriminate between accelerated insertion and delayed clearance of carrier protein, the BGT1-expressing *Xenopus* oocytes were treated with 5 μM brefeldin A, which blocks the insertion of new carrier protein into the cell membrane. Following incubation with brefeldin A, the GABA induced current declined at a faster rate in oocytes expressing BGT1 alone

Fig. 4. Effects of Brefeldin A on GABA induced currents in *Xenopus* oocytes expressing BGT1 with or without wild-type TTBK2. Arithmetic means \pm SEM (n = 9-12) of GABA (1 mM) induced current in *Xenopus* oocytes injected with cRNA encoding BGT1, without (white bars) and with (black bars) wild-type TTBK2 in the absence (left bars) and presence of 5 μ M Brefeldin A for 12 hours (middle bars) or 24 hours (right bars) prior to the measurement. ** (p<0.01) indicates statistically significant difference from the absence of wild-type TTBK2.



than in oocytes expressing BGT1 together with wild-type TTBK2 (Fig. 4). The ratio of BGT1 induced current in oocytes expressing BGT1 together with wild-type TTBK2 over that of oocytes expressing BGT1 alone increased significantly from 223% \pm 40% (n = 12) prior to treatment with brefeldin A to 370% \pm 62% (n = 9-10) following a 12 hours treatment with brefeldin A and to 680% \pm 191% (n = 10-11) following a 24 hours treatment with brefeldin A. The observations suggest that wild-type TTBK2 delays the clearance of carrier protein from the cell membrane.

Discussion

The present observations reveal a completely novel regulator of the Na⁺/Cl⁻-coupled betaine/GABA transporter BGT1. The tau-tubulin-kinase 2 (TTBK2) up-regulates BGT1 activity, an effect at least in part due to stabilization of carrier protein in the cell membrane. The effect of TTBK2 requires functional kinase activity and is disrupted by mutations truncating the C-terminus noncatalytic domain of TTBK2 protein (TTBK2^[1-450]). The truncation has previously been shown to cause spinocerebellar degeneration [2].

In theory, TTBK2 could be effective by modifying the function of the microtubular network, which does participate in the regulation of transport processes [46-48]. Moreover, TTBK2 is critically important for the formation of cilia [49], which in turn are known to participate in the deranged transport regulation of polycystic kidney disease [50]. Clearly, additional experimental effort is needed to elucidate the mechanisms linking TTBK2 activity and carrier protein abundance in the cell membrane.

The regulation of BGT1 or a related carrier could well participate in the protection against neurodegeneration by TTBK2. Osmolyte transporters counteract cell shrinkage, which is known to parallel and facilitate suicidal cell death [51-60]. BGT1 participates in the cellular accumulation of organic osmolytes [9, 10], which are well known to foster cell survival [8, 26]. Thus, betaine has previously been shown to counteract apoptosis [61-64].

Carriers related to BGT1 include the transporters for creatine CreaT [12] and taurine TauT [14]. Those carriers may similarly be candidates for regulation by TTBK2. Genetic defects affecting the creatine transporter CreaT result in mental retardation with seizures [65-82]. Defective cellular taurine uptake by TauT fosters apoptosis [83-91].

At least in theory, TTBK2 sensitive BGT1 activity could modify neuronal function and survival in addition by transport of GABA. BGT1 mediates the cellular uptake of GABA, a function which was expected to decrease extracellular GABA concentration and thus to enhance neuronal excitation. Other carriers belonging to the superfamily of Na⁺/Cl⁻ coupled transporters include various transporters involved in the cellular uptake of further neurotransmitters, such as dopamine, serotonin, norepinephrine and amino

acids [11]. Additional experiments will be needed to explore whether TTBK2 influences neurotransmitter transport and thus neuronal function and survival by regulating one or more of those related transporters.

In conclusion, the present study discloses that TTBK2 is a powerful regulator of the Na⁺,Cl⁻-coupled betaine/GABA transporter BGT1 and thus participates in the regulation of cellular betaine and GABA uptake. The effect may contribute to the cytoprotective role of this widely expressed kinase.

Acknowledgements

The authors gratefully acknowledge the meticulous preparation of the manuscript by T. Loch and L. Subasic, and the technical support by E. Faber. This work was supported by the DFG (GRK 1302/1, SE 1077/3 and SFB 773) and Open Access Publishing Fund of Tuebingen University.

References

- 1 Kitano-Takahashi M, Morita H, Kondo S, Tomizawa K, Kato R, Tanio M, Shirota Y, Takahashi H, Sugio S, Kohno T: Expression, purification and crystallization of a human tau-tubulin kinase 2 that phosphorylates tau protein. *Acta Crystallogr Sect F Struct Biol Cryst Commun* 2007;63:602-604.
- 2 Houlden H, Johnson J, Gardner-Thorpe C, Lashley T, Hernandez D, Worth P, Singleton AB, Hilton DA, Holton J, Revesz T, Davis MB, Giunti P, Wood NW: Mutations in TTBK2, encoding a kinase implicated in tau phosphorylation, segregate with spinocerebellar ataxia type 11. *Nat Genet* 2007;39:1434-1436.
- 3 Tomizawa K, Omori A, Ohtake A, Sato K, Takahashi M: Tau-tubulin kinase phosphorylates tau at Ser-208 and Ser-210, sites found in paired helical filament-tau. *FEBS Lett* 2001;492:221-227.
- 4 Takahashi M, Tomizawa K, Ishiguro K, Takamatsu M, Fujita SC, Imahori K: Involvement of tau protein kinase I in paired helical filament-like phosphorylation of the juvenile tau in rat brain. *J Neurochem* 1995;64:1759-1768.
- 5 Alesutan IS, Sopjani M, Dörmaku-Sopjani M, Munoz C, Voelkl J, Lang F: Upregulation of Na⁺-coupled glucose transporter SGLT1 by Tau tubulin Kinase 2. *Cell Physiol Biochem* 2012;30:458-465.
- 6 Bender C, Ullrich A: PRKX, TTBK2 and RSK4 expression causes Sunitinib resistance in kidney carcinoma- and melanoma-cell lines. *Int J Cancer* 2012;131:E45-E55.
- 7 Bouskila M, Esoof N, Gay L, Fang EH, Deak M, Begley MJ, Cantley LC, Prescott A, Storey KG, Alessi DR: TTBK2 kinase substrate specificity and the impact of spinocerebellar-ataxia-causing mutations on expression, activity, localization and development. *Biochem J* 2011;437:157-167.
- 8 Alfieri RR, Cavazzoni A, Petronini PG, Bonelli MA, Caccamo AE, Borghetti AF, Wheeler KP: Compatible osmolytes modulate the response of porcine endothelial cells to hypertonicity and protect them from apoptosis. *J Physiol* 2002;540:499-508.
- 9 Handler JS, Kwon HM: Regulation of renal cell organic osmolyte transport by tonicity. *Am J Physiol* 1993;265:C1449-C1455.
- 10 Kempson SA, Montrose MH: Osmotic regulation of renal betaine transport: transcription and beyond. *Pflugers Arch* 2004;449:227-234.
- 11 Christie DL: Functional insights into the creatine transporter. *Subcell Biochem* 2007;46:99-118.
- 12 Nash SR, Giros B, Kingsmore SF, Rochelle JM, Suter ST, Gregor P, Seldin MF, Caron MG: Cloning, pharmacological characterization, and genomic localization of the human creatine transporter. *Receptors Channels* 1994;2:165-174.
- 13 Takenaka M, Bagnasco SM, Preston AS, Uchida S, Yamauchi A, Kwon HM, Handler JS: The canine betaine gamma-amino-n-butyric acid transporter gene: diverse mRNA isoforms are regulated by hypertonicity and are expressed in a tissue-specific manner. *Proc Natl Acad Sci U S A* 1995;92:1072-1076.

- 14 Uchida S, Kwon HM, Yamauchi A, Preston AS, Marumo F, Handler JS: Molecular cloning of the cDNA for an MDCK cell Na(+)- and Cl(-)-dependent taurine transporter that is regulated by hypertonicity. *Proc Natl Acad Sci U S A* 1992;89:8230-8234.
- 15 Madsen KK, White HS, Schousboe A: Neuronal and non-neuronal GABA transporters as targets for antiepileptic drugs. *Pharmacol Ther* 2010;125:394-401.
- 16 Zhou Y, Holmseth S, Hua R, Lehre AC, Olofsson AM, Poblete-Naredo I, Kempson SA, Danbolt NC: The betaine-GABA transporter (BGT1, slc6a12) is predominantly expressed in the liver and at lower levels in the kidneys and at the brain surface. *Am J Physiol Renal Physiol* 2012;302:F316-F328.
- 17 Peters-Regehr T, Bode JG, Kubitz R, Haussinger D: Organic osmolyte transport in quiescent and activated rat hepatic stellate cells (Ito cells). *Hepatology* 1999;29:173-180.
- 18 Weik C, Warskulat U, Bode J, Peters-Regehr T, Haussinger D: Compatible organic osmolytes in rat liver sinusoidal endothelial cells. *Hepatology* 1998;27:569-575.
- 19 Zaidi S, Gallos G, Yim PD, Xu D, Sonett JR, Panettieri RA Jr, Gerthoffer W, Emala CW: Functional expression of gamma-amino butyric acid transporter 2 in human and guinea pig airway epithelium and smooth muscle. *Am J Respir Cell Mol Biol* 2011;45:332-339.
- 20 Kempson SA: Differential activation of system A and betaine/GABA transport in MDCK cell membranes by hypertonic stress. *Biochim Biophys Acta* 1998;1372:117-123.
- 21 Miyakawa H, Rim JS, Handler JS, Kwon HM: Identification of the second tonicity-responsive enhancer for the betaine transporter (BGT1) gene. *Biochim Biophys Acta* 1999;1446:359-364.
- 22 Nadkarni V, Gabbay KH, Bohren KM, Sheikh-Hamad D: Osmotic response element enhancer activity. Regulation through p38 kinase and mitogen-activated extracellular signal-regulated kinase kinase. *J Biol Chem* 1999;274:20185-20190.
- 23 Kempson SA, Edwards JM, Sturek M: Inhibition of the renal betaine transporter by calcium ions. *Am J Physiol Renal Physiol* 2006;291:F305-F313.
- 24 Massari S, Vanoni C, Longhi R, Rosa P, Pietrini G: Protein kinase C-mediated phosphorylation of the BGT1 epithelial gamma-aminobutyric acid transporter regulates its association with LIN7 PDZ proteins: a post-translational mechanism regulating transporter surface density. *J Biol Chem* 2005;280:7388-7397.
- 25 Neuhofer W, Steinert D, Fraek ML, Beck FX: Prostaglandin E2 stimulates expression of osmoprotective genes in MDCK cells and promotes survival under hypertonic conditions. *J Physiol* 2007;583:287-297.
- 26 Moeckel GW, Zhang L, Fogo AB, Hao CM, Pozzi A, Breyer MD: COX2 activity promotes organic osmolyte accumulation and adaptation of renal medullary interstitial cells to hypertonic stress. *J Biol Chem* 2003;278:19352-19357.
- 27 Basham JC, Chabrierie A, Kempson SA: Hypertonic activation of the renal betaine/GABA transporter is microtubule dependent. *Kidney Int* 2001;59:2182-2191.
- 28 Bricker JL, Chu S, Kempson SA: Disruption of F-actin stimulates hypertonic activation of the BGT1 transporter in MDCK cells. *Am J Physiol Renal Physiol* 2003;284:F930-F937.
- 29 Kempson SA, Edwards JM, Osborn A, Sturek M: Acute inhibition of the betaine transporter by ATP and adenosine in renal MDCK cells. *Am J Physiol Renal Physiol* 2008;295:F108-F117.
- 30 Rowley NM, Smith MD, Lamb JG, Schousboe A, White HS: Hippocampal betaine/GABA transporter mRNA expression is not regulated by inflammation or dehydration post-status epilepticus. *J Neurochem* 2011;117:82-90.
- 31 Lehre AC, Rowley NM, Zhou Y, Holmseth S, Guo C, Holen T, Hua R, Laake P, Olofsson AM, Poblete-Naredo I, Rusakov DA, Madsen KK, Clausen RP, Schousboe A, White HS, Danbolt NC: Deletion of the betaine-GABA transporter (BGT1; slc6a12) gene does not affect seizure thresholds of adult mice. *Epilepsy Res* 2011;95:70-81.
- 32 Matskevitch I, Wagner CA, Stegen C, Broer S, Noll B, Risler T, Kwon HM, Handler JS, Waldegger S, Busch AE, Lang F: Functional characterization of the Betaine/gamma-aminobutyric acid transporter BGT-1 expressed in *Xenopus* oocytes. *J Biol Chem* 1999;274:16709-16716.
- 33 Munoz C, Sopjani M, Dermaku-Sopjani M, Almilaji A, Foller M, Lang F: Downregulation of the osmolyte transporters SMIT and BGT1 by AMP-activated protein kinase. *Biochem Biophys Res Commun* 2012;422:358-362.

- 34 Bauer P, Stevanin G, Beetz C, Synofzik M, Schmitz-Hubsch T, Wullner U, Berthier E, Ollagnon-Roman E, Riess O, Forlani S, Mundwiler E, Durr A, Schols L, Brice A: Spinocerebellar ataxia type 11 (SCA11) is an uncommon cause of dominant ataxia among French and German kindreds. *J Neurol Neurosurg Psychiatry* 2010;81:1229-1232.
- 35 Bohmer C, Sopjani M, Klaus F, Lindner R, Laufer J, Jeyaraj S, Lang F, Palmada M: The serum and glucocorticoid inducible kinases SGK1-3 stimulate the neutral amino acid transporter SLC6A19. *Cell Physiol Biochem* 2010;25:723-732.
- 36 Mohamed MR, Alesutan I, Foller M, Sopjani M, Bress A, Baur M, Salama RH, Bakr MS, Mohamed MA, Blin N, Lang F, Pfister M: Functional analysis of a novel I71N mutation in the GJB2 gene among Southern Egyptians causing autosomal recessive hearing loss. *Cell Physiol Biochem* 2010;26:959-966.
- 37 Rexhepaj R, Dermaku-Sopjani M, Gehring EM, Sopjani M, Kempe DS, Foller M, Lang F: Stimulation of electrogenic glucose transport by glycogen synthase kinase 3. *Cell Physiol Biochem* 2010;26:641-646.
- 38 Eckey K, Strutz-Seebohm N, Katz G, Fuhrmann G, Henrion U, Pott L, Linke WA, Arad M, Lang F, Seebohm G: Modulation of human ether a gogo related channels by CASQ2 contributes to etiology of catecholaminergic polymorphic ventricular tachycardia (CPVT). *Cell Physiol Biochem* 2010;26:503-512.
- 39 Alesutan IS, Ureche ON, Laufer J, Klaus F, Zurn A, Lindner R, Strutz-Seebohm N, Tavare JM, Boehmer C, Palmada M, Lang UE, Seebohm G, Lang F: Regulation of the glutamate transporter EAAT4 by PIKfyve. *Cell Physiol Biochem* 2010;25:187-194.
- 40 Hosseinzadeh Z, Bhavsar SK, Sopjani M, Alesutan I, Saxena A, Dermaku-Sopjani M, Lang F: Regulation of the glutamate transporters by JAK2. *Cell Physiol Biochem* 2011;28:693-702.
- 41 Dermaku-Sopjani M, Sopjani M, Saxena A, Shojaiefard M, Bogatikov E, Alesutan I, Eichenmuller M, Lang F: Downregulation of NaPi-IIa and NaPi-IIb Na-coupled phosphate transporters by coexpression of Klotho. *Cell Physiol Biochem* 2011;28:251-258.
- 42 Alvarez C, Sztul ES: Brefeldin A (BFA) disrupts the organization of the microtubule and the actin cytoskeletons. *Eur J Cell Biol* 1999;78:1-14.
- 43 Hunziker W, Whitney JA, Mellman I: Brefeldin A and the endocytic pathway. Possible implications for membrane traffic and sorting. *FEBS Lett* 1992;307:93-96.
- 44 Staub O, Gautschi I, Ishikawa T, Breitschopf K, Ciechanover A, Schild L, Rotin D: Regulation of stability and function of the epithelial Na⁺ channel (ENaC) by ubiquitination. *EMBO J* 1997;16:6325-6336.
- 45 Strutz-Seebohm N, Pusch M, Wolf S, Stoll R, Tapken D, Gerwert K, Attali B, Seebohm G: Structural basis of slow activation gating in the cardiac I Ks channel complex. *Cell Physiol Biochem* 2011;27:443-452.
- 46 Cheng J, Wang H, Guggino WB: Regulation of cystic fibrosis transmembrane regulator trafficking and protein expression by a Rho family small GTPase TC10. *J Biol Chem* 2005;280:3731-3739.
- 47 Goswami C, Dreger M, Jahnle R, Bogen O, Gillen C, Hucho F: Identification and characterization of a Ca²⁺-sensitive interaction of the vanilloid receptor TRPV1 with tubulin. *J Neurochem* 2004;91:1092-1103.
- 48 Zheng Y, Sarr MG: Translocation of transfected GLUT2 to the apical membrane in rat intestinal IEC-6 cells. *Dig Dis Sci* 2012;57:1203-1212.
- 49 Goetz SC, Liem KF Jr, Anderson KV: The spinocerebellar ataxia-associated gene Tau tubulin kinase 2 controls the initiation of ciliogenesis. *Cell* 2012;151:847-858.
- 50 Simons M, Walz G: Polycystic kidney disease: cell division without a c(l)ue? *Kidney Int* 2006;70:854-864.
- 51 Lang F, Busch GL, Ritter M, Volkl H, Waldegger S, Gulbins E, Haussinger D: Functional significance of cell volume regulatory mechanisms. *Physiol Rev* 1998;78:247-306.
- 52 Andersen AD, Bentzen BH, Salling H, Klingberg H, Kannevorff M, Grunnet M, Pedersen SF: The cardioprotective effect of brief acidic reperfusion after ischemia in perfused rat hearts is not mimicked by inhibition of the Na⁺/H⁺ exchanger NHE1. *Cell Physiol Biochem* 2011;28:13-24.
- 53 Bortner CD, Cidlowski JA: Life and death of lymphocytes: a volume regulation affair. *Cell Physiol Biochem* 2011;28:1079-1088.
- 54 Bozeat ND, Xiang SY, Ye LL, Yao TY, Duan ML, Burkin DJ, Lamb FS, Duan DD: Activation of volume regulated chloride channels protects myocardium from ischemia/reperfusion damage in second-window ischemic preconditioning. *Cell Physiol Biochem* 2011;28:1265-1278.
- 55 Gatidis S, Zelenak C, Fajol A, Lang E, Jilani K, Michael D, Qadri SM, Lang F: p38 MAPK activation and function following osmotic shock of erythrocytes. *Cell Physiol Biochem* 2011;28:1279-1286.
- 56 Haussinger D, Reinehr R: Osmotic regulation of bile acid transport, apoptosis and proliferation in rat liver. *Cell Physiol Biochem* 2011;28:1089-1098.

- 57 Hoffmann EK: Ion channels involved in cell volume regulation: effects on migration, proliferation, and programmed cell death in non adherent EAT cells and adherent ELA cells. *Cell Physiol Biochem* 2011;28:1061-1078.
- 58 Lambert IH, Hansen DB: Regulation of taurine transport systems by protein kinase CK2 in mammalian cells. *Cell Physiol Biochem* 2011;28:1099-1110.
- 59 Lewis R, Feetham CH, Barrett-Jolley R: Cell volume regulation in chondrocytes. *Cell Physiol Biochem* 2011;28:1111-1122.
- 60 Lionetto MG, Giordano ME, Calisi A, Caricato R, Hoffmann E, Schettino T: Role of BK channels in the apoptotic volume decrease in native eel intestinal cells. *Cell Physiol Biochem* 2010;25:733-744.
- 61 Graf D, Kurz AK, Reinehr R, Fischer R, Kircheis G, Haussinger D: Prevention of bile acid-induced apoptosis by betaine in rat liver. *Hepatology* 2002;36:829-839.
- 62 Horio M, Ito A, Matsuoka Y, Moriyama T, Orita Y, Takenaka M, Imai E: Apoptosis induced by hypertonicity in Madin Darley canine kidney cells: protective effect of betaine. *Nephrol Dial Transplant* 2001;16:483-490.
- 63 Ji C, Kaplowitz N: Betaine decreases hyperhomocysteinemia, endoplasmic reticulum stress, and liver injury in alcohol-fed mice. *Gastroenterology* 2003;124:1488-1499.
- 64 Kharbanda KK, Rogers DD, Mailliard ME, Siford GL, Barak AJ, Beckenhauer HC, Sorrell MF, Tuma DJ: Role of elevated S-adenosylhomocysteine in rat hepatocyte apoptosis: protection by betaine. *Biochem Pharmacol* 2005;70:1883-1890.
- 65 Alcaide P, Rodriguez-Pombo P, Ruiz-Sala P, Ferrer I, Castro P, Ruiz MY, Merinero B, Ugarte M: A new case of creatine transporter deficiency associated with mild clinical phenotype and a novel mutation in the SLC6A8 gene. *Dev Med Child Neurol* 2010;52:215-217.
- 66 Alcaide P, Merinero B, Ruiz-Sala P, Richard E, Navarrete R, Arias A, Ribes A, Artuch R, Campistol J, Ugarte M, Rodriguez-Pombo P: Defining the pathogenicity of creatine deficiency syndrome. *Hum Mutat* 2011;32:282-291.
- 67 Ardon O, Amat di San FC, Salomons GS, Longo N: Creatine transporter deficiency in two half-brothers. *Am J Med Genet A* 2010;152A:1979-1983.
- 68 Battini R, Chilosi A, Mei D, Casarano M, Alessandri MG, Leuzzi V, Ferretti G, Tosetti M, Bianchi MC, Cioni G: Mental retardation and verbal dyspraxia in a new patient with de novo creatine transporter (SLC6A8) mutation. *Am J Med Genet A* 2007;143A:1771-1774.
- 69 Battini R, Chilosi AM, Casarano M, Moro F, Comparini A, Alessandri MG, Leuzzi V, Tosetti M, Cioni G: Language disorder with mild intellectual disability in a child affected by a novel mutation of SLC6A8 gene. *Mol Genet Metab* 2011;102:153-156.
- 70 Braissant O, Beard E, Torrent C, Henry H: Dissociation of AGAT, GAMT and SLC6A8 in CNS: relevance to creatine deficiency syndromes. *Neurobiol Dis* 2010;37:423-433.
- 71 Braissant O, Henry H, Beard E, Uldry J: Creatine deficiency syndromes and the importance of creatine synthesis in the brain. *Amino Acids* 2011;40:1315-1324.
- 72 Hahn KA, Salomons GS, Tackels-Horne D, Wood TC, Taylor HA, Schroer RJ, Lubs HA, Jakobs C, Olson RL, Holden KR, Stevenson RE, Schwartz CE: X-linked mental retardation with seizures and carrier manifestations is caused by a mutation in the creatine-transporter gene (SLC6A8) located in Xq28. *Am J Hum Genet* 2002;70:1349-1356.
- 73 Jensen LR, Chen W, Moser B, Lipkowitz B, Schroeder C, Musante L, Tzschach A, Kalscheuer VM, Meloni I, Raynaud M, van Esch H, Chelly J, de Brouwer AP, Hackett A, van der HS, Henn W, Gecz J, Riess O, Bonin M, Reinhardt R, Ropers HH, Kuss AW: Hybridisation-based resequencing of 17 X-linked intellectual disability genes in 135 patients reveals novel mutations in ATRX, SLC6A8 and PQBP1. *Eur J Hum Genet* 2011;19:717-720.
- 74 Longo N, Ardon O, Vanzo R, Schwartz E, Pasquali M: Disorders of creatine transport and metabolism. *Am J Med Genet C Semin Med Genet* 2011;157:72-78.
- 75 Mancardi MM, Caruso U, Schiaffino MC, Baglietto MG, Rossi A, Battaglia FM, Salomons GS, Jakobs C, Zara F, Veneselli E, Gaggero R: Severe epilepsy in X-linked creatine transporter defect (CRTR-D). *Epilepsia* 2007;48:1211-1213.
- 76 Mercimek-Mahmutoglu S, Connolly MB, Poskitt KJ, Horvath GA, Lowry N, Salomons GS, Casey B, Sinclair G, Davis C, Jakobs C, Stockler-Ipsiroglu S: Treatment of intractable epilepsy in a female with SLC6A8 deficiency. *Mol Genet Metab* 2010;101:409-412.

- 77 Puusepp H, Kall K, Salomons GS, Talvik I, Mannamaa M, Rein R, Jakobs C, Ounap K: The screening of SLC6A8 deficiency among Estonian families with X-linked mental retardation. *J Inher Metab Dis* 2009; doi:<http://link.springer.com/article/10.1007/s10545-008-1063-y>.
- 78 Rosenberg EH, Martinez MC, Betsalel OT, van Dooren SJ, Fernandez M, Jakobs C, deGrauw TJ, Kleefstra T, Schwartz CE, Salomons GS: Functional characterization of missense variants in the creatine transporter gene (SLC6A8): improved diagnostic application. *Hum Mutat* 2007;28:890-896.
- 79 Salomons GS, van Dooren SJ, Verhoeven NM, Marsden D, Schwartz C, Cecil KM, deGrauw TJ, Jakobs C: X-linked creatine transporter defect: an overview. *J Inher Metab Dis* 2003;26:309-318.
- 80 Skelton MR, Schaefer TL, Graham DL, deGrauw TJ, Clark JF, Williams MT, Vorhees CV: Creatine transporter (CrT; SLC6A8) knockout mice as a model of human CrT deficiency. *PLoS One* 2011;6:e16187.
- 81 Stockler S, Schutz PW, Salomons GS: Cerebral creatine deficiency syndromes: clinical aspects, treatment and pathophysiology. *Subcell Biochem* 2007;46:149-166.
- 82 van de Kamp JM, Mancini GM, Pouwels PJ, Betsalel OT, van Dooren SJ, de K, I, Steenweg ME, Jakobs C, van der Knaap MS, Salomons GS: Clinical features and X-inactivation in females heterozygous for creatine transporter defect. *Clin Genet* 2011;79:264-272.
- 83 Bachmann MF: Taurine: energy drink for T cells. *Eur J Immunol* 2012;42:819-821.
- 84 Han X, Yue J, Chesney RW: Functional TauT protects against acute kidney injury. *J Am Soc Nephrol* 2009;20:1323-1332.
- 85 Hansen DB, Guerra B, Jacobsen JH, Lambert IH: Regulation of taurine homeostasis by protein kinase CK2 in mouse fibroblasts. *Amino Acids* 2011;40:1091-1106.
- 86 Ito T, Kimura Y, Uozumi Y, Takai M, Muraoka S, Matsuda T, Ueki K, Yoshiyama M, Ikawa M, Okabe M, Schaffer SW, Fujio Y, Azuma J: Taurine depletion caused by knocking out the taurine transporter gene leads to cardiomyopathy with cardiac atrophy. *J Mol Cell Cardiol* 2008;44:927-937.
- 87 Nishimura T, Sai Y, Fujii J, Muta M, Iizasa H, Tomi M, Deureh M, Kose N, Nakashima E: Roles of TauT and system A in cytoprotection of rat syncytiotrophoblast cell line exposed to hypertonic stress. *Placenta* 2010;31:1003-1009.
- 88 Tabuchi H, Sugiyama T, Tanaka S, Tainaka S: Overexpression of taurine transporter in Chinese hamster ovary cells can enhance cell viability and product yield, while promoting glutamine consumption. *Biotechnol Bioeng* 2010;107:998-1003.
- 89 Tastesen HS, Holm JB, Moller J, Poulsen KA, Moller C, Sturup S, Hoffmann EK, Lambert IH: Pinpointing differences in cisplatin-induced apoptosis in adherent and non-adherent cancer cells. *Cell Physiol Biochem* 2010;26:809-820.
- 90 Zeng K, Xu H, Mi M, Chen K, Zhu J, Yi L, Zhang T, Zhang Q, Yu X: Effects of taurine on glial cells apoptosis and taurine transporter expression in retina under diabetic conditions. *Neurochem Res* 2010;35:1566-1574.
- 91 Zhang LY, Zhou YY, Chen F, Wang B, Li J, Deng YW, Liu WD, Wang ZG, Li YW, Li DZ, Lv GH, Yin BL: Taurine inhibits serum deprivation-induced osteoblast apoptosis via the taurine transporter/ERK signaling pathway. *Braz J Med Biol Res* 2011;44:618-623.