

University of New Mexico

UNM Digital Repository

Faculty and Staff Publications

Mathematics

2016

USES OF SAMPLING TECHNIQUES & INVENTORY CONTROL WITH CAPACITY CONSTRAINTS

Florentin Smarandache

University of New Mexico, smarand@unm.edu

Sachin Malik

Neeraj Kumar

Follow this and additional works at: https://digitalrepository.unm.edu/math_fsp



Part of the [Agricultural and Resource Economics Commons](#), [Economics Commons](#), [Food Studies Commons](#), [International and Area Studies Commons](#), [Mathematics Commons](#), [Public Affairs, Public Policy and Public Administration Commons](#), and the [Science and Technology Studies Commons](#)

Recommended Citation

Sachin Malik, Neeraj Kumar, F. Smarandache (eds.) USES OF SAMPLING TECHNIQUES & INVENTORY CONTROL WITH CAPACITY CONSTRAINTS. Brussels: Pons Editions, 2016.

This Book is brought to you for free and open access by the Mathematics at UNM Digital Repository. It has been accepted for inclusion in Faculty and Staff Publications by an authorized administrator of UNM Digital Repository. For more information, please contact amywinter@unm.edu, lsloane@salud.unm.edu, sarahrk@unm.edu.

The cover features a decorative border on the left and right sides, consisting of a vertical strip of hands in various colors (red, orange, yellow, green, blue) and white doves. The background is a solid orange color. Two horizontal orange bars are positioned above and below the editors' names.

Sachin Malik

Neeraj Kumar

Florentin Smarandache

(Editors)

USES OF SAMPLING
TECHNIQUES
&
INVENTORY CONTROL
WITH CAPACITY
CONSTRAINTS

Sachin Malik, Neeraj Kumar, Florentin Smarandache

Editors

USES OF SAMPLING TECHNIQUES
&
INVENTORY CONTROL WITH CAPACITY
CONSTRAINTS

Peer-Reviewers:

Nassim Abbas, Communicating and Intelligent System Engineering Laboratory, Faculty of Electronics and Computer Science University of Science and Technology Houari Boumediene 32, El Alia, Bab Ezzouar, 16111, Algiers, Algeria.

Mumtaz Ali, Department of Mathematics, Quaid-i-Azam University, Islamabad, 44000, Pakistan.

Said Broumi, Faculty of letters and Humanities, Hay El Baraka Ben M'sik Casablanca, B.P. 7951, University of Hassan II, Casablanca, Morocco.

Jun Ye, Department of Electrical and Information Engineering, Shaoxing University, No. 508 Huancheng West Road, Shaoxing, Zhejiang Province 312000, P. R. China.

**USES OF SAMPLING TECHNIQUES
&
INVENTORY CONTROL WITH CAPACITY
CONSTRAINTS**

Editors:

Sachin Malik, Neeraj Kumar

Department of Mathematics, SRM University
Delhi NCR, Sonapat - 131029, India

Florentin Smarandache

Department of Mathematics, University of New Mexico,
Gallup, NM 87301, USA



Pons Editions

Brussels, 2016

ISBN 978-1-59973-484-2



Pons asbl
Quai du Batelage no. 5
1000 Bruxelles
Belgium

Contents

	<i>Preface</i>	7
Rajesh Singh, Sachin Malik, Florentin Smarandache	An Improved Suggestion in Stratified Random Sampling Using Two Auxiliary Variables	9
Rajesh Singh, Sachin Malik, Florentin Smarandache	Some Ratio and Product Estimators Using Known Value Of Population Parameters	19
Rajesh Singh, Sachin Malik	An Unbiased Estimator For Estimating Population Mean In Simple Random Sampling Auxiliary Attribute	30
Neeraj Kumar, Sanjey Kumar, Florentin Smarandache	Two Warehouse Fuzzy Inventory Model With K-Release Rule	53
Sanjey Kumar, Pallavi Agarwa, Neeraj Kumar	A Two-Warehouse Inventory Model for Deteriorating Items with stock Dependent Demand, Inflation and Genetic Algorithm	69

Preface

The main aim of the present book is to suggest some improved estimators using auxiliary and attribute information in case of simple random sampling and stratified random sampling and some inventory models related to capacity constraints.

This volume is a collection of five papers, written by six co-authors (listed in the order of the papers): Dr. Rajesh Singh, Dr. Sachin Malik, Dr. Florentin Smarandache, Dr. Neeraj Kumar, Mr. Sanjey Kumar & Pallavi Agarwal.

In the first chapter authors suggest an estimator using two auxiliary variables in stratified random sampling for estimating population mean. In second chapter they proposed a family of estimators for estimating population means using known value of some population parameters. In Chapter third an almost unbiased estimator using known value of some population parameter(s) with known population proportion of an auxiliary variable has been used. In Chapter four the authors investigate a fuzzy economic order quantity model for two storage facility. The demand, holding cost, ordering cost, storage capacity of the own - warehouse are taken as trapezoidal fuzzy numbers. And in Chapter five a two-warehouse inventory model deals with deteriorating items, with stock dependent demand rate and model affected by inflation under the pattern of time value of money over a finite planning horizon. Shortages are allowed and partially backordered depending on the waiting time for the next replenishment. The purpose of this model is to minimize the total inventory cost by using the genetic algorithm.

This book will be helpful for the researchers and students who are working in the field of sampling techniques and inventory control.

An Improved Suggestion in Stratified Random Sampling Using Two Auxiliary Variables

Rajesh Singh, Sachin Malik* and Florentin Smarandache**

*Department of Mathematics, SRM University

Delhi NCR, Sonapat- 131029, India

Department of Statistics, Banaras Hindu University

Varanasi-221005, India

**Department of Mathematics, University of New Mexico

Gallup, NM 87301, USA

*Corresponding Author, sachinkurava999@gmail.com

Abstract

In this paper, we suggest an estimator using two auxiliary variables in stratified random sampling following Malik and Singh [12]. The propose estimator has an improvement over mean per unit estimator as well as some other considered estimators. Expressions for bias and MSE of the estimator are derived up to first degree of approximation. Moreover, these theoretical findings are supported by a numerical example with original data.

Key words: Study variable, auxiliary variable, stratified random sampling, bias and mean squared error.

1. Introduction

The problem of estimating the population mean in the presence of an auxiliary variable has been widely discussed in finite population sampling literature. Out of many ratio, product and regression methods of estimation are good examples in this context. Diana [2] suggested a class of estimators of the population mean using one auxiliary variable in the stratified random sampling and examined the MSE of the estimators up to the k^{th} order of approximation. Kadilar and Cingi [3], Singh et al. [7], Singh and Vishwakarma [8], Koyuncu and Kadilar [4] proposed estimators in stratified random sampling. Singh [9] and Perri [6] suggested some ratio cum

product estimators in simple random sampling. Bahl and Tuteja [1] and Singh et al. [11] suggested some exponential ratio type estimators. In this chapter, we suggest some exponential-type estimators using the auxiliary information in the stratified random sampling.

Consider a finite population of size N and is divided into L strata such that $\sum_{h=1}^L N_h = N$ where N_h is the size of h^{th} stratum ($h=1,2,\dots,L$). We select a sample of size n_h from each stratum by simple random sample without replacement sampling such that $\sum_{h=1}^L n_h = n$, where n_h is the stratum sample size. A simple random sample of size n_h is drawn without replacement from the h^{th} stratum such that $\sum_{h=1}^L n_h = n$. Let (y_{hi}, x_{hi}, z_{hi}) denote the observed values of y , x , and z on the i^{th} unit of the h^{th} stratum, where $i=1, 2, 3, \dots, N_h$.

To obtain the bias and MSE, we write

$$\bar{y}_{st} = \sum_{h=1}^L w_h \bar{y}_h = \bar{Y}(1 + e_0), \quad \bar{x}_{st} = \sum_{h=1}^L w_h \bar{x}_h = \bar{X}(1 + e_1), \quad \bar{z}_{st} = \sum_{h=1}^L w_h \bar{z}_h = \bar{Z}(1 + e_2)$$

Such that,

$$E(e_0) = E(e_1) = E(e_2) = 0$$

$$V_{rst} = \sum_{h=1}^L w_h^{r+s+t} \frac{E\left[(\bar{y}_h - \bar{Y})^r (\bar{x}_h - \bar{X})^s (\bar{z}_h - \bar{Z})^t\right]}{\bar{Y}^r \bar{X}^s \bar{Z}^t}$$

where,

$$\bar{y}_{st} = \sum_{h=1}^L w_h \bar{y}_h, \quad \bar{y}_h = \frac{1}{n_h} \sum_{i=1}^{n_h} y_{hi}, \quad \bar{Y}_h = \frac{1}{N_h} \sum_{i=1}^{N_h} Y_{hi}$$

$$Y = \bar{Y}_{st} = \sum_{h=1}^L w_h \bar{Y}_h, \quad w_h = \frac{N_h}{N}$$

and

$$V(\bar{y}_{st}) = \bar{Y}^2 V_{200} \tag{1.1}$$

Similar expressions for X and Z can also be defined.

$$\text{And } E(e_0^2) = \frac{\sum_{h=1}^L W_h^2 f_h S_{yh}^2}{\bar{Y}^2} = V_{200},$$

$$E(e_1^2) = \frac{\sum_{h=1}^L W_h^2 f_h S_{xh}^2}{\bar{X}^2} = V_{020},$$

$$E(e_2^2) = \frac{\sum_{h=1}^L W_h^2 f_h S_{zh}^2}{\bar{Z}^2} = V_{002},$$

$$E(e_0 e_1) = \frac{\sum_{h=1}^L W_h^2 f_h S_{yxh}^2}{\bar{Y}\bar{X}} = V_{110},$$

$$E(e_0 e_2) = \frac{\sum_{h=1}^L W_h^2 f_h S_{yzh}^2}{\bar{Y}\bar{Z}} = V_{101},$$

$$\text{and } E(e_1 e_2) = \frac{\sum_{h=1}^L W_h^2 f_h S_{xzh}^2}{\bar{X}\bar{Z}} = V_{011},$$

where ,

$$S_{yh}^2 = \sum_{i=1}^{N_h} \frac{(\bar{y}_h - \bar{Y}_h)^2}{N_h - 1},$$

$$S_{xh}^2 = \sum_{i=1}^{N_h} \frac{(\bar{x}_h - \bar{X}_h)^2}{N_h - 1}$$

$$S_{zh}^2 = \sum_{i=1}^{N_h} \frac{(\bar{z}_h - \bar{Z}_h)^2}{N_h - 1},$$

$$S_{yxh} = \sum_{i=1}^{N_h} \frac{(\bar{x}_h - \bar{X}_h)(\bar{y}_h - \bar{Y}_h)}{N_h - 1}$$

$$S_{yzh} = \sum_{i=1}^{N_h} \frac{(\bar{z}_h - \bar{Z}_h)(\bar{y}_h - \bar{Y}_h)}{N_h - 1},$$

$$S_{xzh} = \sum_{i=1}^{N_h} \frac{(\bar{x}_h - \bar{X}_h)(\bar{z}_h - \bar{Z}_h)}{N_h - 1}$$

And,

$$f_h = \frac{1}{n_h} - \frac{1}{N_h}$$

2. Estimators in literature

In order to have an estimate of the study variable y , assuming the knowledge of the population proportion P , Naik and Gupta [5] and Singh et al. [11] respectively proposed following estimators

$$t_1 = \bar{y}_{st} \left(\frac{\bar{X}}{\bar{x}_{st}} \right) \tag{2.1}$$

$$t_2 = \bar{y}_{st} \exp \left(\frac{\bar{X} - \bar{x}_{st}}{\bar{X} + \bar{x}_{st}} \right) \tag{2.2}$$

The MSE expressions of these estimators are given as

$$\text{MSE}(t_1) = \bar{Y}^2 [V_{200} + V_{020} - 2V_{110}] \quad (2.3)$$

$$\text{MSE}(t_2) = \bar{Y}^2 \left[V_{200} + \frac{V_{020}}{4} - V_{110} \right] \quad (2.4)$$

When the information on the two auxiliary variables is known, Singh [10] proposed some ratio cum product estimators in simple random sampling to estimate the population mean of the study variable y .

Motivated by Singh [10] and Singh et al. [7], Singh and kumar propose some estimators in stratified sampling as

$$t_3 = \bar{y}_{st} \exp \left[\frac{\bar{X} - \bar{x}_{st}}{\bar{X} + \bar{x}_{st}} \right] \exp \left[\frac{\bar{Z} - \bar{z}_{st}}{\bar{Z} + \bar{z}_{st}} \right] \quad (2.5)$$

$$t_4 = \bar{y}_{st} \exp \left[\frac{\bar{x}_{st} - \bar{X}}{\bar{x}_{st} + \bar{X}} \right] \exp \left[\frac{\bar{z}_{st} - \bar{Z}}{\bar{z}_{st} + \bar{Z}} \right] \quad (2.6)$$

$$t_5 = \bar{y}_{st} \exp \left[\frac{\bar{X} - \bar{x}_{st}}{\bar{X} + \bar{x}_{st}} \right] \exp \left[\frac{\bar{z}_{st} - \bar{Z}}{\bar{z}_{st} + \bar{Z}} \right] \quad (2.7)$$

$$t_6 = \bar{y}_{st} \exp \left[\frac{\bar{x}_{st} - \bar{X}}{\bar{x}_{st} + \bar{X}} \right] \exp \left[\frac{\bar{Z} - \bar{z}_{st}}{\bar{Z} + \bar{z}_{st}} \right] \quad (2.8)$$

The MSE equations of these estimators can be written as

$$\text{MSE}(t_3) = \bar{Y}^2 \left[V_{200} + \frac{V_{020}}{4} + \frac{V_{002}}{4} - V_{110} - V_{101} + \frac{V_{011}}{2} \right] \quad (2.9)$$

$$\text{MSE}(t_4) = \bar{Y}^2 \left[V_{200} + \frac{V_{020}}{4} + \frac{V_{002}}{4} + V_{110} + V_{101} + \frac{V_{011}}{2} \right] \quad (2.10)$$

$$\text{MSE}(t_5) = \bar{Y}^2 \left[V_{200} + \frac{V_{020}}{4} + \frac{V_{002}}{4} - V_{110} + V_{101} - \frac{V_{011}}{2} \right] \quad (2.11)$$

$$\text{MSE}(t_6) = \bar{Y}^2 \left[V_{200} + \frac{V_{020}}{4} + \frac{V_{002}}{4} + V_{110} - V_{101} - \frac{V_{011}}{2} \right] \quad (2.12)$$

When there are two auxiliary variables, the regression estimator of \bar{Y} will be

$$t_7 = \bar{y}_{st} + b_{1h}(\bar{X} - \bar{x}_{st}) + b_{2h}(\bar{Z} - \bar{z}_{st}) \quad (2.13)$$

Where $b_{1h} = \frac{s_{yx}}{s_x^2}$ and $b_{2h} = \frac{s_{yz}}{s_z^2}$. Here s_x^2 and s_z^2 are the sample variances of x and z respectively,

s_{yx} and s_{yz} are the sample covariance's between y and x and between z respectively. The MSE expression of this estimator is:

$$\text{MSE}(t_7) = \sum_{h=1}^L W_h^2 f_h S_{yh}^2 (1 - \rho_{yxh}^2 - \rho_{yzh}^2 + 2\rho_{yxh}\rho_{yzh}\rho_{xzh}) \quad (2.14)$$

3. The proposed estimator

Following Malik and Singh [12], we propose an estimator using information on two auxiliary attributes as

$$t_p = \bar{y}_{st} \exp \left[\frac{\bar{X} - \bar{x}_{st}}{\bar{X} + \bar{x}_{st}} \right]^{m_1} \exp \left[\frac{\bar{Z} - \bar{z}_{st}}{\bar{Z} + \bar{z}_{st}} \right]^{m_2} + b_{1h}(\bar{X} - \bar{x}_{st}) + b_{2h}(\bar{Z} - \bar{z}_{st}) \quad (3.1)$$

Expressing equation (3.1) in terms of e's, we have

$$\begin{aligned} t_p &= \bar{Y} (1 + e_0) \left\{ \exp \left(\frac{-e_1}{2 + e_1} \right)^{m_1} \exp \left(\frac{-e_2}{2 + e_2} \right)^{m_2} \right\} - b_{1h} \bar{X} e_1 - b_{2h} \bar{Z} e_2 \\ &= \bar{Y} \left[1 + e_0 - \frac{m_1 e_1}{2} + \frac{m_1^2 e_1^2}{4} - \frac{m_2 e_2}{2} - \frac{m_1 m_2 e_1 e_2}{4} + \frac{m_2^2 e_2^2}{4} - \frac{m_2 e_0 e_2}{2} - \frac{m_1 e_0 e_1}{2} \right] \\ &\quad - b_{1h} e_1 \bar{X} - b_{2h} e_2 \bar{Z} \end{aligned} \quad (3.2)$$

Squaring both sides of (3.2) and neglecting the term having power greater than two, we have

$$(t_p - \bar{Y})^2 = \left\{ \bar{Y} \left[e_0 - \frac{m_1 e_1}{2} - \frac{m_2 e_2}{2} \right] - b_{1h} e_1 \bar{X} - b_{2h} e_2 \bar{Z} \right\}^2 \quad (3.3)$$

Taking expectations of both the sides of (3.3), we have the mean squared error of t_p up to the first degree of approximation as

$$\text{MSE}(t_p) = \bar{Y}^2 [V_{200} + P_1] + P_2 - \bar{Y} P_3 \quad (3.4)$$

Where,

$$\left. \begin{aligned} P_1 &= \frac{m_1^2 V_{020}}{4} + \frac{m_2^2 V_{002}}{4} + \frac{m_1 m_2 V_{011}}{2} - m_1 V_{110} - m_2 V_{101} \\ P_2 &= B_{1h}^2 V_{020} + B_{2h}^2 V_{002} + 2B_{1h} B_{2h} V_{011} \\ P_3 &= -2B_{1h} V_{110} - 2B_{2h} V_{101} + m_1 B_{1h} V_{020} + m_1 B_{2h} V_{011} + m_2 B_{1h} V_{011} + m_2 B_{2h} V_{002} \end{aligned} \right\} \quad (3.5)$$

$$\text{Where, } B_{1h} = \frac{\sum_{h=1}^L W_h^2 f_h \rho_{yxh} S_{yh} S_{xh}}{\sum_{h=1}^L W_h^2 f_h S_{xh}^2} \text{ and } B_{2h} = \frac{\sum_{h=1}^L W_h^2 f_h \rho_{yzh} S_{yh} S_{zh}}{\sum_{h=1}^L W_h^2 f_h S_{zh}^2}$$

The optimum values of m_1 and m_2 will be

$$\left. \begin{aligned} m_1 &= \frac{4[B_{1h} V_{011} V_{002} + B_{2h} V_{011}^2 - B_{1h} V_{020} V_{002} - B_{2h} V_{011} V_{002}]}{\bar{Y}[V_{020} V_{002} - V_{011}^2]} \\ m_2 &= \frac{4[B_{1h} V_{011} V_{020} + B_{2h} V_{011}^2 - B_{1h} V_{011} V_{020} - B_{2h} V_{002} V_{020}]}{\bar{Y}[V_{020} V_{002} - V_{011}^2]} \end{aligned} \right\} \quad (3.6)$$

Putting optimum values of m_1 and m_2 from (3.6), we obtained min MSE of proposed estimator t_p .

4. Efficiency comparison

In this section, the conditions for which the proposed estimator t_p is better than \bar{y}_{st} , t_1 , t_2 , t_3 ,

t_4 , t_5 , t_6 , and t_7 .

The variance is given by

$$V(\bar{y}_{st}) = \bar{Y}^2 V_{200} \quad (4.1)$$

To compare the efficiency of the proposed estimator with the existing estimator, from (4.1) and (2.3), (2.4), (2.9), (2.10), (2.11), (2.12) and (2.14), we have

$$V(\bar{y}_{st}) - \text{MSE}(t_p) = \bar{Y}^2 P_1 + P_2 - \bar{Y}P_3 \geq 0 \quad (4.2)$$

$$\text{MSE}(t_1) - \text{MSE}(t_p) = \bar{Y}^2 [V_{020} - 2V_{110}] - \bar{Y}^2 P_1 - P_2 + \bar{Y}P_3 \geq 0 \quad (4.3)$$

$$\text{MSE}(t_2) - \text{MSE}(t_p) = \bar{Y}^2 \left[\frac{V_{020}}{4} - V_{110} \right] - \bar{Y}^2 P_1 - P_2 + \bar{Y}P_3 \geq 0 \quad (4.4)$$

$$\text{MSE}(t_3) - \text{MSE}(t_p) = \bar{Y}^2 \left[\frac{V_{020}}{4} + \frac{V_{002}}{4} - V_{110} - V_{101} + \frac{V_{011}}{2} \right] - \bar{Y}^2 P_1 - P_2 + \bar{Y}P_3 \geq 0 \quad (4.5)$$

$$\text{MSE}(t_4) - \text{MSE}(t_p) = \bar{Y}^2 \left[\frac{V_{020}}{4} + \frac{V_{002}}{4} + V_{110} + V_{101} + \frac{V_{011}}{2} \right] - \bar{Y}^2 P_1 - P_2 + \bar{Y}P_3 \geq 0 \quad (4.6)$$

$$\text{MSE}(t_5) - \text{MSE}(t_p) = \bar{Y}^2 \left[\frac{V_{020}}{4} + \frac{V_{002}}{4} - V_{110} + V_{101} + \frac{V_{011}}{2} \right] - \bar{Y}^2 P_1 - P_2 + \bar{Y}P_3 \geq 0 \quad (4.7)$$

$$\text{MSE}(t_6) - \text{MSE}(t_p) = \bar{Y}^2 \left[\frac{V_{020}}{4} + \frac{V_{002}}{4} + V_{110} - V_{101} + \frac{V_{011}}{2} \right] - \bar{Y}^2 P_1 - P_2 + \bar{Y}P_3 \geq 0 \quad (4.8)$$

Using (4.2) - (4.8), we conclude that the proposed estimator outperforms than the estimators considered in literature.

5. Empirical study

In this section, we use the data set in Koyuncu and Kadilar [4]. The population statistics are given in Table 3.2.1. In this data set, the study variable (Y) is the number of teachers, the first auxiliary variable (X) is the number of students, and the second auxiliary variable (Z) is the number of classes in both primary and secondary schools.

Table 5.1: Data Statistics of Population

N ₁ =127	N ₂ =117	N ₃ =103
N ₄ =170	N ₅ =205	N ₆ =201

$n_1=31$	$n_2=21$	$n_3=29$
$n_4=38$	$n_5=22$	$n_6=39$
$S_{y_1} = 883.835$	$S_{y_2} = 644$	$S_{y_3} = 1033.467$
$S_{y_4} = 810.585$	$S_{y_5} = 403.654$	$S_{y_6} = 711.723$
$\bar{Y}_1 = 703.74$	$\bar{Y}_2 = 413$	$\bar{Y}_3 = 573.17$
$\bar{Y}_4 = 424.66$	$\bar{Y}_5 = 267.03$	$\bar{Y}_6 = 393.84$
$S_{x_1} = 30486.751$	$S_{x_2} = 15180.760$	$S_{x_3} = 27549.697$
$S_{x_4} = 18218.931$	$S_{x_5} = 8997.776$	$S_{x_6} = 23094.141$
$\bar{X}_1 = 20804.59$	$\bar{X}_2 = 9211.79$	$\bar{X}_3 = 14309.30$
$\bar{X}_4 = 9478.85$	$\bar{X}_5 = 5569.95$	$\bar{X}_6 = 12997.59$
$S_{xy_1} = 25237153.52$	$S_{xy_2} = 9747942.85$	$S_{xy_3} = 28294397.04$
$S_{xy_4} = 14523885.53$	$S_{xy_5} = 3393591.75$	$S_{xy_6} = 15864573.97$
$\rho_{xy_1} = 0.936$	$\rho_{xy_2} = 0.996$	$\rho_{xy_3} = 0.994$
$\rho_{xy_4} = 0.983$	$\rho_{xy_5} = 0.989$	$\rho_{xy_6} = 0.965$
$S_{z_1} = 555.5816$	$S_{z_2} = 365.4576$	$S_{z_3} = 612.9509281$
$S_{z_4} = 458.0282$	$S_{z_5} = 260.8511$	$S_{z_6} = 397.0481$
$\bar{Z}_1 = 498.28$	$\bar{Z}_2 = 318.33$	$\bar{Z}_3 = 431.36$
$\bar{Z}_4 = 498.28$	$\bar{Z}_5 = 227.20$	$\bar{Z}_6 = 313.71$
$S_{yz_1} = 480688.2$	$S_{yz_2} = 230092.8$	$S_{yz_3} = 623019.3$
$S_{yz_4} = 364943.4$	$S_{yz_5} = 101539$	$S_{yz_6} = 277696.1$
$S_{xz_1} = 15914648$	$S_{xz_2} = 5379190$	$S_{xz_3} = 164900674.56$
$S_{xz_4} = 8041254$	$S_{xz_5} = 2144057$	$S_{xz_6} = 8857729$
$\rho_{yz_1} = 0.978914$	$\rho_{yz_2} = 0.9762$	$\rho_{yz_3} = 0.983511$
$\rho_{yz_4} = 0.982958$	$\rho_{yz_5} = 0.964342$	$\rho_{yz_6} = 0.982689$

We have computed the pre relative efficiency (PRE) of different estimators of \bar{Y}_{st} with respect to \bar{y}_{st} and compiled in table 5.2:

Table 5.2: Percent Relative Efficiencies (PRE) of estimator

S.No.	Estimators	PRE'S
1	\bar{y}_{st}	100
2	t_1	1029.46
3	t_2	370.17
4	t_3	2045.43
5	t_4	27.94
6	t_5	126.41
7	t_6	77.21
8	t_7	2360.54
9	t_p	4656.35

6. Conclusion

In this paper, we proposed a new estimator for estimating unknown population mean of study variable using information on two auxiliary variables. Expressions for bias and MSE of the estimator are derived up to first degree of approximation. The proposed estimator is compared with usual mean estimator and other considered estimators. A numerical study is carried out to

support the theoretical results. In the table 5.2, the proposed estimator performs better than the usual sample mean and other considered estimators.

References

- [1] Bahl, S. and Tuteja, R.K. (1991): Ratio and product type exponential estimator. *Infrm.andOptim. Sci.*, XII, I, 159-163.
- [2] Diana, G. (1993). A class of estimators of the population mean in stratified random sampling. *Statistica*53(1):59–66.
- [3] Kadilar,C. and Cingi,H. (2003): Ratio Estimators in Straitified Random Sampling. *Biometrical Journal* 45 (2003) 2, 218-225.
- [4] Koyuncu, N. and Kadilar, C. (2009) : Family of estimators of population mean using two auxiliary variables in stratified random sampling. *Comm. In Stat. – Theory and Meth.*, 38:14, 2398-2417.
- [5] Naik,V.D and Gupta, P.C., 1996: A note on estimation of mean with known population proportion of an auxiliary character. *Jour. Ind. Soc. Agri. Stat.*, 48(2), 151-158.
- [6] Perri, P. F. (2007). Improved ratio-cum-product type estimators.*Statist. Trans.* 8:51–69.
- [7] Singh, R., Kumar, M., Singh,R.D.,and Chaudhary, M.K.(2008): Exponential Ratio Type Estimators in Stratified Random Sampling. Presented in International Symposium On Optimisation and Statistics (I.S.O.S).-2008 held at A.M.U ., Aligarh, India, during 29-31 Dec 2008.
- [8] Singh, H., P. and Vishwakarma, G. K. (2008): A family of estimators of population mean using auxiliary information in stratified sampling. *Communication in Statistics Theory and Methods*, 37(7), 1038-1050.
- [9] Singh, M. P. (1965): On the estimation of ratio and product of population parameter. *Sankhya* 27 B, 321-328.
- [10] Singh, M.P. (1967): Ratio-cum-product method of estimation. *Metrika* 12, 34- 72.
- [11] Singh, R.,Chauhan, P., Sawan, N. and Smarandache,F. (2007): Auxiliary information and a priory values in construction of improved estimators. *Renaissance High press.*
- [12] Malik and Singh (2013): An improved estimator using two auxiliary attributes. *Applied Mathematics and Computation*, 219(2013), 10983-10986

Some Ratio and Product Estimators Using Known Value of Population Parameters

Rajesh Singh, Sachin Malik* and Florentin Smarandache**

*Department of Mathematics, SRM University

Delhi NCR, Sonapat- 131029, India

Department of Statistics, Banaras Hindu University

Varanasi-221005, India

**Department of Mathematics, University of New Mexico

Gallup, NM 87301, USA

*Corresponding Author, sachinkurava999@gmail.com

Abstract

In the present article, we proposed a family of estimators for estimating population means using known value of some population parameters. Khoshnevisan et al. [1] proposed a general family of estimators for estimating population means using known value of some population parameter(s) which after some substitutions led to some ratio and product estimators initially proposed by Sisodia and Dwivedi [2], Singh and Tailor [3], Pandey and Dubey [4], Adewara et al. [5], yadav and Kadilar [6]. The present family of estimators provides us significant improvement over previous families in theory. An empirical study is carried out to judge the merit of the proposed estimator.

Keywords: Ratio Estimator, Product Estimator, Population Parameter, Efficiency, Mean Square Error.

1. Introduction

The problem of estimating the population mean in the presence of an auxiliary variable has been widely discussed in finite population sampling literature. Ratio, product and difference methods of estimation are good examples in this context. Ratio method of estimation is quite effective when there is high positive correlation between study and auxiliary variables. On the other hand, if correlation is negative (high), the product method of estimation can be employed efficiently.

In recent years, a number of research papers on ratio-type, exponential ratio-type and regression-type estimators have appeared, based on different types of transformations. Some important contributions in this area are due to Singh and Tailor [3], Shabbir and Gupta [7,8], Kadilar and Cingi [9,10], Khosnevisan et. al.(2007).

Khoshnevisan et al. [1] defined their family of estimators as

$$t = \bar{y} \left[\frac{a\bar{X} + b}{\alpha(a\bar{x} + b) + (1-\alpha)(a\bar{X} + b)} \right]^g$$

where $a(\neq 0)$, b are either real numbers or the functions of the known parameters of the auxiliary variable x such as standard deviation (σ_x), Coefficient of Variation (C_x), Skewness ($\beta_1(x)$), Kurtosis ($\beta_2(x)$) and Correlation Coefficient (ρ).

(i). When $\alpha=0$, $a=0=b$, $g=0$, we have the mean per unit estimator, $t_0 = \bar{y}$ with

$$MSE(t_0) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 C_y^2 \quad (1.1)$$

(ii). When $\alpha=1$, $a=1$, $b=0$, $g=1$, we have the usual ratio estimator, $t_1 = \bar{y} \left(\frac{\bar{X}}{\bar{x}} \right)$ with

$$MSE(t_1) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 (C_y^2 + C_x^2 - 2\rho\rho_x C_y) \quad (1.2)$$

(iii). When $\alpha=1$, $a=1$, $b=0$, $g=-1$, we have the usual product estimator, $t_2 = \bar{y} \left(\frac{\bar{x}}{\bar{X}} \right)$ with

$$MSE(t_2) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 (C_y^2 + C_x^2 + 2\rho\rho_x C_y) \quad (1.3)$$

(iv). When $\alpha=1$, $a=1$, $b=C_x$, $g=1$, we have Sisodia and Dwivedi [2] ratio estimator,

$$t_3 = \bar{y} \left(\frac{\bar{X} + C_x}{\bar{x} + C_x} \right) \text{ with}$$

$$MSE(t_3) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 \left(C_y^2 + \left(\frac{\bar{X}}{\bar{x} + C_x} \right)^2 C_x^2 - 2 \left(\frac{\bar{X}}{\bar{x} + C_x} \right) \rho\rho_x C_y \right) \quad (1.4)$$

(v). When $\alpha=1$, $a=1$, $b=C_x$, $g=-1$

we have Pandey and Dubey [4] product estimator, $t_4 = \bar{y} \left(\frac{\bar{X} + C_x}{\bar{X} + C_x} \right)$ with

$$\text{MSE}(t_4) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 (C_y^2 + \left(\frac{\bar{X}}{\bar{X} + C_x} \right)^2 C_x^2 + 2 \left(\frac{\bar{X}}{\bar{X} + C_x} \right) \rho \rho_x C_y) \quad (1.5)$$

(vi). When $\alpha=1$, $a=1$, $b=\rho$, $g=1$, we have Singh and Taylor [3] ratio estimator, $t_5 = \bar{y} \left(\frac{\bar{X} + \rho}{\bar{X} + \rho} \right)$

with

$$\text{MSE}(t_5) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 (C_y^2 + \left(\frac{\bar{X}}{\bar{X} + \rho} \right)^2 C_x^2 - 2 \left(\frac{\bar{X}}{\bar{X} + \rho} \right) \rho \rho_x C_y) \quad (1.6)$$

(vii). When $\alpha=1$, $a=1$, $b=\rho$, $g=-1$, we have Singh and Taylor [3] product estimator,

$t_6 = \bar{y} \left(\frac{\bar{X} + \rho}{\bar{X} + \rho} \right)$ with

$$\text{MSE}(t_6) = \left(\frac{N-n}{Nn} \right) \bar{Y}^2 (C_y^2 + \left(\frac{\bar{X}}{\bar{X} + \rho} \right)^2 C_x^2 + 2 \left(\frac{\bar{X}}{\bar{X} + \rho} \right) \rho \rho_x C_y) \quad (1.7)$$

There are other ratio and product estimators from these families that are not inferred here but this paper will be limited to those ones that made use of Coefficient of Variation (C_x) and Correlation Coefficient (ρ) since the conclusion obtained here can also be inferred on all others that made use of other population parameters such as the standard deviation (σ_x), Skewness ($\beta_1(x)$) and Kurtosis ($\beta_2(x)$) in the same family.

2. On the Modified Ratio and Product Estimators.

Adopting Adewara (2006), Adewara et al. (2012) proposed the following estimators as

$$t_1^* = \bar{y}^* \left(\frac{\bar{X}}{\bar{X}^*} \right), \quad (2.1)$$

$$t_2^* = \bar{y}^* \left(\frac{\bar{X}^*}{\bar{X}} \right), \quad (2.2)$$

$$t_3^* = \bar{y}^* \left(\frac{\bar{X} + C_x}{\bar{X}^* + C_x} \right), \quad (2.3)$$

$$t^*_4 = \bar{y}^* \left(\frac{\bar{x}^* + C_x}{\bar{X} + C_x} \right), \quad (2.4)$$

$$t^*_5 = \bar{y}^* \left(\frac{\bar{X} + \rho}{\bar{x}^* + \rho} \right) \text{ and} \quad (2.5)$$

$$t^*_6 = \bar{y}^* \left(\frac{\bar{x}^* + \rho}{\bar{X} + \rho} \right), \quad (2.6)$$

Where \bar{x}^* and \bar{y}^* are the sample means of the auxiliary variables and variable of interest yet to be drawn with the relationships (i) $\bar{X} = f\bar{x} + (1-f)\bar{x}^*$ and (ii). $\bar{Y} = f\bar{y} + (1-f)\bar{y}^*$. Srivenkataramana and Srinath [12].

The Mean Square Errors of these estimators t^*_i , $i = 1, 2, \dots, 6$ are as follows:

$$(i). \text{MSE}(t^*_1) = \left(\frac{n}{N-n} \right)^2 \text{MSE}(t_1) \quad (2.7)$$

$$(ii). \text{MSE}(t^*_2) = \left(\frac{n}{N-n} \right)^2 \text{MSE}(t_2) \quad (2.8)$$

$$(iii). \text{MSE}(t^*_3) = \left(\frac{n}{N-n} \right)^2 \text{MSE}(t_3) \quad (2.9)$$

$$(iv). \text{MSE}(t^*_4) = \left(\frac{n}{N-n} \right)^2 \text{MSE}(t_4) \quad (2.10)$$

$$(v). \text{MSE}(t^*_5) = \left(\frac{n}{N-n} \right)^2 \text{MSE}(t_5) \quad (2.11)$$

$$(vi). \text{MSE}(t^*_6) = \left(\frac{n}{N-n} \right)^2 \text{MSE}(t_6) \quad (2.12)$$

Following Adewara et al [5], Yadav and Kadilar [6] proposed some improved ratio and product estimators for estimating the population mean of the study variable as follows

$$\eta^*_1 = k\bar{y}^* \left(\frac{\bar{X}}{\bar{x}^*} \right), \quad (2.13)$$

$$\eta^*_2 = k\bar{y}^* \left(\frac{\bar{x}^*}{\bar{X}} \right), \quad (2.14)$$

$$\eta^*_3 = k \bar{y}^* \left(\frac{\bar{X} + C_x}{\bar{x}^* + C_x} \right), \quad (2.15)$$

$$\eta^*_4 = k \bar{y}^* \left(\frac{\bar{x}^* + C_x}{\bar{X} + C_x} \right), \quad (2.16)$$

$$\eta^*_5 = k \bar{y}^* \left(\frac{\bar{X} + \rho}{\bar{x}^* + \rho} \right) \quad (2.17)$$

$$\eta^*_6 = k \bar{y}^* \left(\frac{\bar{x}^* + \rho}{\bar{X} + \rho} \right), \quad (2.18)$$

The mean square error of these estimators η^*_i , $i=1,2,\dots,6$ are as follows

$$\text{MSE}(\eta^*_1) = \bar{Y}^2 \left[h^2 (k_1^2 \lambda C_y^2 + \{3k_1^2 - 2k_1\} \lambda C_x^2 - 2\{2k_1^2 - k_1\} \lambda C_{yx}) + \{k_1 - 1\}^2 \right] \quad (2.19)$$

$$\text{MSE}(\eta^*_2) = \bar{Y}^2 \left[h^2 (k_2^2 \lambda C_y^2 + k_2^2 \lambda C_x^2 + 2\{2k_2^2 - k_2\} \lambda C_{yx}) + \{k_2 - 1\}^2 \right] \quad (2.20)$$

$$\text{MSE}(\eta^*_3) = \bar{Y}^2 \left[h^2 (k_3^2 \lambda C_y^2 + \{3k_3^2 - 2k_3\} v_1^2 \lambda C_x^2 - 2v_1 \{2k_3^2 - k_3\} \lambda C_{yx}) + \{k_3 - 1\}^2 \right] \quad (2.21)$$

$$\text{MSE}(\eta^*_4) = \bar{Y}^2 \left[h^2 (k_4^2 \lambda C_y^2 + k_4^2 v_1^2 \lambda C_x^2 + 2v_1 \{2k_4^2 - k_4\} \lambda C_{yx}) + \{k_4 - 1\}^2 \right] \quad (2.22)$$

$$\text{MSE}(\eta^*_5) = \bar{Y}^2 \left[h^2 (k_5^2 \lambda C_y^2 + \{3k_5^2 - 2k_5\} v_2^2 \lambda C_x^2 - 2v_2 \{2k_5^2 - k_5\} \lambda C_{yx}) + \{k_5 - 1\}^2 \right] \quad (2.23)$$

$$\text{MSE}(\eta^*_6) = \bar{Y}^2 \left[h^2 (k_6^2 \lambda C_y^2 + k_6^2 v_2^2 \lambda C_x^2 + 2v_2 \{2k_6^2 - k_6\} \lambda C_{yx}) + \{k_6 - 1\}^2 \right] \quad (2.24)$$

Where,

$$\lambda = \frac{N-n}{Nn}, h = \frac{n}{N-n}, C_y^2 = \frac{S_y^2}{\bar{Y}^2}, C_x^2 = \frac{S_x^2}{\bar{X}^2}, C_{yx} = \frac{S_{yx}}{\bar{Y}\bar{X}}, v_1 = \frac{\bar{X}}{\bar{X} + C_x}, v_2 = \frac{\bar{X}}{\bar{X} + C_x} \text{ and } \rho = \frac{S_{yx}}{S_y S_x}$$

$$\text{And } k_1 = \frac{h^2 [\lambda C_x^2 - \lambda C_{yx}] + 1}{h^2 [3C_x^2 \lambda - 4C_{yx} \lambda + \lambda C_y^2] + 1}, k_2 = \frac{h^2 \lambda C_{yx} + 1}{h^2 [C_x^2 \lambda + 4C_{yx} \lambda + \lambda C_y^2] + 1},$$

$$k_3 = \frac{h^2 [\lambda v_1^2 C_x^2 - v_1 \lambda C_{yx}] + 1}{h^2 [3v_1^2 C_x^2 \lambda - 4v_1 C_{yx} \lambda + \lambda C_y^2] + 1}, k_4 = \frac{h^2 [\lambda v_1 \lambda C_{yx}] + 1}{h^2 [3v_1^2 C_x^2 \lambda + 4v_1 C_{yx} \lambda + \lambda C_y^2] + 1}$$

$$k_5 = \frac{h^2 [\lambda v_2^2 C_x^2 - v_2 \lambda C_{yx}] + 1}{h^2 [3v_2^2 C_x^2 \lambda - 4v_2 C_{yx} \lambda + \lambda C_y^2] + 1}, \text{ and } k_6 = \frac{h^2 [\lambda v_2 \lambda C_{yx}] + 1}{h^2 [3v_2^2 C_x^2 \lambda + 4v_2 C_{yx} \lambda + \lambda C_y^2] + 1}$$

3. The Proposed family of estimators

Following Malik Singh [14], we define the following class of estimators for population mean \bar{Y} as

$$t_M = \left\{ m_1 \bar{y}^* + m_2 [\bar{X} - \bar{x}^*] \right\} \left(\frac{\psi \bar{X} + \delta}{\psi \bar{x}^* + \delta} \right)^\alpha \exp \left[\frac{(\omega \bar{X} + \mu) - (\omega \bar{x}^* + \mu)}{(\omega \bar{X} + \mu) + (\omega \bar{x}^* + \mu)} \right]^\beta \quad (3.1)$$

Where m_1 and m_2 are suitably chosen constants. ψ, δ, ω , and μ are either real numbers or function of known parameters of the auxiliary variable. The scalar α and β takes values +1 and -1 for ratio and product type estimators respectively.

To obtain the MSE, let us define

$$\bar{y} = \bar{Y}(1 + e_0), \quad \bar{x} = \bar{X}(1 + e_1)$$

such that $E(e_i) = 0, i=0,1$ and

$$E(e_0^2) = \lambda C_y^2, \quad E(e_1^2) = \lambda C_x^2, \quad E(e_0 e_1) = \lambda \rho C_y C_x$$

expressing equation (3.1) in terms of e 's and retaining only terms up to second degree of e 's, we have

$$\begin{aligned} t_M &= \left[m_1 \bar{Y}(1 - h e_0) + m_2 \bar{X} h e_1 \right] \left\{ \frac{\psi \bar{X} + \delta}{\psi \bar{X}(1 - h e_1) + \delta} \right\}^\alpha \exp \left\{ \frac{\omega \bar{X} h e_1}{2\omega \bar{X} + 2\mu - \omega \bar{X} h e_1} \right\}^\beta \\ &= \left[m_1 \bar{Y}(1 - h e_0) + m_2 \bar{X} h e_1 \right] \{1 - R_1 h e_1\}^{-\alpha} \exp \left\{ \beta R_2 h e_1 \left(1 - \frac{R_2 h e_1}{2} + \frac{R_2^2 h^2 e_1^2}{4} \right) \right\} \\ &= m_1 \bar{Y} \left[\begin{aligned} &1 + \alpha h e_1 + \frac{\alpha(\alpha + 1) h^2 e_1^2}{2} + \frac{\beta h e_1}{2} + \frac{\alpha \beta h^2 e_1^2}{2} + \frac{\beta^2 h^2 e_1^2}{8} + \frac{\beta h^2 e_1^2}{4} - h e_0 \\ &- \alpha h^2 e_0 e_1 - \frac{\beta h^2 e_0 e_1}{2} \end{aligned} \right] \\ &\quad + m_2 \bar{X} \left[h e_1 + \alpha h^2 e_1^2 + \frac{\beta h^2 e_1^2}{2} \right] \end{aligned} \quad (3.2)$$

where, $R_1 = \frac{\psi\bar{X}}{\psi\bar{X} + \delta}, R_2 = \frac{\omega\bar{X}}{\omega\bar{X} + \mu}$

Subtracting \bar{Y} from both the sides of (3.2), we have

$$(t_M - \bar{Y}) = m_1 \bar{Y} [1 - h e_0 + L_1 e_1 + L_2 e_1^2 - L_3 e_0 e_1] + m_2 \bar{X} [h e_1 + L_4 e_1^2] - \bar{Y} \tag{3.3}$$

where,

$$L_1 = \alpha R_1 h + \frac{\beta h R_2}{2}$$

$$L_2 = \frac{\alpha(\alpha + 1)h^2 R_1^2}{2} + \frac{\alpha\beta h^2 R_1 R_2}{2} + \frac{\beta^2 h^2 R_2^2}{8} + \frac{\beta h^2 R_2^2}{4}$$

$$L_3 = \alpha h^2 R_1 + \frac{\beta h^2 R_2}{2}$$

$$L_4 = \alpha h^2 R_1 + \frac{\beta h^2 R_2}{2}$$

Squaring both sides of (3.3) and neglecting terms of e's having power greater than two, we have

$$MSE(t_M) = \bar{Y}^2 [1 + m_1^2 T_1 + m_2^2 T_2 + 2m_1 m_2 T_3 - 2m_1 T_4 - 2m_2 T_5] \tag{3.4}$$

where,

$$\left. \begin{aligned} T_1 &= \bar{Y}^2 [1 + \lambda h^2 C_y^2 + L_1^2 \lambda C_x^2 - 2hL_1 \lambda \rho C_y C_x + 2L_2 \lambda C_x^2 - 2L_3 \lambda \rho C_y C_x] \\ T_2 &= h^2 \lambda \bar{X}^2 C_x^2 \\ T_3 &= \bar{Y} \bar{X} [L_4 \lambda C_x^2 + L_1 \lambda h C_x^2 - h^2 \lambda \rho C_y C_x] \\ T_4 &= \bar{Y}^2 [1 + L_2 \lambda C_x^2 - L_3 \lambda \rho C_y C_x] \\ T_5 &= \bar{Y} \bar{X} L_4 \lambda C_x^2 \end{aligned} \right\}$$

minimization of (3.4) with respect to m_1 and m_2 yields optimum values as

$$m_1 = \frac{(T_2 T_4 - T_3 T_5)}{T_1 T_2 - T_3^2}, \quad m_2 = \frac{(T_1 T_5 - T_3 T_4)}{T_1 T_2 - T_3^2}$$

4. Empirical Study:

Population I: Kadilar and Cingi [9]

$N = 106, n = 20, \rho = 0.86, C_y = 5.22, C_x = 2.1, \bar{Y} = 2212.59$ and $\bar{X} = 27421.70$

Population II: Maddala [13]

$N = 16, n = 4, \rho = -0.6823, C_y = 0.2278, C_x = 0.0986, \bar{Y} = 7.6375$ and $\bar{X} = 75.4313$

4. Results:

Table 4.1: Showing the estimates obtained for both the Khoshnevisan et al. [1] estimators and Adewara et al. [5] estimators

Estimator	Population I ($\rho > 0$)	Population II ($\rho < 0$)
t_0	5411349	0.5676
t_1	2542740	-
t_2	-	0.3387
t_3	2542893	-
t_4	-	0.3388
t_5	2542803	-
t_6	-	0.3376
t_1^*	137519.8	-
t_2^*	-	0.03763
t_3^*	137528	-
t_4^*	-	0.03765
t_5^*	137523.1	-
t_6^*	-	0.03751

Table 4.2: Showing the estimates obtained for Yadav and Kadilar [6] estimators

Estimator	Population I ($\rho > 0$)	Population II ($\rho < 0$)
η_1^*	136145.37	-
η_2^*	-	0.03762
η_3^*	136138.05	-
η_4^*	-	0.03764
η_5^*	136107.94	-
η_6^*	-	0.03750

Table 4.3: MSE of suggested estimators with different values of constants

m_1	m_2	α	β	ψ	δ	ω	μ	estimator	MSE	
									PopI	PopII
1	0	1	0	1	0	-	-	t_1^*	137519.8	-
1	0	-1	0	1	0	-	-	t_2^*	-	0.03763
1	0	1	0	1	C_x	-	-	t_3^*	137528	-
1	0	-1	0	1	C_x	-	-	t_4^*	-	0.03765
1	0	1	0	1	ρ	-	-	t_5^*	137523.1	-
1	0	-1	0	1	ρ	-	-	t_6^*	-	0.03751
m_1	0	1	0	1	0	-	-	η_1^*	136145.37	-
m_1	0	-1	0	1	0	-	-	η_2^*	-	0.03762

m_1	0	1	0	1	C_x	-	-	η_3^*	136138.05	-
m_1	0	-1	0	1	C_x	-	-	η_4^*	-	0.03764
m_1	0	1	0	1	ρ	-	-	η_5^*	136107.94	-
m_1	0	-1	0	1	ρ	-	-	η_6^*	-	0.03750
m_1	m_2	1	1	1	1	1	1	t_M	75502.23	-
m_1	m_2	-1	-1	1	1	1	1	t_M	-	0.03370

Since conventionally, for ratio estimators to hold, $\rho > 0$ and also for product estimators to hold, $\rho < 0$. Therefore two data sets are used in this paper, one to determine the efficiency of the modified ratio estimators and the other to determine that of the product estimators as stated below.

5. Conclusion

In this paper, we have proposed a new family of estimator for estimating unknown population mean of study variable using auxiliary variable. Expressions for the MSE of the estimator are derived up to first order of approximation. The proposed family of estimator is compared with the several existing estimators in literature. From table 4.3, we observe that the new family of estimators performs better than the other estimators considered in this paper for both of the data sets.

References

- [1] Khoshnevisan, M., Singh, R., Chauhan, P., Sawan, N. and Smarandache, F.(2007):
A general family of estimators for estimating population mean using known value of some population parameter(s). Far East Jour. of theor. statist. 22(2), 181 – 191.
- [2] Sisodia, B.V.S. and Dwivedi, V.K.(1981): A modified ratio estimator using coefficient of variation of auxiliary variable. Journal Ind. Soc. Agril. Statist.,33(2), 13 – 18.
- [3] Singh, H.P. and Tailor, R. (2003) : Use of known correlation coefficient in estimating the finite population mean. Statist. in Trans. 6(4). 555 – 560.

- [4] Pandey, B.N. and Dubey, Vyas (1988): Modified product estimator using coefficient of variation of auxiliary variate, *Assam Statistical Rev.*, 2(2), 64 – 66
- [5] A.A. Adewara, R. Singh, M. Kumar(2012) : Efficiency of some modified ratio and product estimators using known value of some population parameters, *International Journal of Applied Science and Technology* 2 (2) 76–79.
- [6] Yadav, S.K. and Kadilar, C. (2013): Improved class of ratio and product estimators. *Applied mathematics and computation*.
- [7] Shabbir, J., Gupta, S. (2005). : Improved ratio estimators in stratified sampling. *Amer. J. Mathemat. Manag. Sci.* 25:293–311
- [8] Shabbir, J. and Gupta, S. (2006) : A new estimator of population mean in stratified sampling, *Commun. Stat. Theo. Meth.* 35: 1201–1209
- [9] Kadilar, C. and Cingi, H. (2004) : Ratio Estimators in Simple Random Sampling. *Appl. Mathe. and Comput.* 151, 893-902
- [10] Kadilar, C. and Cingi, H. (2006).: Ratio estimators for the population variance in sample and stratified random sampling. *Appl. Mathe. and Comput.* 173, 1047 – 1059.
- [11] Adewara, A.A. (2006). : Effects of improving both the auxiliary and variable of interest in ratio and product estimators. *Proc. Pakistan Acad. Sci.*43(4): 275 – 278.
- [12] Srivenkataramana, T. and Srinath, K.P. (1976) : Ratio and Product methods of estimation in sample surveys when the two variables are moderately correlated. *Vignana Bharathi* 2: 54 – 58
- [13] Maddala, G.S. (1977): *Econometrics*. “McGraw Hills Pub. Co.” New York .
- [14] Malik and Singh (2015): Estimation of population mean using information on auxiliary attribute in two-phase sampling. *Applied Mathematics and computation*, 261 (2015) 114-1

An Unbiased Estimator for Estimating Population Mean in Simple Random Sampling Using Auxiliary Attribute

Rajesh Singh and Sachin Malik*

*Department of Mathematics, SRM University

Delhi NCR, Sonapat- 131029, India

Department of Statistics, Banaras Hindu University

Varanasi-221005, India

*Corresponding Author, sachinkurava999@gmail.com

Abstract

In this paper we have proposed an almost unbiased estimator using known value of some population parameter(s) with known population proportion of an auxiliary variable. A class of estimators is defined which includes Naik and Gupta [1], Singh and Solanki [2] and Sahai and Ray [3] estimators. Under simple random sampling without replacement (SRSWOR) scheme the expressions for bias and mean square error (MSE) are derived. Numerical illustrations are given in support of the present study.

Key words: Auxiliary information, bias, mean square error, unbiased estimator.

Introduction

It is well known that the precision of the estimates of the population mean or total of the study variable y can be considering improved by the use of known information on an auxiliary variable x which is highly correlated with the study variable y . Out of many methods ratio, product and regression methods of estimation are good illustrations in this context. Using known values of certain populations parameters several authors have proposed improved estimators including Singh and Tailor [4], Kadilar and Cingi [5], Gupta and Shabbir [6,7], Khoshnevisan et al. [8], Singh et al. [9], Singh et al. [10], Koyuncu and Kadilar [11], Diana et al. [12], Upadhyaya et al. [13] and Singh and Solanki [2].

In many practical situations, instead of existence of auxiliary variables there exist some auxiliary attributes ϕ (say), which are highly correlated with the study variable y , such as

- i. Amount of milk produced (y) and a particular breed of cow (ϕ).
- ii. Sex (ϕ) and height of persons (y) and
- iii. Amount of yield of wheat crop and a particular variety of wheat (ϕ) etc. (see Jhajj et al. [15]).

Many more situations can be encountered in practice where the information of the population mean \bar{Y} of the study variable y in the presence of auxiliary attributes assumes importance. For these reasons various authors such as Naik and Gupta [1], Jhajj et al. [14], Abd-Elfattah et al. [15], Grover and Kaur [16], Malik and Singh [17] and Singh and Solanki [2] have paid their attention towards the improved estimation of population mean \bar{Y} of the study variable y taking into consideration the point bi-serial correlation between a variable and an attribute.

Let $A = \sum_{i=1}^N \phi_i$ and $a = \sum_{i=1}^n \phi_i$ denote the total number of units in the population and sample possessing attribute ϕ respectively, $p = \frac{A}{N}$ and $p = \frac{a}{n}$ denote the proportion of units in the population and sample, respectively, possessing attribute ϕ .

Define,

$$e_y = \frac{(\bar{y} - \bar{Y})}{\bar{Y}} \quad e_\phi = \frac{(p - P)}{P},$$

$$E(e_i) = 0, (i = y, \phi)$$

$$E(e_y^2) = fC_y^2, \quad E(e_\phi^2) = fC_p^2, \quad E(e_y e_\phi) = f\rho_{pb} C_y C_p.$$

Where

$$f_1 = \left(\frac{1}{n} - \frac{1}{N} \right), \quad C_y^2 = \frac{S_y^2}{\bar{Y}^2}, \quad C_p^2 = \frac{S_p^2}{P^2},$$

and $\rho_{pb} = \frac{S_{y\phi}}{S_y S_\phi}$ is the point bi-serial correlation coefficient.

Here,

$$S_y^2 = \frac{1}{N-1} \sum_{i=1}^N (y_i - \bar{Y})^2, \quad S_\phi^2 = \frac{1}{N-1} \sum_{i=1}^N (\phi_i - P)^2 \quad \text{and} \quad S_{y\phi} = \frac{1}{N-1} \left(\sum_{i=1}^N y_i \phi_i - NP\bar{Y} \right)$$

In order to have an estimate of the study variable y , assuming the knowledge of the population proportion P , Naik and Gupta [1] proposed following estimate

$$t_{NGR} = \bar{y} \left(\frac{P}{p} \right) \quad (1.1)$$

$$t_{NGP} = \bar{y} \left(\frac{p}{P} \right) \quad (1.2)$$

following Naik and Gupta [1], we propose the following estimator

$$t_1 = \bar{y} \left(\frac{K_1 P + K_2 K_3}{K_1 p + K_2 K_3} \right)^\alpha \quad (1.3)$$

The Bias and MSE expression's of the estimator t_1 up to the first order of approximation are, respectively, given by

$$B(t_1) = \bar{Y} f_1 C_p^2 \left[\frac{\alpha(\alpha+1)V_1^2}{2} - \alpha V_1 K_p \right] \quad (1.4)$$

$$MSE(t_1) = \bar{Y}^2 f_1 \left[C_y^2 + C_p^2 (\alpha^2 V_1^2 - 2\alpha V_1 K_p) \right] \quad (1.5)$$

Also following Singh and Solanki [2], we propose the following estimator

$$t_2 = \bar{y} \left\{ 2 - \left(\frac{p}{P} \right)^\beta \exp \left[\lambda \left(\frac{(K_4 P + K_5) - (K_4 p + K_5)}{(K_4 P + K_5) + (K_4 p + K_5)} \right) \right] \right\} \quad (1.6)$$

The Bias and MSE expression's of the estimator t_2 up to the first order of approximation are, respectively, given by

$$B(t_2) = \bar{Y} f_1 C_p^2 \left[\frac{\lambda V_2 \beta}{2} - \frac{\beta(\beta-1)}{2} - \frac{\lambda(\lambda+2)V_2^2}{8} - \beta K_p + \frac{\lambda V_2 K_p}{2} \right] \quad (1.7)$$

$$MSE(t_2) = \bar{Y}^2 f_1 \left[C_y^2 + C_p^2 \left(\beta^2 + \frac{\lambda^2 V_2^2}{4} - \beta \lambda V_2 \right) - 2K_p C_p^2 \left(\beta - \frac{\lambda V_2}{2} \right) \right] \quad (1.8)$$

α , λ and β are suitable chosen constants. Also K_1, K_3, K_4, K_5 are either real numbers or function of known parameters of the auxiliary attributes ϕ such as C_p , $\beta_2(\phi)$, ρ_{pb} and K_p . K_2 is an integer which takes values +1 and -1 for designing the estimators and

$$\left. \begin{aligned} V_1 &= \frac{K_1 P}{K_1 P + K_2 K_3} \\ V_2 &= \frac{K_4 P}{K_4 P + K_5} \end{aligned} \right\}$$

We see that the estimators t_1 and t_2 are biased estimators. In some applications bias is disadvantageous. Following these estimators we have proposed almost unbiased estimator of \bar{Y} .

2. Almost unbiased estimator

$$\text{Suppose } t_0 = \bar{y}, t_1 = \bar{y} \left(\frac{K_1 P + K_2 K_3}{K_1 p + K_2 K_3} \right)^\alpha, t_2 = \bar{y} \left\{ 2 - \left(\frac{p}{P} \right)^\beta \exp \left[\lambda \left(\frac{(K_4 P + K_5) - (K_4 p + K_5)}{(K_4 P + K_5) + (K_4 p + K_5)} \right) \right] \right\}$$

Such that $t_0, t_1, t_2 \in W$, where W denotes the set of all possible estimators for estimating the population mean \bar{Y} . By definition, the set W is a linear variety if

$$t_p = \sum_{i=0}^3 w_i t_i \in W \tag{2.1}$$

$$\text{Such that, } \sum_{i=0}^3 w_i = 1 \text{ and } w_i \in \mathbb{R} \tag{2.2}$$

where $w_i (i = 0, 1, 2, 3)$ denotes the constants used for reducing the bias in the class of estimators, H denotes the set of those estimators that can be constructed from $t_i (i = 0, 1, 2, 3)$ and \mathbb{R} denotes the set of real numbers.

Expressing t_p in terms of e 's, we have

$$t_p = \bar{Y} \left[1 + e_y + w_1 \left(\frac{\alpha(\alpha+1)V_1^2 e_\phi^2}{2} - \alpha V_1 e_\phi - \alpha V_1 e_y e_\phi \right) \right]$$

$$+ w_2 \left(-\beta e_\phi - \frac{\beta(\beta-1)e_\phi^2}{2} + \frac{\lambda V_2 e_\phi}{2} + \frac{\lambda V_2 \beta e_\phi^2}{2} - \frac{\lambda(\lambda+2)V_2^2 e_\phi^2}{8} - \beta e_y e_\phi + \frac{\lambda V_2 e_y e_\phi}{2} \right) \quad (2.3)$$

Subtracting \bar{Y} from both sides of equation (2.3) and then taking expectation of both sides, we get the bias of the estimator t_6 up to the first order of approximation, as

$$B(t_p) = \bar{Y} f_1 w_1 C_p^2 \left(\frac{\alpha(\alpha+1)V_1^2}{2} - \alpha V_1 K_p \right) + \bar{Y} f_1 w_2 C_p^2 \left(\frac{\lambda V_2 \beta}{2} - \frac{\beta(\beta-1)}{2} - \frac{\lambda(\lambda+2)V_2^2}{8} - \beta K_p + \frac{\lambda V_2 K_p}{2} \right) \quad (2.4)$$

From (2.3), we have

$$(t_p - \bar{Y}) = \bar{Y} \left[e_0 - w_1 \alpha V_1 e_\phi - w_2 \left(\beta e_\phi + \frac{\lambda V_2 e_\phi}{2} \right) \right] \quad (2.5)$$

Squaring both sides of (2.5) and then taking expectation, we get the MSE of the estimator t_6 up to the first order of approximation, as

$$MSE(t_p) = \bar{Y}^2 f_1 [C_y^2 + C_p^2 (Q^2 - 2QK_p)] \quad (2.6)$$

Where

Which is minimum when

$$Q = K_p \quad (2.7)$$

$$\text{Where } Q = w_1 \alpha V_1 + w_2 \left(\beta - \frac{\lambda V_2}{2} \right) \quad (2.8)$$

Putting the value of $Q = K_p$ in (2.6) we have optimum value of estimator as t_p (optimum).

Thus the minimum MSE of t_p is given by

$$\min. MSE(t_p) = \bar{Y}^2 f_1 C_y^2 (1 - \rho_{pb}^2) \quad (2.9)$$

Which is same as that of traditional linear regression estimator.

from (2.2) and (2.8), we have only two equations in three unknowns. It is not possible to find the unique values for w_i 's, $i=0,1,2$. In order to get unique values of w_i 's, we shall impose the linear restriction

$$\sum_{i=0}^2 w_i B(t_i) = 0 \quad (2.10)$$

where $B(t_i)$ denotes the bias in the i^{th} estimator.

Equations (2.2), (2.8) and (2.10) can be written in the matrix form as

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & \alpha V_1 & \beta - \frac{\lambda V_2}{2} \\ 0 & B(t_1) & B(t_2) \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 1 \\ k_p \\ 0 \end{bmatrix} \quad (2.11)$$

Using (2.11), we get the unique values of w_i 's, $i=0,1,2$ as

$$\left. \begin{aligned} w_0 &= \frac{\alpha V_1 [\alpha V_1 A_2 - A_1 X_1] - X_1 K_p A_1 - X_2 \alpha V_1 [\alpha V_1 A_2 - A_1 X_1] - \alpha V_1 K_p A_1}{\alpha V_1 [\alpha V_1 A_2 - A_1 X_1]} \\ w_1 &= \frac{X_1 K_p A_1}{\alpha V_1 [\alpha V_1 A_2 - A_1 X_1]} + X_2 \\ w_2 &= \frac{K_p A_1}{[\alpha V_1 A_2 - A_1 X_1]} \end{aligned} \right\}$$

where,

$$\left. \begin{aligned} A_1 &= \frac{\alpha(\alpha+1)V_1^2}{2} - \alpha V_1 K_p \\ A_2 &= \frac{\lambda V_2 \beta}{2} - \frac{\beta(\beta-1)}{2} - \frac{\lambda(\lambda+2)V_2^2}{8} - \beta K_p + \frac{\lambda V_2 K_p}{2} \\ X_1 &= A_1 \left[\beta - \frac{\lambda V_2}{2} \right] \\ X_2 &= \frac{K_p}{\alpha V_1} \end{aligned} \right\}$$

Use of these w_i 's, $i=0,1,2$ remove the bias up to terms of order $o(n^{-1})$ at (2.1).

3. Empirical study

For empirical study we use the data sets earlier used by Sukhatme and Sukhatme [18], (p.256) (population 1) and Mukhopadhyaya [19] (p.44) (population 2) to verify the theoretical results.

Data statistics:

Population	N	n	\bar{Y}	P	C_y	C_p	ρ_{pb}	$\beta_2(\phi)$
Population 1	89	20	3.360	0.1236	0.60400	2.19012	0.766	6.2381
Population 2	25	10	9.44	0.400	0.17028	1.27478	-0.387	4.3275

Table 3.1 : Values of w_i 's,

w_i 's,	Population 1	Population 2
w_0	-3.95624	1.124182
w_1	5.356173	0.020794
w_2	0.39993	-0.14498

Table 3.2: PRE of different estimators of \bar{Y} with respect to \bar{y}

Choice of scalars												Estimator	PRE (POPII)	PRE (POPII)
w_0	w_1	w_2	K_1	K_2	K_3	K_4	K_5	α	β	λ				
1	0	0										\bar{y}	100	100
0	1	0	1	1	0			1				t_{NGR}	11.63	1.59
			1	1	0			-1				t_{NGP}	5.075	1.94

0	0	1							1	0	$t_{1(1,0)}$	12.88	1.59
									-1	0	$t_{1(-1,0)}$	5.43	1.95
						1	0		1	1	$t_{2(1,1)}$	73.59	0.84
						1	0		1	-1	$t_{2(1,-1)}$	4.94	8.25
						1	0		0	1	$t_{2(0,1)}$	14.95	8.25
						1	0		0	-1	$t_{2(0,-1)}$	73.48	5.58
w_0	w_1	w_2	1	1	1	1	1	1	1	1	t_p optimum	241.98	117.61

4. Proposed estimators in two phase sampling

In some practical situations when P is not known a priori, the technique of two-phase sampling is used. Let p' denote the proportion of units possessing attribute ϕ in the first phase sample of size n' ; p denote the proportion of units possessing attribute ϕ in the second phase sample of size $n' > n$ and \bar{y} denote the mean of the study variable y in the second phase sample.

In two-phase sampling the estimator t_p will take the following form

$$t_{pd} = \sum_{i=0}^3 h_i t_{id} \in H \quad (4.1)$$

$$\text{Such that, } \sum_{i=0}^3 h_i = 1 \text{ and } h_i \in R \quad (4.2)$$

Where,

$$t_{0d} = \bar{y}, t_{1d} = \bar{y} \left(\frac{K_1 p' + K_2 K_3}{K_1 p + K_2 K_3} \right)^m, t_{2d} = \bar{y} \left\{ 2 - \left(\frac{p}{p'} \right)^n \exp \left[\gamma \left(\frac{(K_4 p' + K_5) - (K_4 p + K_5)}{(K_4 p' + K_5) + (K_4 p + K_5)} \right) \right] \right\}$$

The Bias and MSE expression's of the estimator t_{1d} and t_{2d} up to the first order of approximation are, respectively, given by

$$B(t_{1d}) = \bar{Y} \left[\frac{m(m-1)R_1^2 f_2 C_p^2}{2} + \frac{m(m+1)R_1^2 f_1 C_p^2}{2} - m^2 R_1^2 f_2 C_p^2 + m R_1 f_3 k_p C_p^2 \right] \quad (4.3)$$

$$MSE(t_1) = \bar{Y}^2 [f_1 C_y^2 + m^2 R_1^2 f_3 C_p^2 - 2m R_1 k_p f_3 C_p^2] \quad (4.4)$$

$$B(t_{2d}) = \bar{Y} \left[-\frac{n(n-1)f_1 C_p^2}{2} + \frac{n(n+1)f_2 C_p^2}{2} + n f_2 k_p C_p^2 + n^2 f_2 C_p^2 + f_3 \gamma R_2 k_p C_p^2 + f_3 \gamma R_2 n C_p^2 \right] \quad (4.5)$$

$$MSE(t_{2d}) = \bar{Y}^2 [f_1 C_y^2 + L_1^2 f_3 C_p^2] \quad (4.6)$$

Where,

$$\left. \begin{aligned} R_1 &= \frac{K_1 P}{K_1 P + K_2 K_3} \\ R_2 &= \frac{K_4 P}{2[K_4 P + K_5]} \\ L_1 &= n - \gamma A_2 \end{aligned} \right\} \quad (4.7)$$

Expressing (4.1) in terms of e's, we have

$$t_p = \bar{Y} \left[1 + e_y + w_1 \left(\frac{m(m+1)R_1^2 e_\phi^2}{2} - m R_1 e_\phi - m R_1 e_y e_\phi + m R_1 e_y e_\phi + \frac{m(m-1)R_1^2 e_\phi^2}{2} + m R_1 e_y e'_\phi \right) \right. \\ \left. + w_2 \left(-n e_\phi - \frac{n(n-1)e_\phi^2}{2} + n e'_\phi + n^2 e_\phi e'_\phi - \frac{n(n+1)e_\phi^2}{2} - \gamma R_2 (e'_\phi - e_\phi) + \gamma R_2 (e_y e_\phi - e_y e'_\phi) - n e_y e_\phi \right) \right]$$

Subtracting \bar{Y} from both sides of equation (2.3) and then taking expectation of both sides, we get the bias of the estimator t_6 up to the first order of approximation, as

$$B(t_p) = \bar{Y} [B(t_{1d}) + B(t_{2d})] \quad (4.8)$$

Also,

$$(t_p - \bar{Y}) = \bar{Y} \left[e_0 + w_1 \left[mR_1 e'_\phi - mR_1 e_\phi \right] + w_2 \left(-ne_\phi + ne'_\phi - \gamma R_2 e'_\phi + \gamma R_2 e_\phi \right) \right] \quad (4.9)$$

Squaring both sides of (4.9) and then taking expectation, we get the MSE of the estimator t_p up to the first order of approximation, as

$$\text{MSE}(t_p) = \bar{Y}^2 f_1 C_y^2 + L_2^2 f_3 C_p^2 - 2L_2 f_3 k_p C_p^2 \quad (4.10)$$

Where

Which is minimum when

$$L_2 = K_p \quad (4.11)$$

$$\text{Where } L_2 = w_1 mR_1 + w_2 (n - \gamma R_2) \quad (4.12)$$

Putting the value of $L_2 = K_p$ in (4.10), we have optimum value of estimator as t_p (optimum).

Thus the minimum MSE of t_p is given by

$$\text{min.MSE}(t_p) = \bar{Y}^2 C_y^2 (f_1 - f_3 \rho_{pb}^2) \quad (4.13)$$

Which is same as that of traditional linear regression estimator.

from (4.2) and (4.12), we have only two equations in three unknowns. It is not possible to find the unique values for h_i 's, $i=0,1,2$. In order to get unique values of h_i 's, we shall impose the linear restriction

$$\sum_{i=0}^2 h_i B(t_i) = 0 \quad (4.14)$$

where $B(t_i)$ denotes the bias in the i^{th} estimator.

Equations (4.2), (4.12) and (4.14) can be written in the matrix form as

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & mR_1 & n - \gamma R_2 \\ 0 & B(t_1) & B(t_2) \end{bmatrix} \begin{bmatrix} h_0 \\ h_1 \\ h_2 \end{bmatrix} = \begin{bmatrix} 1 \\ k_p \\ 0 \end{bmatrix} \quad (4.15)$$

Using (4.15), we get the unique values of w_i 's, $i=0,1,2$ as

$$\left. \begin{aligned} h_0 &= 1 - h_1 - h_2 \\ h_1 &= \frac{k_p}{mR_1} - \frac{P_1 K_p (n - \gamma R_2)}{P_1 n - mR_1 P_2 - P_1 \gamma R_2} \\ h_2 &= \frac{K_p P_1}{[P_1 n - mR_1 P_2 - P_1 \gamma R_2]} \end{aligned} \right\}$$

Where,

$$\left. \begin{aligned} P_1 &= \frac{m(m-1)R_1^2 f_2 C_p^2}{2} + \frac{m(m+1)R_1^2 f_1 C_p^2}{2} - m^2 R_1^2 f_2 C_p^2 + mR_1 f_3 k_p C_p^2 \\ P_2 &= \left[-\frac{n(n-1)f_1 C_p^2}{2} + \frac{n(n+1)f_2 C_p^2}{2} + n f_2 k_p C_p^2 + n^2 f_2 C_p^2 + f_3 \gamma R_2 k_p C_p^2 + f_3 \gamma R_2 n C_p^2 \right] \end{aligned} \right\}$$

Use of these h_i 's, $i=0,1,2$ remove the bias up to terms of order $o(n^{-1})$ at (4.1).

5. Empirical Study

For empirical study we use the data sets earlier used by Sukhatme and Sukhatme [18] (, p.256) (population 1) and Mukhopadhyaya [19] (p.44) (population 2) to verify the theoretical results.

5.1 Data statistics:

Pop.	N	n	\bar{Y}	P	p'	C_y	C_p	ρ_{pb}	n'
Pop. 1	89	23	1322	0.1304	0.13336	0.69144	2.7005	0.408	45
Pop. 2	25	7	7.143	0.294	0.308	0.36442	1.3470	-0.314	13

Table 5.2: PRE of different estimators of \bar{Y} with respect to \bar{y}

Choice of scalars											Estimator	PRE (POPII)	PRE (POPII)
w_0	w_1	w_2	K_1	K_2	K_3	K_4	K_5	M	n	γ			
1	0	0									\bar{y}	100	100
0	1	0	1	1	0			1			t_{NGR}	11.13	8.85
			1	1	0			-1			t_{NGP}	7.48	12.15
0	0	1							1	0	$t_{1d(1,0)}$	26.84	5.42
									-1	0	$t_{1d(-1,0)}$	23.75	5.87
						1	0		1	1	$t_{2d(1,1)}$	82.55	1.23
						1	0		1	-1	$t_{2d(1,-1)}$	8.56	8.46
						1	0		0	1	$t_{2d(0,1)}$	22.54	6.57
						1	0		0	-1	$t_{2d(0,-1)}$	82.56	7.45
w_0	w_1	w_2	1	1	1	1	1	1	1	1	t_p optimum	112.55	106.89

Conclusion

In this paper, we have proposed an unbiased estimator t_p and t_{pd} using information on the auxiliary attribute(s) in case of single phase and double phase sampling respectively. Expressions for bias and MSE's of the proposed estimators are derived up to first degree of approximation.

From theoretical discussion and empirical study we conclude that the proposed estimators t_p and t_{pd} under optimum conditions perform better than other estimators considered in the article.

Appendix A.

Some members of the proposed family of estimators -

Some members (ratio-type) of the class t_1					
When $w_0 = 0, w_1 = 1, w_2 = 0$					
$\alpha = 1,$					
K_1	K_3	Estimators ($K_2 = 1$)	Estimators ($K_2 = -1$)	PRE'S $K_2 = 1$	PRE'S $K_2 = -1$
1	C_p	$t_{1a1} = \bar{y} \left[\frac{P + C_p}{p + C_p} \right]$	$t_{1b1} = \bar{y} \left[\frac{P - C_p}{p - C_p} \right]$	134.99	72.50
1	$\beta_2(\phi)$	$t_{1a2} = \bar{y} \left[\frac{P + \beta_2(\phi)}{p + \beta_2(\phi)} \right]$	$t_{1b2} = \bar{y} \left[\frac{P - \beta_2(\phi)}{p - \beta_2(\phi)} \right]$	111.62	89.34
$\beta_2(\phi)$	C_p	$t_{1a3} = \bar{y} \left[\frac{P\beta_2(\phi) + C_p}{p\beta_2(\phi) + C_p} \right]$	$t_{1b3} = \bar{y} \left[\frac{P\beta_2(\phi) - C_p}{p\beta_2(\phi) - C_p} \right]$	226.28	12.99
C_p	$\beta_2(\phi)$	$t_{1a4} = \bar{y} \left[\frac{PC_p + \beta_2(\phi)}{pC_p + \beta_2(\phi)} \right]$	$t_{1b4} = \bar{y} \left[\frac{PC_p - \beta_2(\phi)}{pC_p - \beta_2(\phi)} \right]$	126.66	77.93
1	ρ_{pb}	$t_{1a5} = \bar{y} \left[\frac{P + \rho_{pb}}{p + \rho_{pb}} \right]$	$t_{1b5} = \bar{y} \left[\frac{P - \rho_{pb}}{p - \rho_{pb}} \right]$	207.46	39.13

NP	S_ϕ	$t_{1a6} = \bar{y} \left[\frac{NP^2 + S_\phi}{NPp + S_\phi} \right]$	$t_{1b6} = \bar{y} \left[\frac{NP^2 - S_\phi}{NPp - S_\phi} \right]$	18.14	6.86
NP	f	$t_{1a7} = \bar{y} \left[\frac{NP^2 + f}{NPp + f} \right]$	$t_{1b7} = \bar{y} \left[\frac{NP^2 - f}{NPp - f} \right]$	13.79	10.85
$\beta_2(\phi)$	K_{pb}	$t_{1a8} = \bar{y} \left[\frac{\beta_2(\phi)P + K_{pb}}{\beta_2(\phi)p + K_{pb}} \right]$	$t_{1b8} = \bar{y} \left[\frac{\beta_2(\phi)P - K_{pb}}{\beta_2(\phi)p - K_{pb}} \right]$	24.15	5.40
NP	K_{pb}	$t_{1a9} = \bar{y} \left[\frac{NP^2 + K_{pb}}{NPp + K_{pb}} \right]$	$t_{1b9} = \bar{y} \left[\frac{NP^2 - K_{pb}}{NPp - K_{pb}} \right]$	18.62	7.78
N	1	$t_{1a10} = \bar{y} \left[\frac{NP + 1}{Np + 1} \right]$	$t_{1b10} = \bar{y} \left[\frac{NP - 1}{Np - 1} \right]$	15.93	9.26
N	C_p	$t_{1a11} = \bar{y} \left[\frac{NP + C_p}{Np + C_p} \right]$	$t_{1b11} = \bar{y} \left[\frac{NP - C_p}{Np - C_p} \right]$	19.79	6.86
N	ρ_{pb}	$t_{1a12} = \bar{y} \left[\frac{NP + \rho_{pb}}{Np + \rho_{pb}} \right]$	$t_{1b12} = \bar{y} \left[\frac{NP - \rho_{pb}}{Np - \rho_{pb}} \right]$	15.18	9.78
N	S_ϕ	$t_{1a13} = \bar{y} \left[\frac{NP + S_\phi}{Np + S_\phi} \right]$	$t_{1b13} = \bar{y} \left[\frac{NP - S_\phi}{Np - S_\phi} \right]$	12.34	10.96
N	f	$t_{1a14} = \bar{y} \left[\frac{NP + f}{Np + f} \right]$	$t_{1b14} = \bar{y} \left[\frac{NP - f}{Np - f} \right]$	12.99	11.54
N	$g=1-f$	$t_{1a15} = \bar{y} \left[\frac{NP + g}{Np + g} \right]$	$t_{1b15} = \bar{y} \left[\frac{NP - g}{Np - g} \right]$	15.81	9.34
N	K_{pb}	$t_{1a16} = \bar{y} \left[\frac{NP + K_{pb}}{Np + K_{pb}} \right]$	$t_{1b16} = \bar{y} \left[\frac{NP - K_{pb}}{Np - K_{pb}} \right]$	13.52	11.10

n	ρ_{pb}	$t_{1a17} = \bar{y} \left[\frac{nP + \rho_{pb}}{np + \rho_{pb}} \right]$	$t_{1b17} = \bar{y} \left[\frac{nP - \rho_{pb}}{np - \rho_{pb}} \right]$	25.13	4.86
n	S_ϕ	$t_{1a18} = \bar{y} \left[\frac{nP + \rho_{pb}}{np + \rho_{pb}} \right]$	$t_{1b18} = \bar{y} \left[\frac{nP - \rho_{pb}}{np - \rho_{pb}} \right]$	14.98	8.81
n	f	$t_{1a19} = \bar{y} \left[\frac{nP + f}{np + f} \right]$	$t_{1b19} = \bar{y} \left[\frac{nP - f}{np - f} \right]$	13.38	11.20
n	$g=1-f$	$t_{1a20} = \bar{y} \left[\frac{nP + g}{np + g} \right]$	$t_{1b20} = \bar{y} \left[\frac{nP - g}{np - g} \right]$	29.13	3.68
n	K_{pb}	$t_{1a21} = \bar{y} \left[\frac{nP + K_{pb}}{np + K_{pb}} \right]$	$t_{1b21} = \bar{y} \left[\frac{nP - K_{pb}}{np - K_{pb}} \right]$	15.87	9.39
$\beta_2(\phi)$	P	$t_{1a22} = \bar{y} \left[\frac{\beta_2(\phi)P + P}{\beta_2(\phi)p + P} \right]$	$t_{1b22} = \bar{y} \left[\frac{\beta_2(\phi)P - P}{\beta_2(\phi)p - P} \right]$	16.80	7.63
NP	P	$t_{1a23} = \bar{y} \left[\frac{NP^2 + P}{NPp + P} \right]$	$t_{1b23} = \bar{y} \left[\frac{NP^2 - P}{NPp - P} \right]$	15.93	9.26
N	P	$t_{1a24} = \bar{y} \left[\frac{NP + P}{Np + P} \right]$	$t_{1b24} = \bar{y} \left[\frac{NP - P}{Np - P} \right]$	13.23	11.32
n	P	$t_{1a25} = \bar{y} \left[\frac{nP + P}{np + P} \right]$	$t_{1b25} = \bar{y} \left[\frac{nP - P}{np - P} \right]$	14.51	10.28

Appendix B.

Some members (product-type) of the class t_1					
When $w_0 = 0, w_1 = 1, w_2 = 0$					
$\alpha = -1,$					
K_1	K_3	Estimators ($K_2 = 1$)	Estimators ($K_2 = -1$)	PRE'S $K_2 = 1$	PRE'S $K_2 = -1$
1	C_p	$t_{1c1} = \bar{y} \left[\frac{p + C_p}{P + C_p} \right]$	$t_{1d1} = \bar{y} \left[\frac{p - C_p}{P - C_p} \right]$	35.54	9.93
1	$\beta_2(\phi)$	$t_{1c2} = \bar{y} \left[\frac{p + \beta_2(\phi)}{P + \beta_2(\phi)} \right]$	$t_{1d2} = \bar{y} \left[\frac{p - \beta_2(\phi)}{P - \beta_2(\phi)} \right]$	110.12	101.54
$\beta_2(\phi)$	C_p	$t_{1c3} = \bar{y} \left[\frac{p\beta_2(\phi) + C_p}{P\beta_2(\phi) + C_p} \right]$	$t_{1d3} = \bar{y} \left[\frac{p\beta_2(\phi) - C_p}{P\beta_2(\phi) - C_p} \right]$	6.09	0.127
C_p	$\beta_2(\phi)$	$t_{1c4} = \bar{y} \left[\frac{pC_p + \beta_2(\phi)}{PC_p + \beta_2(\phi)} \right]$	$t_{1d4} = \bar{y} \left[\frac{pC_p - \beta_2(\phi)}{PC_p - \beta_2(\phi)} \right]$	99.38	82.52
1	ρ_{pb}	$t_{1c5} = \bar{y} \left[\frac{p + \rho_{pb}}{P + \rho_{pb}} \right]$	$t_{1d5} = \bar{y} \left[\frac{p - \rho_{pb}}{P - \rho_{pb}} \right]$	0.00135	5.42
NP	S_ϕ	$t_{1c6} = \bar{y} \left[\frac{NPp + S_\phi}{NP^2 + S_\phi} \right]$	$t_{1d6} = \bar{y} \left[\frac{NPp - S_\phi}{NP^2 - S_\phi} \right]$	2.53	1.23
NP	f	$t_{1c7} = \bar{y} \left[\frac{NPp + f}{NP^2 + f} \right]$	$t_{1d7} = \bar{y} \left[\frac{NPp - f}{NP^2 - f} \right]$	2.03	1.52

$\beta_2(\phi)$	K_{pb}	$t_{1c8} = \bar{y} \left[\frac{\beta_2(\phi)p + K_{pb}}{\beta_2(\phi)P + K_{pb}} \right]$	$t_{1d8} = \bar{y} \left[\frac{\beta_2(\phi)P - K_{pb}}{\beta_2(\phi)p - K_{pb}} \right]$	1.83	1.68
NP	K_{pb}	$t_{1c9} = \bar{y} \left[\frac{NPp + K_{pb}}{NP^2 + K_{pb}} \right]$	$t_{1d9} = \bar{y} \left[\frac{NPp - K_{pb}}{NP^2 - K_{pb}} \right]$	1.89	1.63
N	1	$t_{1c10} = \bar{y} \left[\frac{Np + 1}{NP + 1} \right]$	$t_{1d10} = \bar{y} \left[\frac{Np - 1}{NP - 1} \right]$	2.37	1.30
N	C_p	$t_{1c11} = \bar{y} \left[\frac{Np + C_p}{NP + C_p} \right]$	$t_{1d11} = \bar{y} \left[\frac{Np - C_p}{NP - C_p} \right]$	2.50	1.23
N	ρ_{pb}	$t_{1c12} = \bar{y} \left[\frac{Np + \rho_{pb}}{NP + \rho_{pb}} \right]$	$t_{1d12} = \bar{y} \left[\frac{Np - \rho_{pb}}{NP - \rho_{pb}} \right]$	1.79	1.70
N	S_ϕ	$t_{1c13} = \bar{y} \left[\frac{Np + S_\phi}{NP + S_\phi} \right]$	$t_{1d13} = \bar{y} \left[\frac{Np - S_\phi}{NP - S_\phi} \right]$	2.16	1.44
N	f	$t_{1c14} = \bar{y} \left[\frac{Np + f}{NP + f} \right]$	$t_{1d14} = \bar{y} \left[\frac{Np - f}{NP - f} \right]$	1.98	1.56
N	$g=1-f$	$t_{1c15} = \bar{y} \left[\frac{Np + g}{NP + g} \right]$	$t_{1d15} = \bar{y} \left[\frac{Np - g}{NP - g} \right]$	2.34	1.32
N	K_{pb}	$t_{1c16} = \bar{y} \left[\frac{Np + K_{pb}}{NP + K_{pb}} \right]$	$t_{1d16} = \bar{y} \left[\frac{Np - K_{pb}}{NP - K_{pb}} \right]$	1.93	1.60
n	ρ_{pb}	$t_{1c17} = \bar{y} \left[\frac{np + \rho_{pb}}{nP + \rho_{pb}} \right]$	$t_{1d17} = \bar{y} \left[\frac{np - \rho_{pb}}{nP - \rho_{pb}} \right]$	1.49	1.96
n	S_ϕ	$t_{1c18} = \bar{y} \left[\frac{np + \rho_{pb}}{nP + \rho_{pb}} \right]$	$t_{1d18} = \bar{y} \left[\frac{np - \rho_{pb}}{nP - \rho_{pb}} \right]$	2.65	1.14

n	f	$t_{1c19} = \bar{y} \left[\frac{np + f}{nP + f} \right]$	$t_{1d19} = \bar{y} \left[\frac{np - f}{nP - f} \right]$	2.06	1.51
n	g=1-f	$t_{1c20} = \bar{y} \left[\frac{np + g}{nP + g} \right]$	$t_{1d20} = \bar{y} \left[\frac{np - g}{nP - g} \right]$	3.29	0.84
n	K_{pb}	$t_{1c21} = \bar{y} \left[\frac{np + K_{pb}}{nP + K_{pb}} \right]$	$t_{1d21} = \bar{y} \left[\frac{np - K_{pb}}{nP - K_{pb}} \right]$	1.88	1.63
$\beta_2(\phi)$	P	$t_{1c22} = \bar{y} \left[\frac{\beta_2(\phi)p + P}{\beta_2(\phi)P + P} \right]$	$t_{1d22} = \bar{y} \left[\frac{\beta_2(\phi)p - P}{\beta_2(\phi)P - P} \right]$	2.99	0.97
NP	P	$t_{1c23} = \bar{y} \left[\frac{NPp + P}{NP^2 + P} \right]$	$t_{1d23} = \bar{y} \left[\frac{NPp - P}{NP^2 - P} \right]$	2.37	1.30
N	P	$t_{1c24} = \bar{y} \left[\frac{Np + P}{NP + P} \right]$	$t_{1d24} = \bar{y} \left[\frac{Np - P}{NP - P} \right]$	2.11	1.47
n	P	$t_{1c25} = \bar{y} \left[\frac{np + P}{nP + P} \right]$	$t_{1d25} = \bar{y} \left[\frac{np - P}{nP - P} \right]$	2.49	1.23

Appendix C.

Some members (product-type) of the class t_2			
When $w_0 = 0, w_1 = 0, w_2 = 1$			
K_4	K_5	Estimators $(\beta = 1, \lambda = -1)$	PRE'S
1	C_p	$t_{21} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{p-P}{p+P+2C_p} \right] \right)$	12.42
1	$\beta_2(\phi)$	$t_{22} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{p-P}{p+P+2\beta_2(\phi)} \right] \right)$	11.92
$\beta_2(\phi)$	C_p	$t_{23} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{\beta_2(\phi)(p-P)}{\beta_2(\phi)(p+P)+2C_p} \right] \right)$	16.29
C_p	$\beta_2(\phi)$	$t_{24} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{C_p(p-P)}{C_p(p+P)+2\beta_2(\phi)} \right] \right)$	12.53
1	ρ_{pb}	$t_{25} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{(p-P)}{(p+P)+2\rho_{pb}} \right] \right)$	13.86
NP	S_ϕ	$t_{26} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{NP(p-P)}{NP(p+P)+2S_\phi} \right] \right)$	44.46
NP	f	$t_{27} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{NP(p-P)}{NP(p+P)+2f} \right] \right)$	61.84

$\beta_2(\phi)$	K_{pb}	$t_{28} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{\beta_2(\phi)(p-P)}{\beta_2(\phi)(p+P) + 2K_{pb}} \right] \right)$	40.17
NP	K_{pb}	$t_{29} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{NP(p-P)}{NP(p+P) + 2K_{pb}} \right] \right)$	48.09
N	1	$t_{210} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2} \right] \right)$	54.10
N	C_p	$t_{211} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2C_p} \right] \right)$	44.84
N	ρ_{pb}	$t_{212} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2\rho_{pb}} \right] \right)$	56.48
N	S_ϕ	$t_{213} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2S_\phi} \right] \right)$	62.40
N	f	$t_{214} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2f} \right] \right)$	65.67
N	$g=1-f$	$t_{215} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2g} \right] \right)$	54.47
N	K_{pb}	$t_{216} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2K_{pb}} \right] \right)$	63.21
N	ρ_{pb}	$t_{217} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{n(p-P)}{n(p+P) + 2\rho_{pb}} \right] \right)$	38.59

N	S_ϕ	$t_{218} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{n(p-P)}{n(p+P) + 2S_\phi} \right] \right)$	52.19
N	f	$t_{219} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{n(p-P)}{n(p+P) + 2f} \right] \right)$	63.74
N	g=1-f	$t_{220} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{n(p-P)}{n(p+P) + 2g} \right] \right)$	35.33
N	K_{pb}	$t_{221} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{n(p-P)}{n(p+P) + 2K_{pb}} \right] \right)$	54.68
$\beta_2(\phi)$	P	$t_{222} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{\beta_2(\phi)(p-P)}{\beta_2(\phi)(p+P) + 2P} \right] \right)$	47.53
NP	P	$t_{223} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{NP(p-P)}{NP(p+P) + 2P} \right] \right)$	54.10
N	P	$t_{224} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{N(p-P)}{N(p+P) + 2P} \right] \right)$	64.43
N	P	$t_{225} = \bar{y} \left(2 - \frac{p}{P} \exp \left[\frac{n(p-P)}{n(p+P) + 2P} \right] \right)$	58.91

In addition to above estimators a large number of estimators can also be generated from the proposed estimators just by putting different values of constants w_i 's, $K_1, K_2, K_3, K_4, K_5, \alpha, \beta$ and λ .

References

- [1] Naik, V.D., Gupta, P.C. (1996): A note on estimation of mean with known population proportion of an auxiliary character. *Journal of the Indian Society of Agricultural Statistics* 48(2) 151-158.
- [2] Singh, H. P., Solanki, R.S. (2012): Improved estimation of population mean in simple random sampling using information on auxiliary attribute. *Applied Mathematics and Computation* 218 (2012) 7798–7812.
- [3] Sahai A., S. K. Ray S. K. (1980) “An efficient estimator using auxiliary information,” *Metrika*, vol. 27, no. 4, pp. 271–275.
- [4] Singh, H.P. and Tailor, R. (2003): Use of known correlation coefficient in estimating the finite population mean. *Statistics in Transition*, 6,4,555-560.
- [5] Kadilar,C. and Cingi,H. (2003): Ratio Estimators in Stratified Random Sampling. *Biometrical Journal* 45 (2003) 2, 218-225.
- [6] Shabbir, J. and Gupta, S. (2007): A new estimator of population mean in stratified sampling, *Commun. Stat. Theo. Meth.* 35: 1201–1209.
- [7] Gupta, S. and Shabbir, J. (2008) : On improvement in estimating the population mean in simple random sampling. *Jour. Of Applied Statistics*, 35, 5, 559-566.
- [8] Khoshnevisan, M. Singh, R., Chauhan, P., Sawan, N. and Smarandache, F. (2007): A general family of estimators for estimating population mean using known value of some population parameter(s). *Far east journal of theoretical statistics*, 22(2), 181-191.
- [9] Singh, R.,Cauhan, P., Sawan, N. and Smarandache,F. (2007): Auxiliary information and a priory values in construction of improved estimators. *Renaissance High press*.
- [10] Singh, R., Kumar, M. and Smarandache, F. (2008): Almost Unbiased Estimator for Estimating Population Mean Using Known Value of Some Population Parameter(s). *Pak. J. Stat. Oper. Res.*, 4(2) pp63-76.
- [11] Koyuncu, N. and Kadilar, C. (2009) : Family of estimators of population mean using two auxiliary variables in stratified random sampling. *Comm. In Stat. – Theory and Meth.*, 38:14, 2398-2417.
- [12] Diana G., Giordan M. Perri P.(2011): An improved class of estimators for the population mean. *Stat Methods Appl* (2011) 20:123–140
- [13] Upadhyaya LN, Singh HP, Chatterjee S, Yadav R (2011) Improved ratio and product exponential type estimators. *J Stat Theory Pract* 5(2):285–302
- [14] Jhaji, H. S., Sharma, M.K. and Grover, L.K. (2006): A family of estimators of population mean using information on auxiliary attribute. *Pakistan journal of statistics*, 22(1), 43-50 (2006).
- [15] Abd-Elfattah, A.M. El-Sherpieny, E.A. Mohamed, S.M. Abdou, O. F. (2010): Improvement in estimating the population mean in simple random sampling using information on auxiliary attribute. *Appl. Mathe. and Compt.* doi:10.1016/j.amc.2009.12.041

- [16] Grover L., Kaur P.(2011): An improved exponential estimator of finite population mean in simple random sampling using an auxiliary attribute. *Applied Mathematics and Computation* 218 (2011) 3093–3099.
- [17] Malik S., Singh R. (2012): A Family Of Estimators Of Population Mean Using Information On Point Bi-Serial and Phi Correlation Coefficient.
- [18] Sukhatme, P.V. and Sukhatme, B.V.,(1970) : *Sampling theory of surveys with applications*. Iowa State University Press, Ames, U.S.A.
- [19] Mukhopadhyaya, P.(2000): *Theory and methods of survey sampling*. Prentice Hall of India, New Delhi, India.

Two-Warehouse Fuzzy Inventory Model with K-Release Rule

Neeraj Kumar*, Sanjey Kumar* and Florentin Smarandache**

*Department of Mathematics, SRM University

Delhi NCR, Sonapat, Haryana

**Department of Mathematics, University of New Mexico

Gallup, NM 87301, USA

*Corresponding Author, sanjeysrm1984@gmail.com

ABSTRACT: Fuzzy set theory is primarily concerned with how to quantitatively deal with imprecision and uncertainty, and offers the decision maker another tool in addition to the classical deterministic and probabilistic mathematical tools that are used in modeling real-world problems. The present study investigates a fuzzy economic order quantity model for two storage facility. The demand, holding cost, ordering cost, storage capacity of the own - warehouse are taken as a trapezoidal fuzzy numbers. Graded Mean Representation is used to defuzzify the total cost function and the results obtained by this method are compared with the help of a numerical example. Sensitivity analysis is also carried out to explore the effect of changes in the values of some of the system parameters. The proposed methodology is applicable to other inventory models under uncertainty.

Keywords: Inventory, Two – warehouse system, Fuzzy Variable, Trapezoidal Fuzzy Number, Graded mean representation method and K – release rule.

1. INTRODUCTION

In most of the inventory models that had been proposed in the early literature, the associated costs are assumed to be precise, although the real-world inventory costs usually exist with imprecise components. In this case, customer demand as one of the key parameters and source of uncertainty have been most often treated by a probability distribution. However, the probability-based approaches may not be sufficient enough to reflect all uncertainties that may arise in a real-world inventory system. Modelers may face some difficulties while trying to build a valid model of an inventory system, in which the related costs cannot be determined precisely. For

example, costs may be dependent on some foreign monetary unit. In such a case, due to a change in the exchange rates, the costs are often not known precisely.

Fuzzy set theory, originally introduced by Zadeh [1], provides a framework for considering parameters that are vaguely or unclearly defined or whose values are imprecise or determined based on subjective beliefs of individuals. Petrovic et al. [2] presented newsboy problem assuming that demand and backorder cost are fuzzy numbers. Kaufmann and Gupta [3] introduced to fuzzy arithmetic: theory and application. The application of fuzzy theory to inventory problem has been proposed by Kacprzyk and Staniewski [4]. Roy and Maiti [5] presented a fuzzy inventory model with constraint. Roy and Maiti [6] developed a fuzzy EOQ model with demand-dependent unit cost under limited storage capacity. Ishii and Konno [7] introduced fuzziness of shortage cost explicitly into classical inventory problem. Chen and Hsieh [8] established a fuzzy economic production model to treat the inventory problem with all the parameters and variables, which are fuzzy numbers. Hsieh [9] presented a fuzzy production inventory model. Yao and Chiang [10] presented an inventory model without backorder with fuzzy total cost and fuzzy storing cost defuzzified by centroid and signed distance. Dutta et al. [11] developed a single-period inventory model with fuzzy random variable demand. In that study, they have applied graded mean integration representation method to find the optimum order quantity. Chen and Chang [12] presented an optimization of fuzzy production inventory model. In this study, they have used 'Function Principle' as arithmetical operations of fuzzy total production inventory cost and also used the 'Graded Mean Integration Representation method' to defuzzify the fuzzy total production and inventory cost. Mahata and Goswami [13] presented a fuzzy inventory model for deteriorating items with the help of fuzzy numbers and so on.

Most of the classical inventory models discussed in the literature deals with the situation of a single warehouse. Because of capacity limitation a single warehouse would not be always sufficient. Additional warehouse are necessary to store excess items. Therefore due to the limited capacity of the existing warehouse (Rented warehouse, RW) is acquired to keep excess items. In practice, large stock attracts the management due to either an attractive price discount for bulk purchase or the acquisition cost being higher than the holding cost in RW. The actual service to the customer is done at OW only. Usually the holding cost is greater in RW than in OW. So in order to reduce the holding cost. The stock of rented warehouse is transferred to the own warehouse. Hartley [14] was discussed a model under the assumption that the cost of transporting a unit from RW to OW is not significantly high. It was as the case with two levels of storage. Sarma [15] extended the model with two levels of storage given by Hartley, by

considering the transportation cost of a unit from rented warehouse to own warehouse. Maurdeswar and Sathe [16] discussed this model by relaxing the condition on production rate (finite production rate). Dave [17] considered it for finite and infinite replenishment, assuming the cost of transportation depending on the quantity to be transported. Pakkala and Achary [18] developed a model for deteriorating items with two warehouses. They extended it with bulk release rule, after words, Gowsami and Chaudhari [19] formulated models for time dependent demand. Kar et al. [20] suggested a two level inventory model for linear trend in demand. Yang [21] considered a two-warehouse inventory models for deteriorating items with shortages under inflation. Singh et al. [22] presented two-warehouse inventory model without shortage for exponential demand rate and an optimum release rule. Jaggi and Verma [23] developed a deterministic order level inventory model with two storage facilities. It has been observed in supermarkets that the demand rate is usually influenced by the amount of stock level, that is, the demand rate may go up or down with the on-hand stock level. Singh et al. [24] developed a deterministic two-warehouse inventory model for deteriorating items with stock-dependent demand and shortages. Neeraj et al. [25] developed three echelon supply chain inventory model with two storage facility. Neeraj et al. [26] presented a two-warehouse inventory model with K-release rule and learning effect. Neeraj et al. [27] considered effect of salvage value on a two-warehouse inventory model. Recently, Kumar and Kumar [28] developed an inventory model with stock dependent demand rate for deterioration items.

Here, in this paper the cost of transporting a unit is considered to be significant and the effect of releasing the stocks of RW in n shipments with a bulk size of K units per shipment, instead of withdrawing an arbitrary quantity, is assumed. Here, K is to be decided optimally and is call this as K-release rule. This problem is to decide the optimal values of Q and C, which minimize the sum of ordering, holding and transportation costs of the system. Here, we assumed that the storage capacity of the own – warehouse, the holding cost in both warehouses and ordering cost is fuzzy in nature. The associated total cost minimization is illustrated by numerical example and sensitivity analysis is carried out by using *MATHEMATICA*–5.2 for the feasibility and applicability of our model.

2. ASSUMPTIONS AND NOTATIONS:

The following assumptions are used to analyze this inventory model:

1. D is the constant demand rate.
2. W is the storage capacity of the OW.

3. A is the fixed set – up cost per order.
4. $C(Q)$ is the cost function.
5. Q is the highest inventory level.
6. H is the holding cost in OW.
7. F is the holding cost in RW.
8. \tilde{D} is the fuzzy demand rate.
9. \tilde{A} is the fuzzy set – up cost per order.
10. \tilde{H} is the fuzzy holding cost in OW.
11. \tilde{F} is the fuzzy holding cost in RW.
12. $\tilde{C}(Q)$ is the fuzzy cost function.
13. \tilde{W} is the fuzzy storage capacity of the OW.
14. The holding cost per unit in OW is higher than in RW.
15. The storage capacity of OW as W and that of RW is unlimited.
16. The transportation cost of K units from RW to OW is C_t at a time, which is constant over time.
17. The items of RW are transferred to OW in ‘ n ’ shipments of which K ($K \leq W$) units are transported in each shipment.
18. Replenishment rate is infinite.
19. Lead-time is zero.
20. Consumption takes place only in OW.

3. FUZZY SETS, MEMBERSHIP FUNCTION, DEFUZZIFYING APPROACH AND ARITHMETICAL OPERATIONS

3.1. Fuzzy Sets

A fuzzy set is a class of objects with a continuum of grades of membership. Such a set is characterized by a membership (characteristic) function which assigns to each object a grade of membership ranging between zero and one. Let $X=\{x\}$ denote a space of objects. Then a fuzzy set A in X is a set of ordered pairs:

$$A = \{ x, \mu_A(x) \}, x \in X$$

Where, $\mu_A(x)$ is termed “ the grade of the membership of x in A ”. For simplicity, $\mu_A(x)$ is a number in the interval $[0, 1]$, with the grades of unity and zero respectively, full membership and non-membership in the fuzzy set. An object (point) P contained in a set (class) Q is an element of Q ($P \subset Q$).

3.2. Membership Function

Membership Function




Fig. 1 Membership function for triangle number

At the outset it would be prudent introduce the concept of membership function. There are different shapes of membership function in the inventory control such as the triangle and trapezoid. The shapes of the triangle membership function and the trapezoid membership function are shown in Fig. 1 and 2.

Membership Function




Fig. 2 Membership function for trapezoid number

\tilde{A} is assumed as a fuzzy number. If \tilde{A} is a triangle number, \tilde{A} can be represented as $\tilde{A} = [k_1, k_2, k_3]$ subject to the constraint $0 < k_1 \leq k_2 \leq k_3$. While \tilde{A} is a trapezoid fuzzy number, $\tilde{A} = [k_1, k_2, k_3, k_4]$ subject to the constraint that $0 < k_1 \leq k_2 \leq k_3 \leq k_4$. Membership function of the triangle and trapezoid fuzzy numbers can be defined as follows:

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & x < k_1, x > k_3 \\ L(x) = \frac{x - k_1}{k_2 - k_1} & k_1 \leq x < k_2 \\ R(x) = \frac{k_3 - x}{k_3 - k_2} & k_2 \leq x < k_3 \end{cases}$$

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & x < k_1, x > k_4 \\ L(x) = \frac{x - k_1}{k_2 - k_1} & k_1 \leq x < k_2 \\ 1 & k_2 \leq x < k_3 \\ R(x) = \frac{k_4 - x}{k_4 - k_3} & k_3 \leq x \leq k_4 \end{cases}$$

where $\mu_{\tilde{A}}(x)$ is a membership function.

3.3. Graded Mean Integration Representation Method

In this study, generalized fuzzy number \tilde{A} was denoted in Fig. 6.1 as $\tilde{A} = (c, a, b, d, \omega_A)_{LR}$. When $\omega_A = 1$, we simplify the notation as $\tilde{A} = (c, a, b, d)_{LR}$. Chen and Hsieh (1999) introduced the graded mean integration representation method of generalized fuzzy number based on the integral value of graded mean h -level of generalized fuzzy number. Its meaning is as follows:

Let L^{-1} and R^{-1} are inverse function of L and R respectively, then the graded mean h -level value of generalized fuzzy number $\tilde{A} = (c, a, b, d, W_A)_{LR}$ is $h \left(L^{-1}(h) + R^{-1}(h) \right) / 2$ as Fig. 3.

Then the graded mean integration representation of \tilde{A} is

$$P(\tilde{A}) = \int_0^{W_A} \frac{h \left(L^{-1}(A) + R^{-1}(h) \right)}{2} dh \Bigg/ \int_0^{W_A} h dh ,$$

where $0 < h \leq W_A$ and $0 < W_A \leq 1$.




Fig. 3 The graded mean h -level of generalized fuzzy number $A = (c, a, b, d, W_A)_{LR}$

In the present, the generalized trapezoidal fuzzy number has been used as the type of all fuzzy parameters in our proposed inventory models. The very popular generalized trapezoidal fuzzy number B is a special case of generalized fuzzy number and can be denoted as $\tilde{B} = (c, a, b, d; W_B)$ its' corresponding graded mean integration representation is

$$P(\tilde{B}) = \int_0^{W_B} \frac{h(c+d + (a-c-d+b)h/W_B)dh}{2} \Bigg/ \int_0^{W_B} h dh = \frac{c+2a+2b+d}{6}$$

where a, b, c, d are any real numbers.

3.4. Properties of Second Function Principle

Chen (1985) proposed second function principal to be as the fuzzy arithmetical operations between generalized trapezoidal fuzzy numbers. Because it does not change the type of membership function of generalized fuzzy number after arithmetical operations. It reduces the trouble and tediousness of operations. Furthermore, Chen already proved the properties of fuzzy arithmetical operations under second function principle. Here some properties of the fuzzy arithmetical operations have been described as follows:

Suppose $\tilde{A}_1 = (c_1, a_1, b_1, d_1)$ and $\tilde{A}_2 = (c_2, a_2, b_2, d_2)$ are two generalized trapezoidal fuzzy numbers. Then

1. The addition of \tilde{A}_1 and \tilde{A}_2 is $\tilde{A}_1 \oplus \tilde{A}_2 = (c_1 + c_2, a_1 + a_2, b_1 + b_2, d_1 + d_2)$
2. The multiplication of \tilde{A}_1 and \tilde{A}_2 is $\tilde{A}_1 \otimes \tilde{A}_2 = (c_1 c_2, a_1 a_2, b_1 b_2, d_1 d_2)$
3. $-\tilde{A}_2 = (-d_2, -b_2, -a_2, -c_2)$ Then the subtraction of \tilde{A}_1 and \tilde{A}_2 is $\tilde{A}_1 \ominus \tilde{A}_2 = (c_1 - d_2, a_1 - b_2, b_1 - a_2, d_1 - c_2)$
4. $1/\tilde{A}_2 = A^{-1}_2 = \left(\frac{1}{d_2}, \frac{1}{b_2}, \frac{1}{a_2}, \frac{1}{c_2} \right)$ where c_2, a_2, b_2 and d_2 are all positive real numbers. If $c_1, a_1, b_1, d_1, c_2, a_2, b_2$ and d_2 are all non zero positive real numbers, then the division of \tilde{A}_1 and \tilde{A}_2 is $\tilde{A}_1 \oslash \tilde{A}_2 = \left(\frac{c_1}{d_2}, \frac{a_1}{b_2}, \frac{b_1}{a_2}, \frac{d_1}{c_2} \right)$.

4. MODEL DEVELOPMENT

Initially the company ordered Q units of the item, out of which W units is kept in OW and Z units are kept in RW, where $Z = (Q - W)$. Initially, demand is satisfied using the stocks of OW until the stock level drops to $(W-K)$ units. At this stage, K units from RW are transported to OW to meet further demand and this process is repeated 'n' times until the stocks of RW are exhausted. The remaining $(W-K)$ units in OW are used again at this stage. The inventory situation in RW and OW are shown in the figure 1.

The inventory units in RW can be seen to be equal.

$$A_t = t_{ik} \left[Z + (Z - K) + (Z - 2K) + \dots + (Z - (n-1)K) \right] = t_{ik} \frac{Z(n+1)}{2} \quad (4.1)$$

Where $t_{ik} = K/D$, the time taken for the consumption of K units, since $Z = (Q - W)$ and the holding cost in RW is F(i), we have-

$$FA_t = Ft_{ik} \frac{Z(n+1)}{2} = \frac{FK}{D} (Q - W) \frac{(n+1)}{2} = \frac{FK(n+1)(Q - W)}{2D} \quad (4.2)$$

The cost of transporting the units from RW to OW in 'n' shipments is given by

$$nC_t = (Z/K)C_t \tag{4.3}$$

Since $n = Z/K$




Fig. 1: Graphical representation of two-storage inventory model

When K units are drawn from RW in each shipment, more are carried in OW for a period of t_k and hence account for a holding cost of $KH (i) t_k / 2$. Since there are 'n' such shipments and taking into consideration, the initial K units of OW, the holding cost for these items is $(n+1)HKt_k/2 = (n+1)HK^2/2D$ (4.4)

A quantity of $(W - K)$ units is kept unused in OW for a period of $t_{i(W - K)} = (n+1)t_k$ and an average inventory during usage in OW is $(W - K)/2$ units for a period $(t - t_{i(W - K)})$. Hence the inventory holding cost in OW for these items is

$$H[K(W-K)(n+1)/D + (W - K)^2/2D]. \tag{4.5}$$

The fixed ordering cost per order is A . Then the total inventory cost for the system using (4.2) to (4.5) becomes

$$C = A + (n+1) \frac{FK(Q-W)}{2D} + \frac{K^2H}{2D} + \frac{HK(W-K)}{2D} + nC_t + \frac{(W-K)^2 H}{2D} \quad (4.6)$$

The average inventory cost

$$C(Q, K) = C / t$$

But we have $t = Q / D$, $Z = Q - W$ and $n = Z / K = Q - W/K$

Total average cost becomes

$$C(Q, K) = \frac{AD}{Q} + \frac{FQ}{2} - W(F-H) + \frac{K}{2}(F-H) - \frac{KW}{2Q}(F-H) + C_t \frac{(Q-W)D}{QK} + \frac{W^2}{2Q}(F-H) \quad (4.7)$$

Fuzzy Model: Due to uncertainly in the environment it is not easy to define all the parameters precisely, accordingly we assume some of these parameters namely \tilde{D} , \tilde{F} , \tilde{H} , \tilde{A} and \tilde{W} may change within some limits. Let $\tilde{D} = [d_1, d_2, d_3, d_4]$, $\tilde{F} = [f_1, f_2, f_3, f_4]$, $\tilde{H} = [h_1, h_2, h_3, h_4]$, $\tilde{A} = [a_1, a_2, a_3, a_4]$, $\tilde{W} = [w_1, w_2, w_3, w_4]$ are as trapezoidal fuzzy numbers. In this case, the total fuzzy cost per unit time is given by

$$\begin{aligned} \tilde{C}(Q, K) = & \left((\tilde{A} \otimes \tilde{D}) \oslash Q \right) \oplus \left((\tilde{F} \otimes Q) \oslash 2 \right) \ominus \left(\tilde{W} \otimes (\tilde{F} \oslash \tilde{H}) \right) \oplus \left((K \otimes (\tilde{F} \oslash \tilde{H})) \oslash 2 \right) \ominus \left((K \otimes (\tilde{W} \otimes (\tilde{F} \oslash \tilde{H}))) \oslash 2Q \right) \\ & \oplus \left((C_t \otimes (Q \oslash \tilde{W})) \otimes \tilde{D} \right) \oslash QK \oplus \left((\tilde{W} \otimes (\tilde{W} \otimes (\tilde{F} \oslash \tilde{H}))) \oslash 2Q \right) \end{aligned} \quad (4.8)$$

By second function principal, one has

$$\begin{aligned} \tilde{C}(Q, K) = & \left(\frac{a_1 d_1}{Q} + \frac{f_1 Q}{2} - (f_4 - h_1) w_4 + \frac{K(f_1 - h_4)}{2} - \frac{K(f_4 - h_1) w_4}{2Q} + \frac{C_t(Q - w_4) d_1}{QK} + \frac{(f_1 - h_4) w_1^2}{2Q}, \right. \\ & \frac{a_2 d_2}{Q} + \frac{f_2 Q}{2} - (f_3 - h_2) w_3 + \frac{K(f_2 - h_3)}{2} - \frac{K(f_3 - h_2) w_3}{2Q} + \frac{C_t(Q - w_3) d_2}{QK} + \frac{(f_2 - h_3) w_1^2}{2Q}, \\ & \frac{a_3 d_3}{Q} + \frac{f_3 Q}{2} - (f_2 - h_3) w_2 + \frac{K(f_3 - h_2)}{2} - \frac{K(f_2 - h_3) w_2}{2Q} + \frac{C_t(Q - w_2) d_3}{QK} + \frac{(f_3 - h_2) w_1^2}{2Q}, \\ & \left. \frac{a_4 d_4}{Q} + \frac{f_4 Q}{2} - (f_1 - h_4) w_1 + \frac{K(f_4 - h_1)}{2} - \frac{K(f_1 - h_4) w_1}{2Q} + \frac{C_t(Q - w_1) d_4}{QK} + \frac{(f_4 - h_1) w_1^2}{2Q} \right) \end{aligned}$$

Now we defuzzify the total cost per unit time, using graded mean integration representation method, the result is

$$\begin{aligned}
 P(\tilde{C}(Q,K)) = & \frac{1}{6} \left[\left(\frac{a_1 d_1}{Q} + \frac{f_1 Q}{2} - (f_4 - h_1) w_4 + \frac{K(f_1 - h_4)}{2} - \frac{K(f_4 - h_1) w_4}{2Q} + \frac{C_t(Q - w_4) d_1}{QK} + \frac{(f_1 - h_4) w_1^2}{2Q} \right. \right. \\
 & + 2 \left(\frac{a_2 d_2}{Q} + \frac{f_2 Q}{2} - (f_3 - h_2) w_3 + \frac{K(f_2 - h_3)}{2} - \frac{K(f_3 - h_2) w_3}{2Q} + \frac{C_t(Q - w_3) d_2}{QK} + \frac{(f_2 - h_3) w_1^2}{2Q} \right) \\
 & + 2 \left(\frac{a_3 d_3}{Q} + \frac{f_3 Q}{2} - (f_2 - h_3) w_2 + \frac{K(f_3 - h_2)}{2} - \frac{K(f_2 - h_3) w_2}{2Q} + \frac{C_t(Q - w_2) d_3}{QK} + \frac{(f_3 - h_2) w_1^2}{2Q} \right) \\
 & \left. \left. + \frac{a_4 d_4}{Q} + \frac{f_4 Q}{2} - (f_1 - h_4) w_1 + \frac{K(f_4 - h_1)}{2} - \frac{K(f_1 - h_4) w_1}{2Q} + \frac{C_t(Q - w_1) d_4}{QK} + \frac{(f_4 - h_1) w_1^2}{2Q} \right) \right] \quad (4.9)
 \end{aligned}$$

The optimal values of Q and K, which minimizes (4.8), are obtained by solving

$$\frac{\partial P(\tilde{C}(Q,K))}{\partial Q} = 0 \text{ and } \frac{\partial P(\tilde{C}(Q,K))}{\partial K} = 0 \quad (4.10)$$

we get

$$Q = \left[\frac{2a_1 d_1 + 4a_2 d_2 + 4a_3 d_3 + 2a_4 d_4 - K[(f_4 - h_1) w_4 + (f_3 - h_2) w_3 + (f_2 - h_3) w_2 + (f_1 - h_4) w_1]}{-\frac{2C_t}{K}(w_4 d_1 + w_3 d_2 + w_2 d_3 + w_1 d_4) + (f_4 - h_1) w_4^2 + (f_3 - h_2) w_3^2 + (f_2 - h_3) w_2^2 + (f_1 - h_4) w_1^2} \right]^{1/2} \quad (4.11)$$

and

$$K = \sqrt{\frac{\frac{C_t}{Q}[(Q - w_4) + 2(Q - w_3) + 2(Q - w_2) + (Q - w_1)]}{\frac{(f_1 - h_4)}{2} + (f_2 - h_3) + (f_3 - h_2) + \frac{(f_4 - h_1)}{2}} - \frac{(f_1 - h_4) w_1}{2Q} - \frac{(f_2 - h_3) w_2}{Q} - \frac{(f_3 - h_2) w_3}{Q} - \frac{(f_4 - h_1) w_4}{2Q}}{\quad}} \quad (4.12)$$

5. COST-REDUCTION DUE TO K-RELEASE RULE

The unit cost of transportation with K-release rule is $C_t' = C_t/K$. Suppose the unit cost of transportation is C_t^* without bulk transportation. The bulk transportation will be economical only if $C_t^* > C_t'$. Hence without K-release rule, the cost function becomes-

$$C(Q) = \frac{AD}{Q} + \frac{FQ}{2} + \frac{W^2(F-H)}{2Q} - W(F-H) + \frac{(Q-W)C_t^*D}{Q} \quad (5.1)$$

Fuzzy Model: Due to uncertainty in the environment it is not easy to define all the parameters precisely, accordingly we assume some of these parameters namely $\tilde{D}, \tilde{F}, \tilde{H}, \tilde{A}$ and \tilde{W} may change within some limits.

Let $\tilde{D} = [d_1, d_2, d_3, d_4], \tilde{F} = [f_1, f_2, f_3, f_4], \tilde{H} = [h_1, h_2, h_3, h_4], \tilde{A} = [a_1, a_2, a_3, a_4],$
 $\tilde{W} = [w_1, w_2, w_3, w_4]$ are as trapezoidal fuzzy numbers. In this case, the total fuzzy cost per unit time is given by

$$\begin{aligned} \tilde{C}(Q) = & \left((\tilde{A} \otimes \tilde{D}) \oslash Q \right) \oplus \left((\tilde{F} \otimes Q) \oslash 2 \right) \oplus \left((\tilde{W} \otimes (\tilde{W} \otimes (\tilde{F} \oslash \tilde{H}))) \oslash 2Q \right) \oslash (\tilde{W} \otimes (\tilde{F} \oslash \tilde{H})) \\ & \oplus \left(\left((Q \oslash \tilde{W}) \otimes C_t^* \right) \otimes \tilde{D} \right) \oslash Q \end{aligned} \quad (5.2)$$

By second function principal, one has

$$\begin{aligned} \tilde{C}(Q) = & \left(\frac{a_1 d_1}{Q} + \frac{f_1 Q}{2} - (f_4 - h_1) w_4 + \frac{C_t^* (Q - w_4) d_1}{Q} + \frac{(f_1 - h_4) w_1^2}{2Q}, \right. \\ & \frac{a_2 d_2}{Q} + \frac{f_2 Q}{2} - (f_3 - h_2) w_3 + \frac{C_t^* (Q - w_3) d_2}{Q} + \frac{(f_2 - h_3) w_1^2}{2Q}, \\ & \frac{a_3 d_3}{Q} + \frac{f_3 Q}{2} - (f_2 - h_3) w_2 + \frac{C_t^* (Q - w_2) d_3}{Q} + \frac{(f_3 - h_2) w_1^2}{2Q}, \\ & \left. \frac{a_4 d_4}{Q} + \frac{f_4 Q}{2} - (f_1 - h_4) w_1 + \frac{C_t^* (Q - w_1) d_4}{Q} + \frac{(f_4 - h_1) w_1^2}{2Q} \right) \end{aligned}$$

Now we defuzzify the total cost per unit time, using graded mean integration representation method, the result is

$$P(\tilde{C}(Q)) = \frac{1}{6} \left\{ \begin{aligned} & \left(\frac{a_1 d_1}{Q} + \frac{f_1 Q}{2} - (f_4 - h_1) w_4 + \frac{C_t^* (Q - w_4) d_1}{Q} + \frac{(f_1 - h_4) w_1^2}{2Q} \right) \\ & + 2 \left(\frac{a_2 d_2}{Q} + \frac{f_2 Q}{2} - (f_3 - h_2) w_3 + \frac{C_t^* (Q - w_3) d_2}{Q} + \frac{(f_2 - h_3) w_1^2}{2Q} \right) \\ & + 2 \left(\frac{a_3 d_3}{Q} + \frac{f_3 Q}{2} - (f_2 - h_3) w_2 + \frac{C_t^* (Q - w_2) d_3}{Q} + \frac{(f_3 - h_2) w_1^2}{2Q} \right) \\ & + \left(\frac{a_4 d_4}{Q} + \frac{f_4 Q}{2} - (f_1 - h_4) w_1 + \frac{C_t^* (Q - w_1) d_4}{Q} + \frac{(f_4 - h_1) w_1^2}{2Q} \right) \end{aligned} \right\} \quad (5.3)$$

The optimal value of Q, which minimizes (5.1), is obtained by $\frac{dP(\tilde{C}(Q))}{dQ} = 0$

$$Q = \left[\frac{-a_1 d_1 - 2a_2 d_2 - 2a_3 d_3 - a_4 d_4 + C_t (w_4 d_1 + 2w_3 d_2 + 2w_2 d_3 + w_1 d_4)}{-\frac{(f_4 - h_1)}{2} w_4^2 - (f_3 - h_2) w_3^2 - (f_2 - h_3) w_2^2 - \frac{(f_1 - h_4)}{2} w_1^2} \right]^{1/2} \\ - \left(\frac{f_1}{2} + f_2 + f_3 + \frac{f_4}{2} \right) \quad (5.4)$$

The proposed K-released rule will be economical if

$$[C(Q) - C(Q, K)] > 0$$

From equation (4.7) and (4.8) we see that-

$$[C(Q) - C(Q, K)] = \left(1 - \frac{W}{Q} \right) \left[D(C_t^* - C_t') - \frac{K}{2}(F - H) \right] \quad (5.5)$$

and hence the inequality

$$\left(1 - \frac{W}{Q} \right) \left[D(C_t^* - C_t') - \frac{K}{2}(F - H) \right] > 0 \\ \Rightarrow (C_t^* - C_t') > \frac{K(F - H)}{2D} \quad (5.6)$$

must be satisfied.

Thus for a given situation, if the unit cost of transportation with bulk release rule satisfies the inequality (5.6), K-release rule must be economical.

6. NUMERICAL EXAMPLE

Consider an inventory system with following parametric values:

Crisp Model: demand rate $D = 2000$, $C_t = 0.5$, $F = 8.5$, $H = 7.5$, $W = 100$, $A = 150$. With the help of the above values, we find the optimal values of ordering quantity and total cost with and without K- release which is given as:

With K – release rule: $Q = 221.62$ & $C(Q, K) = 3456.46$

And without K – release rule: $Q = 216.68$ & $C(Q, K) = 3585.43$

Fuzzy Model: $\tilde{D} = [1900, 2000, 2000, 1900]$, $\tilde{F} = [8.075, 8.5, 8.5, 8.075]$, $\tilde{H} = [7.125, 7.5, 7.5, 7.125]$, $\tilde{A} = [142.5, 150, 150, 142.5]$, $\tilde{W} = [95, 100, 100, 95]$. The optimal values of ordering quantity and total cost with and without K- release which is given as:

With K – release rule: $Q = 225.62$ & $C(Q, K) = 3458.46$

And without K – release rule: $Q = 210.68$ & $C(Q, K) = 3587.43$

7. CONCLUSION

Two storage inventory models discussed in this paper and developed under the assumption that the distribution of the items to the customers takes place at OW only. Because of the distance factor, it is natural to consider the transportation cost associated with the transfer of items from RW to OW. Further, the concept of K-release rule is more pragmatic, as holding large inventory in RW is every expensive. With the help of numerical examples, it is clear that the effect of fuzzy cannot be ignored. We can earn more profit by consider the effect of fuzzy on ordering and holding cost in each lot. This model gives the direction to decision makers to take account of fuzzy effect while taking decision and by taking account of this; he/she earn more profit for the organization.

A future extension is to discuss model in more realistic situation by consider impreciseness in different inventory related cost and taking different form of demand pattern likes as time dependent, ram-type demand with inflation and permissible delay.

8. REFERENCE

1. Zadeh, L.A. (1965) 'Fuzzy sets', *Information and Control*, Vol. 8, pp.338–353.
2. Petrovic, D., Petrovic, R. and Vujosevic, M. (1996) 'Fuzzy models for the newsboy problem', *Int. J. Production Economics*, Vol. 45, pp.435–441.
3. Kaufmann, A. and Gupta, M.M. (1985) *Introduction to Fuzzy Arithmetic: Theory and Application*. New York: Van Nostrand Reinhold.
4. Kacprzyk, J. and Staniewski, P. (1982) 'Long term inventory policy-making through fuzzy decision-making model', *Fuzzy Sets and Systems*, Vol. 8, No. 2, pp.117–132.
5. Roy, T.K. and Maiti, M. (1995) 'A fuzzy inventory model with constraint', *Opsearch*, Vol. 32, No. 4, pp.287–298.
6. Roy, T.K. and Maiti, M. (1997) 'Fuzzy EOQ model with demand-dependent unit cost under limited storage capacity', *European Journal of Operational Research*, Vol. 99, pp.425–432.
7. Ishii, H. and Konno, T. (1998) 'A stochastic inventory problem with fuzzy shortage cost', *European Journal of Operational Research*, Vol. 106, pp.90–94
8. Hsieh, C.H. (2002) 'Optimization of fuzzy production inventory models', *Information Sciences*, Vol. 146, pp.29–40.
9. Dutta, P., Chakraborty, D. and Roy, A.R. (2005) 'A single-period inventory model with fuzzy random variable demand', *Mathematical and Computer Modelling*, Vol. 41, pp.915–922.
10. Yao, J.S. and Chiang, J. (2003) 'Inventory without backorder with fuzzy total cost and fuzzy storing cost defuzzified by centroid and signed distance', *European Journal of Operational Research*, Vol. 148, pp.401–409.
11. Chen, S.H. and Chang, S.M. (2008) 'Optimization of fuzzy production inventory model with unrepairable defective products', *Int. J. Production Economics*, Vol. 113, No. 2, pp.887–894
12. Mahata, G.C. and Goswami, A. (2009) 'A fuzzy replenishment policy for deteriorating items with ramp type demand rate under inflation', *Int. J. Operational Research*, Vol. 5, No. 3, pp.328–348.
13. Hartley, R.V. (1976) *Operations Research – A Managerial Emphasis*, pp.315–317, Good Year Publishing Company, California.
14. Sarma, K.V.S. (1983) 'A deterministic inventory model with two levels of storage and an optimum release rule', *Opsearch*, Vol. 20, No. 3, pp.174–180.
15. Murdeshwar, T.A. and Sathe, Y.S. (1985) 'Some aspects of lot size model with two levels of storage', *Opsearch*, Vol. 22, No. 1, pp.255–262.
16. Dave, U. (1988) 'On the EOQ models with two level of storage', *Opsearch*, Vol. 25, No. 3, pp.190–196.
17. Pakkala, T.M.P. and Achary, K.K. (1992) 'A deterministic inventory model for deteriorating items with two warehouses and finite replenishment rate', *European Journal of Operational Research*, Vol. 57, No. 1, pp.71–76.

18. Goswami, A. and Chaudhur, K.S. (1992) 'An economic order quantity model for items with two levels of storage for a linear trend in demand', *Journal of Operational Research Society*, Vol. 43, No. 2, pp.157–167.
19. Kar, S.K., Bhunia, A.K. and Maiti, M. (2001) 'Deterministic inventory model with two levels of storage, a linear trend in demand and a fixed time horizon', *Computer and Operations Research*, Vol. 28, No. 13, pp.1315–1331.
20. Yang, H.L. (2004) 'Two-warehouse inventory models for deteriorating items with shortages under inflation', *European Journal of Operational Research*, Vol. 157, No. 2, pp.344–356.
21. Singh, S.R., Kumar, N. and Kumari, R. (2008) 'Two-warehouse inventory model without shortage for exponential demand rate and an optimum release rule', *Ultra Scientist of Physical Sciences*, Vol. 20, No. 2-M, pp.395–402.
22. Jaggi, C.K. and Verma, P. (2010) 'A deterministic order level inventory model for deteriorating items with two storage facilities under FIFO dispatching policy', *International Journal of Procurement Management*, Vol. 3, No. 3, pp.265–278.
23. Singh, S.R., Kumari, R. and Kumar, N. (2011) 'A deterministic two-warehouse inventory model for deteriorating items with stock-dependent demand and shortages under the conditions of permissible delay', *International Journal of Mathematical Modelling and Numerical Optimisation*, Vol. 11, No. 3, pp.357–375.
24. Kumar, N., Singh, S. R., Kumari, R. (2012) 'Three echelon supply chain inventory model for deteriorating items with limited storage facility and lead-time under inflation', *Int. J. Services and Operations Management*, Vol. 13, No. 1, pp. 98 – 118.
25. Kumar, N., Singh, S. R., Kumari, R. (2013) 'Two-warehouse inventory model with K-release rule and learning effect', *Int. J. Procurement Management*, Vol. 6, No.1, pp. 76 – 91.
26. Kumar, N., Singh, S. R., Kumari, R. (2014) 'Effect of salvage value on a two-warehouse inventory model for deteriorating items with stock-dependent demand rate and partial backlogging', *Int. J. Operational Research*, Vol. 19, No. 4, pp. 479 – 494.
27. Kumar N and Kumar S (2016) Effect of learning and salvage worth on an inventory model for deteriorating items with inventory-dependent demand rate and partial backlogging with capability constraints. *Uncertain Supply Chain Management* 4: 123-136

A Two-Warehouse Inventory Model for Deteriorating Items with Stock Dependent Demand, Inflation and Genetic Algorithm

Sanjeey Kumar¹, Pallavi Agarwal² & Neeraj Kumar^{3*}

^{1,3} Department of Mathematics, SRM University, Sonapat, Haryana

² Department of Computer Science & Engineering, SRM University, Sonapat, Haryana

*Corresponding Author, nkneerajkapil@gmail.com

Abstract

In this article, the a two-warehouse inventory model deals with deteriorating items, with stock dependent demand rate and model affected by inflation under the pattern of time value of money over a finite planning horizon. Shortages are allowed and partially backordered depending on the waiting time for the next replenishment. The purpose of this model is to minimize the total inventory cost by using Genetic algorithm. Also, a numerical example along with sensitivity analysis is given to explore the model numerically. Some observations are presented on the basis of sensitivity analysis.

Keywords: Two-warehouse, Genetic algorithm, partial backlogging, stock-dependent demand, Inflation, Deterioration, shortages;

1. Introduction

In the busy markets like super market, municipality market etc. the storage area of items is limited. When an attractive price discount for bulk purchase is available or the cost of procuring goods is higher than the other inventory related cost or demand of items is very high or there are some problems in frequent procurement, management decide to purchase a large amount of items at a time. These items cannot be accommodated in the existing storehouse (viz. the Own Warehouse, OW) located at busy market place. In the present senerio, suppliers proposed price discounts for gathering purchases or if the goods are seasonal, the retailers possibly will buy the superfluous goods that can be stored in own warehouse (OW). And rented warehouse (RW) is used as a store over the certain capacity W_1 of the own warehouse. Generally, the rented warehouse may have a costly superior unit holding cost than the own warehouse due to surplus cost of maintenance, material handling, etc. Hartely [1] was the original instigator to consider the

impact of a two-warehouse model in inventory research and developed an inventory model with a RW storage principle. Sarma [2] developed a two-warehouse model for deteriorating items with an infinite replenishment rate and shortages. Sarma and Sastry [3] introduced a deterministic inventory model with an infinite production rate, permissible shortage and two levels of storage. Pakkala and Achary [4] considered a two-warehouse model for deteriorating items with finite replenishment rate and shortages. Lee and Ma [5] compared an optimal inventory policy for deteriorating items with two-warehouse and time-dependent demand. Yang [6] produced a two-warehouse inventory model with constant deteriorating items, constant demand rate and shortages under inflation. Yang [7] investigated the two-warehouse partial backlogging inventory models for deteriorating items under inflation. Kumar et.al [8] developed a Two-Warehouse inventory model without shortage for exponential demand rate and an optimum release rule. Kumar et al. [9] produced a Deterministic Two-warehouse Inventory Model for Deteriorating Items with Stock-dependent Demand and Shortages under the conditions of permissible delay. [10] Analyzed two-warehouse partial backlogging inventory models with three-parameter Weibull distribution deterioration under inflation. Sett et al. [11] introduced a two warehouse inventory model with increasing demand and time varying deterioration. Kumar et al. [12] developed Learning effect on an inventory model with two-level storage and partial backlogging under inflation. Yang and Chang [13] perused a two-warehouse partial backlogging inventory model for deteriorating items with permissible delay in payment under inflation. Guchhaita et al. [14] investigated a two storage inventory model of a deteriorating item with variable demand under partial credit period. Kumar and Singh [15] discussed Effect of Salvage Value on a Two-Warehouse Inventory Model for Deteriorating Items with Stock-Dependent Demand Rate and Partial Backlogging. Deterioration performed a most important contribution in lots of inventory systems. Normally, an inventory model understands with non-deteriorating items and instantaneous deteriorating items. Major part of goods undergo, waste or deterioration over time, examples being medicines, volatile liquids, blood banks, and so on. Therefore, waste or deterioration of physical goods in stock is a more realistic factor and there is a big need to consider inventory modeling. The primary effort to describe the optimal ordering policies for such items was prepared by Ghare and Schrader [16]. They presented an EOQ model for an exponentially decaying inventory. Philip [17] developed an inventory model with three parameter Weibull distribution rate without considering shortages. Deb and Chaudhari [18] derived inventory model with time-dependent deterioration rate. A meticulous assessment of deteriorating inventory literatures is given by Goyal and Giri [19]. Liao [20] studied an EOQ model with non- instantaneous receipt and exponential deteriorating item under two level trade

credits. Chung [21] derived a complete proof on the solution procedure for non-instantaneous deteriorating items with permissible delay in payment. Chang et al. [22] framed optimal replenishment policies for non-instantaneous deteriorating items with stock-dependent demand. Dye [23] investigated the effect of .0preservation technology investment in a non-instantaneous deteriorating inventory model.

Due to high inflation and consequent sharp decline in the purchasing power of money in the developing countries like Brazil, Argentina, India, Bangladesh etc., the financial situation has been completely changed and so it is not possible to ignore the effect of inflation and time value of money any further. Following Buzacott [24] and Misra [25] have extended their approaches to different inventory models by considering the time value of money, different inflation rates for the internal and external costs, finite replenishment, shortages, etc. Datta and Pal [26] considered the effects of inflation and time value of money of an inventory model with a linear time-dependent demand rate and shortages. Sarker and Pan [27] considered a finite replenishment model when the shortage is allowed. Chung [28] developed an algorithm with finite replenishment and infinite planning horizon. Tolgari et al. [29] studied an inventory model for imperfect items under inflationary conditions by considering inspection errors. Guria et al. [30] formulated an inventory policy for an item with inflation induced purchasing price, selling price and demand with immediate part payment. In the case of perishable product, the retailer may need to backlog demand to avoid costs due to deterioration. When the shortage occurs, some customers are willing to wait for back order and others would turn to buy from other sellers. Inventory model of deteriorating items with time proportional backlogging rate has been developed by Dye et al. [31]. Wang [32] studied shortages and partial backlogging of items. Recently, Kumar and Kumar [33] developed an inventory model with stock dependent demand rate for deterioration items.

In this study, we have developed an inventory model for non-instantaneous deteriorating items with stock-dependent under the impact of inflation with genetic algorithm. Shortages are allowed and partially backordered depending on the waiting time for the next replenishment. The main objective of this work is minimizing the total inventory cost and finding the new optimal interval and the optimal order quantity. The model shows the effect of the genetic algorithm due to changes in various parameters by taking suitable numerical examples and sensitivity analysis.

2. Notations and Assumptions

2.1 Notations

The following notations are used throughout this paper:

- A The ordering cost per order

- C_{hr} The ordering cost per item in RW
- C_{ho} The holding cost per item in OW, $C_{hr} > C_{ho}$
- C_2 The deterioration cost per unit per unit cycle
- C_3 The shortage cost for backlogged items per unit per unit cycle
- C_4 The unit cost of lost sales per unit per cycle
- p The purchasing cost per unit
- s The selling price per unit, with $s > p$
- μ_1 The life time of the items in OW
- μ_2 The life time of the items in RW, $\mu_1 < \mu_2$
- α The deterioration rate in OW, $0 \leq \alpha < 1$
- β The deterioration rate in RW, $0 \leq \beta < 1, \alpha > \beta$
- T The length of the order cycle (decision variable)
- H The planning horizon
- m The number of replenishment during planning horizon, $m = H/T$ (decision variable)
- W_1 The capacity of OW
- W_2 The maximum inventory level in RW (decision variable)
- S The maximum inventory level per cycle (decision variable)
- Bl The maximum amount of shortage demand to be backlogged (decision variable)
- Q The $2^{nd}, 3^{rd}, \dots, m^{th}$ order size (Decision variable)
- r The discount rate represents the time value of money.
- f The inflation rate
- R The net discount rate of inflation i.e. $R = r - f$
- $q_r(t)$ The inventory level in RW at time t
- $q_o(t)$ The inventory level in OW at time t
- $q_s(t)$ The negative inventory level at time t
- T_j The total time that elapsed upto and including the j th replenishment cycle ($j = 1, 2, 3, \dots$)
- t_r Length of period during which inventory level reaches to zero in RW

t_j The time at which the inventory level in OW in the j^{th} replenishment cycle drop to zero
($j = 1, 2, \dots, m$).

$T_j - t_j$ The time period when shortage occurs ($j = 1, 2, \dots, m$)

TC_j The total cost for first replenishment cycle

TC The total cost of the system over a finite planning horizon H

2.2 Assumptions:

To develop the mathematical model, the following assumptions are being made:

1. A single item is considered over the prescribed period of planning horizon.
2. There is no replacement or repair of deteriorated items takes place in a given cycle.
3. The lead time is zero.
4. Deterioration takes place after the life time of items. That is, during the fixed period, the Product has no deterioration. After that, it will deteriorate with constant rate.
5. The replenishment takes place at an infinite rate.
6. The effects of inflation and time value of money are considered.
7. The demand rate ($a + bq_r(t)$) is a stock dependent.
8. Shortages are allowed and partially backlogged. During the stock out period, the backlogging rate is variable and is dependent on the length of the waiting time for the next replenishment. So the Backlogging rate of negative inventory is, $1 / (1 + \delta(T - t))$, where δ is backlogging parameter $0 \leq \delta \leq 1$ and $(T - t)$ is waiting time ($t_j \leq t \leq T$), ($j = 1, 2, \dots, m$). The remaining fraction $(1 - B(t))$ is lost.
9. The OW has limited capacity of W_1 units and the RW has unlimited capacity. For economic reasons, the items of RW are consumed first and next the items of OW.

3. Formulation and solution of the model

Suppose with the purpose of the planning horizon H is divided into m equal parts of length $T = H/m$. Hence the reorder times over the planning horizon H are $T_j = jT$ ($j=0, 1, 2, \dots, m$). When the inventory is positive, demand rate is stock dependent, whereas for negative inventory, the demand is partially backlogged. The period for which there is no-shortage in each interval $[jT, (j+1)T]$ is a fraction of the scheduling period T and is equal to kT ($0 < k < 1$). Shortages occur at time $t_j = (k+j-1)T$, ($j = 1, 2, \dots, m$) and are build up until time $t = jT$ ($j = 1, 2, \dots, m$) before they are backordered. This model is demonstrated in Figure-1. The first replenishment lot size of S is replenished at $T_0 = 0$. W_1 units are kept in OW and the rest is stored in RW. The items of OW are consumed only after consuming the goods kept in RW. In the RW, during the time interval $[0,$

μ_2], the inventory level is decreasing only due to demand rate and the inventory level is dropping to zero owing to demand and deterioration during the time interval $[\mu_2, t_r]$. In OW, during the time interval $[0, \mu_1]$, there is no change in the inventory level. However, the inventory W_1 decreases during $[\mu_1, t_r]$ due to deterioration only, but during $[t_r, t_1]$, the inventory is depleted due to both demand and deterioration. By the time t_1 , both warehouses are empty. Finally, during the interval $[t_1, T]$, shortages occur and accumulate until $t = T_1$ before they are partially backlogged.




Fig. 3.1. Graphical representation of the two warehouse inventory system

Based on the above explanation during the time interval $[0, \mu_2]$, the inventory level in RW is decreasing only due to demand rate and the differential equation representing the inventory status is given by

$$\frac{dq_r(t)}{dt} = -(a + bq_r(t)) \quad 0 \leq t \leq \mu_2 \quad (3.1)$$

With the condition $q_r(0) = W_2$, the solution of equation (3.1) is

$$q_r(t) = \frac{-a}{b} + \left(\frac{a}{b} + W_2 \right) e^{-bt} \quad (3.2)$$

In the second interval $[\mu_2, t_r]$ in RW, the inventory level decreases due to demand and deterioration. Thus, the differential equation below represents the inventory status

$$\frac{dq_r(t)}{dt} + \beta q_r(t) = -(a + bq_r(t)) \quad \mu_2 \leq t \leq t_r \quad (3.3)$$

$$\frac{dq_r(t)}{dt} + \beta q_r(t) + bq_r(t) = -a \quad \mu_2 \leq t \leq t_r \quad (3.4)$$

With the condition $q_r(t_r) = 0$, we get the solution of equation (3.4) is

$$q_r(t) = \frac{-a}{\beta + b} + \frac{a}{\beta + b} e^{(\beta+b)(t_r-t)} \quad \mu_2 \leq t \leq t_r \quad (3.5)$$

Put $t = \mu_2$ in equations (3.2) and (3.5) we get the value of W_2 as

$$W_2 = \frac{a\beta}{b(\beta + b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r-\mu_2)+b\mu_2} \right) \right\} \quad (3.6)$$

Putting the value of W_2 in equation (3.2) we get

$$q_r(t) = \frac{a}{b} [-1 + e^{-bt}] + \frac{a\beta}{b(\beta + b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r-\mu_2)+b\mu_2} \right) \right\}, \quad 0 \leq t \leq \mu_2 \quad (3.7)$$

In OW, during the interval $[0, \mu_1]$, there is no change in the inventory level and during $[\mu_1, t_r]$ the inventory W_1 only decreases due to deterioration.

Therefore the rate of change in the inventory is given by

$$\frac{dq_o(t)}{dt} = 0 \quad 0 \leq t \leq \mu_1 \quad (3.8)$$

$$\frac{dq_o(t)}{dt} + \alpha q_o(t) = 0 \quad \mu_1 \leq t \leq t_r \quad (3.9)$$

With the conditions $q_0(0) = W_1$ and $q_0(\mu_1) = W_1$, the solutions of equations (3.8) and (3.9) are

$$q_o(t) = W_1 \quad 0 \leq t \leq \mu_1 \quad (3.10)$$

$$q_o(t) = W_1 e^{\alpha(\mu_1-t)} \quad \mu_1 \leq t \leq t_r \quad (3.11)$$

In the interval $[t_r, t_1]$ in own warehouse, the inventory level decreases due to demand and deterioration. Thus, the differential equation is

$$\frac{dq_o(t)}{dt} + \alpha q_o(t) = -(a + bq_r(t)) \quad t_r \leq t \leq t_1 \quad (3.12)$$

With the condition $q_o(t_1) = 0$, we get the solution of equation (3.12) is

$$q_o(t) = \frac{a}{\alpha + b} \left(e^{(\alpha+b)(t_1-t)} - 1 \right) \quad t_r \leq t \leq t_1 \quad (3.13)$$

Put $t = t_r$ in equations (3.11) and (3.13) we get,

$$t_r = \frac{\left(W_1 \alpha e^{\alpha \mu_1} + a e^{(\alpha+b)t_1} \right) \pm \sqrt{\left(W_1 \alpha e^{\alpha \mu_1} + a e^{(\alpha+b)t_1} \right)^2 + \frac{4a\alpha b}{\alpha + b} \left(W_1 e^{\alpha \mu_1} - \frac{a}{\alpha + b} e^{(\alpha+b)t_1} + \frac{a}{\alpha + b} \right)}}{2 \left(\frac{-a\alpha b}{\alpha + b} \right)} \quad (3.14)$$

During the interval $[t_1, T]$, shortages occurred and the demand is partially backlogged. That is, the inventory level at time t is governed by the following differential equation

$$\frac{dq_s(t)}{dt} = \frac{-a}{1 + \delta(T-t)} \quad t_1 \leq t \leq T \quad (3.15)$$

With the condition $q_s(t_1) = 0$ the solution of equation (3.15) is

$$q_s(t) = a(t_1 - t) \left[1 - \delta T + \frac{\delta}{2}(t_1 + t) \right] \quad t_1 \leq t \leq T \quad (3.16)$$

Therefore the maximum inventory level and maximum amount of shortage demand to be backlogged during the first replenishment cycle are

$$S = W_1 + \frac{a\beta}{b(\beta + b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r - \mu_2) + b\mu_2} \right) \right\} \quad (3.17)$$

$$BI = \frac{aH(1-k)}{2m^2} [2m - \delta H(1-k)] \quad (3.18)$$

There are m cycles during the planning horizon. Since, inventory is assumed to start and end at zero, an extra replenishment at $T_m = H$ is required to satisfy the backorders of the last cycle in the planning horizon. Therefore, there are $m + 1$ replenishments in the entire planning horizon H .

The first replenishment lot size is S .

The 2, 3, ..., m^{th} replenishment order size is:- $Q = S + BI$ (3.19)

The last or $(m + 1)^{\text{th}}$ replenishment lot size is BI .

Since replenishment in each cycle is done at the start of each cycle, the present value of ordering cost during the first cycle is $OC = A$ (3.20)

The holding cost for the RW during the first replenishment cycle is

$$HC_r = C_{hr} \left[\int_0^{\mu_2} q_r(t) e^{-Rt} dt + \int_{\mu_2}^{t_r} q_r(t) e^{-Rt} dt \right] \quad (3.21)$$

$$= C_{hr} \left[-\frac{a}{b} \left\{ \frac{1 - e^{-R\mu_2}}{R} \right\} - \left(\frac{a}{b} + W_2 \right) \left\{ \frac{e^{-(R+b)\mu_2} - 1}{R+b} \right\} \right] \\ + C_{hr} \left[\frac{a}{\beta + b} \left\{ e^{-Rt_r} \left(\frac{\beta + b}{R(\beta + b + R)} \right) + e^{-R\mu_2} \left(\frac{e^{(t_r - \mu_2)(\beta + b)}}{\beta + b + R} - \frac{1}{R} \right) \right\} \right] \quad (3.22)$$

The holding cost for the OW during the first replenishment

$$HC_o = C_{ho} \left[\int_0^{\mu_1} q_o(t) e^{-Rt} dt + \int_{\mu_1}^{t_r} q_o(t) e^{-Rt} dt + \int_{t_r}^{\mu_1} q_o(t) e^{-Rt} dt \right] \\ = C_{ho} \left[\frac{W_1}{R} (1 - e^{-R\mu_1}) + \frac{W_1 e^{\alpha\mu_1}}{\alpha + R} (e^{-(\alpha + R)\mu_1} - e^{-(\alpha + R)t_r}) \right] \\ + \frac{a}{\alpha + b} \left\{ e^{-Rt_1} \left(\frac{\alpha + b}{R(\alpha + b + R)} \right) + e^{-Rt_r} \left(\frac{e^{(\alpha + b)(t_1 - t_r)}}{\alpha + b + R} - \frac{1}{R} \right) \right\} \quad (3.23)$$

The deteriorating cost for RW during first replenishment cycle is

$$DC_r = C_2 \int_{\mu_2}^{t_r} \beta q_r(t) e^{-Rt} dt$$

$$= C_2 \frac{a\beta}{\beta+b} \left\{ e^{-Rt_r} \left(\frac{\beta+b}{R(\beta+b+R)} \right) + e^{-R\mu_2} \left(\frac{e^{(t_r-\mu_2)(\beta+b)}}{\beta+b+R} - \frac{1}{R} \right) \right\} \quad (3.24)$$

The deteriorating cost for OW during first replenishment cycle is

$$\begin{aligned} DC_o &= C_2 \alpha \left[\int_{-\mu_1}^{t_r} q_o(t) e^{-Rt} dt + \int_{t_r}^{t_1} q_o(t) e^{-Rt} dt \right] \\ &= C_2 \alpha \left[\frac{W_1 e^{\alpha\mu_1}}{\alpha+R} \left(e^{-\mu_1(\alpha+R)} - e^{-(\alpha+R)t_r} \right) + \frac{a}{\alpha+b} e^{-Rt_1} \left(\frac{\alpha+b}{R(\alpha+b+R)} \right) \right. \\ &\quad \left. + \frac{a}{\alpha+b} e^{-Rt_r} \left(\frac{e^{(t_1-t_r)(\alpha+b)}}{\alpha+b+R} - \frac{1}{R} \right) \right] \end{aligned} \quad (3.25)$$

Total shortage cost during the first replenishment cycle is

$$\begin{aligned} SC &= -C_3 \int_{t_1}^T q_s(t) e^{-Rt} dt \\ &= \frac{-C_3 a}{R} \left\{ e^{\frac{-RH}{m}} \left[\frac{KH^2 \delta}{m^2} \left(1 - \frac{K}{2} - \frac{m}{\delta H} \right) + \frac{H}{m} \left(1 - \frac{\delta H}{2m} \right) + \frac{1}{R} \left(1 + \frac{\delta}{R} \right) \right] \right. \\ &\quad \left. + \frac{e^{\frac{-RKH}{m}}}{R} \left[\frac{\delta H}{m} \left(1 - K - \frac{m}{HR} \right) - 1 \right] \right\} \end{aligned} \quad (3.26)$$

The lost sale cost during the first replenishment cycle is

$$\begin{aligned} LC &= C_4 \int_{t_1}^T \left(1 - \frac{1}{1 + \delta(T-t)} \right) a e^{-Rt} dt \\ &= \frac{-C_4 a \delta}{R^2} \left[e^{\frac{-RKH}{m}} + e^{\frac{-RKH}{m}} \left(\frac{RH}{m} (1-K) - 1 \right) \right] \end{aligned} \quad (3.27)$$

Replenishment is done at $t = 0$ and T . The present value of purchasing cost PC during the first replenishment cycle is:-

$$PC = pS + pe^{-RT} (BI)$$

$$= p \left\{ W_1 + \frac{a\beta}{b(\beta+b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r - \mu_2) + b\mu_2} \right) \right\} + ae^{-\frac{RH}{m}} \frac{H}{M} (1-k) \left[1 + \frac{\delta H}{2m} (K-1) \right] \right\} \quad (3.28)$$

So, the total cost = Ordering cost + inventory holding cost in RW + inventory holding cost in OW + deterioration cost in RW + deterioration cost in OW + shortage cost + lost sales cost + purchasing cost. $TC_F = OC + HC_r + HC_0 + DC_r + DC_0 + SC + LC + PC$

So, the present value of the total cost of the system over a finite planning horizon H is

$$TC(m, k) = \sum_{j=0}^{m-1} TC_f e^{-RjT} + Ae^{-RH} = TC_F \left(\frac{1 - e^{-RH}}{1 - e^{-\frac{RH}{m}}} \right) + Ae^{-RH} \quad (3.29)$$

Where $T = H/m$ and TC_F derived by substituting equations (3.21) to (3.28) in equation (3.29).

On simplification we get

$$\begin{aligned}
 TC(m, k) = & Ae^{-RH} + G \left[A + C_{hr} \left\{ \frac{a}{b} \left(\frac{e^{-R\mu_2} - 1}{R} \right) + \left(\frac{a}{b} + W_2 \right) \left(\frac{e^{-(R+b)\mu_2} - 1}{R+b} \right) \right\} \right. \\
 & + \left(\frac{C_{hr}a}{\beta+b} + \frac{C_2a\beta}{\beta+b} \right) \left\{ \frac{e^{-Rt_r}(\beta+b)}{R(\beta+b+R)} + e^{-R\mu_2} \left(\frac{e^{(t_r-\mu_2)(\beta+b)} - 1}{\beta+b+R} - \frac{1}{R} \right) \right\} \\
 & + C_{ho} \left[\frac{W_1}{R} (1 - e^{-R\mu_1}) + \frac{W_1 e^{\alpha\mu_2}}{\alpha+R} (e^{-(\alpha+R)\mu_1} - e^{-(\alpha+R)t_r}) \right] \\
 & + (C_{ho} + C_2\alpha) \left[\frac{W_1 e^{\alpha\mu_2}}{\alpha+R} (e^{-\mu_1(\alpha+R)} - e^{-(\alpha+R)t_r}) \right] \\
 & + C_2\alpha \left[\frac{a}{R(\alpha+b+R)} e^{\frac{-RKH}{m}} + \frac{a}{\alpha+b} \left(\frac{e^{\left(\frac{KH}{m}-t_r\right)(\alpha+b)}}{\alpha+b+R} - \frac{1}{R} \right) e^{-Rt_r} \right] \\
 & - \frac{C_3a}{R} \left[e^{\frac{-RH}{m}} \left\{ \frac{KH^2\delta}{m^2} \left(1 - \frac{K}{2} - \frac{m}{\delta H} \right) + \frac{H}{m} \left(1 - \frac{\delta H}{2m} \right) + \frac{1}{R} \left(1 + \frac{\delta}{R} \right) \right\} \right. \\
 & \left. + \frac{e^{\frac{-RKH}{m}}}{R} \left\{ \frac{\delta H}{m} \left(1 - K - \frac{m}{HR} \right) - 1 \right\} \right] - \frac{C_4a\delta}{R^2} \left[e^{\frac{-RKH}{m}} - e^{\frac{-RKH}{m}} \left(\frac{RH}{m} (1-K) - 1 \right) \right] \\
 & + p \left[W_1 + \frac{a\beta}{b(\beta+b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r-\mu_2)+b\mu_2} \right) \right\} \right] \\
 & + ae^{\frac{-RH}{m}} \frac{H}{m} (1-K) \left(1 - \frac{\delta H}{2m} (K-1) \right) \Big] \quad \text{Where } G = \left(\frac{1 - e^{RH}}{1 - e^{-RH/m}} \right) \quad (3.30)
 \end{aligned}$$

4. Solution Procedure

The present value of total cost TC (m, k) is a function of two variables m and k where m is a discrete variable and k is a continuous variable. For a given value of m, the necessary condition for TC(m, k) to be minimized is $dTC(m, k) / dk = 0$ which gives

$$\begin{aligned}
 \frac{dTC(m, k)}{dk} = & C_2\alpha \left[\frac{aH e^{\frac{kH}{m}(\alpha+b)}}{m(\alpha+b+R)} - \frac{aH e^{\frac{-RHk}{m}}}{m(\alpha+b+R)} \right] + C_3 \left(\frac{aH}{Rm} e^{\frac{-RH}{m}} \right) + \frac{H^2\delta}{m^2} e^{\frac{-RHk}{m}} (k-1) \\
 & + \frac{H\delta}{mR} e^{\frac{-RHk}{m}} + \frac{H}{m} e^{\frac{-RHk}{m}} - C_4 \frac{\delta a}{R^2} \left[\frac{R^2 H^2}{m^2} e^{\frac{-RHk}{m}} (1-k) - \frac{RH}{m} e^{\frac{-RHk}{m}} \right] \\
 & + P \left[(1-k) \frac{aH}{m} e^{\frac{-RH}{m}} \left\{ 1 - \frac{H\delta}{m} (1+k) \right\} \right] \quad (3.31)
 \end{aligned}$$

$$\begin{aligned} \frac{d^2TC(m,k)}{dk^2} = & C_2\alpha \left[\frac{aH^2}{m^2} \left(\frac{R e^{-\frac{RHk}{m}}}{(\alpha+b+R)} + (\alpha+b)e^{\frac{kH}{m}(\alpha+b)} \right) \right] + \frac{H^3\delta R}{m^3} e^{-\frac{RHk}{m}} (1-k) \\ & - C_4 \frac{\delta a}{R^2} \left[\frac{R^3H^3}{m^3} e^{-\frac{RHk}{m}} (k-1) \right] + P \left[\frac{a\delta H^2}{2m^2} \left(e^{-\frac{RH}{m}} - 1 \right) + \frac{aH}{m} e^{-\frac{RH}{m}} \left(\frac{\delta kH}{m} - 1 \right) \right] \end{aligned} \quad (3.32)$$

Furthermore, the equation (32) shows that TC (m, k) is convex with respect to k. So, for a given positive integer m, the optimal value of k can be obtained from (31).

Algorithm

Step 1: Start with m = 1.

Step 2: Using (31) solve for k. Then substitute the solution obtained for (31) into (30) to compute the total inventory cost.

Step 3: Increase m by 1 and repeat step 2.

Step 4: Repeat step 2 and step 3 until TC (m, k) increases. The value of m which corresponds to the increase of TC for the first time is taken as the optimal value of m (denoted by m*) and the corresponding k (denoted by k*) is the optimal value of k.

Using the optimal solution procedure described above, we can find the optimal order quantity and maximum inventory levels to be

$$W_2 = \frac{a\beta}{b(\beta+b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r - \mu_2) + b\mu_2} \right) \right\}$$

$$S^* = W_1 + \frac{a\beta}{b(\beta+b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r - \mu_2) + b\mu_2} \right) \right\}$$

$$Q^* = W_1 + \frac{a\beta}{b(\beta+b)} \left\{ e^{b\mu_2} - 1 - \frac{b}{\beta} \left(1 - e^{(\beta+b)(t_r - \mu_2) + b\mu_2} \right) \right\} + \frac{aH(1-k)}{2m^2} [2m - \delta H(1-k)]$$

$$\text{Where } t_r = \frac{\left(W_1\alpha e^{\alpha\mu_1} + a e^{(\alpha+b)t_1} \right) \pm \sqrt{\left(W_1\alpha e^{\alpha\mu_1} + a e^{(\alpha+b)t_1} \right)^2 + \frac{4a\alpha b}{\alpha+b} \left(W_1 e^{\alpha\mu_1} - \frac{a}{\alpha+b} e^{(\alpha+b)t_1} + \frac{a}{\alpha+b} \right)}}{2 \left(\frac{-a\alpha b}{\alpha+b} \right)}$$

5. Numerical Examples

Example 1

Consider an inventory system with the following data: $D = 100$ units; $W_1 = 50$ units; $p = \$4$; $s =$

\$15; $A = \$150$; $C_{hr} = \$2$; $C_{ho} = \$1.2$; $C_2 = \$1.5$; $C_3 = \$5$; $C_4 = \$10$; $\alpha = 0.8$; $\beta = 0.2$; $\delta = 0.008$; $\mu_1 = 5/12$ year; $\mu_2 = 8/12$ year; $R = 0.2$; $H = 20$ years.

6. Implementation of Genetic Algorithm

A genetic algorithm (GA) is a based on natural selection process to optimized tools that minimizes the total costs in supply chain management. It is a evolutionary computation method to solve a inventory problems. This is the more effective methods to find the optimized solution. The genetic algorithm uses three main types of rules at each step to create the next generation from the current population.

The basic steps to find the optimized solution:

Step 1. First one is *Selection rules*, In this we select the individuals, called *parents* that contribute to the population at the next generation.

Step 2. Next one is *Crossover rules*, In this we perform crossover operation between two parents to form children for the next generation.

Step 3. Last one is *Mutation rules*, In mutation we apply some random changes to individual parents.

We will perform these steps till we will not get our optimized solution.

GA is not a method to find the exact solution of problem it only help to find the best or optimized solutions.

Here we are implementing Genetic Algorithm in Table 1: Optimal total cost with respect to m and Table 2: Optimal total cost with respect to m when $\mu_1=0$ and $\mu_2=0$

Table 1: Optimal total cost with respect to m

M	k(m)	t _r	t ₁	T	Q	TC (m,
1	0.4485	8.9519	8.8552	19	3922	26496
2	0.4993	4.8536	4.6241	9.5	1482	11060
M	k(m)	t _r	t ₁	T	Q	TC (m,
1	0.3582	1.9312	1.6842	2.76	452	5658
2	0.2995	0.8767	1.3582	2.653	409	5594

Before Crossover:

		C1 ↓		C2 ↓		
M	k(m)	t_r	t_1	T	Q	TC (m,
1	0.4485	8.9519	8.8552	19	3922	26496
2	0.4993	4.8536	4.6241	9.5	1482	11060

After Crossover

		C1 ↓		C2 ↓		
M	k(m)	t_r	t_1	T	Q	TC (m,
1	0.4485	4.8536	4.6241	19	3922	26597
2	0.4993	8.9519	8.8552	9.5	1482	10090

Mutation

Before mutation

		M1 ↓		M2 ↓		
M	k(m)	t_r	t_1	T	Q	TC (m,
1	0.4485	4.8536	4.6241	19	3922	26597

After mutation

		M1 ↓		M2 ↓		
M	k(m)	t_r	t_1	T	Q	TC (m,
1	0.4485	4.6241	4.8536	19	3922	24384

Before Crossover:

		C1 ↓	C2 ↓			
M	k(m)	t_r	t₁	T	Q	TC (m, k)
1	0.3582	1.9312	1.6842	2.76	452	5658
2	0.2995	0.8767	1.3582	2.653	409	5594

After Crossover

		C1 ↓	C2 ↓			
M	k(m)	t_r	t₁	T	Q	TC (m, k)
1	0.3582	0.8767	1.3582	2.76	452	5703
2	0.2995	1.9312	1.6842	2.653	409	5385

Mutation

Before mutation

		M1 ↓	M2 ↓			
M	k(m)	t_r	t₁	T	Q	TC (m, k)
1	0.3582	0.8767	1.3582	2.76	452	5703

After mutation

		M1 ↓	M2 ↓			
M	k(m)	t_r	t₁	T	Q	TC (m, k)
1	0.3582	1.3582	0.8767	2.76	452	5492

Table 2: Optimal total cost with respect to m when $\mu_1=0$ and $\mu_2=0$

M	k(m)	t_r	t_1	T	Q	TC (m, k)
1	0.4170	8.4331	8.6052	19	4022	27496
2	0.4593	4.4536	4.6281	9	1532	11865
M	k(m)	t_r	t_1	T	Q	TC (m, k)
1	0.2782	0.8312	2.0042	1.763	612	6758
2	0.2098	0.5167	1.3249	1.953	456	6690

Before Crossover:

C1 ↓ C2 ↓

M	k(m)	t_r	t_1	T	Q	TC (m, k)
1	0.4170	8.4331	8.6052	19	4022	27496
2	0.4593	4.4536	4.6281	9	1532	11865

After Crossover

C1 C2
↓ ↓

M	k(m)	t_r	t_1	T	Q	TC (m, k)
1	0.4170	4.4536	4.6281	19	4022	26585
2	0.4593	8.4331	8.6052	9	1532	10336

Mutation

Before mutation

M1 M2
↓ ↓

M	k(m)	t_r	t_1	T	Q	TC (m, k)
1	0.4170	4.4536	4.6281	19	4022	26585

After mutation

M1

M2




M	k(m)	t _r	t ₁	T	Q	TC (m, k)
1	0.4170	4.6281	4.4536	19	4022	24348

Before Crossover:



M	k(m)	t _r	t ₁	T	Q	TC (m, k)
1	0.2782	0.8312	2.0042	1.763	612	6758
2	0.2098	0.5167	1.3249	1.953	456	6690

After Crossover



M	k(m)	t _r	t ₁	T	Q	TC (m, k)
1	0.2782	0.5167	1.3249	1.763	612	6597
2	0.2098	0.8312	2.0042	1.953	456	6385

Mutation

Before mutation



M	k(m)	t _r	t ₁	T	Q	TC (m, k)
1	0.2782	0.5167	1.3249	1.763	612	6597

After mutation

M1

M2



M	k(m)	t_r	t₁	T	Q	TC (m, k)
1	0.2782	1.3249	0.5167	1.763	612	6376

It is clearly visible the changes in total cost after the mutation methods and before crossover and it are very effective cost.

Based on our numerical results, we obtain the following decision-making phenomena:

(i) The total cost is increasing while the time horizon H is increasing, the order quantity is decreasing. Furthermore it is observed that when the time horizon increases, the number of replenishment also increases. Therefore the ordering cost increases. In order to minimize the total cost, the retailer should decrease the time horizon.

(ii) When the total cost of the retailer and the order quantity are also increasing then the deterioration rate β in RW increasing. When the deterioration rate will increase, the total cost of the retailer will increase.

3. If backlogging parameter δ is increased then the total cost and the order quantity will be decreased. But there is no change in the number of replenishment. If the backlogging parameter increases, then the ordering quantity will decrease.

4. When the inflation rate increases, the number of replenishment also increases. then the net discount rate of inflation R is increasing, the optimal cost is decreasing and the order quantity is also decreasing.

5. When the purchasing price p is increasing, the total optimal cost and the order quantity are highly increasing. When the purchase price increases, the number of replenishment decreases. Also, the increasing of purchasing price will increase the total cost of the retailer.

7. Conclusion:

In the present model, we have considered two warehouse inventory models depending on the waiting time for the next replenishment. And shortages and partially backlogging are allowed. In this model, demand rate considered as stock dependent with inflation and model affected by

Genetic Algorithm. An algorithm is designed to find the optimum solution of the proposed model. The model shows that the minimum time horizon will minimize the total cost of the retailer. Furthermore, sensitivity analysis is carried out with respect to the key parameters and helpful decision-making insights are obtained. The graphical illustrations are also given to analyze the efficiency of the model clearly. The proposed model incorporates some practical features that are likely to be linked with some kinds of inventory. additionally this model can be adopted in the inventory control of retail business such as food industries, seasonable cloths, domestic goods, automobile, electronic components etc. The proposed model can be extended in several ways. Like incorporate some more realistic features, such as quantity discount, multi item, trade credit strategy, etc., when the net discount rate of inflation and the backlogging rate are increased then the optimal total cost will be decreased. Also, sensitivity analysis of the model with respect to numerous system parameters has been carried out a number of decision-making.

References

1. Hartley R.V. (1976). *Operations Research—A Managerial Emphasis*. Good Year Publishing Company, California, 315-317
2. Sarma KVS (1983) A deterministic inventory model with two levels of storage and an optimum release rule. *Opsearch* 20:175-180
3. Sarma KVS, Sastry MP (1988) Optimum inventory for systems with two levels of storage. *Industrial Engineering Journal* 8:12-19
4. Pakkala, T.P.M., Achary, K.K., (1992). A deterministic inventory model for deteriorating items with two warehouses and finite replenishment rate. *European Journal of Operational Research*, 57(1), 71-76
5. Lee C, Ma C (2000) optimal inventory policy for deteriorating items with two-warehouse and timedependent demands. *Production Planning and Control* 11:689–696
6. Yang HL (2004) Two-warehouse inventory models for deteriorating items with shortages under inflation. *European Journal of Operational Research* 157:344–356
7. Yang HL (2006) Two-warehouse partial backlogging inventory models for deteriorating items under inflation. *International Journal of Production Economics* 103:362–370
8. Kumar N, Singh S R, Kumari R (2008) a Two-Warehouse inventory model without shortage for exponential demand rate and an optimum release rule. *Ultra Scientist of Physical Sciences* 20:395-402
9. Kumar N, Singh S R, Kumari R (2011) a deterministic Two-warehouse Inventory Model for Deteriorating Items with Sock-dependent Demand and Shortages under the Conditions of

- Permissible Delay. *International Journal of Mathematical Modeling and Numerical Optimizations* 11:357-375
10. Yang HL (2012) Two-warehouse partial backlogging inventory models with three-parameter Weibull distribution deterioration under inflation. *International Journal of Production Economics* 138:107–116
 11. Sarkar T, Ghosh SK, Chaudhuri KS (2012) An optimal inventory replenishment policy for a deteriorating item with time-quadratic demand and time-dependent partial backlogging with shortages in all cycles. *Applied Mathematics and Computation* 218:9147–9155
 12. Kumar N, Singh S R, Kumari R (2013) Learning effect on an inventory model with two-level storage and partial backlogging under inflation. *International Journal of Services and Operations Management* 16:105 – 122
 13. Yang HL, Chang CT (2013) A two-warehouse partial backlogging inventory model for deteriorating items with permissible delay in payment under inflation. *Applied Mathematical Modelling* 37:2717–2726
 14. Guchhaita, P., Maiti ,M.K., & Maiti, M., (2013). Two storage inventory model of a deteriorating item with variable demand under partial credit period. *Applied Soft Computing*, 13(1), 428–448
 15. Kumar N, Singh S R, Kumari R (2014) Effect of Salvage Value on a Two-Warehouse Inventory Model for Deteriorating Items with Stock-Dependent Demand Rate and Partial Backlogging .*International Journal of Operational Research* 19: 479 - 496
 16. Ghare, P.M., Schrader, G.H., (1963). A model for exponentially decaying inventory system. *International Journal of Production Research*, 21, 449-460
 17. Philip GC (1974) A generalized EOQ model for items with weibull distribution. *AIIE Transactions* 6:159-162
 18. Deb, M., Chaudhuri, K.S., (1986). An EOQ model for items with finite rate of production and variable rate of deterioration. *Opsearch*, 23(1), 175-181
 19. Goyal, S.K., Giri, B.C., (2001). Recent trends in modeling of deteriorating inventory. *European Journal of Operational Research*, 134(1), 1-16
 20. Liao J (2008) An EOQ model with non-instantaneous receipt and exponential deteriorating item under two-level trade credit. *International Journal of Production Economics* 113:852–861
 21. Chung, K.J., (2009). A complete proof on the solution procedure for non-instantaneous deteriorating items with permissible delay in payment. *Computers & Industrial Engineering*, 56(1), 267–273
 22. Chang, CT., Teng, JT., & Goyal, SK., (2010). Optimal replenishment policies for non-instantaneous deteriorating items with stock-dependent demand. *International Journal of Production Economics*, 123(1), 62–68
 23. Dye, C.Y., (2013). The effect of preservation technology investment on a non-instantaneous deteriorating inventory model. *Omega*, 41(5), 872–880

24. Buzacott, JA. , (1975). Economic order quantities with inflation. *Operations Research Quarterly*, 26(3), 553–558
25. Misra RB (1979) A note on optical inventory management under inflation. *Naval Research Logistics Quarterly* 26:161–165
26. Data, T.K., Pal, A.K., (1991). Effects of inflation and time value of money on an inventory model with linear time dependent demand rate and shortages. *European Journal of Operational Research*, 52(3), 326–333
27. Sarker BR, Pan H (1994) Effects of inflation and time value of money on order quantity and allowable shortage. *International Journal of Production Economics* 34:65–72
28. Cheng, M., Zhang, B., & Wang, G., (2011). Optimal policy for deteriorating items with trapezoidal type demand and partial backlogging. *Applied Mathematical Modelling*, 35(7), 3552–3560
29. Tolgari JT, Mirzazadeh A, Jolai A (2012) An inventory model for imperfect items under inflationary conditions with considering inspection errors. *Computers and Mathematics with Applications* 63:1007– 1019
30. Guria, A., Das, B., Mondal, S.,& Maiti, M., (2013). Inventory policy for an item with inflation induced purchasing price, selling price and demand with immediate part payment. *Applied Mathematical Modelling*, 37(1), 240–257
31. Dye, C.Y., Ouyang, L.Y., & Hsieh, T.P., (2007). Deterministic inventory model for deteriorating items with capacity constraint and time-proportional backlogging rate. *European Journal of Operational research*, 178(3), 789-807
32. Wu KS, Ouyang LY, Yang CT (2006) An optimal replenishment policy for non-instantaneous deteriorating items with stock-dependent demand and partial backlogging. *International Journal of Production Economics* 101:369-384
33. Kumar N and Kumar S (2016) Effect of learning and salvage worth on an inventory model for deteriorating items with inventory-dependent demand rate and partial backlogging with capability constraints. *Uncertain Supply Chain Management* 4: 123-136

The main aim of the present book is to suggest some improved estimators using auxiliary and attribute information in case of simple random sampling and stratified random sampling and some inventory models related to capacity constraints.

This volume is a collection of six papers, written by five co-authors (listed in the order of the papers): Dr. Rajesh Singh, Dr. Sachin Malik, Dr. Florentin Smarandache, Dr. Neeraj Kumar, Mr. Sanjey Kumar & Pallavi Agarwal.

In the first chapter authors suggest an estimator using two auxiliary variables in stratified random sampling for estimating population mean. In second chapter they proposed a family of estimators for estimating population means using known value of some population parameters. In Chapter third an almost unbiased estimator using known value of some population parameter(s) with known population proportion of an auxiliary variable has been used. In Chapter four authors investigate a fuzzy economic order quantity model for two storage facility. The demand, holding cost, ordering cost, storage capacity of the own - warehouse are taken as a trapezoidal fuzzy numbers and in Chapter five a two-warehouse inventory model deals with deteriorating items, with stock dependent demand rate and model affected by inflation under the pattern of time value of money over a finite planning horizon. Shortages are allowed and partially backordered depending on the waiting time for the next replenishment. The purpose of this model is to minimize the total inventory cost by using Genetic algorithm.

This book will be helpful for the researchers and students who are working in the field of sampling techniques and inventory control.

